Effects of ancient anthropogenic clam gardens on the growth, survival, and transcriptome of Pacific littleneck clams (*Leukoma staminea*)

Monique R. Raap[1,2,a], Helen J. Gurney-Smith[1,2,3,b], Sarah E. Dudas[1,2,3,a], Christopher M. Pearce[4,5], Jong S. Leong[1], Ben J. G. Sutherland[1,6], Ben F. Koop[1]

[1] Department of Biology, University of Victoria, Victoria, British Columbia, Canada, V8W 2Y2

[2] Hakai Institute, Quadra Island, British Columbia, Canada, V0P 1H0

[3] Department of Biology and Department of Fisheries and Aquaculture, Vancouver Island University, Nanaimo, British Columbia, Canada, V9R 5S5

[4] Pacific Biological Station, Fisheries and Oceans Canada, Nanaimo, British Columbia, Canada, V9T 6N7

[5] Department of Geography, University of Victoria, Victoria, British Columbia, Canada, V8W 2Y2

[6] Sutherland Bioinformatics, Lantzville, British Columbia, Canada, V0R 2H0

[a] Present address: Pacific Biological Station, Fisheries and Oceans Canada, Nanaimo, British Columbia, Canada, V9T 6N7

[b] Present address: St. Andrews Biological Station, Fisheries and Oceans Canada, St. Andrews, New Brunswick, Canada, E5B 0E4

Author for correspondence: HJG-S

St. Andrews Biological Station, Fisheries and Oceans Canada, St. Andrews, New Brunswick, Canada, E5B 0E4

Email: Helen.Gurney-Smith@dfo-mpo.gc.ca

27   Data Deposition: Raw RNA-sequencing data have been uploaded to SRA under BioProject

28   PRJNA818991, BioSamples SAMN26893882–SAMN26893909.

Abstract

Clam gardens traditionally established and maintained by coastal Indigenous Peoples of northwest North America are habitat modifications to enhance intertidal clam productivity for reliable local food production. In this study, phenotypic and transcriptomic responses of Pacific littleneck clams (*Leukoma staminea*) were evaluated 16 weeks after transplantation to either unmaintained clam gardens or reference (unmodified) clam beaches. Beach sediment characteristics including grain size and organic content were examined across all beaches. Large differences in abiotic characteristics and phenotypic responses were observed among beaches; however, differences were not related to the clam garden/reference beach effect. Clam survival and growth were negatively associated with small rocks, very fine sand, and silt, along with carbonate and organic content, and positively associated with coarse sand, sand, and fine sand. To investigate molecular responses to unmaintained clam gardens, a *de novo* transcriptome containing 52,000 putative transcripts was assembled for *L. staminea* and was used to test for differential expression between transplanted clams in unmaintained clam gardens or on reference beaches. As expected, given the lack of significant phenotypic differences between treatments, transcriptomic responses to unmaintained clam gardens were minor, although several weakly-associated transcripts were identified. By contrast, the strong survival gradient across beaches was used to identify genes associated with survival and, combined with characterization of tissue-specific expression in the gill and digestive gland, contributes to our understanding of molecular processes in this non-model species.

Keywords: clam garden; *Leukoma staminea*; Pacific littleneck clam; sediment carbonate content; transcriptomics

Introduction

Coastal shorelines have been altered to increase productivity and facilitate the harvesting of natural resources for centuries (Erlandson et al., 2008). Shoreline constructions such as clam gardens and fishing weirs have long been used to provide productive and predictable local food resources for coastal Indigenous Peoples (Neudorf et al., 2017) and provide important insights into cultural food practices, traditional technologies, economies, values, and ancestral practices of coastal communities (Deur et al., 2015; Jackley et al., 2016; Lepofsky et al., 2020; Smith et al., 2019). Construction of clam gardens in North America is estimated to have begun approximately 3,500 years ago (Smith et al., 2019) and clam gardens are found from Alaska (USA), through British Columbia (BC, Canada), to Washington State (USA) (Groesbeck et al., 2014).

Construction of clam gardens involves rolling rocks and boulders to the low-tide line and building a wall parallel to the shoreline (Deur et al., 2015), allowing sediment deposition between the wall and the high-tide mark. This creates a wider intertidal shellfish habitat area with a reduced slope (Neudorf et al., 2017) where shellfish such as Pacific littleneck clams (*Leukoma staminea*) typically occur (Deur et al., 2015; Groesbeck et al., 2014). The reduced slope allows for a thin layer of seawater to be retained on the accumulated sediment, reducing desiccation risk, keeping clams in shallow accessible portions of the intertidal zone, and maximizing submersion time for feeding and therefore growth (Deur et al., 2015). Furthermore, the exterior, ocean-facing side of the wall creates rocky reef habitat for a wide variety of other marine invertebrates, many of which can also be harvested for consumption (Caldwell et al., 2012; Deur et al., 2015; Lepofsky et al., 2017). Traditionally, clam gardens were maintained and tended with practices that included predator exclusion, selective harvesting, and the addition of gravel and crushed shell or shell hash (Deur et al., 2015). The latter may provide settlement cues to larval shellfish as well as protect young recruits from seawater acidity and predators (Green et al., 2012; Sponaugle & Lawton, 1990).

Research on the effects of clam gardens is an area of active interest. Many clam garden walls, although maintained into the 20[th] century (Williams, 2006), now remain as unmaintained structures, which could be expected to provide only partial effects relative to a fully tended clam garden. Nonetheless, untended clam gardens have been found to harbour higher densities of Pacific littleneck clams and butter clams (*Saxidomus giganteus*), to have increased recruit survival, and higher growth rates relative to unmodified beaches (Groesbeck et al., 2014; Jackley et al., 2016). Clam gardens also support distinct

biological communities with increased abundances of other infaunal taxa including Nematoda,
Harpacticoida, and Chironomidae (Cox et al., 2019), and epifaunal taxa such as *Chthamalus dalli*, *Lottia*
persona, *Balanus crenatus*, and *Littorina scutulata*, (Cox et al., 2024a). These increases in diversity and
density were positively correlated with the quantity of gravel and shell hash present (Cox et al., 2024a;
Cox et al., 2019). Sediment composition can be altered in clam gardens, for example through increased
carbonate and reduced silt (Salter, 2018), as well as more gravel and shell hash (Groesbeck et al., 2014)
relative to unmodified beaches, which are often primarily composed of silt, sand, and mud (Jackley et al.,
2016). Such differences in sediment composition are attributed to the tending practices of Indigenous
Peoples, which included the addition of gravel and crushed whole clam shells to clam gardens (Groesbeck
et al., 2014; Williams, 2006).

The effect of sediment shell hash on clam productivity is not definitive. Sediment carbonate
saturation state can affect settlement and recruitment of larval and juvenile bivalves, respectively (Green et
al., 2012). Seawater calcium carbonate ($CaCO_3$) saturation state is important to the formation and
maintenance of shell thickness and integrity of marine molluscs and other calcifiers (Chadwick et al.,
2019; Evans et al., 2014; Green et al., 2012; Waldbusser & Salisbury, 2014). Increased sediment shell
hash has been observed to increase clam settlement and productivity (Green et al., 2012; Groesbeck et al.,
2014; Salter, 2018), but observations have also been made showing negative (Munroe, 2016) or neutral
(Greiner et al., 2018) effects on early post-settlement bivalve growth and recruitment.

Genomic tools such as transcriptomics can capture early signals of environmental stressors and are
increasingly used to evaluate abiotic impacts on aquatic species (Milan et al., 2011; Sutherland et al.,
2012). Transcriptomic data can be associated with phenotypic (*e.g.*, survival, growth) and environmental
data and used to predict potential physiological outcomes (Evans et al., 2011; Miller et al., 2017). An
additional benefit is that in some studies, metatranscriptomics can provide additional taxonomic
identification of microbiota in samples (Bourlat et al., 2013; Sutherland et al., 2022). The present study
aims to determine whether Pacific littleneck clam health and productivity—as assessed by growth,
survival, and transcriptomic response—are affected when clams are transplanted to unmaintained clam
gardens relative to unmodified clam beaches, and whether these unmaintained clam gardens contain
significantly different abiotic conditions. To this end, we generated and assembled a *de novo* reference
transcriptome and characterized Pacific littleneck clam transcriptome responses in several unmaintained
clam gardens and unmodified beaches in the same area. Beaches used in the study were in northern

Quadra Island in coastal BC, an area with a very high density of historic clam gardens, covering

approximately 35% of the shoreline (Groesbeck et al., 2014; Lepofsky et al., 2020; Neudorf et al., 2017).

To evaluate the effects of the unmaintained clam gardens on transplanted clams after a 16-week growth

period, measured environmental parameters, clam phenotypic responses (growth and survival), and clam

transcriptomic responses were considered. The combination of abiotic measurements and organismal and

molecular phenotypes with the characterized transcriptome provides valuable context for both the effects

of different environmental variables on Pacific littleneck clams and on the functions of response genes in

this non-model, ecologically and culturally important species.

Methods

Positionality Statement

The authors of this paper are academic, government, and independent scientific researchers who conduct

research on climate change, ecology, invertebrate physiology, bioinformatics, and genomics. The authors

were educated and trained in North American and European institutions prior to the positions in Canada

that they hold at time of submission, with five reaching doctorate status and two with master's degrees.

This research was conducted in British Columbia, Canada. None of the authors who have provided

heritage information for this statement indicated First Nation or other Indigenous background, and

therefore we respectfully acknowledge the potential for unconscious bias on the presented subject matter.

2.1 | Study sites

The intertidal coastlines of Kanish Bay and adjoining Small Inlet on northwest Quadra Island, BC (Figure

1), were chosen for this study due to an abundance of historic clam gardens where sea level has remained

relatively consistent through time, ensuring walls were still at an ecologically optimal height, and due to

the proximity to the Hakai Institute research station that supported field logistics. Also, complimentary

research has been conducted on clam gardens in the same region and on many of the same sites (Cox et

al., 2024a; Cox et al., 2024b; Groesbeck et al., 2014; Lepofsky et al., 2020).

Quadra Island is located off the northeast coast of Vancouver Island and is in the traditional

territories of the Kwakwaka'wakw (Laich-kwil-tach) and northern Coast Salish Peoples (Smith et al.,

2019). Three unmaintained clam garden beaches (CG, designated as A, C, and E) were selected for the

study along with three unmodified reference clam beaches (Ref, designated as B, D, and F) (Figure 1).

Beaches with similar exposure, slope, and sediment (all determined in the field visually) were sought to
reduce the influence of other variables as much as possible. Once three reference sites were identified,
nearby clam gardens with intact walls were selected. Additionally, all sites were pre-surveyed for the
presence of live bivalve populations using test digs. After the study began, beach B was identified as a
damaged clam garden with over 90% of its wall destroyed most likely due to logging activities in the early
1900s that involved dragging log bundles down the beach into the ocean (Twindle, 1918). Now only a
small remnant of the wall is visible at the very edges at tidal heights below Mean Lower Low Water
(MLLW). The clam gardens are not believed to have been tended as per Indigenous practices since the
early-to-mid 1900s (Deur et al., 2015).

2.2 | Clam transplantation experiment
Juvenile Pacific littleneck clams (N = 400) of 1–2 cm in shell height (hinge to shell margin) were hand
dug from a site in Kanish Bay (Figure 1) between 1 and 2 m MLLW level on May 7, 2016. Each clam was
individually wet weighed, measured for shell height and length (*i.e.*, the widest part of the shell at 90° to
the shell height), and then haphazardly placed into 18 groups of 20 clams each. Clams were held in a pearl
net below the low intertidal zone in Heriot Bay (Quadra Island) until deployment. Clams on beach A were
deployed on May 10, those on beaches B, C, and D on May 11, and those on beaches E and F on May 12,
2016, different deployment days being required due to tidal heights and weather conditions. At low tide at
each of the six beaches, the 1.5–1.8 m intertidal zone was demarcated and three plastic mesh cubes (length
x diameter x height: 20 x 20 x 4 cm; made from black high-density polyethylene, Vexar™), each
containing 20 clams, were buried in the top 20 cm of the sediment, 5 m apart and parallel to the shore in
the middle of the demarcated zone. Each mesh cube was defined as one replicate plot (P) and uniquely
numbered: Beach A with P1–P3, Beach B with P10–P12, Beach C with P4–P6, Beach D with P7–P9,
Beach E with P13–P15; and Beach F with P16–P18.
Transplanted clams were left *in situ* for 16 weeks and collected on August 30–31, 2016. Each mesh
cube, still containing both sediment and clams, was carefully removed from the beach with minimal
disturbance. The cubes were then individually placed in bags and transported in coolers to the Hakai
Institute laboratory on Quadra Island. Upon arrival at the laboratory, each individual was processed as
quickly and carefully as possible. Immediately after wet weight, shell height, and shell length were

measured, and the status (alive or dead) recorded, tissue samples were collected for gene expression
analysis.

Growth and survival per plot were calculated as:

Percent growth = (mean final height – mean initial height) / mean initial height x 100

Percent survival = number of survivors / initial number of clams x 100

In addition, some visual observations were made on animal condition, following the protocols developed

for the Canadian National Animal Aquatic Health Program (NAAHP), including observations of animal

state (*i.e.*, health/vitality, condition) and digestive gland colouration.

2.3 | Sediment collection and analysis

Sediment-core samples (length x diameter: 15 x 5 cm) were collected adjacent to each deployed cube (*n* =

3 cores per beach) to assess grain size. Additional sediment-core samples (10 x 3.5 cm) were collected

adjacent to each cube to assess percent carbonate and percent organic matter content (*n* = 3). Each core

was placed individually in labeled sample bags and frozen at -20°C until analysis.

To determine grain-size distribution, the sediment samples were dried to constant weight at 100°C for

24 hr, weighed, transferred to a series of nested sieves (mesh size: 4.75 mm, 2 mm, 1 mm, 500 µm, 250

µm, 125 µm, and 63 µm), and shaken in a sediment shaker for 15 min. Each size fraction was then

weighed and expressed as a percentage of the total sediment weight. A scale very similar to the

Wentworth scale of sediment size classes was used to classify the grain sizes: pebble and larger gravel

called "rocks" in the present study (> 4.75 mm), granule gravel called "small rocks" in the present study

(2.00 – 4.75 mm), very coarse sand (1.00 – 2.00 mm), coarse sand (0.50 – 1.00 mm), sand (250 – 500

µm), fine sand (125 – 250 µm), very fine sand (63 – 125 µm), and silt (< 63 µm) (Wentworth, 1922).

The loss on ignition method (Heiri et al., 2001) was used to determine sediment percent organic

matter and carbonate content. Briefly, the sediment samples were dried to constant weight at 100°C for 48

hr in previously-ashed (450°C for 8 hr) crucibles, and the dry weights measured. Dried sediment samples

were then ashed in a muffle furnace at 435°C for 8 hr. After cooling the samples in a desiccator for 1 hr,

the weights of the ashed samples were recorded. The percent organic matter was then calculated using the

following equation:

Percent organic matter = (dry weight – ash weight) / dry weight x 100

Sediment percent carbonate content was determined following 2 hr at 950°C and 2 hr of cooling in a

desiccator, and calculated using the equation:

Percent carbonate content = (ash weight$_{435}$ – ash weight$_{950}$) / ash weight$_{435}$ x 100

2.4 | Data analysis: clam size, growth and survival, and sediment characteristics

Data were analyzed in R v.4.4.2. (R.Core.Team, 2025). Model assumptions of normality were assessed

using histograms, quantile-quantile plots of the data, and the Shapiro-Wilk test. Homogeneity of the

residuals was assessed using plots of model residuals versus fitted values and Bartlett's test for

homogeneity of variance. Nested ANOVAs (plots nested within beach location), using the linear mixed-

effects function in the *nlme* package, were used to test for differences between clam garden and

unmodified reference beaches (*i.e.*, beach type, two levels) in initial and final clam shell heights, wet

weights, percent growth, and percent survival as well as individual sediment characteristics (*i.e.*, rocks,

small rocks, very coarse sand, coarse sand, sand, fine sand, very fine sand, silt, percent organic matter, and

percent carbonate concentration), with beach location (six levels) included as a random factor.

Significance was considered when $p \leq 0.05$. ANOVAs, linear mixed-effects models, and post-hoc least

squares means omitting beach type were then used to test for significant differences among beaches in

percent growth, percent survival, sediment carbonates, organics, and individual grain sizes. Principal

component analysis (PCA) and biplot figures were generated in R using the prcomp function in the *stats*

package and the *ggbiplot* function/package. Pearson correlation between the variables was evaluated with

the cor function of the *stats* package of R and inspected with *corrplot* v.0.92 (Wei & Simko, 2021). All

code for the analysis is available (see *Data Availability*).

2.5 | RNA sampling and extraction

For gene-expression analysis, small sections (~2 x 2 mm) of gill and digestive gland tissues were excised

from each surviving clam using sterile techniques and stored in RNAlater as per manufacturer's (Ambion)

protocols. These two tissues were selected because the bivalve gill is often used to study rapid

environmental responses and the digestive gland is an accumulatory organ often used in toxicological and

immunology studies for longer-term effects of environmental change and exposure (Gosling, 2008; Milan

et al., 2011). Total RNA from tissue sections of 25–30 mg was individually extracted from each tissue by

homogenization in 2-mL tubes containing Lysing Matrix D (MP Biomedicals) in a Tissuelyser II (Qiagen)

at 25 Hz for 2 min followed by purification through RNeasy columns (Qiagen). To eliminate DNA

contamination, a DNase protocol was applied (Turbo DNA-free Kits, Ambion). After DNase treatment,

total RNA was quantified by spectrophotometry (NanoDrop 1000, Thermo Fisher Scientific).

2.6 | Library preparation for RNA sequencing

Up to three pooled samples per beach were created for each of the six beaches by combining five

haphazardly selected individuals from each plot within the beach. Plots were only used if they contained at

least five surviving clams. This resulted in three plots being used per beach for all locations except

beaches E and F, which only had five or more survivors in a single plot per beach. This resulted in 14

RNA pools per tissue (N = 28 pools total) to be used for library synthesis and sequencing. Five total RNA

samples were normalized to generate equimolar pools of RNA for each tissue for the 14 plots across the

six beaches. Pooled RNA quality and quantity were evaluated using the RNA 6000 Nano chip on a 2100

Bioanalyzer (Agilent).

Libraries were generated using 250 ng of pooled total RNA using an mRNA enrichment with the

NEBNext Poly(A) Magnetic Isolation Module (New England Biolabs) and cDNA synthesis was

conducted using NEBNext RNA First Strand Synthesis and NEBNext Ultra Directional RNA Second

Strand Synthesis Modules. Remaining library preparation used an NEBNext Ultra II DNA Library Prep

Kit for Illumina, with adapters and PCR primers (NEB). The remaining steps of library preparation were

taken using the Quant-iT^{TM} PicoGreen$^®$ dsDNA Assay Kit (Life Technologies) and the Kapa Illumina GA

with Revised Primers-SYBR Fast Universal Kit (Kapa Biosystems (Pty) Ltd.). Average fragment size was

determined using a LabChip GX instrument (PerkinElmer). Libraries were sequenced across four lanes of

an Illumina HiSeq4000 PE 100 platform (Illumina) and the mean ($\pm$ SD) read pairs per library was 54.4 $\pm$

9.5 million. All library preparation and sequencing steps were undertaken at the Genome Québec

Innovation Centre (Montreal, Canada).

2.7 | Transcriptome assembly and annotation

The 28 sequenced libraries were used to generate a *de novo* transcriptome assembly using Trinity 2.5.1 (--

min_kmer_cov 2) (Grabherr et al., 2011). Before assembly, reads were trimmed to remove adapters, low

quality bases, and too-short reads using Trimmomatic 0.36 (ILLUMINACLIP:TruSeq3-PE-2.fa:2:30:10

LEADING:3 TRAILING:3 SLIDINGWINDOW: 4:15 MINLEN:36) (Bolger et al., 2014). The

TransDecoder 5.0.1 pipeline (Haas et al., 2013) was used to predict likely coding sequences in the

resultant assembly using homology (pfam-a, Sept. 2017 release, UniProt) and open reading frame (ORF)

information. A minimum cutoff for ORFs was 30 amino acids instead of the default 100 amino acids. The

best representative transcript was selected for each gene based on TransDecoder predicted 'complete'

ORF type, which required protein homology. In cases where there were multiple transcripts with complete

ORFs for a gene, the transcript with the largest ORF was chosen. Further filtering was carried out to

choose only transcripts ≥ 100 base pairs to remove small transcripts that could cause analysis errors, and

scanning pfam-a and Uniprot annotations were used to remove repetitive element-related keywords to

produce the final assembly, where each transcript putatively represents a single gene.

2.8 | Data analysis: RNA sequencing data

Raw reads were imported to the repository *Simple_reads_to_counts* (see *Data Availability*) for

quantitation. Raw reads were gently trimmed for quality and to remove adapters using Trimmomatic

(Bolger et al., 2014), with the following flags: ILLUMINACLIP:$VECTORS:2:30:10;

SLIDINGWINDOW:20:2; LEADING:2; TRAILING:2; MINLEN:80. Both the raw and the trimmed data

were inspected using FastQC (Andrews, 2010) and multiQC (Ewels et al., 2016). The *de novo* reference

transcriptome (see above) was indexed using bowtie2 (Langmead & Salzberg, 2012) and reads were

aligned against it in local mode, allowing 40 alignments to be retained per read (-k 40). To compare with

these results, the Manila clam (*Ruditapes philippinarum*) genome (GCA_009026015.1_ASM902601v1)

(Yan et al., 2019) was also tested as a potential reference by indexing the genome using hisat2 (Kim et al.,

2015) and aligning the Pacific littleneck clam reads against the Manila clam genome (also retaining a

maximum of 40 alignments per read; -k 40). The alignments against the Pacific littleneck clam *de novo*

reference transcriptome were quantified using eXpress using default parameters (Roberts & Pachter,

2013). The alignments against the Manila clam genome were also quantified using eXpress default

parameters but with the flag *--rf-stranded*. Resultant effective read counts quantified against the *de novo*

reference transcriptome were used in downstream analyses.

Transcriptome analyses were all conducted as described in the repository *ms_clam_garden* (see *Data*

Availability). Transcript annotation, effective counts, and phenotypic data were all imported into R to be

analyzed with *edgeR* v.3.36.0 (Robinson et al., 2010). Separate datasets were constructed including all

samples, or including only gill or only digestive gland (DG) samples ($n = 3$ datasets). Each dataset was

input into a DGEList, then filtered for low expression by requiring at least 10 reads to be aligned to the

transcript in at least five individuals (*i.e.*, counts per million (cpm) > 0.82–1.03, depending on the dataset).

Samples were normalized by TMM normalization as applied by *calcNormFactors()* and multidimensional

scaling (MDS) plots were produced for each of the three datasets.

Tissue-specific expression was determined by identifying transcripts expressed in one tissue and not

in the other. The normalized counts for these transcripts were then obtained from the all-sample dataset

and 200 of the highest expressed tissue-specific transcripts from both tissues were plotted using *heatmap()*

of the *stats* package in R. Gene Ontology (GO) enrichment was conducted using the tissue-specific

transcripts from both tissues compared to all expressed transcripts for that tissue. Enrichment analysis was

conducted in DAVID Bioinformatics (Huang et al., 2007).

Differential expression analysis was conducted using *limma* v.3.50.1 (Ritchie et al., 2015). Both

tissues were analyzed separately and differentially expressed genes were identified using beach type (*i.e.*,

CG vs. Ref) as an explanatory variable using a gene-wise negative binomial generalized linear model with

quasi-likelihood tests (glmQLFit) and retaining all transcripts with $p \leq 0.001$. Differential expression,

based on beach-type with a Benjamini-hochberg multiple test correction, was also investigated.

Differentially expressed genes with $p \leq 0.001$ were also identified based on survivorship, where the

percent survival of each plot was used to bin the plot into one of high, medium, or low survival based on

the fourth quartile, second and third quartiles, and the first quartile of percent survival, respectively.

Statistical significance was conducted as above, but using contrasts between each survival group. All other

analyses and plots were conducted in R and are available in the repository listed above.

Results

3.1 | Clam growth and survival

There were no significant differences between unmaintained clam garden (CG) and reference (Ref)

beaches in any measured growth or survival variable (ANOVAs, $p \geq 0.168$, Table 1). The mean (± SD)

initial clam wet weight and shell height for CG and Ref beaches was 1.5 ± 0.2 g and 1.8 ± 0.3 g and 1.5 ±

0.1 cm and 1.6 ± 0.1 cm (N = 180), respectively (Table 1). After 16 weeks *in situ*, these increased to 3.9 ±

1.3 g and 3.8 ± 2.9 g and 1.8 ± 0.3 cm and 1.9 ± 0.4 cm, which in terms of height, was an increase in

growth of 20.5 ± 15.3% and 18.0 ± 22.6%, respectively. Calculating percent growth using shell height

includes both the alive and dead clams and is therefore considered to be a more accurate measurement of

growth compared to using wet weights. Wet weight only includes surviving clams, and therefore does not

capture the reduced growth that may have occurred in clams that eventually died. Clam survival was 63.3

$\pm$ 28.3% and 58.3 $\pm$ 37.0% in CG and Ref beaches, respectively.

Beach location regardless of beach type significantly affected all growth and survival variables

(ANOVAs, $p \leq 0.03$, Table 2, *SI* Figure S1). An unplanned significant difference occurred in the initial

size of clams among the six beaches (initial wet weight and shell height ANOVAs $p = 0.031$ and $p =$

0.044, respectively, Table 2, *SI* Figure S2). This difference was likely due to significantly larger clams

inadvertently being planted on beach D (Table 2, *SI* Figure S2). Percent growth was highest on beaches A

(31.2 $\pm$ 7.2%), C (29.7 $\pm$ 2.8%), and D (47.0 $\pm$ 8.2%) and lowest on beaches B (6.6 $\pm$ 6.8%), E (0.8 $\pm$

0.9%), and F (0.3 $\pm$ 0.6%) (Table 2). Percent survival was highest on beaches A (88.3 $\pm$ 12.6%), B (71.7 $\pm$

27.5%), C (68.3 $\pm$ 12.6%), and D (86.7 $\pm$ 5.8%) and lowest on beaches E (33.3 $\pm$ 23.6%) and F (16.7 $\pm$

24.7%) (Table 2).

Gross observations of the clam showed that most of the survivors on beaches A (CG), B (Ref), C

(CG), and D (Ref) were considered healthy (Table 3). These beaches had 68–88% survival, with generally

low percentages of clams considered weak (0–5%), emaciated (0–5%), or having pale or very pale

digestive glands (0–22%) (beach D was a slight exception, where although survival was high, 22% of

clams had pale digestive glands) (Table 3). In contrast, survivors from low-survival (17–33%) beaches E

(CG) and F (Ref) had more clams considered weak (13–15%), emaciated (15–22%), and with digestive

glands being very pale (13–22%). Beaches E and F are therefore considered poorer performing sites.

3.2 | Beach sediment characteristics

There were no significant differences between unmaintained CG and Ref beaches for any of the sediment

characteristics (ANOVAs, $p \geq 0.392$, Table 1). However, beach location regardless of beach type

significantly affected all sediment characteristics except percentage of small rocks (Table 2). A PCA of

sediment characteristics explained 66.0% (dimension 1: 40.8%, dimension 2: 25.6%) of the variation in

the data and separated beach locations into three groups (Figure 2), where pairs of beaches were observed

to be clustered based on similar geographic locations within Kanish Bay (Figure 1). Beach A (CG), on a

west-facing bay, was exposed to wave action from within and outside Kanish Bay. Beach A grouped with

beach B (Ref), an east-facing bay that was also exposed to wave action within Kanish Bay (Figure 1).

349 These beaches are the two sites that are closest to, but on opposite sides of, the entrance of Kanish Bay

350 and show correlation with very course sand, coarse sand, and sand (Figure 2). Beach C (CG) is a south-

351 facing sheltered beach within Small Inlet and groups with beach D (Ref), a north-facing beach on the

352 opposite side of Small Inlet. Beaches C and D correlated with rocks, fine sand, and final weight/shell

353 height (Figure 2). Beaches E (CG) and F (Ref), two proximate beaches on the south side of Kanish Bay,

354 face north and are exposed to moderate wave action. These sites were associated with carbonates,

355 organics, and small rocks, and showed an inverse correlation with percent survival (Figure 2). Notably,

356 these beach groupings in the PCA remain constant when only abiotic characteristics are included in the

357 PCA, or when only survival and growth variables are included (*data not shown*). The PCA was also run

358 with all data, but excluding beach D due to the aforementioned initial size difference of this beach, and

359 this showed similar associations and groupings to the above, but beach C was on its own (*SI* Figure S3).

360  Most sand types, silt, and rocks were either positively or negatively correlated with growth and

361 survival variables (*SI* Figure S4). To investigate correlations between sediment characteristics and clam

362 growth and survival further, beach D was again excluded as described above (*SI* Figure S5). Abiotic

363 variables with significant positive correlations with growth included sand ($p = 0.018$, $R^2 = 0.31$) and fine

364 sand ($p = 0.022$, $R^2 = 0.29$), with those having negative associations being silt ($p = 0.002$, $R^2 = 0.49$) and

365 small rocks ($p = 0.014$, $R^2 = 0.34$). Variables with significant positive correlations with survival were

366 coarse sand ($p = 0.014$, $R^2 = 0.34$) and sand ($p = 0.002$, $R^2 = 0.48$), with those having negative associations

367 being carbonates ($p = 0.008$, $R^2 = 0.39$) and organic content ($p = 0.035$, $R^2 = 0.24$), as well as small rocks

368 ($p = 0.021$, $R^2 = 0.30$), very fine sand ($p = 0.002$, $R^2 = 0.49$), and silt ($p = 0.002$, $R^2 = 0.51$) (*SD* Additional

369 File S1, without Beach D).

371 **3.3 | Transcriptomic overview and *de novo* reference transcriptome**

372 The gill and digestive gland (DG) libraries produced on average ($\pm$ SD) 53.8 ± 10.7 and 55.0 ± 8.3 M read

373 pairs per library, respectively. All 28 libraries, comprising 305 GB of high-quality sequence data, were

374 used to create a *de novo* transcriptome assembly (Table 4; see *Data Availability*). The initial Trinity

375 assembly resulted in 1,695,678 transcripts, of which 1,277,478 were likely coding sequences as predicted

376 by TransDecoder. Following filtering of the *de novo* assembly (see *Methods*), there were 52,000

377 representative transcripts in the assembly, each putatively representing a single gene, and 42,708 had

378 associated Uniprot identifiers.

379 Alignments of trimmed reads resulted in a mean ($\pm$ SD) alignment rate of $27.87 \pm 2.48\%$ against the

380 Pacific littleneck clam *de novo* transcriptome. For comparison, the alignment rate against the Manila clam

381 genome (*i.e.*, the closest relative with a reference genome at the date of the present analysis) resulted in

382 very low alignments of $1.10 \pm 0.17\%$ and $2.59 \pm 1.00\%$ for gill and DG, respectively. All downstream

383 analyses used the *de novo* transcriptome quantified transcripts. After filtering the all-sample dataset (N =

384 28 pooled samples), 33,825 transcripts (67.7%) were expressed in at least five samples above the applied

385 threshold (see *Methods*).

386 An MDS plot of samples from both tissue types showed that gill samples were distinctly separate

387 from DG samples across the first dimension, explaining 65% of the variation in the dataset (Figure 3A).

388 The second dimension, which explained only 3% of the variation, separated the DG samples, but not the

389 gill ones. One outlier sample, P12, was observed in the gill data.

391 **3.4 | Tissue-specific expression**

392 Samples from each tissue were then analyzed separately by conducting tissue-specific filtration and

393 normalization. The gill and DG samples had on average ($\pm$ SD) 12.9 ± 3.0 M and 15.5 ± 2.9 M reads

394 aligning per library and the datasets were found to have 28,391 and 24,699 transcripts expressed,

395 respectively, after filtering for low expression. Inspecting transcripts that were present in only one of the

396 tissue types identified 8,795 (31.0%) gill-specific transcripts and 5,103 (20.7%) DG-specific transcripts

397 (Table 5; *SD* Additional File S2). The highest expressed tissue-specific transcripts are shown in *SI* Figure

398 S6 (N = 200 for each tissue).

399 Tissue-specific transcripts were compared to all expressed genes in that tissue to identify enriched

400 GO categories in the tissue-specific list. Gill-specific transcripts were enriched for response to stimulus (N

401 = 1,839), signal transduction (N = 1,184), biological adhesion (N = 496), ion transport (N = 494), defense

402 response (N = 402), and G-protein coupled receptor signalling pathways (N = 210) (all $p < 0.0001$),

403 among others (*SD* Additional File S3). DG-specific transcripts were enriched for biological adhesion (N =

404 308), defense response (N = 263), immune response (N = 223), ion transport (N = 201), innate immune

405 response (N = 149), organonitrogen compound catabolic process (N = 130), defense response to bacteria

406 (N = 82), and humoral immune response (N = 58) (all $p < 0.0001$), among others (*SD* Additional File S3).

408 **3.5 | Effect of unmaintained clam gardens on clam gene expression**

The gill dataset showed only 13% and 11% of the total variation in an MDS analysis explained by

dimensions 1 and 2, respectively (Figure 3B). No clear clustering was observed in the gill samples by

beach type (*i.e.*, CG vs. Ref) nor by any other evaluated variables (*e.g.*, survival, sand, silt, or carbonates).

There were two outlier samples (P12 from beach B and P7 from beach D) in the MDS plot, both of which

were Ref beaches and had 45% and 80% survival, respectively.

The DG dataset showed only 14% and 9% of the total variation in an MDS analysis explained by

dimensions 1 and 2, respectively (Figure 3C). Some grouping was observed along dimension 2 with CG

samples clustering together, except for P15 from beach E. Ref samples were more spread out along both

axes. A gradient in survival was observed across dimension 1, with high survival being positioned in

negative dimension 1, medium survival near 0-1, and low survival in positive dimension 1, although the

trend was not universal for all samples (*e.g.*, P5). Percent carbonate, which was correlated with survival

(see above), also trended along dimension 1 in the opposite direction to survival (*i.e.*, high survival being

associated with low percent carbonate, *SI* Figure S7).

Differential expression between CG and Ref was evaluated for each tissue separately (Table 5). The

gill had 54 transcripts differentially expressed (FC > 1.5, $p \leq 0.001$, no multiple test correction (MTC))

(see Discussion; *SD* Additional File S2). Of these transcripts, 35 had identifiers recognized by DAVID

Bioinformatics, but no Gene Ontology (GO) enrichment was identified ($p > 0.01$). Most of these

transcripts were overexpressed in the CG samples (N = 40), including several transcripts associated with

immune/apoptotic or stress-related functions, such as two transcripts annotated as *tumour necrosis factor*

(TNF)-related (FC > 1.5), one as *death effector domain (DED)*-related (*i.e.*, top overexpressed transcript,

FC > 71.5), and *universal stress protein A-like protein* (FC > 4.7). In total, 14 transcripts were

underexpressed in the CG samples, including a *heat shock 70 kDa protein 12a* (FC = 17.9). A full list of

differentially expressed genes is available in *SD* Additional File S2.

The DG had 37 transcripts differentially expressed between CG and Ref samples. This included 11

overexpressed and 26 underexpressed transcripts in the CG samples (*SD* Additional File S2). Using the

full, bidirectional list of differentially expressed transcripts, several enriched GO categories were observed

(*SD* Additional File S3), including biological process categories iron ion homeostasis (N = 3, $p = 0.008$)

and negative regulation of hydrolase activity (N = 5, $p < 0.001$). The highest overexpressed transcript in

the CG samples was *replicase polyprotein* annotated from the *cricket paralysis virus* (FC > 500), which

suggests that viral proteins were also captured in the study and that they were differentially abundant

between beach types. Multiple complement-, iron-, and protein-folding-related transcripts were observed

as differentially expressed in the DG samples, including overexpression of a transcript annotated as *c1q*

domain (2.3-fold), *thrombospondin-1* (2-fold), *soma ferritin*, and *cytochrome P450* (FC > 1.5). Transcripts

under-expressed in the DG samples in the CG plots included *interferon-induced very large GTPase 1* (FC

> 8.5), *apoptosis inhibitor IAP* (4.3-fold), *low affinity immunoglobulin epsilon Fc receptor* (FC > 2), *c-*

type mannose receptor 2, and *platelet endothelial aggregation receptor 1* (FC > 1.5), among others (*SD*

Additional File S2).

Consistent responses to the CG plots were investigated between the tissues, and two of the 91 unique

differentially expressed transcripts were found to be differentially expressed in both tissues. This is

notable given that the datasets were normalized, filtered, and analyzed separately. This included the

underexpression in the CG plots of a transcript annotated as *von Willebrand factor type D domain (vwd)*

and of a transcript annotated as *platelet endothelial aggregation receptor 1* (*pear1*; Figure 4) in both

tissues.

3.6 | Gene expression responses and differential survival

Given the differential survival observed among the beaches and plots, genes correlating with survival were

investigated to determine what might be driving these differences and to potentially identify genes that are

related to survival or mortality in this non-model species. Genes were of particular interest if they were

differentially expressed between low and medium survival, between medium and high survival, or

incrementally differentially expressed at each step between high, medium, and low survival (see *Methods*

for beach survivorship classification).

The majority of genes significantly associated with survival in the gill were differentially expressed

between the low and medium survival groups, rather than between medium and high survival (Table 5).

Transcripts overexpressed in the low survival group relative to the medium survival group (N = 26) were

involved in functions such as heat-shock, interferon response, and ligase activity (*SD* Additional Files S2,

S3). For example, *heat-shock protein (hsp) 90* (FC > 60) and three different transcripts annotated as

subunits of *hsp70* (FC > 2) were overexpressed specifically in the low survival group. Transcripts

overexpressed in the medium survival group relative to the high survival group included six transcripts of

various functions (*SD* Additional File S2). Fewer transcripts were overexpressed in the higher survival

group (*i.e.*, high vs. medium, N = 11; medium vs. low, N = 14) and these generally were involved in

immune functions, but also various other functions. Overexpression of *toll-like receptor 1* (FC > 2) and

complement C1q-like protein 4 (FC > 3.5) was observed in medium survivors relative to low survivors

(Figure 5).

In the DG, the majority of significant survival-associated transcripts were overexpressed in low

relative to medium surviving clams (N = 34) and were generally involved in immune system, heat-shock,

or transport functions. Transcripts annotated as *hsp70 12a* and *12b*, as well as several aminopeptidases,

were overexpressed in low relative to medium and in medium relative to high survival (shown correlated

with survival in Figure 5). All the overexpressed genes in medium relative to high surviving clams (N =

18) were also overexpressed in the low relative to medium survival group comparison. Very few genes

were overexpressed in the high survival clams relative to medium (N = 4) or the medium relative to low

survival (N = 7, and these transcripts had various functions, including a transcript annotated as *mucin-2*

protein (Figure 5).

Discussion

4.1 | Effects of unmaintained clam gardens on clam and beach characteristics

Results from the present study showed that there was no significant enhancement of growth or survival on

unmaintained clam garden (CG) beaches relative to reference (Ref) clam beaches. Using clams of similar

sizes that were collected on the same days and location as the present work, Salter (Salter, 2018) also

reported no significantly enhanced growth or survival on unmaintained CG compared to reference clam

beaches in Kanish and Waiatt Bays. It is important to note that in the present study the clam gardens were

unmaintained and therefore were not being tended according to traditional tending practices, which may

influence the impact of the garden on clam survival, growth, and transcriptomic profile, as well as various

abiotic characteristics (*i.e.*, sediment grain size, carbonate content, organic content). Salter (2018),

however, did include some CG traditional tending practices (*i.e.*, adding a 1 cm-thick layer of crushed

shell mixture three times throughout the summer) on replicate treatment plots on both beach types and

reported a positive effect of the shell hash treatment on growth and survival in both CG and reference

clam beaches. Cox et al., reported that even without traditional tending clam gardens contained greater

estimated bivalve biomass, distinct infaunal and epifaunal communities and increased topographic

complexity compared to reference sites (Cox et al., 2024a; Cox et al., 2024b; Cox et al., 2019).

Interestingly, Cox et al., also found that a combination of 45-60% gravel and less than 31% bivalve shells

promotes the biologically diverse epifaunal communities observed within clam gardens (Cox et al.,
2024a). These results confirm the importance of historical Indigenous practices in maintaining clam
gardens and promoting an optimal environment for clam growth and survival, and demonstrate that it is
not simply the physical structure of the clam garden wall that creates the ideal clam habitat.

Some trends were observed in the transcriptomic data that suggest that the unmaintained CGs had a
transcriptomic effect on the clams, albeit not a strong one. First, the CG and Ref samples showed a slight
separation in the PCA of DG tissue, but not gill tissue. Second, although very few genes were
differentially expressed, the same differentially expressed genes were consistently identified in both
tissues, including transcripts annotated as *von Willebrand factor type D domain (vwd)* and *platelet*
endothelial aggregation receptor 1 (*pear1*) both of which were downregulated in CG. The *vwd* domain is
present in many proteins, including mucins and other extracellular glycoproteins, and one *vwd* transcript in
particular was up-regulated in response to heat shock in the Pacific oyster (*Crassostrea gigas*) (Zhang et
al., 2015). The function of *pear1* in shellfish remains unknown; in vertebrates it is a cell membrane protein
involved in platelet aggregation. Homologous genes for platelet activation are present in the Hong Kong
oyster (*Crassostrea hongkongensis*) and are regulated in response to hyposalinity (Xiao et al., 2018). In
terms of the other genes differentially expressed by clams in CGs, the general functions were not clear.
Immune genes and stress-response genes were found both over- and under-expressed in CGs versus Ref
beaches (*e.g.*, a heat-shock protein transcript was found to be underexpressed in the CG and a universal
stress protein-annotated transcript was overexpressed in the CG). In general, the functions of these
transcripts in the Pacific littleneck clam responding to the CG factor will be useful to continue to
characterize as the genomic information develops for this species and additional transcriptomic studies are
conducted in relation to environmental stress in maintained or unmaintained clam gardens. These
molecular phenotypes may indicate a response to the unmaintained CG from the 16-week *in situ* study
period that is below the level of phenotypic growth and survival responses, as these were not found to
differ based on beach type. Further work with longer study periods would be required to determine
whether these transcriptome signatures precede macro-phenotypic responses.

Sediment grain sizes, percent carbonate content, and percent organic matter did not differ
significantly between unmaintained CG and Ref beaches in the present study, although there was
significant variation within both beach types among locations. For example, carbonate content varied from

4 to 10% in CGs and from 2 to 15% in Ref beaches, while silt varied from 0 to 4% and from 1 to 5%,

respectively. This is somewhat contrary to Salter's (2018) observations that untreated CGs contained 2.8 –

12.9 times more carbonate and a smaller percentage of silt than non-walled (Ref) beaches (Salter, 2018).

The reason for this discrepancy is not known and would be valuable to address with future studies. The

lack of differences observed in the present study in growth and survival, and minor effects in

transcriptome responses due to unmaintained CGs may be due to the lack of tending of the clam gardens,

as is traditionally conducted. Future studies would benefit from fully examining the impacts of tending

practices on sediment characteristics alongside clam growth and survival, which would involve clam

garden beach restoration and active maintenance for long periods, with further analysis of impacts on

shellfish molecular phenotypes and productivity.

4.2 | Abiotic variation among beach location and effects on clam growth and survival

Across the six sampling locations, beaches differed in most of the variables examined (*i.e.*, biotic response

variables: growth, survival; abiotic characteristics: organic level, carbonate concentration, grain sizes).

Beaches clustered based on sediment characteristics, which correlated to beaches in similar geographic or

wave-influenced locations, and differential growth and survival were noted among the beach locations.

Given these trends, abiotic factors potentially involved in survival as well as molecular responses to

survival were characterized.

The sedimentary environment is very important for clam growth and survival (Joo et al., 2021).

Littleneck clams typically live as infaunal burrowers between 5 and 15 cm depth, in soft sand or sandy

mud in the intertidal zone (Lazo, 2004). In the present study, growth was positively affected by sand, and

fine sand, and negatively affected by small rocks, and silt, while survival was positively affected by coarse

sand, and sand, and negatively affected by carbonates, organics, small rocks, very fine sand, and silt Joo et

al. (2021) found that grain size and sorting of the sediment determined the amount of pore-water dissolved

oxygen and organic matter, both affecting growth and survival of juvenile Manila clams, with optimal

survival and growth linked to poorly-sorted sediment with an average grain size of medium sand (180 –

335 µm). Similarly, the grain size of the sand in the present study was 250 – 500 µm, and this grainsize

had a significant positive effect on survival. A clam tending revitalization project in Burrard Inlet

observed that water flow had a positive effect on native clam density and biomass while sediment grain

size had little effect (Guttmann, 2022). Salter (2018) found increased water residency time positively

affected clam biomass and growth and reported a weak negative effect of silt on clam density and

biomass. Contrary to the present study, Salter (2018) also found a positive effect of increased sediment silt

content on clam biomass in clam transplant experiments. A multi-year study looking at environmental

factors affecting growth in the hard clam *Venus mercenaria* (now *Mercenaria mercenaria*) found a

consistent decrease in growth with increasing sediment silt-clay content, with all results showing a

negative relationship between growth and fineness of the sediment (Pratt & Campbell, 1956). Permeability

of the sediment is an important factor for pumping activity of burrowing filter feeders, which is essential

for respiration, excretion, and nutrition and Pratt et al. (1956) found permeability to be an inverse function

of the sediment silt-clay content. In addition, commercial Manila clam beds studied in Spain had higher

survival of clams planted in sand-gravel than in those placed in mud (Cigarria & Fernandez, 2000).

Concurring, the clam *Gomphina veneriformis*, which was buried in sand and silt sediment mixtures, had

increased mortality with increasing silt (Li et al., 2021). Likewise, the juvenile bamboo clam (*Solen*

grandis) reared in different sediments had, as also seen in the present study, decreased growth and survival

rates with increasing fine silt (Chen et al., 2008).

Sediment carbonate content had a significant negative effect on clam survival in the present study,

where sediment with more than 8% carbonates had markedly decreased growth and survival. Similarly,

Munroe (2016) found that the average daily growth of early post-settled Manila clams over two years was

negatively correlated with sediment carbonate content (Munroe, 2016). This is not a universally observed

trend in the literature (Groesbeck et al., 2014; Salter, 2018) and counters expectations given the expected

importance of carbonate for clam shell growth and early recruitment. The addition of crushed shell hash

alters sediment calcium-carbonate saturation states, as it increases surface sediment porewater aragonite

content and pH (Green et al., 2009). For instance, there was a three-fold increase in hard clam burrowing

and recruitment by adding crushed shell hash to a final concentration of 8% of wet sediment weight

(Green et al., 2012). Salter (2018) increased carbonate content significantly (1.9–7.2 times) by addition of

shell hash, which increased clam growth and tissue biomass in both clam garden and control beaches after

six months (Salter, 2018). Guttman (2022) when examining clam tending practices in Burrard Inlet found

that sediment carbonate had a positive effect on native clam density and biomass although when compared

to water flow it was imprecise and relatively less important (Guttmann, 2022).

Crushed shell hash also changes the physical structure of the sediment as, like gravel, it can create

interstitial spaces in the sediment, thereby increasing pore-water dissolved oxygen, organic-matter content,

and substrate stability, as well as providing protection from predators for juvenile clams (Joo et al., 2021;

Sponaugle & Lawton, 1990; Thompson, 1995). Thus, there may be multiple benefits of adding shell

material for bivalve growth/survival that are not linked to carbonate concentration *per se*. The effect of

shell hash or crushed shell on bivalve growth and survival is likely dependent on a wide variety of factors

including amount of shell, size and compactness of shell particles, species generating the shell hash,

species/size of bivalve grown, stocking densities, tidal height, temperature, and types/densities of

predators. For example, the application of very finely crushed shell hash could have a negative effect by

decreasing interstitial spaces in the sediment and thus, as seen with silt, decreasing sediment permeability.

Recent studies investigating the efficacy of shell hash to mitigate acidification of intertidal sediments

found it unsuccessful at raising pore-water pH of acidic sediments (Beal et al., 2020; Doyle & Bendell,

2022). Doyle et al. (2022) found the addition of shell hash to be site dependent as it reduced variation in

pH at an intertidal site with pH 8.03, but did not mitigate the pH of mud flats with pH 7.59. More research

is required to comprehensively understand the effects of shell hash on bivalve growth and survival.

4.3 | Pacific littleneck clam transcriptomics

In non-model organisms, such as the Pacific littleneck clam, advances in molecular ecological and

physiological understanding of the species can be advanced through transcriptomic tools and the results

these tools generate. Here it was clear that the reference transcriptome generated *de novo* was necessary

for the study, given that the closest relative genome (*i.e.*, Manila clam) resulted in a very low alignment

rate with the Pacific littleneck clam reads. Further, the use of different tissues in the present study revealed

high levels of tissue-specific expression (*i.e.*, 21‑31% of the transcripts expressed in the tissue). As

multiple individuals were included in each sample (*i.e.*, pools), the true number of individuals sampled for

the tissue-specific expression analysis is greater than the number of libraries, increasing the reliability of

the tissue-specific transcripts. The expression analysis indicated that for both tissues, tissue-specific

transcripts were involved in signal transduction, defense and immune response, and ion transport, among

other processes. Therefore, although the specific transcripts were different for each tissue, similar

functions were enriched in the tissue-specific gene lists for both the gill and the DG, although exceptions

were observed.

The presence of differential survival among the locations provided a valuable opportunity for

analyzing transcripts that are associated with mortality/survival. These characterizations, over time

through multiple transcriptome studies generating more evidence, can begin to characterize specific

transcripts that are involved with specific stress types (Evans et al., 2011; Miller et al., 2017; Sutherland et

al., 2012). This can enable the development of Ecological Association Ontology data collection for non-

model species (Pavey et al., 2012) and in the future may be used to characterize underlying factors

associated with observed mortality or morbidity (*e.g.*, abiotic stress compared to bacterial or viral

infection). Although the Pacific littleneck clam remains largely uncharacterized at the molecular level,

some of the genes identified in the present study correlated with survival, including differential expression

in the gill and DG of heat-shock proteins, immune response genes, and other functional categories.

Additional transcriptomic studies on the species in different environmental conditions over various

temporal and spatial scales should help to improve these categorizations and identify stressor-specific or

generalized response genes.

Differential expression was observed in transcripts involved in immune defense. For example,

transcripts annotated as complement C1q protein complex or involved in cell adhesion and oxidation-

reduction processes had high expression in high surviving beaches and may indicate increased activity of

immune defense responses. Complement C1q proteins are an important component of the innate immune

system (Zhang et al., 2014) and are upregulated in response to bacterial challenges (McDowell et al.,

2014) and salinity stress (Zhao et al., 2012). Cell adhesions are critical for the development, maintenance,

and function of multicellular organisms and are essential for invertebrate immunity (Johansson, 1999).

In the present study—potentially due to the relatively low sample size from pooling individuals

and/or due to having fewer plots with a sufficient number of survivors at some beaches, and/or due to the

high number of expressed transcripts—the impacts of multiple test corrections were severe on the dataset

and, therefore, a stringent *p*-value ($p < 0.001$) was applied without a multiple test correction. Associations

of genes to CGs or to differential survival should be considered as preliminary observations and further

studies evaluating the functions of these transcripts will be beneficial to understand their role in Pacific

littleneck clam transcriptomics.

4.4 | Viral presence and clam health

The use of RNA-sequencing for the present analysis identified a differentially expressed virus transcript in

the CG versus Ref samples, which could have unexpected effects on either gene expression activity or

phenotypic responses, as could any other uncharacterized factor influencing clam growth/physiology (*e.g.*,

temperature, beach exposure). The top overexpressed transcript in the DG from CGs was a *replicase*
polyprotein from the *cricket paralysis virus* (CPV), which is a positive-sense, single-stranded RNA virus
from the family Dicistroviridae in the order Picornavirales. Members of the Dicistroviridae family are
widely distributed in nature with the broadest host range of all small RNA viruses in invertebrates
(Bonning, 2009). As the Dicistroviridae viral sequences isolated in the present study ranged in size from
1479 to 7371 nucleotides, one can deduce that they represent viral mRNA, instead of viral genomes. Some
dicistroviruses can be non-pathogenic or cause subtle impacts (reduced longevity and fecundity), while
others result in rapid paralysis (Bonning, 2009).

Even though there was transcriptomic evidence of viral presence in all the beaches, most of the
Pacific littleneck clams from the medium- and high-survival groups did not exhibit any signs of
pathophysiology. Other clams (mostly from the low-survival group), however, were emaciated or watery
and had slightly to extremely pale-coloured digestive glands, an indicator of reduced feeding and
potentially compromised function and health. Pale digestive glands could also be the result of paramyxean
parasites responsible for disease in marine mollusks, with infection being linked to digestive tropism
interfering with food adsorption and causing pale-yellowish digestive glands with thin, watery flesh in
both mussels and oysters (Alfjorden, 2017; Carella et al., 2011). In the present study, however, there was
no significant parasite-derived gene expression observed and histopathology would be needed to confirm
any paramyxean parasite infections. It is also worth noting that the single time point of sample collection
in the present study and the analysis of only surviving clams may have reduced the likelihood of viewing
viral or parasitic gene expression in these samples.

Although very pale digestive glands and emaciation were observed in clams from beaches with lower
growth and survival, histopathology or genetic identification would again be required to confirm any viral
gene expression linkages to observed impacts on organismal health. Health status is most likely linked to
multiple stressors comprised of abiotic (*e.g*., temperature, sediment composition, water residency) and
biotic factors (*e.g*., bacteria, viruses, harmful algae). RNA sequencing combined with *de novo*
transcriptome assembly revolutionizes virus detection and allows for the screening of a broad range of
symptomatic and asymptomatic virus species in a non-discriminatory manner, thereby displaying a
broader picture of the environmental factors present (Nagano et al., 2015). The current dataset has yet to
be fully characterized for viral sequences and could be mined further at a later date, particularly as more
resources are developed for this species.

Conclusions

The *de novo* transcriptome assembly of the Pacific littleneck clam enabled the investigation of genes
differentially expressed based on clam presence in unmaintained clam gardens relative to reference
beaches. No abiotic characteristic differences were observed between the two beach types, no phenotypic
responses were observed, and all observed transcriptome responses were minor. It is possible that impacts
would be greater if the clam gardens were actively tended. However, significant survival differences
among beaches independent of clam garden presence/absence, as well as abiotic characteristic differences,
indicated several sediment characteristics significantly correlated with growth and survival, including the
unexpected result of carbonate content negatively correlated with survival. The survival differences also
enabled the detection of gene expression transcripts putatively related to stress response, and along with
tissue-specific characterization of the gill and digestive gland, this study advances our molecular
understanding of the Pacific littleneck clam, a non-model bivalve species.

Acknowledgements

We acknowledge the K'ómoks (Comox) Nation, the Xwémalhkwu (Homalco) Nation, the We Wai Kai
(Cape Mudge) Nation, the We Wai Kum (Campbell River) Nation, the Snuneymuxw Nation,
the lək̓ʷəŋən (Songhees) Nation, the WSÁNEĆ (Saanich) Nation, and the Wyomilth (Esquimalt)
Nation, who have stewarded and protected the territory we worked within since time immemorial. We
thank the Hakai Institute, the Tula Foundation, and the Clam Garden Network for facilitating the research
and for their continued support. We also thank Caitlin Smith, Morgan Black, Kayla Suhan, Kayla Long,
and Kieran Cox for their invaluable help in the field, Dr. Eric Rondeau for his assistance in the laboratory,
and Allie Byrne for creating the map of the experimental sites.

Funding

705   Funding for this study was provided by the Hakai Institute. This research was also supported by the
706   Canada Research Chair Program, Canada Foundation for Innovation and British Columbia Knowledge
707   Development Fund (Sarah Dudas).

Competing Interests

710 BJGS is affiliated with Sutherland Bioinformatics. The author has no competing financial interests to

711 declare. The other authors have no relevant financial or non-financial interests to disclose.

Data Availability

Raw RNA-sequencing data have been uploaded to SRA under BioProject PRJNA818991, BioSamples

SAMN26893882–SAMN26893909.

Supplemental materials, including all abiotic and phenotypic site metadata needed for data analysis,

transcript annotation, quantified gene expression counts, sample phenotypes:

Reference transcriptome: https://doi.org/10.6084/m9.figshare.19735753

RNA-seq bioinformatics pipeline: https://github.com/bensutherland/Simple_reads_to_counts

RNA-seq analysis and physiological/abiotic analysis pipeline: https://github.com/raapm/ms_clam_garden

Additional Files

Additional File S1. Abiotic variables with survival and growth linear model p and R^2 values, including and

excluding beach D.

Additional File S2. Differentially expressed genes and background gene lists for both tissues in Pacific

littleneck clams. Gene lists include clam garden vs. reference, survival groups, and tissue-specific

expression.

Additional File S3. Gene Ontology enrichment results for the clam garden analysis, the survival analysis,

and the tissue-specific expression.

Supplemental Results provided, including Figures S1–S7.

References

Alfjorden, A., M. Areskog, D. Bruno, R. Carnegie, D. Cheslett, S. Feist, S. Ford, S. Jones, A. Lillehaug, L.
Madsen, T. Renault, N. Ruane and P. Vennerström. (2017). *New Trends in Important Diseases*
Affecting the Culture of Fish and Molluscs in the ICES Area 2002 – 2015 (In: Anderson, E.D., N.
Ruane, R. Carnegie (eds.) ICES Cooperative Research Report No. 337, Issue.

Andrews, S. (2010). *FastQC: A quality control tool for high throughput sequence data. Available online*
at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc. In

Beal, B. F., Coffin, C. R., Randall, S. F., Goodenow, C. A., Pepperman, K. E., & Ellis, B. W. (2020).
Interactive effects of shell hash and predator exclusion on 0-year class recruits of two infaunal
intertidal bivalve species in Maine, USA. *Journal of Experimental Marine Biology and Ecology*,
530-531. https://doi.org/10.1016/j.jembe.2020.151441

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics*, *30*(15), 2114-2120. https://doi.org/10.1093/bioinformatics/btu170

Bonning, B. C. (2009). The Dicistroviridae: An emerging family of invertebrate viruses. *Virologica*
Sinica, *24*(5), 415-427. https://doi.org/10.1007/s12250-009-3044-1

Bourlat, S. J., Borja, A., Gilbert, J., Taylor, M. I., Davies, N., Weisberg, S. B., Griffith, J. F., Lettieri, T.,
Field, D., Benzie, J., Glockner, F. O., Rodriguez-Ezpeleta, N., Faith, D. P., Bean, T. P., & Obst,
M. (2013). Genomics in marine monitoring: new opportunities for assessing marine health status.
Mar Pollut Bull, *74*(1), 19-31. https://doi.org/10.1016/j.marpolbul.2013.05.042

Caldwell, M. E., Lepofsky, D., Combes, G., Washington, M., Welch, J. R., & Harper, J. R. (2012). A
Bird's Eye View of Northern Coast Salish Intertidal Resource Management Features, Southern
British Columbia, Canada. *The Journal of Island and Coastal Archaeology*, *7*(2), 219-233.
https://doi.org/10.1080/15564894.2011.586089

Carella, F., Aceto, S., Maiolino, P., & De Vico, G. (2011). What is your diagnosis? Pale yellowish
digestive gland and watery tissues in Mediterranean mussels. *Vet Clin Pathol*, *40*(2), 273-274.
https://doi.org/10.1111/j.1939-165X.2011.00316.x

Chadwick, M., Harper, E. M., Lemasson, A., Spicer, J. I., & Peck, L. S. (2019). Quantifying susceptibility
of marine invertebrate biocomposites to dissolution in reduced pH. *R Soc Open Sci*, *6*(6), 190252.
https://doi.org/10.1098/rsos.190252

Chen, A., Zhang, Z., Yao, G.-i., Zhang, C., Liu, G., Wu, X., & Shen, H. (2008). The influence of different
environmental factors on the growth and survival rate of juvenile of bamboo clam *Solen grandis*.
Journal of Shanghai Fisheries University/Shanghai Shuichan Daxue Xuebao, *17*(5), 559-563.
https://www.proquest.com/scholarly-journals/influence-different-environmental-factors-
on/docview/21139392/se-2?accountid=27308

Cigarria, J., & Fernandez, J. M. (2000). Mangement of manila clam beds I. Influence of seed size, type of
substratum and protection on initial mortality. *Aquaculture*, *182*, 173-182.

Cox, K. D., Davies, H. L., Millard-Martin, B., Black, M., Hessing-Lewis, M., Smith, N. F., Juanes, F., &
Dudas, S. E. (2024a). Ancestral and contemporary intertidal mariculture practices support marine
biodiversity in the northeast Pacific. *Communications Earth & Environment*, *5*(1).
https://doi.org/10.1038/s43247-024-01515-3

Cox, K. D., Davies, H. L., Millard☐Martin, B., Black, M., Hessing☐Lewis, M., Smith, N. F., Juanes, F.,
& Dudas, S. E. (2024b). Ancient and contemporary shellfish cultivation practices bolster bivalve
communities and diversity☐biomass relationships. *People and Nature*, *7*(1), 81-98.
https://doi.org/10.1002/pan3.10724

Cox, K. D., Gerwing, T. G., Macdonald, T., Hessing-Lewis, M., Millard-Martin, B., Command, R. J.,
Juanes, F., Dudas, S. E., & Birchenough, S. (2019). Infaunal community responses to ancient clam
gardens. *ICES Journal of Marine Science*, *76*(7), 2362-2373.
https://doi.org/10.1093/icesjms/fsz153
Deur, D., Dick, A., Recalma-Clutesi, K., & Turner, N. J. (2015). Kwakwaka'wakw "Clam Gardens".
Human Ecology, *43*(2), 201-212. https://doi.org/10.1007/s10745-015-9743-3
Doyle, B., & Bendell, L. I. (2022). An evaluation of the efficacy of shell hash for the mitigation of
intertidal sediment acidification. *Ecosphere*, *13*(3). https://doi.org/10.1002/ecs2.4003
Erlandson, J. M., Rick, T. C., Braje, T. J., Steinberg, A., & Vellanoweth, R. L. (2008). Human impacts on
ancient shellfish: a 10,000 year record from San Miguel Island, California. *Journal of
Archaeological Science*, *35*(8), 2144-2152. https://doi.org/10.1016/j.jas.2008.01.014
Evans, T. G., Hammill, E., Kaukinen, K., Schulze, A. D., Patterson, D. A., English, K. K., Curtis, J. M., &
Miller, K. M. (2011). Transcriptomics of environmental acclimatization and survival in wild adult
Pacific sockeye salmon (*Oncorhynchus nerka*) during spawning migration. *Mol Ecol*, *20*(21),
4472-4489. https://doi.org/10.1111/j.1365-294X.2011.05276.x
Evans, W., Mathis, J. T., & Cross, J. N. (2014). Calcium carbonate corrosivity in an Alaskan inland sea.
Biogeosciences, *11*(2), 365-379. https://doi.org/10.5194/bg-11-365-2014
Ewels, P., Magnusson, M., Lundin, S., & Kaller, M. (2016). MultiQC: summarize analysis results for
multiple tools and samples in a single report. *Bioinformatics*, *32*(19), 3047-3048.
https://doi.org/10.1093/bioinformatics/btw354
Gosling, E. (2008). *Bivalve Molluscs: Biology, Ecology and Culture*. Wiley.
https://books.google.ca/books?id=mnQtr7GAjy0C
Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., Adiconis, X., Fan, L.,
Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke, A., Rhind, N., di
Palma, F., Birren, B. W., Nusbaum, C., Lindblad-Toh, K.,…Regev, A. (2011). Full-length
transcriptome assembly from RNA-Seq data without a reference genome. *Nat Biotechnol*, *29*(7),
644-652. https://doi.org/10.1038/nbt.1883
Green, M. A., Waldbusser, G. G., .L., R. S., Emerson, K., & O'Donnell, S. (2009). Death by dissolution
Sediment saturation state as a mortality factor for juvenile bivalves. *American Society of
Limnology and Oceanography*, *54*(4), 1037-1047.
Green, M. A., Waldbusser, G. G., Hubazc, L., Cathcart, E., & Hall, J. (2012). Carbonate Mineral
Saturation State as the Recruitment Cue for Settling Bivalves in Marine Muds. *Estuaries and
Coasts*, *36*(1), 18-27. https://doi.org/10.1007/s12237-012-9549-0
Greiner, C. M., Klinger, T., Ruesink, J. L., Barber, J. S., & Horwith, M. (2018). Habitat effects of
macrophytes and shell on carbonate chemistry and juvenile clam recruitment, survival, and
growth. *Journal of Experimental Marine Biology and Ecology*, *509*, 8-15.
https://doi.org/10.1016/j.jembe.2018.08.006
Groesbeck, A. S., Rowell, K., Lepofsky, D., & Salomon, A. K. (2014). Ancient clam gardens increased
shellfish production: adaptive strategies from the past can inform food security today. *PLoS One*,
9(3), e91235. https://doi.org/10.1371/journal.pone.0091235
Guttmann, M. (2022). *"I'd like to see it come back": The revitalization of Tsleil-Waututh Nation clam
tending in Burrard Inlet,* Simon Fraser University. Burnaby, B.C.
Haas, B. J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P. D., Bowden, J., Couger, M. B.,
Eccles, D., Li, B., Lieber, M., MacManes, M. D., Ott, M., Orvis, J., Pochet, N., Strozzi, F., Weeks,
N., Westerman, R., William, T., Dewey, C. N.,…Regev, A. (2013). De novo transcript sequence

reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nat*
Protoc, *8*(8), 1494-1512. https://doi.org/10.1038/nprot.2013.084
Heiri, O., Lotter, A. F., & Lemcke, G. (2001). Loss on ignition as a method for estimating organic and
carbonate content in sediments: reproducibility and comparability of results. *Journal of*
Paleolimnology, *25*, 101-110.
Huang, D. W., Sherman, B. T., Tan, Q., Kir, J., Liu, D., Bryant, D., Guo, Y., Stephens, R., Baseler, M. W.,
Lane, H. C., & Lempicki, R. A. (2007). DAVID Bioinformatics Resources: expanded annotation
database and novel algorithms to better extract biology from large gene lists. *Nucleic Acids Res*,
35(Web Server issue), W169-175. https://doi.org/10.1093/nar/gkm415
Jackley, J., Gardner, L., Djunaedi, A. F., & Salomon, A. K. (2016). Ancient clam gardens, traditional
management portfolios, and the resilience of coupled human-ocean systems. *Ecology and Society*,
21(4). https://doi.org/10.5751/es-08747-210420
Johansson, M. W. (1999). Cell adhesion molecules in invertebrate immunity. *Dev Comp Immunol*, *23*(4-
5), 303-315. https://www.ncbi.nlm.nih.gov/pubmed/10426424
Joo, S., Jo, K., Bae, H., Seo, H., & Kim, T. (2021). Optimal sediment grain size and sorting for survival
and growth of juvenile Manila clams, *Venerupis philippinarum*. *Aquaculture*, *543*.
https://doi.org/10.1016/j.aquaculture.2021.737010
Kim, D., Langmead, B., & Salzberg, S. L. (2015). HISAT: a fast spliced aligner with low memory
requirements. *Nat Methods*, *12*(4), 357-360. https://doi.org/10.1038/nmeth.3317
Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat Methods*, *9*(4),
357-359. https://doi.org/10.1038/nmeth.1923
Lazo, D. G. (2004). Bivalve Taphonomy: Testing the effect of life habits on the shell condition of the
Littleneck clam *Protothaca (Protothaca) staminea* (Mollusca: Bivalvia). *PALAIOS 19 (5)*(Friday
Harbor Laboratory Special Issue), 451-459.
Lepofsky, D., Smith, N. F., Cardinal, N., Harper, J., Morris, M., Gitla, Bouchard, R., Kennedy, D. I. D.,
Salomon, A. K., Puckett, M., & Rowell, K. (2017). Ancient Shellfish Mariculture on the
Northwest Coast of North America. *American Antiquity*, *80*(02), 236-259.
https://doi.org/10.7183/0002-7316.80.2.236
Lepofsky, D., Toniello, G., Earnshaw, J., Roberts, C., Wilson, L., Rowell, K., & Holmes, K. (2020).
Ancient Anthropogenic Clam Gardens of the Northwest Coast Expand Clam Habitat. *Ecosystems*.
https://doi.org/10.1007/s10021-020-00515-6
Li, X., Chen, L., Zhou, Z., Li, B., & Liu, X. (2021). The behavioral and antioxidant response of the
bivalve *Gomphina veneriformis* to sediment burial effect. *Acta Oceanologica Sinica*, *40*(6), 75-82.
https://doi.org/10.1007/s13131-020-1690-1
McDowell, I. C., Nikapitiya, C., Aguiar, D., Lane, C. E., Istrail, S., & Gomez-Chiarri, M. (2014).
Transcriptome of American oysters, *Crassostrea virginica*, in response to bacterial challenge:
insights into potential mechanisms of disease resistance. *PLoS One*, *9*(8), e105097.
https://doi.org/10.1371/journal.pone.0105097
Milan, M., Coppe, A., Reinhardt, R., Cancela, L., Leite, R. B., Saavedra, C. C., Chelazzi, G., Bortoluzzi,
S., & Bargelloni, L. (2011). Transcriptome sequencing and microarray development for the
Manila clam, *Ruditapes philippinarum:* genomic tools for environmental monitoring. *BMC*
Genomics, *12*, 234.
Miller, K. M., Gunther, O. P., Li, S., Kaukinen, K. H., & Ming, T. J. (2017). Molecular indices of viral
disease development in wild migrating salmon (dagger). *Conserv Physiol*, *5*(1), cox036.
https://doi.org/10.1093/conphys/cox036

Munroe, D. (2016). Habitat effects on early post-settlement growth of intertidal clams,*Venerupis philippinarum* (A. Adams & Reeve, 1850). *Journal of Molluscan Studies*, *82*(4), 507-514. https://doi.org/10.1093/mollus/eyw014

Nagano, A. J., Honjo, M. N., Mihara, M., Sato, M., & Kudoh, H. (2015). Detection of plant viruses in natural environments by using RNA-Seq. *Methods Mol Biol*, *1236*, 89-98. https://doi.org/10.1007/978-1-4939-1743-3_8

Neudorf, C. M., Smith, N., Lepofsky, D., Toniello, G., & Lian, O. B. (2017). Between a rock and a soft place: Using optical ages to date ancient clam gardens on the Pacific Northwest. *PLoS One*, *12*(2), e0171775. https://doi.org/10.1371/journal.pone.0171775

Pavey, S. A., Bernatchez, L., Aubin-Horth, N., & Landry, C. R. (2012). What is needed for next-generation ecological and evolutionary genomics? *Trends Ecol Evol*, *27*(12), 673-678. https://doi.org/10.1016/j.tree.2012.07.014

Pratt, D. M., & Campbell, D. A. (1956). Environmental Factors Affecting Growth in *Venus mercenaria*. *Limnology Oceanography* (January).

R.Core.Team. (2025). *R: A language and environment for statistical computing*. In https://www.r-project.org/

Ritchie, M. E., Phipson, B., Wu, D., Hu, Y., Law, C. W., Shi, W., & Smyth, G. K. (2015). limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res*, *43*(7), e47. https://doi.org/10.1093/nar/gkv007

Roberts, A., & Pachter, L. (2013). Streaming fragment assignment for real-time analysis of sequencing experiments. *Nat Methods*, *10*(1), 71-73. https://doi.org/10.1038/nmeth.2251

Robinson, M. D., McCarthy, D. J., & Smyth, G. K. (2010). edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*, *26*, 139-140.

Salter, N. (2018). *Ancient clam gardens magnify bivalve production by moderating ambient termperature and enhancing sediment carbonate* (Publication Number 709) Simon Fraser University]. Vancouver, B.C.

Smith, N. F., Lepofsky, D., Toniello, G., Holmes, K., Wilson, L., Neudorf, C. M., & Roberts, C. (2019). 3500 years of shellfish mariculture on the Northwest Coast of North America. *PLoS One*, *14*(2), e0211194. https://doi.org/10.1371/journal.pone.0211194

Sponaugle, S., & Lawton, P. (1990). Portunid crab predation on juvenile hard clams: effects of substrate type and prey density. *Marine Ecology progress series*, *67*, 43-53.

Sutherland, B. J., Jantzen, S. G., Yasuike, M., Sanderson, D. S., Koop, B. F., & Jones, S. R. (2012). Transcriptomics of coping strategies in free-swimming *Lepeophtheirus salmonis* (Copepoda) larvae responding to abiotic stress. *Mol Ecol*, *21*(24), 6000-6014. https://doi.org/10.1111/mec.12072

Sutherland, B. J. G., Finke, J. F., Saunders, R., Warne, S., Schulze, A. D., Strohm, J. H. T., Chan, A. M., Suttle, C. A., & Miller, K. M. (2022). Metatranscriptomics reveals a shift in microbial community composition and function during summer months in a coastal marine environment. *bioRxiv preprint doi: https://doi.org/10.1101/2022.01.21.477286; this version posted August 15, 2022.* https://doi.org/10.1101/2022.01.21.477286

Thompson, D. S. (1995). Substrate additive studies for the development of hardshell clam habitat in waters of Puget Sound in Washington State: An analysis of effects on recruitment, growth, and survival of the Manila clam, *Tapes philippinarum*, and on the species diversity and abundance of existing benthic organisms. *Estuarine Research Federation*, *18*(No. 1A), 91-107.

Twindle, H. (1918). *Loading logs at Kanish Bay on Quadra Island*.
https://gallery.crmuseum.ca/photograph/loading-logs-kanish-bay-quadra-island
Waldbusser, G. G., & Salisbury, J. E. (2014). Ocean acidification in the coastal zone from an organism's
perspective: multiple system parameters, frequency domains, and habitats. *Ann Rev Mar Sci*, *6*,
221-247. https://doi.org/10.1146/annurev-marine-121211-172238
Wei, T., & Simko, V. (2021). *R package 'corrplot': Visualization Matrix (Version 0.92). Available from*
http://github/taiyun/corrplot
Wentworth, C. K. (1922). A scale of grade and class terms for clastic sediments. *Journal of Geology*, 377 -
393.
Williams, J. (2006). *Clam Gardens: Aboriginal Mariculture on Canada's West Coast* (Vol. 15). New Star
Books Ltd.
Xiao, S., Wong, N. K., Li, J., Lin, Y., Zhang, Y., Ma, H., Mo, R., Zhang, Y., & Yu, Z. (2018). Analysis of
in situ Transcriptomes Reveals Divergent Adaptive Response to Hyper- and Hypo-Salinity in the
Hong Kong Oyster, *Crassostrea hongkongensis*. *Front Physiol*, *9*, 1491.
https://doi.org/10.3389/fphys.2018.01491
Yan, X., Nie, H., Huo, Z., Ding, J., Li, Z., Yan, L., Jiang, L., Mu, Z., Wang, H., Meng, X., Chen, P., Zhou,
M., Rbbani, M. G., Liu, G., & Li, D. (2019). Clam Genome Sequence Clarifies the Molecular
Basis of Its Benthic Adaptation and Extraordinary Shell Color Diversity. *iScience*, *19*, 1225-1237.
https://doi.org/10.1016/j.isci.2019.08.049
Zhang, L., Li, L., Zhu, Y., Zhang, G., & Guo, X. (2014). Transcriptome analysis reveals a rich gene set
related to innate immunity in the Eastern oyster (*Crassostrea virginica*). *Mar Biotechnol (NY)*,
16(1), 17-33. https://doi.org/10.1007/s10126-013-9526-z
Zhang, Y., Sun, J., Mu, H., Li, J., Zhang, Y., Xu, F., Xiang, Z., Qian, P. Y., Qiu, J. W., & Yu, Z. (2015).
Proteomic basis of stress responses in the gills of the Pacific oyster *Crassostrea gigas*. *J Proteome*
Res, *14*(1), 304-317. https://doi.org/10.1021/pr500940s
Zhao, X., Yu, H., Kong, L., & Li, Q. (2012). Transcriptomic responses to salinity stress in the Pacific
oyster *Crassostrea gigas*. *PLoS One*, *7*(9), e46244. https://doi.org/10.1371/journal.pone.0046244
937
938
939

940   **FIGURES**
941

942
943   **FIGURE 1.** Locations of unmaintained clam gardens (A, C, and E; triangles) and unmodified reference
944   clam beaches (B, D, and F; circles) used in the study. The collection site used as source for the clams used
945   in the study is designated by a diamond. Inset map shows the location of the study site in western Canada.

946   A

947
948
949   **FIGURE 2.** Principal component analysis biplot based on abiotic and biotic phenotypes showing beach
950   groupings and correlations. Each beach is denoted as A through F as per Figure 1 locations, and each
951   replicate plot within the beach as 1–3. Colours of plots and ellipses are for visualization purposes only.

FIGURE 3. (A) MDS plot for all samples showing clustering based on tissue type (gill or digestive gland; dig) for Pacific littleneck clams sampled from plots P1–P18 on Beaches A–F**.** MDS plots for (B) gill and (C) digestive gland tissues individually where each datapoint is plotted as an unmaintained clam garden (CG, circles) or reference beach (Ref, triangles), coloured by beach identifier, and sized by the percentage of survival at the beach. The gill sample plot below -3 in dimension 2 is from plot P12, and the digestive gland sample that is below -3 in dimension 2 is from plot P9.

961
962
963 **FIGURE 4.** Box plots showing expression of genes of interest identified in both digestive glands and gills
964 of Pacific littleneck clams, which were underexpressed in clam gardens (CG) relative to reference beaches
965 (Ref). vwd = *von Willebrand factor type D domain*; pear1 = *platelet endothelial aggregation receptor 1.*

967
968
FIGURE 5. Linear regression of differential expression of selected genes of interest versus percent
survival in Pacific littleneck clams. Transcript expression and their correlation to survival include (top left
to bottom right) *heat shock protein family A member 12A* (*hsp70-12a*), *aminopeptidase*, and *mucin-2* in
the digestive gland (DG) and *hsp70-12a*, *hsp90*, and *toll-like receptor 1* in the gill. Note: adjusted R^2 and
p-values presented here are from linear models of log2 counts per million for each transcript against
percent survival as a numeric variable.

v

TABLES

TABLE 1. Mean (± SD) clam size or survival and sediment characteristic variables for clam garden (CG) and reference (Ref) beaches and associated linear model *p*-values (with beach location as a nested effect). No variables were significantly affected by the CG/Ref factor. All variables were found to have a normal distribution.

	Variable	CG Mean ± SD	Ref Mean ± SD	*p*-value
Clam size or survival	Initial weight (g)	1.5 ± 0.2	1.8 ± 0.3	0.200
	Initial shell height (cm)	1.5 ± 0.1	1.6 ± 0.1	0.168
	Final weight (g)	3.9 ± 1.3	3.8 ± 2.9	0.943
	Final shell height (cm)	1.8 ± 0.3	1.9 ± 0.4	0.864
	Growth in height (%)	20.5 ± 15.3	18.0 ± 22.6	*0.892[a]*
	Survival (%)	63.3 ± 28.3	58.3 ± 37.0	0.860
Sediment characteristics	Carbonates (%)	7.6 ± 1.8	7.2 ± 5.2	0.927
	Organics (%)	1.3 ± 0.6	1.1 ± 0.5	0.723
	Rocks (%)	15.6 ± 7.5	15.0 ± 5.4	0.912
	Small rocks (%)	12.1 ± 5.0	13.0 ± 3.3	0.765
	Very coarse sand (%)	12.2 ± 3.4	13.0 ± 3.6	0.794
	Coarse sand (%)	14.2 ± 3.4	13.2 ± 4.5	*0.788[a]*
	Sand (%)	20.0 ± 5.7	17.6 ± 4.2	0.575
	Fine sand (%)	18.6 ± 6.4	17.0 ± 2.9	0.715
	Very fine sand (%)	5.6 ± 2.1	8.3 ± 4.4	*0.393[a]*
	Silt (%)	2.0 ± 1.2	3.1 ± 1.5	*0.392[a]*

[a] Significant Bartlett's test indicates heteroscedasticity.

985 **TABLE 2.** Mean (± SD) clam size/survival and sediment characteristic variables for each beach and associated linear model *p*-values.

	Variable	Beach A Mean ± SD	Beach B Mean ± SD	Beach C Mean ± SD	Beach D Mean ± SD	Beach E Mean ± SD	Beach F Mean ± SD	*p*-value
Clam size/survival	Initial weight (g)	1.6 ± 0.1	1.8 ± 0.2	1.5 ± 0.3	2.0 ± 0.1	1.5 ± 0.2	1.5 ± 0.2	0.03
	Initial height (cm)	1.5 ± 0.0	1.6 ± 0.1	1.5 ± 0.1	1.7 ± 0.1	1.5 ± 0.1	1.5 ± 0.1	0.044
	Final weight (g)	4.2 ± 0.5	2.5 ± 0.7	5.1 ± 0.6	7.5 ± 0.8	2.5 ± 1.1	1.4 ± 1.3	< 0.0001
	Final height (cm)	2.0 ± 0.1	1.7 ± 0.2	2.0 ± 0.2	2.4 ± 0.2	1.5 ± 0.1	1.5 ± 0.1	< 0.00001
	Growth in height (%)	31.2 ± 7.2	6.6 ± 6.8	29.7 ± 2.8	47.0 ± 8.2	0.8 ± 0.9	0.3 ± 0.6	< 0.000001[a]
	Survival (%)	88.3 ± 12.6	71.7 ± 27.5	68.3 ± 12.6	86.7 ± 5.8	33.3 ± 23.6	16.7 ± 24.7	0.003
Sediment characteristics	Carbonates	8.0 ± 1.0	5.3 ± 2.5	6.0 ± 2.0	2.7 ± 0.6	8.7 ± 1.5	13.7 ± 1.2	< 0.0001
	Organics	0.7 ± 0.1	0.9 ± 0.3	1.4 ± 0.3	0.8 ± 0.2	1.8 ± 0.6	1.7 ± 0.3	0.005
	Rocks	8.0 ± 5.6	11.7 ± 6.1	19.0 ± 4.6	20.7 ± 1.5	19.7 ± 6.8	12.7 ± 2.1	0.037
	Small rocks	11.3 ± 5.1	11.7 ± 4.6	8.3 ± 2.3	11.7 ± 1.2	16.7 ± 3.8	15.7 ± 2.1	0.11
	Very coarse sand	13.7 ± 2.1	17.0 ± 2.0	6.7 ± 2.5	9.7 ± 2.5	14.3 ± 2.5	12.3 ± 1.2	0.005
	Coarse sand	18.0 ± 2.0	19.0 ± 1.7	11.7 ± 3.1	10.7 ± 1.5	13.0 ± 0.0	10.0 ± 0.0	< 0.005[a]
	Sand	25.7 ± 5.8	22.3 ± 3.5	19.7 ± 2.1	16.0 ± 1.0	14.7 ± 1.5	14.3 ± 1.5	0.003
	Fine sand	19.3 ± 4.7	14.7 ± 3.1	24.7 ± 3.2	17.7 ± 3.1	11.7 ± 0.6	18.7 ± 1.5	0.004
	Very fine sand	3.3 ± 0.6	3.0 ± 0.0	7.0 ± 1.7	10.7 ± 3.5	6.3 ± 1.5	11.3 ± 1.5	< 0.0005
	Silt	0.7 ± 0.6	1.7 ± 0.6	2.0 ± 0.0	3.0 ± 1.0	3.3 ± 0.6	4.7 ± 0.6	< 0.0001[a]

986 [a] Significant   Bartlett's   test   indicates   heteroscedasticity.

987 **TABLE 3.** Gross observations (health, condition, and digestive gland coloration) of surviving clams in
988 each beach (A–F).

TMetric	Beach A	Beach B	Beach C	Beach D	Beach E	Beach F
Health (% weak)	0	5	0	2	13	15
Condition (% emaciated)	0	5	0	2	22	15
Digestive gland (% pale)	8	7	5	22	0	0
Digestive gland (% very pale)	0	0	0	2	22	13

989
990
991 **TABLE 4.** Overview of RNA-seq reads and quality scores. Totals that include both reads in the pair are
992 indicated by (R1 + R2). Values are shown in billion (B) or million (M).
993

Leukoma staminea assembly	Value
Total input reads (R1 + R2)	3.0 B
Total input nucleotides (R1 + R2)	305 B
Average (± SD) input reads per:	
single HiSeq4000 lane (R1 + R2)	762 ± 13.0 M
library (read pairs)	54.4 ± 9.5 M
gill library (read pairs)	53.8 ± 10.7 M
digestive-gland library (read pairs)	55.0 ± 8.3 M
Assembly contig length range	102 – 29,916
Mean (± SD) Phred quality score	38.9 ± 0.3
Base-calling accuracy	> 99.9%

995
996 **TABLE 5.** Differentially expressed gene overview comparing beach type (clam garden (CG) vs.
997 reference beach (Ref)), beach survival (high vs. medium vs. low), tissue type (gill vs. digestive gland), or
998 expressed in background. Note that here differential expression is considered when $p \leq 0.001$ and log2
999 fold-change > 0.58 (1.5-fold overexpressed) or < -0.58 (1.5-fold underexpressed).
1000

Comparison	Result	# Transcripts in gills	# Transcripts in digestive gland
Tissue type (gill vs. digestive gland)	Gill vs. digestive gland	8,795	5,103
Beach type (CG vs. Ref)	Overexpressed CG	40	11
	Underexpressed CG	14	26
Beach survival (high vs. medium vs. low)	Overexpressed high vs. medium	11	4
	Overexpressed medium vs. low	14	7
	Overexpressed low vs. medium	26	34
	Overexpressed medium vs. high	6	18
Background	N/A	28,391	24,699

1001