Effects of ancient anthropogenic clam gardens on the growth, survival, and

transcriptome of Pacific littleneck clams (*Leukoma staminea*)

Monique R. Raap[1,2,a], Helen J. Gurney-Smith[1,2,3,b], Sarah E. Dudas[1,2,3,a], Christopher M. Pearce[4,5],

Jong S. Leong[1], Ben J.G. Sutherland[1,6], Ben F. Koop[1]

[1]Department of Biology, University of Victoria, Victoria, British Columbia, Canada, V8W 2Y2

[2]Hakai Institute, Quadra Island, British Columbia, Canada, V0P 1H0

[3]Department of Biology and Department of Fisheries and Aquaculture, Vancouver Island

University, Nanaimo, British Columbia, Canada, V9R 5S5

[4]Pacific Biological Station, Fisheries and Oceans Canada, Nanaimo, British Columbia, Canada,

V9T 6N7

[5]Department of Geography, University of Victoria, Victoria, British Columbia, Canada, V8W

2Y2

[6]Sutherland Bioinformatics, Lantzville, British Columbia, Canada, V0R 2H0

[a]Present address: Pacific Biological Station, Fisheries and Oceans Canada, Nanaimo, British

Columbia, Canada, V9T 6N7

[b]Present address: St. Andrews Biological Station, Fisheries and Oceans Canada, St. Andrews,

New Brunswick, Canada, E5B 0E4

Author for correspondence: HJG-S

St. Andrews Biological Station, Fisheries and Oceans Canada, St. Andrews, New Brunswick,

Canada, E5B 0E4

Email: Helen.Gurney-Smith@dfo-mpo.gc.ca

Abstract

Clam gardens traditionally established and maintained by coastal Indigenous Peoples of northwest

North America are habitat modifications that enhance intertidal clam productivity and therefore provide

secure and reliable local food resources. In this study, transcriptomic and phenotypic responses of Pacific

littleneck clams (*Leukoma staminea*) were investigated in relation to transplantation to either clam gardens

or unmodified clam beaches and growth for 16 weeks. Sediment characteristics (e.g., grain-size,

carbonate, and organic content) were also evaluated and considered in the response. Large differences in

phenotypic and abiotic characteristics were observed among beaches but did not differ based on

unmaintained clam garden presence. A *de novo* transcriptome for *L. staminea* containing 52,000 putative

transcripts was assembled and used to identify differential expression in response to the clam gardens.

This identified a relatively small effect, but found two transcripts that were differentially expressed in both

the gill and digestive gland tissues. In addition, differential expression along survival gradients, as well a

tissue-specific expression analysis provide insight into the characteristics of the transcriptome and its

ecological associations of this non-model organism. Across the beaches, abiotic characteristics with

negative effects on growth and/or survival included small rocks, very fine sand, silt, carbonate, and

organic content, whereas positive effects were observed from coarse sand, sand and fine sand. In

conclusion, here it was found that localized environmental factors are likely to have a greater influence on

Pacific littleneck clam physiology, growth, and survival than the presence or absence of unmaintained

clam gardens.

KEYWORDS: clam gardens; *Leukoma staminea*; Pacific littleneck clam; sediment carbonate content;

transcriptomics

1 | INTRODUCTION

Coastal shorelines have been altered to increase productivity and facilitate the harvesting of natural

resources for centuries (Erlandson et al. 2008). Ancient clam gardens and fishing weirs were shoreline

constructions built to increase clam habitat and trap fish, respectively, providing productive and

predictable local food resources for coastal Indigenous Peoples (Neudorf et al. 2017). These coastal

shoreline modifications provide important insights into cultural food practices, traditional technologies,

economies, values, and ancestral practices of coastal communities (Deur et al. 2015, Jackley et al. 2016,

Lepofsky et al. 2020, Smith et al. 2019).

Construction of clam gardens in North America is estimated to have begun approximately 3,500

years ago (Smith et al. 2019). These gardens were created by rolling rocks and boulders to the low-tide

line and building a wall parallel to the shoreline (Deur et al. 2015), which subsequently allowed for the

deposition of sediment between the wall and the high-tide mark. This created a wider intertidal shellfish

habitat area with a reduced slope (Neudorf et al. 2017) at a tidal height of 1–2 m above chart datum, where

shellfish such as Pacific littleneck clams (*Leukoma staminea*), a traditional food item for coastal

Indigenous Peoples, typically occur (Deur et al. 2015, Groesbeck et al. 2014). This reduced beach slope

also allowed a thin layer of seawater to be retained on the accumulated sediment, reducing desiccation

risk, facilitating clams to remain in shallow accessible portions of the intertidal zone, and maximizing

submersion time for feeding and therefore growth (Deur et al. 2015). The exterior (ocean) side of the clam

garden wall creates rocky reef habitat for other marine invertebrates including octopuses, sea cucumbers,

whelks, chitons, red turban snails, Dungeness crabs, and red rock crabs, many of which are also consumed

by humans (Caldwell et al. 2012, Deur et al. 2015, Lepofsky et al. 2017).

Clam gardens are found in the intertidal zone from Alaska (USA), throughout British Columbia (BC,

Canada), and into Washington State (USA) (Groesbeck et al. 2014). Northern Quadra Island, off the east

coast of Vancouver Island, has perhaps the highest density of clam gardens on the northwest coast of

North America (Neudorf et al. 2017). On Quadra Island there are 45 clam gardens on the northwest side of

the island in Kanish Bay and 49 on the northeast side in Waiatt Bay (Groesbeck et al. 2014). In these two

bays, clam gardens cover approximately 35% of the shoreline, increasing clam habitat by 26–35%

(Lepofsky et al. 2020).

Traditionally, clam gardens were maintained and tended with practices that included predator

exclusion, selective harvesting, and the addition of gravel and crushed shell (*i.e.*, shell hash) (Deur et al.

2015). Shell hash may provide larval shellfish settlement cues and protect the young recruits from water

acidity and predators (Green et al. 2012, Sponaugle and Lawton 1990). Clam gardens were tended into the

20[th] century, but most of them, including all the beaches in the present study, were not maintained at the

time of the present study (Williams 2006).

Research on untended clam gardens in Kanish Bay and Waiatt Bay found significantly higher

densities of Pacific littleneck clams and butter clams (*Saxidomus giganteus*), increased survival of clam

recruits, and higher Pacific littleneck clam growth rates compared to unmodified beaches (Groesbeck et al.

2014). Studies on butter clams on the central coast of BC found that densities were 2.44 times greater on

average in clam gardens compared to densities on unmodified beaches (Jackley et al. 2016). The higher

density was thought to be due to the creation of a wider intertidal area in the clam gardens.

Community composition may also be influenced by clam gardens (Jackley et al. 2016). Greater

abundances of many taxa occur on clam garden beaches relative to unmodified beaches, including

Nematoda, Harpacticoida, Chironomidae, and Bivalvia (*e.g.*, butter clams and Pacific littleneck clams).

The abundance of these taxa has been shown to be positively associated with the amount of surface shell

and gravel present (Cox et al. 2019). A study comparing grain sizes and carbonate content in beach

sediment on Quadra Island found 2.8 – 12.9 times more carbonate, and on average a smaller percentage of

silt, in sediment from clam gardens compared to that from unmodified beaches (Salter 2018). Similarly,

greater amounts of gravel and shell hash have been observed in clam gardens relative to unmodified

beaches (Groesbeck et al. 2014). Unmodified beaches nearby clam gardens are, by contrast, primarily

composed of silt, sand, and mud (Jackley et al. 2016). Such differences in sediment composition are

attributed to the tending practices of Indigenous Peoples, which included the addition of gravel and

crushed whole clam shells to clam gardens (Groesbeck et al. 2014, Williams 2006).

The effect of sediment shell hash on clam productivity is not definitive. Sediment carbonate

saturation state is a significant seawater variable that can affect the settlement and recruitment of larval

and juvenile infaunal bivalves (Green et al. 2012). Seawater calcium carbonate ($CaCO_3$) saturation state is

important in the formation and maintenance of shell thickness and integrity (Chadwick et al. 2019). Ocean

acidification results in a decrease in seawater pH and an alteration in the bioavailability of $CaCO_3$. A

decrease in $CaCO_3$ saturation state can negatively affect the formation and maintenance of shells of marine

molluscs and other calcifiers (Evans et al. 2014, Green et al. 2012, Waldbusser and Salisbury 2014). Some

studies have shown that beaches with increased levels of sediment shell hash have increased clam

settlement and productivity (Green et al. 2012, Groesbeck et al. 2014, Salter 2018), while others have

reported negative (Munroe 2016) or neutral (Greiner et al. 2018) correlations with early post-settlement

bivalve growth and recruitment. A study on Quadra Island found that the addition of crushed shell had a

positive effect on transplanted littleneck clam growth and final tissue biomass in both unmaintained clam

gardens and unmodified beaches (Salter 2018). Most of these studies looked at variables such as growth

and survival over a relatively short period of time (*i.e.*, 30 days to six months, the exception being Munroe

(2016) where the trial ran for two years), whereas the effects of these variables may take a longer time to

manifest as physiological changes in clams.

Genomic tools such as transcriptomics offer the ability to capture early signals of environmental

stressors and are increasingly used to evaluate their biological effects on aquatic life (Milan et al. 2011,

Sutherland et al. 2012). Gene-expression data can then be linked with survival, growth, and environmental

information and used to predict possible physiological effects (Evans et al. 2011, Miller et al. 2017).

Transcriptomics can not only yield faster results but may also be used for taxonomic identifications

(Bourlat et al. 2013, Sutherland et al. 2022). The present study aims to determine whether Pacific

littleneck clam health and productivity, as assessed by growth and survival as well as transcriptomic

profiling, are affected by the presence of unmaintained clam gardens and various sediment characteristics

(e.g., grain-size distribution, organics, and carbonates). To this end, we assembled a *de novo* reference

transcriptome and characterized Pacific littleneck clam transcriptome responses in replicated unmaintained

clam garden beaches and unmodified reference beaches in the same area. Measured environmental

parameters and clam phenotypes (e.g., growth, survival) were considered while characterizing the

transcriptomic responses. The combination of abiotic measurements and organismal and molecular

phenotypes with the characterized transcriptome provides valuable context for both the effects of different

environmental variables on Pacific littleneck clams and on the functions of response genes in this non-

model ecologically and culturally important species.

2 | MATERIALS AND METHODS

A positionality statement is provided here for transparency with respect to any real or perceived research or interpretation biases of the authors due to their backgrounds. The authors of this paper are experienced academic, government, and independent scientific researchers on climate change, ecology, invertebrate physiology, bioinformatics and genomics. The authors were educated and trained in North American and European institutions prior to the positions in Canada that they hold at time of submission, with five reaching doctorate status and two with postgraduate masters degrees. This research was conducted in British Columbia, Canada. None of the authors who have provided heritage information for this statement indicated First Nations or other Indigenous background, and therefore we respectfully acknowledge the potential for unconscious bias on the presented subject matter.

2.1 | Study sites

The intertidal coastlines of Kanish Bay and adjoining Small Inlet on Quadra Island were chosen for this study due to an abundance of historic clam gardens. Quadra Island is located off the northeast coast of Vancouver Island, in southwest BC, and is in the traditional territories of the Kwakwaka'wakw (Laich-kwil-tach) and northern Coast Salish Peoples (Smith et al. 2019). Three unmaintained clam garden beaches (CG, designated as A, C, and E) and three unmodified reference sites (Ref, designated as B, D, and F) were selected in Kanish Bay and Small Inlet (Figure 1). Beach B was later discovered to be a damaged clam garden with ~ 75% of the wall destroyed (possibly from logs being dragged down the beach). In this study beach B is still considered a reference beach as it does not appear to be an outlier from the other reference beaches in any of the sediment characteristics (Table 2). These clam gardens have

likely not been tended in accordance with Indigenous practices since the early-to-mid 1900s (Deur et al.

2015).

2.2 | Clam transplantation experiment

Juvenile Pacific littleneck clams (N = 400), between 1 and 2 cm in shell height (hinge to shell margin),

were hand collected from a site in Kanish Bay (Figure 1) between 1 and 2 m Mean Lower Low Water

level on May 7, 2016, with all field work occurring that year. Each was individually wet weighed,

measured for shell height and length (i.e., the widest part of the shell at 90° to the shell height), and then

haphazardly placed into 18 groups of 20 clams each. Clams were held in a pearl net that was deployed

below the low intertidal zone in Heriot Bay (Quadra Island) until deployment. Clams on beach A were

deployed on May 10; those on beaches B, C, and D on May 11; and those on beaches E and F on May 12.

The different deployment days were required due to tidal heights and weather conditions. At low tide at

each of the six beaches, the 1.5–1.8 m intertidal zone was demarcated and three plastic mesh cubes (length

x diameter x height: 20 x 20 x 4 cm; made from high-density polyethylene, Vexar™), each containing 20

clams, were buried in the top 20 cm of the sediment, 5 m apart and parallel to the shore in the middle of

the demarcated zone. Each mesh cube was defined as one replicate plot (P). Each plot was numbered so

that Beach A contained P1–P3; Beach B: P10-P12; Beach C: P4-P6; Beach D: P7-P9; Beach E: P13–P15;

Beach F: P16–P18.

Transplanted clams were left *in situ* for 16 weeks and collected on August 30–31, 2016. Upon

retrieval, clams were quickly transported in bags in coolers to the Hakai Institute laboratory on Quadra

Island. The wet weight, shell height, shell length, and dead/alive status were recorded for each individual.

Growth and survival per plot were calculated as:

Percent growth = (mean final height – mean initial height) / mean initial height x 100

Percent survival = number of survivors / initial number of clams x 100

In addition, some visual observations were made on animal condition, following the protocols developed

for the Canadian National Animal Aquatic Health Program (NAAHP), including observations of animal

state (*i.e.*, health/vitality, condition) and digestive gland colouration.

2.3 | Sediment collection and analysis

Sediment-core samples (length x diameter: 15 x 5 cm) were collected adjacent to each cube ($n = 3$ cores

per beach) to assess grain size of core. Additional sediment-core samples (10 x 3.5 cm) were also collected

adjacent to each cube to assess percent carbonate and percent organic matter content. Each core was

placed individually in labeled sample bags and frozen at -20°C until analysis.

To determine grain-size distribution, the sediment samples were dried at 100°C for 24 hr, weighed,

transferred to a series of nested sieves (mesh size: 4.75 mm, 2 mm, 1 mm, 500 µm, 250 µm, 125 µm, and

63 µm), and shaken in a sediment shaker for 15 min. Each size fraction was then weighed and expressed

as a percentage of the total sediment weight. A scale very similar to the Wentworth scale of sediment size

classes was used to classify the grain sizes: pebble and larger gravel called "rocks" in the present study (>

4.75 mm), granule gravel called "small rocks" in the present study (2.00 – 4.75 mm), very coarse sand

(1.00 – 2.00 mm), coarse sand (0.50 – 1.00 mm), sand (250 – 500 µm), fine sand (125 – 250 µm), very

fine sand (63 – 125 µm), and silt (< 63 µm) (Wentworth 1922).

The loss on ignition method (Heiri et al. 2001) was used to determine sediment percent organic

matter and carbonate content. Briefly, the sediment samples were dried at 100°C for 48 hr in previously-

ashed (450°C for 8 hr) crucibles, and the dry weights measured. Dried sediment samples were then ashed

in a muffle furnace at 435°C for 8 hr. After cooling the samples in a desiccator for 1 hr, the weights of the

ashed samples were recorded. The percent organic matter was then calculated using the following

equation:

Percent organic matter = (dry weight – ash weight) / dry weight x 100

Sediment percent carbonate content was determined following 2 hr at 950°C and 2 hr of cooling in a

desiccator, and calculated using the equation:

Percent carbonate content = (ash weight$_{435}$ – ash weight$_{950}$) / ash weight$_{435}$ x 100

2.4 | Data analysis: clam size, growth and survival, and sediment characteristics

Data were analyzed in R v.4.0.2 (R Core Team 2022). Model assumptions of normality were assessed

using histograms, quantile-quantile plots of the data, and the Shapiro-Wilk test. Homogeneity of the

residuals was assessed using plots of model residuals versus fitted values and Bartlett's test for

homogeneity of variance. Nested ANOVAs (plots nested within beach location), using the linear mixed-

effects function in the *nlme* package, were used to test for differences between clam garden and

unmodified reference beaches (*i.e.*, beach type, two levels) in initial and final clam shell heights, wet

weights, percent growth, and percent survival as well as individual sediment characteristics (*i.e.*, rocks,

small rocks, very coarse sand, coarse sand, sand, fine sand, very fine sand, silt, percent organic matter, and

percent carbonate concentration), with beach location (six levels) included as a random factor.

Significance was considered when $p \leq 0.05$. ANOVAs, linear mixed-effects models, and post-hoc least

squares means omitting beach type were used to test for significant differences among beaches in percent

growth, percent survival, sediment carbonates, organics, and individual grain sizes. Principal component

analysis (PCA) and biplot figures were generated in R using the prcomp function in the default *stats*

package and the *ggbiplot* function/package. Pearson correlation between the variables was evaluated with

the cor function of the *stats* package of R and inspected with *corrplot* v.0.92 (Wei and Simko 2021). All

code is available (see *Data Availability*).

2.5 | RNA sampling and extraction

For gene-expression analysis, small (~2 x 2 mm) sections of gill and digestive gland tissues were excised

from each surviving clam using sterile techniques and stored in RNAlater as per manufacturer protocols

(Ambion). Bivalve gills are a target tissue in studies of rapid environmental responses, whereas digestive

glands are accumulatory organs used in toxicological and immunology investigations of longer-term

effects of environmental change and exposure (Gosling 2008, Milan et al. 2011). RNA from tissue

sections of 25–30 mg was individually extracted from both tissues using RNeasy kits (Qiagen) after

homogenization in 2-mL tubes filled with Lysing Matrix D (MP Biomedicals) in a Tissuelyser II (Qiagen)

at 25 Hz for 2 min. To eliminate DNA contamination, a DNase protocol was applied (Turbo DNA-free

Kits, Ambion) that inactivates and removes the DNase enzyme after DNA digestion. RNA concentrations

were quantified on a NanoDrop 1000 Spectrophotometer (Thermo Fisher Scientific) after DNase

treatment.

2.6 | Library preparation for RNA sequencing

RNA sequencing pools were created for each of the six locations (beaches) using five haphazardly

selected individuals from each plot within the beach. Plots were only used if they had at least five

surviving clams. This resulted in three plots per beach being used for all locations except for beaches E

and F, which only had five or more survivors in a single plot. This produced 14 RNA pools per tissue (N =

28 pools total) to be used for library synthesis and sequencing. RNA quantities were normalized, and

equimolar pools of RNA were created for both tissues of the five selected individuals from each plot.

RNA quality and quantity of each pool were determined using the RNA 6000 Nano chip on a 2100

Bioanalyzer (Agilent).

Following RNA quality control checks (NanoDrop and Bioanalyzer), mRNA libraries were generated

from 250 ng of total RNA using an mRNA enrichment with the NEB Next Poly(A) Magnetic Isolation

Module (New England Biolabs) and cDNA synthesis was conducted using NEBNext RNA First Strand

Synthesis and NEBNext Ultra Directional RNA Second Strand Synthesis Modules. Remaining library

preparation used NEBNext Ultra II DNA Library Prep Kit for Illumina, with adapters and PCR primers

(NEB). The remaining steps of library preparation were taken using the Quant-iT™ PicoGreen® dsDNA

Assay Kit (Life Technologies) and the Kapa Illumina GA with Revised Primers-SYBR Fast Universal Kit

(Kapa Biosystems (Pty) Ltd.). Average fragment size was determined using a LabChip GX instrument

(PerkinElmer). Libraries were run across four lanes of an Illumina HiSeq4000 PE 100 platform (Illumina)

and the mean (± SD) read pairs per library (both tissues combined) was 54.4 ± 9.5 million. All library

preparation and sequencing steps were undertaken at the Genome Québec Innovation Centre (Montreal,

Canada).

2.7 | Transcriptome assembly and annotation

For the *de novo* clam transcriptome assembly, Trinity 2.5.1 (--min_kmer_cov 2) (Grabherr et al. 2011)

was used to assemble the 28 libraries. Before running the assembly, Trimmomatic 0.36

(ILLUMINACLIP:TruSeq3-PE-2.fa:2:30:10 LEADING:3 TRAILING:3 SLIDINGWINDOW: 4:15

MINLEN:36) was used to clean the read data to obtain a more accurate assembly (Bolger et al. 2014).

With the resulting Trinity assembly (1,695,678 sequences), the TransDecoder 5.0.1 pipeline (Haas et al.

2013) was used to predict likely coding sequences (1,277,478 sequences), which uses homology (pfam-a,

UniProt) and open reading frame (ORF) information. The best representative transcript was chosen for

each gene, as predicted by TransDecoder, that had an ORF type of 'complete', which required protein

homology. In cases where there were multiple transcripts with complete ORFs for a gene, the transcript

with the largest ORF was chosen. Further filtering was carried out to choose only transcripts $\geq$ 100 bp (to

remove small transcripts that could cause analysis error), followed by scanning of pfam-a and Uniprot

annotations to remove repetitive element-related keywords to finally produce a transcriptome of 52,000

transcripts, each putatively representing a single gene of which 42,708 had Uniprot IDs.

2.8 | Data analysis: RNA sequencing data

Raw reads were imported to the repository *Simple_reads_to_counts* (see *Data Availability*) for

quantitation. Raw reads were gently trimmed for quality and adapters removed using Trimmomatic

(Bolger   et   al.   2014)   with   the   following   flags: ILLUMINACLIP:$VECTORS:2:30:10;

SLIDINGWINDOW:20:2; LEADING:2; TRAILING:2; MINLEN:80. Both the raw and the trimmed data

were inspected using FastQC (Andrews 2010) and multiQC (Ewels et al. 2016). The reference

transcriptome was indexed using bowtie2 (Langmead and Salzberg 2012) and reads were aligned against it

in local mode, allowing 40 alignments to be retained per read (-k 40). For comparison with the results

from   the   reference   transcriptome,   the   Manila   clam   (*Ruditapes   philippinarum*)   genome

(GCA_009026015.1_ASM902601v1) (Yan et al. 2019) was also tested as a potential reference by

indexing the genome using hisat2 (Kim et al. 2015) and aligning the Pacific littleneck clam reads against

the Manila clam genome (also retaining a maximum of 40 alignments per read; -k 40). The littleneck clam

alignments against the littleneck reference transcriptome were quantified using eXpress using default

parameters (Roberts and Pachter 2013). The alignments against the Manila clam genome in the test were

also quantified using eXpress default parameters but with the flag *--rf-stranded*. Resultant effective read

counts quantified against the *de novo* reference transcriptome were used in downstream analyses.

Transcriptome analyses were all conducted as described in the repository *ms_clam_garden* (see *Data*

Availability). Transcript annotation, effective counts, and phenotypic data were all imported into R to be

analyzed with *edgeR* v.3.36.0 (Robinson et al. 2010). Datasets were constructed for all samples or only gill

or only digestive gland (DG) samples (N = 3 datasets). Each dataset was input into a DGEList, then

filtered for low expression by requiring at least 10 reads to be aligned to the transcript in at least five

individuals (*i.e.*, counts per million (cpm) > 0.82–1.03, depending on the dataset). Samples were

normalized by TMM normalization as applied by *calcNormFactors()* and multidimensional scaling (MDS)

plots were produced for each dataset (*i.e.*, both tissues combined, gill tissue only, DG tissue only).

Tissue-specific expression was determined by identifying transcripts expressed in one tissue and not

the other. The normalized counts for these transcripts were then obtained from the all-sample dataset and

200 of the highest expressed tissue-specific transcripts from both tissues were plotted using *heatmap()* of

the *stats* package in R. Gene Ontology (GO) enrichment was conducted using the tissue-specific

transcripts from both tissues compared to all expressed transcripts for that tissue. Enrichment analysis was

conducted in DAVID Bioinformatics (Huang et al. 2007).

Differential expression analysis was conducted using *limma* v.3.50.1 (Ritchie et al. 2015). Both

tissues were analyzed separately and differentially expressed genes were identified using beach type (*i.e.*,

CG vs. Ref) as an explanatory variable using a gene-wise negative binomial generalized linear model with

quasi-likelihood tests (glmQLFit) and retaining all transcripts with $p \leq 0.001$. Differentially expressed

genes were also identified based on survivorship, where the percent survival of each plot was used to bin

the plot into one of high, medium, or low survival based on the fourth quartile, second and third quartiles,

or the first quartile of percent survival, respectively. Statistical significance was conducted as above but

using contrasts between each survival group. All other analyses and plots were conducted in R and are

available in the repository listed above.

3 | RESULTS

3.1 | Clam growth and survival

There were no significant differences between CG and Ref beaches in any measured growth or survival variable (ANOVAs, $p \geq 0.168$, Table 1). The mean ($\pm$ SD) initial clam wet weight and shell height for CG and Ref beaches (N = 180) was 1.5 ± 0.2 g and 1.8 ± 0.3 g and 1.5 ± 0.1 cm and 1.6 ± 0.1 cm, respectively (Table 1). After 16 weeks *in situ*, these increased to 3.9 ± 1.3 g and 3.8 ± 2.9 g and 1.8 ± 0.3 cm and 1.9 ± 0.4 cm, which in terms of height, was an increase in growth of $20.5 \pm 15.3\%$ and $18.0 \pm 22.6\%$, respectively. Calculating percent growth using height includes both the alive and dead clams and is therefore considered to be a more accurate measurement of growth compared to using wet weights. Wet weights only include surviving clams, and therefore does not capture the reduced growth that may have occurred in clams that eventually died. Clam survival was $63.3 \pm 28.3\%$ in CG and $58.3 \pm 37.0\%$ in Ref beaches, respectively.

Beach location (independent of the CG/Ref factor), however, significantly affected all growth and survival variables (Table 2, *SI* Figure S1). An unplanned difference occurred in the initial size of clams among the six beaches (initial wet weight and shell height ANOVAs $p = 0.031$ and $p = 0.044$, respectively, Table 2, *SI* Figure S2). This difference was specifically due to significantly larger clams inadvertently planted on beach D (Table 2, *SI* Figure S2). Percent growth was highest on beaches A ($31.2 \pm 7.2\%$), C ($29.7 \pm 2.8\%$), and D ($47.0 \pm 8.2\%$) and lowest in beaches B ($6.6 \pm 6.8\%$), E ($0.8 \pm 0.9\%$), and F ($0.3 \pm 0.6\%$) (Table 2), while percent survival was highest in beaches A ($88.3 \pm 12.6\%$), B ($71.7 \pm 27.5\%$), C ($68.3 \pm 12.6\%$), and D ($86.7 \pm 5.8\%$) and lowest in beaches E ($33.3 \pm 23.6\%$) and F ($16.7 \pm 24.7\%$) (Table 2).

Gross observations showed that most of the survivors on beaches A (CG), B (Ref), C (CG), and D

(Ref) were considered healthy (Table 3). These beaches had 68–88% survival, with the percentage of

clams considered weak (0–5%), emaciated (0–5%), or having very pale digestive glands (0–22%) being

generally low, with the exception of beach D, where 22% of clams had pale digestive glands (Table 3). In

contrast, survivors from beaches E (CG) and F (Ref) had only 17–33% survival, and had more clams

considered weak (13–15%), emaciated (15–22%), and with digestive glands being very pale (13–22%)

than the other beaches. Beaches E and F are therefore considered poorer performing sites.

3.2 | Beach sediment characteristics

There were no significant differences between CG and Ref beaches for any of the sediment characteristics

(ANOVAs, $p \geq 0.392$, Table 1). Beach location (independent of the CG/Ref factor), however, significantly

affected all sediment characteristics except percentage of small rocks (Table 2). A PCA of these sediment

characteristics separates the beach locations into three groups (Figure 2), where pairs of beaches are

clustered based on geographical locations (Figure 1). Beach A (CG), on a west-facing bay, was exposed to

wave action from within and outside Kanish Bay. Beach A grouped with beach B (Ref), an east-facing bay

that was also exposed to wave action within Kanish Bay (Figure 1). These beaches are the two sites that

are closest to, but on opposite sides of, the entrance of Kanish Bay and show correlation with coarse sand

and sand (Figure 2). Beach C (CG) is a south-facing sheltered beach within Small Inlet and groups with

beach D (Ref), a north-facing beach on the opposite side of Small Inlet. Beaches C and D correlated with

rocks, fine sand, and final weight/shell height (Figure 2). Beach E (CG) and F (Ref), two proximate

beaches on the south side of Kanish Bay, face north and are exposed to wave action. These sites correlated

with carbonates, organics, and small rocks, and are opposite to percent survival (Figure 2). Notably, these

beach groupings remain constant when including only abiotic characteristics in the PCA or when

including only survival and growth variable (data not shown). The PCA was also run on the data but

excluding beach D (due to the aforementioned initial size difference of this beach), and this showed

similar associations and groupings, although beach C grouping on its own (*SI* Figure S3).

Most sand types, silt, and rocks were either positively or negatively correlated with growth and

survival variables (*SI* Figure S4). To investigate correlations between sediment characteristics and clam

growth and survival further, beach D was again excluded as described above (*SI* Figure S5). Abiotic

variables with significant positive associations with growth included sand ($p = 0.018$, $R^2 = 0.31$), and fine

sand ($p = 0.022$, $R^2 = 0.29$). Negative associations with growth were silt ($p = 0.002$, $R^2 = 0.49$), and small

rocks ($p = 0.014$, $R^2 = 0.34$). Variables with significant positive associations with survival were coarse

sand ($p = 0.014$, $R^2 = 0.34$), and sand ($p = 0.002$, $R^2 = 0.48$). Negative associations with survival included

carbonates ($p = 0.008$, $R^2 = 0.39$), organics ($p = 0.035$, $R^2 = 0.24$), as well as small rocks ($p = 0.021$, $R^2 =$

0.30), very fine sand ($p = 0.002$, $R^2 = 0.49$), and silt ($p = 0.002$, $R^2 = 0.51$) (*SD* Additional File S1, without

Beach D). .

3.3 | Transcriptomic overview and *de novo* reference transcriptome

The gill and DG libraries produced on average ($\pm$ SD) 53.8 ± 10.7 and 55.0 ± 8.3 M read pairs per library

respectively. All 28 libraries, comprising 305 GB of high-quality sequence data, were used to create a *de*

novo transcriptome assembly (Table 4; see *Data Availability*). This assembly was comprised of 50,000

contigs (*i.e.*, putative transcripts), of which 42,708 were given annotation by sequence similarity to the

SwissProt database.

Alignments of the quality-controlled reads resulted in a mean ($\pm$ SD) overall alignment rate of 27.87

$\pm 2.48\%$ against the Pacific littleneck clam *de novo* transcriptome. For comparison, the alignment rate

against the Manila clam (*i.e.*, the closest relative with a reference genome) resulted in very low alignments

of 1.10 ± 0.17% (gill), 2.59 ± 1.00% (DG). All downstream analyses used the *de novo* transcriptome

quantified transcripts. After filtering the dataset containing all 28 samples, 33,825 transcripts (67.7%)

were expressed in at least five samples above the applied threshold (see *Materials and Methods*).

An MDS plot of samples from both tissue types showed that gill samples were distinctly separate

from DG samples across the first dimension, explaining 65% of the variation in the dataset (Figure 3). The

second dimension, which explained only 3% of the variation separated the DG samples, but not the gill

samples. One outlier sample was observed in the gill (P12).

3.4 | Tissue-specific expression

Samples from each tissue were then analyzed separately after conducting tissue-specific filtration and

normalization. The gill and DG samples on average (± SD) had 12.9 ± 3.0 M and 15.5 ± 2.9 M reads

aligning per library and the datasets were found to have 28,391 and 24,699 transcripts expressed,

respectively, after the low expression filter. Inspecting transcripts that were present in only one of the

tissue types identified 8,795 (31.0%) gill-specific transcripts and 5,103 (20.7%) DG-specific transcripts

(Table 5; *SD* Additional File S2). The highest expressed tissue-specific transcripts are shown in *SI* Figure

S6 (n = 200 shown for each tissue).

Tissue-specific transcripts were compared to all expressed genes in that tissue to identify enriched

GO categories in the tissue-specific list. Gill-specific transcripts were enriched for biological adhesion (N

= 496), ion transport (N = 494), signal transduction (N = 1,184), response to stimulus (N = 1,839), G-

protein coupled receptor signalling pathways (N = 210), and defense response (N = 402, $p < 0.0001$),

among others (*SD* Additional File S3). DG-specific transcripts were enriched for biological adhesion (N =

308), defense response (N = 263), immune response (N = 223), innate immune response (N = 149),

humoral immune response (N = 58), organonitrogen compound catabolic process (N = 130), defense

response to bacteria (N = 82), and ion transport (N = 201, $p < 0.0001$), among others (*SD* Additional File

S3).

3.5 | Effect of clam gardens on gene expression

The gill dataset analyzed by multidimensional scaling (MDS) has 13% and 11% of the total variation

explained by dimensions 1 and 2, respectively (Figure 4A). In general, in the gill samples, no clear

clustering was observed by beach type (*i.e.*, CG vs. Ref) nor any other variables (*e.g.*, survival, sand, silt,

or carbonates). There were two outlier samples (P12 from beach B and P7 from beach D) in the MDS plot,

both of which were Ref beaches and had 45% and 80% survival, respectively.

The DG dataset analyzed by MDS resulted in 14% and 9% of the total variation being explained by

dimensions 1 and 2, respectively (Figure 4B). Some grouping was observed along dimension 2 with CG

samples clustering together, except for sample P15 from beach E. Ref samples were more spread out along

both axes. A gradient was observed across dimension 1 along survival, with high survival being positioned

in negative dimension 1, medium survival near 0‑1, and low survival in positive dimension 1, although

the trend was not universal for all samples. Percent carbonate, which was correlated with survival (*see*

above), also trended along dimension 1 in the opposite direction to survival (*i.e.*, high survival was

associated with low percent carbonate, *SI* Figure S7).

Differential expression between CG and Ref was evaluated for each tissue separately (Table 5). The

gill had 56 transcripts differentially expressed as a result of CG (FC > 1.5, $p \leq 0.001$, no multiple test

correction (MTC)) (see Discussion; *SD* Additional File S2). Of these transcripts, 35 had identifiers

recognized by DAVID Bioinformatics, but no Gene Ontology (GO) enrichment was identified ($p > 0.01$).

Most of these transcripts were overexpressed in the CG samples (N = 42), including several transcripts

associated with immune/apoptotic or stress-related functions—two transcripts annotated as *tumour*

necrosis factor (TNF)-related (FC > 1.5), one as *death effector domain (DED)*-related (*i.e.*, top

overexpressed transcript, FC > 71.5), and *universal stress protein A-like protein* (FC > 4.7). In total, 14

transcripts were underexpressed in the CG, including a *heat shock 70 kDa protein 12a* (FC = 17.9). A full

list of differentially expressed genes is available in *SD* Additional File S2.

The DG had 37 transcripts differentially expressed between CG and Ref samples. This included 11

overexpressed and 26 underexpressed transcripts in CG (*SD* Additional File S2). Using the full,

bidirectional list of differentially expressed transcripts, several enriched GO categories were observed (*SD*

Additional File S3), including biological process categories iron ion homeostasis (N = 3, p = 0.008) and

negative regulation of hydrolase activity (N = 5, p < 0.001). The highest overexpressed transcript in the

CG samples was *replicase polyprotein* annotated from the *cricket paralysis virus* (FC > 500), which

suggests that viral proteins were also captured in the study and that they were differentially abundant

between beach types. Multiple complement-, iron-, and protein-folding-related transcripts were observed

in the DG, including overexpression of a transcript annotated as *c1q domain* (2.3-fold), *thrombospondin-1*

(2-fold), *soma ferritin*, and *cytochrome P450* (FC > 1.5). Transcripts under-expressed in the DG included

interferon-induced very large GTPase 1 (FC > 8.5), *apoptosis inhibitor IAP* (4.3-fold), *low affinity*

immunoglobulin epsilon Fc receptor (FC > 2), *c-type mannose receptor 2*, and *platelet endothelial*

aggregation receptor 1 (FC > 1.5), among others (*SD* Additional File S2).

Consistency was observed between the tissues in the CG response, where two of the 91 unique

differentially expressed transcripts were differentially expressed in both tissues. This is notable given that

the datasets were normalized, filtered, and analyzed separately. This included the under expression in the

CG samples of a transcript annotated as *von Willebrand factor type D domain (vwd)* and of a transcript

annotated as *platelet endothelial aggregation receptor 1* (*pear1*; Figure 5) in both tissues.

3.6 | Gene expression responses and differential survival

Given the differential survival observed among the beaches and plots, genes correlating with survival were investigated to determine what might be driving these differences and to potentially identify genes that are related to survival or mortality in this species. Genes were of particular interest that were differentially expressed between low and medium survival, between medium and high survival, or incrementally differentially expressed at each step between high, medium, and low survival (see *Materials and Methods* for beach survivorship classification).

The majority of differentially expressed genes by survival in the gill were overexpressed between the low and medium survival groups, rather than between medium and high survival (Table 5). Transcripts overexpressed in the low survival group relative to the medium survival group (N = 26) were involved in functions such as heat-shock, interferon response, and ligase activity (*SD* Additional Files S2, S3). For example, *heat-shock protein (hsp) 90* (FC > 60) and three different transcripts annotated as subunits of *hsp70* (FC > 2) were overexpressed specifically in the low survival group. Transcripts overexpressed in the medium survival group relative to the high survival group included six transcripts of various functions (*SD* Additional File S2). Fewer transcripts were overexpressed in the higher survival group (*i.e.*, high vs. medium, N = 11; medium vs. low, N = 14) and these generally were involved in immune functions, but also various other functions. Overexpression of *toll-like receptor 1* (FC > 2) and *complement C1q-like protein 4* (FC > 3.5) was observed in medium survivors relative to low survivors.

Similar to the gill, in the DG the majority of the differentially expressed transcripts by survival were overexpressed in low relative to medium surviving clams (N = 34) and were generally involved in immune system or heat-shock or transport. Transcripts annotated as *hsp70 12a* and *12b*, as well as several aminopeptidases, were overexpressed in low relative to medium and in medium relative to high survival (Figure 6). All the overexpressed genes in medium relative to high surviving clams (N = 18) were also

overexpressed in the low relative to medium survival group comparison. Very few genes were

overexpressed in the high survival clams relative to medium (N = 4) or the medium survival clams relative

to low survival (N = 8), and these transcripts had various functions, including a transcript annotated as

mucin-2 protein (Figure 6).

4 | DISCUSSION

4.1 | Effects of clam gardens on clam and beach characteristics

Results from the present study showed that there was no significantly-enhanced growth or survival on

unmaintained clam garden (CG) versus reference (Ref) non-walled beaches. Using similarly-sized clams

that were collected on the same days and at the same location as the present work Salter (2018) also

reported no significantly enhanced growth or survival on unmaintained CG compared to non-walled

beaches (controls) in Kanish and Waiatt Bays. It should be noted that the clam gardens in the present study

were not being maintained according to traditional tending practices and this maintenance may influence

clam survival, growth, and transcriptomic profile, as well as various abiotic characteristics (*i.e.*, sediment

grain size, carbonate content, organic content). Salter (2018), however, also included some CG traditional

tending practices (*i.e.*, adding a 1 cm-thick layer of crushed shell mixture three times throughout the

summer) on replicate treatment plots on both beach types and reported a positive effect of the shell hash

treatment on growth and survival in both CG and non-walled beaches. Therefore, both the present study

and Salter (2018) did not observe a positive effect of CG on growth and survival in the absence of tending

but Salter (2018) reported a positive effect with the addition of crushed shell hash in both beach types.

Some trends were observed in the transcriptomic data that suggest that the CG had a transcriptomic

effect on the clams, albeit not a strong effect. Most notably, consistent genes were identified in two tissues

even though very few genes were found to be differentially expressed, and the CG and Ref samples

showed a slight separation in the PCA of DG tissue. The two genes that were identified as underexpressed

in the CG in both tissues were annotated as *von Willebrand factor type D domain (vwd)* and *platelet*

endothelial aggregation receptor 1 (pear1). The former is present in many proteins, including mucins and

other extracellular glycoproteins, with one in particular being reported to be up-regulated in response to

heat shock in the Pacific oyster, *Crassostrea gigas* (Zhang et al. 2015). The function of the latter in

shellfish remains unknown; in vertebrates it is a cell membrane protein involved in platelet aggregation.

Homologous genes of platelet activation are present in the Hong Kong oyster (*Crassostrea*

hongkongensis) and are regulated in response to hyposalinity (Xiao et al. 2018). In terms of the other

genes differentially expressed by clams in CG, the general functions were not clear. Immune genes and

stress-response genes were found both over- and under-expressed in CG versus Ref (*e.g.*, a heat-shock

protein transcript was found to be underexpressed in the CG and a universal stress protein-annotated

transcript was overexpressed in the CG). In general, the functions of these transcripts in the Pacific

littleneck clam responding to the CG factor will be useful to continue to characterize as the genomic

information develops for this species and additional transcriptomic studies are conducted in relation to

environmental stress and maintained or unmaintained clam gardens. These molecular phenotypes may

indicate a response to the CG from the 16-week *in situ* study period that is below the level of growth and

survival phenotypic responses. Further work with longer study periods would be required to determine

whether these transcriptome signatures precede macro-phenotypic responses.

Sediment grain sizes, percent carbonate content, and percent organic matter did not differ between

CG and Ref in the present study, although there was a significant level of variation with these factors

within both beach types and among locations. For example, carbonate content varied from 4 to 10% in CG

and from 2 to 15% in Ref beaches, while silt varied from 0 to 4% and from 1 to 5%, respectively. This is

somewhat contrary to Salter's (2018) study that in untreated plots found that CG contained 2.8 – 12.9

times more carbonate and a smaller percentage of silt than non-walled beaches (Salter 2018). The reason

for this discrepancy is not known and would be valuable to address with future studies. The lack of

differences observed in the present study in growth and survival, as well as only a minor effect being

observed in the clam molecular phenotypes, from being grown in CG relative to Ref beaches, may in part

be due to the lack of tending of the clam gardens, as is traditionally conducted. Future studies would

benefit from fully examining the impacts of tending practices on sediment characteristics alongside clam

growth and survival, which would involve clam garden beach restoration and active maintenance for long

periods, with further analysis of impacts on shellfish molecular phenotypes and productivity.

4.2 | Abiotic variation among beach location and effects on clam growth and survival

Across the six sampling locations, beaches differed in most of the variables examined (*i.e.*, biotic response

variables: growth, survival; abiotic characteristics: organic level, carbonate concentration, grain sizes).

Beaches clustered based on these sediment characteristics and differential growth and survival were noted

among the locations. Given these trends, abiotic factors potentially involved in survival as well as

molecular responses to survival were characterized.

The sedimentary environment is very important for clam growth and survival (Joo et al. 2021).

Littleneck clams typically live as infaunal burrowers between 5 and 15 cm depth, in soft sand or sandy

mud in the intertidal zone (Lazo 2004). In the present study, growth was affected by small rocks, sand,

fine sand and silt. Survival was affected by carbonates, organics, small rocks, coarse sand, sand, very fine

sand, and silt. Joo et al. (2021) also found that grain size and sorting of the sediment determined the

amount of pore-water dissolved oxygen and organic matter, both affecting growth and survival of juvenile

Manila clams, with optimal survival and growth linked to poorly-sorted sediment with an average grain

size of medium sand (180 – 335 µm). Similarly, the grain size of the sand in the present study, which had

a significant positive effect on survival, was 250 – 500 µm. Salter (2018) found increased water residency

time positively affected clam biomass and growth and reported a weak negative effect of silt on clam

density and biomass in bivalve surveys and contrary to the present study also found a positive effect of

increased sediment silt content on clam biomass, in clam transplant experiments. An early multi-year

study looking at environmental factors affecting growth in the hard clam *Venus mercenaria* (now

Mercenaria mercenaria) found a consistent decrease in growth with increasing sediment silt-clay content

and (Pratt and Campbell 1956). All their results showed a negative relationship between growth and

fineness of the sediment (Pratt and Campbell 1956). Permeability of the sediment is an important factor

for burrowing filter feeders' pumping activity, which is essential for respiration, excretion, and nutrition

and Pratt et al. (1956) found permeability to be an inverse function of the sediment silt-clay content. In

addition, commercial Manila clam beds studied in Spain had higher survival of clams planted in sand-

gravel than in those placed in mud (Cigarria and Fernandez 2000).

Sediment carbonate content had a significant negative effect on clam survival in the present study,

where sediment with more than 8% carbonates had markedly decreased growth and survival. Similarly,

Munroe (2016) found that the average daily growth of early post-settled Manila clams over two years was

negatively correlated with sediment carbonate content (Munroe 2016). This is not a universally observed

trend in the literature (Groesbeck et al. 2014, Salter 2018) and counters expectations given the expected

importance of carbonate for clam shell growth and early recruitment. The addition of crushed shell hash

alters sediment calcium-carbonate saturation states, as it increases surface sediment porewater aragonite

content and pH (Green et al. 2009). For instance, there was a three-fold increase in hard clam burrowing

and recruitment by adding crushed shell hash to a final concentration of 8% of wet sediment weight

(Green et al. 2012). Salter (2018) increased carbonate content significantly (1.9–7.2 times) by addition of

shell hash, which increased clam growth and tissue biomass in both clam garden and control beaches after

six months (Salter 2018).

Crushed shell hash also changes the physical structure of the sediment as, like gravel, it can create

interstitial spaces in the sediment, thereby increasing pore-water dissolved oxygen, organic-matter content,

and substrate stability, as well as providing protection from predators for juvenile clams (Joo et al. 2021,

Sponaugle and Lawton 1990, Thompson 1995). Thus, there may be potentially multiple benefits of adding

shell material for bivalve growth/survival that are not linked to carbonate concentration. The effect of shell

hash or crushed shell on bivalve growth and survival is likely dependent on a wide variety of factors

including amount of shell, size and compactness of shell particles, species generating the shell hash,

species/size of bivalve grown, stocking densities, tidal height, temperature, and types/densities of

predators. For example, the application of very finely crushed shell hash could have a negative effect by

decreasing interstitial spaces in the sediment and thus, as seen with silt, decreasing sediment permeability.

Recent studies investigating the efficacy of shell hash to mitigate acidification of intertidal sediments

found it unsuccessful at raising pore-water pH of acidic sediments (Beal et al. 2020, Doyle and Bendell

2022). Doyle et al. (2022) found the addition of shell hash to be site dependent as it reduced variation in

pH at an intertidal site with pH 8.03 but did not mitigate the pH of mud flats with pH 7.59. At this point it

appears that more research is required to comprehensively understand the effects of shell hash on bivalve

growth and survival.

4.3 | Pacific littleneck clam transcriptomics

In non-model organisms, such as the Pacific littleneck clam, advances in molecular ecological and

physiological understanding of the species can be advanced through transcriptomic tools and the results

these tools generate. Here it was clear that the reference transcriptome generated *de novo* was necessary

for the study, given that the closest relative genome (*i.e.*, Manila clam) resulted in a very low alignment

rate from the Pacific littleneck clam reads. Further, the use of different tissues in the present study

revealed high levels of tissue-specific expression (*i.e.*, 21–31% of the transcripts expressed in the tissue).

As multiple individuals were included in each sample (*i.e.*, pools), the true number of individuals sampled

for the tissue-specific expression analysis is greater than the number of libraries, increasing the reliability

of the tissue-specific genes. The tissue-specific expression analysis indicated that for both tissues, tissue-

specific transcripts were involved in signal transduction, defense and immune response, and ion transport,

among other processes. Therefore, although the specific transcripts are different, similar functions are

enriched in the tissue-specific gene lists for both the gill and the DG, although exceptions exist.

The presence of differential survival among the locations provided a valuable opportunity for

analyzing transcripts that are associated with mortality/survival. These characterizations, over time

through multiple transcriptome studies generating more evidence, can begin to characterize specific

transcripts that are involved with specific stress types (Evans et al. 2011, Miller et al. 2017, Sutherland et

al. 2012). This can enable the development of Ecological Association Ontology data collection for non-

model species (Pavey et al. 2012) and in the future may be used to characterize underlying factors

associated with observed mortality or morbidity (*e.g.*, abiotic stress compared to bacterial or viral

infection). Although the Pacific littleneck clam remains largely uncharacterized at the molecular level,

some of the genes identified in the present study correlated with survival, including differential expression

in the gill and DG of heat-shock proteins, immune response genes, and other functional categories.

Additional transcriptomic studies on the species in different environmental conditions over different

temporal and spatial scales should help to improve these categorizations and identify stressor-specific or

generalized response genes.

615 Differential expression was observed in transcripts involved in immune defense. For example,

616 transcripts annotated as complement C1q protein complex or involved in cell adhesion and oxidation-

617 reduction processes had high expression in high surviving beaches and may indicate increased activity of

618 immune defense responses. Complement C1q proteins are an important component of the innate immune

619 system (Zhang et al. 2014) and are upregulated in response to bacterial challenges (McDowell et al. 2014)

620 and salinity stress (Zhao et al. 2012). Cell adhesions are critical for the development, maintenance, and

621 function of multicellular organisms and are essential for invertebrate immunity (Johansson 1999).

622 In the present study, potentially due to the relatively low sample size from pooling individuals and

623 due to having fewer plots with a sufficient number of survivors at some beaches and the high number of

624 expressed transcripts, it was not possible to apply multiple test corrections. Therefore, a more stringent *p*-

625 value was applied ($p < 0.001$), but the associations of genes to CG or differential survival should be taken

626 with caution. Further studies confirming these functions will be beneficial to understand their role in clam

627 transcriptomics.

629 **4.4 | Viral presence and clam health**

630 The use of RNA-sequencing for the present analysis identified a differentially expressed virus transcript in

631 the CG versus Ref samples, which could have unexpected effects on either gene expression activity or

632 phenotypic responses, as could any other uncharacterized factor influencing clam growth/physiology (*e.g.*,

633 temperature, beach exposure). The top overexpressed transcript in the DG from CG was a *replicase*

634 *polyprotein* from the *cricket paralysis virus* (CPV), which is a positive-sense, single-stranded RNA virus

635 from the family Dicistroviridae in the order Picornavirales. Members of the Dicistroviridae family are

636 widely distributed in nature with the broadest host range of all small RNA viruses in invertebrates

637 (Bonning 2009). As the Dicistroviridae viral sequences isolated in the present study ranged in size from

1479 to 7371 nucleotides, one can deduce that they represent viral mRNA, instead of viral genomes. Some

dicistroviruses can be non-pathogenic or cause subtle impacts (reduced longevity and fecundity), while

others result in rapid paralysis (Bonning 2009).

Even though there was transcriptomic evidence of viral presence in all the beaches, most of the

Pacific littleneck clams from the medium- and high-survival groups did not exhibit any signs of

pathophysiology. Other clams (mostly from the low-survival group), however, were emaciated or watery

and had slightly to extremely pale-coloured digestive glands, an indicator of reduced feeding and

potentially compromised function and health. Pale digestive glands could also be the result of paramyxean

parasites responsible for disease in marine mollusks, with infection being linked to digestive tropism

interfering with food adsorption and causing pale-yellowish digestive glands with thin, watery flesh in

both mussels and oysters (Alfjorden 2017, Carella et al. 2011). In the present study, however, there was no

significant parasite gene expression at 16 weeks and histopathology would be needed to confirm any

paramyxean parasite infections. It is also worth noting that the single time point of sample collection in the

present study and the analysis of only surviving clams may have reduced the likelihood of viewing viral or

parasitic gene expression in these samples.

Although very pale digestive glands and emaciation was observed in clams from beaches with lower

growth and survival, histopathology or genetic identification would again be required to confirm any viral

gene expression linkages to observed impacts on organismal health. Health status is most likely linked to

multiple stressors comprised of abiotic (*e.g.*, temperature, sediment composition, water residency) and

biotic factors (*e.g.*, bacteria, viruses, harmful algae). RNA sequencing combined with *de novo*

transcriptome assembly revolutionizes virus detection and allows for the screening of a broad range of

symptomatic and asymptomatic virus species in a non-discriminatory manner, thereby displaying a

broader picture of the environmental factors present (Nagano et al. 2015). The current dataset has yet to be

fully characterized for viral sequences and could be mined further at a later date, particularly as more

resources are developed for this species.

CONCLUSIONS

The *de novo* assembly of a Pacific littleneck clam reference transcriptome enabled the detection of a small

but significant differentially expression response in clams from CG and Ref beaches. This was observed

even though no significant phenotypic differences in growth or survival were observed, nor differences in

sediment characteristics between the two unmaintained clam gardens or reference sites. However,

significant phenotypic differences in growth and survival and in sediment characteristics between beaches

allowed for the discovery of several sediment characteristics significantly correlated with growth and

survival. The unexpected result of carbonates having a negative effect on survival may partially be due to

the lack of clam garden tending practices and future work would benefit from examining the effects these

practices and shell-hash have on bivalve health. Transcripts related to stress response were identified by

conducting differential expression analysis among beaches of largely differing survival. The characterized

transcriptome provides a valuable resource for future transcriptomic studies investigating genetic and

phenotypic responses of clams in different habitats and environmental conditions.

Data Deposition: Raw RNA-sequencing data have been uploaded to SRA under BioProject
PRJNA818991, BioSamples SAMN26893882–SAMN26893909.
680

ACKNOWLEDGEMENTS

Funding for this study was provided by the Hakai Institute. We acknowledge the K'ómoks (Comox)
Nation, the Xwémalhkwu (Homalco) Nation, the We Wai Kai (Cape Mudge) Nation, the We Wai Kum
(Campbell River) Nation, the Snuneymuxw Nation, the lək̓ʷəŋən (Songhees) Nation, the WSÁNEĆ
(Saanich) Nation and the Wyomilth (Esquimalt) Nation, which have stewarded and protected the territory
we worked within since time immemorial. We thank the Hakai Institute, the Tula Foundation, and the
Clam Garden Network for facilitating the research and for their continued support. We also thank Caitlin

DATA AVAILABILITY

693 Supplemental materials, including all abiotic and phenotypic site metadata needed for data analysis,
694 transcript annotation, quantified gene expression counts, sample phenotypes:
695 Reference transcriptome: https://doi.org/10.6084/m9.figshare.19735753
696 RNA-seq bioinformatics pipeline: https://github.com/bensutherland/Simple_reads_to_counts
697 RNA-seq analysis and physiological/abiotic analysis pipeline: https://github.com/raapm/ms_clam_garden
698

SUPPLEMENTAL DATA

700 **Additional File S1.** Abiotic variables with survival and growth linear model p and R^2 values, including
701 and excluding beach D.
702
703 **Additional File S2.** Differentially expressed genes and background gene lists for both tissues in Pacific
704 littleneck clams. Gene lists include clam garden vs. reference, survival groups, and tissue-specific
705 expression.
706
707 **Additional File S3.** Gene Ontology enrichment results for the clam garden analysis, the survival analysis,
708 and the tissue-specific expression.
709

REFERENCES

Alfjorden, A., M. Areskog, D. Bruno, R. Carnegie, D. Cheslett, S. Feist, S. Ford, S. Jones, A. Lillehaug, L. Madsen, T. Renault, N. Ruane and P. Vennerström. (2017) New Trends in Important Diseases Affecting the Culture of Fish and Molluscs in the ICES Area 2002 – 2015, p. 50 pp, International Council for the Exploration of the Sea, Conseil International pour l'Exploration de la Mer, Copenhagen, Denmark.

Andrews, S. (2010) FastQC: A quality control tool for high throughput sequence data. Available online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc.

Beal, B.F., Coffin, C.R., Randall, S.F., Goodenow, C.A., Pepperman, K.E. and Ellis, B.W. (2020) Interactive effects of shell hash and predator exclusion on 0-year class recruits of two infaunal intertidal bivalve species in Maine, USA. Journal of Experimental Marine Biology and Ecology 530-531.

Bolger, A.M., Lohse, M. and Usadel, B. (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics 30(15), 2114-2120.

Bonning, B.C. (2009) The Dicistroviridae: An emerging family of invertebrate viruses. Virologica Sinica 24(5), 415-427.

Bourlat, S.J., Borja, A., Gilbert, J., Taylor, M.I., Davies, N., Weisberg, S.B., Griffith, J.F., Lettieri, T., Field, D., Benzie, J., Glockner, F.O., Rodriguez-Ezpeleta, N., Faith, D.P., Bean, T.P. and Obst, M. (2013) Genomics in marine monitoring: new opportunities for assessing marine health status. Mar Pollut Bull 74(1), 19-31.

Caldwell, M.E., Lepofsky, D., Combes, G., Washington, M., Welch, J.R. and Harper, J.R. (2012) A Bird's Eye View of Northern Coast Salish Intertidal Resource Management Features, Southern British Columbia, Canada. The Journal of Island and Coastal Archaeology 7(2), 219-233.

Carella, F., Aceto, S., Maiolino, P. and De Vico, G. (2011) What is your diagnosis? Pale yellowish digestive gland and watery tissues in Mediterranean mussels. Vet Clin Pathol 40(2), 273-274.

Chadwick, M., Harper, E.M., Lemasson, A., Spicer, J.I. and Peck, L.S. (2019) Quantifying susceptibility of marine invertebrate biocomposites to dissolution in reduced pH. R Soc Open Sci 6(6), 190252.

Cigarria, J. and Fernandez, J.M. (2000) Mangement of manila clam beds I. Influence of seed size, type of substratum and protection on initial mortality. Aquaculture 182, 173-182.

Cox, K.D., Gerwing, T.G., Macdonald, T., Hessing-Lewis, M., Millard-Martin, B., Command, R.J., Juanes, F., Dudas, S.E. and Birchenough, S. (2019) Infaunal community responses to ancient clam gardens. ICES Journal of Marine Science 76(7), 2362-2373.

Deur, D., Dick, A., Recalma-Clutesi, K. and Turner, N.J. (2015) Kwakwaka'wakw "Clam Gardens". Human Ecology 43(2), 201-212.

Doyle, B. and Bendell, L.I. (2022) An evaluation of the efficacy of shell hash for the mitigation of intertidal sediment acidification. Ecosphere 13(3).

Erlandson, J.M., Rick, T.C., Braje, T.J., Steinberg, A. and Vellanoweth, R.L. (2008) Human impacts on ancient shellfish: a 10,000 year record from San Miguel Island, California. Journal of Archaeological Science 35(8), 2144-2152.

Evans, T.G., Hammill, E., Kaukinen, K., Schulze, A.D., Patterson, D.A., English, K.K., Curtis, J.M. and Miller, K.M. (2011) Transcriptomics of environmental acclimatization and survival in wild adult Pacific sockeye salmon (*Oncorhynchus nerka*) during spawning migration. Mol Ecol 20(21), 4472-4489.

Evans, W., Mathis, J.T. and Cross, J.N. (2014) Calcium carbonate corrosivity in an Alaskan inland sea. Biogeosciences 11(2), 365-379.

Ewels, P., Magnusson, M., Lundin, S. and Kaller, M. (2016) MultiQC: summarize analysis results for multiple tools and samples in a single report. Bioinformatics 32(19), 3047-3048.

Gosling, E. (2008) Bivalve Molluscs: Biology, Ecology and Culture, Wiley.

Grabherr, M.G., Haas, B.J., Yassour, M., Levin, J.Z., Thompson, D.A., Amit, I., Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke, A., Rhind, N., di Palma, F., Birren, B.W., Nusbaum, C., Lindblad-Toh, K., Friedman, N. and Regev, A. (2011) Full-length transcriptome assembly from RNA-Seq data without a reference genome. Nat Biotechnol 29(7), 644-652.

Green, M.A., Waldbusser, G.G., .L., R.S., Emerson, K. and O'Donnell, S. (2009) Death by dissolution Sediment saturation state as a mortality factor for juvenile bivalves. American Society of Limnology and Oceanography 54(4), 1037-1047.

Green, M.A., Waldbusser, G.G., Hubazc, L., Cathcart, E. and Hall, J. (2012) Carbonate Mineral Saturation State as the Recruitment Cue for Settling Bivalves in Marine Muds. Estuaries and Coasts 36(1), 18-27.

Greiner, C.M., Klinger, T., Ruesink, J.L., Barber, J.S. and Horwith, M. (2018) Habitat effects of macrophytes and shell on carbonate chemistry and juvenile clam recruitment, survival, and growth. Journal of Experimental Marine Biology and Ecology 509, 8-15.

Groesbeck, A.S., Rowell, K., Lepofsky, D. and Salomon, A.K. (2014) Ancient clam gardens increased shellfish production: adaptive strategies from the past can inform food security today. PLoS One 9(3), e91235.

Haas, B.J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P.D., Bowden, J., Couger, M.B., Eccles, D., Li, B., Lieber, M., MacManes, M.D., Ott, M., Orvis, J., Pochet, N., Strozzi, F., Weeks, N., Westerman, R., William, T., Dewey, C.N., Henschel, R., LeDuc, R.D., Friedman, N. and Regev, A. (2013) De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. Nat Protoc 8(8), 1494-1512.

Heiri, O., Lotter, A.F. and Lemcke, G. (2001) Loss on ignition as a method for estimating organic and carbonate content in sediments: reproducibility and comparability of results. Journal of Paleolimnology 25, 101-110.

Huang, D.W., Sherman, B.T., Tan, Q., Kir, J., Liu, D., Bryant, D., Guo, Y., Stephens, R., Baseler, M.W., Lane, H.C. and Lempicki, R.A. (2007) DAVID Bioinformatics Resources: expanded annotation database and novel algorithms to better extract biology from large gene lists. Nucleic Acids Res 35(Web Server issue), W169-175.

Jackley, J., Gardner, L., Djunaedi, A.F. and Salomon, A.K. (2016) Ancient clam gardens, traditional management portfolios, and the resilience of coupled human-ocean systems. Ecology and Society 21(4).

Johansson, M.W. (1999) Cell adhesion molecules in invertebrate immunity. Dev Comp Immunol 23(4-5), 303-315.

Joo, S., Jo, K., Bae, H., Seo, H. and Kim, T. (2021) Optimal sediment grain size and sorting for survival and growth of juvenile Manila clams, *Venerupis philippinarum*. Aquaculture 543.

Kim, D., Langmead, B. and Salzberg, S.L. (2015) HISAT: a fast spliced aligner with low memory requirements. Nat Methods 12(4), 357-360.

Langmead, B. and Salzberg, S.L. (2012) Fast gapped-read alignment with Bowtie 2. Nat Methods 9(4), 357-359.

Lazo, D.G. (2004) Bivalve Taphonomy: Testing the effect of life habits on the shell condition of the
Littleneck clam *Protothaca (Protothaca) staminea* (Mollusca: Bivalvia). PALAIOS 19 (5)(Friday
Harbor Laboratory Special Issue), 451-459.
Lepofsky, D., Smith, N.F., Cardinal, N., Harper, J., Morris, M., Gitla, Bouchard, R., Kennedy, D.I.D.,
Salomon, A.K., Puckett, M. and Rowell, K. (2017) Ancient Shellfish Mariculture on the
Northwest Coast of North America. American Antiquity 80(02), 236-259.
Lepofsky, D., Toniello, G., Earnshaw, J., Roberts, C., Wilson, L., Rowell, K. and Holmes, K. (2020)
Ancient Anthropogenic Clam Gardens of the Northwest Coast Expand Clam Habitat. Ecosystems.
McDowell, I.C., Nikapitiya, C., Aguiar, D., Lane, C.E., Istrail, S. and Gomez-Chiarri, M. (2014)
Transcriptome of American oysters, *Crassostrea virginica*, in response to bacterial challenge:
insights into potential mechanisms of disease resistance. PLoS One 9(8), e105097.
Milan, M., Coppe, A., Reinhardt, R., Cancela, L., Leite, R.B., Saavedra, C.C., Chelazzi, G., Bortoluzzi, S.
and Bargelloni, L. (2011) Transcriptome sequencing and microarray development for the Manila
clam, *Ruditapes philippinarum:* genomic tools for environmental monitoring. BMC Genomics 12,
234.
Miller, K.M., Gunther, O.P., Li, S., Kaukinen, K.H. and Ming, T.J. (2017) Molecular indices of viral
disease development in wild migrating salmon(dagger). Conserv Physiol 5(1), cox036.
Munroe, D. (2016) Habitat effects on early post-settlement growth of intertidal clams,*Venerupis
philippinarum*(A. Adams & Reeve, 1850). Journal of Molluscan Studies 82(4), 507-514.
Nagano, A.J., Honjo, M.N., Mihara, M., Sato, M. and Kudoh, H. (2015) Detection of plant viruses in
natural environments by using RNA-Seq. Methods Mol Biol 1236, 89-98.
Neudorf, C.M., Smith, N., Lepofsky, D., Toniello, G. and Lian, O.B. (2017) Between a rock and a soft
place: Using optical ages to date ancient clam gardens on the Pacific Northwest. PLoS One 12(2),
e0171775.
Pavey, S.A., Bernatchez, L., Aubin-Horth, N. and Landry, C.R. (2012) What is needed for next-generation
ecological and evolutionary genomics? Trends Ecol Evol 27(12), 673-678.
Pratt, D.M. and Campbell, D.A. (1956) Environmental Factors Affecting Growth in *Venus mercenaria*.
Limnology Oceanography (January).
R.Core.Team (2022) R: A language and environment for statistical computing. Computing, R.F.f.S. (ed),
Vienna, Austria.
Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W. and Smyth, G.K. (2015) limma powers
differential expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res
43(7), e47.
Roberts, A. and Pachter, L. (2013) Streaming fragment assignment for real-time analysis of sequencing
experiments. Nat Methods 10(1), 71-73.
Robinson, M.D., McCarthy, D.J. and Smyth, G.K. (2010) edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. Bioinformatics 26, 139-140.
Salter, N. (2018) Ancient clam gardens magnify bivalve production by moderating ambient termperature
and enhancing sediment carbonate, Simon Fraser University, Vancouver, B.C.
Smith, N.F., Lepofsky, D., Toniello, G., Holmes, K., Wilson, L., Neudorf, C.M. and Roberts, C. (2019)
3500 years of shellfish mariculture on the Northwest Coast of North America. PLoS One 14(2),
e0211194.
Sponaugle, S. and Lawton, P. (1990) Portunid crab predation on juvenile hard clams: effects of substrate
type and prey density. Marine Ecology progress series 67, 43-53.

Sutherland, B.J., Jantzen, S.G., Yasuike, M., Sanderson, D.S., Koop, B.F. and Jones, S.R. (2012)
Transcriptomics of coping strategies in free-swimming *Lepeophtheirus salmonis* (Copepoda)
larvae responding to abiotic stress. Mol Ecol 21(24), 6000-6014.
Sutherland, B.J.G., Finke, J.F., Saunders, R., Warne, S., Schulze, A.D., Strohm, J.H.T., Chan, A.M.,
Suttle, C.A. and Miller, K.M. (2022) Metatranscriptomics reveals a shift in microbial community
composition and function during summer months in a coastal marine environment. bioRxiv
preprint doi: https://doi.org/10.1101/2022.01.21.477286; this version posted August 15, 2022.
Thompson, D.S. (1995) Substrate additive studies for the development of hardshell clam habitat in waters
of Puget Sound in Washington State: An analysis of effects on recruitment, growth, and survival
of the Manila clam, *Tapes philippinarum*, and on the species diversity and abundance of existing
benthic organisms. Estuarine Research Federation 18(No. 1A), 91-107.
Waldbusser, G.G. and Salisbury, J.E. (2014) Ocean acidification in the coastal zone from an organism's
perspective: multiple system parameters, frequency domains, and habitats. Ann Rev Mar Sci 6,
221-247.
Wei, T. and Simko, V. (2021) R package 'corrplot': Visualization Matrix (Version 0.92). Available from
http://github/taiyun/corrplot
Wentworth, C.K. (1922) A scale of grade and class terms for clastic sediments. Journal of Geology, 377 -
393.
Williams, J. (2006) Clam Gardens: Aboriginal Mariculture on Canada's West Coast, New Star Books Ltd,
Canada.
Xiao, S., Wong, N.K., Li, J., Lin, Y., Zhang, Y., Ma, H., Mo, R., Zhang, Y. and Yu, Z. (2018) Analysis of
in situ Transcriptomes Reveals Divergent Adaptive Response to Hyper- and Hypo-Salinity in the
Hong Kong Oyster, *Crassostrea hongkongensis.* Front Physiol 9, 1491.
Yan, X., Nie, H., Huo, Z., Ding, J., Li, Z., Yan, L., Jiang, L., Mu, Z., Wang, H., Meng, X., Chen, P., Zhou,
M., Rbbani, M.G., Liu, G. and Li, D. (2019) Clam Genome Sequence Clarifies the Molecular
Basis of Its Benthic Adaptation and Extraordinary Shell Color Diversity. iScience 19, 1225-1237.
Zhang, L., Li, L., Zhu, Y., Zhang, G. and Guo, X. (2014) Transcriptome analysis reveals a rich gene set
related to innate immunity in the Eastern oyster (*Crassostrea virginica*). Mar Biotechnol (NY)
16(1), 17-33.
Zhang, Y., Sun, J., Mu, H., Li, J., Zhang, Y., Xu, F., Xiang, Z., Qian, P.Y., Qiu, J.W. and Yu, Z. (2015)
Proteomic basis of stress responses in the gills of the Pacific oyster *Crassostrea gigas*. J Proteome
Res 14(1), 304-317.
Zhao, X., Yu, H., Kong, L. and Li, Q. (2012) Transcriptomic responses to salinity stress in the Pacific
oyster *Crassostrea gigas*. PLoS One 7(9), e46244.
877
878
879

TABLES

TABLE 1. Average (± SD) clam size/survival and sediment characteristic variables for clam garden (CG) and reference (Ref) beaches and associated linear model *p*-values (with beach location as a nested effect). No variables were significantly affected by the CG/Ref factor or showed a significant deviation from normality as measured by Shapiro Wilk test.
[a] Significant Bartlett's test therefore failing the linear model assumption due to heteroscedasticity.

	Variable	CG Mean ± SD	Ref Mean ± SD	*p*-value
Clam size/survival	Initial weight (g)	1.5 ± 0.2	1.8 ± 0.3	0.200
	Initial shell height (cm)	1.5 ± 0.1	1.6 ± 0.1	0.168
	Final weight (g)	3.9 ± 1.3	3.8 ± 2.9	0.943
	Final shell height (cm)	1.8 ± 0.3	1.9 ± 0.4	0.864
	Growth (%) in height	20.5 ± 15.3	18.0 ± 22.6	*0.892[a]*
	Survival (%)	63.3 ± 28.3	58.3 ± 37.0	0.860
Sediment characteristics	Carbonates (%)	7.6 ± 1.8	7.2 ± 5.2	0.927
	Organics (%)	1.3 ± 0.6	1.1 ± 0.5	0.723
	Rocks (%)	15.6 ± 7.5	15.0 ± 5.4	0.912
	Small rocks (%)	12.1 ± 5.0	13.0 ± 3.3	0.765
	Very coarse sand (%)	12.2 ± 3.4	13.0 ± 3.6	0.794
	Coarse sand (%)	14.2 ± 3.4	13.2 ± 4.5	*0.788[a]*
	Sand (%)	20.0 ± 5.7	17.6 ± 4.2	0.575
	Fine sand (%)	18.6 ± 6.4	17.0 ± 2.9	0.715
	Very fine sand (%)	5.6 ± 2.1	8.3 ± 4.4	*0.393[a]*
	Silt (%)	2.0 ± 1.2	3.1 ± 1.5	*0.392[a]*

890 **TABLE 2:** Average (± SD) clam size/survival and sediment characteristic variables for each beach and associated linear model *p*-values.

	Variable	Beach A Mean ± SD	Beach B Mean ± SD	Beach C Mean ± SD	Beach D Mean ± SD	Beach E Mean ± SD	Beach F Mean ± SD	*p*-value
Clam size/survival	Initial weight (g)	1.6 ± 0.1	1.8 ± 0.2	1.5 ± 0.3	2.0 ± 0.1	1.5 ± 0.2	1.5 ± 0.2	3.1E-02
	Initial height (cm)	1.5 ± 0.0	1.6 ± 0.1	1.5 ± 0.1	1.7 ± 0.1	1.5 ± 0.1	1.5 ± 0.1	4.4E-02
	Final weight (g)	4.2 ± 0.5	2.5 ± 0.7	5.1 ± 0.6	7.5 ± 0.8	2.5 ± 1.1	1.4 ± 1.3	2.5E-05
	Final height (cm)	2.0 ± 0.1	1.7 ± 0.2	2.0 ± 0.2	2.4 ± 0.2	1.5 ± 0.1	1.5 ± 0.1	7.5E-06
	Growth (%) in height	31.2 ± 7.2	6.6 ± 6.8	29.7 ± 2.8	47.0 ± 8.2	0.8 ± 0.9	0.3 ± 0.6	*5.1E-07*[a]
	Survival (%)	88.3 ± 12.6	71.7 ± 27.5	68.3 ± 12.6	86.7 ± 5.8	33.3 ± 23.6	16.7 ± 24.7	3.1E-03
Sediment characteristics	Carbonates	8.0 ± 1.0	5.3 ± 2.5	6.0 ± 2.0	2.7 ± 0.6	8.7 ± 1.5	13.7 ± 1.2	5.3E-05
	Organics	0.7 ± 0.1	0.9 ± 0.3	1.4 ± 0.3	0.8 ± 0.2	1.8 ± 0.6	1.7 ± 0.3	4.8E-03
	Rocks	8.0 ± 5.6	11.7 ± 6.1	19.0 ± 4.6	20.7 ± 1.5	19.7 ± 6.8	12.7 ± 2.1	3.7E-02
	Small rocks	11.3 ± 5.1	11.7 ± 4.6	8.3 ± 2.3	11.7 ± 1.2	16.7 ± 3.8	15.7 ± 2.1	1.1E-01
	Very coarse sand	13.7 ± 2.1	17.0 ± 2.0	6.7 ± 2.5	9.7 ± 2.5	14.3 ± 2.5	12.3 ± 1.2	5.3E-03
	Coarse sand	18.0 ± 2.0	19.0 ± 1.7	11.7 ± 3.1	10.7 ± 1.5	13.0 ± 0.0	10.0 ± 0.0	*1.1E-04*[a]
	Sand	25.7 ± 5.8	22.3 ± 3.5	19.7 ± 2.1	16.0 ± 1.0	14.7 ± 1.5	14.3 ± 1.5	3.2E-03
	Fine sand	19.3 ± 4.7	14.7 ± 3.1	24.7 ± 3.2	17.7 ± 3.1	11.7 ± 0.6	18.7 ± 1.5	3.8E-03
	Very fine sand	3.3 ± 0.6	3.0 ± 0.0	7.0 ± 1.7	10.7 ± 3.5	6.3 ± 1.5	11.3 ± 1.5	3.8E-04
	Silt	0.7 ± 0.6	1.7 ± 0.6	2.0 ± 0.0	3.0 ± 1.0	3.3 ± 0.6	4.7 ± 0.6	*7.5E-05*[a]

891 [a] Significant Bartlett's test therefore failing the linear model assumption due to heteroscedasticity

892 **TABLE 3.** Gross observations (health, condition and digestive gland coloration) of surviving clams in
893 each beach (A-F).

Metric	Beach A	Beach B	Beach C	Beach D	Beach E	Beach F
Health (% weak)	0	5	0	2	13	15
Condition (% emaciated)	0	5	0	2	22	15
Digestive gland (% pale)	8	7	5	22	0	0
Digestive gland (% very pale)	0	0	0	2	22	13

894
895 **TABLE 4.** Overview of RNA-seq reads and quality scores. Totals that include both reads in the pair are
896 indicated by (R1+R2). Values are shown in billion (B) or million (M).
897

Leukoma staminea assembly	Value
Total input reads (R1 + R2)	3.0 B
Total input nucleotides (R1 + R2)	305 B
Average (± SD) input reads per:	
single HiSeq4000 lane (R1 + R2)	762 ± 13.0 M
library (read pairs)	54.4 ± 9.5 M
gill library (read pairs)	53.8 ± 10.7 M
digestive-gland library (read pairs)	55.0 ± 8.3 M
Assembly contig length range	102 – 29,916
Average (± SD) Phred quality score	38.9 ± 0.3
Base-calling accuracy	> 99.9%

899
900 **TABLE 5.** Differentially expressed gene overview comparing beach type (clam garden (CG) vs.
901 reference beach (Ref)), beach survival (high vs. medium vs. low), tissue type (gill vs. digestive gland), or
902 expressed in background. Note that here differential expression is considered when $p \leq 0.001$ and log2
903 fold-change > 0.58 (1.5-fold overexpressed) or < -0.58 (1.5-fold underexpressed).
904

Comparison	Result	# Transcripts in gills	# Transcripts in digestive gland
Beach type (CG vs. Ref)	Overexpressed CG	40	11
	Underexpressed CG	14	26
Beach survival (high vs. medium vs. low)	Overexpressed high vs. medium	11	4
	Overexpressed medium vs. low	14	8
	Overexpressed low vs. medium	26	34
	Overexpressed medium vs. high	6	18
Tissue type (gill vs. digestive gland)	Gill vs. digestive gland	8,795	5,103
Background	N/A	28,391	24,699

906 **FIGURES**
907

908
909 **FIGURE 1.** Locations of experimental clam gardens (A, C, and E, designated by triangles), unmodified
910 (reference) beaches (B, D, and E, designated by circles), and clam collection site (designated by a star) on
911 Quadra Island, British Columbia, Canada.

912   A

913
914
915   **FIGURE 2.** Principal component analysis biplot showing beach groupings including beach D and
916   correlations in terms of percent survival and growth of Pacific littleneck clams and various beach
917   sediment characteristics. Each beach is shown as A–F (as per Figure 1 locations) and plots within beach
918   location as 1–3. Colours of plots and ellipses are for visualization purposes only.

920
921
922 **FIGURE 3.** All sample MDS plot showing groupings of gill (G) and digestive gland (DG) tissues of
923 Pacific littleneck clams sampled from plots P1–P18 (*i.e.*, Beach A: P1–P3; Beach B: P10–P12; Beach C:
924 P4–P6; Beach D: P7–P9; Beach E: P15; Beach F: P18).
925

926 **A**

927
928 **B**

929
930
931 **FIGURE 4.** MDS plots for (A) gill and (B) digestive gland tissues of Pacific littleneck clams where each
932 datapoint is plotted as clam garden (CG) or reference beach (Ref) with the following information: %
933 survival, % carbonates, beach location identifier (A–E), and plot identifier (P1–P18) (*i.e.*, Beach A: P1–
934 P3; Beach B: P10–P12; Beach C: P4–P6; Beach D: P7–P9; Beach E: P15; Beach F: P18).
935

936
937
938 **FIGURE 5.** Box plots showing expression of genes of interest identified in both digestive glands and
939 gills of Pacific littleneck clams, which were underexpressed in clam gardens (CG) relative to reference
940 beaches (Ref). Vwd = *von Willebrand factor type D domain*; pear1 = *platelet endothelial aggregation*
941 *receptor 1*.

943
944
945 **FIGURE 6.** Linear regression of differential expression of selected genes of interest versus percent
946 survival in Pacific littleneck clams. Transcript expression and their correlation to survival include (top left
947 to bottom right) *heat shock protein family A member 12A* (*hsp70-12a*), *aminopeptidase*, and *mucin-2* in
948 the digestive gland (DG) and *hsp70-12a*, *hsp90*, and *toll-like receptor 1* in the gill.
949

950 **SUPPORTING INFORMATION**
951
952

953
954
955 **FIGURE S1.** Boxplots of (A) percent survival and (B) percent growth of Pacific littleneck clams by
956 beach (A–F).
957

958
959
960 **FIGURE S2.** Boxplots of (A) input weight and (B) input height of Pacific littleneck clams by beach (A–
961 F).

963
964 **FIGURE S3.** Principal component analysis biplot showing beach groupings and correlations in terms of
965 survival and growth of Pacific littleneck clams and various beach sediment characteristics excluding
966 beach D. Each beach plot is shown as A-F (as per Figure 1 locations) and 1-3 depending on the plot
967 within the beach location. Colours of plots and ellipses are for visualization purposes only.

981
982 **FIGURE S4.** Pearson correlations between biotic and abiotic variables including growth and survival
983 including beach D.
984

985
986 **FIGURE S5.** Pearson correlations between biotic and abiotic variables including growth and survival of
987 Pacific littleneck clams excluding beach D.
988

989
990
991

992
993
994 **FIGURE S6.** Highest expressed tissue-specific transcripts in the gill (G) and digestive gland (DG) tissues
995 of Pacific littleneck clams within each plot (P1–P18) (*i.e.*, Beach A: P1–P3; Beach B: P10–P12; Beach C:
996 P4–P6; Beach D: P7–P9; Beach E: P15; Beach F: P18). Samples were clustered based on expression of
997 these transcripts, with similarly expressing sample grouping. The higher the expression, the darker the
998 colour of the cell, with each row as an individual transcript.

1004

1005
1006
1007   **FIGURE S7.** Percent survival of Pacific littleneck clams versus percent carbonate in sediments. Beach D
1008   is included in this figure, and when it is not included, the adjusted R2 = 0.39 (p = 0.008), see *SD*
1009   Additional File S1.