Motor and cognitive deficits limit the ability to flexibly modulate spatiotemporal gait features in older adults with mild cognitive impairment

Michael C. Rosenberg[1*], Alexandra Slusarenko[2], Ke Cao[3], J. Lucas McKay[4,5], Laura Emmery[6],
Trisha M. Kesar[7], Madeleine E. Hackney[3,7-10]

[1]Neuromechanics Laboratory, Department of Biomedical Engineering, Emory University & Georgia
Institute of Technology, Atlanta, GA, USA

[2]College of Arts and Sciences, Emory University, Atlanta, GA, USA

[3]Department of Medicine, Division of Geriatrics and Gerontology, Emory University School of
Medicine, Atlanta, GA, USA

[4]Department of Neurology, Emory University School of Medicine, Atlanta, GA, USA

[5]Department of Biomedical Informatics, Emory University School of Medicine, Atlanta, GA, USA

[6]Department of Music, Emory University College of Arts and Sciences, Atlanta, GA, USA

[7]Department of Rehabilitation Medicine, Emory University School of Medicine, Atlanta, GA, USA

[8]Emory University School of Nursing, Atlanta, GA, USA

[9]Atlanta VA Center for Visual & Neurocognitive Rehabilitation, Atlanta, GA, USA

[10]Birmingham/Atlanta VA Geriatric Research Education and Clinical Center, Atlanta, GA, USA

* Correspondence:
Michael C. Rosenberg
mcrose3@emory.edu

Keywords: mild cognitive impairment, dance therapy, gait analysis, biomechanics, music,
rhythm, gait modifications

Abstract

Dance-based therapies are an emerging form of movement therapy aiming to improve motor and
cognitive function in older adults with mild cognitive impairments (MCIs). Despite promising effects
of dance-based therapies on function, it remains unclear how age-related declines in motor and
cognitive function affect movement capacity and influence which movements and rhythms maximize
dance therapy efficacy. Here, we evaluated the effects of age and MCI on the ability to accurately
modulate spatial (*i.e.*, joint kinematics), temporal (*i.e.*, step timing), and spatiotemporal features of
gait to achieve spatial and temporal targets during walking. We developed novel rhythmic movement
sequences - nine spatial, nine temporal, and four spatiotemporal - that deviated from typical spatial
and temporal features of walking. Healthy young adults (HYA), healthy older adults (HOA), and
adults with MCI were trained on each gait modification before performing the modification
overground, with kinematic data recorded using wearable sensors. HOA performed spatial (p =
0.010) and spatiotemporal (p = 0.048) gait modifications less accurately than HYA. Individuals with
MCI performed spatiotemporal gait modifications less accurately than HOA (p = 0.017). Spatial
modifications to the swing phase of gait (p = 0.006, Cohen's *d* = -1.3), and four- and six-step *Duple*

rhythms during temporal modifications ($p \leq 0.030$, Cohen's $d \geq 0.9$) elicited the largest differences in
gait performance in HYA vs. HOA and HOA vs. MCI, respectively. These findings suggest that age-
related declines in strength and balance reduce the ability to accurately modulate spatial gait features,
while declines in working memory in individuals with MCI may reduce the ability to perform longer
temporal gait modification sequences. Differences in rhythmic movement sequence performance
highlight motor and cognitive factors potentially underlying deficits in gait modulation capacity,
which may guide therapy personalization and provide more sensitive indices to track intervention
efficacy.

Introduction

Dance-based therapy, which purposefully sets movements to music, is an emerging form of
movement therapy for individuals with mild cognitive impairments (MCIs), a precursor to dementia
or Alzheimer's involving reduced attention, executive function, spatial cognition, and working
memory, that elicit improvements in cognitive function in individuals with MCI (Gauthier et al.,
2006; Lazarou et al., 2017; Sanford, 2017; Zhu et al., 2018; Zhu et al., 2020). Dance-based therapy is
a logical approach for improving cognitive function, as exercise or general physical activity are
among the few interventions shown to improve cognitive function in individuals with MCI (Geda et
al., 2010; Thom and Clare, 2011). Specifically, Thom and Clare (2011) noted that physical exercise
programs that target functional tasks, aerobic endurance, and balance are likely needed to maximize
the functional benefits of therapy, all of which are challenged during dance (Thom and Clare, 2011).
Currently, it is unclear how to personalize dance-based therapy parameters for individuals with MCI.
Studies of dance-based therapy use different music, dance types, and study designs (Hackney and
Earhart, 2009; McKee and Hackney, 2013; Lazarou et al., 2017; Zhu et al., 2020). Improving our
understanding of the relationships between individual-specific deficits associated with MCI and the
ability to perform different movements during dance-based therapy is needed to inform the objective
design of personalized dance-based therapies for individuals with MCI.

While a major goal of dance-based therapies is to maximize the motor and cognitive benefits of
moving to music, *how* to do so for a given population needs better clarity. Music type and tempo, and
the dance type or steps are basic parameters that can influence the difficulty of performing the
therapy routine. Therapies often rely on trial-and-error to refine a protocol, rather than being
grounded in a principled data-driven understanding of myriad factors that may impact dance-therapy
performance, including individual differences in motor control and cognitive function (McKee and
Hackney, 2013; Lazarou et al., 2017; Zhu et al., 2018). Selecting an appropriate level of challenge is
fundamental to maximizing the motor and cognitive effects of therapy (Guadagnoli and Lee, 2004).
Given the heterogeneity of cognitive deficits in MCI (Gauthier et al., 2006) and age-related declines
in motor function (Booth et al., 1994), understanding how both motor and cognitive deficits impact
the ability to perform challenging rhythmic movement patterns may both enable objective
personalization of dance-based therapy parameters and provide mechanistic insights into the impacts
of MCI on mobility.

Though not yet quantified in the context of dance-based therapy, dance can be considered a series of
Rhythmic Movement Sequences (RMS): rhythmic patterns involving modifications to spatial and
temporal components of normal stepping (Rallis et al., 2018). Importantly, spatial and temporal
features of movement correspond to, and may challenge, distinct neural representations (*e.g.,* the
premotor cortex) and multiple aspects of biomechanics function (*e.g.*, strength, balance, joint
coordination) (Wilson and Kwon, 2008; Kornysheva and Diedrichsen, 2014). A central premise of
this work is that evaluating the capacity for modulating spatial and temporal movement features will

shed light on motor and cognitive mechanisms that may inform future therapy individualization and
enhance the efficacy of dance and music-based movement therapy in individuals with MCI. Spatial
modifications of movement during dance involve altering the magnitude and coordination of hip,
knee, and ankle kinematics (Ivanenko et al., 2005; Wilson and Kwon, 2008). Voluntarily altering
coordination relative to highly stereotyped patterns during locomotion requires the motor capacity to
achieve joint excursions and increased cortical demands (Ivanenko et al., 2005; Cohen et al., 2016;
Hortobágyi et al., 2016; Rucco et al., 2017). Consequently, both motor and cognitive deficits may
impact individuals' abilities to perform modifications to the spatial components of movement.
Muscle function and motor control change with age, and correspond to reductions in overall motor
function in older adults (Booth et al., 1994; Schloemer et al., 2017). Therefore, age-related deficits in
motor function may impact individuals' abilities to modify spatial components of movement.
Because spatial modifications also require mapping perceived biomechanical targets to motor
actions, we expect these modifications to challenge aspects of cognitive function that are impaired in
individuals with MCI, such as spatial attention, executive function, and spatial cognition (Gauthier et
al., 2006; Rucco et al., 2017). However, how these motor and cognitive deficits impact individuals'
abilities to voluntarily modulate spatial aspects of movement remains unclear.

Temporal modifications of movement involve altering step timing to produce rhythmic sequences of
steps, which may challenge cognitive function (Kornysheva and Diedrichsen, 2014). During dance
therapy, temporal modifications can be elicited by altering the music's rhythm and meter, two
fundamental components of music (Hackney et al., 2007a; Hackney and Earhart, 2009). In the
context of stepping or walking, rhythm defines the pattern of temporal stimuli, such as taking two
quick (q) steps followed by two slow (S) steps. Meter is a perceived quantity that individuals
anticipate and to which they entrain their movements. Unimpaired walking typically involves evenly
recurring steps that are approximately rhythmically and metrically constant, such that modifying
temporal features during RMS would represent a deviation from normal gait rhythm and meter.
Because temporal modifications require attention and the ability to perceive rhythmic cues and map
them to motor actions, we expect it to challenge aspects of cognitive function that are impaired in
individuals with MCI, such as attention and working memory (Gauthier et al., 2006; Grahn, 2012;
Sanford, 2017). Because temporal modifications can be performed with normal joint coordination
patterns, they are less likely to challenge motor function than spatial modifications. Therefore, we
expect spatial modifications to movement during RMS to challenge both motor and cognitive
function, while temporal modifications should primarily challenge cognitive function. However, the
extent to which age-related motor and cognitive deficits impact the ability to modulate spatial and
temporal aspects of movement is unclear.

In this study, we leveraged biomechanical analysis, prototypical dance movements, and music theory
to systematically investigate the effects of age-related differences in motor and cognitive function on
the ability to perform spatial and temporal modifications to walking in individuals with MCI. We
hypothesized that differences in motor and cognitive function would impact individuals' abilities to
accurately modulate spatial and temporal features of walking, similar to those performed during
dance-based therapies. Specifically, we predicted that, when performing RMS gait modifications of
varying difficulty, healthy older adults (HOA) would perform spatial and spatiotemporal gait
modifications significantly less accurately than healthy young adults (HYA), suggesting an effect
associated with aging-related differences in motor function. Similarly, we predicted that individuals
with MCI would perform temporal and spatiotemporal RMS less accurately than HOA, suggesting an
effect associated with impaired cognitive function.

Materials and Methods

This study was approved by the Emory University Institutional Review Board (STUDY00003507).
All participants gave written informed consent prior to participation.

Study design and visits

We performed an observational cross-sectional study in healthy younger and older adults, and older
adults with MCI. All outcomes were collected during a single 2-4 hour study visit for each
participant. Participants who were unable to complete the protocol in a single visit completed it
during one additional visit within one week.

Participants

A total of 37 participants were recruited for the study, including younger adults (HYA, N=13), older
adults (HOA, N=12), and older adults with MCI (N=12). One HYA participant failed to complete the
protocol and was excluded from analysis. For all participants, inclusion criteria included the ability to
walk 20m without an assistive device, completion of at least 6 years of education or good work
history, no hospitalizations in the last 60 days, and proficiency in the English language. HYA
participants ages 18-35 years were included in this study and HOA participants ages 55 years and
older were included. Additional inclusion criteria for participants with MCI included amnestic MCI
defined using the Alzheimer's Disease Neuroimaging Initiative criteria and reduced executive
function, working memory, and spatial cognition according to standard clinical assessments (Phillips
et al., 2001; Vandierendonck et al., 2004; Mueller et al., 2005; Bowie and Harvey, 2006; Hackney et
al., 2013).

Motor and cognitive assessments

To characterize participants' motor and cognitive function, trained personnel administered the
following motor and cognitive assessments: Montreal Cognitive Assessment (MoCA), Reverse Corsi
Blocks test, Tower of London test, Trail Making Test, Four-Square Step Test, Body Position Spatial
Task, simple, manual, and cognitive Timed Up-and-Go (TUG), and the 30-second Chair Stand
(Lundin-Olsson et al., 1998; Jones et al., 1999; Morris et al., 2001; Phillips et al., 2001; Dite and
Temple, 2002; Vandierendonck et al., 2004; Nasreddine et al., 2005; Bowie and Harvey, 2006;
Hackney et al., 2013). The average (± 1SD) scores for each clinical assessment and corresponding p-
values denoting the probability of statistical differences in each score were computed using the
software package R. Older adults and individuals with MCI were well-matched in age and sex. As
expected, the MCI participants performed worse on the tests of motor and cognitive function. HYA
participants performed better than HOA or MCI participants on all assessments of motor and
cognitive function. For comparison to the tempi of temporal and spatiotemporal gait modifications,
we also recorded participants' stride frequencies at their preferred walking speeds.

RMS gait modifications

Participants performed normal walking at their self-selected and fast speeds, and 22 RMS consisting
of 9 spatial, 9 temporal, and 4 spatiotemporal gait modifications, described below. Participants were
randomly assigned to perform either the spatial or temporal modifications first. Spatiotemporal
modifications always occurred after spatial and temporal modification trials. Experimental conditions
within each sub-group (spatial, temporal, or spatiotemporal) were randomized.

We defined 64 spatial, temporal, and spatiotemporal gait modifications, constituting a movement
library, from which we selected 9 spatial, 9 temporal, and 4 spatiotemporal gait modifications of

varying levels of difficulty. The selected biomechanical (*i.e.,* spatial) and temporal features of each
modification are described in **Figure 1A**. The figure shows the kinematic configuration of each
movement and lists the target variables. Briefly, *spatial* modifications altered the magnitude of hip,
knee, and ankle movements and their coordination relative to normal walking (Ivanenko et al., 2005).
We selected three stance-phase modifications (ST), three swing-phase modifications (SW), and three
combined modifications in both the swing and stance phases (SWST). Participants were provided
target joint angles for each modification, such as flexing the hip and knee to 90 degrees during swing;
and watched a video of an expert dancer performing the movements. During ST and SW trials,
participants modified stance and swing-phase gait kinematics, respectively, in both legs
asymmetrically. During SWST trials, participants modified both swing and stance only in the left leg.

A) *Spatial gait modifications*

Swing		Stance	
Biomechanical Target	Gait modification	Biomechanical Target	Gait modification
▪ 45° knee flexion ▪ Max. ankle plantarflexion ▪ Toe at stance - leg ankle	Coupé

▪ 0° knee flexion ▪ Max. ankle plantarflexion throughout stance	Piqué
▪ 90° knee flexion ▪ Max. knee flexion ▪ Toe at stance - leg knee	Passé	▪ 30° knee flexion ▪ 30° ankle dorsiflexion	Tombé
▪ 90° hip flexion ▪ 90° knee flexion ▪ Max. ankle plantarflexion	Attitude	▪ 0° knee flexion ▪ Max. ankle dorsiflexion	Heel strike
▪ 90° hip flexion ▪ Max. early - swing knee flexion ▪ 0° late-swing knee flexion ▪ Max. ankle plantarflexion	Developpé	▪ 0° knee flexion ▪ Max. ankle plantarflexion in mid-stance	Relevé
▪ 90° hip flexion ▪ 0° knee flexion ▪ Max. ankle plantarflexion	Battement		

B) *Attitude*

Figure 1: Descriptions of subcomponents of the spatial gait modifications. or rhythmic movement sequences (RMS) evaluated in the current study. A) Spatial gait modifications were derived from ballet steps, noted in the boxes containing movement depictions. The left two columns correspond to swing-phase modifications, while the right two columns correspond to stance-phase modifications. Each depiction shows the modification where biomechanical targets were applied. Colored lines on each person denote biomechanical targets at the hip (purple), knee (orange), and ankle (red) kinematics that were used to evaluate movement performance. Bulleted lists describe the biomechanical targets for each gait modification, which were used to quantify movement performance. Bullets denote the biomechanical targets for each gait modification. Modifications were performed in pairs during spatial trials. B) Example of biomechanical targets for the *Attitude* gait modification. Black dashed lines denote biomechanical target values and gold stars denote where the target value was estimated. For example, during the *Attitude* modification, the swing-leg hip (left) and knee (middle) should be flexed to 90 degrees, and the ankle (right) should be maximally plantarflexed. Colored lines denote exemplary HYA (orange), HOA (blue), and MCI (red) participants. These participants highlight reduced RMS performance at the hip and knee in MCI and at the ankle in HOA and MCI

Temporal modifications altered the rhythmic patterns of movement, consisting of two, four, or six
steps taken at half-beat ("and;" denoted $\&$), one-beat ("quick;" denoted q), and two-beat ("slow;"
denoted S) intervals (**Figure 2**). The music notation used in **Figure 2** indicates the sequence and
duration of steps, with closed (black) notes corresponding to a step spanning one beat and open
(white) notes denoting steps spanning two beats. Performance was evaluated by each participant's
ability to take steps matching the beat and in the correct sequence of quick and slow steps.

A) Simple Duple (*2-count*)

B) Complex Duple (*2-count*)

C) Waltz (*3-count*)

Figure 2: Step sequences for temporal and spatiotemporal gait modifications. Step sequences consisted of 2, 4, or 6 steps. Each sequence specified a combination of very quick (*&* = half-beat), quick (*q* = one beat), and slow (*S* = two beats) steps. The numbers above the musical bars show the beat count for each rhythm. A) Simple duple sequences used a two-count rhythm in which the strong beat was the downbeat. B) Complex duple sequences also used a two-count rhythm, but the strong beat was not the downbeat. C) Waltz sequences used a three-count rhythm, such that sequences spanned either one or two measures.

Temporal modifications were split into three sub-groups: three simple duple, three complex duple,
and three waltz modifications. *Duple* music (Figure 2A & B) had a tempo of 100 beats per minute

(bpm) for all participants, while Waltz music (Figure 2C) had a tempo of 134 bpm for HYA but was
reduced to 104 bpm for the HOA and MCI groups. The faster Waltz tempo for the HYA group was
due to an error in coding the tempi for the first 8 participants. We corrected the tempo for the HOA
and MCI to ensure that all participants could perform the temporal modifications. We used the same
tempi across all participants, rather than personalizing the tempi to each participant's preferred
cadence to avoid giving participants different amounts of time to process and remember auditory
input, which could alter cognitive demands for each participant (Gauthier et al., 2006; Persad et al.,
2008; Montero-Odasso et al., 2012). The *simple duple* meter has a two-count grouping (1-2) where
the downbeat is the strong beat. In more *complex duple* meters, the emphasized beat falls on the weak
beat, which defies Western music listeners' expectations. Consequently, we expected the complex
duple to be more challenging than the simple duple. The *waltz* is in a compound duple meter (a fast
two-count where each is subdivided into three sub-beats; 1-2-3, 1-2-3) or simple triple (three-count;
1-2-3) meter. *Waltz* meters, unlike *simple* and *complex duple* meters, in which synchronization of
movement to music is easier because the same foot always lands on the downbeat, the three-count of
waltz rhythms causes the foot landing on the downbeat to alternate every measure (three steps).
Consequently, individuals accustomed to Western music traditions may have greater difficulty
synchronizing movements to the triple meter of *waltz* than to duple meters. The movements within
each sub-group varied in difficulty. During duple trials, participants were cued by a simplified
Libertango (Astor Piazzolla, 1974), with superfluous musical accents and cues eliminated to improve
the ability of participants with limited musical experience to identify temporal patterns in the music.
During waltz trials, participants were cued by a modified version of *Shostakovich Waltz No. 2*
(Dmitri Shostakovich, 1938). These musical pieces are much more complex than a metronome beat
used in locomotion studies, as the attention required to dissociate meter and rhythm from superfluous
auditory information can alter individuals' abilities to entrain to the tempo (Styns et al., 2007;
Moumdjian et al., 2019). For each modification, participants were provided a sequence of "quick"
and "slow" beats and practiced the sequence by following an instructional video: first clapping, then
tapping the foot, then shifting weight between the legs, then stepping in place, then walking
(Hackney and Earhart, 2010).

Spatiotemporal modifications combined spatial and temporal modifications of varying difficulty
(**Figure 1** & **Figure 2**). We selected these movements to vary the relative motor and cognitive
challenges of the movements. Participants were allowed to review spatial and temporal instructional
videos if needed. Following instructions for all modifications, the assessors would briefly practice (3-
5 sequences) the modifications with the participant to confirm understanding. Data were collected
immediately after understanding was confirmed. Names and descriptions of all gait modifications can
be found in the *MovementLibrary.csv,* with reference to **Figure 1** & **Figure 2**, and are described in
Supplemental – S1.

During each trial, participants wore 15 inertial measurement units (IMUs) in a standard configuration
(APDM, Inc., Portland, USA). We used validated commercial software associated with the IMUs to
estimate lower-limb hip, knee, and ankle flexion angles over the gait cycle (Mancini and Horak,
2016; Washabaugh et al., 2017). Participants began each trial in quiet stance until hearing a tone,
then began walking in a modified gait pattern defined by the trial's spatial, temporal, or
spatiotemporal RMS. When physically capable, participants walked overground along an 11m
walkway four times for each trial, providing 15-30 (spatial) or 15-60 (temporal) strides for each trial.
For participants with reduced physical capacity, we iteratively reduced the number of walkway
lengths traveled from four lengths down to one length based on each participant's physical capacity.
We ensured that each participant performed at least 15 strides for spatial modifications and 10
sequences for temporal and spatiotemporal modifications. However, all participants performed the

same gait modifications, regardless of the distance walked. Participants were allowed to rest as
needed and were provided snacks and water between trials. All assessments were administered by
trained personnel.

RMS performance

For each modification, we defined biomechanical (spatial and spatiotemporal trials) and temporal
(temporal and spatiotemporal trials) targets, defined *a priori* in our movement library (**Figure 1A**).
Biomechanical targets matched those provided during instruction. Temporal targets corresponded to
the step sequence of quick and slow steps. RMS performance was defined as a percent error relative
to biomechanical and temporal targets: the error relative to each target value, normalized by the
magnitude of the corresponding target (**Figure 1B**). For example, the hip flexion angle target for the
attitude movement, the star in **Figure 1B**, is 90 degrees during the swing phase of gait. In swing, the
knee should be flexed to 90 degrees, and the ankle maximally plantarflexed. The targets for each
spatial modification are listed in **Figure 1A**. Therefore, RMS performance reflected the accuracy
with which each participant performed a gait modification, with lower percent error reflecting more
accurate (*i.e.*, better) performance. For each trial and participant, we averaged across each trial's
biomechanical and temporal targets (Equation (1). For each gait modification class (*spatial,
temporal, spatiotemporal*), we computed the error the i^{th} participant made on the j^{th} trial of the k^{th}
target variable (e.g., peak knee flexion angle) as follows:

$$error_i = \frac{100}{J} \sum_j \left(\frac{1}{K} \sum_k \left| \frac{x_{i,j,k} - x_{target,j,k}}{x_{target,j,k}} \right| \right), \tag{1}$$

where $x_{i,j,k}$ denotes the i^{th} participant's outcome for the j^{th} trial and k^{th} target variable (*e.g.*, peak knee
flexion angle), and is a vector of outcomes over all available strides for that trial. The variable $x_{target,j,k}$
denotes the value of the j^{th} trial and k^{th} target variable. The variable J represents the number of trials
included in each modification class and K represents the number of pre-specified target variables for
each gait modification (*spatial*: 9 modifications, 2-3 target variables per trial; *temporal*: 9
modifications, 1 target variable per trial; *spatiotemporal*: 4 modifications, 3-4 target variables per
trial). Percent error for each target (multiple targets per trial; see Figure 1 & Figure 2) was averaged
to compute a single error value for the trial. When multiple trials were included in a single analysis
(*e.g.*, all *spatial* trials), the percent errors were averaged across trials. We computed RMS
performance outcomes using all available data from each trial, regardless of trial length.

Statistical analysis

Group differences in demographic and clinical characteristics were assessed with omnibus analysis of
variance (ANOVA) and Chi-squared test, as appropriate. Significant results ($\alpha = 0.05$) were followed
by post-hoc independent sample t-tests for differences between the HYA and HOA groups, and
between the HOA and MCI groups. Based on our hypotheses, we only compare the HYA and HOA
groups, and the HOA and MCI groups.

We examined differences in RMS performance across groups using separate fixed effects linear
models for each outcome variable. To test for group differences in RMS performance between the
HYA and HOA groups, and between HOA and MCI groups, we used linear models containing
indicator variables denoting the HYA and MCI groups, with HOA as the reference group (Equation

(2). Significant coefficients (Wald Tests, α = 0.05) corresponding to the HYA (β_{HYA}) or MCI
variables (β_{MCI}) would indicate significant differences in performance between the HOA group and
the HYA or MCI groups, respectively. Separate linear models were fit for each of the three gait
modification classes (*spatial, temporal, spatiotemporal*). The sample size for each model was N = 36.
Because we were not interested in comparisons between HYA and MCI, we parameterized linear
models so that the coefficients would directly test the contrasts of interest.

$$error = (\beta_0 + \beta_{HYA} \cdot I_{HYA} + \beta_{MCI} \cdot I_{MCI} + \epsilon), \qquad (2)$$

Where *error* is a 36x1 vector with one sample for each participant. The variable β_0 denotes the
average percent error of the HOA group, β_{HYA} and β_{MCI} are coefficients corresponding to the
magnitude of differences between the HYA and HOA groups and the HOA and MCI groups,
respectively. Variables I_{HYA} and I_{MCI} are indicator variables corresponding to the respective
coefficients. The variable ϵ is an error term. Linear models were fit using *fitlm* in MATLAB 2021b
(Mathworks Ltd, Natick, USA), which estimates coefficients using least-squares regression and
determines the coefficients' p-values using Wald tests. To determine whether specific types of gait
modifications were differentiated between groups, we repeated the above analysis for gait
modification subclasses (*e.g., swing, simple duple*) and individual gait modifications (*e.g., attitude-
developpe, qqqqSS*; **Figure 1**A & **Figure 2**A).

To determine which movements best distinguished between groups, we computed effects sizes
(Cohen's *d*; vs. HOA) and the average difference (Δ vs. HOA) for each gait modification class
(*spatial, temporal, spatiotemporal*). To determine whether specific gait modifications were
differentiated between groups, we repeated the above analysis for gait modification subclasses (*e.g.,
swing, simple duple*).

As an exploratory investigation to identify potential correlates of RMS performance, we computed
separate Pearson correlations between each of the clinical assessments of motor and cognitive
function against *spatial*, *temporal*, and *spatiotemporal* gait modification performance, with N=36
samples per correlation.

Results

Study compliance and participants

Data were collected in 37 participants (12 per group, plus one additional HYA participant). The
additional HYA participant was unable to follow directions when performing rhythmic movement
sequences and was excluded from analysis. One MCI participant failed to complete the protocol due
to fatigue and completed the protocol in a follow-up visit. One MCI participant was only able to
complete temporal trials due to fatigue and balance concerns. Across the 36 participants included in
analysis, only 19 trials (less than 2% of all trials) were omitted due to sensor errors not detected
during the experiment. Most participants completed the full four walkway lengths of the RMS
protocol. Across groups, 24 participants completed four walkway lengths, 10 participants completed
two-to-three lengths (5 HOA, 5 MCI), and only two participants (1 HOA, 1 MCI) completed a single
walkway length.

Demographic characteristics for each participant group are shown in Table 1. Participants exhibited
similar demographic characteristics, as expected. Clinical characteristics for each participant group
are shown in Table 2. Individuals with an MCI diagnosis trended toward worse performance than

HOA on the Trail Making Test (p = 0.099; independent-samples t-tests), cognitive (p = 0.084), and
manual (p = 0.067) TUG tests. HOA performed worse than HYA on the Reverse Corsi Blocks (p =
0.003), Body Position Spatial Task (p = 0.028), and the Four-Square Step Test (p = 0.002). All three
groups walked with similar cadences.

Table 1: Demographic characteristics of each participant group.

	HYA	**HOA**	**MCI**	**p-value**
N	12	12	12	
Age[*],[†]				< 0.01
Mean (SD)	23.9 (4.2)	67.3 (10.5)	69.5 (7.0)	
Range	18.0 - 30.0	53.0 - 80.0	57.0 - 79.0	
Height (m)				0.73
Mean (SD)	1.65 (0.09)	1.68 (0.09)	1.67 (0.10)	
Range	1.57 - 1.91	1.57 - 1.83	1.52 - 1.83	
Mass (kg)				0.08
Mean (SD)	62.7 (12.8)	72.1 (14.9)	75.2 (12.9)	
Range	50.0 - 90.1	51.9 - 95.5	58.2 - 109.1	
Sex				0.69
F	9 (75%)	8 (67%)	7 (58%)	
M	3 (25%)	4 (33%)	5 (42%)	
Years since MCI	-	-	3.2 (2.2)	-

Abbreviations: N, number of participants; HYA, healthy young adult; HOA, healthy older adult; MCI, mild cognitive
impairment; m, meters; kg, kilograms; SD, standard deviation; F, female; M, male.
[*]Significant between-group differences (α = 0.05) according to omnibus ANOVA or Chi-Squared tests, where
appropriate.
[†]Significant differences (α = 0.05) between HOA and HOA groups according to post-hoc independent-samples t-tests.
[‡]Significant differences (α = 0.05) between HYA and MCI groups according to post-hoc independent-samples t-tests.
P-values reflect omnibus ANOVAs across all three groups.
Note that we did not test for differences between the HYA and MCI groups.

Table 2: Clinical characteristics of each participant group.

	HYA	**HOA**	**MCI**	**p-value**
N	12	12	12	
Montreal Cognitive Assessment (MoCA)[a]				0.07
Mean (SD)	28.4 (2.2)	27.8 (2.1)	26.1 (2.6)	
Range	22.0 - 30.0	24.0 - 30.0	22.0 - 29.0	
Preferred cadence (strides/s)				0.34
Mean (SD)	0.92 (0.04)	0.96 (0.06)	0.95 (0.10)	
Range	0.86 - 0.98	0.98 - 1.04	0.78 - 1.14	
Trail Making Test (B-A)[c]				0.10
Mean (SD)	24.3 (13.9)	22.3 (23.2)	42.1 (29.0)	
Range	7.9 - 50.4	-28.6 - 67.2	-19.7 – 81.3	
Tower of London				0.06
Mean (SD)	20.2 (5.2)	17.3 (2.4)	15.8 (5.0)	
Range	14.0 - 29.0	13.0 - 21.0	6.0 - 25.0	

Reverse Corsi Blocks*,†				< 0.01
Mean (SD)	59.6 (21.2)	34.7 (13.6)	26.2 (14.0)	
Range	15.0 - 98.0	24.0 - 60.0	15.0 - 54.0	
Four-Square Step Test[a],*,†				< 0.01
Mean (SD)	7.3 (1.5)	9.6 (1.5)	10.8 (2.1)	
Range	5.9 - 10.9	7.6 - 13.1	8.0 - 15.4	
Body Position Spatial Task*,†				< 0.01
Mean (SD)	31.2 (15.5)	18.5 (8.9)	15.1 (6.1)	
Range	12.0 - 70.0	9.0 - 35.0	9.0 - 25.0	
Timed Up-and-Go*,†				0.02
Mean (SD)	6.2 (0.9)	7.3 (1.1)	8.4 (3.6)	
Range	4.6 - 7.6	6.0 - 9.2	5.3 - 14.9	
Manual Timed Up-and-Go[b],*				0.01
Mean (SD)	8.3 (0.6)	8.8 (1.5)	11.1 (3.6)	
Range	7.3 - 9.6	6.8 - 11.5	7.9 - 18.8	
Cognitive Timed Up-and-Go[b],*				0.02
Mean (SD)	9.3 (4.7)	11.4 (3.2)	15.2 (6.3)	
Range	6.1 - 23.4	7.2 - 19.6	8.9 - 28.4	
30-second Chair Stand				0.09
Mean (SD)	16.2 (3.0)	13.9 (3.4)	12.8 (4.6)	
Range	10.0 - 21.0	9.0 - 22.0	4.0 - 23.0	

Abbreviations: N, number of participants; HYA, healthy young adult; HOA, healthy older adult; MCI, mild cognitive
impairment; m, meters; kg, kilograms; SD, standard deviation; F, female; M, male.
*Significant between-group differences ($\alpha = 0.05$) according to omnibus ANOVA or Chi-Squared tests, where
appropriate.
†Significant differences ($\alpha = 0.05$) between HOA and HOA groups according to post-hoc independent-samples t-tests.
‡Significant differences ($\alpha = 0.05$) between HYA and MCI groups according to post-hoc independent-samples t-tests.
Superscript letters denote outcomes missing participants: [a]N=35; [b]N=34; [c]N=33.
Note that we did not test for differences between the HYA and MCI groups.

RMS performance was reduced with age

When averaging across gait modifications within each modification class (*spatial, temporal,*
spatiotemporal), regression analysis revealed that the HYA group performed *spatial* gait
modifications significantly more accurately than the HOA group (p = 0.010; $d = 1.1$, $\Delta = 4.0\%$)
(**Figure 3**). HYA participants also performed *spatiotemporal* gait modifications more accurately than
HOA participants (p = 0.048; $d = 0.8$, $\Delta = 2.1\%$). Conversely, HYA participants did not perform
temporal gait modifications significantly more accurately than HOA participants.

Figure 3: Summary of group performance on gait modifications. A) Boxplots showing group performance on spatial, temporal, and spatiotemporal trials. Each sample consists of the average percent error relative to target values across nine gait modifications for spatial and temporal trials and four gait modifications for spatiotemporal trials. B) Effect sizes of differences in performance (Cohen's *d*) between HYA and HOA groups (left) and HOA and MCI groups (right) on spatial (orange), temporal (green), and spatiotemporal (white) gait modifications. Positive values (arrows) imply that the HOA group exhibited lower error on gait modifications. We did not test for differences between the HYA and MCI groups.

Within movement subclasses (*e.g., swing, simple-duple*), differences in spatial performance between
the HYA and HOA groups were largest in modifications to the *swing* phase of gait (p = 0.006, *d* = -
1.3, Δ = 5.8%) (Figure 4A & B). Differences in gait modification performance in both the *stance*
phase (p = 0.074, *d* < 0.8, Δ = 2.2%) and the combined *swing-stance* modifications (p = 0.054, *d* <
0.9, Δ = 2.8%) had smaller effects that did not reach significance. In particular, deficits in gait
modification accuracy in HOA were largest in *swing*-phase modifications that involved large hip
flexion ranges of motion (*i.e.*, *attitude*, *developpé*, and *battement*; see modification descriptions in

Figure 1A and supplemental file *MovementLibrary.csv*), with large effect sizes (p < 0.001; *d* > 0.9; Δ
= 4.5%). These differences are exemplified by the *attitude-developpé* modification shown in **Figure**
4C.

Figure 4: A) Effect sizes of differences in performance (Cohen's *d*) between HYA and HOA
groups (top) and HOA and MCI groups (bottom) in each gait modification subclass. Spatial and
temporal modifications are denoted in orange and green, respectively. Positive values (arrows)
imply that the HOA group exhibited lower error on gait modifications compared to either group. B)
Boxplots showing distributions of gait modification accuracy (percent error vs. targets) for spatial
(top) and temporal (bottom) gait modifications. C) Results from exemplary participants for gait
modifications that elicited the largest between-group effect sizes. The *attitude-developpé*
modification (top) elicited the largest difference in spatial performance between the HYA and
HOA groups. The *simple duple* modification, *qqqqSS*, elicited the largest difference in temporal
performance between the HOA and MCI groups but did not differentiate between the HYA and
HOA groups. We did not test for differences between the HYA and MCI groups.

RMS performance was reduced in older adults with mild cognitive deficits

Unlike differences in the HYA and HOA groups, the HOA group performed only *spatiotemporal* gait
modifications significantly more accurately than the MCI group (p = 0.017; *d* = 1.3, Δ = 2.6%;
Figure 3A). While many participants with MCI exhibited reduced *spatial* and *temporal* modification
accuracy compared to the HOA group, these differences did not reach statistical significance.
Temporal accuracy was highly variable in both groups, with some participants with MCI performing

temporal modifications more accurately than those in the HOA group. However, the HOA and MCI
groups still exhibited a moderate differential effect size (p = 0.16; *d* = 0.6; Δ = 2.4%).

Differences in *spatiotemporal* modification accuracy were driven by HOA performing the *simple*
duple (p = 0.012, *d* = 1.0, Δ = 4.9%), *complex duple* (p = 0.030, *d* = 0.9, Δ = 6.5%), and *stance* (p =
0.032, *d* = 0.9, Δ =2.6%) gait modifications more accurately than MCI participants (**Figure 4A &**
B). An example of group differences in *simple duple* modification accuracy is shown in **Figure 4C**,
in which participants with MCI struggled to perform the six-step temporal sequence *qqqqSS*.
Surprisingly, the MCI group performed the *waltz* gait modifications significantly more accurately
than the HOA group (p = 0.013, *d* = 1.2, Δ = -4.0%).

Associations of motor and cognitive function with RMS performance: preliminary evidence

Exploratory correlation analyses identified non-negligible correlations in 27/29 combinations of
variables tested (Table 3; asterisks designate effect sizes with cutoff values according to Cohen
(1992)). The strongest correlations were between *spatial* modification performance and: Reverse
Corsi Blocks product score (Pearson's *r* = -0.50), Four-Square Step Test completion time (*r* = 0.55),
and Body Position Spatial Task product score (*r* = -0.55) (Table 3). In the table, asterisks denote
effect sizes for each correlation (Cohen, 1992). *Temporal* gait modifications exhibited at most
moderate correlations with clinical assessments: Reverse Corsi Blocks product score (*r* = -0.31),
Four-Square Step Test completion time (*r* = 0.29), and Body Position Spatial Task product score (*r* =
-0.28). *Spatiotemporal* gait modifications exhibited the strongest correlations with the Four-Square
Step Test completion time (*r* = 0.59), and Body Position Spatial Task product score (*r* = -0.69).

Table 3: Correlations (Pearson's *r*) between RMS performance in each of three gait modifications
and clinical outcomes.

	Gait Modification Class		
Assessment	**Spatial**	**Temporal**	**Spatio-temporal**
Trail Making Test (B-A)	0.33**	0.20*	0.30**
Tower of London	-0.35**	-0.19*	-0.49**
Reverse Corsi Blocks	-0.50***	-0.31**	-0.40**
Four-Square Step Test	0.50***	0.29*	0.59***
Body Position Spatial Task	-0.55***	-0.28*	-0.69***
Timed Up-and-Go	0.22*	0.02	0.15*
Manual Timed Up-and-Go	0.47**	0.13*	0.37**
Cognitive Timed Up-and-Go	0.42**	-0.05	0.39**
30-second Chair Stand	-0.25*	-0.05	-0.27*

*Small effect according to Cohen (1992).
**Medium effect
***Large effect

Discussion

This study leveraged biomechanical analysis and music theory to investigate the impacts of age and
cognitive function on individuals' abilities to flexibly modulate spatial and temporal aspects of their
gait. Age-related declines in motor function (HYA vs. HOA) appear to reduce individuals' capacities
to flexibly modulate spatial aspects of gait, though cognitive deficits (MCI vs. HOA) may also
contribute to reductions in this capacity. While temporal modification performance was more

variable, cognitive deficits appear to reduce individuals' abilities to modulate temporal aspects of
gait. Unsurprisingly, both age-related declines in motor and cognitive function appear to reduce
individuals' abilities to flexibly modulate spatiotemporal aspects of gait, as would be done in music
or dance therapy. By testing a broad set of spatial and temporal movement patterns, our novel
experimental framework revealed group differences in movement performance that would not be
revealed by normal walking or by a single set of gait modifications (e.g., backward walking, fast
walking). Given the importance of selecting an appropriate level of challenge during rehabilitation,
our findings suggest that both age and cognitive function, as well as movement types, should be
considered when designing complex movement, music-movement, or dance-based therapies for
individuals with MCI as others have recommended (Guadagnoli and Lee, 2004).

Motor and, possibly, cognitive deficits reduce the ability to accurately modulate spatial gait
features

Reduced performance on spatial and spatiotemporal, but not temporal, gait modifications in HOA
relative to HYA suggests that age-related declines in motor function may limit individuals' abilities
to modulate spatial components of gait. Aging is associated with reductions in strength and balance
(Schloemer et al., 2017; Reimann et al., 2020). These findings are consistent with the observed large
deficits in gait modification performance during *swing*-phase modifications in HOA relative to HYA,
which often involved large hip ranges of motion. Additionally, a post-hoc analysis of individual gait
modifications revealed that *swing* and *swing-stance* modifications comprised five of the six
modifications eliciting the largest difference in performance between HOA and HYA. Reduced *swing*
modification performance contrasts with performance noted during the stance modifications, which
challenged strength more than balance because stance modification targets could be achieved during
double-limb support. Consequently, modifications to the *swing*-phase of gait or dance steps may be
critical to personalizing exercise programs based on individuals' motor function during dance-based
therapy (Bauby and Kuo, 2000; Kuo, 2007). Note that the trend toward reduced spatial modification
performance in the MCI compared to the HOA group supported a potential effect of cognitive
function on the ability to modulate spatial components of movement (Cohen et al., 2016; Rucco et
al., 2017), which may emerge with a larger sample size.

Age-related declines in strength may impact individuals' abilities to perform plantarflexion during
stance modifications (Browne and Franz, 2018): multiple HOA and MCI participants were unable to
generate sufficient plantarflexor moments to raise their heel during single-limb support in the *relevé*
and piqué stance modifications. Plantarflexor muscle properties are altered and activation capacity is
reduced in older adults, which may limit ankle torque production capacity during gait modifications
(Morse et al., 2004; Conway and Franz, 2020). Dance-based movements that require plantarflexor
moment generation during stance may, therefore, challenge a prominent age-related deficit in HOA
and could be good targets for personalization during dance-based therapy.

Our assumption that differences in RMS performance between HYA and HOA stemmed primarily
from differences in motor function was supported by worse performance on assessments of motor
function (*e.g.*, TUG, Four-Square Step Test) in HOA compared to HYA. However, some assessments
involving aspects of cognitive function also differed between HYA and HOA (e.g., Body Position
Spatial Task) and were correlated with spatial modification performance. Therefore, it is possible that
differences in cognitive function also impact individuals' abilities to modulate spatial aspects of gait.

Cognitive deficits may reduce the ability to accurately modulate temporal gait features

We observed a moderate effect toward individuals with MCI exhibiting worse performance on
temporal gait modifications than those in the HOA group, suggesting that cognitive deficits
associated with MCI reduce the ability to flexibly modulate temporal features of gait. Such deficits
correspond to the inability to entrain stepping movements to rhythms during dance classes. For
example, individuals with MCI exhibited worse performance for six-step temporal sequences (vs. 2-
step or 4-step sequences), particularly during the *complex duple* modifications. Accurately
performing longer sequences places greater demands on working memory, which is reduced in
individuals with MCI (see *Reverse Corsi Blocks Test* in **Table 1**) (Vandierendonck et al., 2004;
Montero-Odasso et al., 2012). An alternative explanation for poor performance on some temporal
modifications is that reduced motor-cognitive integration in individuals with MCI may limit
individuals' abilities to use the musical cues to adopt the proper timing for the assigned movement
(Kluger et al., 1997; Hackney et al., 2017).

Waltz modifications represented a surprising case in which individuals with MCI performed
modifications more accurately than HOA. Specifically, individuals with MCI performed the *quick-*
Slow (*qS*) modification better than HOA (post-hoc analysis: Cohen's *d* = 2.8). *Waltz* sequences were
either two- or four-step, reducing their challenge to working memory compared to the six-step *Duple*
sequences and possibly making these modifications easier for individuals with MCI (Montero-
Odasso et al., 2012). Additionally, the three-count of *Waltz* consisted of one *quick* (*q;* 33% of the
stride) and one *Slow* (*S;* 67% of the stride) step in each measure. This division of 33% and 67% of a
stride is similar to the approximate percentages of the swing (40%) and stance (60%) phases during
normal walking, respectively(Winter, 1984). Consequently, *Waltz* may have represented a smaller
deviation from normal walking, enabling greater automaticity and reduced cortical demand than
Duple modifications (Clark, 2015). Increased automaticity, however, would only explain a lack of
deficit in *Waltz* performance in individuals with MCI. Better *Waltz* performance in MCI compared to
HOA may stem from differences in music or dance experience, which represents an interesting area
of future research.

Our decision to use fixed tempi for all participants is important for our interpretation of differences in
temporal modification performance. Using a fixed tempo, rather than setting the tempi relative to
each participant's preferred cadence (Moumdjian et al., 2019), ensured that all participants had the
same amount of time to perceive and respond to the musical cues. Personalizing the tempi to cadence
would likely reduce differences in performance due to cognitive function because individuals with
slower cadences would have more time to perceive the musical cues and determine motor commands
to execute. Rather, we selected tempi near 100 bpm, below most participants' preferred cadences.
Therefore, differences in temporal modification performance are likely driven by participants'
abilities to perceive cues and step in the correct patterns, rather than their physical capacity to match
the tempi. This expectation is supported by HYA performing *Waltz* modifications with similar
accuracy to HOA, despite stepping at a roughly 30% faster tempo. Such a fast cadence could increase
demands on motor function but did not obviously reduce HYA performance. Future studies can build
upon our current findings to evaluate the effect of varying tempi and movement speeds.

Motor and cognitive deficits reduced the ability to perform spatiotemporal modifications in
individuals with MCI

Large deficits in spatiotemporal performance in individuals with MCI compared to HOA were
expected and further emphasize the impacts of cognitive deficits on the ability to voluntarily
modulate gait biomechanics. Spatiotemporal gait modifications appear to elicit larger differences in
movement performance between HOA and MCI than spatial or temporal modifications. Motor-

cognition is adversely affected by MCI, and the requirement to modulate both spatial and temporal
parameters affected those with MCI more than HOA (Hackney et al., 2017). Additionally, reduced
set-shifting capacity in MCI compared to HOA - revealed by large differences in the A and B
components of the Trail Making Task (Bowie and Harvey, 2006) - may explain reduced
spatiotemporal performance in MCI. For example, individuals with MCI may struggle to identify
tempos in the music while remembering gait modifications. Participants were instructed to equally
prioritize the temporal or spatial components of the spatiotemporal gait modifications. However,
some participants in the MCI group reported "giving up" on either the spatial or temporal
components of the modifications. Future investigation into which aspects of RMS gait modifications
are prioritized with age and cognitive decline may provide further insight into mechanisms impacting
specific features of movement performance (Earhart, 2013).

Potential correlates of RMS performance

Our results show that *spatial*, *temporal*, and *spatiotemporal* gait modifications may be correlated
with assessments of visuospatial working memory (Reverse Corsi Blocks), motor-cognitive
integration, balance and mobility (Four-Square Step Test), and whole body spatial memory (Body
Position Spatial Task) (Dite and Temple, 2002; Vandierendonck et al., 2004; Hackney et al., 2013).
These correlations suggest potential motor and cognitive contributions to spatial and temporal aspects
of RMS performance and may be useful as design variables for dance-based therapy classes.
However, these findings should be considered with extreme caution because in this preliminary
study, we were not able to rigorously validate the robustness of these correlations, which will be a
valuable focus of future larger-sample studies. Our findings here can pilot future trials with larger
sample sizes and specific hypotheses about which aspects of motor or cognitive function impact
RMS performance.

Clinical implications

Beyond advancing our understanding of how age and cognition impact the ability to flexibly
modulate gait, our findings have implications for dance-based therapies for individuals with MCI.
Our results showing distinct motor control deficits related to aging versus cognitive impairment
inform clinical understanding of the mechanisms underlying impaired gait performance in HOA and
MCI. Creative dance-based movement therapies aim to challenge both motor and cognitive function
and have shown positive impacts on spatial cognition, executive function, balance, and mobility in
individuals with Parkinson's and MCI (Hackney et al., 2007a; Hackney et al., 2007b; McKee and
Hackney, 2013; Zhu et al., 2020). Our results show that different temporal features of music and
spatial components of movements challenge distinct motor (e.g., strength or balance) and cognitive
(e.g., working memory, set-shifting, or motor-cognitive integration) features (Thom and Clare, 2011;
Montero-Odasso et al., 2012; Earhart, 2013; Browne and Franz, 2018). Consequently, personalizing
music and dance selections during therapy based on an individual's motor and cognitive function
may maximize therapeutic effects. Specifically, selecting music and dance parameters that optimize
the level of challenge for an individual may maximize benefits while retaining motivation
(Guadagnoli and Lee, 2004; Hackney et al., 2007a). For example, our results suggest that teaching
spatial or temporal modifications alone may be an appropriate level of challenge for individuals with
MCI, before progressing to spatiotemporal modifications (Hackney and Earhart, 2010). However, we
evaluated only the immediate relationship between age-related declines in motor and cognitive
function and RMS performance. With repeated practice, individuals with MCI may improve RMS
performance levels to near those of HOA or HYA participants (McKee and Hackney, 2013). These
learning or longitudinal effects are beyond the scope of this study but are interesting areas of future

research that could inform personalized design and data-driven progression of dance-based and
music-based movement therapies.

Limitations

The following limitations constrain the generalizability of our findings. First, our modest sample size
of twelve participants per group lacked the statistical power to identify all differences in movement
performance, particularly during temporal gait modifications. Nonetheless, our data revealed
differences in movement performance for spatial and spatiotemporal gait modifications, and showed
moderate differential effect sizes (HOA vs. MCI) during temporal modification performance.
Secondly, while protocol administrators verified that participants understood each gait modification
similarly before collecting data, understanding may have improved over the first few trials,
potentially improving performance in later trials. However, we used a block randomization strategy
to account for changes in comprehension of the movements.

Our findings should be viewed in the context of a broader set of constructs – including and beyond
the motor and cognitive constraints studied here – that could impact movement performance during
movement therapy. The specific gait modifications selected, sensory stimuli (music type or tempo).
walking speed, or even individuals' relationships to music and dance could impact RMS
performance. For example, music with an easier-to-perceive downbeat may alter the observed
relationships between cognitive function and spatiotemporal RMS performance in individuals with
lesser music experience. Our group-level analyses across diverse movement classes provide an early
step toward understanding how dance-based RMS challenge specific aspects of motor and cognitive
function. Larger sample sizes would enable the discovery of specific factors underlying RMS
performance in individuals with MCI to be discovered from a broader set of constructs affecting
movement.

Conclusions

We developed and implemented an innovative RMS gait modifications protocol to evaluate the
impacts of age-related declines in motor and cognitive function on the ability to accurately modulate
spatial and temporal features of gait. By improving our understanding of motor and cognitive impacts
on the ability to modulate gait, this study represents an early step toward systematically personalized
dance-based movement therapies. Furthermore, this study lays the foundation for future evaluation of
the ability of RMS to 1) help diagnose cognitive-motor impairments by providing insight into
mechanisms of movement performance deficits and 2) track intervention efficacy by quantifying
changes in the accuracy of spatial and temporal features of gait modifications following therapy.
Future studies identifying clinical and mechanistic predictors of RMS performance will advance our
ability to provide quantitative metrics that assist clinicians in selecting movement therapy parameters
to maximize treatment efficacy.

Conflict of Interest

563 *The authors declare that the research was conducted in the absence of any commercial or financial*
564 *relationships that could be construed as a potential conflict of interest.*

Author Contributions

MCR, MEH, JLM, LE, and TMK obtained funding for the study. MEH, TMK, LE, MCR, and AS
defined the study design and data collection protocol. MCR, AS, and KC were skilled assessors who

performed data collection. MEH, JLM, TK, LE, and MCR supervised the study and planned data
analysis. MCR and JLM performed data analysis. MCR drafted the original manuscript. All authors
provided intellectual critique and manuscript revisions.

Funding

Research reported in this manuscript was supported by the National Institute of Child Health and
Human Development and the National Institute on Aging of the National Institutes of Health under
award numbers F32HD108927 and R01AG062691, respectively. This research was also supported
by Emory University through a Goizueta Alzheimer's Disease Research Center CEP Innovation
Accelerator Seed Grant, an award from the Senior Vice President of Research at the Intersection
Fund, and a Synergy II/Nexus Award.

Acknowledgments

We thank the participation of our volunteers, and T. Prusin and C. Carroll-Sauer for their assistance
in data collection and participant recruitment.

Data Availability Statement

The original contributions presented in the study are included in the supplementary material, further
inquiries can be directed to the corresponding author.

References

Bauby, C.E., and Kuo, A.D. (2000). Active control of lateral balance in human walking. *Journal of*
586   *biomechanics* 33(11)**,** 1433-1440.

587   Booth, F.W., Weeden, S.H., and Tseng, B.S. (1994). Effect of aging on human skeletal muscle and
588   motor function. *Medicine and science in sports and exercise* 26(5)**,** 556-560.

589   Bowie, C.R., and Harvey, P.D. (2006). Administration and interpretation of the Trail Making Test.
590   *Nature protocols* 1(5)**,** 2277-2281.

591   Browne, M.G., and Franz, J.R. (2018). More push from your push-off: Joint-level modifications to
592   modulate propulsive forces in old age. *PLoS One* 13(8)**,** e0201407.

593   Clark, D.J. (2015). Automaticity of walking: functional significance, mechanisms, measurement and
594   rehabilitation strategies. *Frontiers in human neuroscience* 9**,** 246.

595   Cohen, J. (1992). A power primer. *Psychological bulletin* 112(1)**,** 155.

596   Cohen, J.A., Verghese, J., and Zwerling, J.L. (2016). Cognition and gait in older people. *Maturitas*
597   93**,** 73-77.

598   Conway, K.A., and Franz, J.R. (2020). Shorter gastrocnemius fascicle lengths in older adults
599   associate with worse capacity to enhance push-off intensity in walking. *Gait & posture* 77**,**
600   89-94.

601   Dite, W., and Temple, V.A. (2002). A clinical test of stepping and change of direction to identify
602   multiple falling older adults. *Archives of physical medicine and rehabilitation* 83(11)**,** 1566-
603   1571.

604   Earhart, G.M. (2013). Dynamic control of posture across locomotor tasks. *Movement disorders*
605   28(11)**,** 1501-1508.

606   Gauthier, S., Reisberg, B., Zaudig, M., Petersen, R.C., Ritchie, K., Broich, K., et al. (2006). Mild
607   cognitive impairment. *The Lancet* 367(9518)**,** 1262-1270. doi: https://doi.org/10.1016/S0140-
608   6736(06)68542-5.

609   Geda, Y.E., Roberts, R.O., Knopman, D.S., Christianson, T.J., Pankratz, V.S., Ivnik, R.J., et al.
610   (2010). Physical exercise, aging, and mild cognitive impairment: a population-based study.
611   *Archives of neurology* 67(1)**,** 80-86.

612   Grahn, J.A. (2012). Neural mechanisms of rhythm perception: current findings and future
613   perspectives. *Topics in cognitive science* 4(4)**,** 585-606.

614   Guadagnoli, M.A., and Lee, T.D. (2004). Challenge point: a framework for conceptualizing the
615   effects of various practice conditions in motor learning. *Journal of motor behavior* 36(2)**,**
616   212-224.

617   Hackney, M.E., and Earhart, G.M. (2009). Effects of dance on movement control in Parkinson's
618   disease: a comparison of Argentine tango and American ballroom. *J Rehabil Med* 41(6)**,** 475-
619   481. doi: 10.2340/16501977-0362.

620   Hackney, M.E., and Earhart, G.M. (2010). Recommendations for implementing tango classes for
621   persons with Parkinson disease. *American Journal of Dance Therapy* 32(1)**,** 41-52.

622   Hackney, M.E., Hall, C.D., Echt, K.V., and Wolf, S.L. (2013). Dancing for balance: feasibility and
623   efficacy in oldest-old adults with visual impairment. *Nursing research* 62(2)**,** 138-143.

Hackney, M.E., Kantorovich, S., and Earhart, G.M. (2007a). A study on the effects of Argentine
tango as a form of partnered dance for those with Parkinson disease and the healthy elderly.
American Journal of Dance Therapy 29(2)**,** 109-127.

Hackney, M.E., Kantorovich, S., Levin, R., and Earhart, G.M. (2007b). Effects of tango on functional
mobility in Parkinson's disease: a preliminary study. *Journal of neurologic physical therapy*
31(4)**,** 173-179.

Hackney, M.E., Nocera, J., Creel, T., Riebesell, M.D., and Kesar, T. (2017). "Exercise and balance in
older adults with movement disorders," in *Locomotion and Posture in Older Adults*.
Springer), 323-346.

Hortobágyi, T., Rider, P., Gruber, A.H., and DeVita, P. (2016). Age and muscle strength mediate the
age-related biomechanical plasticity of gait. *European journal of applied physiology* 116(4)**,**
805-814.

Ivanenko, Y.P., Cappellini, G., Dominici, N., Poppele, R.E., and Lacquaniti, F. (2005). Coordination
of locomotion with voluntary movements in humans. *Journal of Neuroscience* 25(31)**,** 7238-
7253.

Jones, C.J., Rikli, R.E., and Beam, W.C. (1999). A 30-s chair-stand test as a measure of lower body
strength in community-residing older adults. *Research quarterly for exercise and sport* 70(2)**,**
113-119.

Kluger, A., Gianutsos, J.G., Golomb, J., Ferris, S.H., George, A.E., Franssen, E., et al. (1997).
Patterns of motor impairment in normal aging, mild cognitive decline, and early
Alzheimer'Disease. *The Journals of Gerontology Series B: Psychological Sciences and Social*
Sciences 52(1)**,** P28-P39.

Kornysheva, K., and Diedrichsen, J. (2014). Human premotor areas parse sequences into their spatial
and temporal features. *Elife* 3**,** e03043.

Kuo, A.D. (2007). The six determinants of gait and the inverted pendulum analogy: A dynamic
walking perspective. *Human movement science* 26(4)**,** 617-656.

Lazarou, I., Parastatidis, T., Tsolaki, A., Gkioka, M., Karakostas, A., Douka, S., et al. (2017).
International ballroom dancing against neurodegeneration: a randomized controlled trial in
Greek community-dwelling elders with mild cognitive impairment. *American Journal of*
Alzheimer's Disease & Other Dementias® 32(8)**,** 489-499.

Lundin-Olsson, L., Nyberg, L., and Gustafson, Y. (1998). Attention, frailty, and falls: the effect of a
manual task on basic mobility. *Journal of the American Geriatrics Society* 46(6)**,** 758-761.

Mancini, M., and Horak, F.B. (2016). Potential of APDM mobility lab for the monitoring of the
progression of Parkinson's disease. *Expert review of medical devices* 13(5)**,** 455-462.

McKee, K.E., and Hackney, M.E. (2013). The effects of adapted tango on spatial cognition and
disease severity in Parkinson's disease. *J Mot Behav* 45(6)**,** 519-529. doi:
10.1080/00222895.2013.834288.

Montero-Odasso, M., Muir, S.W., and Speechley, M. (2012). Dual-task complexity affects gait in
people with mild cognitive impairment: the interplay between gait variability, dual tasking,
and risk of falls. *Arch Phys Med Rehabil* 93(2)**,** 293-299. doi: 10.1016/j.apmr.2011.08.026.

Morris, S., Morris, M.E., and Iansek, R. (2001). Reliability of measurements obtained with the Timed
"Up & Go" test in people with Parkinson disease. *Physical therapy* 81(2)**,** 810-818.

Morse, C.I., Thom, J.M., Davis, M.G., Fox, K.R., Birch, K.M., and Narici, M.V. (2004). Reduced
plantarflexor specific torque in the elderly is associated with a lower activation capacity.
European journal of applied physiology 92(1)**,** 219-226.

Moumdjian, L., Moens, B., Maes, P.-J., Van Nieuwenhoven, J., Van Wijmeersch, B., Leman, M., et
al. (2019). Walking to music and metronome at various tempi in persons with multiple
sclerosis: a basis for rehabilitation. *Neurorehabilitation and neural repair* 33(6)**,** 464-475.

Mueller, S.G., Weiner, M.W., Thal, L.J., Petersen, R.C., Jack, C., Jagust, W., et al. (2005). The
Alzheimer's disease neuroimaging initiative. *Neuroimaging Clinics* 15(4)**,** 869-877.

Nasreddine, Z.S., Phillips, N.A., Bédirian, V., Charbonneau, S., Whitehead, V., Collin, I., et al.
(2005). The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive
impairment. *Journal of the American Geriatrics Society* 53(4)**,** 695-699.

Persad, C.C., Jones, J.L., Ashton-Miller, J.A., Alexander, N.B., and Giordani, B. (2008). Executive
function and gait in older adults with cognitive impairment. *J Gerontol A Biol Sci Med Sci*
63(12)**,** 1350-1355. doi: 10.1093/gerona/63.12.1350.

Phillips, L.H., Wynn, V., McPherson, S., and Gilhooly, K. (2001). Mental planning and the Tower of
London task. *The Quarterly Journal of Experimental Psychology Section A* 54(2)**,** 579-597.

Rallis, I., Doulamis, N., Doulamis, A., Voulodimos, A., and Vescoukis, V. (2018). Spatio-temporal
summarization of dance choreographies. *Computers & Graphics* 73**,** 88-101.

Reimann, H., Ramadan, R., Fettrow, T., Hafer, J.F., Geyer, H., and Jeka, J.J. (2020). Interactions
between different age-related factors affecting balance control in walking. *Frontiers in Sports*
and Active Living 2**,** 94.

Rucco, R., Agosti, V., Jacini, F., Sorrentino, P., Varriale, P., De Stefano, M., et al. (2017). Spatio-
temporal and kinematic gait analysis in patients with Frontotemporal dementia and
Alzheimer's disease through 3D motion capture. *Gait & posture* 52**,** 312-317.

Sanford, A.M. (2017). Mild Cognitive Impairment. *Clinics in Geriatric Medicine* 33(3)**,** 325-337.
doi: https://doi.org/10.1016/j.cger.2017.02.005.

Schloemer, S.A., Thompson, J.A., Silder, A., Thelen, D.G., and Siston, R.A. (2017). Age-Related
Differences in Gait Kinematics, Kinetics, and Muscle Function: A Principal Component
Analysis. *Annals of Biomedical Engineering* 45(3)**,** 695-710. doi: 10.1007/s10439-016-1713-
4.

Styns, F., van Noorden, L., Moelants, D., and Leman, M. (2007). Walking on music. *Human*
movement science 26(5)**,** 769-785.

Thom, J.M., and Clare, L. (2011). Rationale for combined exercise and cognition-focused
interventions to improve functional independence in people with dementia. *Gerontology*
57(3)**,** 265-275.

Vandierendonck, A., Kemps, E., Fastame, M.C., and Szmalec, A. (2004). Working memory
components of the Corsi blocks task. *British journal of psychology* 95(1)**,** 57-79.

Washabaugh, E.P., Kalyanaraman, T., Adamczyk, P.G., Claflin, E.S., and Krishnan, C. (2017).
Validity and repeatability of inertial measurement units for measuring gait parameters. *Gait &*
posture 55**,** 87-93.

Wilson, M., and Kwon, Y.-H. (2008). The role of biomechanics in understanding dance movement: a
review. *Journal of Dance Medicine & Science* 12(3)**,** 109-116.

Winter, D.A. (1984). Kinematic and kinetic patterns in human gait: variability and compensating
effects. *Human movement science* 3(1-2)**,** 51-76.

Zhu, Y., Wu, H., Qi, M., Wang, S., Zhang, Q., Zhou, L., et al. (2018). Effects of a specially designed
aerobic dance routine on mild cognitive impairment. *Clinical interventions in aging* 13**,** 1691.

Zhu, Y., Zhong, Q., Ji, J., Ma, J., Wu, H., Gao, Y., et al. (2020). Effects of aerobic dance on
cognition in older adults with mild cognitive impairment: a systematic review and meta-
analysis. *Journal of Alzheimer's Disease* 74(2)**,** 679-690.