Title: Clomipramine inhibits dynamin GTPase activity by L-α-phosphatidyl-L-serine stimulation

Authors: Hiroshi Miyoshi[1*&], Masahiro Otomo[2§&] and Kiyofumi Takahashi[2#&]

Affiliations: Department of [1]Microbiology and [2]Neuropsychiatry, St. Marianna University School of Medicine, 2-16-1 Sugao, Miyamae, Kawasaki 216-8511, Japan.

[*]Corresponding author: Department of Microbiology, St. Marianna University School of Medicine, 2-16-1 Sugao, Miyamae, Kawasaki 216-8511, Japan; Tel. +81 44 9778111; E-mail: hmiyoshi@marianna-u.ac.jp

[§] Present address: Shinkohkai izukan-nami hospital, 1694 Hirai, Kan-nami, Tagata, Shizuoka 419-0107 Japan; E-mail: otomo@shinkohkai.jp

[#] Present address: Research Organization for Nano and Life Innovation, Waseda University, Tokyo, Japan; E-mail: kiyofu-min@umin.org

[&] All authors contributed equally.

Abstract

Three dynamin isoforms play critical roles in clathrin-dependent endocytosis. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) enters host cells *via* clathrin-dependent endocytosis. We previously reported that 3-(9-chloro-5,6-dihydrobenzo[b][1]benzazepin-11-yl)-N,N-dimethylpropan-1-amine (clomipramine) inhibits the GTPase activity of dynamin 1, which is in mainly neuron. Therefore, we investigated whether clomipramine inhibits the activity of other dynamin isoforms in this study. We found that, similar to its inhibitory effect on dynamin 1, clomipramine inhibited the L-α-phosphatidyl-L-serine-stimulated GTPase activity of dynamin 2, which is expressed ubiquitously, and dynamin 3, which is expressed in the lung. Inhibition of GTPase activity raises the possibility that clomipramine can suppress SARS-CoV-2 entry into host cells.

Keywords: clomipramine, SARS-CoV-2, dynamin, endocytosis, inhibition

Abbreviations

severe acute respiratory syndrome coronavirus 2, SARS-CoV-2; dynamin, Dyn; myristyl trimethyl ammonium bromide, MiTMAB; L-α-phosphatidyl-L-serine, PS; angiotensin-converting enzyme 2, ACE2; histidine, His; orthophosphate, Pi.

Introduction

Dynamin (Dyn) GTPase plays a critical role in clathrin- and caveolae-dependent endocytosis [1-3]. Mammals express three Dyn isoforms. Although Dyn 2 is expressed ubiquitously, Dyn 1 is assumed to be neuron specific. Dyn 3 is expressed mainly in the testis, brain and lung [4].

The essential function of Dyn GTPase activity is production of mechanical force to drive membrane fission during vesicle budding during endocytosis [1, 5]. Notably, endocytosis driven by Dyn GTPase activity is involved in the intracellular transportation of transferrin [6].

All Dyn isoforms carry four functional domains: a GTPase domain, pleckstrin homology domain, GTPase effector domain, and proline/arginine-rich domain [1]. These functional domains act in concert during vesicle budding [7], and lipid species such as L-α-phosphatidyl-L-serine (PS) potentiate dynamin activity [8]. Dyn GTPase activities are inhibited by cationic surface-active compounds such as myristyl trimethyl ammonium bromide (MiTMAB). MiTMAB competes with PS for binding to the pleckstrin homology domain in Dyn, but it does not compete with GTP to bind Dyn [9].

Cells can internalize many resources from the external cell environment *via* endocytosis, and non-enveloped viruses (e.g., hepatitis C virus) generally leverage related mechanisms to enter a cell following the same endocytic pathway as resources [10]. Many enveloped viruses, such as

severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), enter cells *via* endocytosis. The

SARS-CoV-2 spike glycoprotein binds to host angiotensin-converting enzyme 2 (ACE2), thereby

mediating membrane fusion and enabling virus entry into cells [11, 12].

Recently, Kato *et al.* reported that the tricyclic antidepressant clomipramine suppressed

ACE2-mediated SARS-CoV-2 entry into host cells [13]. Clomipramine has been used in basic

research as a reagent to inhibit endocytosis [14].

We previously reported that several antipsychotic treatments inhibited the GTPase

activity of Dyn 1 and Dyn 2 and that the suppression of dynamin GTPase activity by those

antipsychotic drugs led to inhibition of dynamin-driven endocytosis. Moreover, we explained that

(1*S*)-*cis*-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-*N*-methyl-1-naphthalenamine (sertraline) is a

competitive inhibitor of dynamin I GTPase with respect to both GTP and PS. [15, 16]. In a

previous study, we also reported that clomipramine inhibited the GTPase activity of neuron-

specific Dyn 1 [15]. However, the inhibition of the activity of lung-specific Dyn 3 and activity of

ubiquitous Dyn 2 by clomipramine has not yet been investigated. Therefore, the inhibition of Dyn

isoform GTPase activity by clomipramine was measured to understand the inhibitory mechanism.

Materials and Methods

Materials

The following materials were obtained from the cited sources and used in this study: Pfu Ultra II Fusion HS DNA Polymerase (Agilent Technologies); restriction enzymes and DNA ligation kit ver.2 (Takara Bio); pET21a and *E. coli* Rosetta2 (DE3) (Merck); TALON Metal Affinity Resin (Clontech); Mono Q 5/50 GL (GE Healthcare); L-α-phosphatidyl-L-serine (Sigma Aldrich); clomipramine (Sigma); and HRP–conjugated anti Penta-His antibody (Qiagen). Other chemicals used in this study were of an analytical grade or higher. Clomipramine, sertraline and MiTMAB were prepared in stock solutions consisting of 50% (v/v) DMSO and 30 mM Tris-HCl pH 7.4 and diluted in 30 mM Tris-HCl, pH 7.4, prior to use in the assay.

Plasmid construction, protein expression and purification of Dyn isoforms

All Dyn isoform genes were amplified by PCR as described previously [15, 16]. These Dyn genes were cloned into separate pET21a plasmids to create C-terminal-His$_6$-tagged Dyn isoform protein (Dyns-His$_6$) vectors. We expressed Dyns-His$_6$ from *E. coli* Rosetta2 (DE3) by transforming with those Dyn isoforms-coding pET21a expression plasmids. The expressed Dyns-His$_6$ were purified to homogeneity on a MonoQ column after TALON Metal Affinity Resin purification, as previously reported [15-17]. The purified proteins were confirmed by 15 %

SDS-PAGE.

Preparation of PS-liposomes and performance of a GTPase assay with purified Dyns-His6

Forty microlitres of PS (10 mg ml^{-1}) in chloroform/methanol (95:5) was evaporated to a final

5-µl volume, and then PS was suspended in 1 ml of 30 mM Tris-HCl, pH 7.4, and sonicated for 2

min on ice to yield a working solution of 400 µg ml^{-1}. GTPase assays with the purified Dyns-His$_6$

were performed via sensitive malachite green colorimetric detection of orthophosphate (Pi), as

described previously [15-17]. Phosphate release was quantified and the levels were compared

against a standard curve of sodium dihydrogen orthophosphate monohydrate after each

experiment. KaleidaGraph 4.0 (Synergy Software) was used to plot and analyse curves using non-

linear regression.

Blue native (BN)-PAGE

BN-PAGE was performed as previously reported [17]. Purified Dyn2-His$_6$ (2 µM) were

incubated at room temperature for 30 min with 2 mM PS-liposomes in TBS, 2 mM PS-liposomes

+ 1% SDS in TBS, 1 mM clomipramine + 2 mM PS-liposomes in TBS, or 1 mM clomipramine

+ 2 mM PS-liposomes + 1% SDS in TBS, respectively. The incubation solutions were

supplemented with BN-PAGE sample buffer (5% Coomassie Brilliant Blue G-250; 500 mM 6-

aminocaproic acid; 100 mM Bis–Tris-HCl, pH 7.0; and 1 mM phenylmethylsulfonyl fluoride)

and incubated on ice for 5 min. All samples were resolved in 4 to 16% gels by BN-PAGE and

measured by western blotting with HRP–conjugated anti-Penta-His antibody.

Results

Dyns-His6 was activated by PS-liposomes

We expressed and purified Dyns-His$_6$. Those purified Dyns-His$_6$ were confirmed by 15 % SDS-PAGE as shown in Fig. 1. To confirm the GTPase activities of purified Dyns-His$_6$, malachite green colorimetric detection of released orthophosphate was performed to confirm the GTPase activity levels of purified Dyns-His$_6$ with/without PS-liposome treatment. Purified Dyns-His$_6$ at all concentrations hydrolysed GTP to release Pi in a time-dependent manner, and the GTPase activity was stimulated by the PS-liposomes (Fig. 2). The addition of PS-liposomes increased Dyn1-His$_6$ activity by 5.1-fold, Dyn2-His$_6$ activity by 4.7-fold, and Dyn3-His$_6$ activity by 5.7-fold (Fig. 2 A versus B, 60 min).

Clomipramine inhibits GTPase activity of all dynamin isoforms in the presence of PS-liposomes

GTPase assay was performed to evaluate the inhibition of Dyns-His$_6$ GTPase activity by clomipramine, and the activity levels were compared with those of sertraline and MiTMAB (Fig. 3 and Fig. 4). In the presence of PS-liposomes, clomipramine showed significant inhibition of GTPase activity (Fig.3). The IC$_{50}$ values for Dyn1-His$_6$ were 5.5 ± 1.2 μM (sertraline), 8.1 ± 3.7 μM (MiTMAB) and 22.0 ± 2.2 μM μM (clomipramine) in the presence of PS-liposomes (Fig.

3A). For Dyn2-His$_6$, the IC$_{50}$ values were 4.9 ± 0.6 μM (sertraline), 9.2 ± 2.5 μM (MiTMAB) and

12.8 ± 3.2 μM (clomipramine) in the presence of PS-liposomes (Fig. 3B). The IC$_{50}$ values for

Dyn3-His$_6$ were 4.4 ± 0.5 μM (sertraline), 7.8 ± 3.0 μM (MiTMAB) and 11.3 ± 2.9

(clomipramine) in the presence of PS-liposomes (Fig. 3C). In contrast to MiTMAB, clomipramine

showed no overt inhibition of GTPase activity, similar to sertraline, in the absence of PS-

liposomes (Fig. 4).

Clomipramine did not inhibit the oligomer formation of Dyn2-His$_6$

Dyn GTPase activity is increased by its adoption of helical oligomerization [18]. BN-PAGE

was performed to analyse effect of clomipramine for oligomerization of Dyn2-His$_6$. Several bands

corresponding to Dyn2-His$_6$ oligomers were observed at approximately 300 kDa (Arrow II) and

approximately 600 kDa (Arrow III) after BN-PAGE of Dyn2-His$_6$ treated with/without

clomipramine (Fig. 5 Lanes 1 and 3). Moreover, those lanes showed similar band profiles in the

higher molecular weight area than at Arrow III (Fig. 5 Lanes 1 versus 3). In the presence of 1%

SDS, a denaturant, a single band of approximately 140 kDa, corresponding to the Dyn2-His$_6$

monomer, was observed for Dyn2-His$_6$ with or without clomipramine treatment, (Fig. 5 Lanes 2

and 4, Arrow I).

Discussion

PS-liposomes had been previously reported to stimulate the GTPase activity of Dyn [19]. As shown in Fig. 2, PS-liposomes stimulated all purified Dyns-His$_6$ GTPase activities in this study. This meant that these purified Dyns-His$_6$ can be used to assess the inhibition by clomipramine with GTPase assays. In fact, The IC$_{50}$ values of inhibitors (sertraline, MiTMAB and clomipramine) for Dyn1-His$_6$ obtained from Fig. 3A were not significantly different from those obtained in our previous study [15].

Clomipramine exhibited inhibition of Dyn2 and Dyn3 GTPase activity similar to its effect on Dyn1 GTPase activity in the presence of PS-liposomes (Fig. 3). Although Dyns-His$_6$ GTPase activity in the presence of PS-liposomes was completely lost after 320 μM MiTMAB treatment, Dyns-His$_6$ GTPase activity in the addition of clomipramine and sertraline (320 μM) was incompletely lost (Fig. 3A-C). The residual activity is thought to result from GTPase activity in the absence of phospholipids, that is, the so-called basal intrinsic GTPase activity. In fact, these observations agree with the observation that clomipramine did not strongly inhibit basal intrinsic GTPase activity (Fig. 4A-C). Basal intrinsic GTPase activity tends to decrease after disassembly of the dynamin oligomer [20]. Therefore, it is possible that clomipramine does not affect oligomerization of Dyns-His$_6$. Accordingly, we performed BN-PAGE to examine whether clomipramine affects the stabilization of the Dyn2-His$_6$ oligomer. The addition of clomipramine

did not affect any bands corresponding to Dyns-His$_6$ oligomers (Fig. 5, Lane 1 versus Lane 3).

Taken together, these results demonstrate that inhibition of Dyns-His6 activity did not result from

by clomipramine-induced destabilization of the oligomer. It is known that pleckstrin homology

domain on dynamin binds to lipids [21]. Consequently, clomipramine appears to directly interact

pleckstrin homology domain, so that appears to effect interaction between PS-liposome and

dynamin.

Dyn plays a critical role in clathrin-dependent endocytosis, and SARS-CoV-2 enter into host

cells *via* clathrin-dependent endocytosis [22]. Kato *et al.* mentioned that the tricyclic

antidepressant clomipramine suppressed ACE2-mediated SARS-CoV-2 entry [13, 23]. In this

study, we found that clomipramine inhibited the PS-liposome-stimulated GTPase activity of Dyn

2, which was ubiquitously expressed, and Dyn 3, which was expressed in the lung [4]. Taken

together, inhibition of PS-liposome-stimulated GTPase activity by clomipramine may participate

in suppression of ACE2-mediated SARS-CoV-2 entry.

Kato *et al.* also mentioned that the inhibition mechanism by which clomipramine inhibited

the entry of SARS-CoV-2 differed from that of other antiviral drugs, such as remdesivir, and

suggested the potential for the concomitant use of clomipramine and drugs with other mechanistic

effects for treating COVID-19 patients [23]. Our results reinforce the potential for the

concomitant use of these drugs.

Acknowledgments

This work was supported in part by Grants-in-Aid for Scientific Research from the Ministry of

Education, Science, Sports and Culture of Japan to KT (21791152) and HM (25430118).

Author contributions

HM contributed to data analysis and manuscript writing. MO and KT contributed to the

collection of data and data analysis. All authors have read and approved the final manuscript.

Data availability

The data underlying this article are available in the article.

Conflict of interest

The authors declare that they have no conflict of interest.

References

[1] Moreno-Layseca, P. *et al.* (2021). Cargo-specific recruitment in clathrin- and dynamin-independent endocytosis. *Nature Cell Biology* **23**, 1073-1084.

[2] Weng, L. *et al.* (2014). Regulation of cargo-selective endocytosis by dynamin 2 GTPase-activating protein girdin. *EMBO J* **33**, 2098-2112.

[3] Henley, J.R., Krueger, E.W., Oswald, B.J. and McNiven, M.A. (1998). Dynamin-mediated internalization of caveolae. *J Cell Biol* **141**, 85-99.

[4] Reis, C.R., Chen, P.H., Srinivasan, S., Aguet, F., Mettlen, M. and Schmid, S.L. (2015). Crosstalk between Akt/GSK3β signaling and dynamin-1 regulates clathrin-mediated endocytosis. *EMBO J* **34**, 2132-46.

[5] Ramachandran, R. and Schmid, S.L. (2018). The dynamin superfamily. Current Biology 28, R411-R416.

[6] van Dam, E.M. and Stoorvogel, W. (2002). Dynamin-dependent transferrin receptor recycling by endosome-derived clathrin-coated vesicles. *Mol Biol Cell* **13**, 169-82.

[7] Muhlberg, A.B., Warnock, D.E. and Schmid, S.L. (1997). Domain structure and intramolecular regulation of dynamin GTPase. *EMBO J* **16**, 6676-83.

[8] Bashkirov, P.V., Akimov, S.A., Evseev, A.I., Schmid, S.L., Zimmerberg, J. and Frolov, V.A. (2008). GTPase Cycle of Dynamin Is Coupled to Membrane Squeeze and Release, Leading

to Spontaneous Fission. *Cell* **135**, 1276-1286.

[9] Quan, A. *et al.* (2007). Myristyl Trimethyl Ammonium Bromide and Octadecyl Trimethyl

Ammonium Bromide Are Surface-Active Small Molecule Dynamin Inhibitors that Block

Endocytosis Mediated by Dynamin I or Dynamin II. *Molecular Pharmacology* **72**, 1425-

1439.

[10] Helle, F. and Dubuisson, J. (2007). Hepatitis C virus entry into host cells. *Cellular and*

Molecular Life Sciences **65**, 100.

[11] Li, W. *et al.* (2003). Angiotensin-converting enzyme 2 is a functional receptor for the SARS

coronavirus. *Nature* **426**, 450-4.

[12] Hoffmann, M. *et al.* (2020). SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and

Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* **181**, 271-280 e8.

[13] Kato, Y. *et al.* (2021). Clomipramine suppresses ACE2-mediated SARS-CoV-2 entry.

bioRxiv, 2021.03.13.435221.

[14] Vercauteren, D., Vandenbroucke, R.E., Jones, A.T., Rejman, J., Demeester, J., De Smedt,

S.C., Sanders, N.N. and Braeckmans, K. (2010). The use of inhibitors to study endocytic

pathways of gene carriers: optimization and pitfalls. *Mol Ther* **18**, 561-9.

[15] Otomo, M., Takahashi, K., Miyoshi, H., Osada, K., Nakashima, H. and Yamaguchi, N. (2008).

Some Selective Serotonin Reuptake Inhibitors Inhibit Dynamin I Guanosine Triphosphatase

(GTPase). *Biological and Pharmaceutical Bulletin* **31**, 1489-1495.

[16] Takahashi, K., Miyoshi, H., Otomo, M., Osada, K., Yamaguchi, N. and Nakashima, H. (2010). Suppression of dynamin GTPase activity by sertraline leads to inhibition of dynamin-dependent endocytosis. *Biochem Biophys Res Commun* **391**, 382-7.

[17] Takahashi, K., Otomo, M., Yamaguchi, N., Nakashima, H. and Miyoshi, H. (2012). Replacement of Arg-386 with Gly in Dynamin 1 Middle Domain Reduced GTPase Activity and Oligomer Stability in the Absence of Lipids. *Bioscience, Biotechnology, and Biochemistry* **76**, 2195-2200.

[18] Stowell, M.H.B., Marks, B., Wigge, P. and McMahon, H.T. (1999). Nucleotide-dependent conformational changes in dynamin: evidence for a mechanochemical molecular spring. *Nature Cell Biology* **1**, 27-32.

[19] Tuma, P.L., Stachniak, M.C. and Collins, C.A. (1993). Activation of dynamin GTPase by acidic phospholipids and endogenous rat brain vesicles. *Journal of Biological Chemistry* **268**, 17240-17246.

[20] Warnock, D.E., Baba, T. and Schmid, S.L. (1997). Ubiquitously expressed dynamin-II has a higher intrinsic GTPase activity and a greater propensity for self-assembly than neuronal dynamin-I. *Mol Biol Cell* **8**, 2553-2562.

[21] Vallis, Y., Wigge, P., Marks, B., Evans, P.R. and McMahon, H.T. (1999). Importance of the

pleckstrin homology domain of dynamin in clathrin-mediated endocytosis. *Current Biology* **9**, 257-263.

[22] Yang, H., Yuan, H., Zhao, X., Xun, M., Guo, S., Wang, N., Liu, B. and Wang, H. (2022). Cytoplasmic domain and enzymatic activity of ACE2 are not required for PI4KB dependent endocytosis entry of SARS-CoV-2 into host cells. *Virologica Sinica 37*, 380-389.

[23] Kato, Y., Nishiyama, K., Nishimura, A., Noda, T., Okabe, K., Kusakabe, T., Kanda, Y. and Nishida, M. (2022). Drug repurposing for the treatment of COVID-19. *Journal of Pharmacological Sciences* **149**, 108-114.

Figure legends

Fig. 1 Purified Dyns-His$_6$ in this study.

The purified proteins were resolved by 15 % SDS-PAGE and stained with Coomassie Brilliant Blue.

Fig. 2 All Dyn-His$_6$ isoforms released orthophosphate *via* GTP hydrolysis in a time-dependent manner.

The GTPase activities of Dyn1-His$_6$ (hollow circles), Dyn2-His$_6$ (solid circles), and Dyn3-His$_6$ (triangles) were measured as the concentration of released Pi. (A) GTPase activities of Dyns-His$_6$ isomers in the presence of 5.0 μM PS-liposomes. The GTPase activities of 20 nM Dyns-His$_6$ isomers were measured at 30 °C. (B) GTPase activities of Dyns-His$_6$ isomers in the absence of PS-liposomes. The GTPase activities of 20 nM Dyns-His$_6$ were measured at 30 °C.

Fig. 3 Clomipramine inhibits dynamin GTPase activity by PS-liposome stimulation.

The GTPase activity of purified Dyns-His$_6$ (20 nM) was determined in the presence of various concentrations of MiTMAB (hollow circles), sertraline (solid circles) or clomipramine (triangles). All results are representative of at least 3 independent experiments. (A) GTPase activity of Dyn1-His$_6$ in the presence of 5.0 μM PS-liposomes. (B) GTPase activity of Dyn2-His$_6$ in the presence

of 5.0 μM PS-liposomes. (C) GTPase activity of Dyn3-His$_6$ in the presence of 5.0 μM PS-liposomes.

Fig. 4 Clomipramine do not inhibit basal intrinsic dynamin GTPase activity.

The GTPase activity of purified Dyns-His$_6$ (20 nM) was determined in the presence of various concentrations of MiTMAB (hollow circles), sertraline (solid circles) or clomipramine (triangles). All results are representative of at least 3 independent experiments. (A) GTPase activity of Dyn1-His$_6$ in the absence of PS-liposomes. (B) GTPase activity of Dyn2-His$_6$ in the absence of PS-liposomes. (C) GTPase activity of Dyn3-His$_6$ in the absence of PS-liposomes.

Fig. 5 Clomipramine did not affect the oligomerization of Dyn2-His$_6$

The effect of clomipramine on the oligomerization of Dyn2-His$_6$ was analysed by BN-PAGE. Lane 1, Dyn2-His$_6$ + PS-liposomes; Lane 2, Dyn2-His$_6$ + PS-liposomes + 1% SDS; Lane 3, Dyn2-His$_6$ + PS-liposomes + clomipramine; Lane 4, Dyn2-His$_6$ + PS-liposomes + clomipramine + 1% SDS. Arrow I corresponds to a dynamin monomer. Arrows II and III correspond to dynamin oligomers.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5