The mobility of the cap domain is essential for the substrate promiscuity of a family IV

esterase from sorghum rhizosphere microbiome

Marco Distaso[a,‡], Isabel Cea-Rama[b,‡], Cristina Coscolín[c], Tatyana N. Chernikova[a], Hai Tran[a],

Manuel Ferrer[c,*], Julia Sanz-Aparicio[b,*], Peter N. Golyshin[a,*]

[a]*Centre for Environmental Biotechnology, Bangor University, LL57 2UW Bangor, UK*

[b]*IQFR, CSIC, 28006 Madrid, Spain*

[c]*ICP, CSIC, 28049 Madrid, Spain*

[‡]Equal contribution.

[*]Corresponding authors at: Institute of Physical Chemistry "Rocasolano", CSIC, Serrano 119,

28006 Madrid, Spain (J. Sanz-Aparicio). ICP, CSIC, Marie Curie 2, 28049 Madrid, Spain (M.

Ferrer). Centre for Environmental Biotechnology, Bangor University, LL57 2UW Bangor, UK

(P.N. Golyshin)

E-mail addresses: mferrer@icp.csic.es (M. Ferrer), xjulia@iqfr.csic.es (J. Sanz-Aparicio),

p.golyshin@bangor.ac.uk (P.N. Golyshin)

Phone numbers: +34915854872 (M. Ferrer), +34915619400 (J. Sanz-Aparicio),

+441248383629 (P.N. Golyshin)

Running title: The cap domain of an esterase from sorghum rhizosphere

ABSTRACT Metagenomics offers the possibility to screen for versatile biocatalysts. In this study, the microbial community of the *Sorghum bicolor* rhizosphere was spiked with technical cashew nut shell liquid, and after incubation, the eDNA was extracted and subsequently used to build a metagenomic library. We report the biochemical features and crystal structure of a novel esterase from the family IV, EH_0, retrieved from an uncultured sphingomonad after a functional screen in tributyrin agar plates. EH_0 (T_{opt}, 50 °C; T_m, 55.7 °C; pH_{opt}, 9.5) was stable in the presence of 10-20% v/v organic solvents and exhibited hydrolytic activity against *p*-nitrophenyl esters from acetate to palmitate, preferably butyrate (496 U mg^{-1}), and a large battery of 69 structurally different esters (up to 30.2 U mg^{-1}), including bis(2-hydroxyethyl)-terephthalate (0.16 ± 0.06 U mg^{-1}). This broad substrate specificity contrasts with the fact that EH_0 showed a long and narrow catalytic tunnel, whose access appears to be hindered by a thigth folding of its cap domain. We propose that this cap domain is a highly flexible structure whose opening is mediated by unique structural elements, one of which is the presence of two contiguous proline residues likely acting as possible hinges, that altogether allow for the entrance of the substrates. Therefore, this work provides a new role for the cap domain, which until now was thought to be immobile elements that contain hydrophobic patches involved in substrate pre-recognition and in turn substrate specificity within family IV esterases.

IMPORTANCE A better understanding of structure–function relationships of enzymes allows revealing key structural motifs or elements. Here, we studied the structural basis of the substrate promiscuity of EH_0, a family IV esterase, isolated from a sample of the *Sorghum bicolor* rhizosphere microbiome exposed to technical cashew nut shell liquid. The analysis of EH_0 revealed the potential of the sorghum rhizosphere microbiome as a source of enzymes

with interesting properties, such as pH and solvent tolerance and remarkably broad substrate

promiscuity. Its structure resembled those of homologous proteins from mesophilic

Parvibaculum and *Erythrobacter* spp. and hyperthermophilic *Pyrobaculum* and *Sulfolobus*

spp. and had a very narrow, single-entry access tunnel to the active site, access which is

controlled by a capping domain that includes a number of not conserved proline residues.

These structural markers, distinct from those of other substrate promiscuous esterases, can

help tuning substrate profiles beyond tunnel and active site engineering.

KEYWORDS Crystal structure, esterase, α/β-fold hydrolase, metagenome, promiscuity,

rhizosphere.

Currently, in the biotech industrial sector, there is a constant demand for novel biocatalysts

with enhanced biochemical properties fulfilling industrial requirements to replace low-

performance enzymes and dismiss chemical processes (1). From this perspective, the

enzymatic diversity within the microbial communities, that may reach for example up to 10^3-

10^5 microbial species in 1 g of soil, represents a yet unexplored and exceptional resource for

new biocatalysts (2, 3), and thus, it can be a source of biocatalysts. A straightforward

evaluation of such diversity can be performed by metagenomics, through a number of

bioinformatics and so-called naïve screening methods (2, 4-6).

Carboxylic ester hydrolases (EC 3.1.1.-) are a class of enzymes widely distributed in nature

that have attracted much attention because of their high sequence diversity, structural

versatility, catalytic versatility, robustness and lack of need for cofactors (7-10). Considering

the importance of these classes of enzymes, it is not surprising that they are priority targets

to screen by metagenomics. For example, approximately 280,638 sequences encoding

carboxylic ester hydrolases have been cataloged (10), and more than 4,000 new enzymes

have been discovered through metagenomics (11); few of them show enhanced biophysical

properties, such as stability at extreme temperatures, salinity, pH and solvents conditions,

high catalytic efficiency, broad substrate specificity and regio- and enantioselectivity (6, 12-

14). Additionally, metagenomic studies on soil environments significantly contributed to

broadening the knowledge about this class of enzymes with the reannotation of sequences

known as hypothetical proteins and the definition of new esterase families (15, 16).

Carboxylic ester hydrolases are also the most studied enzymes with respect to promiscuity

(17), including catalytic (18) and substrate (19) promiscuity. The question that arises is, would

it not be more efficient to have just a couple of promiscuous enzymes with the ability to

perform a wide range of reactions (19). Based on this principle, biotechnological industries

demand biocatalysts that are capable of transforming a wide range of substrates. Their

isolation has been solved through the application of metagenomic techniques that have

emerged as robust tools for the discovery of unknown biocatalysts, including ester hydrolases

with a broad substrate scope. This method, together with the understanding of the structural

mechanisms and engineering behind promiscuous enzymes, seems crucial for their industrial

implementation. In this context, our recent study of approximately 145 ester hydrolases has

revealed that members of family IV show higher levels of substrate promiscuity than those of

other families. This is the result of their capacity to adapt the topology of the large but

occluded active sites to a high variety of substrates but also to the presence of a cap domain

next to the entrance of the substrate tunnel (19). The cap domains were thought to be

immobile elements, unlike lid elements in lipases, that contain hydrophobic patches involved

in substrate pre-recognition which could contribute to the substrate promiscuity within

members of family IV (20).

The aim of this study was to mine carboxylic ester hydrolases of the soil rhizosphere of

sorghum plants after enrichment with cashew nut shell liquid (CNSL), an oily substance rich

in phenolics and fatty acids (21), with respect to novel biocatalysts. By applying an activity-

based strategy, four such enzymes were retrieved and characterized. We also reported the

crystal structure of one of these enzymes with an extraordinarily broad substrate specificity.

The structural information, complemented with the experimental analysis of the substrate

specificity of a series of variants, provides additional mechanistic understanding of the

molecular basis for the role of the cap domain in the promiscuous character of members

comprising family IV esterases. Importantly, we propose the role of proline residues at the N-

terminal region comprising $\alpha1$-$\alpha2$ helices as hinges of the movement of the cap domain

allowing substrates to enter the active center.

Results

Identification of a carboxylic ester hydrolase. The search for hydrolytic enzymes in the

metagenomics library SorRhizCNSL3 W (see details in Materials and Methods) was carried out

by applying a metagenomic functional screening approach. The strategy involved the use of

LB agar plates containing tributyrin (22). The pooling of 40 clones per well dispensed into 96-

well microtiter plates (approximately 3.8 k clones in one plate) facilitated the colony screening

at a high throughput. After screening more than 100,000 clones, 1 hit from the SorRhizCNSL3

W library showed lipase/esterase activity. Fosmid insert was extracted using the QUIAGEN

Large-Construct Kit (QIAGEN, Hilden, Germany) according to the manufacturer's protocol,

digested with restriction enzymes for insert size estimation, and the insert sequenced by

Illumina technology. Upon the completion of sequencing, the reads were quality-filtered and

assembled to generate nonredundant meta-sequences, and genes were predicted and

annotated as described previously (12). One gene encoding a predicted carboxylic ester hydrolase was identified. The gene was amplified with specific primers, cloned into the p15TV-L vector and transformed into *E. coli* BL21(DE3) cells for expression of the N-terminal (His)6-tagged proteins. The deduced amino acid sequence of the enzyme (324 amino acids long) was used for homology searches in the taxonomy and functional assignment. A database search indicated that EH_0 showed 99% identity with the α/β hydrolase enzyme from *Sphingomonas pruni* (protein ID WP_066587239), both of which are classified as members of the α/β hydrolase-3 family (PF07859). The typical H-G-G-G motif and the G-X-S-X-G catalytic motif are conserved, and EH_0 clustered together with family IV esterase.

Biochemical characterization. The recombinant protein was successfully expressed in soluble form and purified by nickel affinity chromatography. Purified protein was desalted by ultrafiltration, and its enzymatic activity was assessed. Four model *p*-nitrophenyl (*p*-NP) ester substrates with different chain lengths were used to determine the substrate specificity of the enzymes and therefore determine whether the enzymes are in fact true lipases or esterases. Lipases hydrolyze ester bonds of long-chain triglycerides more efficiently than esterases, which instead exhibit the highest activity toward water soluble esters with short fatty acid chains (23). The substrates used for the hydrolytic test were *p*-NP acetate (C2), *p*-NP butyrate (C4), *p*-NP dodecanoate (C12), and *p*-NP palmitate (C16). The hydrolytic activity was recorded under standard assay conditions (Fig. 1). EH_0 showed a specific activity of 496.5 U mg^{-1} for *p*-NP butyrate, which was the best substrate. Lower levels of activity were observed with longer chain esters (C ≥ 12). The esterase followed the Michaelis–Menten kinetics, and its kinetic parameters are reflected in Table 1. A comparison of the catalytic efficiency values (k_{cat}/K$_M$) indicated a high reactivity toward *p*-NP-butyrate followed by *p*-NP-acetate.

Its voluminous (volume of the active site cavity: 5133 Å^3) but low exposed (solvent

accessible surface area (SASA): 5.07 over 100 dimensionless percentage) active site allows

hydrolysis of a broad range of 68 out of 96 structurally and chemically diverse esters (Table

S1), as determined by a pH indicator assay (pH 8.0, 30 °C). Phenyl acetate (30.23 U mg^{-1}) and

glyceryl tripropionate (29.43 U mg^{-1}), were the best substrates. We also found that EH$_0$

efficiently hydrolyzed bis(2-hydroxyethyl)-terephthalate (BHET; 163.6 ± 6.2 U mg^{-1}), an

intermediate in the degradation of polyethylene terephthalate (PET) (24); HPLC analysis (Fig.

S1), performed as described (25), confirmed the hydrolysis of BHET to mono-(2-

hydroxyethyl)-terephthalic acid (MHET) but not to terephthalic acid (TA). However, using

previously described conditions (25), we found that the enzyme did not hydrolyze large

plastic materials such as amorphous and crystalline PET film and PET nanoparticles from

Goodfellow. According to the number of hydrolyzed esters, EH$_0$ can be thus considered as an

esterase with a wide substrate specificity, similar to other enzymes of the family IV (19, 20).

EH$_0$ showed maximal activity at 50 °C, retaining more than 80% of the maximum activity at

40-55 °C (Fig. 2A), suggesting that it is moderately thermostable. This was confirmed by

circular dichroism analysis, which revealed a denaturing temperature of 55.7 ± 0.2 °C (Fig. 2B).

Its optimal pH for activity was 9.5 (Fig. 2C). The effect on the enzymatic activity of organic

solvents at different concentrations was evaluated (Table S2). An activation effect was

observed for EH$_0$ when 10% methanol (60% activity increase) and 10-20% DMSO (22-40%

increase) were added to the reaction mixture. The presence of bivalent and trivalent cations

did not have a remarkable positive effect on the activity of the enzymes, which showed, in

some cases, tolerance to high concentrations of cations (Table S3). A prominent inhibiting

effect was shown for all cations, except for magnesium, which was well tolerated at 1-10 mM

(<5% inhibition).

EH$_0$ presents tight folding of its cap domain. The crystal structure of wild type EH$_0$ was obtained at 2.01 Å resolution, with the P2$_1$2$_1$2$_1$ space group and two crystallography-independent molecules in the asymmetric unit. Molecular replacement was performed using Est8 as a template (PDB code 4YPV) (26), and the final model was refined to a crystallographic R-factor of 0.1717 and R-free of 0.2019 (Table S4). As with other reported family IV esterases, EH$_0$ has an **α/β** hydrolase fold with two different domains, a cap domain (residues 1-43 and 208-229) and a catalytic domain (residues 44-207 and 230-324), constituted by a total of 9 **α**-helices and 8 **β**-sheets (Fig. 3A). The catalytic domain was composed of a central **β**-sheet with eight parallel **β**-strands (**β**1, **β**3, **β**4, **β**5, **β**6, **β**7, **β**8), except **β**2, which was antiparallel and surrounded by five **α**-helices (**α**3, **α**4, **α**5, **α**8, **α**9). The cap domain involved four **α** helices (**α**1, **α**2, **α**6, **α**7) (Fig. 3B). There were two *cis* peptides, Ala122-Pro123 and Trp127-Pro128, located at the **β**4-**α**4 turn within the catalytic domain.

Analysis of EH$_0$ folding using the *DALI* server (27) was employed to search for homologous proteins. The closest homologs are E53 isolated from *Erythrobacter longus*, with 46% identity and RMSD of 2.3 Å on 296 Cα atoms (PDB code 7W8N) (28); Est8 isolated from *Parvibaculum*, with 38% identity and RMSD of 2.5Å on 291 Cα atoms (PDB code 4YPV) (26); PestE isolated from *Pyrobaculum calidifontis*, with 33% identity and RMSD of 2.6Å on 294 Cα atoms (PDB code 3ZWQ) (29); and EstA isolated from *Sulfolobus islandicus REY15A,* with 31% identity and RMSD of 2.6Å on 281 Cα atoms (PDB code 5LK6). The structural superimposition of these proteins reveals a high conservation of the corresponding catalytic domains, and also a common spatial arrangement of the helices at the cap domains in all the proteins except EH$_0$, where α2 and the long α7 are visibly shifted very close to its EH$_0$ active site and apparently impeding the entrance of substrates (Fig. 3C). This was unexpected considering the broad substrate specificity of esterase EH$_0$, approaching that of most promiscuous ones (19). Thus,

withdrawal of the cap domain seems a necessary requirement for allowing access of the bulky

substrates to the EH_0 catalytic site.

In agreement with this assumption, a substantial rearrangement of the cap domain was

previously described in the homolog esterase EST2 (having 31 % sequence identity), with its

M211S/R215L double variant being trapped in the crystal in a conformation resembling the

open form of lipases (30). However, the authors did not assign any biological relevance to this

issue, considering this state as an artifact derived from the crystal packing. In our study, some

flexibility at this region has been found from the two conformations adopted by loop α1-α2

observed in subunits A and B of EH_0 (Fig. 3D). In addition, the inspection of residues within

the cap domain showed a high number of proline residues (Pro8, Pro21, Pro23, Pro31, Pro46

and Pro47) (Fig. 3B) that are mostly non-conserved and could confer flexibility to the N-

terminal cap domain allowing substrate entrance. This feature will be discussed below.

EH_0 is a dimeric enzyme. While E53 and Est8 are monomeric enzymes, and EstA is a

tetramer, EH_0 is presented as a biological homodimer with approximate dimensions of 6.3 x

4.9 x 2.7 nm, which is assembled in a twofold axis symmetry arrangement (Fig. 4A) that buries

5.6% of its total surface area. Hydrogen bonds mainly involve β8 and α8, while salt bridges

involve motifs β8 and α9 (Fig. 4B and Table S5). Similar to its dimeric homolog PestE,

oligomerization occurs through a tight interaction of β8 strands from both subunits. However,

while only the subsequent helices **α12** were involved in the PestE interface (29) (Fig. 4C), both

the precedent **α8** and the subsequent **α9** helices make the EH_0 interface (Fig. 4D). These

different interactions observed in PestE and EH_0 were reflected in a different orientation

between the monomers that, nevertheless, present a similar distance between the catalytic

serines of 35-38 Å and an equivalent disposition of the tunnels giving access to the catalytic

site at two edges of the dimer (Fig. 4E, 4F). Therefore, as seen in Fig. 4A, the two cap domains

are far from the interface and project out from the dimer, revealing that dimerization is not

affecting the cap function.

The peculiar EH$_0$ active site. EH$_0$ has a long catalytic tunnel with a very narrow entrance

(approximately 16.8 Å depth) (Fig. 5A). The catalytic triad of EH$_0$ is formed by Ser161 (in the

conserved motif ^{159}G-D-S-A-G^{163} in the nucleophilic elbow), Asp253 (in the conserved motif

^{253}D-P-I-R-D^{258}) and His283 (Fig. 5B). To analyze the active site, a series of soaking and

cocrystallization experiments were performed with different suicide inhibitors, all of which

were unsuccessful. A deep inspection of the active site showed that the nucleophilic Ser161

is hydrogen bonded to Glu226 from the close **α7**, and a movement of loop **β3-α3**, including

the oxoanion in the conserved motif ^{87}H-G-G-G^{90,} compared to EH$_0$ homologs (Fig. 5B).

Therefore, to remove this hydrogen link in an attempt to expand access to the active site

enabling ligand binding, the EH$_{0E226A}$ variant was produced by directed mutagenesis.

However, the crystals grown from this variant failed to diffract, suggesting a high level of

disorder, putatively due to an increased mobility of the cap. Interestingly, the activity profile

of this variant reveals a faster hydrolysis of most of the esters (62 esters in total; from 70- to

1.2-fold; average: 6.1-fold). Only 7 esters were hydrolyzed to a lower extent compared to the

wild type (from 1.5- to 17-fold; average: 4.6-fold). Remarkably, the EH$_{0E226A}$ variant was

capable of accepting large and voluminous esters such as dodecanoyl acetate, pentadecyl

acetate, vinyl laurate, methyl 2,5-dihydroxycinnamate, and ethyl 2-chlorobenzoate, which

were not accepted by the wild-type enzyme (Table S1). This apparently supports that removal

of the Ser161-Glu226 hydrogen bond increases cap flexibility and enhances the enzymatic

efficiency toward large esters either at the acyl or alcohol sites.

Indeed, it appears clear that helix α7 must retract from the catalytic site to allow substrate

entrance, but this helix seems fixed by many atomic interactions. Close to the Ser161-Glu226

hydrogen bond, Tyr223 makes additional hydrogen bonds to Asp194 and Lys197 main chain,

from loop β6-α6, and with the side chain of Gln258 from α8 (Fig. 5C). A mutation EH$_{0Y223A}$ was

generated by directed mutagenesis and found, unlike the EH$_{0E226A}$ mutation, little effect on

conversion rates compared to the wild-type enzyme. However, this mutation allowed the

hydrolysis of large and voluminous esters, such as dodecanoyl acetate, pentadecyl acetate,

methyl 2,5-dihydroxycinnamate and ethyl 2-chlorobenzoate, which were also hydrolyzed by

the variant EH$_{0E226A}$. Unlike EH$_{0E226A}$, the EH$_{0Y223A}$ variant was not unable to hydrolyze vinyl

laurate, but it was able to hydrolyze methyl 3-hydroxybenzoate, not accepted by the EH$_{0E226A}$

variant. This suggested that Tyr223 may play a role in accepting large esters, particularly at

the acyl side, but may also play an additional role in substrate specificity different from that

of Glu226.

Additionally, at the beginning of α7, Trp218 is within a hydrophobic pocket surrounded by

residues Phe13, Ile17, Leu28 and Phe219 from the cap domain and Phe91 from the catalytic

domain, which is also anchored by interaction with loop α1-α2 and Pro23 (Fig. 5D). Thus, to

depict how these tight molecular packing may be disrupted by the proposed cap motion,

molecular dynamics were applied to crystallographic refinement through the ensemble

refinement strategy, which is shown to model the intrinsic disorder of macromolecules giving

more accurate structures. The ensemble models obtained for molecules A and B within the

asymmetry unit are shown in Fig. 5E and 5H, respectively. The analysis of the molecule A

conformers (Fig. 5E) revealed that the region comprising α1 and α2 shows a wide spectrum

of possible pathways from more "open" to more "closed" conformations. At one edge, Pro23

is in an extended α1-α2 loop far from Trp218, therefore releasing α7 that consequently could

retract from the catalytic pocket ("open-like conformation") (Fig. 5F). In fact, the ensemble

refinement models a very flexible conformation being even unstructured at regions

corresponding to α1 and α2. At the other edge, the second scenario is the entrapment of Trp218 by Pro23 in loop α1-α2, hindering substrate entrance ("closed-like conformation") (Fig. 5G). This last scenario is equivalent to the 3D structure captured by crystallography (Fig. 5D). Furthermore, three prolines can be found within this α1-α2 loop, Pro21 (at the end of α1), Pro23 (at the middle) and Pro31 (at the beginning of α2), all of them unique to EH_0, which are probably behind the two different conformations observed at this loop in both subunits within the asymmetric unit (Fig. 3B and 3D), and explain the ensemble of conformers modelled for molecule B (Fig. 5H).

Furthermore, as seen in Fig. 3B, Pro46 and Pro47 are potential hinges that would involve flexibility of a larger region of the EH_0 cap domain, including the whole N-terminal peptide chain up to the end of α2. This is consistent with the more "open" conformations resulting from the ensemble refinement shown in Fig. 5F. In fact, the sequence comparison of EH_0 to its closest homologs reveals that only EH_0 has two sequential prolines at this region and that Pro46 is unique to EH_0 (Fig. 6), which could be a reason behind the high EH_0 promiscuity. Interestingly, EST2 also presents the two contiguous Pro residues (Pro38, Pro39), which can also confer high mobility to its cap domain and facilitate the "open-like" conformation captured in the crystal mentioned above (30). Therefore, the mutation EH_{0P46A} was generated by directed mutagenesis and submitted to crystallization experiments to investigate the Pro46 putative role. However, the crystals grown from this variant EH_{0P46A} failed to diffract, suggesting that removal of Pro46 introduces some structural instability to the polypeptide chain resulting in crystal disorder. Moreover, analysis of the activity profile showed that Pro46 is a critical residue for the entry and hydrolysis of bulky substrates, as its mutation by Ala extends the substrate specificity from 68 to 84 esters. Additionally, the hydrolytic rate increased from 1.2- to 18000-fold (average: 335-fold) for most esters. This variant was also

able to hydrolyze large glyceryl trioctanoate and 2,4-dichlorophenyl 2,4-dichlorobenzoate,

which were not hydrolyzed by the wild-type enzyme or the EH_{0E226A} and EH_{0Y223A} variants.

Consequently, although the proposed role of Pro46 and Pro47 as putative hinges enabling the

opening of the cap domain seems appealing, other mechanisms promoting EH_0 plasticity to

bulky substrates may also operate. Furthermore, as seen below, it should be noted that the

two proline residues are located at the entrance of the narrow tunnel giving access to the

active-site, an issue that ascribes a prominent role to both residues in binding activity and

specificity.

Structural details of the EH_0 active site and assignment of the acyl and alcohol moieties

were explored by comparison with its homologs E53 complexed with 4-nitrophenyl

hexanoate (PDB code 6KEU) (28) and EH_{1AB1} in complex with a derivative of methyl 4-

nitrophenylhexylphosphonate (PDB code 6RB0) (31). However, the acyl and alcohol moieties

of these complexes are located in opposite sites (Fig. 5I, 5J). Therefore, as we could not obtain

complexes from EH_0, the activity experiments were crucial to correctly assign acyl/alcohol

moieties. As mentioned above, experimental evidence demonstrated that Tyr223 produces a

steric hindrance at the acyl moiety, and consequently, the acyl and alcohol sites correspond

to those observed in EH_{1AB1} (Fig. 5J). On the basis of this assumption, the acyl binding site

seems to be a small cavity bordered by the Tyr199 and Ile255 side chains, which produce

steric hindrance for substrates with large acyl moieties. The long and narrow alcohol binding

site is surrounded by both hydrophobic (Phe13, Phe91, Val92, Leu288) and hydrophilic

residues (Asp45, His99, Asp160, Tyr190, Thr287), with Pro46 and Pro47 being at the entrance

of the tunnel (Fig. 5K). Most residues at the acyl and alcohol moieties are conserved among

EH_0 homologs, with the exception of Met40, Pro46, Tyr199, Glu226 and Thr287. Remarkably,

the bulky Met40 and Tyr199 residues are substituted by smaller residues in the EH_0 homologs.

As previously mentioned, Pro46 is unique in EH_0, while Glu226 is substituted by a conserved

Asp, and finally, most homologs show an Asn residue instead of Thr287 (Fig. 6). Therefore, as

the retraction of the cap domain must be performed to allow substrate entrance, residues

Phe91, Tyr190 and Tyr199 from the catalytic domain, which are located close to the catalytic

triad, seem essential for substrate specificity (Fig. 5K).

DISCUSSION

In this study, the microbial community of the *Sorghum bicolor* rhizosphere was exposed to

a chemical treatment prior to eDNA extraction to construct a metagenomic library. Plant roots

can secrete exudates composed of a large variety of compounds into the soil, some of which

may play important roles in the rhizosphere (32, 33), and with effects that involve multiple

targets, including soil microorganisms. This is why an amendment of the soil with technical

cashew nut shell liquid (tCNSL), containing a mixture of phenolic compounds with long

aliphatic side chains (up to C22:0), was carried out directly in the rhizosphere and, later, for

three weeks under control laboratory conditions, as we were interested in screening lipolytic-

like activity. By applying metagenomics techniques we retrieved an esterase, EH_0, highly

similar (99% identity) to the predicted α/β hydrolase from the genome of *S. pruni* (acc. nr.

WP_066587239). The most homologous, functionally characterized protein is actually

P95125.1, a carboxylic ester hydrolase LipN from *Mycobacterium tuberculosis* H37Rv, which

shows only 41% amino acid sequence identity with EH_0. That said, given the high identity of

WP_066587239 and EH_0 we expect both hydrolases having similar properties, yet to be

experimentally confirmed. Indeed, we observed that there are only three changes in their

sequences, which are located on the outside and in loops away from the key residues and the

dimerisation interface (Fig. S2).

EH_0 was classified within the previously described hormone-sensitive lipase (HDL) type IV

family, that is one of the at least 35 families and 11 true lipase subfamilies known to date (10,

23, 34). This family is reported to contain ester hydrolases with relative SASA values ranging

from 0 to 10% and high levels of substrate specificity (19). Note that SASA, computed as a

(dimensionless) percentage (0–1 or 0-100) of the ligand SASA in solution (19), is a parameter

that inspect the solvent exposure of the cavity containing the catalytic triad and the capacity

of a cavity to retain/stabilize a substrate (19). For example, a SASA of 40% (over 100%) implies

that 40% of the surface is accessible to the solvent, which facilitates the escape of the

substrate to the bulk solvent; this is the case of enzymes with active site in the surface where

the catalytic triad is highly exposed. By contrast, enzymes that has a larger but almost fully

occluded site that can better maintain and stabilize the substrate inside the cavity, are

characterized by relative SASA values of approximately 0–10%. This is the case of EH_0, which

has a SASA of 5.07% because a large but almost fully occluded active site, an architecture that

is known to better maintain and stabilize a higher number of substrates inside the cavity (19).

Indeed, this enzyme houses a very long and narrow catalytic pocket, where helix α7 is very

close to the catalytic triad with residue Glu226, making a direct hydrogen bond to the catalytic

nucleophile Ser161. Therefore, it appeared clear that the cap domain must retract to allow

substrate entrance to the active site. The movement of this domain was modeled by

combining X-ray diffraction data with molecular dynamics simulation through the ensemble

refinement procedure. This strategy showed a broad range of putative conformations at the

cap domain, with Pro46 and Pro47 likely acting as hinges conferring a high plasticity to the N-

terminal region of the cap. Remarkably, the presence of a number of prolines at this region

particularly these two sequential prolines, is a unique feature of EH_0 compared to its

homologs, and other substrate-promiscuous members of the family IV (19). Mutational

analysis confirmed the role of one of these prolines in the access and hydrolysis of large and

voluminous substrates and thus in the increase in the substrate promiscuity level.

The above structural features differ from those of other substrate promiscuous family IV

esterases, tested over the same set of ester substrates, namely, EH_{1AB1} (31), and EH_3 (35, 36)

capable of hydrolyzing a similar number of esters. The comparison of EH_{1AB1} and EH_3 with EH_0,

shows major differences related to the *lid* (Fig. 7A). Whereas EH_{1AB1} and EH_3 show large and

wide catalytic pockets with two possible access to the binding site (Fig. 7B and C), EH_0 has a

unique, narrow and long entrance to the catalytic active site, as said before (Fig. 7D).

Therefore, in the case of EH_0 only a structural rearrangement of the cap domain would allow

its adaptation to all different substrates, which likely implies that the cap domain of EH_0

exhibits more flexibility than those from EH_{1AB1} and EH_3. This is consistent with the fact that

of the 80 esters that the three enzymes together are able to hydrolyze, 61 (or 76%) were

common to all three, and that EH_0 was the only one able to hydrolyze such bulky substrates

as 2,4-dichlorobenzyl 2,4-dichlorobenzoate or diethyl-2,6-dimethyl 4-phenyl-1,4-dihydro

pyridine-3,5-dicarboxylate.

Biochemical characterization of the novel esterase also revealed that the activity or EH_0

was in most cases stimulated in the presence of 10% organic solvents, particularly in 10%

methanol and DMSO. Such activation is also a characteristic of some lipases. For example, the

analysis of the lipase from *Thermus thermophilus* revealed that although the overall structure

was maintained stable with or without polar organic solvent, the lid region was more flexible

in the presence of the later. The flexible lid facilitates the substrate to access the catalytic site

inside the lipase and the lipase displays enhanced activity in the presence of a polar organic

solvent (37). The use of organic solvents offers more advantages over canonical aqueous

biocatalysis for various reasons: higher solubility of hydrophobic substrates, minor risk of

contamination, and higher thermal stability (38-40). EH_0 has a potential advantage in

applications that require alkaline conditions due to its ability to act at the optimal pH of 9.0.

Temperature-controlled tests indicated a mesophilic/slightly thermophilic profile of the

esterase as expected from the original habitat at moderate temperatures. In addition, the

structure of EH_0 is similar (31-33% identity) to those of extremophiles, namely, *Pyrobaculum*

(3ZWQ) and *Sulfolobus* (5LK6) species.

In summary, the present study evaluated the conformational plasticity of the cap domain

in members of Family IV, and the role of several non-conserved prolines as putative structural

factor regulating their broader substrate specificity compared to other members of the 35

families and 11 true lipase subfamilies reported (10). This high molecular flexibility is markedly

different to that found in other family IV esterases and a family VIII β-lactamase fold hydrolase

(EH_7) which has been recently shown to be highly substrate-promiscuous. In this case, the

broad substrate specificity is given by the presence of a more opened and exposed S1 site

having no steric hindrance for the entrance of substrates to the active site, and more flexible

R1, R2 and R3 regions allowing the binding of a wide spectrum of substrates into the active

site.

CONCLUSIONS

Activity-based metagenomics approach was used to study the microbial enzyme diversity

in rhizosphere soil of *Sorghum* plants amended with CNSL soil. A novel esterase was found,

which possessed a broad substrate promiscuity in combination with a significant pH and

solvent tolerance. This work is crucial for deciphering structural markers responsible for the

outstanding broad substrate specificity of EH_0. Indeed, this work further provides important

insights into the role of cap domains and its contribution to the diverse selectivity profiles and

thus versatility of family IV esterases/lipases toward the conversion of multiple substrates.

MATERIALS AND METHODS

Plant material and outdoor seed germination. Seeds of *Sorghum bicolor* genotype BTx623

were obtained from the Agricultural Research Service of United States Department of

Agriculture (USDA) (Georgia, US) as a gift. Field soil was sampled from the Henfaes Research

Centre (53°14'21.0"N, 4°01'06.5"W, Gwynedd, Wales) in September 2014. The soil sample

was composed of a mixture of five topsoil samples collected from randomly selected positions

in the field. The soil was air-dried, mixed thoroughly, and stored at room temperature for use

in subsequent experiments. Two L pots were filled with soil, and two seeds of *S. bicolor*

BTx623 were planted per pot. Plants were cultivated in a greenhouse at 20 °C, and the soil

moisture content was maintained with tap water.

Enrichment with CNSL. Three grams of technical cashew nut shell liquid (tCNSL) dissolved

in 70% ethanol was added to a pot of 20-day-old plants and thoroughly mixed with the soil.

After 60 days, plants were pulled out of the pot, and the soil was shaken off; samples of

rhizosphere soil attached to the plant roots were then brushed off and collected. tCNSL was

provided by the BioComposites Centre at Bangor University (Wales, UK). Three biological

replicates of laboratory microcosm enrichment were set up in conical 1 L Erlenmeyer flasks

by mixing 10 g of the collected rhizosphere soil with 300 mL of sterile Murashige Skoog Basal

medium (Sigma) and 10 mg/L cycloheximide. tCNSL was dissolved in 70% ethanol and added

to the medium to a final concentration of 0.1 g/L; flasks were incubated at 20 °C in an orbital

shaker. 50 g of oil slurry was sampled every seven days, and fresh tCNSL-containing (0.1 g/kg)

medium was added to replace the volume of the medium.

Extraction of DNA and generation of metagenomic library. Samples collected after 3

weeks of flask microcosm enrichment were used for the construction of fosmid metagenomic

libraries. Environmental DNA was extracted using the Meta-G-Nome DNA Isolation Kit

(Epicenter Biotechnologies; WI, USA) according to the manufacturer's instructions. Briefly, 50

ml of the soil suspension from the flask enrichment was centrifuged at 400 x g for 5 min. The

supernatant was filtered through 0.45 μm and 0.22 μm membrane filters. This procedure was

repeated with the initial soil sample four times: the remaining soil was resuspended in PBS

and centrifuged, and the supernatant was filtered as before. Filters were combined, and the

sediment on the filter was resuspended in extraction buffer and collected. DNA extraction

was carried out according to the protocol described by the manufacturer. The quality of the

extracted DNA was evaluated on agarose gel and quantified with the Quant-iT dsDNA Assay

Kit (Invitrogen) on a Cary Eclipse fluorimeter (Varian/Agilent) according to the manufacturer's

instructions. The extracted metagenomic DNA was used to prepare two different

metagenomic fosmid libraries using the CopyControl™ Fosmid Library Production Kit

(Epicenter). DNA was end-repaired to generate blunt-ended, 5'-phosphorylated double-

stranded DNA using reagents included in the kit according to the manufacturer's instructions.

Subsequently, fragments of 30-40 kb were selected by electrophoresis and recovered from a

low melting point agarose gel using GELase 50X buffer, and GELase enzyme preparation was

also included in the kit. Nucleic acid fragments were then ligated to the linearized CopyControl

pCC2FOS vector in a ligation reaction performed at room temperature for 4 hours, according

to the manufacturer's instructions. After in vitro packaging into phage lambda (MaxPlax™

Lambda Packaging Extract, Epicenter), the transfected phage T1-resistant EPI300™-T1R *E. coli*

cells were spread on Luria-Bertani (LB) agar medium (hereinafter, unless mentioned

otherwise, the agar content was 1.5% w/vol) containing 12.5 μg/ml chloramphenicol and

incubated at 37 °C overnight to determine the titer of the phage particles. The resulting

library, SorRhizCNSL3 W, has an estimated titer of 1.5x10^6 clones. For long-term storage, the

library was plated onto solid LB medium with 12.5 µg/ml chloramphenicol, and after

overnight growth, colonies were washed off from the agar surface using LB broth with 20%

(v/v) sterile glycerol, and aliquots were stored at −80 °C.

Screening metagenomic libraries: agar-based methods. Fosmid clones obtained by plating

the constructed libraries on LB agar plates were arrayed in 384-microtiter plates (1

clone/well) or alternatively in 96-microtiter plates (pools of approximately 40 clones/well)

containing LB medium and chloramphenicol (12.5 µg/ml). The plates were incubated at 37 °C

overnight, and the day after replication, the plates were produced and used in the screening

assay. Glycerol (20% v/v, final concentration) was added to the original plates, which were

stored at -80 °C. Gel diffusion and colorimetric assays were adopted for the screening of the

desired activities. The detection of lipase/esterase activity was carried out on LB agar

supplemented with chloramphenicol (12.5 µg/ml), fosmid autoinduction solution (2 ml/L)

(Epicenter), and 0.3% (v/v) tributyrin emulsified with gum arabic (2:1, v/v) by sonication. The

previously prepared microtiter plates were printed on the surface of large (22.5 cm x 22.5 cm)

LB agar plates using 384-pin polypropylene replicators and incubated for 18-48 hours at 37

°C. Lipolytic activity was identified as a clear zone around the colonies where tributyrin was

hydrolyzed (12).

Extraction of fosmids, DNA sequencing and annotation. The fosmid DNA of the positive

clone was extracted using the QIAGEN Plasmid purification kit (QIAGEN). To reduce the host

chromosomal *E. coli* DNA contamination, the sample was treated with ATP-dependent

exonuclease (Epicenter). The purity and approximate size of the cloned fragment were

assessed by agarose gel electrophoresis after endonuclease digestion simultaneously with

BamHI and *XbaI* (New England Biolabs, in NEBuffer 3.1 at 37° for 1 hour using 1 U of enzyme

per 1 µg DNA). DNA concentration was quantified using the Quant-iT dsDNA Assay Kit

(Invitrogen), and DNA sequencing was then outsourced to Fidelity Systems (NJ, USA) for

shotgun sequencing using the Illumina MiSeq platform. GeneMark software (41) was

employed to predict protein coding regions from the sequences of each assembled contig,

and deduced amino acid sequences were annotated via BLASTP and the PSI-BLAST tool (42).

Cloning, expression and purification of proteins. The selected nucleotide sequence was

amplified by PCR using Herculase II Fusion Enzyme (Agilent, USA) with specific oligonucleotide

primer pairs incorporating p15TV-L adapters. The corresponding fosmid was used as a

template to amplify the target genes. The primers used to amplify the esterase gene

characterized in this study were as follows: EH_{0F},

TTGTATTTCCAGGGCATGACCGAGCTCTTCGTCCGC; EH_{0R},

CAAGCTTCGTCATCATGCCGCCGCCTGTGCCATC. PCR products were visualized on a 1% TAE

agarose gel and purified using the Nucleospin PCR Clean-up Kit (Macherey-Nagel) following

the manufacturer's instructions. Purified PCR products were cloned into the p15TV-L vector,

transformed into *E. coli* NovaBlue GigaSingles™ Competent Cells (Novagen, Germany) and

plated on LB agar with 100 µg/ml ampicillin. The correctness of the DNA sequence was then

verified by Sanger sequencing at Macrogen Ltd. (Amsterdam, The Netherlands). 3-D models

of the proteins were generated by Phyre2. The intensive mode attempts to create a complete

full-length model of a sequence through a combination of multiple template modeling and

simplified ab initio folding simulation (43). The nucleotide and amino acid sequences of the

selected nucleotide sequences are available GenBank under ref. MK791218. For recombinant

protein expression, the plasmids were transformed into *E. coli* BL21(DE3) cells and

subsequently plated on LB agar with 100 µg/ml ampicillin. To confirm the esterase activity of

recombinant proteins, *E. coli* clones harboring the recombinant plasmid were streaked onto

LB agar plates containing 0.5% tributyrin and 0.5 mM IPTG, or purified enzymes were spotted

directly on the agar. The plates were then incubated at 37 °C overnight and visually inspected

for the presence of signs of substrate degradation. *E. coli* clones were grown at 37 °C to an

absorbance of 0.8 at 600 nm, induced with 0.5 mM isopropyl-β-D-galactopyranoside (IPTG),

and allowed to grow overnight at 20 °C with shaking. Cells were harvested by centrifugation

at 5000 x g for 30 min at 4 °C. For purification of recombinant protein, the following protocol

was applied. Cell pellets were resuspended in cold binding buffer (50 mM HEPES pH 7.5, 400

mM NaCl, 5% glycerol, 0.5% Triton-X, 6 mM imidazole pH 7.5, 1 mM β-mercaptoethanol, 0.5

mM PMSF) and extracted by sonication. The lysates were then centrifuged at 22,000 x g for

30 min at 4 °C, and the supernatant was purified by affinity chromatography using Ni-NTA His-

bind resin (Novagen). The column packed with the resin was equilibrated with binding buffer,

and after the addition of supernatant, it was washed with 6 volumes of wash buffer (50 mM

HEPES pH 7.5, 400 mM NaCl, 5% glycerol, 0.5% Triton-X, 26 mM imidazole pH 7.5, 1 mM β-

mercaptoethanol, 0.5 mM PMSF) to remove nonspecifically bound proteins. His-tagged

proteins were then eluted with elution buffer (50 mM HEPES pH 7.5, 400 mM NaCl, 5%

glycerol, 0.5% Triton-X, 266 mM imidazole pH 7.5, 1 mM β-mercaptoethanol, 0.5 mM PMSF).

The size and purity of the proteins were estimated by SDS–PAGE. Protein solutions were

desalted through the Amicon Ultra15 10K centrifugal filter device. Protein concentrations

were determined using the Bradford Reagent (Sigma) and the BioMate^{TM} 3S

spectrophotometer (Thermo Scientific, USA).

Biochemical assays. Hydrolytic activity was determined by measuring the amount of *p*-

nitrophenol released by catalytic hydrolysis of *p*-nitrophenyl (*p*-NP) esters through a modified

method of Gupta et al. (44). Stock solutions of *p*-NP esters (100 mM *p*-NP acetate, 100 mM *p*-

NP butyrate, 20 mM *p*-NP dodecanoate, and 20 mM *p*-NP palmitate) were prepared in

DMSO/acetonitrile (1:1 v/v). Unless stated otherwise, the enzymatic assay was performed

under standard conditions in a 1 ml reaction (50 mM potassium phosphate buffer pH 7.0,

0.3% (v/v) Triton X-100, 1 mM substrate) under agitation in a water bath at 50 °C until

complete substrate solubilization (note: solvents were present in the assays at a

concentration of 1% for 100 mM-stocks of *p*-NP butyrate and acetate, or 5% for 20 mM-stocks

of *p*-NP dodecanoate and palmitate). After that, the multititre plate was pre-incubated at 30

ºC for 5 minutes in a BioMate^{TM} 3S spectrophotometer (Thermo Scientific, USA), which was

set up at this temperature, and then an appropriate volume of purified enzyme containing

0.25 µg was added to start the reaction. The reaction mixture was incubated at 30 °C for 5

min and then measured at 410 nm in BioMate^{TM} 3S spectrophotometer (Thermo Scientific,

USA). The incubation time for *p*-NP dodecanoate and *p*-NP palmitate was extended to 15

minutes. All experiments were performed in triplicate, and a blank with denatured enzyme

was included. The concentration of product was calculated by linear regression equation

given on the standard curve performed by the reference compound *p*-nitrophenol (Sigma).

One unit of enzyme activity was defined as 1 µmol of *p*-nitrophenol produced per minute

under the assay conditions.

Kinetic parameters were determined under standard conditions and calculated by

nonlinear regression analysis of raw data fit to the Michaelis–Menten function using

GraphPad Prism software (version 6.0). For the kinetics with *p*-NP butyrate and acetate the

concentrations were set up to 0.1, 0.5, 1, 2 and 10 mM, using stock concentrations of 100 mM

in DMSO/acetonitrile (a maximum concentration of 5% DMSO/acetonitrile, was used in the

assay). For *p*-NP dodecanoate, a concentration of 0.05, 0.1, 0.5, 1, 2 and 3 mM was used, with

stock concentration of 20 mM substrate (a maximum concentration of 7.5%

DMSO/acetonitrile, was used in the assay). Raw data, and information about precision and

calculations are provided in Table S6.

The optimal pH for enzyme activity was evaluated with *p*-NP butyrate by performing the

assay in different buffers, specifically 20 mM sodium acetate buffer (pH 4.0), sodium citrate

buffer (pH 5.5), potassium phosphate buffer (pH 7.0), and Tris-HCl (pH 8.0-9.0). The enzyme

reactions were stopped by adding 1 ml of cold stop solution (100 mM K-phosphate buffer pH

7.0, 10 mM EDTA) to neutralize the pH and avoid changes in the equilibrium between *p*-

nitrophenol and the deprotonated form *p*-nitrophenoxide, which would result in a decrease

in absorption at the applied wavelength of 410 nm (45). The optimal enzymatic temperature

was investigated with *p*-NP butyrate by performing the hydrolytic assay at different

temperatures under standard conditions (see above). To determine the denaturation

temperature, circular dichroism (CD) spectra were acquired between 190 and 270 nm with a

Jasco J-720 spectropolarimeter equipped with a Peltier temperature controller in a 0.1-mm

cell at 25 °C. The spectra were analyzed, and melting temperature (T_m) values were

determined at 220 nm between 10 and 85 °C at a rate of 30 °C per hour in 40 mM HEPES

buffer at pH 7.0. CD measurements were performed at pH 7.0 and not at the optimal pH (8.5-

9.0) to ensure protein stability. A protein concentration of 0.5 mg·mL^{-1} was used. T_m (and the

standard deviation of the linear fit) was calculated by fitting the ellipticity (mdeg) at 220 nm

at each of the different temperatures using a 5-parameter sigmoid fit with SigmaPlot 13.0.

Stability in organic solvents was assayed with *p*-NP butyrate under standard conditions in

the presence of 10-20-40-60% (v/v) of the water-miscible organic solvents ethanol, methanol,

isopropanol, acetonitrile, DMSO, and a mixture acetonitrile/DMSO (50% each). The effect of

cations was investigated with *p*-NP butyrate under standard conditions by the addition of

$MgCl_2$, $CuCl_2$, $FeCl_3$, $CoCl_2$, $CaCl_2$, $MnCl_2$, and $ZnSO_4$ at concentrations in the range of 1 to 10

mM. In all cases, the measured values were then expressed as the relative activity in

comparison to the control reaction performed under standard conditions.

The hydrolysis of esters other than *p*-NP esters, including bis(2-hydroxyethyl)-

terephthalate (BHET), was assayed using a pH indicator assay in 384-well plates at 30 °C and

pH 8.0 in a Synergy HT Multi-Mode Microplate Reader in continuous mode at 550 nm over 24

h (extinction coefficient of phenol red, 8450 $M^{-1}cm^{-1}$). The acid produced after ester bond

cleavage by the hydrolytic enzyme induced a color change in the pH indicator that was

measured at 550 nm. The experimental conditions were as detailed previously (35), with the

absence of activity defined as at least a twofold background signal. Briefly, the reaction

conditions for 384-well plates (ref. 781162, Greiner Bio-One GmbH, Kremsmünster, Austria)

were as follows: protein, 0.2-2.0 μg per well; ester, 20 mM; T: 30 °C; pH, 8.0 (5 mM 4-(2-

hydroxyethyl)-1-piperazinepropanesulfonic acid (EPPS) buffer, plus 0.45 Phenol Red®);

reaction volume, 40 μl. The reactions were performed in triplicate, and datasets were

collected from a Synergy HT Multi-Mode Microplate Reader with Gen5 2.00 software (Biotek).

One unit (U) of enzyme activity was defined as the amount of enzyme required to transform

1 μmol of substrate in 1 min under the assay conditions. Raw data, and information about

precision and calculations are provided in Table S7.

Crystallization and X-ray structure determination of EH_0. Initial crystallization conditions

were explored by high-throughput techniques with a NanoDrop robot (Innovadyne

Technologies) using 24 mg·mL^{-1} protein concentrations in HEPES (40 mM, pH 7, NaCl 50 mM),

protein reservoir ratios of 1:1, 1.5:1 and 2:1, and commercial screens Crystal Screen I and II,

SaltRx, Index (Hampton Research), JBScreen Classic, JBScreen JCSG, JBScreen PACT (Jena

Bioscience). Further optimizations were carried out, and bar-shaped crystals of EH_0 were

grown after one day of mixing 1.2 μL of a mixture of protein (1 μL, 24 mg·mL^{-1}) and seeds (0.2

μL, 1:100) with guanidine hydrochloride (0.2 μL, 0.1 M) and reservoir (0.5 μL, 11%

polyethylene glycol 8000, 100 mM Bis-Tris pH 5.5, 100 mM ammonium acetate). For data

collection, crystals were transferred to cryoprotectant solution consisting of mother liquour

and glycerol (20% (v/v)) before being cooled in liquid nitrogen. Diffraction data were collected

using synchrotron radiation on the XALOC beamline at ALBA (Cerdanyola del Vallés, Spain).

Diffraction images were processed with XDS (46) and merged using AIMLESS from the CCP4

package (47). The crystal was indexed in the $P2_12_12_1$ space group, with two molecules in the

asymmetric unit and 62% solvent content within the unit cell. The data collection statistics

are given in Table S4. The structure of EH_0 was solved by molecular replacement with MOLREP

(48) using the coordinates from Est8 as a template (PDB Code 4YPV). Crystallographic

refinement was performed using the program REFMAC (49) within the CCP4 suite, with NCS

(noncrystallography symmetry) medium restraints and excluding residues 17-36. The free *R*-

factor was calculated using a subset of 5% randomly selected structure-factor amplitudes that

were excluded from automated refinement. Subsequently, heteroatoms were manually built

into the electron density maps with Coot8 (50), and water molecules were included in the

model, which, combined with more rounds of restrained refinement, reached the *R* factors

listed in Table S4. The figures were generated with PyMOL. The crystallographic statistics of

EH_0 are listed in Table S4. To extract dynamical details from the X-ray data, the coordinates

of EH_0 were first refined using PHENIX (51) and then were used as input models for a time-

averaged molecular dynamics refinement as implemented in the Phenix.ensemble-

refinement routine, which was performed as described previously (52).

Codes and accession numbers. The sequence encoding EH_0 was deposited in GenBank

with the accession number MK791218. The atomic coordinates and structure factors for the

EH_0 structure have been deposited in the RCSB Protein Data Bank with accession codes 7ZR3.

SUPPLEMENTARY MATERIAL

Supplemental material is available online only.

ACKNOWLEDGMENTS

This study was conducted under the auspices of the FuturEnzyme Project funded by the European Union's

Horizon 2020 Research and Innovation Programme under Grant Agreement No. 101000327. We also

acknowledge financial support under Grants PID2020-112758RB-I00 (M.F.), PDC2021-121534-I00 (M.F.), and

PID2019-105838RB-C33 (J.S-A.) from the Ministerio de Ciencia e Innovación, Agencia Estatal de Investigación

(AEI) (Digital Object Identifier 10.13039/501100011033), Fondo Europeo de Desarrollo Regional (FEDER) and the

European Union ("NextGenerationEU/PRTR"), and Grant 2020AEP061 (M.F.) from the Agencia Estatal CSIC. C.

Coscolín thanks the Ministerio de Economía y Competitividad and FEDER for a PhD fellowship (Grant BES-2015-

073829). P.N.G. acknowledges the Sêr Cymru programme partly funded by ERDF through the Welsh Government

for the support of the project BioPOL4Life, the project 'Plastic Vectors' funded by the Natural Environment

Research Council UK (NERC), Grant No. NE/S004548/N and the Centre for Environmental Biotechnology Project

co-funded by the European Regional Development Fund (ERDF) through the Welsh Government. The authors

acknowledge David Almendral and Ruth Matesanz for supporting the circular dichroism analysis, and Jose L.

Gonzalez-Alfonso and Francisco J. Plou for supporting the HPLC analysis. We thank the staff of the Synchrotron

Radiation Source at Alba (Barcelona, Spain) for assistance at the BL13-XALOC beamline.

We declare no conflict of interest.

P.N.G., M.F. and J.S-A. developed the conceptual framework, M.D., C.C., T.N.C. and H.T. developed the

experimental design, I.C-R. and J.S-A. performed the crystallization and X-ray structure determinations, analysis

and interpretation, M.D., M.F. and P.N.G. interpreted the experimental data, P.N.G., M.F. and J.S-A. wrote the

initial manuscript. All authors edited the manuscript. All authors read and approved the final manuscript.

References

1. Intasian P, Prakinee K, Phintha A, Trisrivirat D, Weeranoppanant N, Wongnate T, Chaiyen P. 2021.
Enzymes, in vivo biocatalysis, and metabolic engineering for enabling a circular economy and
sustainability. Chem Rev 121:10367–10451.

2. Ferrer M, Méndez-García C, Bargiela R, Chow J, Alonso S, García-Moyano A, Bjerga GEK, Steen IH,
Schwabe T, Blom C, Vester J, Weckbecker A, Shahgaldian P, De Carvalho CCCR, Meskys R, Zanaroli G,
Glöckner FO, Fernández-Guerra A, Thambisetty S, De La Calle F, Golyshina O V., Yakimov MM, Jaeger KE,
Yakunin AF, Streit WR, McMeel O, Calewaert JB, Tonné N, Golyshin PN. 2019. Decoding the ocean's
microbiological secrets for marine enzyme biodiscovery. FEMS Microbiol Lett 366:1–7.

3. Schloss PD, Handelsman J. 2006. Toward a census of bacteria in soil. PLoS Comput Biol 2:0786–0793.

4. Handelsman J. 2005. Metagenomics: Application of genomics to uncultured microorganisms. Microbiol
Mol Biol Rev 69:195–195.

5. Barzkar N, Sohail M, Tamadoni Jahromi S, Gozari M, Poormozaffar S, Nahavandi R, Hafezieh M. 2021.
Marine bacterial esterases: Emerging biocatalysts for industrial applications. Appl Biochem Biotechnol
193:1187–1214.

6. Martínez-Martínez M, Alcaide M, Tchigvintsev A, Reva O, Polaina J, Bargiela R, Guazzaroni ME, Chicote
Á, Canet A, Valero F, Eguizabal ER, Guerrero M del C, Yakunin AF, Ferrer M. 2013. Biochemical diversity
of carboxyl esterases and lipases from lake arreo (Spain): A metagenomic approach. Appl Environ
Microbiol 79:3553–3562.

7. Jaeger KE, Eggert T. 2002. Lipases for biotechnology. Curr Opin Biotechnol 13:390–397.

8. Jochens H, Hesseler M, Stiba K, Padhi SK, Kazlauskas RJ, Bornscheuer UT. 2011. Protein engineering of
α/β-hydrolase fold enzymes. ChemBioChem 12:1508–1517.

9. De Godoy Daiha K, Angeli R, De Oliveira SD, Almeida RV. 2015. Are lipases still important biocatalysts? A
study of scientific publications and patents for technological forecasting. PLoS One 10:1–20.

10. Bauer TL, Buchholz PCF, Pleiss J. 2020. The modular structure of α/β-hydrolases. FEBS J 287:1035–1053.

11. Ferrer M, Martínez-Martínez M, Bargiela R, Streit WR, Golyshina O V., Golyshin PN. 2016. Estimating the
success of enzyme bioprospecting through metagenomics: Current status and future trends. Microb
Biotechnol 9:22–34.

12.   Placido A, Hai T, Ferrer M, Chernikova TN, Distaso M, Armstrong D, Yakunin AF, Toshchakov S V., Yakimov

678   MM, Kublanov I V., Golyshina O V., Pesole G, Ceci LR, Golyshin PN. 2015. Diversity of hydrolases from

679   hydrothermal vent sediments of the Levante Bay, Vulcano Island (Aeolian archipelago) identified by

680   activity-based metagenomics and biochemical characterization of new esterases and an

681   arabinopyranosidase. Appl Microbiol Biotechnol 99:10031–10046.

682   13.   Tchigvintsev A, Tran H, Popovic A, Kovacic F, Brown G, Flick R, Hajighasemi M, Egorova O, Somody JC,

683   Tchigvintsev D, Khusnutdinova A, Chernikova TN, Golyshina O V., Yakimov MM, Savchenko A, Golyshin

684   PN, Jaeger KE, Yakunin AF. 2015. The environment shapes microbial enzymes: five cold-active and salt-

685   resistant carboxylesterases from marine metagenomes. Appl Microbiol Biotechnol 99:2165–2178.

686   14.   Lopez-Lopez O, Cerdan ME, Gonzalez Siso MI. 2014. New Extremophilic Lipases and Esterases from

687   Metagenomics. Curr Protein Pept Sci 15:445–455.

688   15.   Lee MH, Hong KS, Malhotra S, Park JH, Hwang EC, Choi HK, Kim YS, Tao W, Lee SW. 2010. A new esterase

689   EstD2 isolated from plant rhizosphere soil metagenome. Appl Microbiol Biotechnol 88:1125–1134.

690   16.   Faoro H, Glogauer A, Souza EM, Rigo LU, Cruz LM, Monteiro RA, Pedrosa FO. 2011. Identification of a

691   new lipase family in the Brazilian Atlantic Forest soil metagenome. Environ Microbiol Rep 3:750–755.

692   17.   Hult K, Berglund P. 2007. Enzyme promiscuity: mechanism and applications. Trends Biotechnol 25:231–

693   238.

694   18.   Alcaide M, Tornés J, Stogios PJ, Xu X, Gertler C, Leo RDI, Bargiela R, Lafraya Á, Guazzaroni ME, López-

695   Cortés N, Chernikova TN, Golyshina O V., Nechitaylo TY, Plumeier I, Pieper DH, Yakimov MM, Savchenko

696   A, Golyshin PN, Ferrer M. 2013. Single residues dictate the co-evolution of dual esterases: MCP

697   hydrolases from the α/β hydrolase family. Biochem J 454:157–166.

698   19.   Martínez-Martínez M, Coscolín C, Santiago G, Chow J, Stogios PJ, Bargiela R, Gertler C, Navarro-

699   Fernández J, Bollinger A, Thies S, Méndez-García C, Popovic A, Brown G, Chernikova TN, García-Moyano

700   A, Bjerga GEK, Pérez-García P, Hai T, Del Pozo M V., Stokke R, Steen IH, Cui H, Xu X, Nocek BP, Alcaide M,

701   Distaso M, Mesa V, Peláez AI, Sánchez J, Buchholz PCF, Pleiss J, Fernández-Guerra A, Glöckner FO,

702   Golyshina O V., Yakimov MM, Savchenko A, Jaeger KE, Yakunin AF, Streit WR, Golyshin PN, Guallar V,

703   Ferrer M. 2018. Determinants and prediction of esterase substrate promiscuity patterns. ACS Chem Biol

704   13:225–234.

20.   Höppner A, Bollinger A, Kobus S, Thies S, Coscolín C, Ferrer M, Jaeger KE, Smits SHJ. 2021. Crystal

structures of a novel family IV esterase in free and substrate-bound form. FEBS J 288:3570–3584.

21.   Kumar PP, Paramashivappa R, Vithayathil PJ, Rao PVS, Rao AS. 2002. Process for isolation of cardanol

from technical cashew (*Anacardium occidentale* l.) Nut shell liquid. J Agric Food Chem 50:4705–4708.

22.   Reyes-Duarte D, Ferrer M, García-Arellano H. 2012. Functional-based screening methods for lipases,

esterases, and phospholipases in metagenomic libraries. Lipases and Phospholipases 861:101–113.

23.   Arpigny JL, Jaeger KE. 1999. Bacterial lipolytic enzymes: Classification and properties. Biochem J

343:177–183.

24.   Yoshida S, Hiraga K, Takehana T, Taniguchi I, Yamaji H, Maeda Y, Toyohara K, Miyamoto K, Kimura Y, Oda

K. 2016. A bacterium that degrades and assimilates poly(ethylene terephthalate). Sci (American Assoc

Adv Sci 351:1196–1199.

25.   Bollinger A, Thies S, Knieps-Grünhagen E, Gertzen C, Kobus S, Höppner A, Ferrer M, Gohlke H, Smits SHJ,

Jaeger KE. 2020. A novel polyester hydrolase from the marine bacterium *Pseudomonas aestusnigri* –

Structural and functional insights. Front Microbiol 11:1–16.

26.   Pereira MR, Maester TC, Mercaldi GF, de Macedo Lemos EG, Hyvönen M, Balan A. 2017. From a

metagenomic source to a high-resolution structure of a novel alkaline esterase. Appl Microbiol

Biotechnol 101:4935–4949.

27.   Holm L. 2020. Using Dali for Protein Structure Comparison. Struct Bioinforma 2112:29–42.

28.   Ding Y, Nie L, Yang XC, Li Y, Huo YY, Li Z, Gao Y, Cui HL, Li J, Xu XW. 2022. Mechanism and structural

insights into a novel esterase, E53, isolated From *Erythrobacter longus*. Front Microbiol 12:1–12.

29.   Palm GJ, Fernández-Álvaro E, Bogdanović X, Bartsch S, Sczodrok J, Singh RK, Böttcher D, Atomi H,

Bornscheuer UT, Hinrichs W. 2011. The crystal structure of an esterase from the hyperthermophilic

microorganism *Pyrobaculum calidifontis* VA1 explains its enantioselectivity. Appl Microbiol Biotechnol

91:1061–1072.

30.   De Simone G, Menchise V, Alterio V, Mandrich L, Rossi M, Manco G, Pedone C. 2004. The crystal structure

of an EST2 mutant unveils structural insights on the H group of the carboxylesterase/lipase family. J Mol

Biol 343:137–146.

31.   Alonso S, Santiago G, Cea-Rama I, Fernandez-Lopez L, Coscolín C, Modregger J, Ressmann AK, Martínez-

Martínez M, Marrero H, Bargiela R, Pita M, Gonzalez-Alfonso JL, Briand ML, Rojo D, Barbas C, Plou FJ,

Golyshin PN, Shahgaldian P, Sanz-Aparicio J, Guallar V, Ferrer M. 2020. Genetically engineered proteins

with two active sites for enhanced biocatalysis and synergistic chemo- and biocatalysis. Nat Catal 3:319–

328.

32. Bais HP, Weir TL, Perry LG, Gilroy S, Vivanco JM. 2006. The role of root exudates in rhizosphere

interactions with plants and other organisms. Annu Rev Plant Biol 57:233–266.

33. Bertin C, Yang X, Weston LA. 2003. The role of root exudates and allelochemicals in the rhizosphere.

Plant Soil 256:67–83.

34. Hitch TCA, Clavel T. 2019. A proposed update for the classification and description of bacterial lipolytic

enzymes. PeerJ 2019.

35. Giunta CI, Cea-Rama I, Alonso S, Briand ML, Bargiela R, Coscolín C, Corvini PFX, Ferrer M, Sanz-Aparicio

J, Shahgaldian P. 2020. Tuning the properties of natural promiscuous enzymes by engineering their nano-

environment. ACS Nano 14:17652–17664.

36. Cea-Rama I, Coscolín C, Katsonis P, Bargiela R, Golyshin PN, Lichtarge O, Ferrer M, Sanz-Aparicio J. 2021.

Structure and evolutionary trace-assisted screening of a residue swapping the substrate ambiguity and

chiral specificity in an esterase. Comput Struct Biotechnol J 19:2307–2317.

37. Cao H, Nie K, Xu H, Xiong X, Krastev R, Wang F, Tan T, Liu L. 2016. Insight into the mechanism behind the

activation phenomenon of lipase from *Thermus thermophilus* HB8 in polar organic solvents. J Mol Catal

B Enzym 133:S400–S409.

38. Sellek GA, Chaudhuri JB. 1999. Biocatalysis in organic media using enzymes from extremophiles. Enzyme

Microb Technol 25:471–482.

39. Zaks A, Klibanov AM. 1988. Enzymatic catalysis in nonaqueous solvents. J Biol Chem 263:3194–3201.

40. Sharma S, Kanwar SS. 2014. Organic solvent tolerant lipases and applications. Sci World J 2014.

41. Lukashin A V., Borodovsky M. 1998. GeneMark.hmm: New solutions for gene finding. Nucleic Acids Res

26:1107–1115.

42. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997. Gapped BLAST and

PSI-BLAST: A new generation of protein database search programs. Nucleic Acids Res 25:3389–3402.

43. Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE. 2015. The Phyre2 web portal for protein

modeling, prediction and analysis. Nat Protoc 10:845–858.

44. Gupta N, Rathi P, Gupta R. 2002. Simplified para-nitrophenyl palmitate assay for lipases and esterases.

Anal Biochem 311:98–99.

45.   Bowers GN, McComb RB, Christensen RG, Schaffer R. 1980. High-purity 4-nitrophenol: purification,

765   characterization, and specifications for use as a spectrophotometric reference material. Clin Chem

766   26:724–729.

767   46.   Kabsch W. 2010. research papers XDS research papers. Acta Crystallogr Sect D Biol Crystallogr 66:125–

768   132.

769   47.   Evans PR, Murshudov GN. 2013. How good are my data and what is the resolution? Acta Crystallogr Sect

770   D Biol Crystallogr 69:1204–1214.

771   48.   Vagin A, Teplyakov A. 1997. MOLREP: an Automated Program for Molecular Replacement. J Appl

772   Crystallogr 30:1022–1025.

773   49.   Murshudov GN, Vagin AA, Dodson EJ. 1997. Refinement of Macromolecular Structures by the Maximum-

774   Likelihood Method. Acta Crystallogr Sect D, Biol Crystallogr 53:240–255.

775   50.   Emsley P, Lohkamp B, Scott WG, Cowtan K. 2010. Features and development of Coot. Acta Crystallogr

776   Sect D Biol Crystallogr 66:486–501.

777   51.   Liebschner D, Afonine P V., Baker ML, Bunkoczi G, Chen VB, Croll TI, Hintze B, Hung LW, Jain S, McCoy

778   AJ, Moriarty NW, Oeffner RD, Poon BK, Prisant MG, Read RJ, Richardson JS, Richardson DC, Sammito

779   MD, Sobolev O V., Stockwell DH, Terwilliger TC, Urzhumtsev AG, Videau LL, Williams CJ, Adams PD. 2019.

780   Macromolecular structure determination using X-rays, neutrons and electrons: Recent developments in

781   Phenix. Acta Crystallogr Sect D Struct Biol 75:861–877.

782   52.   Tom Burnley B, Afonine P V., Adams PD, Gros P. 2012. Modelling dynamics in protein crystal structures

783   by ensemble refinement. Elife 2012:1–29.

785   Kinetic parameters of EH$_0$ against pNP-esters. The results are the mean ± SD of triplicates.

Substrate[1]	K$_M$ (mM)	V_{max} (μM min^{-1})	k_{cat} (min^{-1})	k_{cat}/K$_M$ (s^{-1} mM^{-1})
p-NP acetate	1.61 ± 0.12	291.50 ± 7.62	41.64 ± 1.09	0.430
p-NP butyrate	1.08 ± 0.18	252.60 ± 13.84	36.09 ± 1.98	0.558
p-NP dodecanoate	1.00 ± 0.28	25.56 ± 1.80	3.65 ± 0.26	0.061

786   [1]Assays were performed in the presence of 5.0-7.5% DMSO/acetonitrile (see Materials and

787   Methods), concentration at which the enzyme 100% of its activity compared to a control

788   without solvent (Table S2).

 Figure Legends

FIG 1. Substrate selectivity of EH_0 against a set of model p-NP esters of different acyl chain lengths. Specific

activities (mean ± SD of triplicates) are shown. Reactions contained 1 mM of the corresponding p-NP esters and

were conducted in the presence of 1-5% DMSO/acetonitrile (see Methods), under standard conditions described

in Materials and Methods. At the solvent concentration used the enzyme 100% of its activity compared to a

control without solvent (Table S2).

FIG 2. Optimal parameters for the activity and stability of purified EH_0. (A) Temperature profile. (B) The thermal

denaturation curve of EH_0 at pH 7.0 was measured by ellipticity changes at 220 nm and obtained at different

temperatures. (C) The pH profile. The maximal activity was defined as 100%, and the relative activity is shown as

the percentage of maximal activity (mean ± SD of triplicates) determined under standard reaction conditions

with p-NP butyrate as the substrate. Graphics were created with SigmaPlot version 14.0 (the data were not fitted

to any model).

FIG 3. A) Topology diagram of EH_0. The catalytic triad is highlighted in violet. B) EH_0 folding, with the cap domain

in yellow and the catalytic domain in teal. The prolines within the N-terminal cap domain are depicted as violet

sticks. C) Superposition of EH_0 and its homologs: E53 (yellow, PDB code 7W8N), Est8 (violet, PDB code 4YPV),

PestE (raspberry, PDB code 3ZWQ) and EstA (slate, PDB code 5LK6). D) Superposition of subunits A and B showing

different conformations of loop $\alpha1$-$\alpha2$, with chain A in green and chain B in blue. Cap/catalytic domains follow

the color code described in B.

FIG 4. A) EH_0 homodimer showing the catalytic triad as sticks (violet). Both subunits are related by a twofold axis

that is perpendicular to the view shown and situates the two active sites on opposite faces of the dimer. B) A

magnified image of the dimeric interface that covers 722.9Å^2 (approximately 5.6% of the total surface), showing

hydrogen bonds (pink) and salt bridges (green). The Cap domain is colored yellow, whereas the catalytic domain

is shown in teal. C) Dimeric interface of the PestE homolog where the interactions are made by $\beta8$ and $\alpha12$. D)

Dimeric interface of EH$_0$, where the interactions are made by α8, β8 and α9. Disposition of catalytic tunnels from

both subunits in PestE (E) and EH$_0$ (F).

FIG 5. A) A tunnel 16.8 deep giving access to the catalytic triad. B) Hydrogen bond network among the residues

making the catalytic triad of EH$_0$ (teal). Movement of loop β3-α3 in EH$_0$ (housing the oxoanion) with respect to

its homologs E53 complexed with (4-nitrophenyl)hexanoate (yellow), Est8 (violet), PestE (raspberry) and EstA

(slate). C) Hydrogen bond network of residues Glu226 and Tyr223 from α7. D) Hydrophobic patch where Trp218

is located. The EH$_0$ cap domain is shown in yellow and the catalytic domain in teal (C and D.)E) All possible

conformations of the region comprising α1-α2 shown by the ensemble refinement of molecule A. Color code

gives chain mobility, from low (blue) to high (red). The opposed scenarios of the region α1-α2, from an "open

conformation" (F, green) to a "closed conformation" (G, slate). The cap domain seen in the crystal structure is

shown in yellow. H) All conformers shown by the ensemble refinement of molecule B. I and J) Superimposition

of EH$_0$ onto E53 complexed with 4-nitrophenylhexanoate (yellow) and EH$_{1AB1}$ in complex with a derivative of

methyl 4-nitrophenylhexylphosphonate (green), showing only the ligands. The catalytic triad of EH$_0$ is shown as

violet sticks, and residues Glu226 and Ile255 are shown as gray sticks. The EH$_0$ cap domain is shown in yellow and

the catalytic domain in teal. K) Active site cavity of EH$_0$ with acyl and alcohol moieties. Residues from the cap

domain are colored yellow, and residues from the catalytic domain are colored teal. The catalytic triad is shown

in violet.

FIG 6. A) Multiple sequence alignment of EH$_0$ and its homologs E53, Est8, PestE and EstA. Highlighted in boxes

are the conserved motif G-X-S-X-G (in blue), including the catalytic serine, and the D-P-X-X-D (in green), including

the catalytic aspartic acid. The prolines from loop α1-α2 (Pro21, Pro23 and Pro31) and the potential hinges (Pro46

and Pro47) are underlined in purple. The catalytic triad residues are shown with yellow triangles.

FIG 7. A) Superposition of EH$_0$ (teal), EH$_{1AB1}$ (purple) and EH$_3$ (orange). Most differences are found in the cap

domain. Catalytic cavities of EH$_{1AB1}$ (B), EH$_3$ (C) and EH$_0$ (D).

FIG 2.

FIG 6.