Spectral slope and neural complexity as robust markers of task demand and brain state during sleep and wakefulness

Christopher Höhn[1,2], Michael A. Hahn[3], Janna D. Lendner[3] & Kerstin Hoedlmoser[1,2]

[1] Laboratory for Sleep, Cognition and Consciousness Research, Department of Psychology, University of Salzburg, Salzburg, Austria

[2] Centre for Cognitive Neuroscience Salzburg (CCNS), University of Salzburg, Salzburg, Austria

[3] Hertie-Institute for Clinical Brain Research, University Medical Center Tübingen, Tübingen, Germany

Abstract

Spectral slope and neural complexity are affected in many neurophysiological disorders such as ADHD, autism or epilepsy and are modulated by sleep, anesthesia, and aging. Yet, the relationship between these two parameters remains unclear. Here, we evaluated the effects of sleep stage and task demands on spectral slope and neural complexity within a narrow- (30 – 45Hz) and broadband (3 – 45Hz) frequency range in 28 healthy male adults (21.54 ± 1.90 years) over three consecutive recordings with a set of different tasks (resting, attention and memory). We show that the slope steepens, and complexity decreases from wakefulness to N3 sleep. Importantly, slope and complexity are not only modulated by sleep but also differ between tasks, with flatter slopes and higher complexity being associated with more demanding tasks. While the slope and complexity are strongly correlated within 3 – 45Hz, we observe a functional dissociation in the 30 – 45Hz range. Critically, only the narrowband slope is steepest during REM sleep and associated with better task performance, especially in a Go/Nogo task. Our results demonstrate that both markers are powerful indices of sleep depth, task demand and cognitive performance. However, depending on the frequency range, they provide distinct information about the underlying brain state.

Introduction

To date, neural oscillations are still the most prominent electrophysiological signature of human brain activity. For instance, wakeful resting is typically characterized by pronounced alpha-band activity (8 – 12Hz), which is suppressed during active task engagement (Kirstein, 2007; Klimesch et al., 1993; Klimesch, 1999). During sleep, different stages are best described by characteristic oscillatory events like sleep spindles and slow oscillations (Davis et al., 1938; Richard et al., 2012; Terzano et al., 2002). However, recent evidence suggests that irregular and aperiodic brain activity by means of neural complexity (Lempel & Ziv, 1976 and Welch, 1984) and the spectral exponent ß (i.e., the magnitude of decay in power with increasing frequency; He, 2014), also carries meaningful information about different brain states. Specifically, the spectral exponent has been discussed as a marker of the brain's excitation and inhibition (E/I) balance (Gao et al., 2017), which is impaired in a variety of clinical conditions such as the attention deficit hyperactivity disorder (ADHD, Karalunas et al., 2022; Robertson et al., 2019), autism (Gao & Penzes, 2015; Rubenstein & Merzenich, 2003) and epilepsy (Symonds, 1959; Wong, 2010). In addition, epilepsy has further been associated with alterations in neural complexity (Aarabi & He, 2012; Zhu et al., 2017).

Conceptually, the spectral exponent and neural complexity are regarded as two distinct measures as they are derived by different analytical approaches from the underlying electrophysiological signal. Neural complexity computed as Lempel-Ziv-Welch complexity (Welch, 1984) expresses the regularity and compressibility of a signal in time-domain (Lau et al., 2022) and is thought to be still strongly influenced by oscillatory activity (González et al., 2022; Tosun et al., 2019). In contrast, the spectral exponent reflects the absolute value of the slope (i.e., steepness) of a signal's power spectrum in the frequency-domain, which is thought to be mainly aperiodic (Donoghue et al., 2020). In the following, we always use and refer to the spectral slope instead of the spectral exponent (which would be the absolute value of the slope) in order to avoid any ambiguity due to different terms that are often used to describe the same parameter (e.g., 1/f signal and scale free or aperiodic activity).

Despite the apparent differences between the spectral slope and neural complexity, the literature suggests that both can capture changes in brain states in a surprisingly similar fashion. Regarding consciousness and sleep, multiple studies showed that the spectral slope steepens (i.e., becomes more negative) during anesthesia compared to wakefulness (Colombo et al., 2019; Gao et al., 2017; Lendner et al., 2020; Waschke et al., 2021), while others showed the same pattern for neural complexity, which also decreases from wakefulness to anesthetized states (Ferenets et al., 2007; Zhang et al., 2001). This mirrors findings of the transition from wakefulness to sleep, where spectral slope (Lendner et al., 2020; Ma et al., 2018; Miskovic et al., 2019; Pereda et al., 1998) and neural complexity (Andrillon et al., 2016; Schartner et al., 2017) both decrease with increasing sleep depth (i.e., from wakefulness to N3 sleep). Besides changes in consciousness, recent evidence from Waschke et al. (2021) further suggests that the spectral slope can even track the level of attention, whereby higher levels of attention and quicker response times are indexed by flatter slopes. This is in line with findings from other studies, which showed that the slope is indicative of cognitive processing speed (Ouyang et al., 2020; Pathania et al., 2022) and modulated by cognitive decline in ageing (Dave et al., 2018; Voytek et al., 2015; Voytek & Knight, 2015). Interestingly, Mediano et al. (2020) recently showed that higher neural complexity values also relate to faster reaction times on a trial-by-trial basis, thus likewise serving as a proxy of attention or processing speed.

With respect to the influence of different frequency contents on the estimation of the spectral slope and neural complexity, no optimal frequency settings are established yet for any of the two parameters. The heterogeneity of frequency content on which the calculations of both measures are based might be responsible for some disparate results in the current literature, thus further hampering our understanding of the contribution of aperiodic brain activity to healthy brain functioning. For instance, González et al. (2022) suggest that particularly for neural complexity, lower frequencies (≤ 12Hz) are more informative than higher frequencies when differentiating between sleep and wakefulness. For the estimation of the spectral slope, researchers have argued either in favor of broadband (Karalunas et al., 2022; Podvalny et al., 2015; Waschke et al., 2021) or narrowband (Gao et al., 2017; Lendner et al., 2020) frequency ranges. While broadband ranges encompass more of the total signal power and result in better overall slope-fits (Donoghue et al., 2020; Gerster et al., 2022), narrowband ranges are less affected by low-frequency oscillatory activity and are therefore reflecting mostly pure aperiodic activity (Gao et al., 2017; Lendner et al., 2020).

Taken together, the slope and complexity research findings suggest a functional overlap of
neural complexity and spectral slope in tracking different brain states. However, to date a direct
comparison of these two measures is still missing and their actual relationship is unclear. Additionally,
little is known about the sensitivity of the spectral slope and neural complexity to changes in brain activity
during wakefulness. As potential markers of arousal and attention, the two parameters might likely be
additionally affected by varying levels of task demands, which require different amounts of cognitive
resources. Finally, it is unclear how the two measures are affected by selecting different frequency
contents for their calculation.

Here, we leverage an expansive, within-subject design with multiple sleep and wake recordings
to investigate (1) whether the spectral slope and neural complexity are modulated by different brain
states during sleep and wakefulness and (2) to what extent they are related to each other as well as
their functional significance for cognition. Using multiple recording sessions per subject, we try to
overcome a limitation of most previous research that only relied on single session recordings, thus,
limiting insights into the robustness of the observed effects. First, we assess the performance of the
spectral slope and neural complexity in delineating sleep from wakefulness. Second, we investigate the
influence of task demands on both measures by increasing task difficulty form simple resting sessions
to an auditory attention task (Go/Nogo) and a declarative memory task. Third, we analyze the
relationship between the spectral slope and neural complexity across brain states and tasks using either
narrow- or broadband frequency ranges for estimation. Finally, we probe whether the two parameters
track behavioral performance in an auditory Go/Nogo and declarative memory task.

Results

We utilized the data from a previously reported study (Höhn et al., 2021; Schmid et al., 2021)
that investigated the effects of different light conditions on alertness, sleep and memory consolidation.
The subjects underwent the same experimental protocol on three different days under highly controlled
and standardized lighting conditions. On three consecutive experimental nights, multiple tasks were
conducted before and after sleep, including two resting sessions with either eyes closed or open, an
auditory attention task (Go/Nogo) and a declarative memory task (cf., Figure 1A). We calculated spectral
slopes and neural complexity (using Lempel-Ziv complexity) for all sleep stages and tasks in a narrow-
(30 – 45Hz) and broadband (3 – 45Hz) frequency range (cf., Figure 1B and C).

Figure 1. (A): Overview of the experimental protocol. EEG was recorded throughout all tasks and during sleep (with full-night polysomnography) on the experimental days 7, 10 and 13. The tasks, which are highlighted by a dashed, dark-green rectangle were primarily used to analyze the effect of task demand. The adaptation session only served adaptation purposes and was not considered in any of the analyses. Results from the entrance examination questionnaires are presented in Supplementary file – Table 1. (B): Example of the spectral slope estimation during N1 sleep. For illustration purposes, data is shown for the electrode Pz and was averaged over all subjects and sleep recordings. The spectral slope was fitted within 3 – 45Hz (broadband, dashed green line) and 30 – 45Hz (narrowband, dashed red line). (C): Schematic overview of the neural complexity calculation based on a random 4s epoch from electrode Pz of a subject during resting with closed eyes. First, the raw signal, filtered within the broad- or narrowband frequency range (in this case within the 3 – 45Hz broadband range), is Hilbert transformed. Second, the resulting data is binarized around its median amplitude and stored as a vector of zeros and ones. Lastly, the Lempel-Ziv-Welch algorithm (Welch, 1984) is applied on this binary sequence in order to obtain a neural complexity value, which is driven by the number of unique repetitions of ones and zeros.

Spectral slope and neural complexity delineate brain states during sleep

First, we strived for replicating previous findings, which showed that sleep stages could be differentiated solely based on the spectral slope and neural complexity. The effect of sleep stage was assessed for the spectral slope and neural complexity in each frequency range (30 – 45Hz and 3 – 45Hz) with semi-parametric Wald-Type Statistics (WTS; Friedrich et al., 2019) averaged over all electrodes while considering the three repeated measurements.

The narrowband (30 – 45Hz) spectral slope model (*WTS* (4) = 133.57, *p* < .001) and the neural complexity model (*WTS* (4) = 14.11, *p* = .030) both indicated significant modulations by sleep stage. In line with previous research, the narrowband slope was significantly steeper in all sleep stages compared

to wakefulness with the steepest slope during REM sleep. In contrast, the narrowband neural complexity
slightly increased from wake to sleep and showed a diverging pattern in comparison to the spectral
slope (see Figure 2). When the broadband (3 – 45Hz) frequency range was used for estimation, the
effect of sleep stage was much more pronounced in both parameters (spectral slope: WTS (4) = 560.01,
p < .001; neural complexity: WTS (4) = 601.92, p < .001). Both, the broadband slope and complexity
significantly decreased from shallow (N1) to deep NREM sleep (N3). For REM sleep, however, both
markers increased again in remarkably similar ways (see Figure 3), arguably reflecting more wake-like
brain activity in the broadband range.

We found no significant effects of the repeated measurements (all $p_{adj.}$ ≥ .166 after correcting
for multiple comparisons), revealing that the effect of sleep stage robustly emerged in all individual
recordings per subject. To evaluate the topographical distribution of the spectral slope and neural
complexity, we additionally ran a multivariate pattern analysis (MVPA) with multi-class linear
discriminant analyses (LDA). With this MVPA, we quantified how well the sleep stages could be decoded
by taking the topographical distribution of the slope and complexity values into account. In both
frequency ranges and for both parameters, classification accuracies were always significantly above
chance level (20%, p < .001) and in general higher for the broadband (3 – 45Hz) than for the narrowband
(30 – 45Hz) frequency range (WTS (1) = 643.36, p < .001). For both frequency ranges, the spectral
slope was much more informative about the underlying brain state (i.e., yielded higher classification
accuracies) than the neural complexity (WTS (1) = 123.88, p < .001), especially in the narrowband range
(spectral slope: 31.83%, neural complexity: 22.67%; see Figure 3 – Figure Supplement 1).

Figure 2. Spectral slope (green, A) and neural complexity (purple, B) from 30 – 45Hz across sleep, averaged over all lab-sessions per subject. Center figures show the data averaged over all electrodes and topographical maps are provided below (color-coding refers to z-values of slope or complexity computed from the grand average across all sleep stages). In (A), the power spectra in log-log space are provided for each sleep stage to illustrate the narrowband slope differences across different sleep stages. Classification accuracies are shown on the right-hand side. A: The spectral slope decreases from wakefulness across all sleep stages to REM sleep with a small temporary increase during N3 sleep. B: Neural complexity increases from shallow N1 to light N2 sleep and is in general much less modulated by sleep stages than the spectral slope. ***: $p < .001$, **: $p \leq .010$, *: $p \leq .050$, n.s.: $p > .050$; all p-values are adjusted for multiple comparisons; error-bars represent 95% confidence intervals ($N = 27$).

Figure 3. Spectral slope (green, A) and neural complexity (purple, B) from 3 – 45Hz across sleep, averaged over all lab-sessions per subject. Center figures show the data averaged over all electrodes and topographical maps are provided below (color-coding refers to z-values of slope or complexity computed from the grand average across all sleep stages). In (A), the log-log power spectra for each sleep stage are provided to illustrate the broadband slope differences across sleep stages. Classification accuracies are shown on the right-hand side. (A): Spectral slope steepens from wakefulness to N3 sleep but flattens to some extent during REM sleep. (B): Neural complexity shows the same pattern as the spectral slope and likewise decreases from wakefulness to N3 with a subsequent increase in REM sleep. ***: $p < .001$, **: $p \leq .010$, *: $p \leq .050$, n.s.: $p > .050$; p-values are adjusted for multiple comparisons; error-bars represent 95% confidence intervals ($N = 27$).

Figure 3: Supplement 1. Direct comparison of the classification accuracies across sleep between neural complexity and spectral slope for the narrow- (30 – 45Hz) and broadband (3 – 45Hz) frequency ranges.

Spectral slope and neural complexity vary as a function of task demand

Next, we investigated whether spectral slope and neural complexity track varying levels of task demand. We calculated both markers from resting sessions with eyes closed (REC) and eyes open (REO), an auditory Go/Nogo task (GNG), an encoding session (ENC) from a declarative memory task as well as its according retrieval session (RET). For these analyses, we focused on the task data from the evening recordings (see dashed dark-green rectangle in Figure 1A). Theoretically, task demands should have been comparable between resting sessions with closed and open eyes, whereas the Go/Nogo and encoding tasks were considered to be more demanding since they required active task participation and higher cognitive control. The retrieval task was deemed to be the most challenging as it was necessary to make an additional active memory recall.

In the narrowband frequency range (30 – 45Hz), we observed a significant flattening (i.e., values closer to zero) of the slope (WTS (4) = 56.64, $p < .001$) but a decrease in complexity (WTS (4) = 199.55, $p < .001$) with increasing task demands (see Figure 4). The flattening of the narrowband spectral slope was most pronounced when contrasting resting states with the Go/Nogo (REC vs. GNG: WTS (1) = 21.05, $p_{adj.} < .001$; REO vs. GNG: WTS (1) = 16.53, $p_{adj.} = .001$) and encoding (REC vs. ENC: WTS (1) = 20.73, $p_{adj.} < .001$; REO vs. ENC: WTS (1) = 15.56, $p_{adj.} = .001$) sessions. However, further flattening of the slope was observable during the retrieval task (GNG vs. RET: WTS (1) = 6.44, $p_{adj.} = .021$; ENC vs. RET: WTS (1) = 13.66, $p_{adj.} = .001$). The narrowband neural complexity did not differ between the resting and Go/Nogo sessions (all $p_{adj.} > .110$) but decreased from the Go/Nogo to the encoding session (GNG vs. ENC: WTS (1) = 16.64, $p_{adj.} < .001$) and was lowest during retrieval (GNG vs. RET: WTS (1) = 98.74, $p_{adj.} < .001$, ENC vs. RET: WTS (1) = 31.11, $p_{adj.} < .001$).

When investigating the broadband frequency range (3 – 45Hz), we found that the diverging pattern between spectral slope and neural complexity disappeared and both parameters were increasing (i.e., higher complexity values and flatter slopes) across tasks (Slope: WTS (4) = 101.04, $p < .001$; Complexity: WTS (4) = 80.28, $p < .001$; see Figure 5). In the broadband frequency range, neural complexity also differed between the two resting sessions (eyes closed and eyes open), likely reflecting a strong difference in alpha power (8 – 12Hz) between the two conditions, thus supporting a greater influence of oscillations on estimates of neural complexity. Again, we did not observe any effects of the repeated measurements (all $p_{adj.} \geq .252$).

During wakefulness, the MVPA results indicated an above chance classification performance for all tasks (20%, $p < .001$). Similar to the results during sleep, classification accuracy was higher when using the broadband instead of the narrowband frequency range (WTS (1) = 397.08, $p < .001$). The spectral slope was again more informative in the narrowband range (Slope: 35.03%, Complexity: 25.98%, WTS (1) = 71.93, $p < .001$) while the complexity yielded better results in the broadband range (Slope: 48.10%, Complexity: 52.69%, WTS (1) = 13.61, $p = .002$; see Figure 5 – Figure Supplement 1).

Figure 4. Spectral slope (green, A) and neural complexity (purple, B) from 30 – 45Hz across tasks, averaged over all lab-sessions per subject. Center figures show the data averaged over all channels and topographical maps are provided below (color-coding refers to z-values of slope or complexity computed from the grand average across all tasks). In (A), the log-log power spectra for each task are provided to illustrate narrowband slope differences across tasks. Classification accuracies are shown on the right-hand side. (A): The slope flattens with increasing task demands but does not differ between the resting or the Go/Nogo and encoding sessions. (B): Neural complexity decreases across tasks and is minimal during the retrieval session. ***: $p < .001$, **: $p ≤ .010$, *: $p ≤ .050$, n.s.: $p >$ .050; p-values adjusted for multiple comparisons; error-bars show 95% confidence intervals ($N = 28$).

Figure 4: Supplement 1. Slope and complexity from 30 – 45Hz across tasks averaged over all timepoints.

Figure 5. Spectral slope (green, A) and neural complexity (purple, B) from 3 – 45Hz across tasks, averaged over
all lab-sessions per subject. Center figures show the data over all channels and topographical maps are provided
below (color-coding refers to z-values of slope or complexity computed from the grand average over all tasks). In
(A), the log-log power spectra for each sleep stage are provided to illustrate broadband slope differences across
tasks. Classification accuracies are shown on the right-hand side. (A): The slope flattens from the resting to the
Go/Nogo sessions and is flattest during retrieval. (B): Neural complexity increases already from resting with closed
to open eyes and is further elevated in all active tasks, peaking during retrieval. ***: $p < .001$, **: $p \leq .010$, *: $p \leq$
.050, n.s.: $p > .050$; p-values adjusted for multiple comparisons; error-bars show 95% confidence intervals ($N = 28$).
Figure 5: Supplement 1. Comparison of the classification accuracies across tasks between neural complexity and
spectral slope for the narrow- (30 – 45Hz) and broadband (3 – 45Hz) frequency ranges.
Figure 5: Supplement 2. Slope and complexity from 3 – 45Hz across tasks averaged over all timepoints.
Figure 5: Supplement 3. Spectral slope and neural complexity between tasks after correcting for the resting eyes
open condition as baseline.

To control whether the results were confounded by the task order and thus solely reflect an
increase in exhaustion or decrease in motivation, we repeated the analyses with the task data averaged
over all available time points (cf., Figure 1A for all time points at which each task was conducted). These
control analyses confirmed the same patterns as in the original analyses with a very similar flattening of
the broad- and narrowband spectral slopes across tasks and an increase in broadband complexity but
decrease in narrowband complexity with increasing task demands (see Figure 4 – Figure Supplement
1 and Figure 5 – Figure Supplement 2).
An overview of the pairwise classification accuracies for all sleep stage and task pairings is
presented in Supplementary file – Tables 2 and 3. All tasks and sleep stages could be differentiated
above chance-level (50% in this context) and as described above, the classification accuracy was in
general higher for the broadband than the narrowband frequency range. However, in the narrowband
frequency range the accuracies for spectral slope were consistently higher than for neural complexity.
Collectively, the results so far suggest that the spectral slope and neural complexity are both
sensitive markers, which can track brain state changes during sleep and wakefulness due to changes
in sleep stage or because of varying levels of task demand and cognitive load. However, while the two
parameters are modulated in remarkably similar ways when using a broadband frequency range (3 –
45Hz), they express diverging patterns when a restricted narrowband frequency range (30 – 45Hz) is
used. Therefore, we next assessed the relationship between spectral slope and neural complexity.

Relationship between the spectral slope and neural complexity

First of all, we assessed the robustness of the spectral slope and neural complexity estimations
over the different recordings per subject. We correlated each parameter (in the narrow- and broadband
frequency range) with itself between the different lab-sessions for each sleep stage and task. Between
all lab-sessions, the parameters were strongly positively correlated, indicating a substantial overlap of
information over the different recordings (see Supplementary file - Table 4). To identify the relationship
between the spectral slope and neural complexity for each of the two frequency ranges, we further
computed the correlations between the two parameters. In the broadband frequency range, the slope
and complexity were consistently positively correlated across all sleep stages and tasks (see Figure 6A
and B, right columns). However, this relationship vanished in the narrowband frequency range where
the correlations between the two parameters were inconsistent and ranged from significant negative to
positive correlations (see Figure 6A and B, left columns). These results imply that the two parameters
do not share a lot of information in the narrowband range. In contrast, the information is almost entirely
redundant in the broadband frequency range. This fits well to our previous results (cf., Figures 2 – 5)
where only the narrowband slope and complexity were differentially modulated by sleep stage and task
demand.
Taken together, this suggests that the narrowband spectral slope and neural complexity actually
track different features of brain activity that are only explicitly captured when using a restricted frequency
range as for instance 30 – 45Hz. In broader frequency ranges, the dominance of other, especially lower
frequencies might blur these effects, thus making them indistinguishable.

Figure 6. Summary of correlations between the spectral slope and neural complexity from 30 – 45Hz and 3 – 45Hz. The sleep (A) and task (B) data per subject were averaged across all lab-sessions; for task data, only the evening assessments highlighted by the dashed dark-green rectangle in Figure 1 were considered. Significant correlations ($p ≤ .050$ after correcting for false discovery rate) are highlighted with a cross on the topographical maps (color codes for the size and directionality of the correlation coefficients). The 30 – 45Hz slope and neural complexity showed no consistent positive or negative relationship across tasks and sleep stages. In contrast, the 3 – 45Hz slope and neural complexity were consistently positively correlated over all tasks and sleep stages ($N = 28$).

Figure 6: Supplement 1. Correlation of the slope and complexity with themselves in the narrow- or broadband frequency range during sleep (A) and wakefulness (B). Only the spectral slope was consistently positively correlated with itself, whereas the complexity was slightly negatively correlated with itself between the two frequency ranges.

Lastly, we assessed how strongly the spectral slope and neural complexity were correlated with themselves in the different frequency ranges. The narrow- and broadband slope were always positively correlated, whereas the opposite was true for neural complexity (see Figure 6 – Figure Supplement 1). Thus, flatter narrowband slopes were usually associated with flatter slopes in the broadband range, but lower narrowband complexity were often even associated with higher broadband complexity. This suggests that especially the narrowband spectral slope might measure a facet of the underlying brain activity that is not represented in the narrowband neural complexity.

The spectral slope as an electrophysiological marker of task performance

Having established that spectral slope and neural complexity are not only modulated by sleep stage but also differ between tasks during wakefulness in a frequency range specific manner, we next investigated their relationship with task performance. Thus, we correlated the spectral slope and neural complexity from the narrow- and broadband frequency ranges during the Go/Nogo task with the

according performance scores (percentage of correct trials divided by median reaction time) over multiple sessions. Again, this allowed us to test the robustness of any correlations with behavior. Only flatter slopes in the narrowband range (30 – 45Hz) were consistently related to better task performance (see Figure 7). Neural complexity, on the other hand, did not correlate with performance, neither in the narrow- nor in the broadband (3 – 45Hz) range (see Figure 7 and Figure 7 – Figure Supplement 1). In the broadband range, the relationship with task performance was still consistently positive for both parameters but did not reach statistical significance. The fact that this positive relationship was strengthened and actually turned significant only for the slope in the narrowband range again suggests that especially this narrowband slope can be interpreted as a distinct marker of task performance.

Figure 7. Relationship between Go/Nogo task performance and spectral slope (A) or neural complexity (B) within 30 – 45Hz across different assessment times. For the large scatterplots, the data was averaged across all lab-sessions (small scatterplots show the relationship in each lab-session). The topoplots depict the strength of the correlation for each electrode. Electrodes forming a significant cluster are highlighted with asterisks. Those showing a significant correlation after false discovery rate correction but did not from a significant cluster are marked with a cross. Only the narrowband spectral slope showed a consistent positive relationship with task performance (*N* = 26).

Figure 7: Supplement 1. Results when using the broadband (3 – 45Hz) frequency range. No significant relationships emerged for the spectral slope and neural complexity, even though the correlation was consistently positive for both parameters.

Next, we determined whether the narrowband spectral slope can even be used to track memory performance. Therefore, we correlated the spectral slope and neural complexity during the retrieval sessions of a declarative memory task with the recall performance scores (i.e., percentage of correctly recalled word pairs). Even though the overall pattern was similar to the Go/Nogo task, most correlation coefficients only showed a trend towards statistical significance (see Figure 8). Despite the lack of statistical significance on most electrodes, the narrowband spectral slope was again consistently positively correlated with recall performance. This indicates that flatter slopes, especially in the narrowband frequency range, are not only related to better attentional performance but might also benefit declarative memory. In contrast, the narrowband complexity was not positively correlated with memory performance and even expressed a negative relationship on some electrodes. Since we observed a positive relationship between overnight decreases in resting state slopes and memory performance in another study (Lendner et al., 2022), we further assessed whether the overnight change in slope during the retrieval task is also correlated with sleep-dependent memory consolidation. However, we did not obtain a significant relationship, indicating that while flatter slopes during the retrieval were associated with slightly better memory performance in the according session, overnight changes in the slope or complexity were not related to performance changes.

Figure 8. Relationship of declarative memory recall performance and the spectral slope (A) or neural complexity (B) within 30 – 45Hz. Results are shown for the immediate recall during the evening and the delayed recall in the

next morning as well as for the overnight change. For the large scatterplots, the data was averaged across all lab-sessions (small scatterplots show the relationship for each session). The topoplots represent the strength of the correlations on each electrode and color codes for the size and directionality of the correlation coefficients. Even though the spectral slope was consistently positively correlated with recall performance, no electrodes formed a significant cluster. Significant single electrodes that survived false discovery rate correction are highlighted with a cross (N = 28).

Figure 8: Supplement 1. Results when using the broadband 3 – 45Hz frequency range. No relationship observable between recall performance and the slope or complexity.

In the broadband frequency range, both parameters did not show a consistent relationship with recall performance (see Figure 8 – Figure Supplement 1). We analyzed whether the similar results between the Go/Nogo and declarative memory task performance could be traced back to better overall attention and higher task engagement but there was no significant relationship between the performance scores from the two tasks (evening: *rho* = 0.10, *p* = .611; morning: *rho* = 0.06, *p* = .766). Thus, subjects that performed well in the Go/Nogo task did not necessarily achieve a high recall performance score in the declarative memory task.

Discussion

In this study comprising three experimental recordings with multiple measurements per subject, we demonstrated that the spectral slope and neural complexity both reliably delineate sleep stages and are modulated by task demand during wakefulness. Critically, we provided evidence that the correlation between spectral slope and neural complexity strongly depends on the frequency content, which alters their modulation across task demands and sleep stages. The narrowband (30 – 45Hz) spectral slope was best suited to differentiate REM sleep from wakefulness, even though the broadband (3 – 45Hz) slope and neural complexity were more strongly modulated by sleep stages in general. During wakefulness, increasing task demands are associated with flatter slopes in the narrow- and broadband range, but only with higher complexity in the broadband range. Critically, solely the narrowband spectral slope tracked task performance in an auditory attention task (Go/Nogo) as well as in a declarative memory task.

Sleep stage specific alterations of spectral slope and neural complexity

Our findings corroborate previous research which demonstrated that the spectral slope and neural complexity are sensitive markers of sleep depth (Abásolo et al., 2015; Bódizs et al., 2021; Lendner et al., 2020; Pascovich et al., 2022; Schartner et al., 2017; Tagliazucchi et al., 2013). Building upon these findings, we leveraged repeated EEG recordings per subject and confirmed that the two parameters robustly differentiate all sleep stages from wakefulness. Overall, sleep stages could be better delineated when a broad frequency range (3 – 45Hz) was used for calculation of the spectral slope and neural complexity. This is probably due to the fact that the broadband range encompasses the frequencies typically used for traditional sleep staging, such as slow wave activity (0.5 – 4Hz) and sleep spindles (11 – 15Hz; Dijk, 1995), thereby increasing the sleep stage specific information in the underlying signal. However, only the spectral slope from the narrowband frequency range (30 – 45Hz) clearly distinguished REM sleep from all other sleep stages, which is in line with recent findings by

Lendner et al. (2020). This behavior of the narrowband spectral slope contradicted the overall
modulation of the slope and complexity in the broadband range, where both parameters showed a
relative increase during REM sleep. Since REM sleep (sometimes called 'paradoxical sleep'; Peigneux
et al., 2001 or Siegel, 2011) is characterized by wake-like, non-oscillatory brain activity (Blumberg et al.,
2020; Peever & Fuller, 2017), the disparate results between the two frequency ranges suggest that the
narrowband slope mainly measures non-oscillatory, aperiodic brain activity. The relative increase in
complexity during REM sleep has been attributed to higher levels of conscious content that accompany
vivid dreaming and thus require more complex brain activity than deeper, mostly dreamless sleep stages
(Lau et al., 2022; Mateos et al., 2018).

Recent modeling work has also linked especially the narrowband spectral slope with the
excitation to inhibition (E/I) balance in the brain (Gao et al., 2017). Within this framework, steeper slopes
during REM sleep potentially reflect stronger inhibitory brain activity. This might allow the brain to
decouple from its environment and by maintaining muscle atonia to enable the consolidation of
emotional memories and the experience of vivid dreams (Aime et al., 2022) without the danger of acting
them out. The narrowband (30 – 45Hz) complexity, however, expressed a diverging pattern compared
to the narrowband slope and stayed almost constant across all sleep stages with even a slight increase
from N1 to N2 sleep. Even though our study is the first to directly compare spectral slope and neural
complexity during sleep, the congruency of both measures within the broadband frequency range might
not be surprising, since previous studies investigating the parameters individually have shown their
decrease across sleep (Aamodt et al., 2021; Lendner et al., 2020; Miskovic et al., 2019; Pereda et al.,
1998; Schartner et al., 2017). Although we were able to classify sleep stages consistently above chance
level with both parameters, it should be noted that our classifier was trained and tested only on our data.
Furthermore, we did not compare the performance of the spectral slope and neural complexity to other
potentially powerful biomarkers. Therefore, it would be interesting to see how accurate sleep stages can
be scored exclusively by means of the slope or complexity and how these two markers perform in
comparison to other indices of sleep depth like heart rate variability and blood pressure (Kuula &
Pesonen, 2021; Mitsukura et al., 2020; Radha et al., 2019; van de Borne et al., 1994) or accelerometric
data from actigraphy (Lüdtke et al., 2021; Sadeh et al., 1989) and multisensor consumer-wearables
(Ameen et al., 2019; Boe et al., 2019; Roberts et al., 2020; Tal et al., 2017).

Spectral slope and neural complexity are modulated by task demands

In addition to our findings during sleep, we demonstrate that the spectral slope and neural
complexity track different levels of task demands. That the slope and complexity are in general also
modulated during wakefulness is in line with other research (Jacob et al., 2021; Mediano et al., 2021;
Sheehan et al., 2018; Waschke et al., 2021), however, to our best knowledge this is the first study
assessing the effect of task demand and the influence of different frequency ranges on the two
parameters. Similar to sleep, we observed a homogenous modulation of the broadband (3 – 45Hz) slope
and complexity, where flatter slopes and higher complexity were associated with an increase in task
demands. This pattern was identical for the narrowband slope but was inverted for the narrowband
complexity, where higher task demands were accompanied by decreasing levels of complexity. In the
E/I balance framework, flatter narrowband slopes are the result of higher excitation in the brain (Chini et

al., 2022; Gao et al., 2017). Thus, our observed pattern of a flattening of the spectral slope with
increasing task demands seems plausible as more difficult tasks require more cognitive resources and
therefore lead to stronger excitatory brain activity (Harris & Thiele, 2011; He, 2011; Kanashiro et al.,
2017). Unlike Waschke et al. (2021), who reported a stronger occipital flattening of the slope for a visual
compared to an auditory task, we did not observe clear topographical differences between modalities,
even though the attentional Go/Nogo task was entirely auditory except for a fixation-cross whereas the
declarative memory task mainly relied on visual content. However, this lack of topographical
distinctiveness might be due to a partial overlap between involved brain areas since both auditory
discrimination and learning involve frontotemporal brain regions (Ackerman, 1992; Halsband, 1998).

Differential contributions of narrow- and broadband frequency ranges

Based on the results from the broadband frequency range, it is tempting to assume that the
spectral slope and neural complexity are indexing the same or at least very similar features of brain
activity. However, the divergence between the narrow- and broadband slope and complexity during
sleep and wakefulness clearly demonstrates that the two parameters cannot be used interchangeable
but instead track different facets of the underlying brain activity. Here, we reveal that the selected
frequency range dramatically influences the information that the two parameters provide and therefore
also their interrelation. Using a narrowband frequency range from 30 – 45Hz for estimation decreases
the relationship between the spectral slope and neural complexity. During wakefulness, different
contributions of oscillatory and aperiodic brain activity to the slope and complexity could account for their
diverging patterns in the narrowband range. Although it appears paradoxical that flatter narrowband
slopes, representing an increase in aperiodic activity, should be accompanied by a decrease in neural
complexity, others have also reported this type of counterintuitive behavior of neural complexity.
Mediano et al. (2021) showed that in MEG within 0.5 – 30Hz, active tasks actually exhibited lower
complexity values compared to rested wakefulness. In addition, a recent review from Lau et al. (2022)
discussed several studies that reported apparently contradicting modulations of neural complexity in
different clinical conditions, where some report lower and others higher levels of complexity. Thus, the
question whether higher neural complexity can always be clearly interpreted as more complex or
irregular brain activity remains unclear. So far, the best explanation for the contradictory findings in the
neural complexity literature is that higher complexity values can both represent either more complex or
more random systems (La Torre-Luque et al., 2016), which makes it difficult to argue whether higher
complexity always represents a healthier neurophysiological brain state. Interestingly, other studies
showed a strong relationship between different complexity or entropy measures and the spectral slope
(Colombo et al., 2019; Miskovic et al., 2019; Waschke et al., 2017), thus, it would be interesting to
investigate in the future what drives their shared information and under which circumstances (i.e.,
frequency ranges) this relationship vanishes.

The narrowband spectral slope as a unique marker of task performance

When relating the spectral slope and neural complexity to behavioral outcomes, we observed
that only the narrowband slope within 30 – 45Hz correlated with attentional task performance in an
auditory Go/Nogo task across all recordings per subject. Thus, it appears that the narrowband slope

serves as a particularly sensitive marker for task-dependent fluctuations in brain states associated with
behavioral performance. This association between adaptively flatter slopes and better task performance
might even translate to more general cognitive tasks that do not solely rely on attention since we also
observed a consistent positive but weaker relationship with memory performance. In larger scale studies
that rely on databases or in multicenter studies, which commonly have higher statistical power, however,
the broadband slope and complexity were also significantly correlated with task performance. For
instance, Mediano et al. (2021) and Waschke et al. (2021) found an association between task-specific
attention levels and spectral slope or neural complexity in a broader frequency range. As in our study
the correlation between the broadband slope and complexity with the Go/Nogo task performance was
also consistently positive but too weak to reach statistical significance, these findings do not necessarily
contradict our claim that the narrowband spectral slope is even more sensitive to adaptive task-
dependent changes in brain state. In contrast, this shows that lower statistical power might suffice for
the narrowband slope to index robust relationships with behavioral performance.

Limitations

It should be noted that the cognitive tasks were not specifically designed for the analyses of
varying levels of task demand as the data presented here was obtained from a study that was originally
designed for the investigation of short-wavelength light effects on sleep, attention and memory
performance (cf., Höhn et al., 2021 and Schmid et al., 2021). In the future, it might be promising to
contrast tasks that exclusively rely on different cognitive resources, i.e., auditory vs. visual to assess
how spectral slope and neural complexity adapt topographically to different modalities. Even though we
used only 11 scalp electrodes, we still robustly detected the effects of sleep stage and task demand,
providing evidence for the power of the spectral slope and neural complexity as indices of different brain
states. Nevertheless, research with high-density or intracranial EEG setups might further contribute to
the understanding of which cortical areas are most influential in driving changes in the slope or
complexity across different brain states. Finally, we only recruited healthy male adults in a restricted age
range (18 – 25 years) in order to avoid potential sex differences and hormonal effects (Kozhemiako et
al., 2021; Plamberger et al., 2021). Therefore, it is unclear to what extent our results generalize to other
populations. While sex does not necessarily affect the spectral slope or neural complexity when
controlling for overall signal amplitude (Bódizs et al., 2021; Tosun et al., 2019), age does seem to play
an important role in terms of developmental changes in the spectral slope and decorrelation of brain
activity, which begins during early childhood (Chini et al., 2022; Schaworonkow & Voytek, 2021) and
lasts until late adulthood (Dave et al., 2018). While this task-independent flattening of the slope in older
subjects has been associated with decline in cognitive functioning (Voytek et al., 2015), our results
suggest that task-dependent increases in excitation (expressed by flatter slopes) might be beneficial for
behavioral performance. Thus, an adaptive task-specific modulation of the slope in healthy individuals
appears to be associated with better task performance and might index cognitive adaptability.

Conclusion

Taken together, our results demonstrate that the EEG spectral slope and neural complexity are
powerful indices of different brain states during sleep and wakefulness. We provide robust evidence

from multiple recordings of three within-subjects measurements, showing that sleep stages and task

demands are reliably indexed by both the spectral slope and neural complexity. Critically, we show that

the selected frequency range has a strong impact on the interpretability and functional relevance of the

two parameters. When trying to distinguish different brain states, the broadband spectral slope and

neural complexity are more sensitive, however, only the narrowband spectral slope within 30 – 45Hz

turned out to be a powerful index of behavioral performance and best suited to differentiate REM sleep

from wakefulness and all other sleep stages.

Materials and methods

Key resources table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Software, algorithm	Brain Vision Analyzer 2.2	Brain Products GmbH	RRID: SCR_002356	https://www.brainproducts.com
Software, algorithm	Adobe Illustrator CS6	Adobe Inc.	RRID: SCR_010279	
Software, algorithm	Fieldtrip (obob_ownft)	Oostenveld et al., 2011	RRID: SCR_004849	https://gitlab.com/obob/obob_ownft/
Software, algorithm	FOOOF (specparam)	Donoghue et al., 2020		https://fooof-tools.github.io/fooof/
Software, algorithm	ggplot-2	Wickham, 2016	RRID: SCR_014601	https://cran.r-project.org/web/packages/ggplot2/index.html
Software, algorithm	Lempel-Ziv complexity	Comsa, 2019		https://github.com/iulia-m-comsa/EEG/tree/master/Lempel-Ziv%20complexity
Software, algorithm	MANOVA.RM	Friedrich et al., 2019		https://cran.r-project.org/web/packages/MANOVA.RM/index.html
Software, algorithm	MATLAB 2018b	MathWorks Inc.	RRID: SCR_001622	
Software, algorithm	MVPA-light toolbox	Treder, 2020	RRID: SCR_022173	https://github.com/treder/MVPA-Light
Software, algorithm	Psychtoolbox PTB-3	Kleiner et al., 2007	RRID: SCR_002881	http://psychtoolbox.org/
Software, algorithm	RStudio 2021.09	RStudio Team	RRID: SCR_000432	

| Software, | Somnolyzer | Koninklijke | https://www.philips.co.in |
| algorithm | 24 x 7 | Philips N.V. | |

Participants and inclusion criteria

We recorded data from 28 male participants (18 – 25 years; mean age 21.54 ± 1.90 years). Final sample
sizes varied for each analysis between N = 26 – 28 as some participants had missing data for specific
tasks or timepoints (the exact sample size for each analysis is provided in the corresponding results
section). All participants were free of medication and did not suffer from a mental or physiological illness
or sleep problems. They adhered to a regular sleep-wake rhythm (i.e., regular bedtimes with about 8
hours of sleep per night) and refrained from drug abuse and above-average caffeine consumption (more
than three cups of coffee per day) during participation. For screening purposes, all subjects filled in an
entrance questionnaire in which we checked for sleep quality, mood, anxiety, perceived stress level and
chronotype (see Supplementary file – Table 1). Written informed consent was obtained from every
participant and all subjects were remunerated with either 100€ and 16 hours course credit or 50€ and
24 hours course credit. The study was approved by the local ethics committee of the University of
Salzburg (EK-GZ: 16/2014) and conducted in agreement with the Declaration of Helsinki.

Experimental protocol

Study design

Each subject participated over a time span of 14 days, with an entrance examination marking day one
(an outline of the study protocol is presented in Figure 1). From that day on, participants wore an
actigraph (MotionWatch 8; CamNtech Ltd, Cambridge, England) and filled in daily online sleep protocols
(LimeSurvey GmbH, Hamburg, Germany), which we used to check for compliance with a regular sleep-
wake rhythm.
The first recording was scheduled on day four and was implemented only for adaptation purposes in
order to avoid potential first night effects (Browman & Cartwright, 1980; Curcio et al., 2004). After
placement of all EEG, ECG, EMG and EOG electrodes, the participants were familiarized with the
resting and Go/Nogo tasks. Bedtime was scheduled for approximately 11:00 pm and the participants
were woken up 8 hours after lights out and left the laboratory at approximately 9:00 am.
The experimental recordings were scheduled on days 7, 10 and 13. Participants arrived at 6:00 pm and
EEG, ECG, EMG and EOG electrodes were mounted. The recordings started with an initial resting
session (3min eyes closed and 3min eyes open) and the Go/Nogo task (10min), which was followed by
the encoding sessions (two times 14min) of a declarative memory task. Before the first cued recall,
another resting and Go/Nogo session were conducted. Afterwards, the participants had a 1.5 hours
break from the tasks, in which they read stories under different light conditions (for details cf., Schmid
et al., 2021). Before going to bed at approximately 11:00 pm, participants completed the last resting and
Go/Nogo session of the day. After awakening, a morning session of resting and the Go/Nogo task as
well as another recall from the declarative memory task were performed. During all wake-recordings,
daylight mimicking room lights (provided by Emilum GmbH, Oberalm, Austria) were dimmed to 4.5
photopic lux and room temperature was adjusted via air conditioning based on participant's preferences.

Go/Nogo task

To assess objective levels of attention, we implemented an auditory version of the Go/Nogo paradigm (Donders, 1969) via the Psychophysics Toolbox (PTB-3; Kleiner et al., 2007) in MATLAB (Release 2018b, The MathWorks Inc., Natick, MA). Participants were asked to react as quickly as possible with a button press on a response time box (RTBox v5/6; Ohio State University, Columbus, OH) whenever they heard a 'Go' sound and needed to inhibit their reaction when a 'Nogo' sound was played. The task comprised 400 trials with Go sounds being presented in 80% of the trials and Nogo sounds occurring in the remaining 20% of trials (the order of Go and Nogo sounds was randomized each time). The two stimuli used for the Go and Nogo sounds were low- (1000Hz) and high-pitched (1500Hz) tones, which were presented for 50ms with a varying interstimulus interval (1480 – 1880ms). Whether the low- or high-pitched sound represented the Go-signal was determined by chance at the beginning of each session. Participants had to react within 500ms for the response to be considered valid, but responses were recorded until 1000ms post-stimulus with reaction times longer than 500ms being regarded as attentional lapses. From each session, the performance score was computed by dividing the percentage of correct trials by the median reaction time of all valid responses (≤ 500ms, no errors) in milliseconds (Figueiro et al., 2016; Höhn et al., 2021).

Declarative memory task

Participants encoded a set of 80 word pairs on days 7, 10 and 13. To avoid learning effects over time, a different but similarly difficult set of 80 word pairs was presented on each of the three days. The order of the sets was randomized across subjects. Each set was presented twice for 14min during encoding and the data from both encoding sessions was pooled for further analyses. Each word pair was presented for 1500ms and was followed by a fixation-cross for 8500ms. Participants were instructed to encode the word pair as vividly as possible during the presentation of the fixation-cross by imagining a semantic connection between the two words. During the cued recall sessions, only the first word of a pair was presented, and participants were asked to press a button on the response time box as soon as they remembered the second word. Whenever a button was pressed, the participant was instructed to name the missing word and a fixation-cross appeared for 3500ms while the experimenter noted the answer. When no button was pressed, the fixation-cross appeared automatically after 6500ms.

EEG recording and analyses

All electrophysiological data were recorded with a sampling rate of 500Hz via the BrainVision Recorder software (Version 2.11, Brain Products GmbH, 2015) using a 32 channel BrainAmp system (Brain Products GmbH, Munich, Germany). We placed 11 gold-cup electrodes (Grass Technologies, Astro-Med GmbH, Rodgau, Germany) according to the international 10-20 system on the positions: F3, Fz, F4, C3, Cz, C4, P3, Pz, P4, O1 and O2. The average of positions A1 and A2 on the left and right mastoids was used for offline re-referencing as the data were online referenced against Cz. Fpz was used as ground electrode. Additionally, two EMG electrodes were placed on the musculus mentalis for measuring muscle activity during sleep and four EOG electrodes around the eyes to record horizontal

572   and vertical eye movements. ECG was recorded with an electrode on the right clavicular and another

573   electrode on the lowest left costal arch. Impedances were always kept below 10kΩ.

Polysomnography

The time in bed was standardized for all polysomnography recordings and comprised exactly 8 hours.

For sleep staging, the data were first low-pass filtered at 30Hz and re-referenced to contralateral

mastoids with the BrainVision Analyzer software (Version 2.2.0.7383, Brain Products GmbH, 2019).

Physio-channels were referenced in a bipolar manner and the data were down-sampled to 128Hz before

sleep stages were classified for each 30 second epoch with the Somnolyzer 24 x 7 algorithm (Koninklijke

Philips N.V.; Eindhoven, The Netherlands) in accordance with the criteria of the American Academy of

Sleep Medicine (Richard et al., 2012). The results were finally verified by a human expert scorer. The

general sleep architecture of each night is presented descriptively in Supplementary file – Table 5.

EEG preprocessing

In a first step, the data were processed with the BrainVision Analyzer software, and we applied a 0.3Hz

high-pass as well as a 50Hz notch filter. The EEG channels were re-referenced to linked mastoids and

the online reference Cz was restored. We corrected for eye movements with the Gratton & Coles method

(Gratton et al., 1983; only implemented for data during wakefulness) and ran an automatic artifact

detection. The data were then down-sampled to 250Hz and exported for further analyses in MATLAB.

The continuous data were de-trended and subsequently segmented into epochs of 4s for each task and

sleep stage using the fieldtrip toolbox (Oostenveld et al., 2011). To be able to compare all task- and

sleep-data, we decided to set the epoch-length to 4s as this enabled the best tradeoff between a

sufficient number of epochs even for the shortest tasks (3min resting sessions) and sufficient frequency

resolution within 0.5 – 45Hz. All artifact-containing epochs (defined as > 1% being detected as artifact)

were removed for the following analyses. Since the remaining number of clean epochs from the different

tasks (resting, Go/Nogo, encoding and retrieval) and sleep-stages (WAKE, N1, N2, N3 and REM) varied

dramatically due to different recording lengths, we balanced the number of epochs across tasks and

sleep-stages for the multivariate pattern analyses (MVPA) to ensure the validity of the classification

results. In more detail, we set the maximum number of epochs to the highest possible number of epochs

from the shortest task (i.e., 45 epochs as the resting sessions only comprised 3min). To do so, we drew

a random subset of 45 epochs from all data that contained more than 45 clean epochs. For all other

analyses we used all available data to maximize the signal to noise ratio wherever possible (for the

number of epochs used per task and sleep stage see Supplementary file – Table 6).

Spectral Slope

To obtain the spectral slope, we first calculated power-spectra between 0.5 – 45Hz from the

preprocessed, 4s segmented data via the *mtmfft* method in Fieldtrip (Oostenveld et al., 2011) using a

multi-taper approach (1Hz frequency smoothing; Lendner et al., 2020). To extract the spectral slope

information as a maker for arousal, we applied robust linear fits (using the *robust fit* MATLAB function)

in log-log space between 30 – 45Hz based on a previously established method (Lendner et al., 2020).

We decided to use robust linear fits instead of using the *FOOOF* algorithm (alternatively known as

specparam; Donoghue et al., 2020) for the narrowband frequency range since this approach has already
been established to yield a sensitive aperiodic marker of arousal by Lendner et al. (2020) and because
in this frequency range also the *FOOOF* would approximate a linear fit, thus leading to highly comparable
results. However, for the broadband frequency range (3 – 45Hz), we applied the *FOOOF* algorithm to
extract the slope as the linear fits would have been skewed by oscillatory bumps in the power spectrum.

Neural Complexity

We followed previous approaches (Mediano et al., 2021; Schartner et al., 2015) and calculated the
Lempel-Ziv-Welch complexity (Lempel & Ziv, 1976; Welch, 1984) as a proxy for neural complexity per
channel and epoch. To obtain the neural complexity in the same frequency ranges in which we
calculated the spectral slope, we applied additional 3 or 30Hz high-pass and 45Hz low-pass filters to
ensure that the underlying signal contained the same frequencies as for the spectral slope. As Rivolta
et al. (2014) demonstrated that 1000 datapoints are sufficient for reliable Lempel-Ziv complexity
analyses during sleep, we used the same 4s segmented data (which translates to 1000 sampling points
per epoch) for the neural complexity analyses that we used for the spectral slope. We then applied a
Hilbert-transformation on each epoch to obtain the instantaneous amplitude. Afterwards, we binarized
the resulting single epoch data around its median amplitude and transformed it into a binary sequence.
Values of 1 were given for amplitude samples above the median and values of 0 for amplitudes below
(or equal with) the median. This binary sequence of ones and zeros was finally subjected to the Lempel-
Ziv-Welch complexity algorithm (Comsa, 2019) in MATLAB. To make sure that our results were not
affected by different algorithm implementations, we additionally ran the original Lempel-Ziv algorithm
(LZ76, implemented in MATLAB by Thai, 2012) but did not obtain any different results. Thus, only the
results from the Comsa (2019) algorithm are reported.

Statistical analyses

Statistics were calculated in R-Studio (Version 4.1.2.; RStudio Team, 2021). MATLAB functions from
the Fieldtrip toolbox and the ggplot-framework (Wickham, 2016) in R were adapted for data visualization.

Factorial analyses and correlations

All analyses involved three repeated measurements (on days 7, 10 and 13; cf., Figure 1) and therefore
at least two factors (lab-session and task or sleep stage). Since in most cases at least one assumption
for parametrical testing was violated, we decided to compute a more conservative semi-parametrical
analyses with the MANOVA.RM package (Friedrich et al., 2019). For these semi-parametrical factorial
analyses, data were averaged over all EEG electrodes to facilitate interpretation of the results. In the
statistical results, we always refer to the Wald-Type-Statistics (WTS) with empirical p-values obtained
from permutation resampling procedures and 10.000 iterations. Whenever multiple comparisons were
conducted for follow-up testing, p-values were corrected for alpha error inflation with the Benjamini-
Hochberg procedure (Benjamini & Hochberg, 1995).
For correlation analyses, we computed Spearman rho coefficients instead of Pearson correlations
whenever the normality assumption was significantly violated (indicated by Shapiro-Wilk tests) and in

general for all cluster correlations on the whole scalp level. For the cluster corrected correlation
approach, we used the Monte-Carlo method and 10.000 iterations to assess the relationship between
the EEG parameters per channel and the behavioral measures.

Multivariate pattern analyses (MVPA)

Since it is difficult to take topographical patterns into account in classical factorial designs, we computed
multivariate pattern analyses using the MVPA-Light toolbox (Treder, 2020) in MATLAB to further exploit
the information present in the complexity and slope data as patterns across electrodes. For each task
and sleep stage, the complexity and slope from every epoch and electrode was fed to the classifier.
Thus, the single epochs per subject were used for training and testing while the complexity and slope
patterns over electrodes represented the multivariate information. For comparisons between more than
two tasks or sleep stages, multiclass linear-discriminant analyses (LDA) were used and regular LDA for
two-condition comparisons. We calculated classifier accuracies per subject via leave-one-out cross
validation (LOO-CV) to account for the restricted amount of data available for training and testing in our
sample. Since no effects regarding the different lab-sessions emerged, we pooled the data from the
different lab-sessions for each subject to improve the reliability of the MVPA analyses.

Acknowledgements

This research was funded by the Austrian Science Fund (FWF, P32028) and the Centre for Cognitive
Neuroscience Salzburg (CCNS). C.H. further received funding from the Doctoral College "Imaging the
Mind" (FWF; W1233-B). J.D.L. received a grant from the German Research Foundation (DFG LE
3863/2-1). We would like to thank all volunteering subjects for their time and effort. Further, we are very
grateful for the support from Sarah R. Schmid, Selina Schindlmayr, Daniela Niebler, Lucy Matthews,
Marina Thierauf, Leoni Bernstorf, Lorenz Rapp, Henrik Rheinwald and Leonard van Dyck regarding the
data collection process and the recruitment of participants.

Additional Information

Competing interests

The authors do not declare any competing interests.

Funding

Funder	Grant reference number	Author
Austrian Science Fund (FWF)	W 1233-G17	Christopher Höhn
German Research Foundation (DFG)	LE 3863/2-1 HE 8329/2-1	Janna D. Lendner, Michael A. Hahn
Austrian Science Fund (FWF)	P 32028	Kerstin Hoedlmoser
Jung Foundation for Science and Research: Ernst Jung Career Advancement Award, awarded to Randolph Helfrich		Janna D. Lendner, Michael A. Hahn

Ethics

This study was conducted in accordance with the guidelines from the Declaration of Helsinki. Written approval was additionally provided by the local ethics committee of the University of Salzburg (EK-GZ: 16/2014).

References

Aamodt, A., Nilsen, A. S., Thürer, B., Moghadam, F. H., Kauppi, N., Juel, B. E., & Storm, J. F. (2021). Eeg Signal Diversity Varies With Sleep Stage and Aspects of Dream Experience. *Frontiers in Psychology*, *12*, 655884. https://doi.org/10.3389/fpsyg.2021.655884

Aarabi, A., & He, B. (2012). A rule-based seizure prediction method for focal neocortical epilepsy. *Clinical Neurophysiology*, *123*(6), 1111–1122. https://doi.org/10.1016/j.clinph.2012.01.014

Abásolo, D., Simons, S., Da Morgado Silva, R., Tononi, G., & Vyazovskiy, V. V. (2015). Lempel-Ziv complexity of cortical activity during sleep and waking in rats. *Journal of Neurophysiology*, *113*(7), 2742–2752. https://doi.org/10.1152/jn.00575.2014

Ackerman, S. (1992). *Discovering the Brain*. National Academies Press. http://gbv.eblib.com/patron/FullRecord.aspx?p=3376517

Aime, M., Calcini, N., Borsa, M., Campelo, T., Rusterholz, T., Sattin, A., Fellin, T., & Adamantidis, A. (2022). Paradoxical somatodendritic decoupling supports cortical plasticity during REM sleep. *Science*, *376*(6594), 724–730. https://doi.org/10.1126/science.abk2734

Ameen, M. S., Cheung, L. M., Hauser, T., Hahn, M. A [Michael A.], & Schabus, M. (2019). About the Accuracy and Problems of Consumer Devices in the Assessment of Sleep. *Sensors (Basel, Switzerland)*, *19*(19). https://doi.org/10.3390/s19194160

Andrillon, T., Poulsen, A. T., Hansen, L. K., Léger, D., & Kouider, S. (2016). Neural Markers of Responsiveness to the Environment in Human Sleep. *The Journal of Neuroscience*, *36*(24), 6583–6596. https://doi.org/10.1523/JNEUROSCI.0902-16.2016

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)*, *57*(1), 289–300. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Blumberg, M. S., Lesku, J. A., Libourel, P.-A., Schmidt, M. H., & Rattenborg, N. C. (2020). What Is REM Sleep? *Current Biology : CB*, *30*(1), R38-R49. https://doi.org/10.1016/j.cub.2019.11.045

Bódizs, R., Szalárdy, O., Horváth, C., Ujma, P. P., Gombos, F., Simor, P., Pótári, A., Zeising, M., Steiger, A., & Dresler, M. (2021). A set of composite, non-redundant EEG measures of NREM sleep based on the power law scaling of the Fourier spectrum. *Scientific Reports*, *11*(1), 2041. https://doi.org/10.1038/s41598-021-81230-7

Boe, A. J., McGee Koch, L. L., O'Brien, M. K., Shawen, N., Rogers, J. A., Lieber, R. L., Reid, K. J., Zee, P. C., & Jayaraman, A. (2019). Automating sleep stage classification using wireless, wearable sensors. *NPJ Digital Medicine*, *2*, 131. https://doi.org/10.1038/s41746-019-0210-1

Browman, C. P., & Cartwright, R. D. (1980). The first-night effect on sleep and dreams. *Biological Psychiatry*, *15*(5), 809–812.

Chini, M., Pfeffer, T., & Hanganu-Opatz, I. (2022). An increase of inhibition drives the developmental decorrelation of neural activity. *ELife*, *11*. https://doi.org/10.7554/eLife.78811

Colombo, M. A., Napolitani, M., Boly, M., Gosseries, O., Casarotto, S., Rosanova, M., Brichant, J.-F., Boveroux, P., Rex, S., Laureys, S [Steven], Massimini, M., Chieregato, A., & Sarasso, S. (2019). The spectral exponent of the resting EEG indexes the presence of consciousness during unresponsiveness induced by propofol, xenon, and ketamine. *NeuroImage*, *189*, 631–644. https://doi.org/10.1016/j.neuroimage.2019.01.024

Comsa, I.-M. (2019). *Tracking brain dynamics across transitions of consciousness* [Doctoral thesis]. University of Cambridge, Cambridge, England. https://doi.org/10.17863/CAM.37723

Curcio, G., Ferrara, M., Piergianni, A., Fratello, F., & Gennaro, L. de (2004). Paradoxes of the first-night effect: A quantitative analysis of antero-posterior EEG topography. *Clinical Neurophysiology*, *115*(5), 1178–1188. https://doi.org/10.1016/j.clinph.2003.12.018

Dave, S., Brothers, T. A., & Swaab, T. Y. (2018). 1/f neural noise and electrophysiological indices of contextual prediction in aging. *Brain Research*, *1691*, 34–43. https://doi.org/10.1016/j.brainres.2018.04.007

Davis, H., Davis, P. A., Loomis, A. L., Harvey, E. N., & Hobart, G. (1938). Human brain potentials during the onset of sleep. *Journal of Neurophysiology*, *1*(1), 24–38.

Dijk, D.-J. (1995). EEG slow waves and sleep spindles: windows on the sleeping brain. *Behavioural Brain Research*, *69*(1-2), 109–116. https://doi.org/10.1016/0166-4328(95)00007-G

Donders, F. C. (1969). On the speed of mental processes. *Acta Psychologica*, *30*, 412–431. https://doi.org/10.1016/0001-6918(69)90065-1

Donoghue, T., Haller, M., Peterson, E. J., Varma, P., Sebastian, P., Gao, R., Noto, T., Lara, A. H., Wallis, J. D., Knight, R. T., Shestyuk, A., & Voytek, B. (2020). Parameterizing neural power spectra into periodic and aperiodic components. *Nature Neuroscience*, *23*(12), 1655–1665. https://doi.org/10.1038/s41593-020-00744-x

Ferenets, R., Vanluchene, A., Lipping, T., Heyse, B., & Struys, M. M. R. F. (2007). Behavior of entropy/complexity measures of the electroencephalogram during propofol-induced sedation: Dose-dependent effects of remifentanil. *Anesthesiology*, *106*(4), 696–706. https://doi.org/10.1097/01.anes.0000264790.07231.2d

Figueiro, M. G., Sahin, L., Wood, B., & Plitnick, B. (2016). Light at Night and Measures of Alertness and Performance: Implications for Shift Workers. *Biological Research for Nursing*, *18*(1), 90–100. https://doi.org/10.1177/1099800415572873

Friedrich, S., Konietschke, F., & Pauly, M. (2019). Resampling-Based Analysis of Multivariate Data and Repeated Measures Designs with the R Package MANOVA.RM. *The R Journal*, *11*(2), 380–400. https://doi.org/10.32614/RJ-2019-051

Gao, R., & Penzes, P. (2015). Common mechanisms of excitatory and inhibitory imbalance in schizophrenia and autism spectrum disorders. *Current Molecular Medicine*, *15*(2), 146–167. https://doi.org/10.2174/1566524015666150303003028

Gao, R., Peterson, E. J., & Voytek, B. (2017). Inferring synaptic excitation/inhibition balance from field potentials. *NeuroImage*, *158*, 70–78. https://doi.org/10.1016/j.neuroimage.2017.06.078

Gerster, M., Waterstraat, G., Litvak, V., Lehnertz, K., Schnitzler, A., Florin, E., Curio, G., & Nikulin, V. (2022). Separating Neural Oscillations from Aperiodic 1/f Activity: Challenges and Recommendations. *Neuroinformatics.* Advance online publication. https://doi.org/10.1007/s12021-022-09581-8

González, J., Mateos, D., Cavelli, M., Mondino, A., Pascovich, C., Torterolo, P., & Rubido, N. (2022). Low frequency oscillations drive EEG's complexity changes during wakefulness and sleep. *Neuroscience*, *494*, 1–11. https://doi.org/10.1016/j.neuroscience.2022.04.025

Gratton, G., Coles, M. G., & Donchin, E. (1983). A new method for off-line removal of ocular artifact. *Electroencephalography and Clinical Neurophysiology*, *55*(4), 468–484. https://doi.org/10.1016/0013-4694(83)90135-9

Halsband, U. (1998). Encoding and retrieval in declarative learning: a positron emission tomography study. *Behavioural Brain Research*, *97*(1-2), 69–78. https://doi.org/10.1016/S0166-4328(98)00028-X

Harris, K. D., & Thiele, A. (2011). Cortical state and attention. *Nature Reviews. Neuroscience*, *12*(9), 509–523. https://doi.org/10.1038/nrn3084

He, B. J. (2011). Scale-free properties of the functional magnetic resonance imaging signal during rest and task. *The Journal of Neuroscience*, *31*(39), 13786–13795. https://doi.org/10.1523/JNEUROSCI.2111-11.2011

He, B. J. (2014). Scale-free brain activity: Past, present, and future. *Trends in Cognitive Sciences*, *18*(9), 480–487. https://doi.org/10.1016/j.tics.2014.04.003

Höhn, C., Schmid, S. R., Plamberger, C. P [Christina P.], Bothe, K., Angerer, M., Gruber, G., Pletzer, B., & Hoedlmoser, K. (2021). Preliminary Results: The Impact of Smartphone Use and Short-Wavelength Light during the Evening on Circadian Rhythm, Sleep and Alertness. *Clocks & Sleep*, *3*(1), 66–86. https://doi.org/10.3390/clockssleep3010005

Jacob, M. S., Roach, B. J., Sargent, K. S., Mathalon, D. H., & Ford, J. M. (2021). Aperiodic measures of neural excitability are associated with anticorrelated hemodynamic networks at rest: A

combined   EEG-fMRI   study.   *NeuroImage*,   *245*,   118705.
https://doi.org/10.1016/j.neuroimage.2021.118705

Kanashiro, T., Ocker, G. K., Cohen, M. R., & Doiron, B. (2017). Attentional modulation of neuronal variability in circuit models of cortex. *ELife*, *6*. https://doi.org/10.7554/eLife.23978

Karalunas, S. L., Ostlund, B. D., Alperin, B. R., Figuracion, M., Gustafsson, H. C., Deming, E. M., Foti, D., Antovich, D., Dude, J., Nigg, J., & Sullivan, E. (2022). Electroencephalogram aperiodic power spectral slope can be reliably measured and predicts ADHD risk in early development. *Developmental Psychobiology*, *64*(3), e22228. https://doi.org/10.1002/dev.22228

Kirstein, C. (2007). Sleeping and Dreaming. In *xPharm: The Comprehensive Pharmacology Reference* (pp. 1–4). Elsevier. https://doi.org/10.1016/B978-008055232-3.60319-8

Kleiner, M., Brainard, D., Pelli, D., Ingling, A., & Murray, R., Broussard, C. (2007). What's new in psychtoolbox-3. *Perception*, *36*(14), 1–16.

Klimesch, W. (1999). EEG alpha and theta oscillations reflect cognitive and memory performance: a review and analysis. *Brain Research Reviews*, *29*(2-3), 169–195. https://doi.org/10.1016/s0165-0173(98)00056-3

Klimesch, W., Schimke, H., & Pfurtscheller, G. (1993). Alpha frequency, cognitive load and memory performance. *Brain Topography*, *5*(3), 241–251. https://doi.org/10.1007/BF01128991

Kozhemiako, N., Mylonas, D., Pan, J. Q., Prerau, M. J., Redline, S., & Purcell, S. M. (2021). Sources of variation in the spectral slope of the sleep EEG. *BioRxiv.* Advance online publication. https://doi.org/10.1101/2021.11.08.467763

Kuula, L., & Pesonen, A.-K. (2021). Heart Rate Variability and Firstbeat Method for Detecting Sleep Stages in Healthy Young Adults: Feasibility Study. *JMIR MHealth and UHealth*, *9*(2), e24704. https://doi.org/10.2196/24704

La Torre-Luque, A. de, Bornas, X., Balle, M., & Fiol-Veny, A. (2016). Complexity and nonlinear biomarkers in emotional disorders: A meta-analytic study. *Neuroscience and Biobehavioral Reviews*, *68*, 410–422. https://doi.org/10.1016/j.neubiorev.2016.05.023

Lau, Z. J., Pham, T., Chen, S. H. A., & Makowski, D. (2022). Brain entropy, fractal dimensions and predictability: A review of complexity measures for EEG in healthy and neuropsychiatric populations. *The European Journal of Neuroscience.* Advance online publication. https://doi.org/10.1111/ejn.15800

Lempel, A., & Ziv, J. (1976). On the Complexity of Finite Sequences. *IEEE Transactions on Information Theory*, *22*(1), 75–81. https://doi.org/10.1109/TIT.1976.1055501

Lendner, J. D., Helfrich, R. F., Mander, B. A., Romundstad, L., Lin, J. J., Walker, M. P., Larsson, P. G., & Knight, R. T. (2020). An electrophysiological marker of arousal level in humans. *ELife*, *9*. https://doi.org/10.7554/eLife.55092

Lendner, J. D., Mander, B. A., Schuh-Hofer, S., Schmidt, H., Knight, R. T., Walker, M. P., Lin, J., & Helfrich, R. F. (2022). Human REM sleep controls neural excitability in support of memory formation. *BioRxiv.* Advance online publication. https://doi.org/10.1101/2022.05.13.491801

Lüdtke, S., Hermann, W., Kirste, T., Beneš, H., & Teipel, S. (2021). An algorithm for actigraphy-based sleep/wake scoring: Comparison with polysomnography. *Clinical Neurophysiology*, *132*(1), 137–145. https://doi.org/10.1016/j.clinph.2020.10.019

Ma, Y., Shi, W., Peng, C.-K., & Yang, A. C. (2018). Nonlinear dynamical analysis of sleep electroencephalography using fractal and entropy approaches. *Sleep Medicine Reviews*, *37*, 85–93. https://doi.org/10.1016/j.smrv.2017.01.003

Mateos, D., Guevara Erra, R., Wennberg, R., & Perez Velazquez, J. L. (2018). Measures of entropy and complexity in altered states of consciousness. *Cognitive Neurodynamics*, *12*(1), 73–84. https://doi.org/10.1007/s11571-017-9459-8

Mediano, P. A. M., Ikkala, A., Kievit, R. A., Jagannathan, S. R., Varley, T. F., Stamatakis, E. A., Bekinschtein, T. A., & Bor, D. (2021). Fluctuations in Neural Complexity During Wakefulness Relate To Conscious Level and Cognition. *BioRxiv.* Advance online publication. https://doi.org/10.1101/2021.09.23.461002

Mediano, P. A. M., Rosas, F. E., Timmermann, C., Roseman, L., Nutt, D. J., Feilding, A., Kaelen, M., Kringelbach, M. L., Barrett, A. B., Seth, A., Muthukumaraswamy, S. D., Bor, D., & Carhart-Harris, R. L. (2020). Effects of external stimulation on psychedelic state neurodynamics. *BioRxiv.* Advance online publication. https://doi.org/10.1101/2020.11.01.356071

Miskovic, V., MacDonald, K. J., Rhodes, L. J., & Cote, K. A. (2019). Changes in EEG multiscale entropy and power-law frequency scaling during the human sleep cycle. *Human Brain Mapping*, *40*(2), 538–551. https://doi.org/10.1002/hbm.24393

Mitsukura, Y., Fukunaga, K., Yasui, M., & Mimura, M. (2020). Sleep stage detection using only heart rate. *Health Informatics Journal*, *26*(1), 376–387. https://doi.org/10.1177/1460458219827349

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). Fieldtrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational Intelligence and Neuroscience*, *2011*, 156869. https://doi.org/10.1155/2011/156869

Ouyang, G., Hildebrandt, A., Schmitz, F., & Herrmann, C. S. (2020). Decomposing alpha and 1/f brain activities reveals their differential associations with cognitive processing speed. *NeuroImage*, *205*, 116304. https://doi.org/10.1016/j.neuroimage.2019.116304

Pascovich, C., Castro-Zaballa, S., Mediano, P. A. M., Bor, D., Canales-Johnson, A., Torterolo, P., & Bekinschtein, T. A. (2022). Ketamine and sleep modulate neural complexity dynamics in cats. *The European Journal of Neuroscience*, *55*(6), 1584–1600. https://doi.org/10.1111/ejn.15646

Pathania, A., Euler, M. J., Clark, M., Cowan, R. L., Duff, K., & Lohse, K. R. (2022). Resting EEG spectral slopes are associated with age-related differences in information processing speed. *Biological Psychology*, *168*, 108261. https://doi.org/10.1016/j.biopsycho.2022.108261

Peever, J., & Fuller, P. M. (2017). The Biology of REM Sleep. *Current Biology : CB*, *27*(22), R1237-R1248. https://doi.org/10.1016/j.cub.2017.10.026

Peigneux, P., Laureys, S [S.], Fuchs, S., Delbeuck, X., Degueldre, C., Aerts, J., Delfiore, G., Luxen, A., & Maquet, P. (2001). Generation of rapid eye movements during paradoxical sleep in humans. *NeuroImage*, *14*(3), 701–708. https://doi.org/10.1006/nimg.2001.0874

Pereda, E., Gamundi, A., Rial, R., & González, J. (1998). Non-linear behaviour of human EEG: fractal exponent versus correlation dimension in awake and sleep stages. *Neuroscience Letters*, *250*(2), 91–94. https://doi.org/10.1016/S0304-3940(98)00435-2

Plamberger, C. P [Christina Paula], van Wijk, H. E., Kerschbaum, H., Pletzer, B. A., Gruber, G., Oberascher, K., Dresler, M., Hahn, M. A [Michael Andreas], & Hoedlmoser, K. (2021). Impact of menstrual cycle phase and oral contraceptives on sleep and overnight memory consolidation. *Journal of Sleep Research*, *30*(4), e13239. https://doi.org/10.1111/jsr.13239

Podvalny, E., Noy, N., Harel, M., Bickel, S., Chechik, G., Schroeder, C. E., Mehta, A. D., Tsodyks, M., & Malach, R. (2015). A unifying principle underlying the extracellular field potential spectral responses in the human cortex. *Journal of Neurophysiology*, *114*(1), 505–519. https://doi.org/10.1152/jn.00943.2014

Radha, M., Fonseca, P., Moreau, A., Ross, M., Cerny, A., Anderer, P., Long, X., & Aarts, R. M. (2019). Sleep stage classification from heart-rate variability using long short-term memory neural networks. *Scientific Reports*, *9*(1), 14149. https://doi.org/10.1038/s41598-019-49703-y

Richard, B. B., Albertario, C. L., Harding, S. M., Uoyd, R. M., Plante, D. T., Quan, S. F., Troester, M. M., & Vaughn, B. V. (2012). *The AASM Manual for the Scoring of Sleep and Associated Events: Rules, Terminology and Technical Specifications* (2nd ed.). American Academy of Sleep Medicine: Darien, IL. www.aasmnet.org

Rivolta, M. W., Migliorini, M., Aktaruzzaman, M., Sassi, R., & Bianchi, A. M. (2014). Effects of the series length on Lempel-Ziv Complexity during sleep. *Annual International Conference of the IEEE Engineering in Medicine and Biology Society. IEEE Engineering in Medicine and Biology Society. Annual International Conference*, *2014*, 693–696. https://doi.org/10.1109/EMBC.2014.6943685

Roberts, D. M., Schade, M. M., Mathew, G. M., Gartenberg, D., & Buxton, O. M. (2020). Detecting sleep using heart rate and motion data from multisensor consumer-grade wearables, relative to wrist actigraphy and polysomnography. *Sleep*, *43*(7). https://doi.org/10.1093/sleep/zsaa045

Robertson, M. M., Furlong, S., Voytek, B., Donoghue, T., Boettiger, C. A., & Sheridan, M. A. (2019). Eeg power spectral slope differs by ADHD status and stimulant medication exposure in early childhood. *Journal of Neurophysiology*, *122*(6), 2427–2437. https://doi.org/10.1152/jn.00388.2019

Rubenstein, J. L. R., & Merzenich, M. M. (2003). Model of autism: Increased ratio of excitation/inhibition in key neural systems. *Genes, Brain, and Behavior*, *2*(5), 255–267. https://doi.org/10.1034/j.1601-183X.2003.00037.x

Sadeh, A., Alster, J., Urbach, D., & Lavie, P. (1989). Actigraphically based automatic bedtime sleep-wake scoring: Validity and clinical applications. *Journal of Ambulatory Monitoring*(3), 209–216.

Schartner, M., Pigorini, A., Gibbs, S. A., Arnulfo, G., Sarasso, S., Barnett, L., Nobili, L., Massimini, M., Seth, A., & Barrett, A. B. (2017). Global and local complexity of intracranial EEG decreases during NREM sleep. *Neuroscience of Consciousness*, *2017*(1), 1-12. https://doi.org/10.1093/nc/niw022

Schartner, M., Seth, A., Noirhomme, Q., Boly, M., Bruno, M.-A., Laureys, S [Steven], & Barrett, A. B. (2015). Complexity of Multi-Dimensional Spontaneous EEG Decreases during Propofol Induced General Anaesthesia. *PloS One*, *10*(8), e0133532. https://doi.org/10.1371/journal.pone.0133532

Schaworonkow, N., & Voytek, B. (2021). Longitudinal changes in aperiodic and periodic activity in electrophysiological recordings in the first seven months of life. *Developmental Cognitive Neuroscience*, *47*, 100895. https://doi.org/10.1016/j.dcn.2020.100895

Schmid, S. R., Höhn, C., Bothe, K., Plamberger, C. P [Christina P.], Angerer, M., Pletzer, B., & Hoedlmoser, K. (2021). How Smart Is It to Go to Bed with the Phone? The Impact of Short-Wavelength Light and Affective States on Sleep and Circadian Rhythms. *Clocks & Sleep*, *3*(4), 558–580. https://doi.org/10.3390/clockssleep3040040

Sheehan, T. C., Sreekumar, V., Inati, S. K., & Zaghloul, K. A. (2018). Signal Complexity of Human Intracranial EEG Tracks Successful Associative-Memory Formation across Individuals. *The Journal of Neuroscience*, *38*(7), 1744–1755. https://doi.org/10.1523/JNEUROSCI.2389-17.2017

Siegel, J. M. (2011). Rem sleep: A biological and psychological paradox. *Sleep Medicine Reviews*, *15*(3), 139–142. https://doi.org/10.1016/j.smrv.2011.01.001

Symonds, C. (1959). Excitation and inhibition in epilepsy. *Proceedings of the Royal Society of Medicine*, *52*(6), 395–402.

Tagliazucchi, E., Wegner, F. von, Morzelewski, A., Brodbeck, V., Jahnke, K., & Laufs, H. (2013). Breakdown of long-range temporal dependence in default mode and attention networks during deep sleep. *Proceedings of the National Academy of Sciences of the United States of America*, *110*(38), 15419–15424. https://doi.org/10.1073/pnas.1312848110

Tal, A., Shinar, Z., Shaki, D., Codish, S., & Goldbart, A. (2017). Validation of Contact-Free Sleep Monitoring Device with Comparison to Polysomnography. *Journal of Clinical Sleep Medicine*, *13*(3), 517–522. https://doi.org/10.5664/jcsm.6514

Terzano, M. G., Parrino, L., Smerieri, A., Chervin, R., Chokroverty, S., Guilleminault, C., Hirshkowitz, M., Mahowald, M., Moldofsky, H., Rosa, A., Thomas, R., & Walters, A. (2002). Atlas, rules, and recording techniques for the scoring of cyclic alternating pattern (CAP) in human sleep. *Sleep Medicine*, *3*(2), 187–199. https://doi.org/10.1016/S1389-9457(01)00149-6

Tosun, P. D., Dijk, D.-J., Winsky-Sommerer, R., & Abásolo, D. (2019). Effects of Ageing and Sex on Complexity in the Human Sleep EEG: A Comparison of Three Symbolic Dynamic Analysis Methods. *Complexity*, *2019*, 1–12. https://doi.org/10.1155/2019/9254309

Treder, M. S. (2020). Mvpa-Light: A Classification and Regression Toolbox for Multi-Dimensional Data. *Frontiers in Neuroscience*, *14*, 289. https://doi.org/10.3389/fnins.2020.00289

van de Borne, P., Nguyen, H., Biston, P., Linkowski, P., & Degaute, J. P. (1994). Effects of wake and sleep stages on the 24-h autonomic control of blood pressure and heart rate in recumbent men. *The American Journal of Physiology*, *266*(2), 548-554. https://doi.org/10.1152/ajpheart.1994.266.2.H548

Voytek, B., & Knight, R. T. (2015). Dynamic network communication as a unifying neural basis for cognition, development, aging, and disease. *Biological Psychiatry*, *77*(12), 1089–1097. https://doi.org/10.1016/j.biopsych.2015.04.016

Voytek, B., Kramer, M. A., Case, J., Lepage, K. Q., Tempesta, Z. R., Knight, R. T., & Gazzaley, A. (2015). Age-Related Changes in 1/f Neural Electrophysiological Noise. *The Journal of Neuroscience*, *35*(38), 13257–13265. https://doi.org/10.1523/JNEUROSCI.2332-14.2015

Waschke, L., Donoghue, T., Fiedler, L., Smith, S., Garrett, D. D., Voytek, B., & Obleser, J. (2021). Modality-specific tracking of attention and sensory statistics in the human electrophysiological spectral exponent. *ELife*, *10.* https://doi.org/10.7554/eLife.70068

Waschke, L., Wöstmann, M., & Obleser, J. (2017). States and traits of neural irregularity in the age-varying human brain. *Scientific Reports*, *7*(1), 17381. https://doi.org/10.1038/s41598-017-17766-4

Welch (1984). A Technique for High-Performance Data Compression. *Computer*, *17*(6), 8–19. https://doi.org/10.1109/MC.1984.1659158

Wickham, H. (2016). *Ggplot2: Elegant graphics for data analysis* (Second edition). *Use R!* Springer.
https://doi.org/10.1007/978-3-319-24277-4

Wong, M. (2010). Too much inhibition leads to excitation in absence epilepsy. *Epilepsy Currents*, *10*(5),
131–132. https://doi.org/10.1111/j.1535-7511.2010.01379.x

Zhang, X. S., Roy, R. J., & Jensen, E. W. (2001). Eeg complexity as a measure of depth of anesthesia
for patients. *IEEE Transactions on Bio-Medical Engineering*, *48*(12), 1424–1433.
https://doi.org/10.1109/10.966601

Zhu, X., Xu, H., Zhao, J., & Tian, J. (2017). Automated Epileptic Seizure Detection in Scalp EEG Based
on Spatial-Temporal Complexity. *Complexity*, *2017*, 1–8. https://doi.org/10.1155/2017/5674392