Crystal structure of *Vibrio cholerae* (6-4) photolyase with DNA binding region

Baris Cakilkaya[1], Ibrahim Halil Kavakli[1,2*], and Hasan DeMirci[1,3*]

[1]*Koc University, Department of Molecular Biology and Genetics, Rumelifeneri Yolu Sariyer, Istanbul, Türkiye*

[2]*Koc University, Department Chemical and Biological Engineering, Rumelifeneri Yolu Sariyer, Istanbul, Türkiye*

[3]*PULSE Institute, SLAC National Accelerator Laboratory Menlo Park, California 94025, United States*

[*]**Corresponding authors**

e-mails: hdemirci@ku.edu.tr

hkavakli@ku.edu.tr

Abstract

Photolyases (PLs) are the enzymes that reverse UV-Induced DNA damages by using blue-light as an energy source. Among those photolyases (6-4) PLs are the enzymes that repair (6-4) lesioned photoproduct. Here, we determined *Vibrio cholerae* (*Vc*) crystal structure of (6-4) PL at 2.5 Å resolution. Our high-resolution structure revealed that the presence of two well-known cofactors, flavin adenine dinucleotide (FAD) and or 6,7-dimethyl 8-ribityl-lumazin (DMRL), stably interact with an α-helical and an α/β and domains, respectively. Additionally, it has also a third cofactor with distinct electron clouds corresponding to [4Fe-4S] cluster. Asp106 makes a hydrogen bond with the water and DMRL, which indicates further stabilization of the photoantenna DMRL to *Vc*(6-4) PL. Further analysis of *Vc*(6-4) PL structure revealed a possible region responsible for the DNA binding. The region located between residues 478-484 may binds the lesioned DNA and Arg483 forms a salt bridge with DNA to stabilize further *Vc*(6-4) PL with its substrate. Our comparative analysis revealed that DNA lesion could not bind to the *Vc*(6-4) PL in a similar fashion to the *Dm*(6-4) PL without a significant conformational change in the protein. The 23[rd] helix of the bacterial (6-4) PLs seems to have a remarkable plasticity and conformational changes facilitate the DNA binding.

Introduction

The cryptochrome/photolyase family (CPF) is a large family of flavoproteins that absorb near UV–visible light (300–500 nm range) and has diverse functions depending on the type of organism (1,2). Cryptochromes (CRY) are the circadian photoreceptors in Drosophila and plants while they are acting as transcriptional repressor of the circadian clock in mammals (2-4). Photolyases (PLs) are enzymes that repair UV-induced DNA damage (5-7). Although phylogenetic analyses revealed that there are different classes of PLs, there are three different types of PLs based on their repair activities: Cyclobutane pyrimidine dimer (CPD) photolyases (PL), (6–4) photolyases, and Drosophila, Arabidopsis, Synechocystis, human (DASH)-type cryptochrome (CRY-DASH) (8-10). CPD PLs repair mainly cyclobutane pyrimidine dimers (Pyr<>Pyr) while (6–4) PLs repair pyrimidine-pyrimidine photoproducts (Pyr [6–4] Pyr) using blue light (350–500 nm) as the energy source (1,6). CRY-DASHs have quite diverse physiological functions including single-strand DNA photolyase activity (11). PLs are monomeric proteins with molecular masses of 50–61 kDa and possess two chromophores (12). Flavin Adenine Dinucleotide (FAD) is an essential cofactor for the catalytic activity of PLs and acts as a catalytic cofactor in all PLs (5,12). The second chromophore, which acts as photoantenna, varies depending on the organisms. Following photoantenna are being found: methenyltetrahydrofolate (MTHF), 8-hydroxy-7,8-didemethyl-5-deazariboflavin (8-HDF), FAD, flavin mononucleotide (FMN), or 6,7-dimethyl 8-ribityl-lumazin (DMRL) depending on the type of photolyase (13,14). The (6–4) photolyases had previously been thought to exist only in eukaryotes. However, studies with different bacteria revealed the presence of (6–4) PLs (15-17). These bacterial (6–4) PLs are unique having [4Fe-4S] in addition to DMRL and FAD cofactors (15,17).

Crystal structures of various PLs have been determined by X-ray crystallography. First available crystal structure is the *Escherichia coli* CPD PL (18). Subsequent structures have been obtained from different organismal PLs including *Anacystis nidulans* (19), *Thermus thermophilus* (20), (6–4) photolyases of *Arabidopsis thaliana* (21) and *Agrobacterium tumefaciens* (13). Comparison of these crystal structures of various homologous PLs revealed valuable information about their structural features and enzyme mechanisms. PLs are made up of two well-defined domains which are a α/β and an α-helical domains. The α-helical domain is

associated with FAD while α/β domain is the binding site for the second chromophore (12). These two domains are connected by a long flexible loop.

V. cholerae O1 bivar Tor str. N16961 has a CPD photolyase that repairs T<>T dimers and two CRY-DASHs that repair T<>T dimers in single-stranded DNA (11,22). Our previous study with a gene (Vca0809) in V. cholerae highly expressed upon exposure of the organism to blue light (23). Our further characterization of this gene revealed that it encodes (6–4) PL with FAD and DMRL and consists of the [4Fe-4S] domain (15). Here, we determined the crystal structure of Vc(6–4) PL with 2.5 Å resolution. Our crystal structure comparison revealed a putative DNA binding region of Vc(6–4) PL.

Results

Crystal structure of *Vibrio cholerae* (6-4) photolyase

We recently identified a (6-4) PL from *Vibrio cholerae* (*Vc*) which possess catalytic cofactor FAD, an antenna chromophore DMRL, and additional cofactor [4Fe-4S] cluster (15). We determined the crystal structure of Vc(6-4) PL at 2.5 Å resolution at cryogenic temperature. The structure contains all the cofactors with well-defined electron densities and contains a total of 24 α helixes and 5 β sheets (**Figure 1A**). It consists of two domains: an α-helical domain that is associated with FAD and α/β domain which interacts with DMRL cofactor. The sequence alignment of Vc(6-4) PL with structurally available bacterial (6-4) PL with 38.4% sequence similarity with *Agrobacterium tumefaciens* (*At*)(6-4) PL and 43.1% sequence similarity with *Rhodobacter sphaeroides* (*Rs*)(6-4) PL (**Figure 1B**).

DMRL Binding domain

DMRL is a photoantenna chromophore and has higher efficiency to absorb light then FAD for increasing the catalytic efficiency (13). In our crystal structure, there is a very well-defined electron density of the DMRL and interacting amino acid residues (**Figure 2A**). Aromatic ring of DMRL interacts with nitrogen atoms of Gln11, Leu35, Gln39, and oxygen atom of Ala33 (**Figure 2B**). Additionally, aromatic ring of DMRL interacts with Trp488, Ala33 and Ile9. The ribityl group of the DMRL forms hydrogen bonds with oxygen of the Glu38, Tyr41, Asp12 and with amino group of Asp10. When we compare the interaction partners of

DMRL in *Vc*(6-4) PL with other known bacterial (6-4) PLs, almost all the interactions are well-conserved with a few exceptions. First, Cys32 carboxyl group of *Af*(6-4) PL, interacts with aromatic ring in DMRL (PDB ID: 4DJA) while in both *Vc* and *Rs* (6-4) PLs DMRL interact with main chain carbonyl group of Ala33. We identified a critical water molecule (W22) that interact with Asp12, Gln13, Asp106, and DMRL in *Vc*(6-4) PL (**Figure 2C**). Analyses of other bacterial (6-4) PLs have shown similar conserved motifs (**Figure 2C**). Such motif increases the overall stability of the DMRL within the enzyme. However, W22 molecule interacts with Asp106 of the *Vc*(6-4) PL with a distance of 2.8 Å while same interaction occurs with Gly105 of other bacterial (6-4) PLs with a distance of over 3.3 Å. These difference might increase the affinity of DMRL with *Vc*(6-4) PL.

FAD binding domain

FAD is located in the α-helical domain with U shaped conformation (**Figure 3A**). The N5 and O4 of FAD make hydrogen bonds with W7 and additional hydrogen bonds with carbonyl group of Tyr398 and NH1 atom of Arg376 residue (**Figure 3B**). As previously suggested for other PLs (17) these critical interaction may also serves a stabilization function *Vc*(6-4) PL. Unlike *At*(6-4) PL or *Rs*(6-4) PL, *Vc*(6-4) PL Glu410 amino acid faces away from FAD and interacts with Asp395 rather than an His365 residue (**Figure 3B**). This results a much-relaxed interaction between the residues due to hydrogen bonding formation rather than a salt bridge formation in *Vc*(6-4) PL.

Although reaction mechanism of the CPD photolyases is elucidated, (6-4) PLs not yet fully characterized. There are different mechanisms that are proposed for the (6–4) PL mechanism during DNA repair (7,14,24,25). In fact, mutagenesis studies with *Dm* (6-4) PL indicated that replacement of His365 to Asn365 result in loss of its DNA repair activity while mutagenesis of His369 into Met369 result in highly reduced activity of *Dm*(6-4) PL (26). However, mutagenesis studies with corresponding amino acid residues in *Xenopus* (6-4) PL showed complete loss of activity (27). Femtosecond spectroscopy and site-directed mutagenesis suggest initial electron transfer from excited flavin induces transfer of a proton from a histidine on the active site of the enzyme to the (6–4) photoproduct (7,28).

Catalytically active His373 (corresponds His365 of the *Dm*(6-4) PL) is conserved in *Vc*(6-4) PL (**Figure 3C**). In addition, there are other conserved amino acid residues in the motif.

For instance, His372 and Arg376 (corresponds to His364 and Arg368 of the *Dm*(6-4) PL) are conserved in *Vc*(6-4) PL (**Figure 3C**). However, the conformation of the *Vc*(6-4) PL His372 is significantly different than the *Dm*(6-4) PL His364, which may suggest an alternate function of this residue.

[4Fe-4S] cluster

The high-resolution Vc(6-4) PL structure contains an iron-sulfur cluster with an unknown function with a well-defined electron density (**Figure 4A**). Its interactions with the surrounding cysteine amino acids summarized in **Figure 4B**. Cys357, Cys445, Cys448, Cys461 sulfide atoms make bonds with Fe atoms in the cluster with each bond at 2.3 Å distance. The function of [4Fe-4S] cluster is currently unknown, however, its distance to the FAD is 17Å, which indicates it does not have a catalytic role in the DNA repair due to its distance to the active site region. [4Fe-4S] clusters are well-known for their oxygen sensitivity, and under aerobic conditions it quickly decomposes into [3Fe-4S] cluster and can further dissociates into [2Fe-2S]. Mostly the decomposition of the cluster deactivates the cluster containing enzymes and handling and purification of these enzymes require using glovebox to prevent decomposition of the [4Fe-4S] clusters. In case of *Vc*(6-4) PL, the entire expression, purification, and crystallization procedures were performed in atmospheric conditions that took months. In our structure, there was no observable damage in the [4Fe-4S] cluster, which might indicate the cluster in the photolyase is unusually oxygen tolerant.

DNA binding region of *Vc*(6-4) photolyase

The DNA binding region of the *Vc*(6-4) PL has defined electron density covers amino acids Asn178, Phe179, Asp180, Ala181, Asp182, Asn183, Arg184, and Asn185 (**Figure 5A**). In contrast region has not a defined electron density in *At*(6-4) PL structures obtained in either cryogenic (PDB ID:4DJA) or ambient temperature (PDB ID:6DD6) conditions (**Figure 5B** and **C**). A similar analysis were carried out with *Rs*(6-4) PL, where there is a well-defined electron density   in the same region (**Figure 5D**). These results suggest that while this region is highly stable in *Rs*(6-4) PL and *Vc*(6-4) PL, it is flexible and disordered in *At*(6-4) PL structure and, therefore this region has species specific stability differences.

Notably, we have not observed a distinct electron density of amino acid residues between 478 and 485 *Vc*(6-4) PL (**Figure 6A**). However, this region has defined structures in other bacterial PLs (**Figure 6 B,C and D**). Interestingly, this region is adjacent to the 23rd helix, which is one of the positively surface charge helices around the catalytic region (**Figure 7A**). When we superimposed *Vc*(6-4) PL with *Dm*(6-4) PL (PDB ID: 3CVU) with (6-4) T-T, this region and 23rd helix were seen to pass through damaged DNA (**Figure 7B**). This observation suggest that region between 478 and 485 *Vc*(6-4) PL might be DNA binding region and further stabilized after interacting with the DNA lesion. Upon investigation of C-terminus of *Vc*(6-4) PL and *Dm*(6-4) PL, 2 additional helixes were identified on *Vc*(6-4) PL and 23rd helix seems to be blocking the DNA lesion in superimposed structure. Further analysis of superimposed structures indicated that the catalytic cofactor FADs are in similar conformation and their RMSD is 0.7 Å (**Figure 7C**). To DNA lesion to bind the bacterial (6-4) PL, while still interacting with the FAD, there may be a conformational changes on the 23rd helix to provide a gap for DNA lesion. B-factor analysis showed that, 23rd helix has the highest b-factor values in the structure and therefore, it has highest probability to subject a conformational change (**Figure 7D**). According to the superimposition of DNA lesion from *Dm*(6-4) PL with *Vc*(6-4) PL, which is compatible with a prototypical photolyase-DNA binding mode, helix α23 of *Vc*(6-4) PL interacts with the DNA, while helix α24 looks away from the DNA lesion. The surface charge of helix α23 is positive, which is consistent with the role in DNA binding. The last two helixes (α23, α24) are missing in *Dm*(6-4) PL or most of the other PLs, which indicates they may play regulatory role rather than their involvement in repair activity.

Discussion

Photolyases (PLs) belong to a large family of Cryptochrome/Photolyase responsible for repairing the UV-induced DNA damages. Different types of photolyases have been discovered and they are shown to repair two different types of DNA damages: cyclobutane pyrimidine dimers (CPD) (Pyr<>Pyr) and pyrimidine-pyrimidone photoproducts (Pyr[6–4]Pyr). CPD photoproducts are mainly repaired by CPD PLs while (6-4) photoproducts are being repaired by (6-4) PLs (12). The reaction mechanism of CPD photolyases is elucidated at *in vivo* level (1,5,6). On the other hand, different reaction mechanisms are proposed for the (6-4) PL. The (6-4) PLs is initially thought to specifically present in eucaryotes (13). However, studies revealed that

bacteria also possess (6-4) PLs (15,16,29). Comparison of the crystal structure of the *Vc* (6-4) PL with other known bacterial (6-4) PLs indicated that that cofactors binding the region is well conserved with some differences: a water molecule interacts with the Asp106 in *Vc*(6-4) PL with a very short distance compare to the *At,* and *Rs* (6-4) PLs, where water molecule interacts with Gly105 (**Figure 2B**). The FAD binding region of the *Vc*(6-4) PL is highly conserved. However, one notable difference is that Glu410 makes a hydrogen bond with Asp395; however, in the case of Glu399 (corresponds to Glu410 of *Vc*(6-4) PL) makes a salt bridge with His384 (corresponds to Asp395 of *Vc*(6-4) PL) *At*(6-4) PL and *Rs*(6-4) PL (13,16). Whether such structural difference affects its activity or conformation of FAD needs to be further investigated. The His-His-X-X-Arg motif in (6-4) PLs is shown to be important for their catalytic activity (16). This motif is also conserved in the *Vc*(6-4) PL. Further conformational comparison of the *Vc*(6-4) PL and *Dm*(6-4) PL of this motif indicates His373 and Arg376 of *Dm*(6-4) PL have identical configuration while His372 (His364 of *Dm*(6-4) PL) acquired a different configuration, which may reveal similar/different functionalities between species.

One of the major difference between *Vc*(6-4) PL and *At*(6-4) PL structures is their predicted DNA binding site (**Figure 5**). The predicted site (amino acid residues between 178 and 185) of Vc(6-4) PL has distinct structure compared to *At*(6-4) PL, where corresponding region has no defined structure (17). The same region of *Rs*(6-4) PL has also a nice electron density except its Arg residue (16) which indicates species specific stability differences of DNA binding domains. Comparison of *Vc*(6-4) PL and *At*(6-4) PL revealed that *Vc*(6-4) PL has unstructured region with 8 amino acid residues (**Figure 6A**), which might be DNA binding region and also conserved in other bacterial PLs. Arg-483 might form a salt bridge with (6-4) photoproduct that stabilize the disordered region. The general structure of *Vc* and *Dm*(6-4) PLs are highly similar except in C terminus, where *Vc*(6-4) PLs contain additional 2 helixes. Due to position of 23rd helix, DNA lesion cannot bind the *Vc*(6-4) PL similar to *Dm*(6-4) PL (**Figure 7B**). However, the FADs are located extremely similar manner, which indicates a similar DNA lesion binding between the PLs (**Figure 7C**). A conformational change upon DNA binding at the *Vc*(6-4) PL might open a space for DNA to bind. The 23rd helix is the main candidate for the conformational change as it is the major blocker of the DNA and has the largest b-factor (**Figure 7D**).

Experimental procedures

Expression and purification

Full-length (6-4) photolyase from *Vibrio cholerae* in pET28a(+) vector between NdeI and BamHI cut sites purchased from Genscript (Piscataway, NJ, USA) as codon optimized for *Escherichia coli protein expression*. The plasmid transformed into *E. coli* BL21 Rosetta-2 strain and grown in 4.5 liters of LB-Miller with 50 μg/mL Kanamycin and 35 μg/mL Chloramphenicol antibiotics at 37 °C with 110 rpm shaking. When OD_{600} reached 0.8, induction was performed by adding 0.4 mM of isopropyl β-D-thiogalactopyranoside (IPTG) as the final concentration, and induction was performed at 18°C for 24h. The culture was centrifuged at 2850 g for 45 minutes at 4 °C and the pellet was stored at -80 °C. Lysis buffer containing 500 mM NaCl, 50 mM Tris-HCl pH 7.5, %10 glycerol (v/v), 0.1% Triton X-100 (v/v) was added into pellets and sonication (Branson W250 sonifier, USA) was performed at 45 sec at 60% power 3 times. Samples were ultracentrifuged at 35000 rpm with Ti-45 rotor (Beckman, USA) for 1 hour at 4°C, subsequently the supernatant was filtered with 45 μm cellulose mixed ester filter (ISOLAB, Germany), then applied to Ni-NTA agarose resin (QIAGEN, USA). The column was equilibrated with His A buffer containing 200 mM NaCl, 20 mM Tris-HCl pH 7.5 (v/v), 5% glycerol (v/v) then sample was loaded with 2.5 ml/min. Washing of the column was performed with His A buffer, and protein was eluted with His B buffer containing200 mM NaCl, 20 mM Tris-HCl pH 7.5, 250 mM imidazole, 5% glycerol (v/v) into 5 ml of 100% glycerol to prevent protein precipitation. Yellow colored (6-4) photolyase was flash frozen with liquid nitrogen and stored at -80 °C with the final concentration of 25% glycerol (v/v).

Crystallization, Data collection and Processing

Initial crystallization screening experiments were setup with 5 mg/mL final concentration of (6-4) photolyase with the sitting drop, microbatch screening (under oil) method at 4 °C by mixing 0.83 μl of protein with equal volume of crystal screening conditions in 72 well Terasaki plates. The well containing the protein:coctail mixture was covered with 16.6 μl of parafilm oil to allow slow evaporation of solvents. Approximately 3500 commercially available crystal screening conditions were tested, and the yellow-colored crystals were observed at Wizard

Synergy #40 (Rigaku, Japan), which contains 2 M of ammonium citrate/citric acid pH 7.5 and 5% PEG 400 (v/v) after 7 weeks. The conditions were further optimized by mixing 0.5 µl Cryo-Pro #45 (Hampton research, USA), which contain 1 M sodium sulfate decahydrate, 1 µl of protein, 1 µl of Wizard Synergy #40. The crystals were flash frozen with liquid nitrogen, and data collection was performed at 100 K. X-ray diffraction data were collected to 2.0 Å and 2.5 Å resolutions with a wavelength of 1.54 Å at University of Health Sciences (Istanbul, Turkiye) with Rigaku's XtaLAB Synergy Flow X-ray diffractometer (Rigaku, Japan). X-ray generator PhotonJet-R (Rigaku, Japan) was operated at 40 kV and 30 mA with 23% beam intensity. HyPix-Arc-150° detector (Rigaku, Japan) was used with 60 mm detector distance, 1-degree oscillation scan width, and 15 and 30 seconds exposure times. The total run time for the two data collections was 1h 11m 30sec, and 2h 33min 0sec respectively. Profit merge is done by using the data from both crystals and data reduction is performed with CrysAlis Pro (30) software version 171.42.51a. Unit cell dimensions were a=200.751 Å b=200.751 Å c=77.0231 Å α=90 β=90 γ=120, in space group P6$_4$22.

Structure Determination and Refinement

Swiss model (31) was used for building a search model for structure determination that gave the result with PDB ID: 4DJA (13), which was used in the automated molecular replacement program *PHASER* (32) in *PHENIX* (33) software version 1.20.1-4487. Further refinements were performed with *PHENIX*, and the addition of water and cofactors was performed by using *COOT* (34) software version 0.9.6. The final R_{work} is 22.54% and R_{free} is as 28.30%, completeness is 99.60% with 24.250 to 2.500 Å refinement resolution (Table 1). The structure contains 0.00% Ramachandran outliers with 98.01% favored residues. The structure was deposited to the rcsb website with the PDB ID:7YKN. Figures were generated at *PyMOL* (35) Version 2.4.1. Data collection and structure determination information are summarized in Table 1.

Data Availability

All data generated or analyzed during this study are included in this article and the structure was deposited to the RCSB Protein Data Bank website with the PDB ID:7YKN.

Author Contributions

H.D. and I.H.K. designed and coordinated the project. H.D guided B. C. to execute all the sample preparation, data collection, data reduction and refinement of photolyase structure. H.D., I.H.K and B.C analyzed data. H.D., I.H.K and B.C. prepared the manuscript.

Acknowledgements

Authors would like to dedicate this manuscript to the memory of Dr. Albert E. Dahlberg and Dr. Nizar Turker. H.D. acknowledges support from NSF Science and Technology Center grant NSF-1231306 (Biology with X-ray Lasers, BioXFEL). This publication has been produced benefiting from the 2232 International Fellowship for Outstanding Researchers Program of the TÜBİTAK (Project No. 118C270). However, the entire responsibility of the publication belongs to the authors of the publication. The financial support received from TÜBİTAK does not mean that the content of the publication is approved in a scientific sense by TÜBİTAK.

Conflict of Interest Statement

The authors declare that they have no conflicts of interest with the contents of this article.

References

1.   Kavakli, I. H., Baris, I., Tardu, M., Gul, S., Oner, H., Cal, S., Bulut, S., Yarparvar, D., Berkel, C., Ustaoglu, P., and Aydin, C. (2017) The Photolyase/Cryptochrome Family of Proteins as DNA Repair Enzymes and Transcriptional Repressors. *Photochem Photobiol* **93**, 93-103
2.   Sancar, A. (2004) Photolyase and cryptochrome blue-light photoreceptors. *Adv Protein Chem* **69**, 73-100
3.   Chaves, I., Pokorny, R., Byrdin, M., Hoang, N., Ritz, T., Brettel, K., Essen, L. O., van der Horst, G. T., Batschauer, A., and Ahmad, M. (2011) The cryptochromes: blue light photoreceptors in plants and animals. *Annu Rev Plant Biol* **62**, 335-364
4.   Kavakli, I. H., and Sancar, A. (2004) Analysis of the role of intraprotein electron transfer in photoreactivation by DNA photolyase in vivo. *Biochemistry* **43**, 15103-15110
5.   Kavakli, I. H., Ozturk, N., and Gul, S. (2019) DNA repair by photolyases. *Adv Protein Chem Struct Biol* **115**, 1-19
6.   Sancar, A. (2016) Mechanisms of DNA Repair by Photolyase and Excision Nuclease (Nobel Lecture). *Angew Chem Int Ed Engl* **55**, 8502-8527

7.   Zhong, D. (2015) Electron transfer mechanisms of DNA repair by photolyase. *Annu Rev Phys Chem* **66**, 691-715

8.   Asimgil, H., and Kavakli, I. H. (2012) Purification and characterization of five members of photolyase/cryptochrome family from Cyanidioschyzon merolae. *Plant Sci* **185-186**, 190-198

9.   Ozturk, N. (2017) Phylogenetic and Functional Classification of the Photolyase/Cryptochrome Family. *Photochem Photobiol* **93**, 104-111

10.   Ozturk, N., Kao, Y. T., Selby, C. P., Kavakli, I. H., Partch, C. L., Zhong, D., and Sancar, A. (2008) Purification and characterization of a type III photolyase from Caulobacter crescentus. *Biochemistry* **47**, 10255-10261

11.   Selby, C. P., and Sancar, A. (2006) A cryptochrome/photolyase class of enzymes with single-stranded DNA-specific photolyase activity. *Proc Natl Acad Sci U S A* **103**, 17696-17700

12.   Sancar, A. (2003) Structure and function of DNA photolyase and cryptochrome blue-light photoreceptors. *Chem Rev* **103**, 2203-2237

13.   Zhang, F., Scheerer, P., Oberpichler, I., Lamparter, T., and Krauss, N. (2013) Crystal structure of a prokaryotic (6-4) photolyase with an Fe-S cluster and a 6,7-dimethyl-8-ribityllumazine antenna chromophore. *Proc Natl Acad Sci U S A* **110**, 7217-7222

14.   Zhang, M., Wang, L., and Zhong, D. (2017) Photolyase: Dynamics and electron-transfer mechanisms of DNA repair. *Arch Biochem Biophys* **632**, 158-174

15.   Dikbas, U. M., Tardu, M., Canturk, A., Gul, S., Ozcelik, G., Baris, I., Ozturk, N., and Kavakli, I. H. (2019) Identification and Characterization of a New Class of (6-4) Photolyase from Vibrio cholerae. *Biochemistry* **58**, 4352-4360

16.   Geisselbrecht, Y., Fruhwirth, S., Schroeder, C., Pierik, A. J., Klug, G., and Essen, L. O. (2012) CryB from Rhodobacter sphaeroides: a unique class of cryptochromes with new cofactors. *EMBO Rep* **13**, 223-229

17.   Oberpichler, I., Pierik, A. J., Wesslowski, J., Pokorny, R., Rosen, R., Vugman, M., Zhang, F., Neubauer, O., Ron, E. Z., Batschauer, A., and Lamparter, T. (2011) A photolyase-like protein from Agrobacterium tumefaciens with an iron-sulfur cluster. *PLoS One* **6**, e26775

18.   Park, H. W., Kim, S. T., Sancar, A., and Deisenhofer, J. (1995) Crystal structure of DNA photolyase from Escherichia coli. *Science* **268**, 1866-1872

19.   Tamada, T., Kitadokoro, K., Higuchi, Y., Inaka, K., Yasui, A., de Ruiter, P. E., Eker, A. P., and Miki, K. (1997) Crystal structure of DNA photolyase from Anacystis nidulans. *Nat Struct Biol* **4**, 887-891

20.   Komori, H., Masui, R., Kuramitsu, S., Yokoyama, S., Shibata, T., Inoue, Y., and Miki, K. (2001) Crystal structure of thermostable DNA photolyase: pyrimidine-dimer recognition mechanism. *Proc Natl Acad Sci U S A* **98**, 13560-13565

21.   Hitomi, K., DiTacchio, L., Arvai, A. S., Yamamoto, J., Kim, S. T., Todo, T., Tainer, J. A., Iwai, S., Panda, S., and Getzoff, E. D. (2009) Functional motifs in the (6-4) photolyase crystal structure make a comparative framework for DNA repair photolyases and clock cryptochromes. *P Natl Acad Sci USA* **106**, 6962-6967

22.   Worthington, E. N., Kavakli, I. H., Berrocal-Tito, G., Bondo, B. E., and Sancar, A. (2003) Purification and characterization of three members of the photolyase/cryptochrome family blue-light photoreceptors from Vibrio cholerae. *J Biol Chem* **278**, 39143-39154

23.   Tardu, M., Bulut, S., and Kavakli, I. H. (2017) MerR and ChrR mediate blue light induced photo-oxidative stress response at the transcriptional level in Vibrio cholerae. *Sci Rep* **7**, 40817

24.   Faraji, S., and Dreuw, A. (2017) Insights into Light-driven DNA Repair by Photolyases: Challenges and Opportunities for Electronic Structure Theory. *Photochemistry and Photobiology* **93**, 37-50

25.   Matsubara, T., Araida, N., Hayashi, D., and Yamada, H. (2014) Computational Study on the Mechanism of the Electron-Transfer-Induced Repair of the (6-4) T-T Photoproduct of DNA by Photolyase: Possibility of a Radical Cation Pathway. *B Chem Soc Jpn* **87**, 390-399

26. Maul, M. J., Barends, T. R. M., Glas, A. F., Cryle, M. J., Domratcheva, T., Schneider, S., Schlichting, I., and Carell, T. (2008) Crystal Structure and Mechanism of a DNA (6-4) Photolyase. *Angew Chem Int Edit* **47**, 10076-10080

27. Hitomi, K., Nakamura, H., Kim, S. T., Mizukoshi, T., Ishikawa, T., Iwai, S., and Todo, T. (2001) Role of two histidines in the (6-4) photolyase reaction. *Journal of Biological Chemistry* **276**, 10103-10109

28. Li, J., Liu, Z., Tan, C., Guo, X., Wang, L., Sancar, A., and Zhong, D. (2010) Dynamics and mechanism of repair of ultraviolet-induced (6-4) photoproduct by photolyase. *Nature* **466**, 887-890

29. Hendrischk, A. K., Fruhwirth, S. W., Moldt, J., Pokorny, R., Metz, S., Kaiser, G., Jager, A., Batschauer, A., and Klug, G. (2009) A cryptochrome-like protein is involved in the regulation of photosynthesis genes in Rhodobacter sphaeroides. *Mol Microbiol* **74**, 990-1003

30. Diffraction, R. O. (2021) CrysAlisPro Software system, version 1.171.42.51a, Rigaku Corporation, Wroclaw, Poland.

31. Bienert, S., Waterhouse, A., de Beer, T. A., Tauriello, G., Studer, G., Bordoli, L., and Schwede, T. (2017) The SWISS-MODEL Repository-new features and functionality. *Nucleic Acids Res* **45**, D313-D319

32. Mccoy, A. J., Grosse-Kunstleve, R. W., Adams, P. D., Winn, M. D., Storoni, L. C., and Read, R. J. (2007) Phaser crystallographic software. *J Appl Crystallogr* **40**, 658-674

33. Adams, P. D., Afonine, P. V., Bunkoczi, G., Chen, V. B., Davis, I. W., Echols, N., Headd, J. J., Hung, L. W., Kapral, G. J., Grosse-Kunstleve, R. W., McCoy, A. J., Moriarty, N. W., Oeffner, R., Read, R. J., Richardson, D. C., Richardson, J. S., Terwilliger, T. C., and Zwart, P. H. (2010) PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-221

34. Emsley, P., and Cowtan, K. (2004) Coot: model-building tools for molecular graphics. *Acta Crystallogr D Biol Crystallogr* **60**, 2126-2132

35. DeLano, W. L. (2009) PyMOL molecular viewer: Updates and refinements. *Abstr Pap Am Chem S* **238**

Table 1: Statistics for data collection, processing and structure refinement

Resolution range	24.25-2.5 (2.589-2.5)
Space group	P6$_4$ 2 2
Unit cell	200.751 200.751 77.0231 90 90 120
Total reflections	1935882 (189833)

Unique reflections	32043 (3130)
Multiplicity	60.4 (60.6)
Completeness (%)	99.60 (99.97)
Mean I/σ	5.61 (1.70)
Wilson B-factor	15.97
R-merge	1.382 (3.765)
R-meas	1.394 (3.796)
R-pim	0.1776 (0.4855)
CC 1/2	0.955 (0.546)
CC*	0.988 (0.84)
Reflections used in refinement	31969 (3130)
Reflection used for R-free	1068 (105)
R-work	0.2254 (0.2891)
R-free	0.2830 (0.3620)
CC(work)	0.910 (0.712)
CC(free)	0.929 (0.542)
Number of non-hyrdogen atoms	4778
Macromolecules	4120
Ligands	101
solvent	557

Protein residues	507
RMS(bonds)	0.002
RMS(angles)	0.48
Ramachandran favored (%)	98.01
Ramachandran allowed (%)	1.99
Ramachandran outliers (%)	0.00
Rotamer outliers (%)	1.84
Clashscore	5.92
Average B-factor	22.46
macromolecules	22.31
Ligands	17.61
solvent	24.49
Number of TLS groups	8

A

B

Figure 1: Crystal structure and sequence comparison of Vc(6-4) PL. (A) Overview of Vc(6-4) PL structure, pink is α/β-domain, light blue is catalytic domain, and orange is the inter-domain linker region. Cofactor places are indicated by the black line. (B) Sequence alignment of bacterial (6-4) photolyases with available crystal structures and secondary structures of Vc(6-4) PL are shown. Red lines are α helixes and green lines are β sheets. Percentage identity coloring is used in Jalview software.

Figure 2: DMRL interactions and comparisons with other (6-4) photolyases. Every distance is in units of Armstrong (Å). (A) Electron cloud of DMRL at σA-weighted 2 F0-Fc electron density map contoured at 1.0σ in deepsalmon mesh and Vc(6-4) PL cartoon in splitpea color. (B) Possible interaction partners of DMRL. Yellow dashes represent hydrogen bondings, black dashes indicate Van der Waals interactions, and red dots are water molecules. (C) Additional water molecule that might stabilize the DMRL interaction. Deepteal, magenta, and deepsalmon colors indicate Vc(6-4) PL, At(6-4) PL (PDB ID: 4DJA) and Rs(6-4) PL (PDB ID: 3ZXS) respectively. Their superimposed structures have RMSD: 0.834 Å (At(6-4) PL), and RMSD: 0.876 Å (Rs(6-4) PL). Each dashed line represents hydrogen bonding of species with corresponding color.

A

B

C

Figure 3: **FAD interactions and comparisons with other (6-4) photolyas**es. Every distance is in units of Armstrong (Å). (A) Electron cloud of FAD at σA-weighted 2 F0-Fc electron density map contoured at 1.0σ in deepsalmon mesh, and Vc(6-4) PL cartoon in splitpea color. (B) Possible interaction partners of FAD. Yellow dashes represent hydrogen bondings, red dot is a water molecule. (C) Superimposed structure of Vc(6-4) PL and Dm(6-4) PL (PDB ID: 7AYV) at RMSD: 4.350 Å. Teal color represents Vc(6-4) PL and tv_orange color represents Dm(6-4) PL.

A

B

Figure 4: [4Fe-4S] cluster interactions. (A) Electron cloud of [4Fe-4S] cluster at σA-weighted 2 F0-Fc electron density map contoured at 1.0σ in deepsalmon mesh and Vc(6-4) PL cartoon in splitpea color. (B) Direct interactions of [4Fe-4S] cluster with surrounding amino acids of Vc(6-4) PL. Every distance is in units of Armstrong (Å).

Figure 5: Possible DNA binding site comparison with structurally known wildtype bacterial (6-4) photolyases. Possible DNA binding sites are drawn with electron cloud at σA-weighted 2 F0-Fc electron density map contoured at 1.0σ in deepsalmon mesh and (6-4) photolyase cartoon in splitpea color. (A) Vc(6-4) PL, (B) At(6-4) PL cryogenic (PDB ID: 4DJA), (C) At(6-4) PL ambient (PDB ID: 6DD6), (D) Rs(6-4) PL (PDB ID: 3ZXS). In (B), 3 amino acids are missing, which are Ala180, Glu181, Asn182, due to lack of electron cloud.

Figure 6: Possible DNA binding domain comparison with structurally known wildtype bacterial (6-4) photolyase. Each possible DNA binding site's electron cloud is drawn at σA-weighted 2 F0-Fc electron density map contoured at 1.0σ in deepsalmon mesh and (6-4) photolyase cartoon in splitpea color. (A) Vc(6-4) PL, (B) At(6-4) PL cryogenic (PDB ID: 4DJA), (C) At(6-4) PL ambient (PDB ID: 6DD6), (D) Rs(6-4) PL (PDB ID: 3ZXS). In (A), 8 amino acids are missing, which are Leu478, Ala479, Asn480, Asn481, Pro482, Arg483, Ile484, Gly485 due to lack of electron density cloud.

Figure 7: DNA binding comparison between Vc(6-4) PL and Dm(6-4) PL. (A) Generated surface density charge of Vc(6-4) PL, where blue is positive while red is negatively charged. Helixes with positive surface charge are indicated in the structure. (B) Superimposed (RMSD: 3.944 Å) image of Vc(6-4) PL and Dm(6-4) PL with (6-4) DNA lesion (PDB ID: 3CVU). Deepteal is for Vc(6-4) PL, while Tv_orange is for Dm(6-4) PL. (C) FAD comparison with superimposed image of Vc(6-4) PL and Dm(6-4) PL. All atoms RMSD between the structures for FAD cofactor is 0.7 Å. (D) B-factor analysis of C terminal of Vc(6-4) PL, where highest b-factor has yellowish color, while lowest b-factor has darker blue color. α23 has the highest b-factor in the structure.