Tropical high-altitude insects show limited capacity to handle high temperatures

[1]Harshad Vijay Mayekar, [2]Pooran Singh Solanki, [1]Homica Arya, [3]Rajaguru Aradhya,

[3,4]Prashanth Suravajhala, [5]Volker Loeschcke, and [1]Subhash Rajpurohit

1: Division of Biological and Life Sciences, School of Arts and Sciences, Ahmedabad

University, Navrangpura, Ahmedabad 380009 GJ, India

2: Department of Biotechnology and Bioinformatics, Birla Institute of Scientific Research,

Jaipur, 302001, India

3: Amrita School of Biotechnology, Amrita Vishwavidyapeetham, Amritapuri, Kerala 690525,

India

4: Bioclues.org, India

5: Department of Biology, Aarhus University, 8000 Aarhus C, Denmark

18 **Correspondence**: Subhash Rajpurohit and Volker Loeschcke

Abstract

Growing summer season and increased anthropogenic activities pose a continual challenge to resident species. Ectotherms like insects are especially vulnerable to rapid climatic changes. High-altitude tropical insect populations have been rarely examined for their responses to high-temperature. We exposed a tropical out-bred highland population of *Drosophila melanogaster* from the Himalayas to growing summer conditions in outdoor mesocosm units. Population response to thermal changes was tracked over ninety days at both phenotypic and genotypic level. Whole genomic resequencing data suggested a clear seasonal shift in allele frequencies. Interestingly, the general heat responsive genes were missing in the summer due to monsoon allele shift; an atypical response noted for high-altitude tropical populations. Instead, candidates involved in kinases and phosphorylation emerged as key players. Heat-knockdown time decreased over time indicating a limited ability to handle increasing temperature. Merging data from both allelic shifts and heat-knockdown time indicated a limited capacity for high-altitude insects in coping with climate warming.

Keywords: *Drosophila*, Heat, Climate change, Rapid adaptation, High altitudes

1. Introduction

Temperature is a dominant climatic variable with overarching effect on organismal fitness (Stevens 1989). Rising global temperatures raise alarms of extinction for several species which are unable to adapt to climate change at shorter time scales (Turney, Ausseil et al. 2020, Habibullah, Din et al. 2022). Rising temperatures aside, variability in temperatures is also a confounding factor increasing the complexity of assessing the impact of climate change (Duffy, Gouhier et al. 2022) as diverse zones face different climatic regimes. Temperature rise in tropics was considered to be lower than in temperate or polar regions, recent climate models strongly predict tropical zones of Amazonia, Sahel, South-east Asia and India to face increased warming (Holmes, Woollings et al. 2016, Bathiany, Dakos et al. 2018). In Amazonia the effect could be predominantly due to soil drying, but in the Asian tropics e.g. India, it is largely attributed to variation in atmospheric forces. Not surprisingly, the Indian monsoon patterns (Menon, Levermann et al. 2013), have varied greatly in recent years; with recurring floods supporting the correlation with rising temperatures. Yet, conclusions from climatic models could be taken with a caveats since tropical mappings are scarce (Bathiany, Dakos et al. 2018). Even then the global impact of climate change on living organisms cannot be ignored (Root, Price et al. 2003). In India, the increased incidence of heat-waves (1300 people died in 2010 in Ahmedabad alone (Knowlton, Kulkarni et al. 2014)) has been predicted to rise in several parts of north, central and western India with annual temperatures expected to rise from 2.2^0C to 5.5^0C (Kumar, Wiltshire et al. 2013, Dholakia, Mishra et al. 2015). Rising temperatures could be detrimental to survival of organisms and need research attention.

Anticipating thermal breadths could perhaps provide cues to organismal responses to rising temperatures. Janzen (1967) surmised broader thermal tolerance for temperate species than

tropical species since the tropics have evolutionarily experienced stable thermal ranges. Hence

seasonal effects are generally less pronounced in the tropics. However this does not include

places of high elevation/ altitude from the tropics which have cooler temperatures compared to

lowlands. Tropical altitudinal zones are distinct with little spatial overlap. Hence, populations

from higher altitudes could be expected to have responses different from lowlands. Whether high

altitudinal tropical populations mimic temperate/ polar zone patterns or display a different

pattern is not known as tropical species have been scarcely studied for their response to climate

change (Feeley, Stroud et al. 2017).

Ectotherms regulate their body temperatures based on the external environment (Hertz,

Huey et al. 1993) and are ideal to study thermal responses to changing climate and several

species are known to be impacted by climate (Duffy, Gouhier et al. 2022). Insects as notable

ectotherms; are seen to exhibit tremendous populations declines (Halsch, Shapiro et al. 2021).

Coping with climate change has been tested across several insects (Sears et al., 2011; Woods,

2013), but none parallel the diverse trends observed in the vinegar fly, *Drosophila melanogaster.*

D. melanogaster likely originated in Sub-Saharan Africa (Lachaise, Cariou et al. 1988) and

shortly attained a global distribution (David and Capy 1988). Within a few ten thousand years,

the sub-tropical *D. melanogaster* populations adapted worldwide to spatially diverse climatic

selection pressures, e.g. alcohol tolerance (Berry and Kreitman 1993, Fry, Donlon et al. 2008),

diapause (Schmidt, Zhu et al. 2008), circadian rhythm (Svetec, Zhao et al. 2015), thermal stress

and tolerance (Weeks, McKechnie et al. 2002), (Fallis, Fanara et al. 2014), and metabolic load

(Sezgin, Duvernell et al. 2004). However, whole genome sequencing offers more insights into

climatic adaptations in *D. melanogaster* rather than interpretations based on few genes, loci or

makers (Turner, Levine et al. 2008, Fabian, Kapun et al. 2012, Adrion, Hahn et al. 2015).

Surprisingly, genomic analyses revealed parallel trends in adaptation in north American and

Australian *D. melanogaster* populations compared to their southern counterparts respectively

(Reinhardt, Kolaczkowski et al. 2014). Not only spatial but also temporal heterogeneity due to

changing seasons could affect adaptation to climate change (Bergland, Behrman et al. 2014,

Rudman, Greenblum et al. 2022). Thermal variation could be experienced not only at the annual

scale, but also at the diel and within and across generations or even at the seasonal scale (Wang

and Dillon 2014, Kefford, Ghalambor et al. 2022). Typically, tracking thermal performance

curves (Sinclair, Marshall et al. 2016) or investigating traits exhibiting plastic responses to

thermal variation (Seebacher, White et al. 2015, Sgrò, Terblanche et al. 2016) have been

instrumental in understanding climate change. However, temporal genomic studies indicate

seasonally changing allele frequencies are predictable and adaptive (Bergland, Behrman et al.

2014, Machado, Bergland et al. 2021). It should however be noted that most of the spatial and/ or

temporal genomic studies of *Drosophila* thermal adaptations stem from temperate zones

(Rudman, Greenblum et al. 2022).

Tropical organisms living closer to their upper thermal limits could find the pace of

temperature rise challenging (Sørensen, Kristensen et al. 2016). Temperate *D. melanogaster*

populations respond to cold hardening through phenotypic plasticity (Stone, Erickson et al.

2020). Whether tropical ectotherms respond to thermal fluctuations through phenotypic plasticity

or local adaptation is not clear (Ghalambor, Huey et al. 2006). Phenotypic plasticity could either

buffer responses (Muñoz 2022) to climate change or even accelerate adaptive responses

(Harshman and Hoffmann 2000). Yet whether, plastic changes keep pace with variable climatic

patterns needs attention(Kapun and Flatt 2019). Limited capacity to handle thermal stress has

been hinted irrespective of latitude or elevation (Sunday, Bates et al. 2014). Yet, a broad

geographic sampling of insect populations from the tropics and their response patterns to climate

change are currently not known, especially from the Asian continent (particularly South-east

Asian region). Besides, high altitudinal tropical insect populations have rarely been addressed for

genomic changes in response to climate change.

Here, we investigate the genomic and physiological response of a high altitudinal tropical

D. melanogaster population from Western Himalayas to heat stress across a growing summer

season. *D. melanogaster* populations collected were outbred under laboratory conditions,

released into the mesocosm units in a tropical orchard set-up at the Ahmedabad University

Experimental Ecology and Evolution station with overlapping generations, and sub-sampled at

three different time points (TP's) (spanning a growing summer season with a duration of ninety

days. Samples were subjected to whole genome re-sequencing and we wanted to look for

genomic responses to warming i.e. changes in thermal responsive genes (TRGs) (Venkatachalam

and Montell 2007, Matsuura, Sokabe et al. 2009). Since *Drosophila* piggybacks between these

thermal preferences, particularly in the early life stages, state of relative conservation among

species and unrelated temperature changes at species origin was felt necessary to understand the

TRGs oscillations if any. At the physiological level, we tested the heat knock-down times of

populations collected from the three time points across the summer season. We aimed to

investigate the macro physiological pattern of a cold-adapted high altitudinal population to a

tropical growing summer.

2.   Materials and Methods

2.1   *Drosophila* collection and maintenance

D. melanogaster population was collected from higher altitudinal orchards from Western Himalayas (Rohru, Himachal Pradesh, India; 31.2046^0N, 77.7524^0E, Altitude 1554 meters) using banana baits in the year 2018. Each collected female was placed in a separate tube (iso-female line) at the site of collection. Almost 270 such lines were made in the field using females captured during the field trip. Upon 10 to15 hour's arrival to the lab, vials were placed at room temperature (25^0C). Successfully established lines were examined based on the taxonomic keys to reconfirm their identity at the species level. Overall, 40 lines of *D. melanogaster* were successful and were maintained in the laboratory, on standard 'agar-jaggery-yeast-maize' media at 25^0C, with a 12/ 12light/dark cycle. All the 40 lines were used in this work.

2.2   *Out-crossing*

Forty isofemale lines of *D. melanogaster* were outcrossed (founding numbers 800 adults: 400 males + 400 females) in a 0.61m X 0.61m X 0.61m dimension cage made up of plexiglass acrylic sheet. For this, 10 males and 10 females each from 40 isofemale lines were used. Each generation of eggs was collected in twenty half-pint size culture bottles. As soon as the pupae became darker and ready to eclose, bottles were moved to the plexiglass cage and plugs were removed. This allowed random mating in the founding base population. Once all the flies had enclosed in the cage, eggs were collected and transferred to fresh food bottles for the next generation. These lines were outcrossed for 5 generations keeping populations' size between 5-6 thousand individuals every generation.

2.3　*Mesocosm Cages*

To track adaptive changes in real time, and whether or not changes in temperature (increasing temperature over growing tropical summer) lead to changes in allelic frequencies, we tracked populations (3-fold replication) in outdoor mesocosm cages over a period of 90 days (Fig. 1). Outdoor (mesocosm) cages were set up at the Ahmedabad University Experimental Evolution Study Station. Each mesocosm was a 1.5x1.5x1.5 meters dimension (custom designed at Ahmedabad University Fabrication Shop) outdoor insect rearing enclosure with a mature (dwarf) Sapota tree. Three cages were used for this experiment (MESO-1, MESO-2, and MESO-3). Each cage was founded with 1000 males and 1000 females collected from the 6^{th} generation of laboratory cage (i.e. outcrossed population). Every morning at 09.00 am fresh food (50 ml of standard 'agar-jaggery-yeast-maize') in a half-pint bottle was placed in each enclosure for the entire duration of the experiment (15^{th} April 2020 to 15^{th} July 2020). Flies were allowed to oviposit on the fresh food for 24 hours. Each morning, bottles with eggs laid in the last 24 hours, were sealed with cotton plugs and larvae were allowed to develop (inside the same cages); upon eclosion, bottles were opened and adults were released into the cages. Thus populations were cultured under a natural regime of overlapping generations. Temperature and RH in the cages were recorded using HOBO U23 Pro v2 data loggers (Onset Computer Corp., Bourne, MA, USA).

Figure 1. Outdoor mescocosm experimental set-up. Forty *D. melanogaster* isofemale lines
were pooled to generate an outbred population. Lines were outcrossed for 5 generations in
24x24x24 inch dimension cages under laboratory conditions (25^0C). From this outcrossed
population (maintaining the size close to 5000-6000 individuals every generation) we released
500 males and 500 females in each outdoor mesocosm cages (MESO-1, MESO-2, and MESO-3).
For the next 90 days food was changed every day and the previous bottle with eggs was capped
and placed in the cage itself. Upon eclosion adults were released in the existing cage and bottles
were discarded. Cages were sampled at three time-points (TP) for genomic and phenotypic
analysis: at the beginning (TP1), in the middle (TP2) and at the end (TP3).

2.4 *Adaptive tracking of populations*

Samples for genomic and phenotypic measurements were collected at three time-points
for adaptive tracking. Samples for TP1 were collected from the founding adults while the TP2
and TP3 samples were collected on the 45th and 90th day, respectively. Eggs were collected in

30 bottles at lower density (10 bottles for each replicate cage). These bottles were allowed to

develop under a constant temperature (25°C) in the laboratory and phenotypic data was collected

using F3 individuals.

2.5 *DNA sample preparation for genomic analysis*

Pooled genome re-sequencing was performed with 100 *D. melanogaster* females in

triplicate from each outdoor cage, a total of nine samples were collected for whole genome

sequencing analysis. The DNA extraction procedures involved homogenizing the flies (n = 100)

in 200 µL of lysis buffer (100 mM Tris-Cl, 100 mM EDTA, 100 mM NaCL, 0.5% SDS). The

homogenate was incubated at 65°C for 30 minutes. Following this, proteins were precipitated by

adding 800 µL of 2 parts 5M potassium acetate, 5 parts of 6M lithium chloride solution and 15

minutes ice incubation. The supernatant was collected after centrifugation of the homogenate at

12K RPM for 15 minutes at room temperature. DNA was precipitated by adding 800 µL of

isopropanol and centrifugation at 12K RPM for 15 minutes. The pellet was collected and washed

with 70% ethanol. After removing the ethanol, the pellet was allowed to dry at room temperature

and was re-suspended in 100 µL of TE buffer.

2.6 *Library construction and quality control*

1µg of DNA was used as input material for the DNA sample preparations. Sequencing

libraries were generated using NEBNext® DNA Library Prep Kit following manufacturer's

recommendations and indices were added to each sample. The genomic DNA was randomly

fragmented to a size of 350bp by shearing, then DNA fragments were end polished, A-tailed, and

ligated with the NEBNext adapter for Illumina sequencing, and further PCR enriched by P5 and

indexed P7 oligos. The PCR products were purified (AMPure XP system) and resulting libraries

were analyzed for size distribution by Agilent 2100 Bioanalyzer and quantified using real-time

PCR.

2.7 *Next Generation Sequencing*

The whole genome sequencing (WGS) of natural populations was done to check the role

of mutations in functional candidates. Library preparation and tagmentation was done using

DNA-350 bp by default, while the sequencing was performed on a HiSeq PE150. The samples

were mixed for library construction using a PCR-free library preparation guide. The downstream

analysis and annotation was done with raw reads run through an in-house benchmarked pipeline

(Meena, Mathur et al. 2018), tweaked for a whole exome pipeline

(https://github.com/prashbio/WES).

After the allele frequencies were mapped, we read all mutation positions in MESO-1,

MESO-2 and MESO-3 for all the three time point scales (T1|T2|T3) and created a data structure

which contained three types of positions: single, double and triple mutations. For example,

single types of mutations are those that are positioned only once across the 3 different TPs,

double- any two TP scales while triple – across all three of them. This was done for MESO-1,

MESO-2 and MESO-3 (experimental scale) and the resulting files were checked for interpreting

the possible SNPs. However, we asked whether mutations that are identical are present at least

at 2 time points at the same position and finally a composite matrix containing mutational

positions of MESO-1, MESO-2, and MESO-3 was done.

2.8. *Heat-knockdown assay*

We tested F3 females (ca. 3 days old) for their thermal tolerance to an increasing

temperature using the dynamic heat ramp assay. In this assay, temperatures ecologically relevant

to the species are the base-point after which they are steadily increased till the fly is knocked

down (Lutterschmidt and Hutchison 1997). Individual flies were collected in glass tubes of

$0.05m^3$ volume. The tubes were immersed in a water bath set at room temperature (ca. 25°C).

Temperature was increased at the rate of 1°C/ minute with the upper thermal limit being 43°C.

We measured three technical replicates with 24 flies each from three cages across three time

points. Nested ANOVA (experimental cages within time-points) was used to assess the variation

in knock-down time. Post-hoc analysis was performed using the *glht* () function from the

"multcomp" package (Hothorn, Bretz et al. 2008). For genomic analysis, the flies were aspirated

from the cages itself.

3. Results

3.1. Allele frequencies

The MESO-1, 2, & 3 replicates were run through the whole genome sequencing (WGS)

pipeline where we inferred causal SNPs across these sub-populations (Table 1). The minor allele

frequencies were tabulated from the depth with number of reads showing variation (DP4) with

1) forward ref alleles; 2) reverse ref; 3) forward alt; 4) reverse alt alleles, used in variant calling

using the formula, variant allele frequency(VAF) = (forward alt + reverse alt alleles) / (forward

ref alleles + reverse ref + forward alt+ reverse alt alleles). After setting up the MAF<=0.15

assuming that 15% of the SNPs would have this MAF, we further retained heterozygous SNPs.

All 9 raw reads yielded 9,963,857 SNPs per sample. The most common SNPs between the

MESO-1, 2, & 3 cages and the outliers were tabulated by mapping the chromosomal positions to

flybase reference genome (flybase.org last accessed date: September 22, 2021) (Table 1/Fig. 2).

By removing those specific minor allele frequencies (MAFs) above 0.15, we could assume that

there is a clear distribution of contamination. We therefore sought to assess the possibility of

observed TRG with MAF <0.15 which gave us an average of 240 SNPs per sample

(Supplementary File 1). This raises the possibility that allele frequency could be generated by *D.*

melanogaster alleles consistently at various time point scales. We further assessed the frequency

of SNPs that are fixed for alternate alleles and cross checked this with the previously identified

SNPs associated with seasonal changes (Temperature) and tabulated all SNPs for time points

(Table 1).

Table 1: Average number of SNPs after filtering

Total identified SNPs/ Average SNPs per TP sample >=10 DP <=400	89,674,715/9,963,857 5,015,244
Total number of SNPs with MAF <0.15/ Average MAF <0.15/sample	3,573,280/397,031
Total used in analysis/ Average bona fide SNPs per TRG	5,015,244/ 240

a)   b)

Figure 2. (a) Flybase SNPs mapped to our experimental candidates across all chromosomes (b)

The SNPs mapped across the three time point experimental pools. The numbers indicate the

individual time point SNPs apart from the most common SNPs at the intersection.

3.2. Positional mapping with Flybase SNPs

The downstream annotation was processed with chromosomal alias obtaining 1870 total

number of scaffolds with an average length of 17,195,935 per chromosome. The chromosomal

distribution of SNPS is largely at the interface of 2L and X (Data not shown; Supplementary File

2 1, Fig. 3). We also mapped the SNPs to that of reported 193 SNPs from flybase to check if

TRG mutations exist. We found Single TP SNPs: (n=noPOS: 4,666), Double TP SNPs: POS:

(n=2,536) and no Triple TP SNPs (n=POS: 0).The final list of protein coding genes were mapped

across various chromosomes (Supplementary File 1). As the current analysis represented

289 identifying common mutations, we further looked into mapping the existing known SNPs (for

290 heat stress).

3.3. Population comparisons

The time point samples and replicates mapped to the temperature variation across season

regime were treated as T1_MESO-1, T2_MESO-1 and T3_MESO-1 as populations. As we

obtain the data, the coverage across these three sub-populations were filtered for common

mutations or SNPs that have been retained from T1_MESO-1,2&3 through T3_MESO-1,2&3 or

the unique set of SNPs through them (Supplementary File 1). While removing the SNPs with

total counts with the same allelic proportion, we deem that they have not undergone any changes

through the three experiments. On the contrary, those SNPs that have an overlap with changes in

SNPs across the same positions were filtered and we therefore limited the downstream analysis

to such overlapping SNPs from three sub-populations or experiments. As we combined all the

reads, we have checked the significant differences between the experiments for overlapping

SNPs using the chi-square test as a standard practice. The *p*-value heuristics resulting from the

analysis based on three population comparisons were in agreement. Another approach was to

check the SNP mean density which could be done using an existing method (proposed by Burke

et al (2010) and Winbush and Singh (2021) which measures the quantile score but our study is

not a measure of nucleotide diversity from a threshold value instead the number of SNPs

emerging from a whole genome sequencing study.

While we have found SNPs that have been deleted from T1 to T3 (Supplementary File 3),

the MAFs in earlier cases were set to as big as MAF<0.89 which is 89%. Our stringent cutoff

(Winbush and Singh 2021) indicates that highly sensitive SNPs emerge from our analysis and

when these were tried to map to flySNP base, none or not many of them are mapped indicating

that these are novel (data not shown). Nevertheless, we lacked sequencing data on similar scales

where whole genome sequencing (WGS) was done so that we could have identified convergent

or divergent regions contributing between the three sub-populations. For each comparison, we

observed that 77,915 (24.97%), 71,938 (23.06%) and 27,664 (8.8%) SNPs in the respective three

time points existed indicating that the temperature affects the sensitivity of the flies. We then

mapped these SNPs to summarize the genes that have escaped the mutations in T3_MESO-1, 2

&3. When we map them to the table depicting counts for genes associated with these SNPs for

T3_MESO-1, 2 & 3, we observe that a large number of diseased phenotypes are associated with

these conditions. However, despite consistent read depth of coverage and SNPs, we observe a

partial overlap suggesting that the accorded convergence can be attributed to slightly reduced

heterozygosity which may invite a bias. However, this was reconfirmed from our validation set

of founder experiments.

As we identified the genes with diseased phenotypes we were interested to see whether or

not the candidate loci of these genes have an influence on the pathways. To check this, our

panther gene ontology annotation derived enriched genes unique to temperature. In summary,

our analysis identified a large number of candidate genes and SNPs associated with heat

handling wiring machinery. Rather, our approaches indicate the variation across different

subpopulations, as we argue that several diseased phenotypes have emerged from our analysis.

Taken together, temperature sensitivity plays a very important role and serves as a determinant to

mesocosm experimental study. Although genetic drift augurs well for taking temperature as a

context, functional validation for observational variation is presumptive for maintaining such

selection pressure. It may be argued that the function of the genes stemming from allelic

variation has allowed us to determine candidate TRGs.

Figure 3. Variable polymorphisms shown to have a large deviation from time point scale 1 to

time point 3 taking into account read depth and the number of sampled chromosomes. The X-

axis shows the chromosomes with the Y-axis showing the single nucleotide polymorphism

(SNP) positions.

To check genomic signatures affected by seasonal fluctuations, we pooled sequencing

data containing the SNPs by estimating the allele frequencies (AF). From each population and

time point, the three replicate cages were merged and the files were segregated to average read

depth of 20–200× coverage (Supplementary File 1). We observed that these estimated AFs have

been shown to be accurate for understanding the magnitude of genetic variation through the time

point scales. From Fig. 2, we reason that variable polymorphisms had a large deviation from time

point scale 1 to time point 3 taking into account read depth and the number of sampled

chromosomes. Of the 3,119,540 common SNPs tested, we identified substantial numbers falling

between the statistical values, meaning they are called integrated thermal responsive SNPs.

However, the AFs corresponding to selection coefficients per generation were not done with this

statistical power as they are variable. Our rationale was to assess evolutionary features

underlying rapid adaptive states in response to selection pressure. As the data is variable with

the random disturbance and is different across elements of the vector, we asked whether the

SNPs are enriched among functional genetic elements. To find out whether or not the SNPs are

genic, we screened total SNPs present on specific genes common to these time points. A putative

interaction map indicates that the genes are largely spread across with the key pathways

attributing to these factors are MAP kinases and other important signaling pathways (Fig. 4). In

general, we find four important genes and a few un-characterized genes which could be

attributed to pleiotropy and linkage disequilibrium largely due to TRGs/seasonal variation.

Figure 4. A putative interaction map of *Svil*, *Myc* and *MAP205* with interacting partners connected through edges (lines). The pink edges indicate that they are known to be physically interacting with each other while other edge colors denote pathways, domain association, co-localization, neighborhood and genetic interactions if any.

3.4. Heat tolerance

369   Knock-down times of flies significantly differed among all three time-points (df = 2, F= 966.30, P < 0.0001) (Supplementary File 4). Heat knockdown time was significantly lower for flies from T2 (middle time-point) (median = 60.38 minutes, interquartile range (IQR) = 4.55, N = 72) in comparison to early (T1) (median = 73.22 minutes, IQR = 3.07, N = 72) (estimate = -13.151, t = -27.115, P < 0.0001) and late (T3) (median = 65.39 minutes, IQR = 3.71, N = 72)

(estimate = 4.275, t = 8.814, P < 0.0001) time-points. Variation in heat knockdown response is

partially explained by variation in experimental cages too (df = 6, F= 42.12, P< 0.0001) (Fig. 5).

Figure 5: Box-plots representing variation in knock-down times in response to heat ramping

assays, across three time-points (early, middle and late) from three replicate experimental

cages. Flies from the middle time-point (45[th] day) display the significantly lowest tolerance

to increase in temperatures compared to the early (day 0) and late (90[th] day) time-points.

4. Discussion

An organisms' response to changing environmental conditions determines its fitness.

Adaptations to rapid environmental fluctuations are known (Franks and Hoffmann 2012).

Responses to rapid environmental change could be micro-evolutionary or mediated through

phenotypic plasticity (Meyers and Bull 2002, Sgrò, Terblanche et al. 2016). Trait plasticity in

response to climate change has been documented across diverse taxa, e.g. in birds (Charmantier,

McCleery et al. 2008), mammals (Nussey, Wilson et al. 2007), reptiles (Leal and Gunderson

2012) and insects (Boggs 2016, Halsch, Shapiro et al. 2021). Still, studies investigating response

to climate change have been attempted predominantly for temperate (Finn, Grattarola et al. ,

Feeley, Stroud et al. 2017, Sheldon 2019) rather than tropical species (especially south-east Asia)

(but see (Parkash, Tyagi et al. 2005, Rajpurohit, Zhao et al. 2017)). *D. melanogaster* originating

from sub-Saharan Africa showing cosmopolitan distribution worldwide (David and Capy 1988,

Lachaise, Cariou et al. 1988, Sprengelmeyer, Mansourian et al. 2019) has been investigated in

recent years for responses to climate changes (Hoffmann, Hallas et al. 2003). The role of natural

selection in latitudinal clinal variation in *D. melanogaster* across broad demographic zones of

tropics and temperate regions is known (Kolaczkowski, Kern et al. 2011). However, genomic

analyses for climatic adaptation have been studied in North American (Fabian, Kapun et al.

2012), European (Božičević, Hutter et al. 2016) African (Pool, Corbett-Detig et al. 2012,

Sprengelmeyer, Mansourian et al. 2020) and Australian (Hangartner, Hoffmann et al. 2015) *D.*

melanogaster populations. Parallel adaptations to cold climatic conditions are recorded for fly

populations from Africa, Europe and North America (Božičević, Hutter et al. 2016, Pool, Braun

et al. 2017). Here, we investigated the response of *D. melanogaster* collected from colder regions

of the western Himalayas to a growing summer season in outdoor cages (mesocosm). Thus, our

population was sourced from a high altitudinal zone (where maximum temperatures do not

escalate above 15^0C) hence cold adapted, but subjected to a warm/ hotter summer season

(maximum temperatures can soar to 45^0C). We expected thermos-sensitive genes to be

upregulated in response to the growing season. However, we found no such pattern. Laboratory

experiments demonstrate that *D. melanogaster* flies adapting to fluctuating thermal regimes

increase their thermal breadth. Thus, *D. melanogaster* lines exposed to alternate cycles of high

and low temperatures exhibited a generally tolerance to both cold and heat extremes (Tobler,

Hermisson et al. 2015). Cold hardened flies survived better than non-hardened flies after a heat-

stressor treatment. Tolerance to heat could be due to the activation of heat-shock proteins

(Hsp70) which could have been inactive at low temperatures initially (Sejerkilde, Sørensen et al.

2003). Contrarily, cold adapted population from the Himalayas did not exhibit strong tolerance

to summer conditions (Fig. 5) probably because selection for a colder climate could have

reduced the physiological capacity to tolerate higher temperatures.

We used whole genome sequencing and filtered common SNPs for TRGs across

the three time-points (T1 to T3). Results demonstrate considerable allelic variation across the

three time-points. Temperature would increase from T1 to T2 and then decrease from T2 to T3.

We therefore surmise that allelic variation could execute through physiological plasticity in

response to temperature extremes in this tropical species. Higher allelic variation was observed

in the first two time-points in comparison to the third (T3), where temperature could plateau due

to the onset of monsoon. Further studies could determine if the observed allelic variation

corroborates positively with survival and fitness in the peak hot season. In temperate orchards,

considerable allelic variation with respect to seasons (winter versus fall) has been noted

(Bergland, Behrman et al. 2014) maintained through balancing selection. However, in this case,

considerable allelic variation for temperature responsive genes was seen within a span of 90 days

(corresponding to peak summer season). It therefore remains to be analyzed if selection is indeed

strong, relaxed or balancing through seasonal oscillations like temperate counterparts.

While the mechanistic basis of temperature regulating TRGs is still being explored

(Singh, McGoldrick et al. 2019), our study which reveals novel SNP's in *Drosophila* TRGs

opens up new avenues for testing thermal acclimation hypotheses. The pleiotropic effect of

temperature sensation on other functions can be seen through the association of different

pathways integrated through the protein interaction map. Nonetheless, genes associated are novel

and not typical of those associated with thermal acclimation, e.g. heat shock proteins (Sørensen

and Loeschcke 2007).

We found two independent lines of evidence: First, a large number of kinases and

phosphorylases are associated with stress, possibly linked to heat/temperature as evident from

the protein interaction map (Fig. 4). Second, the large number of key regulators and transcription

factors that are associated with cell proliferation and growth. Underlying these, *Myc* emerged as

a key gene with the interactants possibly associated with genetic arrangements, and most

importantly reducing the mutational load or burden. This is also in agreement with the fact that

the *Myc* plays a key role in insufficiency of haplotypes reducing the mutation load which further

is the key factor for extended lifespan or pro-aging (Morrison, Murakami et al. 2000, Greer, Lee

et al. 2013). On the other hand, phosphatidylinositol 5-phosphate 4-kinase (PIP4K) is known to

regulate the growth during fly development (Gupta, Toscano et al. 2013). While its activity is

implicated in cellular responses, regulating the growth factors, the protein interaction networks

clearly reveal that there are transient mutations underpinning the changes in heat/temperature

stress as these changes could be because of the activity of downstream kinases. We envisage that

the pathways associated with PIP4K switch signals modulating the strength of a signal. Taken

together, the kinases and phosphatases contribute to the outcome or ability to acclimate the flies

in these conditions. As they adapt, essential regulatory processes and changes mediating

phosphorylation and kinases are known. Further, our genomic sequencing has emerged as an

approach to understand such "adaptive loci" influencing these signaling pathways.

Mesocosm experiments as ours closely resemble natural conditions are hitherto reported only in temperate populations (e.g. (Rudman, Greenblum et al. 2022)) but not for tropical species. Ours is therefore the first study to track genomic genetic variation at more realistic and natural time-scales in a tropical high altitudinal fruit fly species. We considered a time frame where summer temperature steadily increases, peaks and then wanes. Our results from heat ramp assay corroborate this, in that increase in temperatures of the external environment (cages) also altered the thermal sensitivity. Thus, the peak in the temperature during T2 (Fig. 1) also corresponded with flies from T2 being the most sensitive to thermal stress and hence the faster knock-down time in T2 compared to T1 and T3 (Fig. 5). However, further studies could ascertain if seasonal variation in alleles related to TRGs are connected/ pleiotropic with fitness-related traits. Thus, in the temperate regions, diapause associated genes are known to be up and down regulated corresponding to winter and fall seasonal variation, respectively (Zhao, Bergland et al. 2016) with outcomes in fecundity (ovariole size (Schmidt, Matzkin et al. 2005)). In tropical scenarios, where adaptations like diapause are less known, one could expect sensitivity towards thermal extremes to evolve through alternate gene regulatory networks. Since *Drosophila melanogaster* originated from Africa (David and Capy 1988) and then dispersed to other continents, reaction norms for temperature tolerance could be expected to be broadly similar across the tropics than across temperate regions. Surprisingly, our study populations from higher altitudes (Rohru, western Himalayas) exhibited limited tolerance to higher temperatures, thus re-affirming the speculation that tropical species are indeed sensitive to higher temperature thresholds. It remains to be tested if differential response to temperature extremes does exist across altitudinal populations. Given clinal variation for multiple traits noted for Drosophilids

from the Indian subcontinent (Rajpurohit, Zhao et al. 2017) (Rajpurohit, Oliveira et al. 2013,

Rajpurohit, Zhao et al. 2017), exploring thermal tolerance and associated genomic variation is an

interesting research avenue. Further investigations could involve understanding chromosomal

inversion clines which have been linked with local adaptation in *D. melanogaster* (Kapun,

Fabian et al. 2016, Kapun and Flatt 2019). Inversion clines are common across all continents

where *D. melanogaster* has dispersed, yet they have been less addressed from south-east Asia

(Das and Singh 1990) where the inversion frequency is speculated to be low (Glinka, Stephan et

al. 2005). Surprisingly, inversion polymorphisms linked with climate adaptation were found to

be less frequent at higher altitudes than lowlands in Africa populations (Aulard, David et al.

2002, Pool, Braun et al. 2017). Does the ancestral (sub-Saharan) derived inversion

polymorphism confer selective advantage to tropical high altitudinal *D. melanogaster*? Whether

south-east Asian flies evolved alternative loci/ gene networks as our study reveals needs further

exploration.

Conclusions

Tropical species are likely to be affected by global increase in temperature since the

thermal range to which they could respond is narrow compared to their temperate counterparts.

Our study hints that the narrow temperature tolerance of tropical species could be linked with

alternate molecular pathways for thermal sensitivity. We found a considerable number of newer

candidate alleles whose frequencies are significantly shifting as temperature increases over the

growing season. Our study highlights that genomic signatures could be instrumental to

understand organismal responses to changing environmental conditions. Our findings of allelic

variation have implications not only in tracking fitness outcomes across seasons but also how

organisms are evolved in the face of rapid climate change. Future work could combine molecular

and field based approaches for a comprehensive understanding of climate mediated changes.

Competing interests: None to declare.

References

Adrion, J. R., M. W. Hahn and B. S. Cooper (2015). "Revisiting classic clines in Drosophila

melanogaster in the age of genomics." Trends Genet **31**(8): 434-444.

Aulard, S., J. R. David and F. Lemeunier (2002). "Chromosomal inversion polymorphism in

Afrotropical populations of Drosophila melanogaster." <u>Genetics Research</u> **79**(1): 49-63.

Bathiany, S., V. Dakos, M. Scheffer and T. M. Lenton (2018). "Climate models predict

increasing temperature variability in poor countries." <u>Science Advances</u> **4**(5): eaar5809.

Bergland, A. O., E. L. Behrman, K. R. O'Brien, P. S. Schmidt and D. A. Petrov (2014).

"Genomic Evidence of Rapid and Stable Adaptive Oscillations over Seasonal Time Scales in

Drosophila." <u>PLOS Genetics</u> **10**(11): e1004775.

Berry, A. and M. Kreitman (1993). "Molecular analysis of an allozyme cline: alcohol

dehydrogenase in Drosophila melanogaster on the east coast of North America." <u>Genetics</u>

134(3): 869-893.

Boggs, C. L. (2016). "The fingerprints of global climate change on insect populations." <u>Current</u>

<u>Opinion in Insect Science</u> **17**: 69-73.

Božičević, V., S. Hutter, W. Stephan and A. Wollstein (2016). "Population genetic evidence for

cold adaptation in European Drosophila melanogaster populations." <u>Molecular ecology</u> **25**(5):

1175-1191.

Burke, M. K., J. P. Dunham, P. Shahrestani, K. R. Thornton, M. R. Rose and A. D. Long (2010).

"Genome-wide analysis of a long-term evolution experiment with Drosophila." <u>Nature</u>

467(7315): 587-590.

Charmantier, A., R. H. McCleery, L. R. Cole, C. Perrins, L. E. Kruuk and B. C. Sheldon (2008).

"Adaptive phenotypic plasticity in response to climate change in a wild bird population." <u>Science</u>

320(5877): 800-803.

Das, A. and B. N. Singh (1990). "Chromosome inversions in Indian Drosophila melanogaster."

<u>Genetica</u> **81**(2): 85-88.

David, J. R. and P. Capy (1988). "Genetic variation of Drosophila melanogaster natural

populations." Trends in Genetics **4**(4): 106-111.

Dholakia, H. H., V. Mishra and A. Garg (2015). "Predicted increases in heat related mortality

under climate change in urban India."

Duffy, K., T. C. Gouhier and A. R. Ganguly (2022). "Climate-mediated shifts in temperature

fluctuations promote extinction risk." Nature Climate Change **12**(11): 1037-1044.

Fabian, D. K., M. Kapun, V. Nolte, R. Kofler, P. S. Schmidt, C. Schlötterer and T. Flatt (2012).

"Genome-wide patterns of latitudinal differentiation among populations of Drosophila

melanogaster from North America." Molecular Ecology **21**(19): 4748-4769.

Fallis, L. C., J. J. Fanara and T. J. Morgan (2014). "Developmental thermal plasticity

among*Drosophila melanogaster*populations." Journal of Evolutionary Biology **27**(3): 557-564.

Feeley, K. J., J. T. Stroud and T. M. Perez (2017). "Most 'global' reviews of species' responses

to climate change are not truly global." Diversity and Distributions **23**(3): 231-234.

Finn, C., F. Grattarola and D. Pincheira-Donoso "More losers than winners: investigating

Anthropocene defaunation through the diversity of population trends." Biological Reviews

n/a(n/a).

Franks, S. J. and A. A. Hoffmann (2012). "Genetics of climate change adaptation." Annu Rev

Genet **46**: 185-208.

Fry, J. D., K. Donlon and M. Saweikis (2008). "A worldwide polymorphism in aldehyde

dehydrogenase in Drosophila melanogaster: evidence for selection mediated by dietary ethanol."

Evolution **62**(1): 66-75.

Ghalambor, C. K., R. B. Huey, P. R. Martin, J. J. Tewksbury and G. Wang (2006). "Are

mountain passes higher in the tropics? Janzen's hypothesis revisited." Integr Comp Biol **46**(1): 5-

17.

Glinka, S., W. Stephan and A. Das (2005). "Homogeneity of common cosmopolitan inversion

frequencies in Southeast Asian Drosophila melanogaster." J Genet **84**(2): 173-178.

Greer, C., M. Lee, M. Westerhof, B. Milholland, R. Spokony, J. Vijg and J. Secombe (2013).

"Myc-Dependent Genome Instability and Lifespan in Drosophila." PLOS ONE **8**(9): e74641.

Gupta, A., S. Toscano, D. Trivedi, D. R. Jones, S. Mathre, J. H. Clarke, N. Divecha and P.

Raghu (2013). "Phosphatidylinositol 5-phosphate 4-kinase (PIP4K) regulates TOR signaling and

cell growth during Drosophila development." Proc Natl Acad Sci U S A **110**(15): 5963-5968.

Habibullah, M. S., B. H. Din, S.-H. Tan and H. Zahid (2022). "Impact of climate change on

biodiversity loss: global evidence." Environmental Science and Pollution Research **29**(1): 1073-

1086.

Halsch, C. A., A. M. Shapiro, J. A. Fordyce, C. C. Nice, J. H. Thorne, D. P. Waetjen and M. L.

Forister (2021). "Insects and recent climate change." Proceedings of the National Academy of

Sciences **118**(2): e2002543117.

Hangartner, S. B., A. A. Hoffmann, A. Smith and P. C. Griffin (2015). "A collection of

Australian Drosophila datasets on climate adaptation and species distributions." Scientific Data

2(1): 150067.

Harshman, L. G. and A. A. Hoffmann (2000). "Laboratory selection experiments using

Drosophila: what do they really tell us?" Trends in Ecology & Evolution **15**(1): 32-36.

Hertz, P. E., R. B. Huey and R. D. Stevenson (1993). "Evaluating temperature regulation by

field-active ectotherms: the fallacy of the inappropriate question." Am Nat **142**(5): 796-818.

Hoffmann, A., R. Hallas, J. Dean and M. Schiffer (2003). "Low potential for climatic stress

adaptation in a rainforest Drosophila species." Science **301**(5629): 100-102.

Holmes, C. R., T. Woollings, E. Hawkins and H. De Vries (2016). "Robust future changes in

temperature variability under greenhouse gas forcing and the relationship with thermal

advection." Journal of Climate **29**(6): 2221-2236.

Hothorn, T., F. Bretz and P. Westfall (2008). "Simultaneous inference in general parametric

models." Biom J **50**(3): 346-363.

Janzen, D. H. (1967). "Why mountain passes are higher in the tropics." The American Naturalist

101(919): 233-249.

Kapun, M., D. K. Fabian, J. Goudet and T. Flatt (2016). "Genomic evidence for adaptive

inversion clines in Drosophila melanogaster." Molecular biology and evolution **33**(5): 1317-

1336.

Kapun, M. and T. Flatt (2019). "The adaptive significance of chromosomal inversion

polymorphisms in Drosophila melanogaster." Molecular ecology **28**(6): 1263-1282.

Kefford, B. J., C. K. Ghalambor, B. Dewenter, N. L. Poff, J. Hughes, J. Reich and R. Thompson

(2022). "Acute, diel, and annual temperature variability and the thermal biology of ectotherms."

Global Change Biology **28**(23): 6872-6888.

Knowlton, K., S. P. Kulkarni, G. S. Azhar, D. Mavalankar, A. Jaiswal, M. Connolly, A. Nori-

Sarma, A. Rajiva, P. Dutta and B. Deol (2014). "Development and implementation of South

Asia's first heat-health action plan in Ahmedabad (Gujarat, India)." International journal of

environmental research and public health **11**(4): 3473-3492.

Kolaczkowski, B., A. D. Kern, A. K. Holloway and D. J. Begun (2011). "Genomic

differentiation between temperate and tropical Australian populations of Drosophila

melanogaster." Genetics **187**(1): 245-260.

Kumar, P., A. Wiltshire, C. Mathison, S. Asharaf, B. Ahrens, P. Lucas-Picher, J. H. Christensen,

A. Gobiet, F. Saeed and S. Hagemann (2013). "Downscaled climate change projections with

uncertainty assessment over India using a high resolution multi-model approach." Science of the

Total Environment **468**: S18-S30.

Lachaise, D., M.-L. Cariou, J. R. David, F. Lemeunier, L. Tsacas and M. Ashburner (1988).

"Historical biogeography of the Drosophila melanogaster species subgroup." Evolutionary

biology: 159-225.

Leal, M. and A. R. Gunderson (2012). "Rapid Change in the Thermal Tolerance of a Tropical

Lizard." The American Naturalist **180**(6): 815-822.

Lutterschmidt, W. and V. Hutchison (1997). "The critical thermal maximum: Data to support the

onset of spasms as the definitive end point." Canadian Journal of Zoology **75**: 1553-1560.

Machado, H. E., A. O. Bergland, R. Taylor, S. Tilk, E. Behrman, K. Dyer, D. K. Fabian, T. Flatt,

J. González, T. L. Karasov, B. Kim, I. Kozeretska, B. P. Lazzaro, T. J. S. Merritt, J. E. Pool, K.

O'Brien, S. Rajpurohit, P. R. Roy, S. W. Schaeffer, S. Serga, P. Schmidt and D. A. Petrov

(2021). "Broad geographic sampling reveals the shared basis and environmental correlates of

seasonal adaptation in Drosophila." eLife **10**: e67577.

Matsuura, H., T. Sokabe, K. Kohno, M. Tominaga and T. Kadowaki (2009). "Evolutionary

conservation and changes in insect TRP channels." BMC Evol Biol **9**: 228.

Meena, N., P. Mathur, K. Medicherla and P. Suravajhala (2018). <u>A Bioinformatics Pipeline for</u>

<u>Whole Exome Sequencing: Overview of the Processing and Steps from Raw Data to</u>

<u>Downstream Analysis</u>.

Menon, A., A. Levermann and J. Schewe (2013). "Enhanced future variability during India's

rainy season." <u>Geophysical Research Letters</u> **40**(12): 3242-3247.

Meyers, L. A. and J. J. Bull (2002). "Fighting change with change: adaptive variation in an

uncertain world." <u>Trends in Ecology & Evolution</u> **17**(12): 551-557.

Morrison, D. K., M. S. Murakami and V. Cleghon (2000). "Protein Kinases and Phosphatases in

the Drosophila Genome." <u>Journal of Cell Biology</u> **150**(2): F57-F62.

Muñoz, M. M. (2022). "The Bogert effect, a factor in evolution." <u>Evolution</u> **76**(s1): 49-66.

Nussey, D. H., A. J. Wilson and J. E. Brommer (2007). "The evolutionary ecology of individual

phenotypic plasticity in wild populations." <u>J Evol Biol</u> **20**(3): 831-844.

Parkash, R., P. K. Tyagi, I. Sharma and S. Rajpurohit (2005). "Adaptations to environmental

stress in altitudinal populations of two Drosophila species." <u>Physiological Entomology</u> **30**(4):

353-361.

Pool, J. E., D. T. Braun and J. B. Lack (2017). "Parallel Evolution of Cold Tolerance within

Drosophila melanogaster." <u>Mol Biol Evol</u> **34**(2): 349-360.

Pool, J. E., R. B. Corbett-Detig, R. P. Sugino, K. A. Stevens, C. M. Cardeno, M. W. Crepeau, P.

Duchen, J. J. Emerson, P. Saelao, D. J. Begun and C. H. Langley (2012). "Population Genomics

of sub-saharan Drosophila melanogaster: African diversity and non-African admixture." <u>PLoS</u>

<u>Genet</u> **8**(12): e1003080.

Rajpurohit, S., C. C. Oliveira, W. J. Etges and A. G. Gibbs (2013). "Functional genomic and phenotypic responses to desiccation in natural populations of a desert drosophilid." Mol Ecol **22**(10): 2698-2715.

Rajpurohit, S., X. Zhao and P. S. Schmidt (2017). "A resource on latitudinal and altitudinal clines of ecologically relevant phenotypes of the Indian Drosophila." Scientific Data **4**(1): 170066.

Reinhardt, J. A., B. Kolaczkowski, C. D. Jones, D. J. Begun and A. D. Kern (2014). "Parallel Geographic Variation in Drosophila melanogaster." Genetics **197**(1): 361-373.

Root, T. L., J. T. Price, K. R. Hall, S. H. Schneider, C. Rosenzweig and J. A. Pounds (2003). "Fingerprints of global warming on wild animals and plants." Nature **421**(6918): 57-60.

Rudman, S. M., S. I. Greenblum, S. Rajpurohit, N. J. Betancourt, J. Hanna, S. Tilk, T. Yokoyama, D. A. Petrov and P. Schmidt (2022). "Direct observation of adaptive tracking on ecological time scales in *Drosophila*." Science **375**(6586): eabj7484.

Rudman, S. M., S. I. Greenblum, S. Rajpurohit, N. J. Betancourt, J. Hanna, S. Tilk, T. Yokoyama, D. A. Petrov and P. Schmidt (2022). "Direct observation of adaptive tracking on ecological time scales in Drosophila." Science **375**(6586): eabj7484.

Schmidt, P. S., L. Matzkin, M. Ippolito and W. F. Eanes (2005). "GEOGRAPHIC VARIATION IN DIAPAUSE INCIDENCE, LIFE-HISTORY TRAITS, AND CLIMATIC ADAPTATION IN DROSOPHILA MELANOGASTER." Evolution **59**(8): 1721-1732.

Schmidt, P. S., C.-T. Zhu, J. Das, M. Batavia, L. Yang and W. F. Eanes (2008). "An amino acid polymorphism in the *couch potato* gene forms the basis for climatic adaptation in *Drosophila melanogaster*." Proceedings of the National Academy of Sciences **105**(42): 16207-16211.

Seebacher, F., C. R. White and C. E. Franklin (2015). "Physiological plasticity increases

resilience of ectothermic animals to climate change." <u>Nature Climate Change</u> **5**(1): 61-66.

Sejerkilde, M., J. G. Sørensen and V. Loeschcke (2003). "Effects of cold- and heat hardening on

thermal resistance in Drosophila melanogaster." <u>J Insect Physiol</u> **49**(8): 719-726.

Sezgin, E., D. D. Duvernell, L. M. Matzkin, Y. Duan, C. T. Zhu, B. C. Verrelli and W. F. Eanes

(2004). "Single-locus latitudinal clines and their relationship to temperate adaptation in

metabolic genes and derived alleles in Drosophila melanogaster." <u>Genetics</u> **168**(2): 923-931.

Sgrò, C. M., J. S. Terblanche and A. A. Hoffmann (2016). "What Can Plasticity Contribute to

Insect Responses to Climate Change?" <u>Annual Review of Entomology</u> **61**(1): 433-451.

Sgrò, C. M., J. S. Terblanche and A. A. Hoffmann (2016). "What Can Plasticity Contribute to

Insect Responses to Climate Change?" <u>Annu Rev Entomol</u> **61**: 433-451.

Sheldon, K. S. (2019). "Climate Change in the Tropics: Ecological and Evolutionary Responses

at Low Latitudes." <u>Annual Review of Ecology, Evolution, and Systematics</u> **50**(1): 303-333.

Sinclair, B. J., K. E. Marshall, M. A. Sewell, D. L. Levesque, C. S. Willett, S. Slotsbo, Y. Dong,

C. D. Harley, D. J. Marshall and B. S. Helmuth (2016). "Can we predict ectotherm responses to

climate change using thermal performance curves and body temperatures?" <u>Ecology letters</u>

19(11): 1372-1385.

Singh, A. K., L. L. McGoldrick, L. Demirkhanyan, M. Leslie, E. Zakharian and A. I. Sobolevsky

(2019). "Structural basis of temperature sensation by the TRP channel TRPV3." <u>Nat Struct Mol</u>

<u>Biol</u> **26**(11): 994-998.

Sørensen, J. G., T. N. Kristensen and J. Overgaard (2016). "Evolutionary and ecological patterns

of thermal acclimation capacity in Drosophila: is it important for keeping up with climate

change?" <u>Current Opinion in Insect Science</u> **17**: 98-104.

Sørensen, J. G. and V. Loeschcke (2007). "Studying stress responses in the post-genomic era: its

ecological and evolutionary role." J Biosci **32**(3): 447-456.

Sprengelmeyer, Q. D., S. Mansourian, J. D. Lange, D. R. Matute, B. S. Cooper, E. V. Jirle, M. C.

Stensmyr and J. E. Pool (2019). "Recurrent Collection of Drosophila melanogaster from Wild

African Environments and Genomic Insights into Species History." Molecular Biology and

Evolution **37**(3): 627-638.

Sprengelmeyer, Q. D., S. Mansourian, J. D. Lange, D. R. Matute, B. S. Cooper, E. V. Jirle, M. C.

Stensmyr and J. E. Pool (2020). "Recurrent Collection of Drosophila melanogaster from Wild

African Environments and Genomic Insights into Species History." Mol Biol Evol **37**(3): 627-

638.

Stevens, G. C. (1989). "The latitudinal gradient in geographical range: how so many species

coexist in the tropics." The American Naturalist **133**(2): 240-256.

Stone, H. M., P. A. Erickson and A. O. Bergland (2020). "Phenotypic plasticity, but not adaptive

tracking, underlies seasonal variation in post-cold hardening freeze tolerance of Drosophila

melanogaster." Ecol Evol **10**(1): 217-231.

Sunday, J. M., A. E. Bates, M. R. Kearney, R. K. Colwell, N. K. Dulvy, J. T. Longino and R. B.

Huey (2014). "Thermal-safety margins and the necessity of thermoregulatory behavior across

latitude and elevation." Proceedings of the National Academy of Sciences **111**(15): 5610-5615.

Svetec, N., L. Zhao, P. Saelao, J. C. Chiu and D. J. Begun (2015). "Evidence that natural

selection maintains genetic variation for sleep in Drosophila melanogaster." BMC Evolutionary

Biology **15**(1): 41.

Tobler, R., J. Hermisson and C. Schlötterer (2015). "Parallel trait adaptation across opposing

thermal environments in experimental Drosophila melanogaster populations." Evolution **69**(7):

1745-1759.

Turner, T. L., M. T. Levine, M. L. Eckert and D. J. Begun (2008). "Genomic analysis of adaptive

differentiation in Drosophila melanogaster." Genetics **179**(1): 455-473.

Turney, C., A.-G. Ausseil and L. Broadhurst (2020). "Urgent need for an integrated policy

framework for biodiversity loss and climate change." Nature Ecology & Evolution **4**(8): 996-

996.

Venkatachalam, K. and C. Montell (2007). "TRP channels." Annu. Rev. Biochem. **76**: 387-417.

Wang, G. and M. E. Dillon (2014). "Recent geographic convergence in diurnal and annual

temperature cycling flattens global thermal profiles." Nature Climate Change **4**(11): 988-992.

Weeks, A. R., S. W. McKechnie and A. A. Hoffmann (2002). "Dissecting adaptive clinal

variation: markers, inversions and size/stress associations in Drosophila melanogaster from a

central field population." Ecology Letters **5**(6): 756-763.

Winbush, A. and N. D. Singh (2021). "Genomics of Recombination Rate Variation in

Temperature-Evolved Drosophila melanogaster Populations." Genome Biol Evol **13**(1).

Zhao, X., A. O. Bergland, E. L. Behrman, B. D. Gregory, D. A. Petrov and P. S. Schmidt (2016).

"Global Transcriptional Profiling of Diapause and Climatic Adaptation in Drosophila

melanogaster." Mol Biol Evol **33**(3): 707-720.