Circadian and environmental signal integration in a natural population of *Arabidopsis*

Haruki Nishio[1,2,7], Dora L. Cano-Ramirez[3,4,7], Tomoaki Muranaka[1,5], Luíza Lane de Barros

Dantas[6], Mie N. Honjo[1], Jiro Sugisaka[1], Hiroshi Kudoh[1], Antony N. Dodd[6*].

1. Center for Ecological Research, Kyoto University, Otsu, Shiga, Japan.

2. Data Science and AI Innovation Research Promotion Center, Shiga University, Hikone,

Shiga, Japan.

3. The Sainsbury Laboratory, University of Cambridge, Cambridge, UK.

4. School of Biological Sciences, University of Bristol, Bristol, UK.

5. Graduate School of Bioagricultural Sciences, Nagoya University, Nagoya, Aichi, Japan.

6. Department of Cell and Developmental Biology, John Innes Centre, Norwich, UK.

7. Equal contribution to this research.

* Corresponding author; antony.dodd@jic.ac.uk

Word counts: Main text (inc. methods) 4883; Abstract 178.

Abstract

Plants sense and respond to environmental cues during 24 h fluctuations in their

environment. This requires the integration of internal cues such as circadian timing with

environmental cues such as light and temperature to elicit cellular responses through signal

transduction. However, the integration and transduction of circadian and environmental

signals within plants growing in natural environments remains poorly understood. To gain

insights into the 24 h dynamics of environmental signalling in nature, we performed a field

study of signalling from the nucleus to chloroplasts in a natural population of *Arabidopsis*

halleri. Using advanced modelling approaches to interpret the data, we identified that the

circadian clock and temperature are key regulators of this pathway under natural conditions.

We identified potential time-delay steps between pathway components, and diel fluctuations

in the response of the pathway to temperature cues that are reminiscent of the process of

circadian gating. This approach of combining studies of gene expression in the field with

modelling allowed us to identify the dynamic integration and transduction of environmental

cues, in plant cells, under naturally fluctuating diel cycles.

Introduction

Plants have sophisticated environmental sensing and signalling mechanisms that underpin their responses to the fluctuating environment. Under natural conditions, this requires signalling pathways that integrate dynamic, overlapping and complex environmental stimuli [1, 2]. These environmental fluctuations include an oscillating component with a 24 h period, which arises from the cycle of day and night. Circadian clocks, which are endogenous biological oscillators, provide a cellular estimate of the time of day that coordinates the diel responses of plants with environmental fluctuations by aligning transcription, metabolism and development with the time of day [3-8]. In plants, environmental information including the light and temperature conditions is used to adjust the phase of the circadian oscillator, through the process of entrainment, so that the phase is aligned with the 24 h environmental cycle. This alignment between the circadian oscillator and the 24 h environmental cycle contributes to the fitness of plants [5].

Under natural conditions, environmental cues are a major determinant of temporal programs of gene expression [9]. For example, 97% of diel transcript profiles in field-grown rice can be predicted from meteorological data [9], and temperature cues regulate the alternative splicing of transcripts encoding circadian oscillator components in field-grown sugarcane [10]. Recent studies have provided insights into the diel organization of the transcriptome and metabolism, under field conditions, for several crops and *Arabidopsis* species [9-18]. A question that arises from these studies is how information from the circadian clock and environmental cues are processed to form integrated outputs. The diel dynamics of environmental signalling pathways, with defined inputs and outputs, are less well understood under natural conditions. The diel regulation of processes under controlled laboratory conditions does not necessarily reflect the dynamics in the field, even when using sophisticated replication of field conditions [15, 17]. Therefore, analysing the dynamics of cellular processes within complex fluctuating natural conditions allows us to elucidate how molecular mechanisms identified under controlled laboratory conditions operate in a natural

setting [15]. Understanding this is valuable for translating laboratory studies into crop

improvement. Furthermore, considering that plants comprise the major part of organismal

biomass on Earth, it could contribute to forecasting ecosystem responses to increasingly

unpredictable climates [19, 20].

To study the integration and transduction of circadian and environmental signals under

naturally fluctuating conditions, we selected a well-characterized environmental signalling

pathway as a model system. This model comprises the regulation of three components; a

nuclear-encoded key component of the *Arabidopsis* circadian clock (*CIRCADIAN CLOCK*

ASSOCIATED 1, *CCA1*), a nuclear-encoded regulator of chloroplast transcription (*SIGMA*

FACTOR 5, *SIG5*), and the chloroplast-encoded gene *psbD*, which encodes the D2 protein

of Photosystem II [21-23] (Fig. 1A). *CCA1* transcript abundance can be used as a proxy for

the status of the circadian clock. *SIG5* is regulated closely by the circadian oscillator under

constant conditions [22], and its transcript abundance responds to light, temperature and

other environmental cues [21-33], which made us reason that SIG5 is an integrator of

circadian and environmental information. SIG5 is imported into chloroplasts, and regulates

transcription from the blue light responsive promoter of *psbD* (*psbD* BLRP) [21] (Fig. 1A).

Thus, we assumed that SIG5-mediated signalling to chloroplasts involves a hierarchically-

organized pathway, whereby *AhgCCA1* is positioned upstream from the regulation of

AhgSIG5 transcript accumulation, and *AhgpsbD* BLRP is positioned downstream of

AhgSIG5 activity (Fig. 1A). We also assumed that environmental signals might influence

each component independently (Fig. 1A) [34]. We chose this pathway because it includes

three major components representing distinct regulatory points of signal transduction, and

has a relatively low level of complexity to evaluate circadian and environmental signal

integration and transduction under realistic field conditions.

By targeting this model pathway, we combined time-series gene expression data with

statistical modelling to investigate the temporal dynamics of signal integration and

transduction in a fluctuating natural environment. We conducted the study using a natural

population of a perennial *Arabidopsis* species (*Arabidopsis halleri* subsp. *gemmifera*,

referred to as *A. halleri*) (Fig. 1B-D) [35]. The close-relatedness of *A. halleri* and *A. thaliana*

makes it possible to identify pairs of homologous genes based on the sequence similarity

(indicated by *Ahg* or *At* prefixes to gene names) [36]. The circadian clock-SIG5-*psbD* BLRP

pathway is present in *A. thaliana* and *A. halleri*, and well-conserved across the vascular

plants [37, 38]. We obtained several time series, during two seasons of the year, that

monitored pathway function under representative light and temperature conditions in *A.*

halleri in its natural habitat.

Interpreting time-series transcript data derived from complex fluctuating environments is a

challenging task. We used statistical models, rather than models of biochemical kinetics [39],

because this provides an effective tool for interpreting the influences of external factors on

the response variables. We applied state-space modelling that integrates observations and

the latent true state of a dynamic system into a statistical model. It is often used to interpret

trends and oscillations in a system, and to evaluate the effects of external drivers on time-

series data [40-42]. It explicitly formulates the system noise and observation error as state

and observation equations, respectively. By incorporating an autoregressive trend and

exogenous influences into a state space model, we estimated the dynamics of each pathway

component that arises from its internal dynamics and external factors. Comparable

approaches have allowed the investigation of diel and seasonal changes of transcriptome

dynamics in *A. halleri* [12, 43, 44] and rice [9].

We evaluated evidence for causal regulation between the pathway components using

convergent cross mapping (CCM), which provides a quantitative test for causation in time-

series of coupled dynamical systems [45]. In contrast to our predictive state-space models,

CCM evaluates whether the historical fluctuation in one variable can be used to estimate the

fluctuation in a second variable [45]. This has been used previously to investigate a variety

of ecological processes, including the seasonal regulation of gene expression by epigenetic

marks in a natural *Arabidopsis* population [44] and the seasonal transcriptome in Japanese

beech [46]. Using these approaches, we identified key roles for temperature and the

circadian clock in the regulation of this signalling pathway under natural conditions. To our

knowledge, this is the first investigation of circadian signal transduction in plants in their

natural habitat, and the first use of these analytical approaches to understand circadian

clock-regulated pathway in the field. Our approaches could be applicable to the study of

many circadian-regulated processes under naturally fluctuating conditions.

Results

Biological data underlying models of signal transduction

Under controlled conditions of constant light, *AtCCA1* and *AtSIG5* transcript abundance are

very well correlated (Fig. S1A-D; data from [3, 4, 6, 22, 47]). This correlation between

AtCCA1 and *AtSIG5* transcript abundance is absent under light/dark cycles (Fig. S1E, F;

data from [22, 48]), suggesting that the integration of light and dark cues alters the diel

regulation of *AtSIG5* transcript accumulation. To investigate these processes of signal

integration under natural conditions, we acquired time-series of transcript abundance during

spring (March) and autumn/fall (September), close to the spring or autumn equinox (Fig. S2;

Fig. S3). Although both the spring and autumn equinoxes share 12 h photoperiods, they

provide contrasting temperature regimes (cool and warm, respectively) (Fig. 2A, B; Fig. S4A,

B), with irradiance levels determined by weather conditions (Fig. 2C, D, Fig. S4C, D). This

allowed us to investigate temperature, light and seasonal influences upon SIG5-mediated

signalling to chloroplasts, because this pathway is known to be affected by light and

temperature in *A. thaliana* [21, 23, 24, 28, 33, 37]. We obtained data from areas with open

sky and with vegetational shade, to include within our models the transcriptional responses

to a wider range of irradiance levels (Fig. 2C, D; Fig. S4C, D). The "sun" and "shade"

sampling sites were chosen by measurement of the ratio of red to far-red light (R:FR) (Fig.

S5) and availability of plant patches, because *A. halleri* does not grow in deep shade. The

total light intensity at the sun sampling site was 5 to 10-fold greater during March 2015 than

during September 2015, depending on the time of day, due to weather differences (Fig. 2C,

D; Fig. S4C, D). During March 2015, the study site temperature ranged from 0 °C to 14 °C at

the sun site, and from 0 °C to 13 °C at the shade site (Fig. 2A, B; Fig. S4A). The

temperature was always above 17 °C during September 2015, with greater diel fluctuations

at the sun site (Fig. 2A, B; Fig. S4B).

We monitored a 24 h cycle of abundance of *AhgCCA1*, *AhgSIG5* and *AhgpsbD* BLRP

transcripts under each of these conditions, using RT-qPCR (Fig. S2; Fig. S3). Data from all

the experiments were normalized to a single reference sample collected at midday, to

enable comparability between sampling seasons and conditions. *AhgCCA1* oscillated with a

24 h cycle, reaching peak abundance at or just after astronomical dawn (Fig. S2E, F),

whereas *AhgSIG5* and *AhgpsbD* BLRP reached peak abundance between the middle and

end of the photoperiod, depending on conditions (Fig. S2G-J). The magnitude of oscillation

of each transcript varied according to the sampling season and light conditions. *AhgCCA1*

transcripts underwent a 5- (sun) and 7-fold (shade) change over the 24 h cycle during the

March sampling, compared with a 98-fold (sun) and 190-fold (shade) change during the

September sampling (fold-changes based on peak and trough abundance). This greater

magnitude of oscillation occurred also for *AhgSIG5* during September (sun 38-fold; shade

33-fold) compared with March (sun 2.7-fold, shade 5.0-fold), and also for *AhgpsbD* BLRP

during September (4.3-fold in sun and shade conditions) compared with March (sun 1.7-fold,

shade 1.3-fold). Therefore, the relative magnitude of *AhgCCA1* transcript oscillation aligned

well with that of *AhgSIG5* and *AhgpsbD* BLRP, whereby stronger diel cycling was present in

September than in March, as suggested by the greater amplitude of *AhgCCA1* transcript

during September sampling period compared with March. Furthermore, the magnitude (fold-

change) of the 24 h oscillation of *AhgCCA1* was always greater than for *AhgSIG5,* and even

less for *AhgpsbD* BLRP under any given condition. Whilst *AhgCCA1* underwent a greater

daily fold-change under shade conditions compared with sun conditions, this was not

reflected equivalently for *AhgSIG5* and *AhgpsbD* BLRP.

We compared the pathway dynamics between the spring and autumn sampling periods by

using a smooth trend model (referred to here as STM). This is a type of state-space

modelling that allows inference of a trend over time, where there is sampling noise and

environmental stochasticity [49]. We estimated the parameters of a smooth trend model by

Bayesian inference using the Markov Chain Monte Carlo (MCMC) approach [42, 50] to

visualize the differences in transcript abundance between the spring and autumn sampling

periods. Using this approach, we determined that the greater morning peaks of *AhgCCA1*

expression in autumn compared to those in spring was statistically significant under both

light conditions tested (the shaded areas of the lower panels of Fig. 2E, F represent the 95%

confidence interval of the difference between seasons not containing zero around dawn).

During both sampling seasons, *AhgSIG5* transcripts reached peak abundance between the

middle and end of the photoperiod (Fig. 2G, H). This differs from *AtSIG5* transcript

accumulation in *A. thaliana* under square-wave light/dark cycles under controlled conditions,

where *AtSIG5* transcript abundance peaks around dawn [22]. The peak of the diel fluctuation

of *AhgSIG5* was significantly greater during the March sampling period than during the

September sampling period (95% confidence interval; Fig. 2G, H). Transcripts encoding the

SIG5 regulatory target *AhgpsbD* BLRP (Fig. 1A) had a diel fluctuation during the September

sampling season that reached a much greater peak under shade conditions compared with

sun conditions (Fig. 2I, J), whereas there was very little diel fluctuation of *AhgpsbD* BLRP

during the March sampling season.

AhgCCA1 transcript abundance was significantly greater under shade compared with sun

conditions during the photoperiod, during both sampling seasons (95% confidence interval;

Fig. S4E, F). We did not identify the diminished *AtCCA1* oscillation that occurs under

constant light with a very low R:FR [51] or on the shaded western side of crop fields around

dawn [16]. In comparison, *AhgSIG5* transcript abundance was not altered consistently

between sun and shade conditions (95% confidence interval; Fig. S4G, H). This tendency

was also not observed for *AhgpsbD* BLRP transcript levels (95% confidence interval; Fig.

S4I, J).

Dynamics of environmental regulation of the signalling pathway

We represented the behaviour of the pathway components using a local level model with

exogenous variables (referred to here as LLMX), a type of state-space modelling [52]. This

allows the inclusion of a circadian or diel trend within a regression model that predicts

transcript levels from environmental variables and the upstream regulatory component (Fig.

1A). The output of the Bayesian estimation reproduced well the dynamics of the observed

AhgSIG5 transcript level (Fig. 3A, B), with convergence of MCMC sampling (Fig. S6). The

model estimated a clear diel trend of *AhgSIG5* transcript level that reached the trough level

at dawn and the peak level between the middle and end of the photoperiod (Fig. 3C). The

model also estimated significant negative and positive effects of ambient temperature and

AhgCCA1 on *AhgSIG5* transcript abundance, respectively (the error bars representing the

95% confidence interval of regression coefficients not containing zero; Fig. 3D). There was

no significant effect of irradiance upon the estimation of *AhgSIG5* transcript abundance (Fig.

3D). Under controlled laboratory conditions, a temperature reduction leads to an increase in

SIG5 transcript abundance [33]. Therefore, the estimated negative relationship between

ambient temperature and *AhgSIG5* transcript levels (Fig. 3D) could suggest a similar

regulation in *A. halleri* under naturally fluctuating conditions. If the temperature input to the

model was set to a constant value (the mean temperature across all conditions) and the

estimated parameter values in this model were used, the greater *AhgSIG5* transcript

abundance during March disappeared, whereas the phase of the dynamics was not affected

(compare Fig. S7A, B and Fig. S7C ,D). This suggests that temperature is a key predictor of

the difference in *AhgSIG5* transcript abundance between seasons. Setting the irradiance

input to a constant level (the mean irradiance across all conditions) did not affect the

AhgSIG5 dynamics (compare Fig. S7A, B and Fig. S7E, F). Setting a constant *AhgCCA1*

transcript level shifted the phase of *AhgSIG5* dynamics in September (compare Fig. S7A, B

and Fig. S7G, H), suggesting that *AhgCCA1* input partially contributes to the prediction of

the *AhgSIG5* oscillation.

Diel fluctuations of *AhgpsbD* BLRP transcript abundance were reproduced reasonably well

by the model (Fig. 3E, F), with convergence of MCMC sampling (Fig. S8). The model

predicted a diel trend of *AhgpsbD* BLRP transcript level that increased towards the end of

the photoperiod (Fig. 3G). The model estimated a significant positive effect of ambient

temperature on *AhgpsbD* BLRP transcript abundance (95% confidence interval; Fig. 3H). In

A. thaliana under controlled conditions, *psbD* BLRP transcript accumulation is regulated by

SIG5 [21], and within our model, there was also a significant positive coefficient of

regression between *AhgSIG5* and *AhgpsbD* BLRP transcript levels (95% confidence interval;

Fig. 3H). Setting the temperature input as a constant value (the mean temperature among all

conditions) with estimated parameter values in this model, greater *AhgpsbD* BLRP transcript

abundance in September disappeared, and the diel dynamics became slightly flatter

(compare Fig. S9A, B and Fig. S9C, D). This suggests that temperature is a key predictor of

the difference in *AhgpsbD* BLRP transcript abundance between seasons. Constant

irradiance did not affect the *AhgpsbD* BLRP dynamics (compare Fig. S9A, B and Fig. S9E,

F). Constant *AhgSIG5* transcript level caused apparent flatter *AhgpsbD* BLRP dynamics

(compare Fig. S9A, B and Fig. S9G, H), suggesting that *AhgSIG5* input makes a partial

prediction of the *AhgpsbD* BLRP oscillation.

Overall, this analysis identifies that the ambient temperature, rather than the irradiance, was

important for predicting the seasonal differences in the transcript abundance of *AhgSIG5*

and *AhgpsbD* BLRP under naturally fluctuating conditions. This contrasts evidence from *A.*

thaliana under controlled conditions, where light inputs have a key role [21, 23]. In addition,

the circadian clock (*AhgCCA1*) affected the diel oscillation of *AhgSIG5* transcript abundance

(Fig. 3D; Fig. S8G, H), and *AhgSIG5* affected the diel oscillation of *AhgpsbD* BLRP transcript

abundance (Fig. 3H; Fig. S9G, H). More specifically, seasonal difference in temperature

affected the base levels of *AhgSIG5* and *AhgpsbD* BLRP transcripts, while the diel

oscillation patterns of *AhgSIG5* and *AhgpsbD* BLRP transcripts were affected by the

transcript levels of their upstream genes rather than diel temperature fluctuations.

The significant effects of temperature on *AhgSIG5* and *AhgpsbD* BLRP transcripts (Fig. 3D,

H), and the restriction to specific periods of the day of significant differences between the

transcript dynamics under the different temperature conditions of March and September (Fig.

2E-J), is reminiscent of the concept of circadian gating. Circadian gating is the process

whereby the circadian clock restricts certain biological processes to specific times in the 24 h

cycle [53]. In plants, this can take the form of a circadian rhythm in the magnitude of the

response to identical environmental stimuli given at different times of day [8].

Temporal gating of temperature regulation of SIG5-mediated signalling to

chloroplasts under natural conditions

Our LLMX analysis suggests that under natural conditions, lower ambient temperatures

might upregulate *AhgSIG5* transcript levels, and greater ambient temperatures might

upregulate *AhgpsbD* BLRP transcript levels (Fig. 3D, H). To test this hypothesis and further

examine whether a circadian gating-like process might occur, we applied moderate

temperature manipulations to adjacent patches of *A. halleri* plants, in the field, using custom-

designed equipment (Fig. 4A; Fig. S10A-D). We collected 24-h time-series of RNA samples

from these plant patches (Fig. S10E-G), and interpreted the data with smooth trend models

(STM; Fig. 4B-D). This experiment occurred during September 2016, to take advantage of

the greater magnitude of oscillation of the transcripts observed during experimentation

during September 2015 compared with March 2015 (Fig. 2E-J). *AhgCCA1* transcripts

underwent a substantial fold-change fluctuation over the 24 h cycle, ranging from 210-fold

(moderate temperature reduction) to 339-fold (ambient temperature) and 1,230-fold

(moderate temperature increase) (Fig. S10E). Interestingly, a moderate temperature

reduction treatment during September 2016 did not decrease the fluctuation of *AhgCCA1*

transcripts to a level equivalent to the March 2015 sampling season (5 to 7-fold fluctuation),

suggesting that either there is a tipping point temperature below which *A. halleri CCA1*

transcripts have a substantial loss of rhythmicity, or that the past temperature history of the

plants during each season can influence the amplitude of oscillation of this circadian clock

transcript.

Both the moderate temperature increase and temperature reduction treatment caused a

small significant upregulation of *AhgCCA1* transcript abundance around dawn relative to the

ambient temperature control (significance at > 95% probability; Fig. 4B). The moderate

temperature increase treatment was without effect upon *AhgSIG5* transcript abundance (Fig.

4C). In comparison, the temperature reduction treatment upregulated *AhgSIG5* transcripts

significantly after dawn and around dusk, relative to the ambient temperature control

(significance at > 95% probability; Fig. 4C). This is consistent with the negative coefficient of

regression of temperature on *AhgSIG5* transcript under naturally fluctuating conditions (Fig.

3D), and with the upregulation of *A. thaliana SIG5* by a short cold treatment under laboratory

conditions [21, 33]. The restriction of the response of *AhgSIG5* transcripts to the moderate

temperature reduction to certain times of day (Fig. 4C) supports the notion that there is

temporal gating of the response of *AhgSIG5* to temperature under naturally fluctuating

conditions.

Transcripts for the chloroplast target of SIG5, *AhgpsbD* BLRP, were also altered by

temperature manipulation (Fig. 4D). The moderate temperature elevation significantly

increased *AhgpsbD* BLRP transcript levels relative to the control, whereas the moderate

temperature reduction significantly reduced *AhgpsbD* BLRP transcripts relative to the control

(significance at > 95% probability; Fig. 4D). This is consistent with the positive coefficient of

regression of temperature on *AhgpsbD* BLRP transcript abundance under naturally

fluctuating conditions (Fig. 3H). The response of *AhgpsbD* BLRP transcripts to temperature

manipulations was restricted to the photoperiod, which might be because chloroplast DNA

binding and transcription by PEP generally requires light [54-58].

Evaluation of causal regulation within a circadian-regulated signalling pathway under

natural conditions

We performed CCM analysis of the relationship between *AhgCCA1* and *AhgSIG5*, and of the

relationship between *AhgSIG5* and *AhgpsbD* BLRP, using a variety of time-delays between

each pair of variables. We considered potential time-delays in CCM analysis [59] because

the abundance of each related transcript was monitored at each timepoint, but their

responses to each other might not be instantaneous. For example, in *A. thaliana* under

controlled square-wave light/dark cycle conditions, *AtCCA1* transcript abundance peaks at

dawn, *AtSIG5* approximately 3 h after dawn, and *AtpsbD* BLRP approximately 6 hours after

dawn [22].

To prepare for CCM analysis, we determined the optimal embedding dimension for each

variable (*AhgCCA1*, *AhgSIG5* and *AhgpsbD* BLRP; Fig. S11). A feature of CCM analysis

that can confuse readers is that the direction of successful prediction is *opposite* to that of

causality (Fig. 5A). Subsequent analysis with CCM identified a significant prediction (cross

map skill) of *AhgCCA1* transcript abundance from *AhgSIG5* transcript abundance (Fig. 5B)

when no time delay was incorporated between the variables. Furthermore, a significant

prediction of *AhgSIG5* from *AhgpsbD* BLRP transcript abundance occurred when no time

delay was incorporated, and potentially also with a time delay between *AhgSIG5* and

AhgpsbD BLRP (Fig. 5C). In both cases, the cross map skill exceeded a 95% interval of the

diel surrogate, which imitates the same degree of oscillation of the explanatory variable, but

where the variation is randomized (Fig. 5B, C). When the library size (number of time points

used to reconstruct a state space) was increased, cross map skill was improved for each

time lag, attaining a significant cross map skill (Fig. 5D-F), suggesting convergence (a key

property that distinguishes causation from simple correlation) [45]. Overall, CCM analysis

suggests that there is a causal relationship between fluctuations in *AhgCCA1* and *AhgSIG5*,

and *AhgSIG5* and *AhgpsbD* BLRP, and that there is a potential time-delay from nuclear

genome-encoding *AhgSIG5* to chloroplast genome-encoding *AhgpsbD* BLRP in a natural

plant population.

Discussion

We deployed a set of approaches that allowed us to gain insights into the integration and

transduction of circadian and environmental signals in a natural plant population. Using a

modelling approach, we predicted that the circadian clock and ambient temperature are

potential regulators of *AhgSIG5*-mediated signalling to chloroplasts, in a natural population

of *A. halleri*. There was a significant influence of *AhgCCA1* on *AhgSIG5* transcript

abundance, and of *AhgSIG5* on *AhgpsbD* BLRP transcript abundance (Fig. 3D, H).

Furthermore, convergent cross mapping analysis predicted significant causal relationships

within this signalling pathway in the field (Fig. 5). Therefore, it appears that in a natural plant

population, a signal is communicated from the circadian oscillator (using *AhgCCA1* as a

proxy) to the signalling pathway output of *AhgpsbD* BLRP. One interpretation is that the

pathway couples the circadian oscillator and temperature response processes to chloroplast

gene transcription under naturally fluctuating conditions.

We identified seasonal differences in the maximum accumulation of *AhgCCA1*, *AhgSIG5*

and *AhgpsbD* BLRP. *AhgCCA1* and *AhgpsbD* BLRP had significantly lower peak

accumulation during the spring sampling period compared with the autumn sampling season

(Fig. 2E, F, I, J). In comparison, *AhgSIG5* had significantly greater peak accumulation during

the spring compared with the autumn sampling season (Fig. 2G, H). The difference in

AhgCCA1 dynamics between these sampling seasons likely reflects the decreased

amplitude of the circadian oscillator that occurs under lower temperature conditions, in both

controlled laboratory environments and the field [12, 60-62]. The difference in the seasonal

response of *AhgSIG5* compared with *AhgCCA1* suggests that an additional temperature

input into this pathway occurs between the circadian oscillator and *AhgSIG5*. In *A. thaliana*,

AtSIG5 transcripts are upregulated by short cold temperature treatments [21, 33]. In our field

experiment, *AhgSIG5* transcript accumulation showed a negative coefficient of regression

with temperature (Fig. 3D), with its accumulation increased during the morning and around

dusk by a low temperature manipulation (Fig. 4C). This suggests that under lower

temperature conditions, *AhgSIG5* transcript abundance will increase. Therefore, the lower

temperatures of the spring sampling season compared with the autumn sampling season

(Fig. 2A, B) might explain the significantly greater abundance of *AhgSIG5* transcripts at

many times of day during the spring. A further feature of interest within the data is the

relative robustness of the dawn timing of peak *AhgCCA1* abundance, irrespective of the

environmental conditions (Fig. 2E, F), which is consistent with a previous field study [12]. In

Arabidopsis, *AtCCA1* is light-induced [63, 64], yet under field conditions *AhgCCA1* reached a

peak around astronomical dawn when the irradiance was relatively low, and was

unresponsive to increased light intensity later in the day (Fig. 2C-F). Furthermore, its peak

time was relatively unaffected by seasonal differences in light conditions (Fig. 2C-F). This

reveals an astonishing consistency of the phase of expression of a clock gene in plants

under complex fluctuating environments, which might be accomplished through the

complexity of the *Arabidopsis* circadian oscillator that is thought to confer robustness to its

regulation [65, 66], and perhaps a restriction of light-induced phase shifts to certain times of

day due to the gating of light inputs. Mathematical modelling indicates that the interplay

between different simultaneous circadian entrainment cues might reinforce circadian clocks

[67], although the timing of individual cues applied in combination can also cause changes in

the phase [68]. The processes that align the phase of the plant circadian clock in plants with

growing environments, under naturally fluctuating conditions, represent a fascinating topic

for future investigation.

Because we considered *AhgpsbD* BLRP to represent the ultimate output from the signalling

pathway (Fig. 1A), our analysis suggests that environmental inputs occurred within at least

three positions in the pathway; first, in the regulation of *AhgCCA1* transcript accumulation by

the season or temperature, second, in the regulation of *AhgSIG5* transcript accumulation by

temperature, and third, in the regulation of *AhgpsbD* BLRP transcript accumulation by

temperature. These environmental inputs might occur through several biologically

independent processes. One possibility is temperature inputs to the circadian clock mediated

by temperature-responsive components, such as the evening complex [69, 70]. A further

mechanism could be the regulation of *AhgSIG5* by HY5, which is a known regulator of SIG5

that participates in low-temperature gene regulation [21, 27, 71] and underlies a response of

AtSIG5 transcripts to cold in *A. thaliana* [33]. Furthermore, there might be direct effects of

light upon sigma factor activity in chloroplasts through, for example, redox regulation [72] or

light- and temperature-regulation of chloroplast protein import [73]. We did not consider here

the short- or long-term history of light or temperature conditions upon leaves prior to

experimentation. However, the past temperature conditions can affect *AhgCCA1* oscillations,

since the sensitivity of *AhgCCA1* oscillations to a moderate temperature reduction treatment

during September 2016 was limited compared with its response to the lower ambient

temperature conditions during Spring 2015. Other biotic or abiotic factors such as water

availability, relative humidity, and atmospheric CO_2 concentration might also have effects

that we did not include in our models.

Statistical modelling of the transcriptome of field-grown *Oryza sativa* (rice) concluded that

the main environmental driver of *OsSIG5* (*Os05g0586600*) transcript accumulation is

temperature [9]. In this case, the temperature had a negative regression coefficient with

OsSIG5 [9]. This is consistent with our finding of a negative coefficient of regression of

temperature on *AhgSIG5*. The study of the rice transcriptome in the field [9] did not monitor

chloroplast-encoded transcripts, so a direct comparison between *AhgpsbD* BLRP and our

data is not possible.

A key finding from our work is the detection of a 24 h fluctuation of the influence of the

temperature upon each of the three genes (Fig. 4). One interpretation is that there is a diel

cycle of sensitivity of these pathway components to temperature cues, with their response to

temperature restricted to certain times of day, reminiscent of the process of circadian gating

[53]. Our findings from the field are corroborated by laboratory experiments, which

demonstrate that the circadian oscillator gates its own response to temperature [74], and the

response of *AtSIG5* to blue light and cold temperatures is gated by the circadian oscillator

[22, 33]. Statistical modelling approaches applied to diel cycles of the rice transcriptome in

the field have identified 24 h oscillations in the coefficient that describes the relationship

between temperature and transcript levels, which is also reminiscent of circadian gating [9].

In field experiments, it is difficult to link the fluctuating temperature sensitivity to the circadian

oscillator causatively, although a combination of field and laboratory experiments

investigating the environmental regulation of *FLOWERING LOCUS T* (*FT*) expression

suggest that the circadian clock can gate light quality and temperature responses of *FT*

under field conditions [18, 75]. Taken together, these findings are important because they

suggest that processes leading to circadian gating might operate in natural plant populations

to modulate their environmental responses.

Our investigation provides insights into molecular aspects of signal transduction in plants

under field conditions. This represents a relatively under-studied topic, and we adopted

quantitative approaches to interpret relatively noisy transcript data collected under complex

fluctuating environments, to investigate the functioning of a specific pathway. This allowed

us to infer multiple positions of environmental inputs into the pathway, and temporal gating of

a response to temperature. The approaches used provide a framework to study

environmental signal integration in all organisms under field conditions, providing

opportunities to investigate whether and how mechanisms identified in the laboratory might

function in the field. This could be important to understand rhythmic biological responses

within an increasingly unpredictable climate.

Materials and Methods

Field site and plant material

Our experiments used a naturally-occurring population of *Arabidopsis halleri* subsp.

gemmifera (Matsum.) O'Kane & Al-Shehbaz growing beside a forested stream in Hyogo

Prefecture, Japan (Omoide-gawa site; 35°06' N, 134°55' E, elevation 190–230 m) [12, 43,

76] (Fig. 1B). We selected *A. halleri* as an experimental model for several reasons [35]. First,

it has high nucleotide sequence identity and good synteny with *A. thaliana* [36]. Second,

unlike *A. thaliana*, the perennial life history of *A. halleri* allows investigation of transcriptional

responses across the seasons [2, 76]. Many individuals are clones because the species

propagates by producing clonal rosettes as well as by seeds, which allows repeated

sampling from single genotypes. Furthermore, *A. halleri* is metal tolerant and occurs in

natural habitats that are relatively free from other vegetation due to contamination by heavy

metals, which provides experimentally-convenient sites enriched with many *A. halleri* plants

[76]. *Arabidopsis halleri* subsp. *gemmifera* at this site was previously identified by

examination of museum and herbarium specimens, and a nearby population provided

material for sequencing the *A. halleri* genome [36, 76].

AhgCCA1 (*g097040*) and *AhgSIG5* (*g25274*) were identified from *A. halleri* genome

Version Ahal2.2 [36]. *AhgCCA1* has 94.8% coding sequence identity and 93.3% protein

sequence identity with *AtCCA1*. *AhgSIG5* has 94.9% coding sequence identity and 95.0%

protein sequence identity with *AtSIG5*. Chloroplast-encoded *psbD* is not annotated within

Version Ahal2.2 of the *A. halleri* genome, and we identified this instead within scaffold 2 of

an *A. halleri* reference transcriptome [77]. The *AhgpsbD* BLRP promoter region, which was

our focus, has a 100% sequence identity with *AtpsbD* BLRP [55].

Experimental conditions for field sampling of A. halleri

The first sets of samples were obtained under natural conditions without temperature

manipulation during 24 – 26 March 2015 and 15 – 17 September 2015, which were close to

the spring and autumn equinox at the field site, respectively. We exploited variations in

environmental conditions across the field site, and sampled leaves from the locations

nominated as "sun" and "shade" sites on successive days. At "sun" locations, plants received

direct sunlight during the day, and at "shade" locations plants received sunlight filtered by

surrounding vegetation for most of the day with the sites identified by measurement of the

ratio of red to far red light (Fig. S5; R:FR calculated as the photon irradiance from 660 to

670nm divided by the photon irradiance from 725 to 735nm [78]). During March 2015, plants

received more direct sunlight, whereas during September 2015 the light was scattered

through sky overcast with clouds.

The second sets of samples were obtained under natural conditions with manipulation of the

temperature conditions around patches of plants during 13 – 14 September 2016, which was

close to the autumn equinox at the field site. In addition to control plants that were not

manipulated (Fig. S10A), we applied two temperature treatments. These were (1) a

continuous temperature increase (Fig. S10B), whereby plants were covered with clear

plastic horticultural domes to block air currents and trap warm air; (2) a continuous

temperature reduction, using a custom device that passed air through a duct within a heat-

exchanging ice-filled polystyrene box and expelled the chilled air into a clear horticultural

dome covering the plants, with chilling augmented by small ice packs within the dome (Fig.

S10C).

Field sampling for transcript analysis

Across all experimental conditions, the same sampling and RNA isolation procedures were

used. At 2 h intervals, a fully expanded rosette leaf was excised with dissecting scissors

from 6 replicate plants for each condition. The time-courses using naturally occurring sun

and shade conditions each comprised 13 sampling timepoints over a total of 26 hours (from

14:00 on the first day to 16:00 on the second day). The time-courses involving artificial

temperature manipulations comprised 15 sampling timepoints over a total of 30 hours (from

17:00 on the first day to 21:00 on the second day). The same replicate plants were sampled

repeatedly through each time-series, but different plant patches were used in different

sampling seasons. Sampled leaves were placed immediately into individual microtubes

containing at least 400 µL RNA*later* Stabilization Solution (Thermo Fisher Scientific,

Waltham, MA, USA). Scissors and forceps were cleaned with 70% (w/v) ethanol between

samples. After sampling, tubes were placed temporarily on dry ice for up to 2 hours, at -40

°C for 3 days in a portable freezer during transfer to the laboratory, and then at -80 °C until

RNA isolation. During hours of darkness, sampling occurred using green-filtered head

torches.

We wished to ensure that the abundance of transcripts could be compared between each

sampling season. We normalized all transcript measurements to the transcript levels in one

sample. Therefore, we obtained this reference sample for normalization of all RT-qPCR

experiments in the study by pooling RNA from 10 leaves sampled at noon on 26 March

2015, from 10 healthy plants across the field site that were each separated by at least 1

metre. This provided a reference cDNA sample against which all RT-qPCR analyses from all

sampling seasons were normalized within the ΔΔCt method [79], to allow comparability

between all datasets. The reference sample was collected at midday because all transcripts

under investigation were expressed to some extent at that time point. In all experiments,

dawn and dusk were defined as the astronomical (solar) time of sunrise and sunset.

RNA isolation and RT-qPCR

Frozen samples containing RNA*later* were defrosted in a cold room for 4 hours, the

RNA*later* was removed, and leaf tissue was transferred to new dry tubes and frozen in liquid

nitrogen. Frozen tissue was ground with a TissueLyzer (Qiagen, Hilden, Germany) and total

RNA was isolated from the powdered plant material using Macherey-Nagel Nucleospin II

RNA extraction kits (Thermo Fisher Scientific). RNA concentrations were determined using a

Nanodrop spectrophotometer (Thermo Fisher Scientific). cDNA was synthesized using a

High Capacity cDNA Reverse Transcription Kit (Thermo Fisher Scientific) and random

primers supplemented with RNAase inhibitor (Thermo Fisher Scientific), as described

previously [22, 23]. 1:500 cDNA dilutions were analyzed using Brilliant III Ultra-Fast SYBR

Green QPCR Master Mix (Agilent Technologies, Santa Clara, CA, USA), required primer

pairs (Table S2), and Agilent Mx3005P qPCR instrument. Primers were designed using the

PrimerQuest™ Tool (Integrated DNA Technologies, Coralville, IA, USA). Results were

normalized using the ΔΔCt method to *AhgACTIN2* [22, 23]. *AhgACTIN2* is encoded in *A.*

halleri by locus *g21632* [36] and has 97.8% coding sequence identity with *A. thaliana*

ACTIN2 (*At3g18780*). We selected *AhgACTIN2* as a reference transcript, because it has

been used previously as a reference transcript for experiments involving *A. halleri* at this

field site [43], *AhgACTIN2* does not fluctuate across the seasonal cycle [12], and in our

experiments, the RT-qPCR Ct for *AhgACTIN2* did not oscillate across the diel cycle

(*AhgACTIN2* time-series arrhythmic after analysis with JTK_CYCLE test for rhythmicity [80];

Fig. S12).

Environmental monitoring

The temperature and irradiance were measured beside the plants during sampling. The

temperature at each location, for each temperature manipulation, was monitored with EL-

USB-2 data loggers (Lascar Electronics, Whiteparish, UK) at 5-minute intervals.

Temperature loggers were wrapped in aluminium foil to prevent surface heating by solar

radiation. Irradiance was measured using a CC-3-UV-S cosine corrector connected to a

USB2000+ spectrometer with a QP400-2-UV-VIS fibre optic cable (Ocean Optics, Dunedin,

FL, USA). Ambient light spectra (200 nm to 900 nm) were collected every 5 minutes over the

14 hours of light during each day of sampling using OceanView software (Ocean Optics) on

a laptop PC, controlled by a custom script. The spectrometer and computer were powered

using portable lithium battery packs (Powertraveller, Hampshire, UK).

Smooth trend model

The smooth trend model (STM) allows inference of a trend within time-series data that

contains both sampling noise and biological stochasticity, and also allows a level of

statistical confidence to be applied to that trend. The STM to assess the difference in

transcript abundance between March and September under sun and shade conditions (Fig.

2) was defined by the equations:

$$\mu_{1,t} \sim Normal(2\mu_{1,t-1} - \mu_{1,t-2}, \ \sigma_{\mu1}^2), \tag{1}$$

$$\delta_t \sim Cauchy(\delta_{t-1}, \ \sigma_{\delta}^2), \tag{2}$$

$$\mu_{2,t} = \mu_{1,t} + \delta_t, \tag{3}$$

$$y_{1,t} \sim Normal(\mu_{1,t}, \ \sigma_y^2), \tag{4}$$

$$y_{2,t} \sim Normal(\mu_{2,t}, \ \sigma_y^2), \tag{5}$$

where μ_1 and μ_2 are the smooth trend components in March and September in 2015,

respectively, δ is the time-varying difference between the two seasons, and y_1 and y_2 are

the observed transcript abundance in the two seasons. $t = (1, 2, \cdots, 13)$ is the time point at

two-hour intervals. The same STM was used to analyze the difference in transcript

abundance between sun and shade conditions in March and September (Fig. S4).

For the models of the three (ambient, warm and cool) conditions in the temperature

manipulation experiment (Fig. 4), additional δ, μ and y were considered:

$$\delta_{2,t} \sim Cauchy(\delta_{2,t-1}, \ \sigma_{\delta2}^2), \tag{6}$$

$$\mu_{3,t} = \mu_{1,t} + \delta_{2,t}, \tag{7}$$

$$y_{3,t} \sim Normal(\mu_{3,t}, \ \sigma_y^2). \tag{8}$$

The parameters of the models were estimated by Bayesian inference using the Markov

Chain Monte Carlo (MCMC) approach. We note that unlike classical hypothesis testing

methods, multiple comparisons do not raise a problem in a Bayesian multilevel modelling

[81]. The statistical models were written in the Stan language and the programs were

compiled using CmdStan (v2.24). To operate CmdStan, the cmdstanr package (v0.4.0) in R

was used. After 1,000 warm-up steps, 1,000 MCMC samples were obtained by thinning out

3,000 MCMC samples for each of four parallel chains. Thus, 4,000 MCMC samples were

obtained in total. As prior distributions, we used a positive uniform distribution for σ, cauchy

distribution for δ (equations 2 and 6), and a normal distribution for μ and y (equations 1, 4, 5

and 8). A cauchy distribution of the priors was used to represent the difference between two

time series (δ) because this distribution corrects better for the influence of outliers, due to its

relatively long tails and its efficiency for detecting change points in time series data [82, 83].

Local level model with exogenous variables

The local level model with exogenous variables (LLMX) to analyze a diel trend and the effect

of environmental variables on transcript abundance (Fig. 3) was defined by the equations:

$$\mu_t \sim Normal(\mu_{t-1}, \ \sigma_\mu^2), \tag{9}$$

$$\alpha_{MarSun,t} = \mu_t + \beta_{temp} \cdot temp_{MarSun,t} + \beta_{irrad} \cdot irrad_{MarSun,t} + \beta_{gene} \cdot \tag{10}$$
$$gene_{MarSun,t},$$

$$\alpha_{MarShade,t} = \mu_t + \beta_{temp} \cdot temp_{MarShade,t} + \beta_{irrad} \cdot irrad_{MarShade,t} + \beta_{gene} \cdot \tag{11}$$

$$gene_{MarShade,t},$$

$$\alpha_{SepSun,t} = \mu_t + \beta_{temp} \cdot temp_{SepSun,t} + \beta_{irrad} \cdot irrad_{SepSun,t} + \beta_{gene} \cdot \qquad (12)$$

$$gene_{SepSun,t},$$

$$\alpha_{SepShade,t} = \mu_t + \beta_{temp} \cdot temp_{SepShade,t} + \beta_{irrad} \cdot irrad_{SepShade,t} + \beta_{gene} \cdot \qquad (13)$$

$$gene_{SepShade,t},$$

$$y_{MarSun,t} \sim Normal(\alpha_{MarSun,t}, \; \sigma_y^2), \qquad (14)$$

$$y_{MarShade,t} \sim Normal(\alpha_{MarShade,t}, \; \sigma_y^2), \qquad (15)$$

$$y_{SepSun,t} \sim Normal(\alpha_{SepSun,t}, \; \sigma_y^2), \qquad (16)$$

$$y_{SepShade,t} \sim Normal(\alpha_{SepShade,t}, \; \sigma_y^2), \qquad (17)$$

where μ is the autoregressive trend component that is common among conditions, β is the

regression coefficient, α is the true state of transcript abundance, y is the observed

transcript abundance, and σ^2 is the variance. The subscripts, $temp$, $irrad$, $gene$, Mar, Sep,

Sun and $Shade$ represent temperature, irradiance, an upstream gene, March, September,

sun condition and shade condition, respectively. The mean transcript abundance of the

upstream gene (i.e., *AhgCCA1* in the *AhgSIG5* model and *AhgSIG5* in the *AhgpsbD* BLRP

model) was used as one of the inputs. $t = (1, 2, \cdots, 13)$ is the time point at two-hour intervals.

The parameters of the models were estimated by Bayesian inference using the Markov

Chain Monte Carlo (MCMC) approach (Table S2, S3). The statistical models were written in

the Stan language and the programs were compiled using CmdStan (v2.24). To operate

CmdStan, the cmdstanr package (v0.4.0) in R was used. After 3,000 warm-up steps, 1,000

MCMC samples were obtained for each of the four parallel chains, and thus 4,000 MCMC

samples were obtained in total. As prior distributions, we used a uniform distribution for β, a

positive uniform distribution for σ, and a normal distribution for μ and y (equations 9 and 14-

17).

Embedding dimension

In the empirical dynamic modelling (EDM), time series is embedded into time-lagged series,

a procedure known as state space reconstruction [84, 85]. The embedding dimension *E* is

the dimension (i.e. the number of time-lagged series) used to reconstruct the state space.

Prior to convergent cross mapping (CCM), the optimal *E* value was determined for each of

AhgCCA1, *AhgSIG5* and *AhgpsbD* BLRP in temperature manipulation experiments in

September 2016, by univariate simplex projection [86], using the simplex function of the

rEDM package (v0.7.5) of R. We determined the optimal *E* values showing the maximum

forecast skill ρ (Fig. S11A). The optimal *E* value of 8 for *AhgCCA1* was relatively large,

considering those of *AhgSIG5* and *AhgpsbD* BLRP, and data points available in this

analysis. We adopted *E* value of 2 as the optimal value for *AhgCCA1*, because ρ is higher

than 0.9 for *E* values from 2 to 8 (Fig. S11A) and the model prediction well fits the observed

data at *E* value of 2 (Fig. S11B). The optimal *E* values determined were 2 for *AhgCCA1*, 4

for *AhgSIG5*, and 2 for *AhgpsbD* BLRP. The model predictions well fit the observed data for

these genes at the determined *E* values (Fig. S11B-D).

Convergent cross mapping

When a causal relationship exists from a variable *X* to another variable *Y*, the information of

X can be found in the time series of *Y*, meaning that the prediction of *X* is possible using the

information of *Y*. In convergent cross mapping (CCM) based on simplex projection, one

predicts the nearest neighbors of *X* at time *t* in the reconstructed lagged trajectory, using

their time-corresponding points of *Y* [45]. When this prediction (called cross mapping) is

successful, the causality from *X* to *Y* is assumed. The cross map skill (i.e. prediction skill) is

evaluated by Pearson's correlation coefficient (ρ) between predicted and observed values.

We performed CCM between *AhgCCA1* (*X*) and *AhgSIG5* (*Y*), and between *AhgSIG5* (*X*)

and *AhgpsbD* BLRP (*Y*) for the temperature manipulation experiments (September 2016),

using the ccm function in the rEDM package (v0.7.5) of R. We considered time delay in the

interactions by changing the parameter tp from –6 to 2 (negative, past; positive, future) in the

ccm function. When causality exists, optimal predictability is expected to occur for tp ≤ 0, i.e,

prediction of past values of *X* from *Y* [59]. We used a technique known as multispatial CCM

[87], to utilize the relatively small number of time points of 15/condition. Since we used three

temperature conditions (ambient, warm and cool), we applied the total of 45 time points to

CCM. To test the significance of cross map skill, we produced 1,000 diel surrogate time

series for *X* into which a similar level of oscillation was incorporated, but the sequence of

data deviation from the oscillation was randomized. We used three criteria for significant

causality. First, optimal predictability occurs for tp ≤ 0, second, cross map skill is above the

95 % interval of diel surrogates at the optimal time lag (tp), third, cross map skill is improved

according to the increase in a library size (number of time points used to reconstruct a state

space), known as convergence [45].

Acknowledgements

We thank Noriane M. L. Simon and Tasuku Ito for experimental assistance, and Paige E.

Panter and Deirdre A. Lynch for constructive feedback. This research was funded by

BBSRC (UK) (BB/I005811/2; BB/J014400/1; Institute Strategic Programme GEN

BB/P013511/1; Institute Strategic Programme bRiC BB/X01102X/1), The Royal Society

(IE140501), The Leverhulme Trust (RPG-2018-216), the Japan Society for Promotion of

Science (JSPS KAKENHI, JP21H04977, JP21H05659, JP21K15164), the Japan Science

and Technology Agency (JST, CREST no. JPMJCR1501) and the Bristol Centre for

Agricultural Innovation. DLCR is grateful to Consejo Nacional de Ciencia y Tecnología

(Mexico) for awarding a scholarship. This research was conducted through the Joint Usage

programme of the Center for Ecological Research of Kyoto University.

Author contributions

DLCR, HN, JS, MNH, HK and AND designed and performed experimentation; HN, DLCR,

TM, LLBD, HK and AND analyzed and interpreted data, and DLCR, HN, LLBD, HK and AND

wrote the paper.

Competing interests

The authors declare no competing interests.

Data and code availability

Source data are provided in Spreadsheet S1. Code is provided at

https://github.com/hnishio/SIG5_field3.git.

References

1.   Millar, A.J., *The intracellular dynamics of circadian clocks reach for the light of*
ecology and evolution. Annual Review of Plant Biology, 2016. **67**(1): p. 595-618.
2.   Kudoh, H., *Molecular phenology in plants: in natura systems biology for the*
comprehensive understanding of seasonal responses under natural environments.
New Phytologist, 2016. **210**(2): p. 399-412.
3.   Covington, M.F., et al., *Global transcriptome analysis reveals circadian regulation of*
key pathways in plant growth and development. Genome Biology, 2008. **9**(8): p.
R130.
4.   Edwards, K.D., et al., *FLOWERING LOCUS C mediates natural variation in the high-*
temperature response of the Arabidopsis circadian clock. The Plant Cell, 2006. **18**(3):
p. 639-650.
5.   Dodd, A.N., et al., *Plant circadian clocks increase photosynthesis, growth, survival,*
and competitive advantage. Science, 2005. **309**(5734): p. 630-633.
6.   Harmer, S.L., et al., *Orchestrated transcription of key pathways in Arabidopsis by the*
circadian clock. Science, 2000. **290**(5499): p. 2110-2113.
7.   Graf, A., et al., *Circadian control of carbohydrate availability for growth in Arabidopsis*
plants at night. Proceedings of the National Academy of Sciences, 2010. **107**(20): p.
9458-9463.
8.   Hotta, C.T., et al., *Modulation of environmental responses of plants by circadian*
clocks. Plant, Cell & Environment, 2007. **30**(3): p. 333-349.
9.   Nagano, Atsushi J., et al., *Deciphering and prediction of transcriptome dynamics*
under fluctuating field conditions. Cell, 2012. **151**(6): p. 1358-1369.

10.   Dantas, L.L.B., et al., *Alternative splicing of circadian clock genes correlates with*
661   *temperature in field-grown sugarcane.* Frontiers in Plant Science, 2019. **10**: p. article
662   1614.
663   11.   Dantas, L.L.B., et al., *Rhythms of transcription in field-grown sugarcane are highly*
664   *organ specific.* Scientific Reports, 2020. **10**(1): p. article 6565.
665   12.   Nagano, A.J., et al., *Annual transcriptome dynamics in natural environments reveals*
666   *plant seasonal adaptation.* Nature Plants, 2019. **5**(1): p. 74-83.
667   13.   Matsuzaki, J., Y. Kawahara, and T. Izawa, *Punctual transcriptional regulation by the*
668   *rice circadian clock under fluctuating field conditions.* The Plant Cell, 2015. **27**(3): p.
669   633-648.
670   14.   Annunziata, M.G., et al., *Response of Arabidopsis primary metabolism and circadian*
671   *clock to low night temperature in a natural light environment.* Journal of Experimental
672   Botany, 2018. **69**(20): p. 4881-4895.
673   15.   Annunziata, M.G., et al., *Getting back to nature: a reality check for experiments in*
674   *controlled environments.* Journal of Experimental Botany, 2017. **68**(16): p. 4463-
675   4477.
676   16.   Dantas, L.L.B., et al., *Field microenvironments regulate crop diel transcript and*
677   *metabolite rhythms.* New Phytologist, 2021. **232**(4): p. 1738-1749.
678   17.   Hashida, Y., et al., *Fillable and unfillable gaps in plant transcriptome under field and*
679   *controlled environments.* Plant, Cell & Environment, 2022. **45**(8): p. 2410-2427.
680   18.   Song, Y.H., et al., *Molecular basis of flowering under natural long-day conditions in*
681   *Arabidopsis.* Nature Plants, 2018. **4**(10): p. 824-835.
682   19.   Steed, G., et al., *Chronoculture, harnessing the circadian clock to improve crop yield*
683   *and sustainability.* Science, 2021. **372**(6541): p. eabc9141.
684   20.   O'Neill, C.M., et al., *Vernalization and floral transition in autumn drive winter annual*
685   *life history in oilseed rape.* Current Biology, 2019. **29**(24): p. 4300-4306.e2.
686   21.   Nagashima, A., et al., *The multiple-stress responsive plastid sigma factor, SIG5,*
687   *directs activation of the psbD blue light responsive promoter (BLRP) in Arabidopsis*
688   *thaliana.* Plant and Cell Physiology, 2004. **45**(4): p. 357-368.
689   22.   Noordally, Z.B., et al., *Circadian control of chloroplast transcription by a nuclear-*
690   *encoded timing signal.* Science, 2013. **339**(6125): p. 1316-1319.
691   23.   Belbin, F.E., et al., *Integration of light and circadian signals that regulate chloroplast*
692   *transcription by a nuclear-encoded sigma factor.* New Phytologist, 2017. **213**(2): p.
693   727-738.
694   24.   Onda, Y., et al., *Light induction of Arabidopsis SIG1 and SIG5 transcripts in mature*
695   *leaves: differential roles of cryptochrome 1 and cryptochrome 2 and dual function of*
696   *SIG5 in the recognition of plastid promoters.* The Plant Journal, 2008. **55**(6): p. 968-
697   978.
698   25.   Yamburenko, M.V., Y.O. Zubo, and T. Börner, *Abscisic acid affects transcription of*
699   *chloroplast genes via protein phosphatase 2C-dependent activation of nuclear*
700   *genes: repression by guanosine-3'-5'-bisdiphosphate and activation by sigma factor*
701   *5.* The Plant Journal, 2015. **82**(6): p. 1030-1041.
702   26.   Zhao, P., et al., *ATHB17 enhances stress tolerance by coordinating photosynthesis*
703   *associated nuclear gene and ATSIG5 expression in response to abiotic stress.*
704   Scientific Reports, 2017. **7**(1): p. article 45492.
705   27.   Mellenthin, M., et al., *Expression of the Arabidopsis sigma factor SIG5 is*
706   *photoreceptor and photosynthesis controlled.* Plants, 2014. **3**: p. 359-391.
707   28.   Tsunoyama, Y., et al., *Blue light specific and differential expression of a plastid σ*
708   *factor, Sig5 in Arabidopsis thaliana.* FEBS Letters, 2002. **516**(1-3): p. 225-228.
709   29.   Mochizuki, T., et al., *Two independent light signals cooperate in the activation of the*
710   *plastid psbD blue light-responsive promoter in Arabidopsis.* FEBS Letters, 2004.
711   **571**(1): p. 26-30.
712   30.   Møller, S.G., et al., *PP7 is a positive regulator of blue light signaling in Arabidopsis.*
713   The Plant Cell, 2003. **15**(5): p. 1111-1119.

31.   Davey, M.P., et al., *The UV-B photoreceptor UVR8 promotes photosynthetic*
efficiency in Arabidopsis thaliana exposed to elevated levels of UV-B. Photosynthesis
Research, 2012. **114**(2): p. 121-131.
32.   Brown, B.A. and G.I. Jenkins, *UV-B signaling pathways with different fluence-rate*
response profiles are distinguished in mature Arabidopsis leaf tissue by requirement
for UVR8, HY5, and HYH. Plant Physiology, 2008. **146**(2): p. 576-588.
33.   Cano-Ramirez, D.L., et al., *Low-temperature and circadian signals are integrated by*
the sigma factor SIG5. Nature Plants, 2023. **9**(4): p. 661-672.
34.   Panter, P.E., et al., *Circadian regulation of the plant transcriptome under natural*
conditions. Frontiers in Genetics, 2019. **10**: p. article 1239.
35.   Honjo, M.N. and H. Kudoh, *Arabidopsis halleri: a perennial model system for*
studying population differentiation and local adaptation. AoB PLANTS, 2019. **11**(6):
p. article plz076.
36.   Briskine, R.V., et al., *Genome assembly and annotation of Arabidopsis halleri, a*
model for heavy metal hyperaccumulation and evolutionary ecology. Molecular
Ecology Resources, 2017. **17**(5): p. 1025-1036.
37.   Tsunoyama, Y., et al., *Blue light-induced transcription of plastid-encoded psbD gene*
is mediated by a nuclear-encoded transcription initiation factor, AtSig5. Proceedings
of the National Academy of Sciences of the United States of America, 2004. **101**(9):
p. 3304-3309.
38.   Shimmura, S., et al., *Comparative analysis of chloroplast psbD promoters in*
terrestrial plants. Frontiers in Plant Science, 2017. **8**: p. article 1186.
39.   Locke, J.C.W., A.J. Millar, and M.S. Turner, *Modelling genetic networks with noisy*
and varied experimental data: the circadian clock in Arabidopsis thaliana. Journal of
Theoretical Biology, 2005. **234**(3): p. 383-393.
40.   Kitagawa, G., *Introduction to time series modeling: with applications in R.*
Monographs on Statistics and Applied Probability 166. 2020, New York: CRC Press.
41.   Laine, M., *Introduction to dynamic linear models for time series analysis*, in *Geodetic*
Time Series Analysis in Earth Sciences, J.-P. Montillet and M.S. Bos, Editors. 2020,
Springer International Publishing: Cham. p. 139-156.
42.   Nishio, H., S. Hirano, and Y. Kodama, *Statistical analysis of organelle movement*
using state-space models. Plant Methods, 2023. **19**(1): p. 67.
43.   Aikawa, S., et al., *Robust control of the seasonal expression of the Arabidopsis FLC*
gene in a fluctuating environment. Proceedings of the National Academy of Sciences,
2010. **107**(25): p. 11632-11637.
44.   Nishio, H., et al., *Repressive chromatin modification underpins the long-term*
expression trend of a perennial flowering gene in nature. Nature Communications,
2020. **11**(1): p. article 2065.
45.   Sugihara, G., et al., *Detecting causality in complex ecosystems.* Science, 2012.
338(6106): p. 496-500.
46.   Satake, A., et al., *Field transcriptome revealed a novel relationship between nitrate*
transport and flowering in Japanese beech. Scientific Reports, 2019. **9**(1): p. 4325.
47.   Dodd, A.N., et al., *The Arabidopsis circadian clock incorporates a cADPR-based*
feedback loop. Science, 2007. **318**(5857): p. 1789.
48.   Michael, T.P., et al., *Network discovery pipeline elucidates conserved time-of-day–*
specific cis-regulatory modules. PLOS Genetics, 2008. **4**(2): p. e14.
49.   Ward, E.J., et al., *Smoothed dynamic factor analysis for identifying trends in*
multivariate time series. Methods in Ecology and Evolution, 2022. **13**(4): p. 908-918.
50.   Triantafyllopoulos, K., *Bayesian Inference of state space models.* 2021: Springer.
51.   Fraser, D.P., et al., *Phytochrome A elevates plant circadian-clock components to*
suppress shade avoidance in deep-canopy shade. Proceedings of the National
Academy of Sciences, 2021. **118**(27): p. e2108176118.
52.   Commandeur, J.J.F. and S.J. Koopman, *An introduction to state space time series*
analysis. 2007, Oxford: Oxford University Press.

53.   Pittendrigh, C.S. and D. Skopik Steven, *Circadian Systems, V. The driving oscillation*
and the temporal sequence of development. Proceedings of the National Academy of
Sciences, 1970. **65**(3): p. 500-507.
54.   Finster, S., et al., *Light-dependent, plastome-wide association of the plastid-encoded*
RNA polymerase with chloroplast DNA. Plant Journal, 2013. **76**(5): p. 849-860.
55.   Hoffer, P.H. and D.A. Christopher, *Structure and blue-light-responsive transcription of*
a chloroplast psbD promoter from Arabidopsis thaliana. Plant Physiology, 1997.
115(1): p. 213-222.
56.   Satoh, J., et al., *Developmental stage-specific multi-subunit plastid RNA polymerases*
(PEP) in wheat. Plant Journal, 1999. **18**(4): p. 407-415.
57.   Baena-González, E., et al., *Chloroplast transcription at different light intensities.*
Glutathione-mediated phosphorylation of the major RNA polymerase involved in
redox-regulated organellar gene expression. Plant Physiology, 2001. **127**(3): p. 1044-
1052.
58.   Nakamura, T., et al., *Array-based analysis on tobacco plastid transcripts: Preparation*
of a genomic microarray containing all genes and all intergenic regions. Plant and
Cell Physiology, 2003. **44**(8): p. 861-867.
59.   Ye, H., et al., *Distinguishing time-delayed causal interactions using convergent cross*
mapping. Scientific Reports, 2015. **5**(1): p. 14750.
60.   Bieniawska, Z., et al., *Disruption of the Arabidopsis circadian clock is responsible for*
extensive variation in the cold-responsive transcriptome. Plant Physiology, 2008.
147(1): p. 263-279.
61.   Gould, P.D., et al., *The molecular basis of temperature compensation in the*
Arabidopsis circadian clock. The Plant Cell, 2006. **18**(5): p. 1177-1187.
62.   Gould, P.D., et al., *Network balance via CRY signalling controls the Arabidopsis*
circadian clock over ambient temperatures. Molecular Systems Biology, 2013. **9**(1):
p. article 650.
63.   Wang, Z.-Y. and E.M. Tobin, *Constitutive expression of the CIRCADIAN CLOCK*
ASSOCIATED 1 gene disrupts circadian rhythms and suppresses its swn expression.
Cell, 1998. **93**(7): p. 1207-1217.
64.   Wang, Z.Y., et al., *A Myb-related transcription factor is involved in the phytochrome*
regulation of an Arabidopsis Lhcb gene. The Plant Cell, 1997. **9**(4): p. 491-507.
65.   Troein, C., et al., *Weather and seasons together demand complex biological clocks.*
Current Biology, 2009. **19**(22): p. 1961-1964.
66.   Shalit-Kaneh, A., et al., *Multiple feedback loops of the Arabidopsis circadian clock*
provide rhythmic robustness across environmental conditions. Proceedings of the
National Academy of Sciences, 2018. **115**(27): p. 7147-7152.
67.   Grabe, S., et al., *Synergies of multiple zeitgebers tune entrainment.* Frontiers in
Network Physiology, 2022. **1**.
68.   Breit, A., et al., *Insulin-like growth factor-1 acts as a zeitgeber on hypothalamic*
circadian clock gene expression via glycogen synthase kinase-3B signaling. Journal
of Biological Chemistry, 2018. **293**(44): p. 17278-17290.
69.   Mizuno, T., et al., *Ambient temperature signal feeds into the circadian clock*
transcriptional circuitry through the EC night-time repressor in Arabidopsis thaliana.
Plant and Cell Physiology, 2014. **55**(5): p. 958-976.
70.   Thines, B. and G. Harmon Frank, *Ambient temperature response establishes ELF3*
as a required component of the core Arabidopsis circadian clock. Proceedings of the
National Academy of Sciences, 2010. **107**(7): p. 3257-3262.
71.   Catalá, R., J. Medina, and J. Salinas, *Integration of low temperature and light*
signaling during cold acclimation response in Arabidopsis. Proceedings of the
National Academy of Sciences, 2011. **108**(39): p. 16475-16480.
72.   Shimizu, M., et al., *Sigma factor phosphorylation in the photosynthetic control of*
photosystem stoichiometry. Proceedings of the National Academy of Sciences, 2010.
107(23): p. 10760-10764.

73. Ling, Q. and P. Jarvis, *Regulation of chloroplast protein import by the ubiquitin E3*
ligase SP1 is important for stress tolerance in plants. Current Biology, 2015. **25**(19):
p. 2527-2534.
74. Michael, T.P., P.A. Salomé, and C.R. McClung, *Two Arabidopsis circadian oscillators*
can be distinguished by differential temperature sensitivity. Proceedings of the
National Academy of Sciences, 2003. **100**(11): p. 6878-6883.
75. Lee, N., et al., *The FLOWERING LOCUS T gene expression is controlled by high-*
irradiance response and external coincidence mechanism in long days in
Arabidopsis. New Phytologist, 2023. **239**(1): p. 208-221.
76. Kudoh, H., et al., *The long-term "in natura" study sites of Arabidopsis halleri for plant*
transcription and epigenetic modification analyses in natural environments, in *Plant*
Transcription Factors: Methods and Protocols, N. Yamaguchi, Editor. 2018, Springer
New York: New York, NY. p. 41-57.
77. Kamitani, M., et al., *RNA-Seq reveals virus–virus and virus–plant interactions in*
nature. FEMS Microbiology Ecology, 2016. **92**(11): p. article fiw176.
78. Franklin, K.A., *Shade avoidance.* New Phytologist, 2008. **179**(4): p. 930-944.
79. Livak, K.J. and T.D. Schmittgen, *Analysis of relative gene expression data using real-*
rime quantitative PCR and the 2−ΔΔCT method. Methods, 2001. **25**(4): p. 402-408.
80. Hughes, M.E., J.B. Hogenesch, and K. Kornacker, *JTK_CYCLE: An efficient*
nonparametric algorithm for detecting rhythmic components in genome-scale data
sets. Journal of Biological Rhythms, 2010. **25**(5): p. 372-380.
81. Gelman, A., J. Hill, and M. Yajima, *Why we (usually) don't have to worry about*
multiple comparisons. Journal of Research on Educational Effectiveness, 2012. **5**(2):
p. 189-211.
82. Peluso, S., S. Chib, and A. Mira, *Semiparametric multivariate and multiple change-*
point modeling. Bayesian Analysis, 2019. **14**(3): p. 727-751.
83. Austin, E., et al., *Online non-parametric changepoint detection with application to*
monitoring operational performance of network devices. Computational Statistics &
Data Analysis, 2023. **177**: p. 107551.
84. Takens, F., *Dynamical systems and turbulence*, in *Lecture Notes in Mathematics*, D.
Rand and L.S. Young, Editors. 1981, Springer-Verlag: New York. p. 366-381.
85. Chang, C.-W., M. Ushio, and C.-h. Hsieh, *Empirical dynamic modeling for beginners.*
Ecological Research, 2017. **32**(6): p. 785-796.
86. Sugihara, G. and R.M. May, *Nonlinear forecasting as a way of distinguishing chaos*
from measurement error in time series. Nature, 1990. **344**(6268): p. 734-741.
87. Clark, A.T., et al., *Spatial convergent cross mapping to detect causal relationships*
from short time series. Ecology, 2015. **96**(5): p. 1174-1181.

March (Spring)   September (Autumn)

A *AhgSIG5* transcript (sun)

B *AhgSIG5* transcript (shade)

C Diel trend

D Regression coefficients

Temperature   Irradiance   *AhgCCA1*

E *AhgpsbD BLRP* transcript (sun)

F *AhgpsbD BLRP* transcript (shade)

G Diel trend

H Regression coefficients

Temperature   Irradiance   *AhgSIG5*

— March (Spring)   — September (Autumn)