Circadian and environmental signal transduction in a natural population of

Arabidopsis

Dora L. Cano-Ramirez[1,2,7], Haruki Nishio[3,4,7], Tomoaki Muranaka[3,5], Luíza Lane de Barros

Dantas[6], Mie N. Honjo[3], Jiro Sugisaka[3], Hiroshi Kudoh[3], Antony N. Dodd[6*].

1. Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge, UK.

2. School of Biological Sciences, University of Bristol, Bristol, UK.

3. Center for Ecological Research, Kyoto University, Otsu, Shiga, Japan.

4. Data Science and AI Innovation Research Promotion Center, Shiga University, Hikone,

Shiga, Japan.

5. Department of Environmental Sciences and Technology, Faculty of Agriculture,

Kagoshima University, Korimoto, Kagoshima, Japan.

6. Department of Cell and Developmental Biology, John Innes Centre, Norwich, UK.

7. Equal contribution to this research.

* Corresponding author; antony.dodd@jic.ac.uk

Word counts: Main text (inc. methods) 5524; Abstract 198.

Abstract

Plants sense and respond to environmental cues during 24 h fluctuations in their

environment. This requires the integration of internal cues such as circadian timing with

environmental cues such as light and temperature to elicit cellular responses through signal

transduction. The integration and transduction of circadian and environmental signals within

plants growing in natural environments remain poorly understood. To gain insights into the

24 h dynamics of environmental signalling in nature, we performed a field study of cell

signalling in a natural population of *Arabidopsis halleri.* As a representative model signalling

pathway, we exploited the transduction of circadian and environmental signals from the

nucleus to chloroplasts, by a sigma factor, to study diel cycles of environmental signalling

under natural conditions. Using dynamic linear models to interpret the data, we identified that

circadian regulation and temperature are key regulators of this pathway under natural

conditions. We identified potential time-delay steps between pathway components, and diel

fluctuations in the response of the pathway to temperature cues that are reminiscent of the

process of circadian gating. This approach allowed us to identify dynamic integration and

transduction of environmental cues, in the cells of plants, under naturally fluctuating diel

cycles.

Introduction

Plants have sophisticated environmental sensing and signalling mechanisms that underpin

their responses to the fluctuating environment. Under naturally fluctuating conditions, this

requires signalling pathways that integrate dynamic, overlapping and complex environmental

stimuli [1, 2]. These environmental fluctuations include the 24 h changes in environmental

conditions that arise from the cycle of day and night. The 24 h environmental fluctuations

have selected for the evolution of circadian clocks, which are endogenous biological

oscillators that produce a cellular estimate of the time of day. Over each day, circadian

rhythms structure the responses of plants to environmental fluctuations by aligning

transcription, metabolism and development with the daily fluctuating environment [3-8]. In

plants, environmental information including the light and temperature conditions is used to

adjust the phase of the circadian oscillator, through the process of entrainment, so that the

phase is aligned with the 24 h environmental cycle. This alignment between the circadian

oscillator and the 24 fluctuating environment contributes to the fitness of plants [5].

Under natural conditions, circadian timing information is combined with environmental cues

to establish a temporal program of gene expression [9]. For example, 97% of diel transcript

profiles in field-grown rice can be predicted from meteorological data [9], and temperature

cues regulate the alternative splicing of transcripts encoding circadian oscillator components

in field-grown sugarcane [10]. Recent studies have provided insights into the diel

organization of the transcriptome and metabolism, under field conditions, for several crops

and natural plant populations [9-17]. However, the diel dynamics of environmental signalling

pathways, with defined inputs and outputs, are less well understood under natural

conditions. Understanding signal transduction in plants under natural conditions is a valuable

part of translating laboratory studies into crop improvement. For example, this could

contribute to forecasting the responses of ecosystems and crops to increasingly

unpredictable climates [18]. Experiments conducted in controlled conditions that mimic

components of field conditions remain unable to replicate all aspects of plant gene regulation

under natural conditions [17], so field experiments provide valuable insights into plant

environmental responses.

To study the integration and transduction of circadian and environmental signals under

naturally fluctuating conditions, we selected a well-characterized environmental signalling

pathway as an experimental model. This comprises the regulation by the circadian clock of

SIGMA FACTOR 5 (SIG5), which in turn regulates the transcription of *psbD* (Fig. 1A). *CCA1*

and *SIG5* are nuclear encoded, and *psbD* is chloroplast encoded. We chose this pathway

because it is relatively straightforward, consisting of three major components, each of which

provides distinct regulatory points of signal transduction. Therefore, the pathway provides a

relatively low level of complexity to evaluate circadian and environmental signal integration

and transduction under realistic field conditions. CCA1 is a key component of the

Arabidopsis circadian oscillator, and *CCA1* transcript abundance can be used as a proxy for

the status of the circadian oscillator (Fig. 1A). SIG5 is a nuclear-encoded regulator of

chloroplast transcription, which is regulated closely by the circadian oscillator under constant

conditions [20]. Based on its responses under controlled conditions [19-30], we hypothesized

that under natural conditions SIG5 might integrate information concerning circadian

regulation, light quantity, light quality, temperature, and abiotic stress. Therefore, *SIG5*

transcript abundance presents a read-out several environmental signal integration processes

(Fig. 1A). It is thought that SIG5 is imported into chloroplasts, and communicates the

integrated environmental information to chloroplast gene expression by regulating

transcription from the blue light responsive promoter of *psbD* (*psbD* BLRP) [22] that encodes

the D2 protein of Photosystem II (Fig. 1A). This chloroplast transcript provides an

experimental read-out of a later step in this signalling pathway (Fig. 1A) [19, 20, 22].

We investigated the temporal dynamics of this signalling pathway in a natural habitat of the

perennial *Arabidopsis* species, *Arabidopsis halleri* subsp. *gemmifera* (referred to here as *A.*

halleri) [55]. The close-relatedness of *A. halleri* and *A. thaliana* makes it possible to identify

pairs of homologous genes based on the sequence similarity (indicated by *Ahg* or *At* prefixes

to gene names) [56]. The circadian clock-SIG5-*psbD* BLRP pathway is present in *A. thaliana*

and *A. halleri*, and well-conserved across the vascular plants [31, 32]. We obtained a

number of time series, during two seasons of the year, that monitored pathway function

under representative light and temperature conditions. We interpreted these data using

dynamic linear models, which are a type of state space model derived from control theory. In

these models, the state of the system can be predicted from the prior state of the system,

onto which can be superimposed external effects. This allows the estimation of the dynamics

of the system that arise from its internal dynamics and external factors (such as

environmental cues). Using this approach, we identified key roles for temperature and the

circadian clock in the regulation of this pathway under natural conditions, and obtained

evidence for temporal gating of responses of the pathway to environmental cues. Our

approaches could be applicable to the study of many circadian-regulated processes under

naturally fluctuating conditions.

Results

Biological data underlying models of signal transduction

Under controlled conditions of constant light, *AtCCA1* and *AtSIG5* transcript abundance are

very well correlated (Fig. S1A-D; data from [20]). This correlation between *AtCCA1* and

AtSIG5 transcript abundance is absent under light/dark cycles (Fig. S1E, F), suggesting that

the integration of light and dark cues alters the diel regulation of *AtSIG5* transcript

accumulation [19, 20]. We acquired time-series of transcript abundance during spring

(March) and autumn/fall (September), close to the spring or autumn equinox (Fig. 1B-K; Fig.

S2). Although both the spring and autumn equinoxes share 12-h photoperiods, they provide

contrasting temperature regimes (cool and warm, respectively) (Fig. 1B, C; Fig. S2A, B), with

irradiance levels determined by weather conditions (Fig. 1D, E; Fig. S2C, D). This allowed us

to investigate temperature, light and seasonal influences upon SIG5-mediated signalling to

chloroplasts, because the pathway is known to be affected by light and temperature in *A.*

thaliana [19, 21, 22, 26, 31]. We obtained data from areas with open sky and with

vegetational shade, to include within our models the transcriptional responses to a wider

range of irradiance levels (Fig. 1D, E; Fig. S3). The "sun" and "shade" sampling sites were

chosen by measurement of the ratio of red to far-red light (R:FR) (Fig. S3C, D) and

availability of plant patches, because *A. halleri* does not grow in deep shade at this location.

The total light intensity at the sun sampling site was 5 to 10-fold greater during March 2015

than during September 2015, depending on the time of day, due to weather differences (Fig.

1D, E). During March 2015, the study site temperature at both sun and shade sites ranged

from 0 °C to 17 °C (Fig. 1B, C). The temperature was often above 20 °C during September

2015, with greater diel fluctuations at the sun site (Fig. 1B, C).

We compared the pathway dynamics between the spring and autumn sampling periods by

using a smooth trend model, and identified differences in pathway regulation (Fig. 1F-K and

Fig. S2E-J). We estimated the parameters of a smooth trend model by Bayesian inference to

visualize the differences in transcript abundance between the spring and autumn sampling.

The morning peak accumulation of transcripts encoding the circadian clock component

AhgCCA1 was significantly greater during the autumn sampling compared with the spring,

under both light conditions tested (Fig. 1F, G). During both sampling seasons, *AhgSIG5*

transcripts reached peak abundance between the middle and end of the photoperiod (Fig.

1H, I). This differs from the phase of *AtSIG5* transcript accumulation under square-wave

light/dark cycles under controlled conditions, where *AtSIG5* transcript abundance peaks

around dawn [20]. The peak of the diel fluctuation of *AhgSIG5* was significantly greater

during the March sampling period than during the September sampling period (Fig. 1H, I).

Furthermore, the pre-dawn accumulation of *AhgSIG5* transcripts occurred at an earlier time

during the dark period during September than during March (Fig. 1H, I). This delay in pre-

dawn transcript accumulation might be due to weaker circadian control during the spring

sampling period, as suggested by the significantly decreased peak height of *AhgCCA1*

transcript oscillations during March compared with September (Fig. 1F, G). Transcripts

encoding the SIG5 regulatory target *AhgpsbD* BLRP (Fig. 1A) had a significantly greater

peak of accumulation during the September sampling season compared with the March

season, but only under the shade light conditions (Fig. 1J, K).

AhgCCA1 transcript abundance was significantly greater under shade conditions during the

photoperiod, during both sampling seasons (Fig. S2E, F). This is reminiscent of the greater

AtCCA1 promoter activity that occurs directly after dawn under controlled conditions of far

red light compared with red light [33]. We did not identify the diminished *AtCCA1* oscillation

that occurs under constant light with a very low R:FR [34] or on the shaded western side of

crop fields around dawn [16]. As with *AhgCCA1*, *AhgSIG5* transcript abundance was

significantly greater under shade than sun conditions, with this difference restricted to the

end of the photoperiod (Fig. S2G, H). *AhgpsbD* BLRP transcript levels were unaltered by the

two light environments (Fig. S2I, J).

Time delays between signalling pathway components

We assumed that SIG5-mediated signalling to chloroplasts involves a hierarchically-

organized pathway, whereby *AhgCCA1* is positioned upstream from the regulation of

AhgSIG5 transcript accumulation, and *AhgpsbD* BLRP is positioned downstream of

AhgSIG5 activity (Fig. 1A). We also assumed that environmental signals might influence

AhgCCA1, *AhgSIG5* and *AhgpsbD* BLRP transcript accumulation independently (Fig. 1A)

[35]. To understand the dynamics of this process, we first considered the temporal

relationship between *AhgCCA1*, *AhgSIG5* and *AhgpsbD* BLRP transcript accumulation

under natural conditions (*t*1, *t*2; Fig. 1A). The abundance of each of these related transcripts

was monitored at each timepoint, but their responses to each other might not be

instantaneous. For example, in *A. thaliana* under controlled square-wave light/dark cycle

conditions, *AtCCA1* transcript abundance peaks at dawn, *AtSIG5* approximately 3 h after

dawn, and *AtpsbD* BLRP approximately 6 hours after dawn [20]. Comparable dynamics are

present in our field data, whereby *AhgCCA1* peaks at or after solar dawn (Fig. 1F, G),

AhgSIG5 mid-photoperiod (Fig. 1H, I), and *AhgpsbD* BLRP towards the end of the

photoperiod under those conditions where it is rhythmic (Fig. 1J, K). Therefore, we reasoned

that there would be a time lag in the regulation of *AhgSIG5* by *AhgCCA1* (*t1* in Fig. 1A), and

in the regulation of *AhgpsbD* BLRP by *AhgSIG5* (*t2* in Fig. 1A).

We were interested to quantify these time lags, and use the information arising to construct

models that assess the regulation of the pathway by specific environmental variables. We

developed dynamic linear models that predict the abundance of *AhgSIG5* from *AhgCCA1*

transcript abundance, and predict *AhgpsbD* BLRP from *AhgSIG5* transcript abundance,

together with temperature and irradiance as explanatory variables. In these models, we used

the transcript abundance of the upstream component as an explanatory variable, and the

transcript abundance of the target component as a response variable within this analysis.

Therefore, *AhgSIG5* is the response variable in the first model with *AhgCCA1* as the

explanatory variable, whilst in the second model, *AhgSIG5* is used as the explanatory

variable for *AhgpsbD.* We tested the quality of model fit for a range of time delays (lags)

between the genes in the pathway (Fig. 1A). For this, we compared three model selection

parameters to estimate the time delay that provides the best estimation of the downstream

transcript (response variable). For a prediction of *AhgSIG5* from *AhgCCA1*, a model

containing a 6 h time lag produced the best model fit, according to three model selection

parameters (Fig. 2A, B, C). For a prediction of *AhgpsbD* BLRP from *AhgSIG5*, a model

containing a 4 h time lag produced the best model fit for two out of three model selection

parameters (Fig. 2D, E, F).

We detected differences between the two sampling seasons in the time lags that produced

the best model selection parameters (Fig. S4). During the March sampling season, the best

prediction of either *AhgSIG5* or *AhgpsbD* BLRP arose when time lags of 6-8 h (*AhgSIG5*

prediction from *AhgCCA1*) and 4-8 h (*AhgpsbD* BLRP prediction from *AhgSIG5*) were tested

(Fig. S4A-F). This was relatively longer than during the September sampling season, when

time lags of 0 or 4 h (*AhgSIG5* prediction from *AhgCCA1*) and 4 h (*AhgpsbD* BLRP

prediction from *AhgSIG5*) produced the best model fit estimates (Fig. S4G-L). This suggests

that the low temperature in March delays the speed of signal transduction. Taken together,

these analyses suggest that time delays in signalling pathways are detectable under field

conditions, and that environmental conditions (seasonal differences in temperature, or

seasonal regulation) might affect the speed of signal transduction.

Dynamics of environmental regulation of signalling pathway

Environmental fluctuations are complex, noisy, and occur in simultaneous combinations.

This presents a challenge for interpreting time-series transcript data from the field within the

context of environmental signalling. We elaborated upon our modelling approach to

investigate the relationship between key environmental variables and SIG5-mediated

signalling to chloroplasts under field conditions. We used statistical models, rather than

models of biochemical kinetics [36], because this provides an effective tool for interpreting

diel and seasonal transcriptome dynamics [9, 12, 37, 38]. Comparable approaches have

allowed the investigation of diel and seasonal changes of transcriptome dynamics in *A.*

halleri [12, 37, 38] and rice [9].

We represented the behaviour of the pathway components using dynamic linear models, into

which the time delays that produced the best model fit were incorporated (Fig. 2). The output

of the Bayesian estimation reproduced well the essential dynamics of the observed

AhgCCA1 transcript level (Fig. 3A, B). The model estimated a significant positive relationship

between ambient temperature and *AhgCCA1* transcript abundance between midnight and

midday, with no effect of temperature at other times (Fig. 3C). There was no significant effect

of irradiance upon the estimation of *AhgCCA1* transcript abundance (Fig. 3D).

Diel fluctuations of *AhgSIG5* transcript abundance were reproduced well by the model (Fig.

3E, F). We identified a significant negative correlation between ambient temperature and

AhgSIG5 transcript level towards the end of the light period (Fig. 3G), whereas there was no

significant effect of irradiance upon the prediction of *AhgSIG5* transcript abundance (Fig.

3H). *AhgSIG5* is regulated by the circadian clock, and we included within the model

AhgCCA1 transcript abundance as a proxy for circadian clock dynamics. There was a

significant positive coefficient of regression between *AhgCCA1* and *AhgSIG5* during the dark

period, and around the middle of the photoperiod (Fig. 3I).

Diel fluctuations of chloroplast *psbD* BLRP transcript abundance were also predicted well

(Fig. 3J, K). In this case, there was a significant positive coefficient of regression between

ambient temperature and *psbD* BLRP transcript abundance, which was restricted to the light

period (Fig. 3L). There was no significant coefficient of regression between irradiance and

psbD BLRP transcript abundance (Fig. 3M). *psbD* BLRP transcript accumulation is regulated

in *A. thaliana* by SIG5 [22], and within our model, there was a significant positive coefficient

of regression between *AhgSIG5* and *AhgpsbD* BLRP transcript levels during part of the light

period (Fig. 3N).

Together, this analysis identifies that the ambient temperature, rather than the irradiance,

was important for predicting the dynamics of all pathway components under naturally

fluctuating conditions. In addition, the effect of the circadian clock (*AhgCCA1*) contributed to

the prediction of *AhgSIG5* transcript abundance (Fig. 3I), and the effect of SIG5 contributed

to the prediction of *AhgpsbD* BLRP transcript abundance (Fig. 3N). A feature within these

predictions was the restriction to specific times of day of significant coefficients of regression

between transcripts abundance and certain variables (such as for temperature in the

prediction of *AhgCCA1* (Fig. 3C) and *AhgSIG5* (Fig. 3G)). These 24-h fluctuations in the

coefficient of regression are suggestive of the concept of circadian gating, which is the

process whereby the circadian clock constrains certain biological processes to specific times

in the 24 h cycle [39]. In plants, this often takes the form of a circadian rhythm in the

magnitude of the response to identical environmental stimuli given at different times of day

[8].

251 **Temporal gating of temperature regulation of SIG5-mediated signalling to**

252 **chloroplasts under natural conditions**

253 Our dynamic linear modelling analysis suggested that under natural conditions, greater

254 ambient temperatures upregulate *AhgCCA1* and *AhgpsbD* BLRP transcript levels, whereas

255 lower ambient temperatures upregulate *AhgSIG5* transcript levels (Fig. 3C, G, L). We tested

256 this hypothesis by applying moderate temperature manipulations to adjacent patches of *A.*

257 *halleri* plants, in the field, using custom-designed equipment (Fig. 4A; Fig. S5). We collected

258 24-h time-series of RNA samples from these plant patches, and interpreted the data with

259 smooth trend models. The moderate temperature increase caused a small significant

260 upregulation of *AhgCCA1* transcript abundance after dawn relative to the control, whereas

261 the temperature reduction treatment was without effect (Fig. 4B). The moderate temperature

262 increase was without effect upon *AhgSIG5* transcript abundance, whereas the temperature

263 reduction treatment upregulated *AhgSIG5* transcripts significantly immediately after dawn,

264 relative to the control, and caused a significant reduction in transcript abundance during the

265 afternoon (Fig. 4C). This is consistent with the negative coefficient of regression between

266 *AhgSIG5* and temperature under naturally fluctuating conditions (Fig. 3G), and with the

267 upregulation of *A. thaliana SIG5* by a short cold treatment under laboratory conditions [22].

268 The restriction of the response of *AhgSIG5* transcripts to the moderate temperature

269 reduction (Fig. 4C) is consistent with the 24-h cycle of the magnitude of the coefficient of

270 regression of temperature for *AhgSIG5* transcript abundance (Fig. 3G). This further supports

271 the notion of temporal gating of the influence of temperature upon this pathway under

272 naturally fluctuating conditions.

273 Transcripts for the chloroplast target of SIG5, *AhgpsbD* BLRP, were also altered by

274 temperature manipulation. The moderate temperature elevation significantly increased

275 *AhgpsbD* BLRP transcript levels relative to the control, whereas the moderate temperature

276 reduction significantly reduced *AhgpsbD* BLRP transcripts relative to the control. These

277 significant alterations were restricted to the photoperiod, which might be because chloroplast

DNA binding and transcription by PEP generally requires light [40-44]. Furthermore, the

positive regulation of *AhgpsbD* BLRP transcript abundance by the temperature

manipulations (Fig. 4D) is consistent with the coefficient of regression between *AhgpsbD*

BLRP transcript abundance and temperature under naturally fluctuating conditions (Fig. 3L).

Discussion

We established that circadian regulation and ambient temperature are potential regulators of

AhgSIG5-mediated signalling to chloroplasts, in a natural population of *A. halleri*. Our

analysis identified a significant regulation of *AhgSIG5* by *AhgCCA1* from midnight to morning

(Fig. 3I), and significant regulation of *AhgpsbD* BLRP by *AhgSIG5* towards the end of the

photoperiod (Fig. 3N). These significant relationships suggest that under natural conditions,

a signal is communicated from the circadian oscillator (using *AhgCCA1* as a proxy) to the

signalling pathway output of *AhgpsbD* BLRP. One interpretation is that the pathway couples

the circadian oscillator and temperature response processes to chloroplast gene

transcription under naturally fluctuating conditions.

We identified seasonal differences in the maximum accumulation of *AhgCCA1*, *AhgSIG5*

and *AhgpsbD* BLRP. *AhgCCA1* and *AhgpsbD* BLRP had significantly lower peak

accumulation during the spring sampling period compared with the autumn sampling

season. In comparison, *AhgSIG5* had significantly greater peak accumulation during the

spring compared with the autumn sampling season. The difference in *AhgCCA1* dynamics

between these sampling seasons likely reflects the decreased amplitude of the circadian

oscillator that occurs under lower temperature conditions, in both controlled environments

and the field [12, 45-47]. The difference in dynamics of *AhgSIG5* compared with *AhgCCA1*

suggests that an additional temperature input into this pathway occurs between the circadian

oscillator and *AhgSIG5*. In *A. thaliana*, *SIG5* transcripts are upregulated by short cold

temperature treatments [22] and in our field experiment, *AhgSIG5* transcript accumulation

had a negative coefficient of regression with the temperature (Fig. 3G, Fig. 4C). This

negative coefficient of regression predicts that under lower temperature conditions, *AhgSIG5*

transcript abundance will increase. Therefore, the lower temperatures of the spring sampling

season compared with the autumn sampling season (Fig. 1A, B) might explain the greater

levels of *AhgSIG5* transcript accumulation during the spring.

Because we considered *AhgpsbD* BLRP to represent the ultimate output from the signalling

pathway (Fig. 1A), our analysis suggests that environmental inputs occurred within at least

three positions in the pathway; first, in the regulation of *AhgCCA1* transcript accumulation by

the season or temperature, second, in the regulation of *AhgSIG5* transcript accumulation by

temperature, and, a third environmental input occurring downstream of *AhgSIG5* transcript

accumulation for the regulation of *AhgpsbD* BLRP. This is evidenced by the differences in

temperature responses of *AhgCCA1* (positive relationship), *AhgSIG5* (negative relationship),

and *AhgpsbD* BLRP (positive relationship) (Fig. 3C, G, L). These environmental inputs might

occur through biologically independent processes, such as temperature inputs to the

circadian clock mediated by temperature-responsive components such as the evening

complex [48, 49]. One of these mechanisms could be the regulation of *AhgSIG5* by HY5,

which is a known regulator of SIG5 that participates in low-temperature gene regulation [22,

25, 50] and binds the SIG5 promoter in *A. thaliana* [51]. Furthermore, there might be direct

effects of light upon sigma factor activity in chloroplasts through, for example, redox

regulation [52] or light- and temperature-regulation of chloroplast protein import. We did not

consider here the long history of light or temperature conditions upon leaves prior to

experimentation [53], or other biotic or abiotic factors such as water availability, relative

humidity, and atmospheric CO_2 concentration.

Statistical modelling of the transcriptome of field-grown *Oryza sativa* (rice) concluded that

the main environmental driver of *OsSIG5* (*Os05g0586600*) transcript accumulation is

temperature [9]. In this case, the temperature had a negative regression coefficient with

OsSIG5 [9]. This is consistent with our finding of a negative coefficient of regression

between temperature and *AhgSIG5*. The study of the rice transcriptome in the field [9] did

not monitor chloroplast-encoded transcripts, so a direct comparison between *AhgpsbD*

BLRP and our data is not possible.

A key finding from our work is the detection of a 24 h fluctuation in the coefficient of

regression between the three genes and temperature (Fig. 3C, G, L). One interpretation of

this is that there is a diel cycle of sensitivity of these pathway components to temperature

cues, with their response to temperature restricted to certain times of day. This is

reminiscent of circadian gating, which is the phenomenon whereby the circadian oscillator

restricts the response to a stimulus to certain times of day [39]. These findings are

corroborated by laboratory experiments, which demonstrate that the circadian oscillator

gates its own response to temperature [54], and the response of *AhgSIG5* to blue light is

gated by the circadian oscillator [20]. This is important, because it suggests that processes

of circadian gating might operate under naturally fluctuating conditions to modulate the

environmental responses of plants.

Our investigation provides insights into molecular aspects of signal transduction in plants

under field conditions. This represents a relatively under-studied topic, and we developed

new quantitative approaches to interpret transcript data collected under complex fluctuating

environments, to investigate a specific pathway. This allowed us to identify potential time-

delay steps within a signalling mechanism, multiple positions of environmental inputs, and

temporal gating of a response to temperature. The approaches used provide a framework to

study environmental signal integration in plants and other organisms under field conditions,

which might be valuable to understanding rhythmic biological responses within an

increasingly unpredictable climate.

Materials and Methods

Field site and plant material

Our experiments used a naturally-occurring population of *Arabidopsis halleri* subsp.

gemmifera (Matsum.) growing beside a forested stream in Hyogo Prefecture, Japan

(Omoide-gawa site; 35°06' N, 134°55' E, elevation 190–230 m) [12, 37, 55] (Fig. 1B, C). We

selected *A. halleri* as an experimental model for several reasons [56]. First, it has a high

nucleotide sequence identity and good synteny with *A. thaliana* [57]. Second, unlike *A.*

thaliana, the perennial life history of *A. halleri* allows investigation of transcriptional

responses across the seasons [2, 55]. Many individuals are clones because the species

propagates by producing clonal rosettes as well as by seeds, which allows repeated

sampling from single genotypes. Furthermore, *A. halleri* is metal tolerant and occurs in

natural habitats that are relatively free from other vegetation due to contamination by heavy

metals, which provides experimentally-convenient sites enriched with many *A. halleri* plants

[58]. *Arabidopsis halleri* subsp. *gemmifera* at this site was previously identified by

examination of museum and herbarium specimens, and a nearby population provided

material for sequencing the *A. halleri* genome [55, 57]. Sampling occurred during 24 – 26

March 2015, 15 – 17 September 2015, and 13 - 14 September 2016, where March and

September correspond to spring and autumn (fall) at the field site, respectively.

The *A. halleri* homologs of the *A. thaliana* genes *CCA1* and *SIG5* are loci *g25274* (*AhgSIG5*)

and *g097040* (*AhgCCA1*), respectively. These were identified from *A. halleri* genome

Version Ahal2.2 [57]. *AhgSIG5* has 94.9% coding sequence identity and 95.0% protein

sequence identity with the *A. thaliana* homolog. *AhgCCA1* has 94.8% coding sequence

identity and 93.3% protein sequence identity with the *A. thaliana* homolog. Chloroplast-

encoded *psbDC* is not annotated within Version Ahal2.2 of the *A. halleri* genome, and we

identified this instead within scaffold 2 of an *A. halleri* reference transcriptome [59]. The

AhgpsbD BLRP promoter region, which was our focus, has a 100% sequence identity with

psbD BLRP of *A. thaliana* [41].

Sampling under natural and manipulated conditions

The first sets of samples were obtained under natural conditions without environmental

manipulation. For this, we sampled during two different seasons, March 2015 and

September 2015, on dates that were close to the spring and autumn equinox. We exploited

variations in environmental conditions across the field site, and sampled leaves from the

locations nominated as "sun" and "shade" sites. At "sun" locations, plants received direct

sunlight during the day, and at "shade" locations plants received sunlight filtered by

surrounding vegetation for most of the day with the sites identified by measurement of the

ratio of red to far red light (Fig. S3; R:FR calculated as the photon irradiance from 660 to

670nm divided by the photon irradiance from 725 to 735nm [60]). In each case, sampling

occurred for at least 24 h. During March 2015, plants received more direct sunlight, whereas

during September 2015 the light was scattered through sky overcast with clouds.

We expanded the range of environmental conditions by manipulating the temperature

conditions around patches of plants (September 2016). In addition to control plants that were

not manipulated (Fig. S5A), we applied two temperature treatments. These were (1) a

continuous temperature increase (Fig. S5B), whereby plants were covered with clear plastic

horticultural domes to block air currents and trap warm air; (2) a continuous temperature

reduction, using a custom device that passed air through a duct within a heat-exchanging

ice-filled polystyrene box and expelled the chilled air into a clear horticultural dome covering

the plants, with chilling augmented by small ice packs within the dome (Fig. S5C).

Field sampling for transcript analysis

Across all experimental conditions, the same sampling and RNA isolation procedures were

used. At 2 h intervals, a fully expanded rosette leaf was excised with dissecting scissors

from 6 replicate plants for each condition. The time-courses using naturally occurring sun

and shade conditions each comprised 13 sampling timepoints over a total of 26 hours, and

the time-courses involving artificial temperature manipulations comprised 15 sampling

timepoints over a total of 30 hours. Sampled leaves were placed immediately into individual

microtubes containing at least 400 µL RNALater (Invitrogen). Scissors and forceps were

cleaned with 70% (w/v) ethanol between samples. After sampling, tubes were placed

temporarily on dry ice for up to 2 hours, at -40 °C for 3 days in a portable freezer during

transfer to the laboratory, and then at -80 °C until RNA isolation. During hours of darkness,

sampling occurred using green-filtered head torches. Each sampling timepoint was from the

same set of replicate plants. We obtained a separate reference standard for all RT-qPCR

experiments in the study, by pooling RNA from 10 leaves sampled at midday during March

2015 from healthy plants located randomly across the study site. This provided a reference

cDNA sample against which all RT-qPCR analyses from all sampling seasons were

normalized, to allow comparability between all datasets. This reference RNA sample was

collected during March 2015. In all experiments, dawn and dusk were defined as the

astronomical (solar) time of sunrise and sunset.

RNA isolation and RT-qPCR

Frozen samples containing RNALater were defrosted in a cold room for 4 hours, the

RNALater was removed, and leaf tissue was transferred to new dry tubes and frozen in

liquid nitrogen. Frozen tissue was ground with a TissueLyzer and total RNA was isolated

from the powdered plant material using Macherey-Nagel Nucleospin II RNA extraction kits

(Thermo-Fisher). cDNA was synthesized using a High Capacity cDNA Reverse Transcription

Kit (Applied Biosystems) supplemented with RNAase inhibitor, as described previously [19,

20]. RNA concentrations were determined using a Nanodrop spectrophotometer (Thermo

Scientific). cDNA was synthesized using an ABI High Capacity cDNA Reverse Transcription

Kit (Applied Biosystems) according to the manufacturer's instructions, using random primers

for the cDNA synthesis reaction. 1:500 cDNA dilutions were analysed using Brilliant III Ultra-

Fast SYBR Green QPCR master mix (Agilent Technologies) and required primer pairs

(Table S2). Primers were designed using the PrimerQuest™ Tool from Integrated DNA

Technologies. Results were normalized using the ΔΔCt method to *AhgACTIN2* [19, 20].

AhgACTIN2 is encoded in *A. halleri* by locus *g21632* [57] and has 97.8% coding sequence

identity with *A. thaliana ACTIN2* (*At3g18780*). Statistical comparisons within transcript

abundance data were conducted using the SPSS software package.

Environmental monitoring

The temperature and irradiance were measured beside the plants during sampling. The

temperature at each location, for each environmental manipulation, was monitored with EL-

USB-2 data loggers (Lascar Electronics) at 5-minute intervals. Temperature loggers were

wrapped in aluminium foil to prevent surface heating by solar radiation. Irradiance was

measured using a CC-3-UV-S cosine corrector connected to a USB2000+ spectrometer with

a QP400-2-UV-VIS fibre optic cable (Ocean Optics). Ambient light spectra (200 nm to

900 nm) were collected every 5 minutes over the 14 hours of light during each day of

sampling using OceanView software (Ocean Optics) on a laptop PC, controlled by a custom

script. The spectrometer and computer were powered using portable lithium battery packs

(Powertraveller, Hampshire, UK).

Smooth trend model analysis

The smooth trend model (STM) to analyze the difference in transcript abundance between

March and September under sun and shade conditions (Fig. 1) was defined by the

equations:

$$\mu_{1,t} \sim Normal(2\mu_{1,t-1} - \mu_{1,t-2},\ \sigma_{\mu1}^2), \tag{1}$$

$$\delta_t \sim Cauchy(\delta_{t-1},\ \sigma_\delta^2), \tag{2}$$

$$\mu_{2,t} = \mu_{1,t} + \delta_t, \tag{3}$$

$$y_{1,t} \sim Normal(\mu_{1,t},\ \sigma_y^2), \tag{4}$$

$$y_{2,t} \sim Normal(\mu_{2,t},\ \sigma_y^2), \tag{5}$$

where μ_1 and μ_2 are the smooth trend components in March and September in 2015,

respectively, δ_t is the time-varying difference between the two seasons, and y_1 and y_2 are

the observed transcript abundance in the two seasons. $t = (1, 2, \cdots, 13)$ is the time point at

two-hour intervals. The same STM was used to analyze the difference in transcript

abundance between sun and shade conditions in March and September (Fig. S1).

The parameters of the models were estimated by Bayesian inference. The statistical models

were written in the Stan language and the programs were called by the rstan package (using

version 2.21.0 of R). After 2,000 warm-up steps, 1,000 Markov Chain Monte Carlo (MCMC)

samples were obtained by thinning out 6,000 MCMC samples for each of four parallel

chains. Thus, 4,000 MCMC samples were obtained in total.

For the models of the three (ambient, warm and chill) conditions in the local environment

manipulation experiment, additional δ, μ and y were considered:

$$\delta_{2,t} \sim Cauchy(\delta_{2,t-1},\ \sigma_{\delta 2}^2), \tag{6}$$

$$\mu_{3,t} = \mu_{1,t} + \delta_{2,t}, \tag{7}$$

$$y_{3,t} \sim Normal(\mu_{3,t},\ \sigma_y^2). \tag{8}$$

463 *Dynamic linear model*

464 The dynamic linear model (DLM) to analyze the time-varying effect of environmental

465 variables on transcript abundance (Fig. 2) was defined by the equations:

$$\mu_t \sim Normal(\mu_{t-1},\ \sigma_\mu^2), \tag{9}$$

$$\beta_{temp,t} \sim Normal(\beta_{temp,t-1},\ \sigma_{\beta temp}^2), \tag{10}$$

$$\beta_{light,t} \sim Normal(\beta_{light,t-1},\ \sigma_{\beta light}^2), \tag{11}$$

$$\alpha_{MarSun,t} = \mu_t + \beta_{temp,t} \cdot temp_{MarSun,t} + \beta_{light,t} \cdot light_{MarSun,t}, \tag{12}$$

$$\alpha_{MarShade,t} = \mu_t + \beta_{temp,t} \cdot temp_{MarShade,t} + \beta_{light,t} \cdot light_{MarShade,t}, \tag{13}$$

$$\alpha_{SepSun,t} = \mu_t + \beta_{temp,t} \cdot temp_{SepSun,t} + \beta_{light,t} \cdot light_{SepSun,t}, \tag{14}$$

$$\alpha_{SepShade,t} = \mu_t + \beta_{temp,t} \cdot temp_{SepShade,t} + \beta_{light,t} \cdot light_{SepShade,t}, \tag{15}$$

$$y_{MarSun,t} \sim Normal(\alpha_{MarSun,t},\ \sigma_y^2), \tag{16}$$

$$y_{MarShade,t} \sim Normal(\alpha_{MarShade,t},\ \sigma_y^2), \tag{17}$$

$$y_{SepSun,t} \sim Normal(\alpha_{SepSun,t},\ \sigma_y^2), \tag{18}$$

$$y_{SepShade,t} \sim Normal(\alpha_{SepShade,t},\ \sigma_y^2), \tag{19}$$

where μ is the trend component, β is the time-varying regression coefficient, α is the true

state of transcript abundance, y is the observed transcript abundance, and σ^2 is the

variance. The subscripts, $temp, light, Mar, Sep, Sun$ and $Shade$ represent temperature,

irradiance, March, September, sun condition and shade condition, respectively. $t =$

$(1, 2, \cdots, 13)$ is the time point at two-hour intervals.

In the *AhgSIG5* and *AhgpsbD* BLRP models, the effects of the upstream genes (i.e.,

AhgCCA1 in the *AhgSIG5* model and *AhgSIG5* in the *AhgpsbD* BLRP model) were

additionally considered. Thus, the equations of α are modified as follows:

$$\alpha_{MarSun,t} = \mu_t + \beta_{temp,t} \cdot temp_{MarSun,t} + \beta_{light,t} \cdot light_{MarSun,t} + \beta_{gene,t} \cdot \tag{20}$$

$$gene_{MarSun,t},$$

$$\alpha_{MarShade,t} = \mu_t + \beta_{temp,t} \cdot temp_{MarShade,t} + \beta_{light,t} \cdot light_{MarShade,t} + \beta_{gene,t} \cdot \tag{21}$$

$$gene_{MarShade,t},$$

$$\alpha_{SepSun,t} = \mu_t + \beta_{temp,t} \cdot temp_{SepSun,t} + \beta_{light,t} \cdot light_{SepSun,t} + \beta_{gene,t} \cdot \tag{22}$$

$$gene_{SepSun,t},$$

$$\alpha_{SepShade,t} = \mu_t + \beta_{temp,t} \cdot temp_{SepShade,t} + \beta_{light,t} \cdot light_{SepShade,t} + \beta_{gene,t} \cdot \tag{23}$$

$$gene_{SepShade,t},$$

where $gene$ is the mean transcript abundance of the upstream genes, and the other symbols

are the same as above. The lagged effects of the upstream genes were tested by using

values at previous time points (e.g., using $gene_{MarSun,t-1}$, $gene_{MarSun,t-2}$, $gene_{MarSun,t-3}$ or

$gene_{MarSun,t-4}$ for $\alpha_{MarSun,t}$). The same DLM was used in Fig. 2 and S4.

The parameters of the models were estimated by Bayesian inference. The statistical models

were written in the Stan language and the programs were compiled using CmdStan (version

2.24). To operate CmdStan, the cmdstanr package (version 0.4.0) of R was used. After

3,000 warm-up steps, 1,000 MCMC samples were obtained for each of the four parallel

chains, and thus 4,000 MCMC samples were obtained in total.

Acknowledgements

We thank Noriane M. L. Simon and Tasuku Ito for experimental assistance, and Paige E.

Panter and Deirdre A. Lynch for constructive feedback. This research was funded by

BBSRC (UK) (BB/I005811/2; BB/J014400/1; Institute Strategic Programme GEN

BB/P013511/1), The Royal Society (IE140501), The Leverhulme Trust (RPG-2018-216), the

Japan Society for Promotion of Science (JSPS KAKENHI, JP21H04977, JP21H05659,

JP21K15164), the Japan Science and Technology Agency (JST, CREST no. JPMJCR1501)

and the Bristol Centre for Agricultural Innovation. DLCR is grateful to Consejo Nacional de

Ciencia y Tecnología (Mexico) for awarding a scholarship. This research was conducted

through the Joint Usage programme of the Center for Ecological Research of Kyoto

University.

Author contributions

DLCR, HN, JS, MNH, HK and AND designed and performed experimentation; HN, DLCR,

TM, LLBD, HK and AND analysed and interpreted data, and DLCR, HN, LLBD, HK and AND

wrote the paper.

Competing interests

The authors declare no competing interests

Figure legends

Fig. 1. Components of a circadian signalling pathway have diel fluctuations in a natural plant

population. (A) Potential architecture of a signal transduction pathway underlying SIG5-

mediated signalling to chloroplasts, with environmental inputs occurring at several positions.

t1 and *t2* represent the time taken for signal transduction between each pathway component.

(B-E) Diel fluctuations in (B, C) ambient temperature and (D, E) total irradiance detected

(200-900 nm), measured at 5-minute intervals. (F-K) Bayesian estimation of smooth trend

model (STM) for March and September 2015. The output of STM for (F, G) *AhgCCA1*, (H, I)

AhgSIG5 and (J, K) *AhgpsbD BLRP*. In F-K, the upper graphs show the predicted relative

transcript abundance for March (pink) and September (brown) with the mean of observed

values (dots), and the lower graphs represent the differences in transcript abundance

between March and September. The solid line and the shaded region are the median and

the 95% credible interval of the posterior distribution. When the 95% credible interval of the

difference between March and September does not contain zero, the difference is

considered significant.

Fig. 2. Time-delay steps are predicted within this signalling pathway under naturally-

fluctuating conditions. Lagged effects of variables in Bayesian estimation of dynamic linear

models (DLM) for transcript levels during March and September 2015. (A, B, C) RMSE, log-

likelihood and correlation of the models to predict *AhgSIG5* against the observed values,

incorporating time lags of the upstream *AhgCCA1*. (D, E, F) RMSE, log-likelihood and

correlation of the models to predict *AhgpsbD BLRP* against the observed values,

incorporating time lags of the upstream *AhgSIG5*. The time lags of temperature and

irradiance are set to 0. Asterisks represent (A, D) the lowest RMSE, (B, E) the highest log-

likelihood and (C, F) the highest correlation. Error bars represent the 95% Bayesian credible

intervals.

Fig. 3. The circadian clock and ambient temperature are key regulators of SIG5-mediated

signalling to chloroplasts under naturally-fluctuating conditions. Bayesian estimation of the

dynamic linear model (DLM) for March and September 2015. (A-D) The output of DLM for

AhgCCA1 where relative transcript abundance for (A) sun condition and (B) shade condition,

with the coefficient of regression for (C) temperature and (D) irradiance. (E-I) The output of

DLM for *AhgSIG5*, where (I) the coefficient of regression for *AhgCCA1* is shown, with other

plots the same as (A-D). (J-N) The output of DLM for *AhgpsbD BLRP*, where (N) is the

coefficient of regression for *AhgSIG5* is shown, with other plots the same as (A-D). The

predicted relative transcript abundance for March (orange) and September (blue) are shown

with the mean of observed values (dots). In each graph, the solid line and the shaded region

are the median and the 95% credible interval of the posterior distribution.

Fig. 4. Prediction of diel rhythms of gating of temperature response in a natural plant

population. Bayesian estimation of smooth trend model (STM) for temperature manipulation

experiments in September 2016. (A) Temperature changes during the study period in each

condition. (B-D) The output of STM for (B) *AhgCCA1*, (C) *AhgSIG5* and (D) *AhgpsbD BLRP*.

In each panel, the upper graphs show the predicted relative transcript abundance for

ambient (black), warm (red) and cool (light blue) conditions with the mean of observed

values (dots), and the lower graphs represent the differences in transcript abundance

against the ambient condition. In each graph, the solid line and the shaded region are the

median and the 95% credible interval of the posterior distribution. When the 95% credible

interval of the difference between conditions does not contain zero, the difference is

considered significant.

Fig. S1. Close relationship between *AtCCA1* and *AtSIG5* transcript abundance under free-

running conditions in *A. thaliana* under controlled conditions. (A-D) Relationship between

AtCCA1 and *AtSIG5* transcript abundance under conditions of constant light, from the

transcriptome studies of (A) [3] (B) [53], (C) [4], (D) [6]. (E, F) Relationship between *AtCCA1*

and *AtSIG5* transcript abundance under light/dark cycles with (E) long and (F) short

photoperiods, from the transcriptome study of [54]. Blue lines indicate a regression line.

Pearson's correlation coefficient (R) with p-values testing for the likelihood of a chance

correlation are shown for each plot.

Fig. S2. Components of a circadian signalling pathway have diel fluctuations in a natural

plant population. Bayesian estimation of smooth trend model (STM) comparing sun and

shade conditions, sampled during 2015. (A-D) Diel fluctuations in total irradiance detected

(A, B; 200-900 nm) and (C, D) ambient temperature, measured at 5-minute intervals. (E-J)

The output of STM for (E, F) *AhgCCA1*, (G, H) *AhgSIG5* and (I, J) *AhgpsbD* BLRP. In E-J,

the upper graphs show the predicted relative transcript abundance for sun (orange) and

shade (light grey) conditions, with the mean of observed values (dots). The lower graphs

represent the differences in transcript abundance between sun and shade conditions. The

solid line and the shaded region are the median and the 95% credible interval of the

posterior distribution. When the 95% credible interval of the difference between sun and

shade conditions does not contain zero, the difference is considered significant.

Fig. S3. Effects of the ratio of red to far-red light upon *A. halleri* plants in the field, during

March and September sampling seasons. (A, B) Examples of rosette-stage plants growing

under (A) sun and (B) shade conditions during the September sampling season. (C, D)

Comparison of the ratio of red to far-red light received by plants under the sun- and shade

conditions during (C) March 2015 and (D) September 2015 sampling seasons. The R:FR

varied during the photoperiod during both sampling seasons, and the effect of shade on

R:FR was ameliorated by heavy cloud cover.

Fig. S4. The nature of the time-delay steps within this signalling pathway depends on the

sampling season. Lagged effects of variables in Bayesian estimation of the dynamic linear

model (DLM) for March and September separately in 2015. (A, D, G, J) RMSE, (B, E, H, K)

log-likelihood and (C, F, I, L) correlation of the models to predict *AhgSIG5* in March against

the observed values, incorporating time lags of *AhgCCA1*. (D-F) RMSE, log-likelihood and

correlation of the models to predict *AhgpsbD* BLRP in March against the observed values,

incorporating time lags of *AhgSIG5*. (G-I) RMSE, log-likelihood and correlation of the models

to predict *AhgSIG5* in September against the observed values, incorporating time lags of

AhgCCA1. (J-L) RMSE, log-likelihood and correlation of the models to predict *AhgpsbD*

BLRP in September against the observed values, incorporating time lags of *AhgSIG5*. Time

lags of temperature and irradiance are set to 0. Asterisks represent the lowest RMSE (A, D,

G, J), the highest log-likelihood (B, E, H, K) and the highest correlation (C, F, I, L). Error bars

represent the 95% Bayesian credible intervals.

Fig. S5. Moderate temperature manipulations to adjacent patches of *A. halleri* plants, in the

field, using custom-designed equipment. (A) The representative appearance of plant patches

under naturally fluctuating conditions. (B) Plants covered with a plastic dome to raise

temperature. (C) Plants covered with plastic dome undergoing temperature reduction with a

custom chilling device. In this device, cool air is introduced to enclosed plant patches after

being driven slowly through a heat exchanger, positioned within an expanded polystyrene

box filled with ice.

Fig. S6. Location of field sampling. Photographs of (A) upstream and (B) downstream views

of Omoide river site, which has naturally occurring populations of *A. halleri*. The majority of

plants at ground level on the stony river banks are *A. halleri*.

References

1.   Millar, A.J., *The intracellular dynamics of circadian clocks reach for the light of*
ecology and evolution. Annual Review of Plant Biology, 2016. **67**(1): p. 595-618.
2.   Kudoh, H., *Molecular phenology in plants: in natura systems biology for the*
comprehensive understanding of seasonal responses under natural environments.
New Phytologist, 2016. **210**(2): p. 399-412.
3.   Covington, M.F., et al., *Global transcriptome analysis reveals circadian regulation of*
key pathways in plant growth and development. Genome Biology, 2008. **9**(8): p.
R130.
4.   Edwards, K.D., et al., *FLOWERING LOCUS C mediates natural variation in the high-*
temperature response of the Arabidopsis circadian clock. The Plant Cell, 2006. **18**(3):
p. 639-650.
5.   Dodd, A.N., et al., *Plant circadian clocks increase photosynthesis, growth, survival,*
and competitive advantage. Science, 2005. **309**(5734): p. 630-633.
6.   Harmer, S.L., et al., *Orchestrated transcription of key pathways in Arabidopsis by the*
circadian clock. Science, 2000. **290**(5499): p. 2110-2113.
7.   Graf, A., et al., *Circadian control of carbohydrate availability for growth in Arabidopsis*
plants at night. Proceedings of the National Academy of Sciences, 2010. **107**(20): p.
9458-9463.
8.   Hotta, C.T., et al., *Modulation of environmental responses of plants by circadian*
clocks. Plant, Cell & Environment, 2007. **30**(3): p. 333-349.
9.   Nagano, Atsushi J., et al., *Deciphering and prediction of transcriptome dynamics*
under fluctuating field conditions. Cell, 2012. **151**(6): p. 1358-1369.
10.   Dantas, L.L.B., et al., *Alternative splicing of circadian clock genes correlates with*
temperature in field-grown sugarcane. Frontiers in Plant Science, 2019. **10**: p. article
1614.
11.   Dantas, L.L.B., et al., *Rhythms of transcription in field-grown sugarcane are highly*
organ specific. Scientific Reports, 2020. **10**(1): p. article 6565.
12.   Nagano, A.J., et al., *Annual transcriptome dynamics in natural environments reveals*
plant seasonal adaptation. Nature Plants, 2019. **5**(1): p. 74-83.
13.   Matsuzaki, J., Y. Kawahara, and T. Izawa, *Punctual transcriptional regulation by the*
rice circadian clock under fluctuating field conditions. The Plant Cell, 2015. **27**(3): p.
633-648.
14.   Annunziata, M.G., et al., *Response of Arabidopsis primary metabolism and circadian*
clock to low night temperature in a natural light environment. Journal of Experimental
Botany, 2018. **69**(20): p. 4881-4895.

15. Annunziata, M.G., et al., *Getting back to nature: a reality check for experiments in controlled environments.* Journal of Experimental Botany, 2017. **68**(16): p. 4463-4477.

16. Dantas, L.L.B., et al., *Field microenvironments regulate crop diel transcript and metabolite rhythms.* New Phytologist, 2021. **232**(4): p. 1738-1749.

17. Hashida, Y., et al., *Fillable and unfillable gaps in plant transcriptome under field and controlled environments.* Plant, Cell & Environment, 2022. **45**(8): p. 2410-2427.

18. Steed, G., et al., *Chronoculture, harnessing the circadian clock to improve crop yield and sustainability.* Science, 2021. **372**(6541): p. eabc9141.

19. Belbin, F.E., et al., *Integration of light and circadian signals that regulate chloroplast transcription by a nuclear-encoded sigma factor.* New Phytologist, 2017. **213**(2): p. 727-738.

20. Noordally, Z.B., et al., *Circadian control of chloroplast transcription by a nuclear-encoded timing signal.* Science, 2013. **339**(6125): p. 1316-1319.

21. Onda, Y., et al., *Light induction of Arabidopsis SIG1 and SIG5 transcripts in mature leaves: differential roles of cryptochrome 1 and cryptochrome 2 and dual function of SIG5 in the recognition of plastid promoters.* The Plant Journal, 2008. **55**(6): p. 968-978.

22. Nagashima, A., et al., *The multiple-stress responsive plastid sigma factor, SIG5, directs activation of the psbD blue light responsive promoter (BLRP) in Arabidopsis thaliana.* Plant and Cell Physiology, 2004. **45**(4): p. 357-368.

23. Yamburenko, M.V., Y.O. Zubo, and T. Börner, *Abscisic acid affects transcription of chloroplast genes via protein phosphatase 2C-dependent activation of nuclear genes: repression by guanosine-3'-5'-bisdiphosphate and activation by sigma factor 5.* The Plant Journal, 2015. **82**(6): p. 1030-1041.

24. Zhao, P., et al., *ATHB17 enhances stress tolerance by coordinating photosynthesis associated nuclear gene and ATSIG5 expression in response to abiotic stress.* Scientific Reports, 2017. **7**(1): p. article 45492.

25. Mellenthin, M., et al., *Expression of the Arabidopsis sigma factor SIG5 is photoreceptor and photosynthesis controlled.* Plants, 2014. **3**: p. 359-391.

26. Tsunoyama, Y., et al., *Blue light specific and differential expression of a plastid σ factor, Sig5 in Arabidopsis thaliana.* FEBS Letters, 2002. **516**(1-3): p. 225-228.

27. Mochizuki, T., et al., *Two independent light signals cooperate in the activation of the plastid psbD blue light-responsive promoter in Arabidopsis.* FEBS Letters, 2004. **571**(1): p. 26-30.

28. Møller, S.G., et al., *PP7 is a positive regulator of blue light signaling in Arabidopsis.* The Plant Cell, 2003. **15**(5): p. 1111-1119.

29. Davey, M.P., et al., *The UV-B photoreceptor UVR8 promotes photosynthetic efficiency in Arabidopsis thaliana exposed to elevated levels of UV-B.* Photosynthesis Research, 2012. **114**(2): p. 121-131.

30. Brown, B.A. and G.I. Jenkins, *UV-B signaling pathways with different fluence-rate response profiles are distinguished in mature Arabidopsis leaf tissue by requirement for UVR8, HY5, and HYH.* Plant Physiology, 2008. **146**(2): p. 576-588.

31. Tsunoyama, Y., et al., *Blue light-induced transcription of plastid-encoded psbD gene is mediated by a nuclear-encoded transcription initiation factor, AtSig5.* Proceedings of the National Academy of Sciences of the United States of America, 2004. **101**(9): p. 3304-3309.

32. Shimmura, S., et al., *Comparative analysis of chloroplast psbD promoters in terrestrial plants.* Frontiers in Plant Science, 2017. **8**: p. article 1186.

33. Wenden, B., et al., *Light inputs shape the Arabidopsis circadian system.* The Plant Journal, 2011. **66**(3): p. 480-491.

34. Fraser, D.P., et al., *Phytochrome A elevates plant circadian-clock components to suppress shade avoidance in deep-canopy shade.* Proceedings of the National Academy of Sciences, 2021. **118**(27): p. e2108176118.

35. Panter, P.E., et al., *Circadian regulation of the plant transcriptome under natural*
conditions. Frontiers in Genetics, 2019. **10**: p. article 1239.
36. Locke, J.C.W., A.J. Millar, and M.S. Turner, *Modelling genetic networks with noisy*
and varied experimental data: the circadian clock in Arabidopsis thaliana. Journal of
Theoretical Biology, 2005. **234**(3): p. 383-393.
37. Aikawa, S., et al., *Robust control of the seasonal expression of the Arabidopsis FLC*
gene in a fluctuating environment. Proceedings of the National Academy of Sciences,
2010. **107**(25): p. 11632-11637.
38. Nishio, H., et al., *Repressive chromatin modification underpins the long-term*
expression trend of a perennial flowering gene in nature. Nature Communications,
2020. **11**(1): p. article 2065.
39. Pittendrigh, C.S. and D. Skopik Steven, *Circadian Systems, V. The driving oscillation*
and the temporal sequence of development. Proceedings of the National Academy of
Sciences, 1970. **65**(3): p. 500-507.
40. Finster, S., et al., *Light-dependent, plastome-wide association of the plastid-encoded*
RNA polymerase with chloroplast DNA. Plant Journal, 2013. **76**(5): p. 849-860.
41. Hoffer, P.H. and D.A. Christopher, *Structure and blue-light-responsive transcription of*
a chloroplast psbD promoter from Arabidopsis thaliana. Plant Physiology, 1997.
115(1): p. 213-222.
42. Satoh, J., et al., *Developmental stage-specific multi-subunit plastid RNA polymerases*
(PEP) in wheat. Plant Journal, 1999. **18**(4): p. 407-415.
43. Baena-González, E., et al., *Chloroplast transcription at different light intensities.*
Glutathione-mediated phosphorylation of the major RNA polymerase involved in
redox-regulated organellar gene expression. Plant Physiology, 2001. **127**(3): p. 1044-
1052.
44. Nakamura, T., et al., *Array-based analysis on tobacco plastid transcripts: Preparation*
of a genomic microarray containing all genes and all intergenic regions. Plant and
Cell Physiology, 2003. **44**(8): p. 861-867.
45. Bieniawska, Z., et al., *Disruption of the Arabidopsis circadian clock is responsible for*
extensive variation in the cold-responsive transcriptome. Plant Physiology, 2008.
147(1): p. 263-279.
46. Gould, P.D., et al., *The molecular basis of temperature compensation in the*
Arabidopsis circadian clock. The Plant Cell, 2006. **18**(5): p. 1177-1187.
47. Gould, P.D., et al., *Network balance via CRY signalling controls the Arabidopsis*
circadian clock over ambient temperatures. Molecular Systems Biology, 2013. **9**(1):
p. article 650.
48. Mizuno, T., et al., *Ambient temperature signal feeds into the circadian clock*
transcriptional circuitry through the EC night-time repressor in Arabidopsis thaliana.
Plant and Cell Physiology, 2014. **55**(5): p. 958-976.
49. Thines, B. and G. Harmon Frank, *Ambient temperature response establishes ELF3*
as a required component of the core Arabidopsis circadian clock. Proceedings of the
National Academy of Sciences, 2010. **107**(7): p. 3257-3262.
50. Catalá, R., J. Medina, and J. Salinas, *Integration of low temperature and light*
signaling during cold acclimation response in Arabidopsis. Proceedings of the
National Academy of Sciences, 2011. **108**(39): p. 16475-16480.
51. Burko, Y., et al., *Chimeric activators and repressors define HY5 activity and reveal a*
light-regulated feedback mechanism. The Plant Cell, 2020. **32**(4): p. 967-983.
52. Shimizu, M., et al., *Sigma factor phosphorylation in the photosynthetic control of*
photosystem stoichiometry. Proceedings of the National Academy of Sciences, 2010.
107(23): p. 10760-10764.
53. Ling, Q. and P. Jarvis, *Regulation of chloroplast protein import by the ubiquitin E3*
ligase SP1 is important for stress tolerance in plants. Current Biology, 2015. **25**(19):
p. 2527-2534.

54.   Michael, T.P., P.A. Salomé, and C.R. McClung, *Two Arabidopsis circadian oscillators*
741   *can be distinguished by differential temperature sensitivity.* Proceedings of the
742   National Academy of Sciences, 2003. **100**(11): p. 6878-6883.
743   55.   Kudoh, H., et al., *The long-term "in natura" study sites of Arabidopsis halleri for plant*
744   *transcription and epigenetic modification analyses in natural environments*, in *Plant*
745   *Transcription Factors: Methods and Protocols*, N. Yamaguchi, Editor. 2018, Springer
746   New York: New York, NY. p. 41-57.
747   56.   Honjo, M.N. and H. Kudoh, *Arabidopsis halleri: a perennial model system for*
748   *studying population differentiation and local adaptation.* AoB PLANTS, 2019. **11**(6):
749   p. article plz076.
750   57.   Briskine, R.V., et al., *Genome assembly and annotation of Arabidopsis halleri, a*
751   *model for heavy metal hyperaccumulation and evolutionary ecology.* Molecular
752   Ecology Resources, 2017. **17**(5): p. 1025-1036.
753   58.   Kawagoe, T. and H. Kudoh, *Escape from floral herbivory by early flowering in*
754   *Arabidopsis halleri subsp. gemmifera.* Oecologia, 2010. **164**(3): p. 713-720.
755   59.   Kamitani, M., et al., *RNA-Seq reveals virus–virus and virus–plant interactions in*
756   *nature.* FEMS Microbiology Ecology, 2016. **92**(11): p. article fiw176.
757   60.   Franklin, K.A., *Shade avoidance.* New Phytologist, 2008. **179**(4): p. 930-944.

A

Nucleus — Chloroplast

CCA1 (proxy for circadian oscillator) —*t1*→ *SIG5* —*t2*→ *psbD* BLRP

Environmental cues (light, temperature, seasonality)

B Temperature (sun)

C Temperature (shade)

D Irradiance (sun)

E Irradiance (shade)

F *AhgCCA1* (sun)

G *AhgCCA1* (shade)

H *AhgSIG5* (sun)

I *AhgSIG5* (shade)

J *AhgpsbD* BLRP (sun)

K *AhgpsbD* BLRP (shade)

— March (Spring) — September (Autumn)