Determinants restricting ACE2 recognition of MERS-related coronaviruses in bats

Chengbao Ma[1*],Qing Xiong[1,*], Mengxue Gu[1], Chunli Wang[1], Lulu Shi[1], Chen Liu[1], Junyu Si[1], Fei

Tong[1], Peng Liu[1], Meiling Huang[1], Huan Yan[1✉]

[1]State Key Laboratory of Virology, Institute for Vaccine Research and Modern Virology Research

Center, College of Life Sciences, Wuhan University, Wuhan, Hubei, China.

*These authors contributed equally.

✉e-mail: huanyan@whu.edu.cn

Abstract

Phylogenetically distant coronaviruses have evolved to employ ACE2 as their common

receptors, including NL63 and many Severe acute respiratory syndrome (SARS) coronavirus-related

viruses[1]. Recently, we found two Middle East respiratory syndrome coronaviruses

(MERS-CoV)-related bat coronaviruses, NeoCoV and PDF-2180, also use Angiotensin-converting

enzyme 2(ACE2) but not MERS-CoV receptor dipeptidyl peptidase 4 (DPP4) for entry[2]. Receptor

binding domain (RBD)-binding and pseudovirus entry assays based on a wide range of bat ACE2

orthologs revealed that the two viruses strongly prefer ACE2 from *Yangochiropteran* bats as

compared with *Yinpterochiropteran* bats, which is not observed in NL63 and SARS-CoV-2. Genetic

and structural analyses of the virus-receptor interactions of 50 bat ACE2 orthologs pointed to four

crucial host range determinants in two viral binding loops on ACE2. Subsequent functional

verifications via mutagenesis on representative ACE2 orthologs confirmed the importance of these

determinants on human and bat cells. Remarkably, NeoCoV-T510F, a mutation previously shown to

acquire human ACE2 recognition, displayed an expanded potential host range covering most tested

bat ACE2, probably due to its reinforced interaction with an evolutionary conserved hydrophobic

pocket. Our results elucidated the molecular mechanisms for the species-specific ACE2 usage of

MERS-related viruses, offering basic information for assessing the zoonotic risk of these ACE2

utilizing merbecoviruses.

Keywords: Bat coronavirus, MERS-CoV, NeoCoV, ACE2, bats, host range determinants

Introduction

Emerging coronaviruses associated with several outbreaks in the 21st century impose severe threats to human health[3-6]. Being the most important reservoirs of at least hundreds of α- and β-coronaviruses (CoVs), bats are the crucial hosts of evolutional ancestors and relatives of many high-risk human CoVs, including SARS-CoV, SARS-CoV-2, and MERS-CoV [7-11]. Although the origin of highly pathogenic MERS-CoV remains elusive, hypotheses have been proposed that MERS-CoV may arise from the recombination and evolution of MERS-related bat coronaviruses[12-14]. NeoCoV and PDF-2180 are MERS-related viruses (merbecoviruses subgenus) sampled in vesper bats harbors in South Africa and Southwest Uganda, respectively [15,16]. NeoCoV represents the yet-identified closest relative of MERS-CoV in bats with ~85% whole genome nucleotide similarity[17]. However, its RBD sequence and structure are very different from MERS-CoV[2,15].

Several merbecoviruses use their host's DPP4 as entry receptors, such as MERS-CoV, bat CoV HKU4, and HKU25, whereas the receptors for many other merbecoviruses remain elusive[15,18-20]. Recently, we reported that NeoCoV and PDF-2180 unexpectedly engage bat ACE2 as their receptors [2]. Cryo-electron microscopy (Cryo-EM) analysis of NeoCoV or PDF-2180 RBD in complex with a bat ACE2 from *Pipistrellus pipistrellus* (Ppip) revealed a relatively small and flat ACE2 binding surface characterized by an N-glycosylation mediated protein-glycan interaction, a mode distinct from other ACE2 utilizing viruses.

Receptor recognition is usually species-specific and determines the cross-species transmission potential of CoVs. For example, SARS-CoV and SARS-CoV-2 exhibited varying efficiency to use ACE2 orthologs from different bat species[21]. NeoCoV and PDF-2180 highly preferred ACE2 orthologs from *Yangochiropteran* bats (Yang-bats) for entry[2]. Comparatively, human ACE2 (hACE2) is not efficiently recognized by these viruses, whereas adaptive spike mutations like T510F on receptor binding motif (RBM) enhanced hACE2 binding affinity. Interestingly, most tested ACE2 orthologs from Yin-bats were even more defective in supporting viral binding and entry of the two viruses than wild-type (WT) hACE2, indicative of specific host range determinants among bat species[2].

Here, we extensively explored the determinants restricting ACE2 orthologs from Yin-bats of supporting NeoCoV and PDF-2180 entry. Four species-specific sites within the binding interface were identified and confirmed as potential host range determinants, including two N-glycosylation sites.

Results

Bat-specific receptor usage of ACE2 utilizing viruses

We previously showed that ACE2 orthologs from most tested (14/15) Yin-bats and very few (2/30) Yang-bats (Nlep and Nhum) were deficient in supporting NeoCoV and PDF-2180 RBD binding and pseudovirus entry[2]. In this study, we re-examined the receptor function of 18 Yin-Bat's ACE2 for NeoCoV, including three (Rcor, Raff, and Rmal) we subsequently found capable of supporting NeoCoV and PDF-2180 RBD entry. ACE2 orthologs from four Yang-bats (Nlep, Ppip, Lbor, and Nhum) and hACE2 were also included. Immunofluorescence assay detecting the C-terminal fused 3×Flag tags indicated these receptors were expressed at a similar level (**Extended data figure 1**). The results of the NeoCoV RBD binding and pseudovirus entry assays were generally consistent and further confirmed our previous observation that most, but not all, Yin-bats' ACE2 are deficient in supporting NeoCoV entry (**Fig.1a-b**). We next investigated whether the species-specific bat ACE2 preference can be observed in other representative ACE2 utilizing viruses: PDF-2180, SARS-CoV-2, and NL63. The receptor binding modes of the four viruses are shown in **Fig.1c**, highlighted with distinct RBD binding footprints. NeoCoV and PDF-2180 RBD binding and pseudovirus entry exhibited very similar profiles among the 22 tested bat ACE2 orthologs and are very different from those of SARS-CoV-2 and NL63, consistent with the distinct receptor binding modes of these viruses (**Fig.1d and extended data figure 1 and 2**). SARS-CoV-2, which is supposed to share common ancestors infecting Yin-bats (*Rhinolophus malayanus*, Rmal)[9], can efficiently use 12 ACE2 from Yin-bats. Although the close relatives of NL63 were sampled in *Yang*-bat (*Perimyotis subflavus*)[22], efficient binding and entry of NL63 can also be supported by most (14/18) tested Yin-bats' ACE2. These data indicate that the species-specificity is not strictly correlated with the phylogeny, and additional determinants are affecting the functionality of ACE2 from various bats in mediating NeoCoV and PDF-2180 infection.

Identification of potential host range determinants restricting NeoCoV/PDF-2180 recognition

We next sought to identify the host range determinants by combining genetic, structural, and functional analyses focusing on sequences of the two critical RBD interacting loops, both of which contain an important glycosylation site **(Fig.2a)**. We first conducted multiple sequence alignment and conservation analyses based on 50 bat ACE2 sequences (from 19 Yin- and 31 Yang-bats), which are separated into two groups by their ability to support NeoCoV entry **(Fig.2b and Extended data 3 and 4)**. These results highlighted four candidate sites (from A-D) with contrasting residue frequency across the two groups **(Fig.2c)**. According to the presence of deficient/sub-optional residues, we classified the ACE2 orthologs deficient in viral receptor functions into several corresponding defect types **(Fig.2d).** For example, Raeg ACE2 was designated defect type ABCD as it carries deficient residues on all four predicted sites. We next analyzed the impact of the deficient/sub-optimal ACE2 residues on viral RBD recognition via structural modeling based on the Ppip ACE2-NeoCoV-RBD complex **(Fig.2d)**. Glycosylations sites (N-X-T/S) in A and C involved in in the glycan-protein interactions are missing in several Yin-bats' ACE2. Site A glycosylation (N54 glycosylation) seems more important than site C glycosylation (N330-glycosylation) as it mediated a protein-glycan interaction required for stabilizing RBD in Region 1. In contrast, site C glycosylation only partially contributes to the extensive interactions in region 2 via forming hydrogen bonds with two RBM glycines of NeoCoV but not PDF-2180. Residues in sites B and D of Yin-bats' ACE2 either abolish the polar contacts (e.g., two salt bridges formed by residues E305 and D338, respectively) or introduce steric clashes (e.g., E305K) that reduce the binding affinity. It is worth mentioning that site D, especially the D338, is a critical host range determinant restricting hACE2 from efficiently supporting the binding and entry of the two viruses[2].

Functional verification of host range determinants of NeoCoV/PDF2180 in bat ACE2

To explore the role of candidate determinants, we picked representative ACE2 orthologs of specific defect types for gain-of-function tests. Specifically, a series of point or swap mutagenesis based on ACE2 orthologs from Rsin (type A), Rfer and Rpea (type B), Hgal (type BC), Nlep (type BCD), and Raeg (type ABCD) were conducted to test their positive effects in supporting NeoCoV/PDF2180 RBD binding and entry. Rsin-N55T (type A) with an amended N53 glycosylation

site (corresponding to N54 in Ppip ACE2) showed remarkable improvements in supporting RBD binding and viral entry (**Fig.3a-b and Extended data figure 5**). Rfer-K305E (type B) and Rpea-N305E (type B*), with introduced salt bridges, also exhibited significantly improved receptor functions (**Fig.3c-d and Extended data figure 5 and 6**). Efficient binding and entry supported by bat ACE2 mutants from Hgal (type BC) and Nlep (type BCD) can be observed after corresponding residues are substituted with Ppip ACE2 counterparts (**Extended data figure 6**). Particularity, a gradual gain of receptor function of Raeg ACE2 (type ABCD) can be observed by stepwise increasing substitutions in sites A, AB, ABC, and ABCD, respectively (**Fig.3e-g**). Similar gain-of-function of Rsin-N55T and Raeg ACE2-ABCD   was also confirmed on a Bat cell line Tb 1 Lu (**Extended data figure 7**). We further evaluated the binding affinity between viral RBDs proteins and representative WT and mutant ACE2 by Flow cytometry and Bio-layer interferometry (BLI) assays (**Fig.3h-i**). As expected, the binding affinities between WT Raeg/Rsin ACE2 and NeoCoV/PDF-2180 RBDs were below the detection limit, but the related mutations significantly improved their RBD binding affinities, with apparent binding affinities ($K_{D, app}$) ranging from 2.04×10^{-8} to 6.94×10^{-10} (**Fig.3i**). Together, these results demonstrated that all the tested bat ACE2 orthologs with poor receptor function for NeoCoV and PDF-2180 could be primarily attributed to the deficient/sub-optional residues in the four predicted host range determinants.

RBM mutations broaden the potential host range of NeoCoV

We previously reported that specific mutations of NeoCoV and PDF-2180 RBM confer hACE2 recognition[2]. For NeoCoV, a T510F mutation increasing RBM local hydrophobicity enhanced its interaction with a hydrophobic pocket of ACE2 (**Fig.4a**). Sequence analysis of the 50 bat ACE2 indicated residues constituting this hydrophobic pocket are highly conserved among bat species (**Fig.4b**). Thus, we hypothesized that mutations such as T510F could expand the potential host range by recognizing Yin-bats via increasing hydrophobic interactions with this pocket. As expected, NeoCoV-T510F efficiently binds with most tested ACE2 orthologs and can achieve robust entry by these receptors, except for Rsin and Raeg ACE2 without the critical N53 glycosylations (**Fig.4c-d**). Further experiments demonstrated that Rsin ACE2-N55T and Raeg ACE2-I55T with functional N53 glycosylation sites achieved efficient NeoCoV-T510F entry (**Fig.4e-f**). These results indicate that NeoCoV, PDF-2180, or related viruses may efficiently infect Yin-bats through antigenic

drifts on RBM and further highlight the importance of N53 glycan in species-specific ACE2

recognition.

Discussion

The order Chiroptera comprises more than 1,400 bat species with remarkable diversity and wide

geographic distribution[23]. These bats are considered natural reservoirs of hundreds of known α- and

β-CoVs and important hosts for viral evolution[8]. Critical residues on viral receptors responsible for

spike protein binding exhibit accelerated evolution in bats[24,25], resulting in species-specificity in

supporting coronavirus binding and entry, which has been reported in SARS-CoV, SARS-CoV-2,

and MERS-CoV[21,26]. To date, coronaviruses of three different subgenera have evolved to engage

ACE2 for cellular entry, including NL63 (Setracovirus subgenus, α-CoV), many SARS-related CoVs

(Sarbecovirus subgenus, β-CoV), and two recently reported MERS-related CoVs (Merbecovirus

subgenus, β-CoV) in this study [1,2]. The distinct viral RBD structure and ACE2 binding footprints

suggest convergent evolutionary histories of receptor adaptation of these viruses[2].

In this study, we demonstrated that NeoCoV and PDF-2180 prefer ACE2 orthologs from

Yang-bats but not from most Yin-bats, while NL63 and SARS-CoV-2 displayed very different usage

profiles. This receptor preference is in line with the observation that most Merbecoviruses were

sampled in bats belonging to the family *Vespertilionidae* (vesper bats), the largest family of

Yangochiroptera (**Extended data figure 3**), including the hosts of NeoCoV and PDF-2180[8].

Comparatively, most sarbecoviruses were identified in Yin-bats, particularly *Rhinolophine* or

Hipposideros[8,27]. The differences in host preference likely limited the opportunities for cross-lineage

recombination of these high-risk viruses.

We here revealed that the receptor preference is determined by specific sequences in the viral

binding interface of ACE2 orthologs. Interestingly, glycan-protein interactions play a vital role in

species-specific ACE2 recognition of Merbecoviruses, especially the crucial interaction mediated by

N54-(or N53 in Rsin and Raeg) glycosylation. Similar glycan-protein interaction in receptor

engagement has also been reported in other coronaviruses, including SARS-CoV-2 and

human-infecting CCoV-HuPn-2018[28,29]. It could be interesting to investigate their contribution to

host range determination in other coronaviruses. The other three determinants were close to the N329

(or N330 in some species) glycosylation. Since only some Yang-bats carry glycosylation at this site,

its contribution to binding affinity and species specificity is less important than the N53

glycosylation, probably due to the compensation of nearby protein-protein interactions. Other than

the two glycosylation, site B and D determinants represent two critical sites that participated in

protein-protein interactions required for effective receptor recognition. Both sites of most Yang-bats'

ACE2 form salt bridges with viral RBMs. Although site D has been demonstrated to restrict hACE2

recognition, the interaction mediated by site B (residue 305) seems to play a more significant role in

host range determinants in bats. Most ACE2 from Yin-bats lack an E305 in site B, which forms a salt

bridge with the RBM of the viruses (e.g., K512 in NeoCoV) in most Yang-bat. N305 in several

Yin-bats forms sub-optional hydrogen bonds with the viruses, while K305 seems unable to interact

with the viruses and may even result in steric hindrance.

Receptor recognition is a primary barrier to cross-species transmission of coronaviruses, while

adaptive antigenic drift could break this barrier and lead to spillover in other species[30]. Here we show

that the T510F mutation in the NeoCoV spike, increasing binding affinity via interacting with a

conserved hydrophobic pocket, broadens the potential host range. Our results indicate that NeoCoV

and related viruses hold the potential to break the current host range barrier of ACE2 incompatibility

through antigenic drift or recombination. Moreover, these ACE2-using merbecoviruses may also

have more competent relatives in the wild that are not yet on our radar.

Taken together, our study elucidated the species-specific ACE2 recognizing mechanism of

MERS-related viruses and identified several critical residues on the binding interface for host range

determination. Our study adds knowledge to the long-neglected and potentially high-risk

coronaviruses, which is crucial for virus surveillance in various species to prepare for future

outbreaks.

Methods

Cell lines

HEK293T (CRL-3216) and the bat epithelial cell line Tb 1 Lu (CCL-88) were purchased from

American Type Culture Collection (ATCC). Cells were maintained by Dulbecco's Modified Eagle

Medium (DMEM, Monad, China) with 1% PS (Penicillin/Streptomycin) and 10% Foetal Bovine

Serum. An I1-Hybridoma (CRL-2700) cell line secreting a neutralizing mouse monoclonal antibody

targeting the VSV glycoprotein (VSVG) was cultured in Minimum Essential Medium(MEM) with

Earles's balances salts and 2.0 mM of L-glutamine (Gibico) and 10% FBS. All the cells were

cultured at 37°C with 5% CO_2 with regular passage every 2-3 days.

Plasmids

Plasmids expressing WT or mutated bats ACE2 orthologs were generated by inserting human

codon-optimized sequences with/without specific mutations into a lentiviral transfer vector

(pLVX-EF1a-Puro, Genewiz) with C-terminus 3×Flag (DYKDHD-G-DYKDHD-I-DYKDDDDK).

Human codon-optimized the spike sequences of NeoCoV (AGY29650.2), PDF-2180

(YP_009361857.1), SARS-CoV-2 (D614G) (YP_009724390.1), and NL63(APF29071.1) were

cloned into the PCAGGS vector with C terminal deletions (13-15 aa) to improve the assembly

efficiency. The plasmids expressing the recombinant CoVs RBD-hFc fusion proteins were

constructed by inserting NeoCoV RBD (380-585), PDF-2180 RBD (381-586), SARS-CoV-2 RBD

(331-524) and NL63 RBD (481-616) coding sequences into the pCAGGS vector containing an

N-terminal CD5 signal peptide (MPMGSLQPLATLYLLGMLVASVL) and a C-terminal hFc tag for

purification and detection. The expression plasmids of bats ACE2 ectodomain proteins were

generated by inserting WT or mutated sequences of Rousettus_aegyptiacus (18-740aa),

Rhinolophus_sinicus (18-740aa), and Pipistrellus_kuhlii (20-738 aa) into the pCAGGS vector with

an N-terminal CD5 signal peptide and a C-terminal twin-strep-3×Flag tag

(WSHPQFEKGGGSGGGSGGSAWSHPQFEKGGGRSDYKDHDGDYKDHDIDYKDDDDK).

Bioinformatic and structural analysis

Sequence alignments of different bats ACE2 were performed either by the MUSCLE algorithm by

MEGA-X (version 10.1.8) or ClustalW (https://www.genome.jp/tools-bin/clustalw) software. The

residue usage frequency (sequence logo) and mean hydrophobicity of 50 ACE2 sequences were

generated by the Geneious Prime software. Phylogenetic trees were produced using the maximal

likelihood method in MEGA-X (1000 Bootstraps) and polished with iTOL(v6) (https://itol.embl.de/)

[31]. The structural were shown by ChimeraX based on SARS-CoV-2 RBD & human ACE2(PDB:

6M0J), NL63 RBD & human ACE2(PDB: 3KBH), NeoCoV RBD & Ppip ACE2(PDB: 7WPO) and

PDF-2180 RBD & Ppip ACE2(PDB: 7WPZ). RBD binding footprints and interaction details were

analyzed and demonstrated using the UCSF ChimeraX[32]. Structural representatives of NeoCoV RBD interacting with WT or mutated Ppip ACE2 were analyzed using the UCSF Chimera X. The colored sequence conservation of 50 bats ACE2 was demonstrated by the UCSF ChimeraX based on multi-sequence alignments data generated by MEGA-X.

Protein expression and purification

HEK293T transfected with different protein-expressing plasmids through the GeneTwin reagent (Biomed, TG101-01). At 4-6 hours post-transfection, the medium of the transfected cells was replenished with the SMM 293-TII Expression Medium (Sino Biological, M293TII), and the protein-containing supernatant was harvested every three days 2-3 times. Recombinant RBD-hFc proteins were captured by Pierce Protein A/G Plus Agarose (Thermo Scientific, 20424), washed by wash buffer (100 mM Tris/HCl, pH 8.0, 150 mM NaCl, 1mM EDTA), eluted with pH 3.0 Glycine buffer (100 mM in H2O), and then immediately balanced by 1/10 volume of UltraPure 1M Tris-HCI, pH 8.0 (15568025, Thermo Scientific). Proteins with twin-strep tag were captured by Strep-Tactin XT 4Flow high-capacity resin (IBA, 2-5030-002), washed by wash buffer, and then eluted by buffer BXT (100 mM Tris/HCl, pH 8.0, 150 mM NaCl, 1mM EDTA, 50 mM biotin). The eluted proteins were concentrated by Ultrafiltration tubes and buffer changed with PBS and stored at -80°C. Protein concentrations were determined by the Omni-Easy Instant BCA Protein Assay Kit (Epizyme, ZJ102).

Coronavirus RBD-hFc live-cell binding assays

Different coronavirus RBD-hFc recombinant proteins were diluted in DMEM at indicated concentrations and incubated with live cells 36 hours post-transfection of different bat ACE2 for 30 mins at 37°C. After binding, cells were washed once by Hanks' Balanced Salt Solution (HBSS) and then incubated with 2 μg/mL of Alexa Fluor 488-conjugated goat anti-human IgG (Thermo Fisher Scientific; A11013) diluted in PBS/1% BSA for 1 hour at 37°C. Next, cells were washed once by HBSS and then incubated with Hoechst 33342 (1:10,000 dilution in HBSS) for nucleus staining for 30mins at 37°C. Images were captured by a fluorescence microscope (MI52-N). The relative fluorescence of the stained cells were determined by a Varioskan LUX Multi-well Luminometer (Thermo Scientific). For flow cytometry analysis, the stained cells were detached by 5 mM of

EDTA/PBS and analyzed with a CytoFLEX Flow Cytometer (Beckman). 10,000 events in a gated

live cell population (based on SSC/FSC) were analyzed for all samples. HEK293T cells without

receptor expression were used as negative controls.

Biolayer interferometry (BLI) binding assays

BLI assays were performed on the Octet RED96 instrument (Molecular Devices) under the

manufacturers' instructions. In general, RBD-hFc recombinant proteins (20 μg/mL) were

immobilized on the Protein A (ProA) biosensors (ForteBio, 18-5010) and incubated with 2-fold

serial diluted bat ACE2-ectodomain proteins starting from 500nM in the kinetic buffer (PBST). The

background was set with a kinetic buffer without the ACE2-ectodomain proteins. The kinetic

parameters and binding affinities between the RBD-hFc and different bat ACE2 were analyzed by

Octet Data Analysis software 12.2.0.20 through curve-fitting kinetic or steady-state analysis.

Pseudovirus production and titration

VSV-dG-based pseudoviruses carrying CoVs spike proteins were produced based on a modified

protocol as previously described[33]. In general, HEK293T cells were transfected with CoVs spike

protein expression plasmids. At 24 hours post-transfection, cells were transduced with 1.5×10^6

$TCID_{50}$ VSV-G glycoprotein-deficient VSV expressing GFP and firefly luciferase

(VSV-dG-GFP-fLuc) diluted in DMEM with 8 μg/mL polybrene for 4-6 hours at 37 ℃. After three

times of PBS wash, the culture medium was replenished with DMEM+10% FBS or SMM 293-TII

containing VSV neutralizing antibody (from I1-mouse hybridoma). Twenty-four hours later, the

virus-containing supernatant was clarified through centrifugation at 4,000 rpm for 5 mins at 4℃ and

then stored at -80℃. $TCID_{50}$ of pseudotyped viruses were determined based on a three-fold

serial-dilution based infection assays on HEK293T-bat40ACE2 cells for NeoCoV and PDF-2180 S

pseudotypes, and 293T-hACE2 for NL63 and SARS-CoV-2 S pseudotypes. $TCID_{50}$ were calculated

according to the Reed-Muench method[34,35].

Pseudovirus entry assay

Pseudovirus entry assays were conducted on HEK293T or Tb 1 Lu cells transiently expressing WT

or mutant ACE2 orthologs at 36 hours post-transfection. In general, $3x10^4$ trypsinized cells were

incubated with pseudovirus ($1.5×10^5$ TCID$_{50}$/100μL) in a 96-well plate to allow attachment and viral

entry. TPCK-trypsin (Sigma-Aldrich, T8802) treatment was conducted before NeoCoV and

PDF-2180 pseudovirus entry assay. In this case, pseudovirus produced in FBS-free SMM-293-TII

medium were incubated with 100 μg/mL TPCK-treated trypsin for 10mins at room temperature,

followed by neutralization with 100 μg/mL soybean trypsin inhibitor (Sigma-Aldrich, T6414). The

intracellular luciferase activity was measured by Bright-Glo Luciferase Assay Kit (Promega, E2620)

and detected with a GloMax 20/20 Luminometer (Promega) at 18 hours post-infection. GFP intensity

were analyzed by a fluorescence microscope (Mshot, MI52-N).

Western-Blot

For Western Blot analysis, cells were lysed with 1% TritonX/PBS+1mM PMSF(Beyotime, ST506)

for 10 mins at 4℃, then clarified through centrifugation of 12000 rpm for 5 mins at 4℃. The

clarified cell lysate was mixed with the 1/5 volume of 5×SDS loading buffer and incubated at 98℃

for 10 mins. After gel electrophoresis and membrane transfer, the PVDF-membrane blots were

blocked with 5% skimmed milk in PBST for 2 hours at room temperature and then incubated

1μg/mL anti-Flag mAb (Sigma, F1804) or anti-glyceraldehyde-3-phosphate dehydrogenase

(GAPDH) (AntGene, ANT011) mAb diluted in PBST containing 1% milk overnight at 4℃. After

three times washing with PBST, the blots were incubated with Horseradish peroxidase

(HRP)-conjugated secondary antibody AffiniPure Goat Anti-Mouse IgG (H+L) (Jackson Immuno

Research, 115-035-003) in 1% skim milk in PBST and incubated for one hour at room temperature.

The blots were then washed three times by PBST and then visualized using an Omni-ECL Femto

Light Chemiluminescence Kit (EpiZyme, SQ201) by a ChemiDoc MP Imaging System (Bio-Rad).

Immunofluorescence assay

Immunofluorescence assays were conducted to verify the expression levels of ACE2 with C-terminal

fused 3×Flag. In general, the transfected cells were incubated with 100% methanol for 10 mins at

room temperature for fixation and permeabilization. Cells were then incubated with a mouse

antibody M2 (Sigma-Aldrich, F1804) diluted in PBS/1%BSA for one hour at 37℃, followed by

extensive wash and the incubation of secondary antibody of Alexa Fluor 594-conjugated goat

anti-mouse IgG (Thermo Fisher Scientific, A32742) diluted in 1% BSA/PBS for one hour at 37℃.

330 Before visualization, the nucleus was stained blue with Hoechst 33342 reagent (1:5,000 dilution in

331 PBS). Images were captured and merged with a fluorescence microscope (Mshot, MI52-N).

Statistical Analysis

334 Most experiments were conducted 2-3 times with 3 or 4 biological repeats. Representative results

335 were shown. Data were presented by MEAN±SD or MEAN±SEM as indicated in the figure legends.

336 Unpaired two-tailed t-tests were conducted for all statistical analyses using GraphPad Prism 8.

337 $P < 0.05$ was considered significant. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.005$, and **** $p < 0.001$.

Acknowledgements

340 Huan Yan was supported by China NSFC projects (32270164, 32070160) and Special Fund for

341 COVID-19 Research of Wuhan University. We thank Beijing Taikang Yicai Foundation for their

342 support to this work.

Author contributions

345 H.Y. and C.B.M. conceived and designed the study. C.B.M., Q.X. and H.Y. performed the

346 experiments and analyzed the data. H.Y., C.B.M. and Q.X. interpreted the results. H.Y. and C.B.M.

347 wrote the manuscript. H.Y. C.B.M. and Q.X revised the manuscript. M.X.G., C.L.W., L.L.S., C.L.,

348 J.Y.S., F.T., P.L.and M.L.H. commented on the manuscript.

Competing interests

351 The authors declare no competing interests.

Data availability

354 All data and information supporting the findings of this study are available with the paper and its

355 supplementary information files.

Additional Information:

358 Correspondence and requests for materials should be addressed to H.Y. (huanyan@whu.edu.cn)

Figure legends

Figure 1 Species-specific ACE2 usage of NeoCoV and several other ACE2-utilizing viruses.

a-b, Most but not all ACE2 orthologs from Yin-bats were deficient in supporting NeoCoV and PDF-2180 binding and entry. NeoCoV RBD-hFc binding (**a**) and pseudovirus entry efficiency (**b**) were evaluated on HEK293T cells transiently expressing the indicated ACE2 orthologs. Vector plasmid was used as a background control. Underline indicate species from Yang-bats. **c**, Distinct RBD binding modes of four ACE2 utilizing coronaviruses. RBD footprints of SARS-CoV-2 (PDB: 6M0J, purple), NL63 (PDB: 3KBH, green), NeoCoV (PDB: 7WPO, blue) and PDF-2180 (PDB: 7WPZ, yellow) on indicated ACE2 were highlighted in red. **d,** Heat map of the RBD binding (mean fluorescence value) and pseudovirus entry efficiency (RLU) of ACE2 utilizing coronaviruses on HEK293T cells transiently expressing the indicated ACE2 orthologs. The pseudovirus entry efficiency on hACE2 was set as 1.0 for SARS-CoV-2 and NL63. The efficiency on Lbor ACE2 (highest efficiency) was set as 1.0 for NeoCoV and PDF-2180. Mean±SD for **b** (n=3) ; Mean for **d** (n=3). RLU: relative light units.

Figure 2 Identification of host range determinants restricting NeoCoV/PDF-2180 recognition.

a, The structure of NeoCoV RBD (purple) and Ppip ACE2 (rainbow) binding interface. Region 1 and region 2 indicate two main interaction surfaces formed by two ACE2 loops with glycosylations. **b-c,** Prediction of the potential host range determinants based on sequence analysis. **b,** The sequence logo analyzes the residue conservation of sequences of the two critical ACE2 loops based on 50 ACE2 functionally divided into two groups (**b**). Upper: efficiently support entry. Lower, deficient in supporting entry. The pie charts show the numbers of Yin-bats and Yang-bats in each group. **c,** Residues showing contrasting differences in frequency or contributing critical interactions were defined as site A-D, and their location in full-length ACE2 was indicated below. TM, transmembrane motif. **d,** Summary of defect types of bat and human ACE2 orthologs according to their residue features in site A-D. The predicted sub-optimal residues in site A (54-56), B (305), C (329-330), and D (337-340) were highlighted with red, green, blue, and magenta, respectively. *, sub-optimal but acceptable by forming a hydrogen bond instead of salt bridge. **e-i,** Structure modeling the impact of sub-optimal residue substitution to the interaction between Ppip ACE2 and NeoCoV RBD. **e,** Overall structure of Ppip ACE2 and NeoCoV RBD

complex, with each sites indicated by dashed boxes. **f-i,** Zoomed in structure details of the interactions in sites A-D in WT (upper) and mutated (lower) Ppip ACE2, respectively. All structures are shown as ribbon representations, with key residues rendered as sticks. Salt bridges and hydrogen bonds are shown as red and yellow dashed lines. Mutated residues were highlighted in purple.

Figure 3 Verification of determinants crucial for species-specific receptor usage in bats.

a-g, Effect of sites A-D gain-of-function mutants of Yin-bats' ACE2 orthologs on NeoCoV RBD and PDF-2180 RBD binding (**a, c, f**) and pseudovirus entry (**b, d, g**). **a-b,** Rsin ACE2 (defect type A); **c-d**, Rfer ACE2 (defect type B); **e-g,** Raeg ACE2 (defect type ABCD). **e,** Schematic illustration of Raeg ACE2 swap mutants with PpipACE2 counterparts. **h**, Flow cytometry analysis of NeoCoV and PDF-2180 RBD-hFc binding with HEK293T cells transiently expressing the indicated ACE2 orthologs. The vector is considered a negative control, and the Ppip ACE2 was employed as a positive control. The red dashed lines indicate the background to define the positive cells. **i**, BLI assays analyzing the binding kinetics between NeoCoV-RBD-hFc and PDF-2180-RBD-hFc and Raeg and Rsin ACE2 WT and mutated ectodomain proteins. Ppip ACE2 was used as a positive control. RLU: Relative luciferase unit. Mean±SD for **b, d, g.** n=3 for **b** and **d**, n=4 for **g**.

Figure 4. RBM mutation T510F enhances the cross-species transmission ability of NeoCoV in bats.

a, Structure showing NeoCoV-T510F interacting with a conserved hydrophobic pocket in Ppip ACE2. Residue conservation was indicated by red and blue colors. The side chains of hydrophobic residues that constitute the pocket were indicated as sticks. The blue and red color on the Ppip ACE2 shows the conservation ratio of 50 Bat ACE2 counterparts residues from low to high. **b,** The sequence logo of surrounding residues of the conserved hydrophobic pocket on 50 bat ACE2. The side chain hydrophobicity was shown in blew. **c-d,** NeoCoV RBD-T510F-hFc RBD binding (**c**) and pseudovirus entry (**d**) on HEK293T cells transiently expressing indicated bat ACE2 orthologs. **e.** The immunofluorescence analysis of the expression level of WT and mutated Raeg and Rsin ACE2. **f.** The NeoCoV-T510F pseudovirus entry to Raeg and Rsin ACE2 WT and mutants. RLU: relative light unit. Mean±SD for **d** and **f**, n=3.

Extended data figure 1. The RBD binding efficiency of ACE2 utilizing CoVs in HEK293T cells transiently expressing the indicated bats ACE2.

a, The expression level of ACE2 orthologs examined by immunofluorescence assay detecting the C terminal fused 3×Flag. **b-d,** The PDF-2180 (**b**), SARS-CoV-2 (**c**), and NL63 (**d**) RBD binding (2μg/mL) on HEK293T cells transiently expressing the indicated bats ACE2.

Extended data figure 2. The pseudovirus entry efficiency of ACE2 utilizing CoVs in HEK293T cells transiently expressing the indicated bats ACE2.

a-c. The entry efficiency of PDF-2180 (**a**), NL63 (**b**), and SARS-CoV-2 (**c**) pseudoviruses in HEK293T cells transiently expressing the indicated bats ACE2. RLU: relative luciferase unit.

Extended 3 Phylogenetic tree of the 50 bats ACE2 tested in this study.

Phylogenetic tree was generated by the MEGA-X software based on the ACE2 protein sequences from the 49 bats species (Two ACE2 sequences for Rsin) across the phylogeny. Yin-bats and Yang-bats were highlighted by turquoise and coral backgrounds. The GenBank accession numbers and protein sequences were summarized in supplementary table 1.

Extended data figure 4. The multi-sequence alignments of 50 bats ACE2.

The sequence alignment was conducted based on sequences around sites A, B, C, and site D by the Clustal W. Identical residues were highlighted in red, and similar residues were in the blue frames.

Extended data figure 5. Western blot analysis of the Rsin, Rfer, and Raeg ACE2 expression levels.

a-c. Western blot analysis of the WT and mutant Rsin(**a**), Rfer(**b**), and Raeg(**c**) ACE2 expression levels in HEK293T cells detecting the C-terminal fused 3×Flag tag. Mw: molecular weight.

Extended data figure 6. Verification of mutants on Nlep, Hgal, Rpea, and Rmal ACE2 with improved NeoCoV/PDF-2180 RBD binding and PSV entry efficiency.

a, Expression levels of the WT and mutated Rpea and Rmal ACE2 by immunofluorescence. **b,** The NeoCoV and PDF-2180 PSV entry efficiency supported by WT and mutated Rpea and Rmal ACE2 in HEK293T cells. **c,** NeoCoV or PDF-2180 RBD binding efficiency on HEK293T cells expressing the WT Hgal ACE2 and related mutants. **d,** The NeoCoV, and PDF-2180 PSV entry efficiency into the HEK293T cells transiently expressing WT and mutated Hgal ACE2. **e,** Western blot analysis of the WT and mutated Hgal ACE2 expression in HEK293T cells.**f,** Schematic illustration of Nlep ACE2 swap mutants with PpipACE2 counterparts. **g,** The binding efficiency of NeoCoV and PDF-2180 RBD to the WT and mutated Nlep ACE2. **h,** Entry efficiency of NeoCoV and PDF-2180 spike pseudotyped virus into the Nlep ACE2 mutants transiently expressing on the HEK293T cells. **i,** Western blot analysis of the Nlep ACE2 mutants expression levels in HEK293T cells. RLU: relative luciferase unit. Mean±SD for **c**, **f**, **i** (n=3). Mw: molecular weight.

Extended data figure 7 NeoCoV and PDF-2180 pseudotyped virus entry in Tb1 Lu cells transiently expressing WT and mutated Raeg and Rsin ACE2.

a. ACE2 expression level of WT and mutated Raeg and Rsin ACE2 in HEK293T cells by detecting the C terminal fused 3×Flag tag by immunofluorescence. **b-d,** NeoCoV and PDF-2180 pseudoviruses entry efficiency at 16 h post-infection were indicated by (GFP) (**b**) and luciferase activity (**c-d**).

1   Li, F. Receptor recognition mechanisms of coronaviruses: a decade of structural studies. *J Virol* **89**, 1954-1964, doi:10.1128/JVI.02615-14 (2015).

2   Xiong, Q. *et al. bioRxiv*, doi:10.1101/2022.01.24.477490 (2022).

3   Chen, L. *et al.* RNA based mNGS approach identifies a novel human coronavirus from two individual pneumonia cases in 2019 Wuhan outbreak. *Emerg Microbes Infect* **9**, 313-319, doi:10.1080/22221751.2020.1725399 (2020).

4   Ksiazek, T. G. *et al.* A novel coronavirus associated with severe acute respiratory syndrome. *N Engl J Med* **348**, 1953-1966, doi:10.1056/NEJMoa030781 (2003).

5   Zaki, A. M., van Boheemen, S., Bestebroer, T. M., Osterhaus, A. D. & Fouchier, R. A. Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J Med* **367**, 1814-1820, doi:10.1056/NEJMoa1211721 (2012).

6   Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* **579**, 270-273, doi:10.1038/s41586-020-2012-7 (2020).

7   Ge, X. Y. *et al.* Isolation and characterization of a bat SARS-like coronavirus that uses the ACE2 receptor. *Nature* **503**, 535-538, doi:10.1038/nature12711 (2013).

8   Latinne, A. *et al.* Origin and cross-species transmission of bat coronaviruses in China. *Nat Commun* **11**, 4235, doi:10.1038/s41467-020-17687-3 (2020).

9   Temmam, S. *et al.* Bat coronaviruses related to SARS-CoV-2 and infectious for human cells. *Nature* **604**, 330-336, doi:10.1038/s41586-022-04532-4 (2022).

10   Wang, Q. *et al.* Bat origins of MERS-CoV supported by bat coronavirus HKU4 usage of human receptor CD26. *Cell Host Microbe* **16**, 328-337, doi:10.1016/j.chom.2014.08.009 (2014).

11   Wong, A. C. P., Li, X., Lau, S. K. P. & Woo, P. C. Y. Global Epidemiology of Bat Coronaviruses. *Viruses* **11**, doi:10.3390/v11020174 (2019).

12   Memish, Z. A. *et al.* Middle East respiratory syndrome coronavirus in bats, Saudi Arabia. *Emerg Infect Dis* **19**, 1819-1823, doi:10.3201/eid1911.131172 (2013).

13   Zhang, Z., Shen, L. & Gu, X. Evolutionary Dynamics of MERS-CoV: Potential Recombination, Positive Selection and Transmission. *Sci Rep* **6**, 25049, doi:10.1038/srep25049 (2016).

14   Mohd, H. A., Al-Tawfiq, J. A. & Memish, Z. A. Middle East Respiratory Syndrome Coronavirus (MERS-CoV) origin and animal reservoir. *Virol J* **13**, 87, doi:10.1186/s12985-016-0544-0 (2016).

15   Anthony, S. J. *et al.* Further Evidence for Bats as the Evolutionary Source of Middle East Respiratory Syndrome Coronavirus. *mBio* **8**, doi:10.1128/mBio.00373-17 (2017).

16   Corman, V. M. *et al.* Rooting the phylogenetic tree of middle East respiratory syndrome coronavirus by characterization of a conspecific virus from an African bat. *J Virol* **88**, 11297-11303, doi:10.1128/JVI.01498-14 (2014).

17   Geldenhuys, M. *et al.* A metagenomic viral discovery approach identifies potential zoonotic and novel mammalian viruses in Neoromicia bats within South Africa. *PLoS One* **13**, e0194527, doi:10.1371/journal.pone.0194527 (2018).

18   Corman, V. M. *et al.* Characterization of a novel betacoronavirus related to middle East respiratory syndrome coronavirus in European hedgehogs. *J Virol* **88**, 717-724, doi:10.1128/JVI.01600-13 (2014).

19   Ithete, N. L. *et al.* Close relative of human Middle East respiratory syndrome coronavirus in bat, South Africa. *Emerg Infect Dis* **19**, 1697-1699, doi:10.3201/eid1910.130946 (2013).

20   Lau, S. K. P. *et al.* Identification of a Novel Betacoronavirus (Merbecovirus) in Amur Hedgehogs from China. *Viruses* **11**, doi:10.3390/v11110980 (2019).

21   Yan, H. *et al.* ACE2 receptor usage reveals variation in susceptibility to SARS-CoV and SARS-CoV-2 infection among bat species. *Nat Ecol Evol* **5**, 600-608, doi:10.1038/s41559-021-01407-1 (2021).

22   Huynh, J. *et al.* Evidence supporting a zoonotic origin of human coronavirus strain NL63. *J Virol* **86**, 12816-12825, doi:10.1128/JVI.00906-12 (2012).

23   Wilson, D. E. M., R. A. Handbook of the Mammals of the World.   **Vol. 9: Bats (Lynx, 2019)** (2019).

24   King, S. B. & Singh, M. Comparative genomic analysis reveals varying levels of mammalian adaptation to coronavirus infections. *PLoS Comput Biol* **17**, e1009560, doi:10.1371/journal.pcbi.1009560 (2021).

25   Liu, Y. *et al.* Functional and genetic analysis of viral receptor ACE2 orthologs reveals a broad potential host range of SARS-CoV-2. *Proc Natl Acad Sci U S A* **118**, doi:10.1073/pnas.2025373118 (2021).

26   Letko, M. *et al.* Adaptive Evolution of MERS-CoV to Species Variation in DPP4. *Cell Rep* **24**, 1730-1737, doi:10.1016/j.celrep.2018.07.045 (2018).

27   Hu, B. *et al.* Discovery of a rich gene pool of bat SARS-related coronaviruses provides new insights into

the origin of SARS coronavirus. *PLoS Pathog* **13**, e1006698, doi:10.1371/journal.ppat.1006698 (2017).

28   Mehdipour, A. R. & Hummer, G. Dual nature of human ACE2 glycosylation in binding to SARS-CoV-2

spike. *Proc Natl Acad Sci U S A* **118**, doi:10.1073/pnas.2100425118 (2021).

29   Tortorici, M. A. *et al.* Structure, receptor recognition, and antigenicity of the human coronavirus

CCoV-HuPn-2018 spike glycoprotein. *Cell* **185**, 2279-2291 e2217, doi:10.1016/j.cell.2022.05.019 (2022).

30   Zhang, Y. *et al.* Cross-species tropism and antigenic landscapes of circulating SARS-CoV-2 variants. *Cell*

Rep **38**, 110558, doi:10.1016/j.celrep.2022.110558 (2022).

31   Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and

annotation. *Nucleic Acids Res* **49**, W293-W296, doi:10.1093/nar/gkab301 (2021).

32   Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers, educators, and developers.

Protein Sci **30**, 70-82, doi:10.1002/pro.3943 (2021).

33   Whitt, M. A. Generation of VSV pseudotypes using recombinant DeltaG-VSV for studies on virus entry,

identification of entry inhibitors, and immune responses to vaccines. *J Virol Methods* **169**, 365-374,

doi:10.1016/j.jviromet.2010.08.006 (2010).

34   Biacchesi, S. *et al.* Rapid human metapneumovirus microneutralization assay based on green fluorescent

protein expression. *J Virol Methods* **128**, 192-197, doi:10.1016/j.jviromet.2005.05.005 (2005).

35   Nie, J. *et al.* Quantification of SARS-CoV-2 neutralizing antibody by a pseudotyped virus-based assay. *Nat*

Protoc **15**, 3699-3715, doi:10.1038/s41596-020-0394-5 (2020).

Figure1

d

Summary of residues in sites A-D in ACE2 orthologs

	Species	A	B	C	D	defect types
Yin-bats	Raeg	NII	K	E K	DNDQ	ABCD
	Pale	NIT	K	E K	GNDQ	BCD
	Pgig	NIT	K	E K	GNDQ	BCD
	Ehel	NIT	K	E K	GNDQ	BCD
	Espe	NIT	K	E K	GNDQ	BCD
	Msob	NIT	K	E K	GNDQ	BCD
	Csph	NIT	K	E K	GNDQ	BCD
	Cbra	NIT	K	E K	GNDQ	BCD
	Rfer	NIS	K	N N	GDGR	B
	Rsin	NIN	D	N N	GDGR	A
	Rpea	NIS	N	N N	GDGR	B*
	Harm	NIT	K	E N	GDGR	BC
	Hgal	NIT	I	E N	GDGR	BC
	Hpra	NIT	K	E N	GDGR	BC
	Mlyr	NIT	D	N N	G_GR	D
	Raff	NIS	N	N N	GDGR	B*
	Rcor	NIS	N	N N	GDGR	B*
	Rmal	NIS	N	N N	GDGR	B*
Yang-bats	Nlep	NIT	V	K N	EDDR	BCD
	Ppip	NIT	E	N N	SDGR	
	Lbor	NIT	E	S N	ADGR	
	Nhum	NIT	E	N N	GNGR	D
Human	Human	NIT	Q	E N	GNVQ	CD

Figure3

a

Rsin

WT | N55T | Ppip | Vector

Expression ACE2 (Flag)

100µm

RBD binding NeoCoV

100µm

PDF-2180

100µm

Defect type A

b

NeoCoV
RLU (PSV entry, Log₁₀)
****
WT | N55T | Ppip | Vector
Rsin

PDF-2180
****
WT | N55T | Ppip | Vector
Rsin

c

Rfer

WT | K305E | Ppip | Vector

Expression ACE2 (Flag)

100µm

RBD binding NeoCoV

100µm

PDF-2180

100µm

Defect type B

d

NeoCoV
RLU (PSV entry)
****
WT | K305E | Ppip | Vector

PDF-2180
****
WT | K305E | Ppip | Vector
Rfer

e

54-56 305 329-330 337-340
Ppip ACE2 N NIT E NN SDGR C
Raeg ACE2 N NII K EK DNDQ C
A N T C
AB N T E C
ABC N T E NN C
ABCD N T E NN SDGR C
53-55 305- 329-330 337-340

f

Raeg

WT | A | AB | ABC | ABCD | Ppip | Vector

Expression ACE2 (Flag)

100µm

RBD binding NeoCoV

100µm

PDF-2180

100µm

Defect type ABCD

g

NeoCoV
RLU (PSV entry, Log₁₀)
NS NS ** **** ****
WT | A | AB | ABC | ABCD | Ppip | Vector

PDF-2180
NS *** *** ****
WT | A | AB | ABC | ABCD | Ppip | Vector
Raeg

h

NeoCoV | PDF-2180
Cell counts

0.11±0.03% 1.43±0.18% Vector
87.9±0.5% 73.95±0.35% Ppip ACE2
0.15±0.02% 0.73±0.1% Rsin ACE2
87.7±0.02% 63.6±0.4% Rsin ACE2 (N55T)
0.23±0.08% 1.75±0.2% Raeg ACE2
86.15±0.55% 59.25±0.55% Raeg ACE2 (ABCD)

RBD binding

i

— 500nM — 250nM — 125nM — 62.5nM — 31.25nM — 15.625nM

Response(nm)

NeoCoV RBD

Raeg ACE2 WT | Raeg ACE2 (ABCD) | Rsin ACE2 WT | Rsin ACE2 N55T | Ppip ACE2

Raeg ACE2 (ABCD):
$K_D(M)=2.35E-08$
$k_{on}(Ms^{-1})=5.88E+03$
$k_{off}(S^{-1})=1.38E-04$
$R^2=0.9978$

Rsin ACE2 N55T:
$K_D(M)=2.04E-08$
$k_{on}(Ms^{-1})=3.99E+04$
$k_{off}(S^{-1})=8.12E-04$
$R^2=0.9978$

Ppip ACE2:
$K_D(M)=1.78E-09$
$k_{on}(Ms^{-1})=5.85E+04$
$k_{off}(S^{-1})=1.04E-04$
$R^2=0.9981$

PDF-2180 RBD

Raeg ACE2 (ABCD):
$K_D(M)=6.94E-10$
$k_{on}(Ms^{-1})=1.58E+04$
$k_{off}(S^{-1})=1.10E-05$
$R^2=0.9987$

Rsin ACE2 N55T:
SSG $K_D(M)=6.94E-10$
SSG $R^2=0.9987$

Ppip ACE2:
$K_D(M)=4.17E-10$
$k_{on}(Ms^{-1})=1.33E+05$
$k_{off}(S^{-1})=5.54E-05$
$R^2=0.9920$

Time(s)

Figure 4

a

NeoCoV RBD

Ppip ACE2

F308
T510
W328
M323 F327

Ratio of conservation
Low   High

b

Sequence logo
Residue frequency %

F308   M323   F327 W328

100
50
0

FKEAEKFFvSvGLpNMTeGFW

Mean Hydrophobicity

50 bat ACE2

c NeoCoV-T510F RBD binding

Raeg	Pale	Pgig	Ehel	Espe	Msob	Csph	Cbra
Rfer	Rsin	Rpea	Harm	Hgal	Hpra	Mlyr	Raff
Rcor	Rmal	Nlep	Ppip	Lbor	Nhum	Human	Vector

100μm (×3)

d NeoCoV-T510F pseudovirus entry

RLU (PSV entry, Log$_{10}$)

Raeg Pale Pgig Ehel Espe Msob Csph Cbra Rfer Rsin Rpea Harm Hgal Hpra Mlyr Raff Rcor Rmal | Nlep Ppip Lbor Nhum | Human Vector

Yin-bats   Yang-bats

e

ACE2 expression (Flag)

Raeg
WT A AB ABC ABCD

Rsin
WT N55T   Ppip Lbor Vector

100μm (×2)

f NeoCoV-T510F pseudovirus entry

RLU (PSV entry, Log$_{10}$)

WT A AB ABC ABCD WT N55T Ppip Lbor Vector
*** **** * ****   ****

Raeg   Rsin

a

ACE2 expression-HEK293T cells

b

PDF-2180 RBD binding

c

SARS-CoV-2 RBD binding

d

NL63 RBD binding

a

b

c

Yinpterochiroptera (19)

Megadermatidae   *Hipposideridae*

Rhinolophidae

Pteropodidae(Old world fruit bats)

Emballonuridae

Noctilionidae

Mormoopidae

Phyllostomidae(New world leaf-nosed bats)

Molossidae

Miniopteridae

Vespertilionidae(Vesper bats)

Yangpterochiroptera (31)

Tree scale: 0.1

Extended 4

Site A (54, 56) | Site B (302, 305) | Site C (329, 330) | Site D (337, 340, 343)

Species	Site A	Alignment	Site C	Site D	Code
Rousettus_aegyptiacus	NII	WNAKRIFKEAEKFFVSLGLPNMTE TFWEK	SVLTEP	DNDQKVA	Raeg
Rhinolophus_sinicus-3357	NIN	WDADRIFKEAEKFFVSVGLPNMTE GFWNN	SMLTEP	GDGRKVV	Rsin-3357
Pteropus_giganteus	NIT	WDEKRIFKEAEKFFVSLGLPNMTE KFWEK	SMLTEP	GNDQKVA	Pgig
Hipposideros_galeritus	NIT	WDAIKIFQEAEKFFVSIGLPKMTE GFWEN	SMLTEP	GDGRKVV	Hgal
Rhinolophus_sinicus	NIN	WDADRIFKEAEKFFVSVGLPNMTE GFWNN	SMLTEP	GDGRKVV	Rsin
Macroglossus_sobrinus	NIT	WDAKRIFKEAENFFVSLGLPNMTE KFWEK	SMLTEP	GNDQKVA	Msob
Hipposideros_pratti	NIT	WDAKKIFQEAEKFFVSVGLPNMTK GFWEN	SMLTEP	GDGRKVV	Hpra
Eidolon_helvum	NIT	WDAKRIFKEAEKFFVSLGLPNMTE KFWEK	SMLTEP	GNDQKVA	Ehel
Pteropus_alecto	NIT	WDEKRIFKEAEKFFVSLGLPNMTE KFWEK	SMLTEP	GNDQKVA	Pale
Eonycteris_spelaea	NIT	WSAKRIFKEAEKFFVSLGLPNMTE KFWEK	SMLTEP	GNDQKVA	Espe
Cynopterus_brachyotis	NIT	WDAKRIFKEAEKFFVSLGLPNMTE GFWEK	SMLTEP	GNDQKVA	Cbra
Noctilio_leporinus	NIT	WDAVRIFKEAEKFFVSVGLSNMTQ GFWKN	SMLTKP	EDDRKVV	Nlep
Hipposideros_armiger	NIT	WDAKKIFQEAEKFFVSVGLPNMTK GFWEN	SMLTEP	GDGRKVV	Harm
Rhinolophus_ferrumequinum	NIS	WDAKRIFKEAEKFFVSIGLPNMTE GFWNN	SMLTDP	GDGRKVV	Rfer
Cynopterus_sphinx	NIT	WDAKRIFKEAEKFFVSLGLPNMTE GFWEK	SMLTEP	GNDQKVA	Csph
Megaderma_lyra	NIT	WDADRIFKEAEKFFVSVGLPNMTE GFWNN	SMLTEP	G.DRKVV	Mlyr
Rhinolophus_pearsonii	NIS	WDANRIFKEAEKFFVSVGLPNMTE GFWNN	SMLTEP	GDGRKVV	Rpea
Nycticeius_humeralis	NIT	WDAEKIFKEAEKFYMSVGLPAMTP GFWNN	SMLTEP	GNGRKVV	Nhum
Homo_sapien	NIT	WDAQRIFKEAEKFFVSVGLPNMTQ GFWEN	SMLTDP	GNVQKAV	Human
Rhinolophus_affinis	NIS	WDANRIFKEAEKFFVSVGLPNMTE GFWNN	SMLTEP	GDGRKVV	Raff
Rhinolophus_malayanus-4	NIS	WDANRIFKEAEKFFVSVGLPNMTE GFWEK	SMLTEP	GDGRKVV	Rmal
Rhinolophus_cornutus	NIS	WDANRIFKEAEKFFVSVGLPNMTE GFWNN	SMLTEP	GDGRKVV	Rcor
Tonatia_saurophila	NIT	WDAQRIFKEAEKFFVSVGLFNMTQ GFWDN	SMLTKP	DDGREVV	Tsau
Pteronotus_parnellii	NIT	WDAKKIFKEAEKFFVSVGLFNMTQ GFWDN	SMLTKP	DDGREVV	Ppam
Tadarida_brasiliensis	NIT	WNAERIFKEAEKFFVSIGLPNMTQ EFWNN	SMLTEP	GDGREVV	Tbra
Anoura_caudifer	NIT	WDAQKIFKKAEEFFRSVGLYNMTQ GFWDN	SMLTKP	DDGREVV	Acau
Pipistrellus_kuhlii	NIT	WDADKIFKEAEKFYLSVGLRNMTP GFWNK	SMLTEP	SDGRQVV	Pkuh
Myotis_lucifugus	NIT	WDAEKIFKEAEKFYISVGLPSMTP GFWNN	SMLTEP	GDGRKVV	Mluc
Phyllostomus_discolor	NIT	WDAQRIFKEAEKFFVSVGLFNMTQ GFWDN	SMLTKP	DDGREVV	Pdis
Vampyram_spectrum	NIT	WDAQRIFKEAEKFFVSVGLFNMTQ GFWDN	SMLTKP	DDGREVV	Vspe
Pteronotus_davyi	NIT	WDAEKIFKEAEKFFMSVGLFNMTQ GFWNN	SMLTKP	DDGREVV	Pdav
Eptesicus_fuscus	NIT	WDAEKIFKEAEKFYMSVGLPSMTP GFWNN	SMLTEP	GDGRKVV	Efus
Miniopterus_natalensis	NIT	WSAEKIFKEAEKFYVSVGLPNMTE GFWNN	SMLTEP	GDGRKVV	Mnat
Murina_feae	NIT	WDAEKIFKEAEKFYVSVGLPSMTA GFWNN	SMLTEP	GDGRKVV	Mfea
Myotis_myotis	NIT	WDAEKIFKEAEKFYISVGLPSMTP GFWNN	SMLTEP	GDGRKVV	Mmyo
Artibeus_jamaicensis	NIT	WDAQRIFKEAEKFFMSVGLFNMTQ GFWDN	SMLTKP	DDGREVV	Ajam
Miniopterus_schreibersii	NIT	WSAEKIFKEAEKFYVSVGLPNMTE GFWNN	SMLTEP	GDGRKVV	Msch
Myotis_brandtii	NIT	WDAEKIFKEAEKFYISVGLPSMTP GFWNN	SMLTEP	GDGRKVV	Mbra
Myotis_davidii	NIT	WDAEKIFKEAEKFYISVGLPSMTP GFWKN	SMLTEP	GDGRKVV	Mdav
Sturnira_hondurensis	NIT	WDAQKIFKEAEKFFISVGLYNMTQ GFWDN	SMLTKP	DDGREVV	Shon
Micronycteris_hirsuta	NIT	WDARRIFEEAEKFFRSVGLFNMTQ GFWDN	SMLTKP	DDGREVV	Mhir
Pipistrellus_pipistrellus	NIT	WDAEKIFKEAEKFYLSVGLYSMTQ GFWNN	SMLTEP	SDGRQVV	Ppip
Trachops_cirrhosus	NIT	WDAQRIFKEAEKFFVSVGLFNMTQ GFWDN	SMLTKP	DDGREVV	Tcir
Carollia_perspicillata	NIT	WDAQKIFKEAEKFFVSVGLFSMTQ GFWDN	SMLTKP	DDGREVV	Cper
Desmodus_rotundus	NIT	WDAQRIFKEAEKFFKSVGLFSMTQ GFWDN	SMLTKP	DDGREVV	Drot
Taphozous_melanopogon	NIT	WDAERIFKEAEKFFVSIALPNMTE GFWNK	SMLTEP	SDGRKVV	Tmel
Mormoops_blainvillei	NIT	WNAQMIFKAAEDFFTSIGLPNMTQ GFWDN	SMLTKP	DDGRKVV	Mbla
Molossus_molossus	NIT	WDAEKIFKKAEEFFVSIGLGNMTQ GFWDN	SMLTEP	SDGRKVV	Mmol
Aeorestes_cinereus	NIT	WDAEKIFKEAEKFYVSVGLPRMTP GFWSN	SMLTEPA	DGRKVV	Acin
Antrozous_pallidus	NIT	WDAEKIFKEAEKFYVSVGLPNMTS GFWDN	SMLTEP	GDGRKVV	Apal
Lasiurus_borealis	NIT	WDAEKIFKEAEKFYASVGLPNMTS GFWSN	SMLTEPA	DGRKVV	Lbor

Extended 5

a

Rsin

WT N55T Ppip Vector

Mw(kDa)

ACE2 (Flag)   — 100

GAPDH   — 35

b

Rfer

WT K305E Ppip Vector

Mw(kDa)

— 100

— 35

c

Raeg

WT A AB ABC ABCD Ppip Vector

Mw(kDa)

— 100

— 35

Defect type B*

Defect type BC

Defect type BCD

Tb1 Lu

a

Expression
ACE2 (Flag)

Ppip   Raeg (ABCD)   Rsin N55T   Vector

200μm

b

Pseudovirus entry

NeoCoV

300μm

PDF-2180

300μm

c

NeoCoV pseudovirus entry

NS   ***

RLU (PSV entry, Log_{10})

Raeg (ABCD)   Rsin N55T   Ppip   Vector

d

PDF-2180 pseudovirus entry

NS   ****

RLU (PSV entry, Log_{10})

Raeg (ABCD)   Rsin N55T   Ppip   Vector