Global phylogenomic assessment of *Leptoseris* and *Agaricia* reveals substantial undescribed diversity at mesophotic depths

Gijsbers JC[1]*, Englebert N[2], Prata KE[1,3], Pichon M[4], Dinesen Z[5], Brunner R[2,6], Eyal G[3,7,8], González-Zapata FL[9], Kahng SE[10], Latijnhouwers KRW[12,13], Muir P[2], Radice VZ[2,11], Sánchez JA[9], Vermeij MJA[12,13], Hoegh-Guldberg O[2,3,7], Jacobs SJ[1], Bongaerts P [1,2,12]*

[1] California Academy of Sciences, San Francisco, California 94118, USA

[2] Global Change Institute, The University of Queensland, St Lucia Queensland 4072, Australia

[3] School of Biological Sciences, The University of Queensland, St Lucia, Queensland 4072, Australia

[4] Biodiversity Section, Queensland Museum, Townsville 4811, Australia

[5] Centre for Biodiversity and Conservation Science, The University of Queensland, St Lucia Queensland 4072, Australia

[6] ARC Centre of Excellence for Coral Reef Studies, James Cook University Townsville, Queensland 4811, Australia

[7] ARC Centre of Excellence for Coral Reef Studies, The University of Queensland, St Lucia, Queensland 4072, Australia

[8] The Mina & Everard Goodman Faculty of Life Sciences, Bar-Ilan University, Ramat Gan, 5290002, Israel

[9] Laboratorio de Biología Molecular Marina (BIOMMAR), Departamento de Ciencias Biológicas, Facultad de Ciencias, Universidad de Los Andes, Bogotá 111711, Colombia

[10] Department of Oceanography, University of Hawaii at Manoa, 1000 Pope Road, Honolulu, Hawaii 96822, USA

[11] Department of Biological Sciences, Old Dominion University, Norfolk, Virginia 23529, USA

[12] CARMABI Foundation, PO Box 2090, Piscaderabaai z/n, Willemstad, Curaçao

[13] Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, Science Park 700, 1098 XH Amsterdam, The Netherlands

* Corresponding authors: jgijsbers@calacademy.org, pim@calacademy.org

Keywords (5): *Mesophotic, phylogenetics, Scleractinia, biodiversity, depth-differentiation*

ABSTRACT

Mesophotic coral communities are increasingly gaining attention for the unique biological species they host, exemplified by the numerous mesophotic fish species that continue to be discovered. In contrast, many of the photosynthetic scleractinian corals observed at mesophotic depths are assumed to be depth-generalists, with very few species characterised as mesophotic-specialists. This presumed lack of a specialised community remains largely untested, as phylogenetic studies rarely include mesophotic samples and have long suffered from resolution issues associated with traditional sequence markers. Here, we used reduced-representation genome sequencing to provide a phylogenomic assessment of the two dominant mesophotic genera of plating corals in the Indo-Pacific and Western Atlantic, respectively, *Leptoseris* and *Agaricia*. While these genome-wide phylogenies broadly corroborated the morphological taxonomy, they also exposed substantial undescribed diversity (at least 24 molecular clades across the 8 species), including deep divergences within genera and current taxonomic species. Five of the eight focal species consisted of at least two sympatrically-occurring, genetically distinct clades, which were consistently detected across different methods. The repeated observation of genetically divergent clades associated with mesophotic depths highlights that there are many more mesophotic-specialist coral species than currently acknowledged, and that an urgent assessment of this largely unstudied biological diversity is warranted.

INTRODUCTION

Because mesophotic coral ecosystems (MCEs) occur at depths beyond the limits of regular SCUBA diving (~30-150 m depth), they remain relatively understudied compared to shallow coral reefs, despite equalling or even exceeding the area occupied by the latter [1]. Over the past decades, interest in these deeper coral reef communities has grown due to their potential to act as a refuge against disturbances (for species with large depth distributions) [2-5] and as habitats hosting unique biological communities [7-9]. Their uniqueness is exemplified by the diversity and continuous discovery of depth-specialist fish species at mesophotic depths [6,9], as well as the vast differences observed between shallow and mesophotic fish species assemblages [7]. Although a similar differentiation over depth has been observed for reef-building coral assemblages [7,10], only a few scleractinian coral species dominate the assemblage at lower mesophotic depths [2,11-15], and a small proportion of them are considered deep-specialists [15,16].

Visual assessments (*in situ* or through imagery) of mesophotic coral diversity are challenging due

to the intricate scale of and significant intraspecific variation in morphological traits used to

identify species [16,17]. Because collection-based assessments of coral diversity are rare for

mesophotic reef corals (particularly for depths >~60 m) [10] and because reference collections

contain mostly shallow water coral specimens [15,18], morphological differences among shallow

and mesophotic species can easily go unnoticed. Moreover, genetic identification has been

hampered by the lack of species-level resolution when using traditional sequencing markers in

scleractinian corals [19-22]. Consequently, these methodological challenges have greatly hindered

the ability to differentiate and identify putative new scleractinian coral species associated with

mesophotic depths.

Throughout the tropics, mesophotic coral ecosystems host reef-building scleractinian coral species

with predominantly plating growth forms, which maximize light capture by their symbiotic

dinoflagellates to sustain photosynthesis in low light conditions at greater depths [23-25]. Most of

these plating corals belong to the family Agariciidae, with the genera *Leptoseris* and *Pavona*

dominating mesophotic coral communities in the Indo-Pacific, and *Agaricia* in the Western

Atlantic [11,14,17,26]. In these genera, species can generally occur over wide depth ranges, cover

large areas of substrate, and provide important habitat structure to other reef-associated mesophotic

organisms [13,27,28]. The genus *Leptoseris* is particularly dominant at lower mesophotic depths

(>60 m [24]), and has been reported at depths of 172 m [29]. In Eastern Australia (Great Barrier

Reef and Western Coral Sea) it has been observed down to 125 m depth [30], with four taxonomic

species –*Leptoseris scabra* [31], *L. glabra* ([18], *L. explanata sensu* [32]), *L. mycetoseroides* [33],

and *L. hawaiiensis* [31]– observed to dominate scleractinian coral communities at mesophotic

depths [14,15]. These species also have a wide distribution ranging from the Red Sea [26,34,35]

to the Hawaiian Archipelago (albeit with narrower depth distributions; [17]). The Western Atlantic

genus *Agaricia* comprises seven species, of which four are dominant members of the mesophotic

coral communities [27,36,37]: *Agaricia lamarcki* [38], *A. fragilis* [39], *A. grahamae* [40]*,* and *A.*

undata [41]. These four species occur throughout most of the Caribbean basin, and *A. fragilis* even

extends north to Bermuda [3] and south to the Brazilian coast [42]. The deepest *Agaricia* colony

(*A. grahamae*) was reported from a depth of 119 m [43]. Many of these widespread *Leptoseris* and

Agaricia species occur across large depth ranges, making them important candidates to address

the question to what extent mesophotic coral communities harbour unique depth-specialized

species.

Molecular assessments of the genera *Leptoseris* and *Agaricia* using traditional sequence markers

have exposed polyphyletic patterns, often in discordance with morphology-based taxonomy

[11,13,17,27,34], and frequently were unable to discriminate between some of the well-established

and morphologically distinct agariciid species [11,19,20,22,44-47]. Nonetheless, despite the

pervasive and well-known issues with these markers [21,22], the few molecular studies that have

been undertaken on this critically important family of scleractinian corals (Agariciidae) have

highlighted the potential for undescribed diversity and depth-differentiation [11,13,17,34].

Reduced-representation genome sequencing methods (e.g., sequencing of restriction site-

associated DNA (RAD-seq) or ultra-conserved elements (UCEs)) have demonstrated their

potential to overcome these issues, resulting in phylogenies that have strong support (e.g., [48-

52]), providing a promising way forward to study the evolutionary relationships within the

Agariciidae. The increased resolution of such reduced representation methods was demonstrated

through recent population genomics studies of *Agaricia* species, revealing significant genetic

structuring within all four species dominating Atlantic mesophotic communities [3,37,53,54].

Here, we build on these initial findings and present a phylogenomic assessment of the two

dominant mesophotic genera found within the Agariciidae family (figure 1b-c) in the Indo-Pacific

and Western Atlantic (figure 1a). Focusing on eight species from the genera *Leptoseris* and

Agaricia (figure 1a-b), we evaluate whether current systematics give an accurate reflection of the

species diversity and extent of specialisation (i.e., depth specificity) in mesophotic coral

communities.

Figure 1. Overview of the *Leptoseris* and *Agaricia* study species (a) Map of the sampling
locations for *Leptoseris* in the Indo-Pacific (blue) and *Agaricia* in the Western Atlantic (orange).
(b) *Leptoseris*-dominated coral community at 87 m depth in the Hawaiian Archipelago (Photo:
Hawaiian Undersea Research Laboratory). (c) *Agaricia*-dominated coral community at 40 m in
Curaçao, Southern Caribbean. (d) Phylogenetic tree (RAxML-ng) based on 37,528 concatenated
nextRAD loci (3,361,114 sites) of the overall dataset, highlighting the position of *Leptoseris* (blue)
and *Agaricia* (red) relative to included outgroups. Specimens from the focal species of this study
are represented by coloured branches in the tree, with grey branches representing unidentified
specimens.

METHODS

Sample collection and preparation

Coral specimens (n = 189) from the genera *Leptoseris* and *Agaricia* were collected as part of the "XL Catlin Seaview Survey" during visits to Eastern Australia (Western Coral Sea and Great Barrier Reef) and the Southern Caribbean (Curaçao and Bonaire; figure 1a, table S1). Samples were collected in Eastern Australia with a Seabotix vLBV300 Remotely Operated Vehicle [14,15] and in the Southern Caribbean with the "Curasub" submersible [11] and technical SCUBA. Small fragments (~1 cm²) were stored in 99% EtOH or NaCl 20% DMSO 0.5M EDTA solutions for genomic DNA extraction. Bleached skeletal specimens (all *Leptoseris*, and a subset of *Agaricia*) were deposited at the Queensland Museum Collection or the Invertebrate Zoology collection at the California Academy of Sciences (table S1). Additional tissue samples were acquired from collaborators in the Hawaiian Archipelago, US Minor Outlying Islands, Gulf of Aqaba (Red Sea), San Andrés Island, Cartagena, and Santa Marta (Colombia). The *Agaricia* dataset includes nextRAD sequence data (n = 52) from three published datasets [3,37,53], and we also included *Stephanocoenia intersepta* (n = 3 [3]), *Helioseris cucullata* (n = 5), and *Pachyseris speciosa* (n = 3 [55]) samples to serve as outgroups (table S1).

Molecular dataset preparation

Genomic DNA was extracted as described in Bongaerts et al. [3,55], using the additional centrifugation steps to reduce endosymbiont contamination when sufficiently high gDNA yields were obtained. Extracted gDNA was used to create nextRAD DNA libraries (SNPsaurus, LLC). using selective PCR primers to genotype genomic loci consistently [56]. gDNA was fragmented and ligated with Nextera adapters (Illumina Inc). Once ligated, fragmented DNA was PCR-amplified (26 cycles, 73°C) with the matching primer for the adapter ("GTGTAGAGG"). Final libraries were sequenced (Illumina HiSeq 2500) to generate 100 bp single-end reads. Nextera adapters and low-quality ends (PHRED-quality score below 20) were trimmed using TrimGalore v0.6.4 (https://github.com/FelixKrueger/TrimGalore) to discard reads <30 bp and trim sequences up to 100 bp. IpyRAD v0.9.62 [57] was used for locus clustering and variant calling (85% clustering threshold, minimum coverage of six, minimum four samples per final locus, with all other settings run at default recommended values) (table S3).

Symbiont contamination was identified through a BLASTN comparison of each nextRAD locus

against nextRAD sequence data from isolated Symbiodiniaceae, and four published

Symbiodiniaceae genomes: *Symbiodinium microadriaticum* [58], *Breviolum minutum* [59],

Cladocopium goreaui [60], and *Durusdinium trenchii* [61] removing positive matches (maximum

E-value = 10^{-15}) from the coral nextRAD loci. Potential microbial contamination was identified by

an additional BLASTN comparison against the NCBI non-redundant database. The taxonomic IDs

of positive matches (maximum E-value = 10^{-4}) of non-cnidarian taxa were removed from the

dataset. Three datasets were created from the filtered assembly and used for all downstream

analyses: the *Leptoseris* dataset, composed of individuals from the genus *Leptoseris* with

Helioseris cucullata and *A. fragilis* as outgroups (n = 127), the *Agaricia* dataset, composed of

individuals from the genus *Agaricia* with *H. cucullata* and *L. glabra* as outgroups (n = 70), and

the "*Agalepto*" dataset, composed of individuals from both genera with *H. cucullata*, *S. intersepta*,

and *P. speciosa* as outgroups (n = 201) (figure 1d-e, table S1). NextRAD loci were then trimmed

to 90 bp and filtered to retain loci genotyped for ε10 samples, with the resulting Variant Call

Format (VCF) file filtered to retain only those SNPs genotyped for ε10% of samples.

Phylogenetic and species tree inference

Maximum likelihood (ML) phylogenetic inference was performed using RAxML-ng v.1.0.1 [62]

and a concatenated matrix with complete sequences of all loci. The best fit model of nucleotide

substitution GTR+I+G4 was identified by ModelTest-ng x.y.z [63] and independent searches and

bootstrap replicates were performed on each alignment until convergence was reached. To assess

genealogical concordance, we used the concordance factor analysis as implemented in IQ-Tree

2.1.4-beta [64,65]. This analysis first infers single locus phylogenies (coupled with model

selection) and then calculates the percentage of gene trees and alignment sites that are concordant

with a reference ML topology (gCF and sCF, respectively). The reference ML topology was

inferred in IQ-Tree using the concatenated matrix, applying an edge-linked proportional partition

model [64] and employing standard model selection [66]. Species tree inference was performed

using Tetrad, a coalescent species tree approach based on SVDquartets [67] implemented in

IpyRAD v 0.9.65 [57]. This analysis constructs quartets using SNPs sampled from each locus and

joins these to create a 'supertree' statistically consistent under the multispecies coalescent model.

A single SNP was randomly sampled from each locus, all possible quartets were sampled, and 100

bootstrap replicates performed generating a majority-rule consensus tree and individual trees used

to generate a density tree.

Genetic structure using SNP-based analyses

To assess the genetic structure across our samples while avoiding the bias of *a priori* species

assignment, we used *de novo* Discriminant Analysis of Principal Components (DAPC, [68]), an

unsupervised dimensionality reduction method where novel genetic clusters are defined using a

K-means clustering method and subsequently visualised using Principal Components Analysis

(PCA). The DAPC analysis was conducted in R with the Adegenet package [69], choosing PCs

based upon the optimal a-score and assessing the optimal numbers of clusters running K-means

sequentially with increasing values of K, starting with K equal to the lowest AIC and BIC and then

sequentially increasing K up to the maximum number of clades found in the ML tree.

Comparison with traditional sequence markers

To compare the resolution of nextRAD with traditional mitochondrial markers, *cox*1-1-rRNA

intron sequence data of 101 *Leptoseris*, 9 *Agaricia*, and 1 *Pavona* individuals were amplified using

AGAH/AGAL primer pairs [70]. The PCR amplifications were performed following the approach

of Bongaerts *et al.* [27]. Agarose gels were used to assess the quality of the PCR products, being

then cleaned (ExoSAP-IT) and sequenced in forward and reverse directions (ABI BigDye

Terminator chemistry, Australian Genome Research Facility). Additionally, previously published

*cox*1-1-rRNA intron sequence data of 46 *Leptoseris* individuals [17], 61 *Agaricia* [11,71], 1

Pavona clavus [11], and 1 *L. hawaiiensis* [11] individuals were retrieved from GenBank.

Codoncode Aligner was used to analyse the resulting sequences. ML phylogenies were inferred

using RAxML-ng v.0.9.0 on the concatenated alignment, under the K80+G4 model, with a

calculation of 20 trees and bootstrap support values based on 50,000 and 3,300 replicates for

species from the genera *Leptoseris* and *Agaricia*, respectively.

RESULTS

Phylogenomic patterns across the genera *Leptoseris* and *Agaricia*

Using a reduced-representation sequencing approach (nextRAD), we recovered an average of 2.6

million reads (range: 580K-12.7M) from 201 scleractinian coral specimens; *Leptoseris* samples

averaged 1.8 million reads (figure S1b), and *Agaricia* samples averaged 3.7 million reads (figure
S2b). A maximum likelihood phylogenetic tree of the overall dataset, including outgroup genera
(based on 37,528 nextRAD loci), confirmed that all of our ingroup specimens belonged to either
the *Leptoseris* or *Agaricia* clade (figure 1d-e). One exception was a group of five presumed
"*Leptoseris*" specimens from the Red Sea that grouped with the *Pachyseris* outgroup, and were
later identified as *Pachyseris inatessa* (figure 1e). After separating the datasets and filtering, the
Leptoseris dataset consisted of 15,250 nextRAD loci and 100,270 SNPs, and the *Agaricia* dataset
of 19,902 nextRAD loci and 221,031 SNPs. For the genealogical concordance analysis, we
inferred phylogenies for each single loci containing data for ε10% of the samples, resulting in
10,317 and 30,650 nextRAD loci for *Leptoseris* and *Agaricia*, respectively (figure S3c-d, figure
S4c-d). The phylogenetic analyses supported the current taxonomic species within both genera but
exposed substantial genetic substructure with at least 13 and 12 molecular clades observed within
our focal *Leptoseris* and *Agaricia* species, respectively.

For the genus *Leptoseris*, maximum likelihood phylogenetic inference using both RAxML-ng and
IQTree recovered extremely similar topologies consisting of many well-supported clades,
particularly at deeper nodes (RAxML-ng: 6.92-100%, bootstrap range; IQTree: 35-100%; figure
S3a-b). Gene (gCF) and site concordance factors (sCF) across the IQTree topology were lower
and variable, particularly near the shallow nodes of the tree (gCF: $7.0 \pm 12.2\%$, mean $\pm$ SD; sCF:
$42.1 \pm 16.8\%$; figure S3c-d), indicating the presence of discordant signal across loci and sites (most
likely attributed to the short length of nextRAD loci). The species tree analysis using Tetrad
recovered similar clades as those identified with ML, although the relationship among clades
differs and their support varies (1-100%; $42.3 \pm 30.4\%$; figure S5a). Clustering analysis (*de novo*
DAPC) recovered clusters that corresponded with nodes at varying evolutionary depths in the ML
and species tree topologies, with clusters identified at higher numbers of K corresponding with the
same clades found in common among phylogenetic analyses (figure S6a). The exception was one
group observed in a *L. glabra* clade (with mixed assignment to a spurious cluster that lacked
individuals fully assigned to it).

While the clades recovered across methods for *Leptoseris* were largely composed of a single
taxonomic species, all of the taxonomic focal species were represented by multiple clades. In

several cases, these different clades were separated by relatively deep nodes (e.g. for *L. scabra* and *L. mycetoseroides*; figure 2a), whereas others represent substructuring within major clades (based on the tree topologies and the signatures of admixture at higher values of K from DAPC analysis; figure S6a). Samples belonging to the two divergent clades of *L. mycetoseroides* corresponded to respectively mesophotic samples from WCS (Western Coral Sea; 5 out of 6) and shallow water samples from Hawaii/USOMI (United States Minor Outlying Islands; 5 out of 6) (figure 2a). The former clade was identified as *L.* cf. *mycetoseroides* due to morphological variations from typical *L. mycetoseroides* (Pichon and Dinesen, personal observation). Similarly, while the two clades of *L. scabra* occurred sympatrically on reefs of the WCS and GBR (Great Barrier Reef), these clades were differentially characterised by depth, one clade was composed of a mix of shallow (10 - 20 m; n = 13) and mesophotic (40 - 60 m; n = 19) specimens and the other exclusively composed of mesophotic specimens (40 - 80 m, n = 14). There were three clades composed of *L. glabra* specimens; two had a wide geographic distribution (including Australia and Red Sea) and exclusively contain specimens from mesophotic depths (n = 4 and 6), and the third clade (middle *L. glabra* clade in figure 2a; also observed in Australia and Red Sea) represented a mix of shallow (10 - 20 m; n = 8) and mesophotic specimens (40 - 60 m, n = 10). *L. hawaiiensis* specimens were observed across four clades (figure 2a). One clade comprised samples from primarily lower mesophotic depths on the GBR and WCS, but matched a specimen identified as *Leptoseris* sp. 1 from Hawaii [17]. A second clade had widespread representation containing specimens from Australia, USOMI, and the Red Sea, and originating from depths down to 60 m. In contrast, the third *L. hawaiiensis* clade contained almost exclusively specimens from below 120 m depth in Hawaii, with a closely related individual originating from the GBR collected at 124 m depth. The fourth clade was identified as *L.* cf. *hawaiiensis* due to morphological variations from typical *L. hawaiiensis* (Pichon and Dinesen, personal observation), with samples from 60 m in the WCS. There were also five *Leptoseris* sp. that could not be identified further: one of them (87 m depth; WCS) was related to the former *L.* (cf.) *hawaiiensis* clades. Two samples (40 - 60 m depth; WCS) grouped within two distinct *L. glabra* clades. The remaining two *Leptoseris* sp. originating from shallow depths (9 and 22 m, Hawaii) grouped with the shallow *L. hawaiiensis* and *L. mycetoseroides* clades from Hawaii. The analyses also included a single specimen of *L. amitoriensis* from the Red Sea, which grouped closest to the clade containing *Leptoseris* sp. 1 (figure 2a).

For the genus *Agaricia*, maximum likelihood phylogenetic inference using both RAxML-ng and IQTree also recovered very similar topologies consisting of several well-supported clades (RAxML-ng: 21.4 - 100%, bootstrap range; 80.3 ± 23.6%, mean ± SD; IQTree: 74 - 100%; 98.1 ± 5.3%; figure S4). Both phylogenies supported a major division of the currently acknowledged *Agaricia* genus into two major clades, representing the focal species, *A. grahamae*, *A. fragilis*, *A. lamarcki*, and *A. undata*, and the taxonomic species, *A. agaricites, A. humilis,* and *A. tenuifolia* [39] (figure 2b). Once again, the concordance factor values (gCF and sCF) across the IQTree topology were low and variable, particularly near the shallow nodes of the tree (gCF: 7.2 ± 12.4%, mean ± SD; sCF: 51.6 ± 22.2%; figure S4). The species tree analysis (Tetrad) recovered the same topology, consisting of the same genetic clades as those identified during ML analyses, with high statistical support, especially for the major clades (74 - 100%; 98.1 ± 5.3%; figure S5b). Clustering analysis (DAPC) recovered clusters that corresponded with nodes at varying evolutionary depths in the ML and species tree topologies, with the higher levels of K corresponding with the same clades found in the phylogenetic analyses (figure S6b).

Although each of the observed clades in the *Agaricia* phylogenies was composed of specimens of a single taxonomic species, all four taxonomic focal species were represented across two or more monophyletic subclades, with *A. grahamae* exhibiting further structuring into subclades that were supported by the DAPC clustering analyses (figure S6b). *A. grahamae* specimens were represented by four major sympatric clades (figure 2b). One clade (with substantial substructure) corresponded to the lower-mesophotic (>60 m) specimens from San Andrés (n = 7) and a single specimen from Bonaire upper-mesophotic reefs (figure 2b). The remaining *A. grahamae* clades consisted mostly of specimens from upper-mesophotic depths (<60 m) from Curaçao and Bonaire (n = 14), with one of those clades containing only two samples, one from Colombia and one from San Andrés. *A. lamarcki* consisted of two sympatrically occurring clades with one clade exclusively representing mesophotic specimens (n = 6) and the other mostly shallow water specimens (n = 6). The *A. undata* clade was further split into two clades corresponding to two sampling regions; the Southern Caribbean (Curaçao) containing lower-mesophotic specimens (n = 4) and South-western Caribbean (Colombia and San Andrés) consisting of upper-mesophotic specimens (n = 6), although with DAPC clustering showing admixture between clusters (figure S6b). Three

specimens identified as *Agaricia* sp. from Colombia formed a separate clade grouping with *A.*
fragilis from Bermuda (n = 6), with no further differentiation observed between shallow and upper
mesophotic specimens of the latter. Specimens from *A. agaricites*, *A. tenuifolia*, and *A. humilis*
formed a separate, highly divergent clade from the other *Agaricia* spp. Within this clade, *A. humilis*
is the most divergent, with the *A. tenuifolia* from Colombia (n = 4) and *A. agaricites* from Curaçao
and Bonaire (n = 3) forming closely related groups and exhibiting high levels of admixture in the
DAPC clustering analyses (figure S6b).
320

Comparison with traditional sequence markers

Phylogenetic analysis of the mitochondrial sequence marker *cox*1-1-rRNA using a ML approach
(RAxML-ng) for a subset of the Australian *Leptoseris* specimens (n = 101, figure S7) including
available sequences from Luck *et al.* [17], identified several major molecular clades with high
bootstrap support that exceeded the number of focal taxonomic species (figure S7). Even though
many of these clades consisted of specimens from a single taxonomic species, substantial
polyphyly was observed with clades consisting of representatives from multiple taxonomic taxa.
These clades do not appear to be described by geography nor bathymetric origins, with the
exception of two divergent *L. scabra* clades consisting only of mesophotic representatives (1st: 40
- 101 m, n = 19, 2nd: 40 - 79 m, n = 7), as well as three *L. hawaiiensis* clades containing only *L.*
hawaiiensis specimens mostly from mesophotic depths (1st: 60 m, n = 3; 2nd: 40 - 82 m, n = 8/9,
3rd: 40 - 60 m, n = 16, including an unidentified sample from 124 m, figure S7). For *Agaricia*, a
comparison with available sequences (n = 61) from previous studies [11, 71] and additional
sequences (n = 9) using traditional sequence markers (*cox*1-1-rRNA) distinguished between three
major clades, one consisting of *A. grahamae*, *A. fragilis*, *A. lamarcki*, one consisting of *A. undata*,
and another one composed by *A. agaricites*, and *A. humilis*. However, this marker did not
consistently discriminate between taxonomic species within these two groups (figure S8).

Figure 2. *Leptoseris* and *Agaricia* (a) Phylogenetic tree (RAxML-ng) of the genus *Leptoseris* based on 15,250 concatenated nextRAD loci. **(b)** Phylogenetic tree (RAxML-ng) of the genus *Agaricia* based on 19,902 concatenated nextRAD loci. Colours across the trees represent the different identified clades (across phylogenetic and clustering methods) using blue, turquoise, purple, and green colours to represent the taxonomic species and shades of the different subclades. Columns next to the tree correspond to the taxonomic identification, sampling location, and sampling depth. The summary column on the right outlines the consensus taxonomic identification of each clade, and pie charts summarising the sampling depths and locations.

DISCUSSION

Mesophotic coral ecosystems have gathered great scientific interest over the past decade, with numerous assessments evaluating the similarities and differences of these spatially extensive biological communities as compared to their shallow-water counterparts. From these, it has become clear that mesophotic communities become increasingly distinct with depth and can host unique and diverse species assemblages [7,15,72]. However, despite the growing number of accounts of depth-differentiated populations within scleractinian coral species (e.g., [3,73,74]), most scleractinian coral species at mesophotic depths are characterised as deep-generalists, with only a handful of reef-building scleractinian species recognised as mesophotic-specialists [16]. Through an extensive phylogenomic assessment of plating corals belonging to two dominant mesophotic genera in the Indo-Pacific and Caribbean (*Leptoseris* and *Agaricia*), we uncover substantial undescribed diversity with (1) assumed depth-generalists representing multiple depth-associated taxa, and (2) deep-specialist species consisting of multiple sympatric taxa. Overall, the results indicate that coral communities at mesophotic depths are more speciose and likely more specialised than currently acknowledged, urging for both systematic and ecological studies to capture and better understand this diversity.

Phylogenomic insights into *Leptoseris* diversity

The observed phylogenomic patterns based on nextRAD data confirm that *Leptoseris* represent a taxonomically diverse group and comprise several highly divergent clades. When considering assignments based on the current *Leptoseris* taxonomy [18,32], species appear to be polyphyletic and often even separated by deep nodes (figure 2a), similar to previous studies using a

mitochondrial intergenic spacer region [13,17,34]. Although the mitochondrial marker seemed

initially promising for species delineation based on specimens solely from Hawaii [13,17], it did

show pervasive polyphyly that extended across both *Leptoseris* and *Agaricia* genera [17], and a

lack of genetic variation across morphologically divergent taxonomic species when applied in a

different geographic region [34]. Using the same marker, we obtained similar results for specimens

from Australia, where some of the taxonomic diversity is captured though not consistently, with

taxonomic species spread widely across the tree (figure S7). In contrast, the reduced representation

data shows much more consistent phylogenetic patterns, with all recovered clades consisting

almost exclusively of a single taxonomic species (figure 2a). The patterns corroborate the current

taxonomy based on morphological differences, but with the increased genomic resolution, sample

sizes and geographic range, also unveiling additional diversity associated with specific geographic

regions and bathymetric ranges. Given this congruence with taxonomic species and spatial

distributions, the observed phylogenomic patterns are expected to more closely reflect the

evolutionary relationships within this genus and demonstrate the resolving power of reduced

representation methods.

Within the genus *Leptoseris*, the major split in *L. mycetoseroides* seems to correspond with both

geography and depth (figure 2a). Given that the type specimen originates from the Marshall Islands

(adjacent to USOMI) [75], there are no junior synonyms reported for this species [75], and several

specimens from East Australia were noted as morphologically distinct, it is possible that the

Hawaii/USOMI clade represents *L. mycetoseroides* with the mesophotic specimens in the East

Australian clade representing an undescribed species. One of the clades identified for *L.*

hawaiiensis consisted of predominantly lower mesophotic specimens from the WCS, and matched

with a *Leptoseris* "sp. 1" specimen from Hawaii, where it was identified as a putative new species

based on its distinct micromorphology [17]. In the *cox*1-1-rRNA phylogeny, one of our specimens

from this clade also grouped with the *Leptoseris* "sp. 1" clade from Luck *et al.* [17], although the

other specimens from the same nextRAD clade were widely spread across the tree (figure S7).

This predominantly mesophotic clade is present in Australian waters and may be geographically

widespread, although further investigations are warranted. The other three *L. hawaiiensis* clades

each showed different depth distributions, with one from predominantly >100 m depth in Hawaii,

and another representing morphologically atypical specimens from 60 m depth in Australia.

Related to these clades but branching separately was a single specimen collected at the GBR at
124 m, representing the deepest published report of a zooxanthellate coral collected from the GBR
[30], and indicating that *Leptoseris* clades occurring at the lower boundaries of mesophotic depths
represent uniquely adapted species. We observed a similar partitioning into depth-associated
clades for *L. glabra* and *L. scabra*, with a further subdivision into additional geographically
sympatric clades for the latter. As reported by Luck *et al.* [17], the deeper clade of *L. scabra* is
very divergent compared to the other *Leptoseris* taxa (although not resulting in generic polyphyly),
and with *cox*1-1-rRNA sequences matching those from Hawaii. Detailed morphological
characterization of the specimens (beyond identification according to currently acknowledged
taxonomic species) was beyond the scope of the current study, but is the focus of ongoing work
that aims to determine which of the exposed taxa are cryptic versus morphologically differentiated.
412
Phylogenomic insights into *Agaricia* diversity
For the genus *Agaricia*, we observed a split into two major clades corresponding to species with
deeper (*A. grahamae*, *A. fragilis*, *A. lamarcki*, *A. undata*) and shallower (*A. agaricites*, *A. humilis*,
A. tenuifolia) distributions (figure 2b). This division was also observed using the *cox*1-1-rRNA
region (figure S8) as well as other mitochondrial markers [11,27,71,76-78], although these markers
have lacked the resolution to consistently discern the established taxonomic species within the
clades. The division into two major clades is further corroborated by a split in gross morphology
(unifacial versus bifacial colonies), microskeletal characteristics (including wall thickness and
septocostae orientation; [79]), and genetic relatedness among symbiont associations within each
major clade [27]. Based on a morphological analysis of modern and fossil representatives, Stemann
[79] proposed these should represent two separate genera: *Agaricia* and *Undaria* [80]. Recent
differences in spatial genetic structure observed across taxa further suggest major differences in
reproductive strategies between these major clades [53]. Given the major phylogenetic divergence
observed in this study (figure 2b), and further corroborated by the major locus dropout between
clades (figure S2), we argue that the shallower and deeper *Agaricia* indeed represent two
genetically, morphologically, and biologically divergent groups that might require generic
reassignment.
430

The phylogenetic analyses supported the current taxonomic species within *Agaricia* but revealed
substantial genetic substructure with at least 12 molecular clades observed within our four focal
Agaricia species (figure 2b). This structure was consistently recovered across phylogenetic and
clustering methods and often corresponded to different depths and/or geographic regions,
indicating the role of both environmental and geographical (i.e., allopatric) contributors to the
underlying diversification processes. Our analyses included four focal species previously assessed
in population genomic studies with varying levels of intraspecific genetic structure [3,37,53].
Through a combined phylogenomic analysis, we were able to assess this genetic structuring in the
context of interspecific variation. For example, despite the genome-wide differentiation observed
for shallow and deep *A. fragilis* populations in Bermuda [3], representative specimens of those
populations formed a single clade here (i.e., under a phylogenomic framework), indicating that
these reflect the earlier stages of the divergence continuum. In contrast, the *A. lamarcki* samples
clearly separated into two distinct clades associated with predominantly shallower (15 m) and
upper mesophotic (50 m) depths (figure 2b) despite their geographically sympatric distribution
and ongoing low levels of gene flow [53], and clearly represent distinct evolutionary units.
Similarly, we observed four distinct clades of *A. grahamae* occurring sympatrically in Curaçao
and Bonaire with varying levels of divergence (figure 2b), and while not partitioned by depth, it
indicates undescribed mesophotic taxa that are currently not accounted for. The two clades
observed for *Agaricia undata* corresponded with geography (Curaçao vs Colombia and San
Andrés), whereas several Colombian specimens that could not be reliably identified down to
species through morphological assessment (*Agaricia* sp. clade) grouped with *A. fragilis* from
Bermuda, indicating the presence of this or an undescribed related species in the Southern
Caribbean. In the aforementioned "*Undaria*" clade, *A. humilis* formed a separate clade from *A.*
agaricites and *A. tenuifolia* specimens, corroborating their ecological and reproductive differences
[36]. The separation between *A. agaricites* and *A. tenuifolia* was confounded by distinct
geographic origins for the specimens; however, the lack of support in clustering analyses indicates
that these morphologically similar species warrant further taxonomic investigation.
458

Evolutionary patterns and remaining challenges
As a result of this study, we can now begin to directly compare and contrast the present diversity
of these genera and to hypothesise the existence of associated generative processes. An evident

pattern we observe across both genera is the depth-associated divergence within several focal
species (e.g., *L. scabra*, *L. hawaiiensis*, or *A. lamarcki*). Depth is an important contributor to the
divergence of marine species [73,81,82], with several well-studied examples in scleractinian corals
(e.g., [83-87]). Our results corroborate the expectation that depth has led to divergent species
associated with mesophotic depths [82], and that the underestimated diversity of mesophotic-
specialist species is likely to be a consequence of logistical and taxonomic challenges [16].
However, depth does not immediately explain the sympatric taxa we observed within mesophotic-
specialist species (e.g., *A. grahamae* and the "deep" *L. scabra* clade; lacking clear depth
differences). Their ecological differentiation may be related to finer scale environmental
conditions or, indeed, other factors and should be assessed within a spatially explicit framework
targeted to microhabitat characterisation [88].

Another clear pattern we observe here is the higher levels of taxonomic/phylogenetic discordance
in (often sympatrically-occurring) genetic groups in *Leptoseris*, relative to *Agaricia.* This
difference might be the result of *Leptoseris* having a larger geographic distribution spanning a
variety of ecoregions (Indo-Pacific and Caribbean; [89,90]) in contrast to the relatively restricted
geographic distribution of *Agaricia* in the Western Atlantic and Caribbean [89]. In addition, the
fossil record of *Leptoseris* dates back to the Oligocene (~23 Ma years, [91]), while *Agaricia* is a
younger genus, dating from the Neogene ~12 Ma years [92], and likely diversified over the last ~3
Ma years following the closure of the Central American Seaway [89,92]. These differences suggest
increased opportunities for ecological diversification in *Leptoseris* (compared to *Agaricia*), while
signatures of admixture (figure S6a) suggest a possible history of hybridization and reticulation
(i.e., the process of genetic lineages both merging and/or diverging through time; [93]) among its
lineages. It is important to note, however, that the number of loci and sites supporting some of
these polyphyletic patterns are low (figure S3c), thus additional sequence data (e.g., longer/more
informative loci, whole-genome data) might reveal additional patterns within this genus (see
below).

The reduced representation sequencing (nextRAD) data across multiple analytical approaches,
consistently recovered groupings of individuals (i.e., clades and/or clusters) coherent with
biological, ecological, or morphological evidence. Compared to traditional sequence markers, our

genome-wide sequencing approach resulted in a higher resolving power, confirming the status of current morphologically and ecologically divergent taxonomic species, and allowing us to gain a better insight into the evolutionary history of members of both genera. Despite the consistent topology recovered across approaches, the support for these underlying phylogenies did vary across analytical methods. For example, despite the strong bootstrap support recovered across all ML phylogenies, the gene- and site-concordance analysis revealed variable values in both datasets, indicating, on average, a low fraction of gene trees and a moderate fraction of alignment sites supporting each topology (figure S3 and figure S4). This discrepancy between methods is often observed across different RAD-seq approaches, in most cases, due to the short length of each locus and its associated phylogenetic information [94-97]. Targeted sequence-capture approaches focused on ultra-conserved elements, or exons, can potentially represent an alternative to recover greater concordance across loci. Because conserved regions are less affected by demographic events, and next to these regions, more variable sites can be considered side by side, these genomic regions provide a more holistic view of individual evolutionary histories [98-99]. Compared to RAD-seq, however, the generally greater cost of such sequence-capture approaches often translates to smaller sample sizes and less replication. Ultimately, whole-genome resequencing of representatives across a wide range of depths and geographies from our study will enable us to identify the loci involved in the diversification of *Leptoseris* and *Agaricia* and help assess divergence histories through introgression events.

Conclusions

Our results shed light on the diversity of two key genera of mesophotic ecosystems, *Leptoseris* and *Agaricia*. Using genome-wide sequencing data in a phylogenomic framework, we observed that genomic data corroborate current morpho-taxonomic criteria, but also exposed substantial undescribed diversity associated with mesophotic depths. Distinguishing where these different taxa sit along the speciation continuum remains challenging, particularly as traditional species delimitation methods are known to oversplit when using genomic data [100]. Nonetheless, current taxonomic species were observed to comprise highly divergent clades (e.g., *L. scabra*, *L. mycetoseroides*) or sympatrically occurring but geographically widespread subclades (e.g., *L. glabra*, *A. grahamae*) across phylogenetic and clustering methods, indicating that a reasonable extent of reproductive isolation has evolved and that most of these clades represent distinct species.

Further integrative taxonomic studies are currently being developed to formally describe the uncovered species diversity, and verify whether these taxa are morphologically cryptic, differentiated and/or potentially align with junior synonyms. Overall, our study highlights how our perception of mesophotic coral ecosystems is affected by our shallow knowledge bias, and that studying the ecology and evolution of this newly exposed mesophotic biodiversity should be a priority in order to advance our understanding of these ecosystems.

REFERENCES

1. Pyle RL, Copus JM. 2019 Mesophotic coral ecosystems: introduction and overview. *Coral Reefs of the World*. 3–27. (doi:10.1007/978-3-319-92735-0_1)

2. Bongaerts P, Ridgway T, Sampayo EM, Hoegh-Guldberg O. 2010 Assessing the 'deep reef refugia' hypothesis: focus on Caribbean reefs. *Coral Reefs* **29**, 309–327. (doi:10.1007/s00338-009-0581-x)

3. Bongaerts P, Riginos C, Brunner R, Englebert N, Smith SR, Hoegh-Guldberg O. 2017 Deep reefs are not universal refuges: Reseeding potential varies among coral species. *Sci. Adv.* **3**. (doi:10.1126/sciadv.1602373)

4. Holstein DM, Smith TB, Gyory J, Paris CB. 2015 Fertile fathoms: Deep reproductive refugia for threatened shallow corals. *Sci. Rep.* **5**. (doi:10.1038/srep12407)

5. Holstein DM, Paris CB, Vaz AC, Smith TB. 2015 Modeling vertical coral connectivity and mesophotic refugia. *Coral Reefs* **35**, 23–37. (doi:10.1007/s00338-015-1339-2)

6. Pyle RL. 2000 Assessing undiscovered fish biodiversity on deep coral reefs using advanced self-contained diving technology. *Mar. Technol. Soc. J.* **34**, 82–91. (doi:10.4031/mtsj.34.4.11)

7. Rocha LA, Pinheiro HT, Shepherd B, Papastamatiou YP, Luiz OJ, Pyle RL, Bongaerts P.
2018 Mesophotic coral ecosystems are threatened and ecologically distinct from shallow
water reefs. *Science* **361**, 281–284. (doi:10.1126/science.aaq1614)

8. Eyal G, Pinheiro HT. 2020 Mesophotic ecosystems: The link between shallow and deep-
sea habitats. *Diversity* **12**, 411. (doi:10.3390/d12110411)

9. Pinheiro HT, Eyal G, Shepherd B, Rocha LA. 2019 Ecological insights from environmental
disturbances in mesophotic coral ecosystems. *Ecosphere* **10**, e02666.
(doi:10.1002/ecs2.2666)

10. Lesser MP, Slattery M, Laverick JH, Macartney KJ, Bridge TC. 2019 Global community
breaks at 60 m on mesophotic coral reefs. *Global Ecol. Biogeogr*. **28**, 1403–1416.
(doi:10.1111/geb.12940)

11. Bongaerts P, Frade PR, Hay KB, Englebert N, Latijnhouwers KRW, Bak RPM, Vermeij
MJA, Hoegh-Guldberg O. 2015 Deep down on a Caribbean reef: lower mesophotic depths
harbor a specialized coral-endosymbiont community. *Sci. Rep*. **5**. (doi:10.1038/srep07652)

12. Kahng S, Hochberg E, Apprill A, Wagner D, Luck D, Perez D, Bidigare R. 2012 Efficient
light harvesting in deep-water zooxanthellate corals. *Mar. Ecol. Prog. Ser*. **455**, 65–77.
(doi:10.3354/meps09657)

13. Pochon X, Forsman ZH, Spalding HL, Padilla-Gamiño JL, Smith CM, Gates RD. 2015
Depth specialization in mesophotic corals (*Leptoseris* spp.) and associated algal symbionts
in Hawai'i. *R. Soc. Open Sci*. 2, 140351. (doi:10.1098/rsos.140351)

14. Englebert N, Bongaerts P, Muir PR, Hay KB, Pichon M, Hoegh-Guldberg O. 2017 Lower
mesophotic coral communities (60-125 m Depth) of the northern Great Barrier Reef and
Coral Sea. *PLoS ONE* **12**, e0170336. (doi:10.1371/journal.pone.0170336)

15. Muir PR, Wallace CC, Pichon M, Bongaerts P. 2018 High species richness and lineage
diversity of reef corals in the mesophotic zone. *Proc. R. Soc. B* **285**, 20181987.
(doi:10.1098/rspb.2018.1987)

16. Muir PR, Pichon M. 2019 Biodiversity of reef-Building, scleractinian corals. In
Mesophotic coral ecosystems, *Volume 12: Corals Reefs of the World* (eds. Loya Y, Puglise
KA, Bridge T), 589–620. Charm: Springer. (doi:10.1007/978-3-319-92735-0_33)

17. Luck DG, Forsman ZH, Toonen RJ, Leicht SJ, Kahng SE. 2013 Polyphyly and hidden
species among Hawai'i's dominant mesophotic coral genera, *Leptoseris* and *Pavona*
(Scleractinia: Agariciidae). *PeerJ*. **1**, e132. (doi:10.7717/peerj.132)

18. Dinesen ZD. 1980 A revision of the coral genus *Leptoseris* (Scleractinia: Fungiina:
Agariciidae). Memoirs of the Queensland Museum **20**,181–235

19. van Oppen MJH, McDonald BJ, Willis B, Miller DJ. 2001 The evolutionary history of the
coral genus *Acropora* (Scleractinia, Cnidaria) based on a mitochondrial and a nuclear
marker: reticulation, incomplete lineage sorting, or morphological convergence? *Mol. Biol.*
Evol. **18**, 1315–1329. (doi:10.1093/oxfordjournals.molbev.a003916)

20. Forsman ZH, Barshis DJ, Hunter CL, Toonen RJ. 2009 Shape-shifting corals: Molecular
markers show morphology is evolutionarily plastic in Porites. *BMC Evol. Biol.* **9**, 45.
(doi:10.1186/1471-2148-9-45)

21. Vollmer SV, Palumbi SR. 2004 Testing the utility of internally transcribed spacer
sequences in coral phylogenetics. *Mol. Ecol.* **13**, 2763–2772. (doi:10.1111/j.1365-
294x.2004.02265.x)

22. McFadden CS, Sanchez JA, France SC. 2010 Molecular phylogenetic insights into the
evolution of Octocorallia: a review. *Integr. Comp. Biol.* **50**, 389–410.
(doi:10.1093/icb/icq056)

23. Kühlmann DHH. 1983 Composition and ecology of deep-water coral associations. *Helgolander Meeresunters* **36**, 183–204. (doi:10.1007/bf01983856)

24. Kahng SE, Garcia-Sais JR, Spalding HL, Brokovich E, Wagner D, Weil E, Hinderstein L, Toonen RJ. 2010 Community ecology of mesophotic coral reef ecosystems. *Coral Reefs* **29**, 255–275. (doi:10.1007/s00338-010-0593-6)

25. Kramer N, Tamir R, Ben-Zvi O, Jacques SL, Loya Y, Wangpraseurt D. 2021 Efficient light-harvesting of mesophotic corals is facilitated by coral optical traits. *Funct. Ecol.* **36**, 406–418. (doi:10.1111/1365-2435.13948)

26. Kramer N, Tamir R, Eyal G, Loya Y. 2020 Coral morphology portrays the spatial distribution and population size-structure along a 5–100 m depth gradient. *Front. Mar. Sci.* **7**. (doi:10.3389/fmars.2020.00615)

27. Bongaerts P *et al*. 2013 Sharing the slope: depth partitioning of agariciid corals and associated *Symbiodinium* across shallow and mesophotic habitats (2-60 m) on a Caribbean reef. *BMC Evol. Biol.* **13**, 205. (doi:10.1186/1471-2148-13-205)

28. Edinger EN, Risk MJ. 1995 Preferential survivorship of brooding corals in a regional extinction. *Paleobiology* **21**, 200–219. (doi:10.1017/s0094837300013208)

29. Rouzé H *et al*. 2021 Symbiotic associations of the deepest recorded photosynthetic scleractinian coral (172 m depth). *ISME J.* **15**, 1564–1568. (doi:10.1038/s41396-020-00857-y)

30. Englebert N, Bongaerts P, Muir P, Hay KB, Hoegh-Guldberg O. 2014 Deepest zooxanthellate corals of the Great Barrier Reef and Coral Sea. *Mar. Biodiv.* **45**, 1–2. (doi:10.1007/s12526-014-0221-8)

31. Vaughan TW. 1907 Recent Madreporaria of the Hawaiian Islands and Laysan. US National Museum Bulletin **59**, 1-427

32. Veron JEN, Pichon M. 1979 Scleractinia of Eastern Australia, part III: Families Agariciidae, Siderastreidae, Fungiidae, Oculinidae, Merulinidae, Mussidae, Pectiniidae, Caryophylliidae, Dendrophylliidae. Australian Government Publishing Service. Canberra **4**, 40-42, 44

33. Wells JW. 1954 Bikini and nearby atols: 2. Oceanography (biologic). Recent corals of the Marshall Islands. *Geological Survey Professional Paper* **260-1**, 94-185

34. Terraneo TI, Arrigoni R, Benzoni F, Tietbohl MD, Berumen ML. 2017 Exploring the genetic diversity of shallow-water Agariciidae (Cnidaria: Anthozoa) from the Saudi Arabian Red Sea. *Mar. Biodiv*. **47**, 1065–1078. (doi:10.1007/s12526-017-0722-3)

35. Eyal G, Tamir R, Kramer N, Eyal-Shaham L, Loya Y. 2019 The Red Sea: Israel. In *Mesophotic coral ecosystems*, *Volume 12: Corals Reefs of the World* (eds. Loya Y, Puglise KA, Bridge T), pp. 199-214. Charm: Springer (doi:10.1007/978-3-319-92735-0_11)

36. Van Moorsel G. 1983 Reproductive strategies in two closely related stony corals (*Agaricia*, Scleractinia). *Mar. Ecol. Prog. Ser.* **13**, 273–283. (doi:10.3354/meps013273)

37. Gonzalez-Zapata FL, Bongaerts P, Ramírez-Portilla C, Adu-Oppong B, Walljasper G, Reyes A, Sanchez JA. 2018 Holobiont diversity in a reef-building coral over its entire depth range in the mesophotic zone. *Front. Mar. Sci.* **5**. (doi:10.3389/fmars.2018.00029)

38. Milne Edwards H, Haime J. 1849 Mémoire sur les Polypes appartenant à la famille des Oculinides, au groupe intermédiaire des Pseudastréides et à la famille des Fongides. Comptes rendus hebdomadaires des séances de l'Académie des sciences, Paris **29**, 67-73

677   39. Dana JD. 1846-1849 Zoophytes. United States Exploring Expedition during the years
678   1838-1842. Lea and Blanchard, Philadelphia **7**, 1-740
679

680   40. Wells JW. 1973 New and old scleractinian corals from Jamaica. *Bull. Mar. Sci.* **23**, 16–58
681

682   41. Ellis J, Solander D. 1786 The Natural History of many curious and uncommon Zoophytes,
683   collected from various parts of the Globe. 1-206
684

685   42. Mazzei EF *et al.* 2017 Newly discovered reefs in the southern Abrolhos Bank, Brazil:
686   Anthropogenic impacts and urgent conservation needs. *Mar. Pollut. Bull.* **114**, 123–133.
687   (doi:10.1016/j.marpolbul.2016.08.059)
688

689   43. Reed JK. 1985 Deepest distribution of Atlantic hermatypic corals discovered in the
690   Bahamas. *Proceedings of the Fifth International Coral Reef Congress* **6**, 249-254
691

692   44. Verrill AE. 1901 Variations and nomenclature of Bermudian, West Indian and Brazilian
693   reef corals, with notes on various Indo-Pacific corals. Transactions of the Connecticut
694   Academy of Arts and Sciences **11**, 63-168
695

696   45. Linnaeus C. 1758 Systema Naturae per regna tria naturae, secundum classes, ordines,
697   genera, species, cum characteribus, differentiis, synonymis, locis **1**, 824
698

699   46. Shearer TL, van Oppen MJH, Romano SL, Wörheide G. 2002 Slow mitochondrial DNA
700   sequence evolution in the Anthozoa (Cnidaria). *Mol. Ecol.* **11**, 2475–2487.
701   (doi:10.1046/j.1365-294x.2002.01652.x)
702

703   47. Terraneo TI, Benzoni F, Arrigoni R, Berumen ML. 2016 Species delimitation in the coral
704   genus *Goniopora* (Scleractinia, Poritidae) from the Saudi Arabian Red Sea. *Mol.*
705   *Phylogenet. Evol.* **102**, 278–294. (doi:10.1016/j.ympev.2016.06.003)
706

707 48. Herrera S, Shank TM. 2016 RAD sequencing enables unprecedented phylogenetic
708  resolution and objective species delimitation in recalcitrant divergent taxa. *Mol.*
709  *Phylogenet. Evol.* **100**, 70–79. (doi:10.1016/j.ympev.2016.03.010)

711 49. Rosser NL, Thomas L, Stankowski S, Richards ZT, Kennington WJ, Johnson MS. 2017
712  Phylogenomics provides new insight into evolutionary relationships and genealogical
713  discordance in the reef-building coral genus *Acropora*. *Proc. R. Soc. B* **284**, 2016-2182.
714  (doi:10.1098/rspb.2016.2182)

716 50. Quattrini AM, Wu T, Soong K, Jeng M-S, Benayahu Y, McFadden CS. 2019 A next
717  generation approach to species delimitation reveals the role of hybridization in a cryptic
718  species complex of corals. *BMC Evol. Biol.* **19**. (doi:10.1186/s12862-019-1427-y)

720 51. Cunha RL, Forsman ZH, Belderok R, Knapp ISS, Castilho R, Toonen RJ. 2019 Rare coral
721  under the genomic microscope: timing and relationships among Hawaiian *Montipora*.
722  *BMC Evol. Biol.* **19**. (doi:10.1186/s12862-019-1476-2)

724 52. Arrigoni R *et al.* 2020 Towards a rigorous species delimitation framework for scleractinian
725  corals based on RAD sequencing: the case study of *Leptastrea* from the Indo-Pacific. *Coral*
726  *Reefs* **39**, 1001–1025. (doi:10.1007/s00338-020-01924-8)

728 53. Prata KE, Riginos C, Gutenkunst RN, Latijnhouwers KRW, Sánchez JA, Englebert N, Hay
729  KB, Bongaerts P. 2022 Deep connections: Divergence histories with gene flow in
730  mesophotic *Agaricia* corals. *Mol. Ecol.* **31**, 2511–2527. (doi:10.1111/mec.16391)

732 54. Hammerman NM, Rivera-Vicens RE, Galaska MP, Weil E, Appledoorn RS, Alfaro M,
733  Schizas NV. 2017 Population connectivity of the plating coral *Agaricia lamarcki* from
734  southwest Puerto Rico. *Coral Reefs* **37**, 183–191. (doi:10.1007/s00338-017-1646-x)

736 55. Bongaerts P *et al.* 2021 Morphological stasis masks ecologically divergent coral species
737  on tropical reefs. *Curr. Biol.* **31**, 2286-2298.e8. (doi:10.1016/j.cub.2021.03.028)

56. Russello MA, Waterhouse MD, Etter PD, Johnson EA. 2015 From promise to practice: pairing non-invasive sampling with genomics in conservation. *PeerJ*. **3**, e1106. (doi:10.7717/peerj.1106)

57. Eaton DAR, Overcast I. 2020 Ipyrad: Interactive assembly and analysis of RADseq datasets. *Bioinformatics* **36**, 2592–2594. (doi:10.1093/bioinformatics/btz966)

58. Aranda M *et al.* 2016 Genomes of coral dinoflagellate symbionts highlight evolutionary adaptations conducive to a symbiotic lifestyle. *Sci. Rep*. **6**. (doi:10.1038/srep39734)

59. Shoguchi E, Tanaka M, Shinzato C, Kawashima T, Satoh N. 2013 A genome-wide survey of photoreceptor and circadian genes in the coral, *Acropora digitifera*. *Gene* **515**, 426–431. (doi:10.1016/j.gene.2012.12.038)

60. Liu H *et al.* 2018 *Symbiodinium* genomes reveal adaptive evolution of functions related to coral-dinoflagellate symbiosis. *Commun. Biol.* **1**. (doi:10.1038/s42003-018-0098-3)

61. Dougan KE *et al.* 2022 Whole-genome duplication in an algal symbiont serendipitously confers thermal tolerance to corals. *bioRxiv* (doi:10.1101/2022.04.10.487810)

62. Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A. 2019 RAxML-NG: a fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* **35**, 4453–4455. (doi:10.1093/bioinformatics/btz305)

63. Darriba D, Posada D, Kozlov AM, Stamatakis A, Morel B, Flouri T. 2019 ModelTest-NG: A new and scalable tool for the selection of DNA and protein evolutionary models. *Mol. Biol. Evol*. **37**, 291–294. (doi:10.1093/molbev/msz189)

64. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A,
Lanfear R. 2020 IQ-TREE 2: New models and efficient methods for phylogenetic inference
in the genomic era. *Mol. Biol. Evol.* **37**, 1530–1534. (doi:10.1093/molbev/msaa015)

65. Minh BQ, Hahn MW, Lanfear R. 2020 New methods to calculate concordance factors for
phylogenomic datasets. *Mol. Biol. Evol.* **37**, 2727–2733. (doi:10.1093/molbev/msaa106)

66. Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat. Methods* **14**,
587–589. (doi:10.1038/nmeth.4285)

67. Chifman J, Kubatko L. 2014 Quartet inference from SNP data under the coalescent model.
Bioinformatics **30**, 3317–3324. (doi:10.1093/bioinformatics/btu530)

68. Jombart T, Devillard S, Balloux F. 2010 Discriminant analysis of principal components: a
new method for the analysis of genetically structured populations. *BMC Genet.* **11**, 94.
(doi:10.1186/1471-2156-11-94)

69. Jombart T. 2008 adegenet: a R package for the multivariate analysis of genetic markers.
Bioinformatics **24**, 1403–1405. (doi:10.1093/bioinformatics/btn129)

70. Terraneo TI, Berumen ML, Arrigoni R, Waheed Z, Bouwmeester J, Caragnano A, Stefani
F, Benzoni F. 2014 *Pachyseris inattesa* sp. n. (Cnidaria, Anthozoa, Scleractinia): a new
reef coral species from the Red Sea and its phylogenetic relationships. *ZK.* **433**, 1–30.
(doi:10.3897/zookeys.433.8036)

71. Medina M, Collins AG, Takaoka TL, Kuehl JV, Boore JL. 2006 Naked corals: Skeleton
loss in Scleractinia. *Proc. Natl. Acad. Sci. U.S.A.* **103**, 9096–9100.
(doi:10.1073/pnas.0602444103)

72. Tamir R, Eyal G, Kramer N, Laverick JH, Loya Y. 2019 Light environment drives the
shallow-to-mesophotic coral community transition. *Ecosphere* **10**.
(doi:10.1002/ecs2.2839)

73. Prada C, Hellberg ME. 2013 Long prereproductive selection and divergence by depth in a
Caribbean candelabrum coral. *Proc Natl Acad Sci. U.S.A.* **110**, 3961–3966.
(doi:10.1073/pnas.1208931110)

74. Rippe JP, Dixon G, Fuller ZL, Liao Y, Matz M. 2021 Environmental specialization and
cryptic genetic divergence in two massive coral species from the Florida Keys Reef Tract.
Mol. Ecol. **30**, 3468–3484. (doi:10.1111/mec.15931)

75. Hoeksema BW, Cairns S. 2022. *World List of Scleractinia. Leptoseris mycetoseroides*
Wells, 1954. See: https://www.marinespecies.org/aphia.php?p=taxdetails&id=207283

76. Snell TL, Foltz DW, Sammarco PW. 1998 Variation in morphology vs conservation of a
mitochondrial gene in *Montastraea cavernosa* (Cnidaria, Scleractinia). *goms*. **16**.
(doi:10.18785/goms.1602.08)

77. Fukami H *et al*. 2008 Mitochondrial and nuclear genes suggest that stony corals are
monophyletic but most families of stony corals are not (Order Scleractinia, Class
Anthozoa, Phylum Cnidaria). *PLoS ONE* **3**, e3222. (doi:10.1371/journal.pone.0003222)

78. Shearer TL, Coffroth MA. 2008 DNA Barcoding: Barcoding corals: limited by
interspecific divergence, not intraspecific variation. *Mol. Ecol. Resour.* **8**, 247–255.
(doi:10.1111/j.1471-8286.2007.01996.x)

79. Stemann TA. 1991 Evolution of the reef-coral family Agariciidae (Anthozoa: Scleractinia)
in the Neogene through recent of the Caribbean. Doctoral dissertation, University of Iowa,
Iowa City, IA.

80. Oken L. 1815-1816 Lehrbuch der Naturgeschichte. Dritter Theil: Zoologie. Erste Abtheilung: Fleischlose Thiere. **842**

81. Knowlton N, Weigt LA, Solórzano LA, Mills DK, Bermingham E. 1993 Divergence in proteins, mitochondrial DNA, and reproductive compatibility across the Isthmus of Panama. *Science* **260**, 1629–1632. (doi:10.1126/science.8503007)

82. Sánchez JA, González-Zapata FL, Prada C, Dueñas LF. 2021 Mesophotic gorgonian corals evolved multiple times and faster than deep and shallow lineages. *Diversity* **13**, 650. (doi:10.3390/d13120650)

83. Knowlton N, Weil E, Weigt LA, Guzmán HM. 1992 Sibling species in *Montastraea annularis*, coral bleaching, and the coral climate record. *Science* **255**, 330–333. (doi:10.1126/science.255.5042.330)

84. Carlon DB, Budd AF. 2002 Incipient speciation across a depth gradient in a scleractinian coral? *Evolution* **56**, 2227–2242. (doi:10.1111/j.0014-3820.2002.tb00147.x)

85. Frade PR, Reyes-Nivia MC, Faria J, Kaandorp JA, Luttikhuizen PC, Bak RPM. 2010 Semi-permeable species boundaries in the coral genus *Madracis*: Introgression in a brooding coral system. *Mol. Phylogenet. Evol*. **57**, 1072–1090. (doi:10.1016/j.ympev.2010.09.010)

86. Bongaerts P, Riginos C, Hay KB, van Oppen MJ, Hoegh-Guldberg O, Dove S. 2011 Adaptive divergence in a scleractinian coral: Physiological adaptation of *Seriatopora hystrix* to shallow and deep reef habitats. *BMC Evol. Biol.* **11**, 303. (doi:10.1186/1471-2148-11-303)

87. Costantini F, Rossi S, Pintus E, Cerrano C, Gili J-M, Abbiati M. 2011 Low connectivity and declining genetic variability along a depth gradient in *Corallium rubrum* populations. *Coral Reefs* **30**, 991–1003. (doi:10.1007/s00338-011-0771-1)

88. Bongaerts P, Dubé CE, Prata KE, Gijsbers JC, Achlatis M, Hernandez-Agreda A. 2021
Reefscape genomics: Leveraging advances in 3D imaging to assess fine-scale patterns of
genomic variation on coral reefs. *Front. Mar. Sci.* **8**. (doi:10.3389/fmars.2021.638979)

89. Veron, JEN. 2000 *Corals of the world*. Australian Institute of Marine Science and CRR
Qld. Pty. Ltd. Australia. 3, 490 (ISBN: 0642322384)

90. Spalding MD *et al.* 2007 Marine ecoregions of the world: a bioregionalization of coastal
and shelf areas. *BioScience* **57**, 573–583. (doi:10.1641/b570707)

91. Mihaljević M. 2019 Oligocene-Miocene scleractinians from the central Indo-Pacific:
Malaysian Borneo and the Philippines. *Palaeontol Electron*. (doi:10.26879/978)

92. Budd AF, Stemann TA, Johnson KG. 1994 Stratigraphic distributions of genera and species
of Neogene to recent Caribbean reef corals. *J. Paleontol*. **68**, 951–977.
(doi:10.1017/s0022336000026585)

93. Veron JEN. 1995 Corals in space and time: The biogeography and evolution of the
Scleractinia. Cornell University Press, New York, NY.

94. Lanier HC, Huang H, Knowles LL. 2014 How low can you go? The effects of mutation
rate on the accuracy of species-tree estimation. *Mol. Phylogenet. and Evol*. **70**, 112–119.
(doi:10.1016/j.ympev.2013.09.006)

95. Liu L, Wu S, Yu L. 2015 Coalescent methods for estimating species trees from
phylogenomic data. *J. Syt. Evol*. **53**, 380–390. (doi:10.1111/jse.12160)

96. Manthey JD, Campillo LC, Burns KJ, Moyle RG. 2016 Comparison of target-capture and
restriction-site associated DNA sequencing for phylogenomics: A test in Cardinalid
Tanagers (Aves, Genus:*Piranga*). *Syst. Biol*. **65**, 640–650. (doi:10.1093/sysbio/syw005)

97. Hühn P, Dillenberger MS, Gerschwitz-Eidt M, Hörandl E, Los JA, Messerschmid TFE,
Paetzold C, Rieger B, Kadereit G. 2022 How challenging RADseq data turned out to favor
coalescent-based species tree inference. A case study in Aichryson (Crassulaceae). *Mol.*
Phylogenet. Evol. **167**, 107342. (doi:10.1016/j.ympev.2021.107342)

98. Harvey MG, Smith BT, Glenn TC, Faircloth BC, Brumfield RT. 2016 Sequence capture
versus restriction site associated DNA sequencing for shallow systematics. *Syst. Biol.* **65**,
910–924. (doi:10.1093/sysbio/syw036)

99. Quattrini AM *et al.* 2018 Universal target-enrichment baits for anthozoan (Cnidaria)
phylogenomics: New approaches to long-standing problems. *Mol. Ecol. Resour.* **18**, 281–
295. (doi:10.1111/1755-0998.12736)

100.   Hundsdoerfer AK, Lee KM, Kitching IJ, Mutanen M. 2019 Genome-wide SNP data
reveal an overestimation of species diversity in a group of Hawkmoths. *Genome Biol. and*
Evol. **11**, 2136–2150. (doi:10.1093/gbe/evz113)

Acknowledgements

We thank for logistical support: CARMABI Research Station, Substation Curaçao, CIEE Bonaire
Research Station, STINAPA Bonaire, CORALINA, PNN "Corales de Profundidad", Hawaii
Undersea Research Laboratory (HURL), Reef Connections, Mike Ball Dive Expeditions, SY
Ethereal, and Waitt Foundation.

Funding

This work was funded by the XL Catlin Seaview Survey (funded by the XL Catlin Group in
partnership with Underwater Earth and The University of Queensland), "The Explorers Club -
Eddie Bauer Grant for Expeditions", an Australian Research Council Discovery Early Career
Researcher Award (DE160101433), and the Hope for Reefs Initiative at the California Academy
of Sciences.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.