Metadata preservation and stewardship for genomic data is possible, but must happen now

Eric D. Crandall[1], Rachel H. Toczydlowski[2,3], Libby Liggins[4], Ann E. Holmes[5], Maryam Ghoojaei[6], Michelle R. Gaither[6], Briana E. Wham[7], Andrea L. Pritt[8], Cory Noble[4], Tanner J. Anderson[9], Randi L. Barton[10,11], Justin T. Berg[12,13], Sofia G. Beskid[14], Alonso Delgado[15], Emily Farrell[6], Nan Himmelsbach[16], Samantha R. Queeno[9], Thienthanh Trinh[6], Courtney Weyand[17], Andrew Bentley[18], John Deck[19], Cynthia Riginos[20], Gideon S. Bradburd[2,20], Robert J. Toonen[22]

1. Department of Biology, Pennsylvania State University, University Park, Pennsylvania, PA 16801
2. Department of Integrative Biology - Ecology, Evolution, and Behavior Program, Michigan State University, East Lansing, MI 48824
3. Current Affiliation: Institute for Applied Ecosystem Studies, Northern Research Station, United States Forest Service, Rhinelander, WI, 54501, U.S.
4. School of Natural Sciences, Massey University, Auckland, Aotearoa New Zealand
5. Department of Animal Science, University of California-Davis, Davis, CA 95616
6. Department of Biology, University of Central Florida, Orlando, FL 32816
7. Department of Research Informatics and Publishing, The Pennsylvania State University Libraries, University Park, Pennsylvania, PA 16801
8. Madlyn L. Hanes Library, The Pennsylvania State University Libraries, Pennsylvania State University, Harrisburg Campus, 351 Olmsted Drive, Middletown, PA 17057
9. Department of Anthropology, University of Oregon, Eugene, OR 97403
10. California State University Monterey Bay, 100 Campus Circle, Seaside, CA, 93955
11. Moss Landing Marine Laboratories, Moss Landing, CA 95039
12. UOG Marine Laboratory, University of Guam, Mangilao, Guam 96910
13. Current Affiliation: Department of Oceanography, University of Hawai'i at Mānoa, Kane'ohe, HI 96744
14. Department of Integrative Biology, University of Texas at Austin, Austin, TX 78712
15. Department of Evolution, Ecology, and Organismal Biology, The Ohio State University; Columbus, OH 43210
16. Department of Natural Science, Hawai'i Pacific University, Honolulu, HI 96813
17. Department of Biological Sciences, Auburn University, Auburn, AL 36849
18. Biodiversity Institute, University of Kansas, Lawrence, KS, 66045
19. Berkeley Natural History Museums, University of California, Berkeley, California

20. School of Biological Sciences, The University of Queensland, Brisbane, Australia
21. Current affiliation: Department of Ecology and Evolutionary Biology, University of Michigan 1105 N. University Ave, Ann Arbor, MI, 48109
22. Hawai'i Institute of Marine Biology, University of Hawai'i at Mānoa, Kane'ohe, HI 96744

Abstract

Genetic diversity within species represents a fundamental yet underappreciated level of biodiversity. Because genetic diversity can indicate species and population resilience to changing climate, its measurement is relevant to many national and global conservation policy targets. Many studies of evolutionary biology, molecular ecology and conservation genetics produce large amounts of genome-scale genetic diversity data for wild populations. While open data policies have ensured an abundance of freely available genomic data stored in the databases of the International Nucleotide Sequence Database Collaboration (INSDC), only about 13% of current accessions have the associated spatial and temporal metadata in INSDC necessary to be reused in monitoring programs, macrogenetic studies, or for acknowledging the sovereignty of nations or Indigenous Peoples. We undertook a "distributed datathon" to quantify the availability of these missing metadata in sources external to the INSDC and to test the hypothesis that these metadata decay with time. We also worked to remediate these missing metadata by extracting them, when present, from associated published papers, online repositories, and/or from direct communication with authors. Starting with 848 programmatically identified candidate datasets (INSDC BioProjects), we manually determined that 561 contained samples from wild populations. We successfully restored spatiotemporal metadata (locality name and/or geospatial coordinates and collection year) for 78% of these 561 datasets (N = 440 BioProjects comprising 45,105 individuals or BioSamples from 762 species in 17 phyla). We also quantified the availability of 33 additional categories of metadata in sources external to the INSDC. Information about associated publications and the type of habitat from which the samples were taken was the most easily found; information about sampling permits was the most challenging to locate. Looking at papers and online repositories was much more fruitful than contacting authors, who only replied to our email requests 45% of the time. Overall, 23% of our email queries to authors discovered useful metadata. Importantly, we found that the probability of retrieving spatiotemporal metadata declines significantly with the age of the dataset, with a 13.5% yearly decrease for metadata located in published papers or online repositories and up to a 22% yearly decrease for metadata that were only available from authors. This observable metadata decay, mirrored in studies of other types of biological data, should motivate swift updates to data sharing policies and researcher practices to ensure that the valuable context provided by metadata is not lost forever.

Introduction

Genetic diversity is the foundational layer of biodiversity. Just as ecosystem health and resilience depends on the diversity of its component species, the health and resilience of each species depends on its genomic diversity (Clark, 2010; Reusch, Ehlers, Hämmerli, & Worm, 2005). Without genetic diversity in the form of standing allelic variation, populations and species cannot adapt to a rapidly changing climate and other anthropogenically-induced stresses (Blanchet et al., 2020; Raffard, Santoul, Cucherousset, & Blanchet, 2019). Local or global extinctions of species in turn threaten the ecosystems upon which the quality of human lives depend (Brauman et al., 2020; Des Roches, Pendleton, Shapiro, & Palkovacs, 2021). Concerningly, genetic diversity, like all levels of biodiversity, is declining rapidly during the Anthropocene across the tree of life (Exposito-Alonso et al., 2022; Leigh, Hendry, Vázquez-Domínguez, & Friesen, 2019; Miraldo et al., 2016; Pinsky & Palumbi, 2014).

Recognizing the vital importance of biodiversity to human well-being and the future of our planet, several international agreements strongly encourage the monitoring and conservation of genetic diversity in both wild and domesticated species. Foremost among these are the United Nations Sustainable Development Goal 2.5 and the international Convention on Biological Diversity (CBD) treaty, which explicitly acknowledge the importance of monitoring and conserving any component of biological diversity (including genetic diversity) that may have "actual or potential use or value for humanity." Moreover, the CBD's article 15 and attendant Nagoya Protocol codify procedures to ensure the sharing of benefits arising from genetic resources (such as digital sequence information; DSI) discovered or accessed within a nation's sovereign borders. The subsequent Strategic Plan for Biodiversity 2011-2020 laid out the 20 Aichi Biodiversity Targets, including target 13, which aims to maintain the "genetic diversity of cultivated plants and farmed and domesticated animals and of wild relatives, including other socio-economically as well as culturally valuable species." Now, even as we are facing shortfalls on all 20 of the Aichi Biodiversity Targets (CBD, 2020; Hoban et al., 2021), there have been calls to broaden genetic diversity targets to include *all* extant species in the New Post-2020 Global Biodiversity Framework, in what has been described as a "moonshot for biology" (Hoban et al., 2020; Laikre et al., 2020; Lewin et al., 2018).

Over the last decade, advances in DNA sequencing technology have enabled the generation of genome-scale datasets of ever larger numbers of individuals, drawn from a growing variety of species (Allendorf, 2017; Hendricks et al., 2018). Researchers are now able to genotype thousands of genomic loci or sequence whole genomes from non-model species for which they have no prior genetic resources (Lou, Jacobs, Wilder, & Therkildsen, 2021; Willette et al., 2014). The shift from genetic to genomic-scale datasets is catalyzing novel conservation insights including: the detection of inbreeding depression (e.g. Kardos, Taylor, Ellegren, Luikart, & Allendorf, 2016), the discovery of subtle, previously undetectable population structure (e.g. Cheng, Gold, Rodriguez, & Barber, 2021; Gaither et al., 2018), reconstruction of demographic histories (Prada et al., 2016), the precise identification of distant pedigree relationships (e.g. Baetscher et al., 2019), uncovering cryptic species (e.g. Quattrini et al., 2019), clues about the genomic basis of local adaptation (e.g. Wilder, Palumbi, Conover, & Therkildsen, 2020) and important traits such as nutritional components (e.g. Kumar et al., 2021). Accordingly, the DSI

82 derived in these studies is highly valued as a resource equivalent to biobanks, providing essential
83 information for conservation (Hoban et al., 2022) as well as ensuring future food security
84 (Castañeda-Álvarez et al., 2016; Halewood et al., 2018).
85
Genomic datasets record the genetic diversity of a species at a particular time and
location, providing a benchmark for how populations are responding to human drivers of
changing environmental conditions, cultivation, and land and sea use, as well as measuring
indicators of progress toward conservation targets and goals (Hoban et al., 2022, 2020) and the
genetic resources available for future cultivation or domestication (Halewood et al., 2018).
However, genomic datasets can only be useful for monitoring global genetic biodiversity and the
sustainable human use of genetic diversity (including benefit-sharing, Cowell et al., 2022) when
archived publicly with accompanying metadata about the spatiotemporal, environmental and
methodological context of the sequenced sample (Riginos et al., 2020; Scholz et al., 2022;
Schriml et al., 2020).
96
The genetics community has long championed open data publication with the
foundational databases of the International Nucleotide Sequence Database Collaboration
(INSDC; Cochrane, Karsch-Mizrachi, Takagi, & INSDC, 2016) forming in the early 1980's. In
2009, the INSDC launched the Sequence Read Archive as a repository dedicated to second-
generation sequence data. It has since grown exponentially to include over 600 terabytes of
freely-available DNA sequence data from over 16,700 wild and domesticated eukaryotic species
as of 2021 (Toczydlowski et al., 2021). Around the same time, the MIxS metadata standards
(Field et al., 2008; Yilmaz et al., 2011) were defined to inform the minimum information about
what (detailed taxonomy), *where* (GPS coordinates and habitat), *when* (collection date), *how*
(sampling and sequencing protocols) and *by whom* a genetic sample was collected. Enabled by
the INSDC infrastructure and encouraged by the Joint Data Archiving Policy (JDAP;
http://datadryad.org/pages/jdap) implemented by top journals in 2011, the proportion of papers
providing open access to their genetic data increased dramatically (Pope, Liggins, Keyse,
Carvalho, & Riginos, 2015). However, the inclusion of accompanying metadata crucial for the
reuse of these data for genetic diversity monitoring, macrogenetic studies, or identifying their
provenance within national boundaries or the lands and waters of Indigenous Peoples, has lagged
behind (Pope et al., 2015; Toczydlowski et al., 2021). As of 2021, out of over 300,000 SRA
BioSamples that are potentially relevant to global genetic biodiversity, only ~13% had metadata
indicating both the time and precise location from which they were sampled (Toczydlowski et
al., 2021).
117
In a timely and welcome update to their policy, INSDC now intends to extend their minimum
metadata requirements to include collection date and country of origin
(https://www.insdc.org/spatio-temporal-annotation-policy-18-11-2021). Although 'country' is
legislatively aligned with the Nagoya Protocol, it is not spatially aligned with the lands and
waters of Indigenous Peoples (e.g. https://native-land.ca/) and does not provide adequate spatial
resolution for conservation monitoring. Moreover, this policy and infrastructure change will take
time to implement (anticipated to be end of 2022), meaning that much of the genomic data
collated over the last ~12 years for past and present populations, of immeasurable value to
understanding and monitoring the biodiversity crisis, are not Findable, Accessible, Interoperable
or Reusable (FAIR; Wilkinson et al., 2016). This absence of appropriate spatio-temporal

metadata represents the effective loss of tens to hundreds of millions of dollars of research effort
for most future purposes (Schriml et al., 2020; Toczydlowski et al., 2021), rendering associated
genetic data invisible to government ministries and non-governmental organizations tasked with
protecting the world's natural environment (Laikre, 2010; Laikre et al., 2020). Moreover,
without spatiotemporal provenance of genomic data enabling connection to the lands and waters
of Indigenous Peoples, these peoples will potentially lose out on benefits (e.g. capacity
development, food security, biomedical advances) arising from genomic information originating
within their territories (Liggins, Hudson, & Anderson, 2021; Marden et al., 2021; McCartney et
al., 2022; Scholz et al., 2022). There is urgency in addressing this metadata gap: previous studies
of morphological (Vines et al., 2014) and genetic (Pope et al., 2015) data suggest that the
probability of existing metadata ever being linked to the genomic data significantly decreases
over time.
140
In the summer of 2020, we convened a distributed remote datathon to (1) assess the
availability of metadata outside of the INSDC, (2) recover and curate metadata missing in
INSDC from external sources (i.e. published research papers, other online repositories, or the
authors themselves), and (3) extend our initial report on the metadata gap (Toczydlowski et al.,
2021) to investigate how the recoverability of these metadata is affected by dataset age and to
document shortfalls and costs of our remedial efforts. In our datathon, 14 graduate students from
the USA and 12 professional academics (authors on this paper) across 3 countries worked
together via Zoom, Slack and Google Sheets as "metadata curators" to establish and execute
curation protocols and infill missing metadata. Collectively, we searched for metadata external to
the INSDC (e.g. associated scientific publications, Dryad, museum collections) for 848 genomic
datasets (INSDC BioProjects) representing 94,416 individual samples (BioSamples). The
BioSamples and associated genetic sequence data in these projects were selected as they were
missing at least latitude and longitude metadata in the INSDC. Our findings underscore the
importance of appropriate and immediate metadata archival going forward. We provide guidance
based on our collective experience gained over the datathon on practices to retain crucial
metadata.
157
158 **Methods**
159
160 *Identifying BioProjects for metadata curation*
161
Our datathon followed a workflow illustrated in Figure 1. We first searched the INSDC to
identify datasets that were potentially relevant to monitoring genetic diversity of wild
populations (including wild relatives of domesticated species) but were lacking critical metadata
about the latitude and longitude of the sampling location (as described in Toczydlowski et al.,
2021). On November 7, 2019, we searched the INSDC BioProject database using the *rentrez* R
package (Winter, 2017) and custom R scripts to locate datasets that contained genomic-level
DNA sequence data from eukaryotic species, excluding human and metagenomic datasets. We
then downloaded all metadata associated with these BioProjects from the INSDC PubMed,
Taxonomy, BioSample and SRA databases. We further filtered the BioProjects to remove
BioSamples (sequenced individuals) from species whose population dynamics and evolution are
largely governed by humans: pathogens and their vectors, model organisms, and domesticated
species. We used custom lists for each category of non-wild organisms (Supplementary

Materials S1). These filters helped to target our efforts at recovering metadata for the datasets
most likely to be of relevance to genetic diversity monitoring of wild populations. Additional
details about the search and filtering steps can be found in Supplemental Materials of
Toczydlowski et al. 2021. These filtering steps yielded 848 BioProjects that entered our datathon
curation pipeline. These BioProjects contained at least 5 individuals that lacked geospatial
coordinates in the INSDC and were potentially sampled from wild populations.
180
We built a blank template to enter metadata into that we located external to the INSDC using
the Genomic Observatories Metadatabase (GEOME; Deck et al., 2017; Riginos et al., 2020)
This template included terms that described collection date and location, habitat etc. Table 1
gives definitions of 16 required and recommended metadata terms, while Supplemental Materials
S2 provides an example template with all 36 metadata terms and their definitions and controlled
vocabularies. We then programmatically filled this template for each of the 848 BioProjects in
our pipeline with the metadata already present in the INSDC using custom R scripts.
188
189   *Recovering and curating metadata external to the INSDC*
190
Metadata curators were each randomly assigned a set of BioProjects. Curators followed a
standard protocol (Figure 1; Supplemental Materials S3) to locate and enter associated metadata
for samples in each BioProject that were missing in the INSDC but reported in external sources
(e.g. associated published scientific papers). Briefly, we first searched for associated publications
by googling the BioProject PRJ accession number and/or key words associated with the
BioProject (e.g. author, affiliation, funding information, species, and locality names). While the
INSDC includes terms to capture associated publications, these were only populated for 7% of
the 848 BioProjects that we examined. We then skimmed each paper to determine if the
BioProject contained sequence data from wild individuals. BioProjects comprising over 50%
BioSamples from non-wild individuals (brood stock, laboratory stock, domesticated species, and
some seed banks) were marked as "not relevant" and removed from further curation. Non-wild
BioSamples that were a minority in their BioProject were designated as such under the
establishmentMeans metadata term, and further metadata recovery was not prioritized for these
samples. BioProjects that pooled their genetic samples without barcodes (precluding estimates of
genetic diversity at the level of individual) were also marked as "not relevant" and removed from
the curation pipeline. Once relevance to our efforts was determined, curators looked in figures,
tables, and supplemental information of associated publications and/or linked online repositories
(e.g. Data Dryad, Github, Zenodo, and databases of biocollections such as museums and
herbaria) for sample-level metadata from 36 metadata terms (most defined as Darwin Core
terms; Wieczorek et al., 2012). When necessary, curators converted metadata to standard formats
(e.g. degree minutes seconds data were converted to decimal degrees), and then added metadata
into the pre-generated GEOME template for that BioProject. After performing quality control,
these metadata could then be easily uploaded to GEOME and potentially cross-walked into the
appropriate INSDC databases.
215
After adding all metadata that could be gleaned from the associated paper(s) into the
GEOME templates, curators made a structured comment on a master spreadsheet (Supplemental
Materials S4) indicating whether metadata for each of the required and recommended terms were
absent for all BioSamples ("none"), present for less than 50% of BioSamples ("some"), present

for greater than 50% of BioSamples ("most"), or iv) present for all BioSamples ("all"). If the
paper was missing information from one of six or seven "required" terms (georeference-able
locality OR [*decimalLatitude* AND *decimalLongitude*], *coordinateUncertaintyInMeters*,
georeferenceProtocol, *habitat*, *environmentalMedium*, *yearCollected*), the curator flagged the
BioProject to initiate author contact. An additional nine metadata terms were considered
"recommended": missing metadata in these fields alone did not trigger an author contact but

Figure 1. Datathon workflow. The number of BioProjects and BioSamples remaining after each step
are given below the step.

226   curators and authors were asked to populate these fields as completely as possible. These
227   recommended terms included *country, establishmentMeans, permitInformation,*
228   *associatedReferences, preservative*, and 4 termss that tracked genetic data derived from the raw
229   reads such as SNP genotypes or sequence alignments. Progress and notes at each curation step
230   were tracked as meta-metadata on the master spreadsheet.
231
After a quality-control step to ensure that author names and email addresses found in papers
were input correctly, corresponding authors of the paper were contacted by email (see
Supplemental Material S5 for email text) using the Yet Another Mail Merge add-on for Google
Sheets (yamm.com). Using YAMM allowed us to track email receipts and responses. If an email
was undeliverable, we used our best efforts to locate an alternate email. We were able to
successfully deliver email queries for all 351 of 492 relevant BioProjects that met the criteria for
author contact. About two weeks after sending the initial email, curators sent reminder emails to
unresponsive authors at least once and at most twice. This process emulated the efforts of a
reasonably persistent researcher to obtain metadata important to their research.
241
Once curators had gathered and entered all available metadata into the GEOME template for
a BioProject (from online sources external to the INSDC and/or directly from authors), curators
validated the metadata at the GEOME website (geome-db.org). This programmatic GEOME
validation ensured that metadata in each field were correctly formatted and within controlled
vocabularies, e.g. for *country* and *habitat* (see lists tab of Supplemental Materials S3 for all
controlled vocabularies). Following validation, MRG, BEW, AP and EDC performed a final
quality-control check (QC; steps described in Supplemental Material S1). Filled and QCed
GEOME templates for each BioProject will be uploaded to the GEOME database.
250
251   *Investigating metadata decay*
252
We investigated the effect of BioProject age on the probability that we were able to recover
metadata information for 11 metadata categories. Previous investigations of metadata have
indicated rapid decay when data are not publicly archived (Roche et al., 2014; Vines et al., 2014,
2013). We used Bayesian logistic regression to fit four distinct models to investigate the
relationships between BioProject age (number of days between publication in the INSDC and
November 7, 2019) and: A) the probability that metadata could be retrieved from INSDC,
associated published papers, and/or repositories, B) the probability that we received an author
response for the 351 BioProjects that triggered an author contact via email, C) the probability
that authors provided any metadata, given that they responded and D) the probability that authors
provided metadata for a majority of samples, given that they responded.
263
Information about the collection date and location of a sample are the most critical pieces of
metadata required to identify the Indigenous provenance of genomic data and make genomic
sequence data repurposable for monitoring and fundamental data synthesis research, so we
focused our investigations on these two categories; we refer to the aggregate as spatiotemporal
metadata. We defined a BioProject as having spatiotemporal data if collection dates, and
latitudes and longitudes and/or locality were present for at least 50% of the BioSamples that it
contained. In model C, we counted a gain in collection year, or place name, or latitude/longitude
for any number of BioSamples as recovery of metadata. In model D, we only counted increases

in metadata where BioProjects had incomplete spatiotemporal metadata for > 50% of its
BioSamples and then had spatiotemporal metadata present for > 50% of BioSamples after
contacting authors. That is, model C assessed the probability of recovering any metadata external
to the INSDC, and model D assessed the probability of recovering metadata for the majority of
samples. In a supplemental analysis, we investigated how metadata within individual
spatiotemporal terms and other important metadata terms (i.e. required and recommended terms,
Table 1) decayed, including: *decimalLongitude and decimalLatitude, collectionYear, country,*
locality, habitat, environmentalMedium, materialSampleID, permitInformation,
associatedReferences (publication DOI)*, preservative*, and derived genetic data terms
(Supplemental Materials S6).
282
We conducted all statistical analyses at the level of BioProject (as opposed to BioSamples or
genomic sequences), because presence/absence of metadata for BioSamples within a given
BioProject was highly correlated (Toczydlowski et al., 2021). Curators also tracked metadata
recovery efforts at the level of BioProject for convenience, and we contacted authors about entire
BioProjects rather than individual BioSamples. In each set of models, we removed BioProjects
that already had complete metadata in the category of interest, and therefore could not gain any
more.
290
We analyzed the effect of BioProject age on our response variables (model A: successful
metadata retrieval; model B: author response; model C: author provided any metadata
conditional on a response to our query; model D: author provided metadata for the majority of
samples conditional on a response to our query) using generalized linear models. In each
analysis, we modeled our response variable as a Bernoulli-distributed variable with a probability
of success that was a linear function of our predictor variable: BioProject age. In each analysis,
the parameters of our model were a global mean probability of success and an effect size of
BioProject age on probability of success for that response variable. Analyses used the canonical
inverse-logit inverse link function. In mathematical notation, our model was:
300

$$Y_i \sim \text{Bernoulli}(p_i)$$
$$p_i = \frac{1}{1 + e^{-\theta_i}}$$
$$\theta_i = \mu + \beta \times X_i$$

301
Where Y_i is the ith outcome (response variable), p_i is the probability of successfully observing that
outcome, μ is the global mean probability of success, and β is the effect of BioProject age on the
transformed probability of success for that outcome (θ_i). We had no strong prior beliefs about the
effect of BioProject age on success in each of the four analyses we ran; to reflect these beliefs,
the priors we placed on our parameters were: $\beta \sim N(0,10)$; $\mu \sim N(0,10)$. All statistical analyses
were performed using Rstan version 2.21.2 [50] running 4 independent chains for 2,000
iterations, thinning to sample only every 4th iteration to reduce autocorrelation, and discarding
the first 1,000 iterations as burn-in. To assess the significance of the effect of BioProject age on

Figure 2. The taxonomic and geographic scope of the datathon. A) A cladogram of 719 of the 762 species from BioProjects that passed through the final quality control step. This is a subtree of the Open Tree of Life (Hinchcliff et al. 2015) generated with the rotl package for R (Michonneau et al. 2016) and visualized with iTOL software (itol.embl.de; Letunic and Bork 2021). B) Heatmap of species for BioSamples for which spatial coordinates were recovered by the datathon. C) Map showing the geographic distribution of broad taxonomic categories of these BioSamples. D) Heatmap of these BioSamples.

success of each outcome, we determined whether the 95% equal-tailed credible interval of the marginal distribution on β contained 0; if it did, the effect of BioProject age was deemed not significant.

Results

We identified 848 INSDC BioProjects, representing 94,416 BioSamples from individual eukaryotic organisms that lacked geospatial coordinates and had at least five putatively wild individuals as determined by our filters. Curators were able to locate associated published scientific papers for 741 of these 848 BioProjects. Reading these papers revealed 561 BioProjects with a majority of relevant, truly wild individuals, comprising 63,684 individuals from 873 species. After scouring associated published papers for metadata and contacting authors, a total of 440 BioProjects with 45,105 BioSamples from 762 species in 17 eukaryotic phyla (Figure 2A) had geospatial data (either coordinates or a locality name) and were passed through quality control for eventual upload to GEOME. BioSamples that passed through the datathon came from all continents and all major oceans (Figure 2B-D).

For the subset of BioProjects that we focused on (those that were missing latitude and longitude), datathon curators were able to recover metadata for a majority of BioSamples in a BioProject as follows (depicted in Figure 3). For geospatial coordinates, nearly 60% could be

found in an associated publication or online repository. While nearly 30% of these BioProjects
did already contain information about collection year in the INSDC, curators were only able to
recover an additional 21% from papers or online repositories. Datathon curators recovered
metadata regarding habitat, environmental medium (the media displaced by the sampled
organism) and publication DOI for over 80% of BioProjects from published papers and their
supplemental information. Additional large gains in BioProjects were made from online sources
external to the INSDC for locality (48.8%), and country name (39.8%). Notably, permit
information was the least available of any of the metadata categories that we explored. There is
no permit metadata term in the INSDC and curators found permit information in papers for only
21% of BioProjects.
340
Contacting authors yielded comparatively less metadata than our search of papers and
supplemental information, although it should be noted that this step was secondary to looking in
papers and online. Out of 351 author contact attempts, we received 158 responses (45% response
rate). Of the 158 responses, 80 (51%) provided at least some missing metadata, yielding an
overall "useful author response rate" of 23%. Through contacting authors, we were able to
recover collection year metadata for an additional 9% of BioProjects, and geospatial coordinates
for an additional 8.5% percent of BioProjects. Gains in other metadata categories were all less
than 5%, with permit information showing only a 1.2% increase from authors.
349
350

Figure 3. Stacked bars showing the percent of BioProjects for which metadata were
found from each of three sources, across 10 priority metadata categories.

The age (time since deposition into the INSDC) of the BioProject had a strong effect on
whether metadata could be recovered. After searching for metadata within the INSDC and within
published papers, we found that spatiotemporal metadata (defined as year AND geospatial
coordinates OR locality), had a mean odds ratio of 0.865 (95% highest posterior density credible
interval [HPD CI]: 0.775 - 0.964; Figure 4A). This indicates that for every year after a
BioProject is published to the SRA, there is about a 13.5% decrease (HPD CI: 3.6% - 22.5%) in
the probability that its metadata can be found in the SRA, in papers or elsewhere online. On the
other hand, there was a strong positive effect of BioProject age on whether an attempt to contact
the authors was answered, with a 25.5% increase in the probability of a reply of any kind for
every year after SRA publication (mean odds ratio of 1.255; 95% HPD CI: 1.120 - 1.412; Figure
4B). In other words, we were more likely to get an email response for older datasets. However,
given a response, the probability that authors furnished any amount of metadata for year OR
coordinates OR locality decreased with BioProject age by 21% per year (odds ratio 0.810; 95%
HPD CI: 0.680 - 0.949; Figure 4C). Similarly, the probability that the authors provided metadata
for year AND coordinates OR locality for a majority of BioSamples decreased by 22% per year
(odds ratio: 0.819; 95% HPD CI: 0.671 - 0.994; Figure 4D).
368
Figures for Bayesian logistic regressions of BioProject age on other metadata categories can
be found in Supplemental Materials S6 (figures) and S7 (tables of β values). In accordance with
the results for spatiotemporal metadata, supplementary analyses indicated that metadata for
collection year (posterior mean slope = -0.133, 95% Credible Interval: -0.233 - -0.034; Table S1,
Figure S6-11A) and preservative used (posterior mean slope = -0.111, 95% HPD: -0.218 - -
0.009; Figure S6-9A) were significantly less likely to be recovered from INSDC, publications,
and/or online repositories with increasing age of a BioProject. Furthermore, and as with
spatiotemporal metadata, the probability that responding authors provided additional metadata

Figure 4. The effect of BioProject age on the probability of recovering spatiotemporal metadata.
Density plots depict posterior distribution for log(Odds Ratio), with black lines showing 95% highest
posterior density (HPD). (A) Probability that metadata were found in the INSDC or in the INSDC or
associated papers and repositories. (B) Probability of receiving a reply from BioProject authors to our
contact email. (C) Probability of receiving any amount of additional metadata for year OR coordinates
OR locality. (D) Probability of receiving metadata for year AND coordinates OR locality for a
majority of BioSamples. 95% HPD intervals exclude 0 in all panels.

for georeferences (decimalLatitude and decimalLongitude; posterior mean slope = -0.151, 95% Credible Interval: -0.386 – -0.05; Figure S6-2C), collection year (posterior mean slope = -0.174, 95% Credible Interval: -0.363 – 0.000; Figure S6-11C), and preservative used (posterior mean slope = -0.438, 95% Credible Interval: -0.873 – -0.081; Figure S6-9C) was significantly greater for younger BioProjects. The provisioning of permit information followed this same trend (although marginally insignificant, posterior mean slope = -0.555, 95% Credible Interval: -1.31 - 0.003; Figure S6-5C), suggesting these metadata are relatively available within the personal data management system of authors.

Concerningly and counter to our result for spatiotemporal metadata, supplementary analyses indicated that metadata for habitat (Table S1, posterior mean slope = 0.141, 95% Credible Interval: 0.006 - 0.285; Figure S6-6C) and environmental medium (posterior mean slope = 0.176, 95% Credible Interval: 0.016 - 0.355; Figure S6-5C) were less frequently recovered from INSDC, publications and/or repositories for younger BioProjects. The reasons for these results are unclear, but may indicate a decline in author attention and appreciation of organismal natural history. Retrieval of these metadata through author contact had no relationship with BioProject age.

Discussion

With this distributed datathon, we have demonstrated that crucial metadata can be restored for many genomic investigations of wild organisms. However, our analyses show that metadata are more difficult to recover the longer we wait, and many are locked in non-standard formats. Because the great majority of publicly available genomic datasets lack important metadata (Toczydlowski et al., 2021), they are not findable, accessible, interoperable nor reusable (FAIR; Wilkinson et al., 2016). Only genomic data that are FAIR will allow systematic monitoring of the fundamental layer of biodiversity (Hoban et al., 2021), and enable assertions regarding provenance for informing CBD Nagoya Protocol obligations. Our results illustrate that: (1) metadata availability is dependent on type: location, publication and habitat metadata are much more available or inferable than metadata about permits and preservatives; (2) with considerable time and paid effort, it is possible to recover some of these important metadata from the non-standardized and non-machine-readable formats in which they are currently being stored; and (3) while metadata archival practices may be improving incrementally, genomic metadata are subject to the same decay processes demonstrated for other types of scientific data (Pope et al., 2015; Vines et al., 2014).

There are likely multiple factors underlying the observed metadata decay. First, it is not surprising that older metadata are less likely to have been archived. Metadata archival practices are gradually improving, with more metadata being recorded into the INSDC, research papers, and online repositories such as Data Dryad (Figure 4A). This is consistent with increasing acknowledgement that these metadata are relevant and important to future research. However, the rate of metadata archival is apparently not keeping up with the rapid growth of genomic datasets (see Figure 1 of Toczydlowski et al., 2021) and it is certainly not closing the gap. Second, we found that authors of recent SRA datasets were significantly less likely to reply to our queries than those of older datasets (Figure 4B), although the overall response rate of 45% was comparable to previous studies (Vines et al., 2014, 2013). This result may indicate that

recent SRA depositions are part of ongoing research projects for which authors are unwilling to
share metadata for fear of getting "scooped" by others working on similar research questions. It
is also true that younger authors are more likely to leave science than older authors (Reithmeier
et al., 2019) and thus may no longer support their publications. Similarly, there may be a cohort
effect in which authors of older studies are more established in their careers and have more time,
and/or are more aware of increasing expectations around FAIR data, and thus more willing to
communicate and share. Finally, of the authors that did reply, there was a significant decrease
with the age of the BioProject in whether partial or complete spatiotemporal metadata were
provided (Figure 4C,D), suggesting that if metadata are not properly archived to public
repositories, they are subject to being lost over time, as previously highlighted for morphological
data (Vines et al., 2014).
434

Taken together, our results support assertions that the current research system overly weights
publications and citations, while underweighting scientific openness and transparency (S. W.
Davies et al., 2021; McNutt et al., 2016; Nosek et al., 2015). Changing the system will likely
require a combination of carrots and sticks (Whitlock, 2011). Carrots can take the form of citable
data publications (Dimitrova et al., 2021), recognition of open data practices by hiring,
promotion and tenure committees, or commendations from professional societies or departments
(Roche, Kruuk, Lanfear, & Binning, 2015; Roche et al., 2014). Sticks in the form of open
metadata mandates, must come from journals (Sibbett, Rieseberg, & Narum, 2020, Gareth
Jenkins, pers. comm.), funding agencies, and data repositories, which all have a responsibility to
respond to the needs of the research community (Lin et al., 2020). While we applaud the
INSDC's new spatiotemporal metadata annotation policy requiring country of origin metadata
and their adoption of the MIxS metadata standards, we call for greater mandated spatial
resolution to include at least a locality name or spatial coordinates (Table 1) with appropriate
uncertainty or additional terms (such as Darwin Core's coordinateUncertaintyInMeters,
informationWithheld; Wieczorek et al., 2012) to protect endangered species or sovereignty of
Indigenous Peoples (Hudson et al., 2020; McCartney et al., 2022).
451

Our datathon provided an unparalleled opportunity to train graduate students in the
importance of proper data curation, and to raise awareness that almost every dataset has a
potential for re-use. We suggest that training in data curation and metadata usage should be part
of reproducible research training in every science graduate program, with emphasis on avoiding
some of the metadata practices that hinder metadata recovery described in Box 1. Additionally,
"datathons" such as that undertaken here could help to close the metadata gap in the short term,
as they are very cost effective. If we assume a mean cost of sequencing of USD 50 per
BioSample (and ignore the much higher, additional cost of sample collection and processing),
this datathon rescued over USD 2.1 million worth of genomic sequence data for future research
purposes. Co-authors of this paper spent about 2,300 hours on this metadata retrieval effort,
which, if valued at an average wage of USD 19 per hour, yields a return on investment of nearly
4,700%, with average costs of remediating a BioSample or BioProject at USD 1.05 and USD 110
respectively. But ultimately, datathons are a stopgap solution.
465

Going forward, the entire biodiversity genomics research community should give the same
priority to sharing metadata that they have given to sharing primary data, because it is only the
metadata that make primary data FAIR. From a process standpoint, the collection of metadata

should begin at the time of sampling, with the assignment of a globally unique identifier (GUID)
to the actual material sample. This identifier, which should be assigned as early as possible after
collection, serves as the root to which all subsequent derived products could be linked in an
extended specimen cloud to establish clear provenance and thereby prevent duplication of data or
effort (N. Davies et al., 2021; Lendemer et al., 2020). Through the use of GUIDs, both physical
and digital products of the sample (digital sequence information, but also DNA or RNA
extractions, subsamples, images, video, audio, CT scans, measurements of morphology, traits,
gut contents, parasites, and other related data and associated metadata) will be linked to their
material sample GUID to provide an extensive, holistic metadata cloud that can be used to better
inform current research endeavors as well as create additional data-intensive research pathways.
GEOME (Deck et al., 2017; Riginos et al., 2020) is an example of an easy-to-use "metadata
broker" platform that can provide spreadsheet templates with definitions that can be filled in
offline when the sample is collected. It can then mint a GUID for any sample that is added to it,
and then harvest the INSDC accession numbers for genomic reads that are submitted to the SRA
through GEOME, thereby maintaining permanent links between the sample metadata and
genomic data.
485
The challenge then is to integrate these metadata downstream into databases (such as
INSDC) which describe data derived from the sample. INSDC enables such linkages to other
metadata platforms through the use of both Structured Voucher
(https://www.ncbi.nlm.nih.gov/biocollections/docs/faq/) and Linkout
(https://www.ncbi.nlm.nih.gov/projects/linkout/) facilities for both Nucleotide and SRA (through
their corresponding BioSample record) datasets respectively (e.g.
https://www.ncbi.nlm.nih.gov/nuccore/KC825472). Through these linkages, metadata
corresponding to the original material sample can be tied to the resulting sequence(s) to both
validate the metadata associated with the sequence record as well as provide updated information
should specimens be reidentified or georeferenced after the lodging of the sequence with INSDC.
Using the INSDC as a long-term repository for metadata about the sample may not make sense,
in part because researchers who submit the sequences to INSDC have sole editing rights to the
sequence record and it is currently quite difficult for others (such as the collections who hold the
vouchers) to keep the INSDC metadata up to date or add additional information. Thus, the
integration of these metadata from an upstream source somewhat negates the necessity for this
information to be duplicated by the sequence depositor and ensures that the metadata are
constantly up to date. This not only supports open, reproducible science (Buckner, Sanders,
Faircloth, & Chakrabarty, 2021) but also exemplifies the Findable and Accessible principles of
FAIR data (Wilkinson et al., 2016).
505
What this piecemeal system currently lacks, however, is support for data Interoperability and
Reusability. This is because of the siloed nature of the data and our inability to compile it into a
single resource for machine readability, data manipulation or downstream use. This shortcoming
is being addressed through various initiatives such as the Extended Specimen Network (ESN;
Lendemer et al., 2020; Thiers et al., 2021), the Digital Extended Specimen
(https://dissco.tech/2020/03/31/what-is-a-digital-specimen/), the Distributed System for
Scientific Collections (DiSCCo; https://www.dissco.eu/), iSamples (N. Davies et al., 2021) and
others. Such a system would require all actors in the data landscape (researchers, collections,
data aggregators, publishers, etc.) to utilize and publish resolvable GUIDs on all specimens,

datasets and products of research to make these linkages possible, and thereby create an
extensive online network of knowledge, and increase the potential for scientific research
questions to be answered.
518
We join others in calling for ambitious policy goals that safeguard genetic diversity and
scientific practices to enable this (Des Roches et al., 2021; Díaz et al., 2020; Laikre et al., 2020).
Swift collective action is required to protect all levels of global biodiversity, and the first step
towards protecting the evolutionary health of eukaryotic species worldwide is to close the
metadata gap highlighted here. Simultaneously, conservation geneticists, molecular ecologists
and evolutionary biologists must engage with global biodiversity assessment programs to ensure
genomic data can be collected, interpreted and archived appropriately (Brodersen & Seehausen,
2014). Several exemplary international networks (e.g. GEOBON Genetic Composition Working
Group, IUCN Conservation Genetics Specialist Group, and EU COST Action Genetic
Biodiversity Knowledge for Ecosystem resilience [GBiKE]) have already made a case for
protecting the genetic diversity of all species (Laikre et al., 2020), proposed indicators to gauge
progress toward goals (Hoban et al., 2020; Laikre et al., 2020). These groups have asserted their
rationale for these changes to stakeholders in policy documents, providing essential clarity in the
use of genetic data, and reporting against targets (Hoban et al., 2021). These actions and
advances encourage the uptake of genetic diversity monitoring by national authorities and
international bodies. The vision for many of these biodiversity monitoring networks is to develop
agile pipelines that intake raw biodiversity data and produce outputs that can directly inform
conservation policies and decisions (Hoban et al., 2021). Yet, without appropriate archival of
genomic data that includes the spatiotemporal metadata, we will be unable to deliver on the
promise of genetic diversity monitoring.
539
The GEOME datathon enabled 13 students and 7 academics from 15 institutions and 4
countries to take account of the growing metadata gap for genomics data and begin to remediate
it. The serendipity of being able to run a remote, distributed datathon due to travel restrictions
and funding reallocation forced by COVID-19, in a time when Indigenous rights, biodiversity
conservation and the value of genetic diversity have been front of mind, has not been lost on the
participants. While our efforts have just begun to address the growing metadata gap, it is our
hope that most researchers will start to ensure the FAIRness of their genomic data and metadata
before or upon publication, thereby honoring the work that went into creating it and providing
limitless opportunities for reuse of their data to help answer the important scientific questions of
the future.

Acknowledgements

This effort arose from an Evolution in Changing Seas Research Coordination Network (RCN) working group (NSF-OCE-1764316, Katie Lotterhos) and was funded by the Diversity of the Indo-Pacific Network RCN (NSF-DEB-1457848, to R.J.T.). We gratefully thank all of the authors who took the time to provide helpful responses to our metadata inquiries. We also thank Neil Davies, Chris Meyer, Beth Davis and Kiersey Nielsen for their input.

Data Availability Statement

The metadata that were recovered by the datathon are available from the Genomic Observatories Metadatabase (GEOME; geome-db.org) https://geome-db.org/workbench/project-overview?projectId=305. Relevant code for analyses in this paper may be found at https://github.com/ericcrandall/geome_metadatathon1. The meta-metadata for BioProjects that were determined to be relevant to the datathon are in Supplemental Materials S4.

Literature Cited

Allendorf, F. W. (2017). Genetics and the conservation of natural populations: Allozymes to genomes. *Molecular Ecology*, *26*(2), 420–430. doi: 10.1111/mec.13948

Baetscher, D. S., Anderson, E. C., Gilbert-Horvath, E. A., Malone, D. P., Saarman, E. T., Carr, M. H., & Garza, J. C. (2019). Dispersal of a nearshore marine fish connects marine reserves and adjacent fished areas along an open coast. *Molecular Ecology*, *1*(2), 0148–13. doi: 10/gf8rkk

Blanchet, S., Prunier, J. G., Paz-Vinas, I., Saint-Pé, K., Rey, O., Raffard, A., … Dubut, V. (2020). A river runs through it: The causes, consequences, and management of intraspecific diversity in river networks. *Evolutionary Applications*, *13*(6), 1195–1213. doi: 10.1111/eva.12941

Brauman, K. A., Garibaldi, L. A., Polasky, S., Aumeeruddy-Thomas, Y., Brancalion, P. H. S., DeClerck, F., … Verma, M. (2020). Global trends in nature's contributions to people. *Proceedings of the National Academy of Sciences*, *117*(51), 32799–32805. doi: 10.1073/pnas.2010473117

Brodersen, J., & Seehausen, O. (2014). Why evolutionary biologists should get seriously involved in ecological monitoring and applied biodiversity assessment programs. *Evolutionary Applications*, *7*(9), 968–983. doi: 10.1111/eva.12215

Buckner, J. C., Sanders, R. C., Faircloth, B. C., & Chakrabarty, P. (2021). The critical importance of vouchers in genomics. *ELife*, *10*, e68264. doi: 10.7554/eLife.68264

Buttigieg, P. L., Morrison, N., Smith, B., Mungall, C. J., Lewis, S. E., & the ENVO Consortium. (2013). The environment ontology: Contextualising biological and biomedical entities. *Journal of Biomedical Semantics*, *4*(1), 43. doi: 10.1186/2041-1480-4-43

Buttigieg, P. L., Pafilis, E., Lewis, S. E., Schildhauer, M. P., Walls, R. L., & Mungall, C. J. (2016). The environment ontology in 2016: Bridging domains with increased scope, semantic density, and interoperation. *Journal of Biomedical Semantics*, *7*(1), 57. doi: 10.1186/s13326-016-0097-6

Castañeda-Álvarez, N. P., Khoury, C. K., Achicanoy, H. A., Bernau, V., Dempewolf, H., Eastwood, R. J., … Toll, J. (2016). Global conservation priorities for crop wild relatives. *Nature Plants*, *2*(4), 1–6. doi: 10.1038/nplants.2016.22

CBD. (2020). *Global Biodiversity Outlook 5*. Montreal: United Nations. Retrieved from United Nations website: https://www.cbd.int/gbo/gbo5/publication/gbo-5-en.pdf

Cheng, S. H., Gold, M., Rodriguez, N., & Barber, P. H. (2021). Genome-wide SNPs reveal complex fine scale population structure in the California market squid fishery (Doryteuthis opalescens). *Conservation Genetics*, *22*(1), 97–110. doi: 10.1007/s10592-020-01321-2

Clark, J. S. (2010). Individuals and the Variation Needed for High Species Diversity in Forest Trees.
Science, *327*(5969), 1129–1132. doi: 10.1126/science.1183506
Cochrane, G., Karsch-Mizrachi, I., Takagi, T., & INSDC. (2016). The International Nucleotide Sequence
Database Collaboration. *Nucleic Acids Research*, *44*(D1), D48–D50. doi: 10.1093/nar/gkv1323
Cowell, C., Paton, A., Borrell, J. S., Williams, C., Wilkin, P., Antonelli, A., … Kersey, P. J. (2022). Uses
and benefits of digital sequence information from plant genetic resources: Lessons learnt from
botanical collections. *PLANTS, PEOPLE, PLANET*, *4*(1), 33–43. doi: 10.1002/ppp3.10216
Davies, N., Deck, J., Kansa, E. C., Kansa, S. W., Kunze, J., Meyer, C., … Lehnert, K. (2021). Internet of
Samples (iSamples): Toward an interdisciplinary cyberinfrastructure for material samples.
GigaScience, *10*(5). doi: 10.1093/gigascience/giab028
Davies, S. W., Putnam, H. M., Ainsworth, T., Baum, J. K., Bove, C. B., Crosby, S. C., … Bates, A. E.
(2021). Promoting inclusive metrics of success and impact to dismantle a discriminatory reward
system in science. *PLOS Biology*, *19*(6), e3001282. doi: 10.1371/journal.pbio.3001282
Deck, J., Gaither, M. R., Ewing, R., Bird, C. E., Davies, N., Meyer, C., … Crandall, E. D. (2017). The
Genomic Observatories Metadatabase (GeOMe): A new repository for field and sampling event
metadata associated with genetic samples. *PLoS Biology*, *15*(8), e2002925. doi: 10/gbqt2p
Des Roches, S., Pendleton, L. H., Shapiro, B., & Palkovacs, E. P. (2021). Conserving intraspecific
variation for nature's contributions to people. *Nature Ecology & Evolution*, *5*(5), 574–582. doi:
10.1038/s41559-021-01403-5
Díaz, S., Zafra-Calvo, N., Purvis, A., Verburg, P. H., Obura, D., Leadley, P., … Zanne, A. E. (2020). Set
ambitious goals for biodiversity and sustainability. *Science*, *370*(6515), 411–413. doi:
10.1126/science.abe1530
Dimitrova, M., Meyer, R., Buttigieg, P. L., Georgiev, T., Zhelezov, G., Demirov, S., … Penev, L. (2021).
A streamlined workflow for conversion, peer review, and publication of genomics metadata as
omics data papers. *GigaScience*, *10*(5). doi: 10.1093/gigascience/giab034
Exposito-Alonso, M., Booker, T. R., Czech, L., Gillespie, L., Hateley, S., Kyriazis, C. C., … Zess, E.
(2022). *Genetic diversity loss in the Anthropocene*. 6. doi: 10.1126/science.abn5642
Field, D., Garrity, G., Gray, T., Morrison, N., Selengut, J., Sterk, P., … Wipat, A. (2008). The minimum
information about a genome sequence (MIGS) specification. *Nature Biotechnology*, *26*(5), 541–
547. doi: 10/dms
Gaither, M. R., Gkafas, G. A., de Jong, M., Sarigol, F., Neat, F., Regnier, T., … Hoelzel, A. R. (2018).
Genomics of habitat choice and adaptive evolution in a deep-sea fish. *Nature Ecology &
Evolution*, *2*(4), 680–687. doi: 10.1038/s41559-018-0482-x
Halewood, M., Lopez Noriega, I., Ellis, D., Roa, C., Rouard, M., & Sackville Hamilton, R. (2018). Using
Genomic Sequence Information to Increase Conservation and Sustainable Use of Crop Diversity
and Benefit-Sharing. *Biopreservation and Biobanking*, *16*(5), 368–376. doi:
10.1089/bio.2018.0043
Hendricks, S., Anderson, E. C., Antao, T., Bernatchez, L., Forester, B. R., Garner, B., … Luikart, G.
(2018). Recent advances in conservation and population genomics data analysis. *Evolutionary
Applications*, *11*(8), 1197–1211. doi: 10/gd5nc2
Hinchliff, C. E., Smith, S. A., Allman, J. F., Burleigh, J. G., Chaudhary, R., Coghill, L. M., … Cranston,
K. A. (2015). Synthesis of phylogeny and taxonomy into a comprehensive tree of life.
Proceedings of the National Academy of Sciences, *112*(41), 12764–12769. doi: 10/f7vmqg
Hoban, S., Archer, F. I., Bertola, L. D., Bragg, J. G., Breed, M. F., Bruford, M. W., … Hunter, M. E.
(2022). Global genetic diversity status and trends: Towards a suite of Essential Biodiversity
Variables (EBVs) for genetic composition. *Biological Reviews*, *Early Online*. doi:
10.1111/brv.12852
Hoban, S., Bruford, M., D'Urban Jackson, J., Lopes-Fernandes, M., Heuertz, M., Hohenlohe, P. A., …
Laikre, L. (2020). Genetic diversity targets and indicators in the CBD post-2020 Global
Biodiversity Framework must be improved. *Biological Conservation*, *248*, 108654. doi:
10/gg32bm

Hoban, S., Bruford, M. W., Funk, W. C., Galbusera, P., Griffith, M. P., Grueber, C. E., … Vernesi, C.
(2021). Global Commitments to Conserving and Monitoring Genetic Diversity Are Now
Necessary and Feasible. *BioScience*, (biab054). doi: 10.1093/biosci/biab054
Hudson, M., Garrison, N. A., Sterling, R., Caron, N. R., Fox, K., Yracheta, J., … Carroll, S. R. (2020).
Rights, interests and expectations: Indigenous perspectives on unrestricted access to genomic
data. *Nature Reviews Genetics*, *21*(6), 377–384. doi: 10.1038/s41576-020-0228-x
Kardos, M., Taylor, H. R., Ellegren, H., Luikart, G., & Allendorf, F. W. (2016). Genomics advances the
study of inbreeding depression in the wild. *Evolutionary Applications*, *9*(10), 1205–1218. doi:
10.1111/eva.12414
Kumar, A., Anju, T., Kumar, S., Chhapekar, S. S., Sreedharan, S., Singh, S., … Lim, Y. P. (2021).
Integrating Omics and Gene Editing Tools for Rapid Improvement of Traditional Food Plants for
Diversified and Sustainable Food Security. *International Journal of Molecular Sciences*, *22*(15),
8093. doi: 10.3390/ijms22158093
Laikre, L. (2010). Genetic diversity is overlooked in international conservation policy implementation.
Conservation Genetics, *11*(2), 349–354. doi: 10.1007/s10592-009-0037-4
Laikre, L., Hoban, S., Bruford, M. W., Segelbacher, G., Allendorf, F. W., Gajardo, G., … Vernesi, C.
(2020). Post-2020 goals overlook genetic diversity. *Science*, *367*(6482), 1083. doi: 10/ggnjfx
Leigh, D. M., Hendry, A. P., Vázquez-Domínguez, E., & Friesen, V. L. (2019). Estimated six per cent
loss of genetic variation in wild populations since the industrial revolution. *Evolutionary*
Applications, *12*(8), 1505–1512. doi: 10.1111/eva.12810
Lendemer, J., Thiers, B., Monfils, A. K., Zaspel, J., Ellwood, E. R., Bentley, A., … Aime, M. C. (2020).
The Extended Specimen Network: A Strategy to Enhance US Biodiversity Collections, Promote
Research and Education. *BioScience*, *70*(1), 23–30. doi: 10.1093/biosci/biz140
Letunic, I., & Bork, P. (2021). Interactive Tree Of Life (iTOL) v5: An online tool for phylogenetic tree
display and annotation. *Nucleic Acids Research*, *49*(W1), W293–W296. doi:
10.1093/nar/gkab301
Lewin, H. A., Robinson, G. E., Kress, W. J., Baker, W. J., Coddington, J., Crandall, K. A., … Zhang, G.
(2018). Earth BioGenome Project: Sequencing life for the future of life. *Proceedings of the*
National Academy of Sciences, *115*(17), 4325–4333. doi: 10/gdh5vz
Liggins, L., Hudson, M., & Anderson, J. (2021). Creating space for Indigenous perspectives on access
and benefit-sharing: Encouraging researcher use of the Local Contexts Notices. *Molecular*
Ecology, *30*(11), 2477–2482. doi: 10.1111/mec.15918
Lin, D., Crabtree, J., Dillo, I., Downs, R. R., Edmunds, R., Giaretta, D., … Westbrook, J. (2020). The
TRUST Principles for digital repositories. *Scientific Data*, *7*(1), 144. doi: 10/ggwrtj
Lou, R. N., Jacobs, A., Wilder, A. P., & Therkildsen, N. O. (2021). A beginner's guide to low-coverage
whole genome sequencing for population genomics. *Molecular Ecology*, *30*(23), 5966–5993. doi:
10.1111/mec.16077
Marden, E., Abbott, R. J., Austerlitz, F., Ortiz-Barrientos, D., Baucom, R. S., Bongaerts, P., … Rieseberg,
L. H. (2021). Sharing and reporting benefits from biodiversity research. *Molecular Ecology*,
30(5), 1103–1107. doi: 10.1111/mec.15702
McCartney, A. M., Anderson, J., Liggins, L., Hudson, M. L., Anderson, M. Z., TeAika, B., … Phillippy,
A. M. (2022). Balancing openness with Indigenous data sovereignty: An opportunity to leave no
one behind in the journey to sequence all of life. *Proceedings of the National Academy of*
Sciences, *119*(4). doi: 10.1073/pnas.2115860119
McNutt, M., Lehnert, K., Hanson, B., Nosek, B. A., Ellison, A. M., & King, J. L. (2016). Liberating field
science samples and data. *Science*, *351*(6277), 1024–1026. doi: 10.1126/science.aad7048
Michonneau, F., Brown, J. W., & Winter, D. J. (2016). rotl: An R package to interact with the Open Tree
of Life data. *Methods in Ecology and Evolution*, *7*(12), 1476–1481. doi: 10.1111/2041-
210X.12593

Miraldo, A., LI, S., Borregaard, M. K., Flórez-Rodríguez, A., Gopalakrishnan, S., Rizvanovic, M., … Nogués-Bravo, D. (2016). An Anthropocene map of genetic diversity. *Science*, *353*(6307), 1532–1535. doi: 10/f9ccgr

Nosek, B. A., Alter, G., Banks, G. C., Borsboom, D., Bowman, S. D., Breckler, S. J., … Yarkoni, T. (2015). Promoting an open research culture. *Science*, *348*(6242), 1422–1425. doi: 10/gcpzwn

Pinsky, M. L., & Palumbi, S. R. (2014). Meta-analysis reveals lower genetic diversity in overfished populations. *Molecular Ecology*, *23*(1), 29–39. doi: 10.1111/mec.12509

Pope, L. C., Liggins, L., Keyse, J., Carvalho, S. B., & Riginos, C. (2015). Not the time or the place: The missing spatio-temporal link in publicly available genetic data. *Molecular Ecology*, *24*(15), 3802–3809. doi: 10.1111/mec.13254

Prada, C., Hanna, B., Budd, A. F., Woodley, C. M., Schmutz, J., Grimwood, J., … Medina, M. (2016). Empty Niches after Extinctions Increase Population Sizes of Modern Corals. *Current Biology*, *26*(23), 3190–3194. doi: 10.1016/j.cub.2016.09.039

Quattrini, A. M., Wu, T., Soong, K., Jeng, M.-S., Benayahu, Y., & McFadden, C. S. (2019). A next generation approach to species delimitation reveals the role of hybridization in a cryptic species complex of corals. *BMC Evolutionary Biology*, *19*(1), 116. doi: 10.1186/s12862-019-1427-y

Raffard, A., Santoul, F., Cucherousset, J., & Blanchet, S. (2019). The community and ecosystem consequences of intraspecific diversity: A meta-analysis. *Biological Reviews*, *94*(2), 648–661. doi: 10.1111/brv.12472

Reithmeier, R., O'Leary, L., Zhu, X., Dales, C., Abdulkarim, A., Aquil, A., … Zou, C. (2019). The 10,000 PhDs project at the University of Toronto: Using employment outcome data to inform graduate education. *PLOS ONE*, *14*(1), e0209898. doi: 10/ggfnvv

Reusch, T. B. H., Ehlers, A., Hämmerli, A., & Worm, B. (2005). Ecosystem recovery after climatic extremes enhanced by genotypic diversity. *Proceedings of the National Academy of Sciences*, *102*(8), 2826–2831. doi: 10.1073/pnas.0500008102

Riginos, C., Crandall, E. D., Liggins, L., Gaither, M. R., Ewing, R. B., Meyer, C., … Deck, J. (2020). Building a global genomics observatory: Using GEOME (the Genomic Observatories Metadatabase) to expedite and improve deposition and retrieval of genetic data and metadata for biodiversity research. *Molecular Ecology Resources*, *20*(6), 1458–1469. doi: 10/ghf4wp

Roche, D. G., Kruuk, L. E. B., Lanfear, R., & Binning, S. A. (2015). Public Data Archiving in Ecology and Evolution: How Well Are We Doing? *PLoS Biology*, *13*(11), e1002295-12. doi: 10.1371/journal.pbio.1002295

Roche, D. G., Lanfear, R., Binning, S. A., Haff, T. M., Schwanz, L. E., Cain, K. E., … Kruuk, L. E. B. (2014). Troubleshooting Public Data Archiving: Suggestions to Increase Participation. *PLoS Biology*, *12*(1), e1001779. doi: 10/ggvr85

Scholz, A. H., Freitag, J., Lyal, C. H. C., Sara, R., Cepeda, M. L., Cancio, I., … Overmann, J. (2022). Multilateral benefit-sharing from digital sequence information will support both science and biodiversity conservation. *Nature Communications*, *13*(1), 1086. doi: 10.1038/s41467-022-28594-0

Schriml, L. M., Chuvochina, M., Davies, N., Eloe-Fadrosh, E. A., Finn, R. D., Hugenholtz, P., … Walls, R. (2020). COVID-19 pandemic reveals the peril of ignoring metadata standards. *Scientific Data*, *7*(1), 188. doi: 10.1038/s41597-020-0524-5

Sibbett, B., Rieseberg, L. H., & Narum, S. (2020). The Genomic Observatories Metadatabase. *Molecular Ecology Resources*, *20*(6), 1453–1454. doi: 10/ghf4wm

Thiers, B., Bates, J., Bentley, A. C., Ford, L. S., Jennings, D., Monfils, A. K., … Pandey, J. L. (2021). Implementing a Community Vision for the Future of Biodiversity Collections. *BioScience*, *71*(6), 561–563. doi: 10.1093/biosci/biab036

Toczydlowski, R. H., Liggins, L., Gaither, M. R., Anderson, T. J., Barton, R. L., Berg, J. T., … Crandall, E. D. (2021). Poor data stewardship will hinder global genetic diversity surveillance. *Proceedings of the National Academy of Sciences*, *118*(34), e2107934118. doi: 10.1073/pnas.2107934118

Vines, T. H., Albert, A. Y. K., Andrew, R. L., Débarre, F., Bock, D. G., Franklin, M. T., … Rennison, D.
J. (2014). The Availability of Research Data Declines Rapidly with Article Age. *Current Biology*,
24(1), 94–97. doi: 10/qpm
Vines, T. H., Andrew, R. L., Bock, D. G., Franklin, M. T., Gilbert, K. J., Kane, N. C., … Yeaman, S.
(2013). Mandated data archiving greatly improves access to research data. *The FASEB Journal*,
27(4), 1304–1308. doi: 10.1096/fj.12-218164
Whitlock, M. C. (2011). Data archiving in ecology and evolution: Best practices. *Trends in Ecology*
& Evolution, *26*(2), 61–65. doi: 10.1016/j.tree.2010.11.006
Wieczorek, J., Bloom, D., Guralnick, R., Blum, S., Döring, M., Giovanni, R., … Vieglais, D. (2012).
Darwin Core: An evolving community-developed biodiversity data standard. *PLoS ONE*, *7*(1),
e29715. doi: 10.1371/journal.pone.0029715
Wilder, A. P., Palumbi, S. R., Conover, D. O., & Therkildsen, N. O. (2020). Footprints of local adaptation
span hundreds of linked genes in the Atlantic silverside genome. *Evolution Letters*, *4*(5), 430–
443. doi: 10.1002/evl3.189
Wilkinson, M. D., Dumontier, M., Aalbersberg, Ij. J., Appleton, G., Axton, M., Baak, A., … Mons, B.
(2016). Comment: The FAIR Guiding Principles for scientific data management and stewardship.
Scientific Data, *3*(1), 1–9. doi: 10/bdd4
Willette, D. A., Allendorf, F. W., Barber, P. H., Barshis, D. J., Carpenter, K. E., Crandall, E. D., … Seeb,
J. E. (2014). So, you want to use next-generation sequencing in marine systems? Insight from the
Pan-Pacific Advanced Studies Institute. *Bulletin Of Marine Science*, *90*(1), 79–122. doi:
10.5343/bms.2013.1008
Winter, D. J. (2017). rentrez: An R package for the NCBI eUtils API. *The R Journal*, *9*(2), 520–526. doi:
10.32614/rj-2017-058
Yilmaz, P., Kottmann, R., Field, D., Knight, R., Cole, J. R., Amaral-Zettler, L., … Glöckner, F. O.
(2011). Minimum information about a marker gene sequence (MIMARKS) and minimum
information about any (x) sequence (MIxS) specifications. *Nature Biotechnology*, *29*(5), 415–
420. doi: 10.1038/nbt.1823

807
808 **Box 1.** Summary of metadata practices encountered during the datathon that hindered metadata recovery
809 and recommended practices to improve future usability of samples and genetic sequence data. In general,
810 we recommend that authors use metadata software (GEOME: geome-db.org, COPO: copo-project.org, or
811 museum database software such as Specify) to organize and archive their sample metadata.
812

Practice	Challenge	Solution
INSDC materialSampleID (SRA) does not match any sample identifiers in associated scientific publication(s)	Metadata external to the INSDC cannot be assigned to genetic samples or metadata are associated with the wrong sample	Use consistent, persistent, globally unique sample identifiers (e.g. Darwin Core *materialSampleID*) across data repositories and publications; if sample identifiers are not consistent, provide an explicit cross-reference table in all associated publications and data repositories
Large amounts of metadata are only available in associated publications in PDF format & lack consistent formatting	Metadata cannot be programmatically converted to standard table formats (e.g. entries formatted column by row, rather than row by column) & require time-consuming manual extraction	Provide metadata in comma- or tab-delimited files (.csv or .txt) using standard column headers (i.e. terms suggested by Darwin Core, MIxS, or GEOME or COPO) & associated vocabulary, where possible
Specimens, BioSamples or metadata are deposited in a biodiversity collection (e.g., museum, herbarium, biobank, zoo), but biodiversity collection accession numbers are not provided in the associated publications or INSDC	Biodiversity collection record searches can be time-consuming & may not yield enough information to link samples in collection databases back to INSDC databases	Use consistent sample identifiers across all databases & publications; or provide a cross-reference table in associated publication(s) that links biodiversity collection accessions to INSDC materialSampleIDs and identifies the biocollection repository by name. The Darwin Core standard accommodate multiple terms
Associated publication references previous publication(s) for details about the sample metadata	Time consuming & challenging to track citation trail back to metadata in original/earlier publication. Sample metadata or identifiers may be absent or in inconsistent formats across associated publications	Compile & include relevant metadata from previous publications in supplementary materials for the publication linked to the INSDC BioProject; if needed, include a column flagging whether data are new to the present study or originated from another source (& identify that source)
Sample collection geospatial coordinates or location name withheld in order to protect endangered species, sensitive habitat, or Indigenous sovereignty	Sample lacks spatial metadata	Provide imprecise geospatial coordinates & use large, defined *coordinateUncertainty* to maintain local anonymity of sensitive collection sites. Provide additional comments using Darwin Core *informationWithheld*
Codes used to abbreviate sample collection locations are inconsistent or hard to find throughout publication & related materials	Sample collection locations cannot be determined or require time-consuming manual curation	Include site codes in the sample identifier (*materialSampleID*). Use consistent site codes throughout associated publications & provide a key with codes & geospatial coordinates in associated publications
Sample collection dates are a range or a season (e.g., winter 2017-2018)	Sample collection date may not be identifiable to a specific year; unclear which samples were collected in which date range	Denote/report which year <u>each</u> sample was collected in (dates that also include month & day are ideal)

No metadata on BioSample relevance to genetic diversity of wild populations/species provided	Unclear which BioSamples (if any) were collected from wild populations versus e.g. brood stock, laboratory stock, domesticated species, artificial selection experiments, non-wild collections in seed banks	Provide metadata denoting which BioSamples were collected from wild populations, using Darwin Core term *establishmentMeans*
Metadata provided for some but not all BioSamples	Some BioSamples lack metadata, unclear why metadata are incomplete	Provide metadata for all BioSamples or list a specific reason for missing metadata (e.g. not collected, metadata lost, sample excluded from study due to misidentification) using Darwin Core *informationWithheld*.
Sampling location provided, but only at a coarse geographic scale i.e., state, province, or country name	Sample lacks spatial metadata at a resolution useful for future monitoring & macrogenetic research questions	Provide geospatial coordinates for sample collection locations. Specific place, state, & country names can be helpful additions to confirm the geospatial coordinates are correct (& to programmatically filter by broader geographic locations)

Table 1: Alphabetized list of required (in bold) and recommended metadata terms for individual organisms and/or derived tissues or DNA sequences included in the datathon. Square brackets in the definition column denote the metadata standard from which the definition comes. Terms with multiple definitions are in order of decreasing specificity. The importance column indicates which terms support the identification of Indigenous provenance and can therefore inform Access and Benefit-Sharing (ABS), and those that can support sample or Digital Sequence Information (DSI) re-use in conservation, according to the study approach definitions of Leigh et al. (2021). Class I studies generate new sequence data, requiring precise information regarding the spatiotemporal context of the collected sample, a unique materialSampleID, as well as the preservative the tissue is held in; Class II studies compile genetic diversity values from published studies, generally requiring less precise spatiotemporal information, but this needs to be associated with a publication (associatedReferences); Class III studies re-analyse digital sequence information, or derived genetic data, requiring precise spatiotemporal information, and a unique materialSampleID. Depending on the objective of re-use, habitat and environmental_medium may also be important for sample/DSI re-use in conservation. Controlled vocabularies refer to standardized lists of acceptable entries, often defined by a standards organization.

Term	Definition	Importance	Controlled Vocabulary	Example
associatedReferences	[GEOME][1] Any associated publications/references pertaining to this individual or its derivative tissues or sequences. The first place it was published is particularly relevant. DOIs in format: https://doi.org/10.1007/s10530-007-9196-8. Multiple DOIs separated by |. [Darwin Core][2] A list (concatenated and separated) of identifiers (publication, bibliographic reference, global unique identifier, URI) of literature associated with the Occurrence.	Indigenous provenance, Class II	None	"https://doi.org/10.1111/j.1365 https://doi.org/10.5343/bms.20
coordinateUncertaintyInMeters	[Darwin Core] The horizontal distance (in meters) from the given decimalLatitude and decimalLongitude describing the smallest circle containing the whole of the locality where the sample could possibly have come from. Value empty if the uncertainty is unknown, cannot be estimated, or is not applicable (because there are no coordinates). Zero is not a valid value for this term.	Class I, III	None	1 km = "1000"

country	[Darwin Core] The name of the country or major administrative unit or exclusive economic zone (for marine samples) in which the locality occurs.	Indigenous provenance, Class II	ISO 3166-1	"Indonesia"
decimalLatitude	[Darwin Core] The geographic latitude (in decimal degrees, using the spatial reference system given in geodeticDatum) of the geographic center of a Location. Positive values are north of the Equator, negative values are south of it. Legal values lie between -90 and 90, inclusive.	Indigenous provenance, Class I, II, III	None	"-6.147183"
decimalLongitude	[Darwin Core] The geographic longitude (in decimal degrees, using the spatial reference system given in geodeticDatum) of the geographic center of a Location. Positive values are east of the Greenwich Meridian, negative values are west of it. Legal values lie between -180 and 180, inclusive.	Indigenous provenance, Class I, II, III	None	"105.46326"
derivedDataFilename	[GEOME] A list (concatenated and separated with |) of the file names for datasets that include data derived from this tissue that are accessible via the 'derivedDataURI'. Could be a compressed archive.	Class III	None	"SDM_snps.tar.gz"
derivedDataFormat	[GEOME] A list (concatenated and separated with |) of the dataset formats relating to the 'derivedDataType' that include data derived from this tissue.	Class III	{microsatellites, Sequence alignment, SNPs,OTUs,ASVs, Other}	"SNPs"
derivedDataType	[GEOME] A list (concatenated and separated with |) of the dataset types that include data derived from this tissue.	Class III	{genepop,FASTA,VCF, nexus,PHYLIP, structure,Other}	"VCF"
derivedDataURI	[GEOME] A URI (preferably a DOI in this format: https://doi.org/10.1007/s10530-007-9196-8) for any datasets that include data derived from this tissue. Multiple URIs/DOIs can be separated by |.	Class III	None	"https://doi.org/10.5061/dryad.
environmental_medium	[MIxS][3] Terms that identify the material displaced by the entity at time of sampling. Recommend subclasses of environmental material [ENVO:00010483]. Multiple terms can be separated by pipes e.g.: a duck might displace fresh water|air.		ENVO[4] Environmental N ENVO_00010483	"sea water"

habitat	[MIxS: Broad-scale environmental context] In this field, report which major environmental system your sample or specimen came from. The systems identified should have a coarse spatial grain, to provide the general environmental context of where sampling was done. [Darwin Core] A category or description of the habitat in which the Event occurred.		ENVO Biomes: ENVO_00000428	"marine benthic biome"
locality	[Darwin Core] The specific name or description of the site or place where the sample was taken as given by the original researchers. This would be the place name that appears in a table next to the coordinates, or the labels for sampling sites on a map. Less specific geographic information can be provided in other geographic terms (continentOcean, country, stateProvince, island). This term may contain information modified from the original to correct perceived errors or standardize the description.	Indigenous provenance, Class I, II, III	None	"Rakata"
materialSampleID	[GEOME] The collector's specimen number. This number must be unique among the IDs within the sheet. [Darwin Core]. An identifier for the MaterialSample (as opposed to a particular digital record of the material sample). In the absence of a persistent global unique identifier, construct one from a combination of identifiers in the record that will most closely make the materialSampleID globally unique.	Class I, III	None	"Rakata_1190.01"
permitInformation	[GEOME] Information regarding the permit acquired to collect this sample. At least the permit number and issuing authority. Multiple values separated by |	Indigenous provenance,	None	"Indonesian Institute of Sciences Permit #'s 1187/ SU/KS/2006 an 04239/SU.3/KS/2006"
preservative	[GEOME] Preservative used on the specimen.	Class I	GEOME List of preserva	"95% Ethanol"
yearCollected	[Darwin Core] The year the collecting event took place	Class I, II, III	None	March 24, 2006 = "2006"

1. Deck et al. (2017); Riginos et al. (2020)

2. Wieczorek et al. (2012)
3. Yilmaz et al. (2011)
4. Buttigieg et al. (2013, 2016)