Two gustatory receptors involved in host plant recognition of silkworm larvae

Haruka Endo[1,2,4,5,*], Kana Tsuneto[2,4], Takayuki Yamagishi[2], Dingze Mang[2,6], Katsuhiko Ito[3],

Shinji Nagata[1] and Ryoichi Sato[2,*]

[1]Department of Integrated Bioscience, Graduate School of Frontier Sciences, The University of

Tokyo, Kashiwa, Chiba 277-8562, Japan

[2]Graduate School of Bio-Applications and Systems Engineering, Tokyo University of Agriculture and

Technology, Koganei, Tokyo 184-8588, Japan

[3]Department of Science of Biological Production, Graduate School of Agriculture, Tokyo University

of Agriculture and Technology, Fuchu,Tokyo 183-8509, Japan

[4]The authors contributed equally.

[5]Present address: RIKEN Center for Brain Science, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan

[6]Present address: College of Life Science, Institute of Life Science and Green Development, Hebei

University, Baoding 071002, China

*Correspondence: haruka.endo@riken.jp (H.E.) or ryoichi@cc.tuat.ac.jp (R.S.).

Abstract

Herbivorous insects can identify their host plants by sensing plant secondary metabolites as chemical

cues. We previously reported the two-factor host acceptance system of the silkworm *Bombyx mori*

larvae. The chemosensory neurons in the maxillary palp (MP) of the larvae detect mulberry secondary

metabolites, chlorogenic acid (CGA), and isoquercitrin (ISQ), with ultrahigh sensitivity, for host plant

recognition and feeding initiation. Nevertheless, the molecular basis for the ultrasensitive sensing of

these compounds remains unknown. In this study, we demonstrated that two gustatory receptors

(Grs), BmGr6 and BmGr9, are responsible for sensing the mulberry compounds with attomolar

sensitivity for host plant recognition by silkworm larvae. Calcium imaging assay using cultured cells

expressing the silkworm putative sugar receptors (BmGr4-10) revealed that BmGr6 and BmGr9 serve

as receptors for CGA and ISQ with attomolar sensitivity in human embryonic kidney 293T cells.

CRISPR/Cas9-mediated knockout (KO) of BmGr6 and BmGr9 resulted in a low probability of making

a test bite of the mulberry leaves, suggesting that they lost the ability to recognize host leaves

specifically. Electrophysiological recordings showed that the loss of host recognition ability in the Gr-

KO strains was due to a drastic decrease in MP sensitivity toward ISQ in BmGr6-KO larvae and

toward CGA and ISQ in BmGr9-KO larvae. Our findings unraveled that the two Grs, which have been

regarded as sugar receptors, are molecules responsible for detecting plant phenolics in host plant

recognition.

Keywords: herbivorous insects, host-plant selection, plant secondary metabolites; silkworm;
gustatory receptors

Introduction

Herbivorous insects and plants have evolved through reciprocal adaptation. Plants have various
physical and chemical defense mechanisms to combat insect herbivory [1]. For instance, plants have
evolved a variety of secondary metabolites, most of which act as feeding deterrents and are often
toxic to herbivorous insects [2]. In turn, herbivorous insects have adapted to some of the chemicals
by developing chemosensory systems, enzymatic detoxication, and other mechanisms, which
enabled them to consume a specific limited range of plants [3,4]. In parallel, they were able to identify
host plants using plant secondary metabolites as sensory cues [5].

Sensory exploration at the leaf surface is essential for host plant selection in larval feeding and adult
ovipositioning [6,7]. We previously reported that the silkworm *Bombyx mori* larvae, well known for
their preference for mulberry leaves, feed on the leaves utilizing a two-factor authentication system
controlled by two taste organs and a set of chemoreceptors with differing sensitivities, operating
sequentially in time [8]. The first step involves probing the leaf surface with the maxillary palp (MP),
which contains ultrasensitive chemosensory neurons sensitive to a set of crucial mulberry compounds,
including secondary metabolites (chlorogenic acid [CGA] and quercetin glycosides) and a phytosterol
(β-sitosterol). The ultrahigh sensitivities in the attomolar and femtomolar ranges enable the larvae to
detect trace amounts of these compounds at the dry leaf surface [6,9] and identify the host leaves. If
these three compounds are detected simultaneously, the larva makes a test bite, which leads to the
second step. Another gustatory organ, the maxillary galea (MG), is used to probe the sugar content
released by the test bite. Sucrose mainly induces continued biting, resulting in feeding. The two-factor
host acceptance system is expected to be conserved among herbivorous insects because of their
typical stepwise feeding behavior, composed of foraging, palpation at the leaf edge, test and
persistent biting, and swallowing [10,11]. Although understanding the behavioral and neural basis of
host acceptance in silkworm larvae has progressed, the molecular basis underlying MP's
ultrasensitive detection of the mulberry compounds remains unknown.

Insects have three prominent chemosensory receptor families: gustatory (Grs), olfactory (Ors), and
ionotropic receptors (Irs) [12]. Most Grs and some Irs are believed to mediate taste. Research on
insect chemosensory receptors is advanced in the fruit fly, *Drosophila melanogaster*. *Drosophila* Gr5a
and Gr64a-f are broad-tuned sugar receptors [13–15]. *Drosophila* Gr43a and its orthologs in other
insect species have been reported as fructose-specific receptors [16,17], whereas some lepidopteran
Grs in the same lineage respond to *myo*-inositol [18,19]. Some Grs in *Drosophila* are receptors for
toxic compounds, including plant secondary metabolites [20], but many of these "bitter" Grs do not
have clear ortholog relationships across insect orders. Some Irs were reported to detect amino acids
and fatty acids in *Drosophila* [21,22].

Some Grs have been reported as receptors for plant secondary metabolites in lepidopteran insects. PxutGr1 from the swallowtail butterfly, *Papilio xuthus*, is responsible for the sensitivity of tarsal taste sensilla to synephrine, which acts as an ovipositioning stimulant [23]. PrapGr28 from the cabbage butterfly, *Pieris rapae*, contributes to sinigrin sensitivity in adult tarsal taste sensilla [24], although sinigrin is a feeding stimulant for their larvae [25]. PxylGr34 from the diamondback moth, *Plutella xylostella*, is involved in response to canonical plant hormones, brassinolide, and 24-epibrassinolide, a feeding and oviposition deterrent, in MG of larvae [26]. BmGr16, BmGr18, and BmGr53 from *B. mori* respond to coumarin and caffeine in human embryonic kidney 293T (HEK293T) cells [27], but their role *in vivo* is unknown. The two Ors in *P. xylostella*, Or35 and Or49, are necessary and sufficient for detecting isothiocyanates, which act as ovipositioning stimulants and its host cruciferous plants [28]. Thus, several examples of low-sensitive chemoreceptors for plant secondary metabolites in the foreleg of adult and MG of larvae have been reported. However, ultrahigh-sensitive chemoreceptors in larval MP underlying the first step in the two-factor host acceptance system are unknown.

We addressed the molecular basis of host recognition in silkworm larvae in the present study. *In vitro* calcium imaging in the cells suggested that BmGr6 and BmGr9 are candidates that confer MP ultrahigh sensitivity for CGA and ISQ. CRISPR/Cas9-mediated KOs demonstrate that the two Grs are responsible for host recognition by sensing CGA and/or ISQ.

Results

BmGr6 and BmGr9 serve as ultrahigh-sensitive receptors for CGA and ISQ in HEK293T cells and sugar receptors

We started with phylogenetic analysis to find closely related silkworm Grs with PxutGr1, a candidate receptor for CGA and ISQ because PxutGr1 is the only Gr reported as a receptor for plant phenolics [23]. Phylogenetic analysis of lepidopteran Grs indicates that PxutGr1 belongs to the same lineage as BmGr9 and BmGr10, *Drosophila* Gr43a orthologs in the silkworm (Fig. 1). Based on the phylogenetic analysis, we focused on BmGr9 and BmGr10 in the Gr43a clade, and BmGr4-8 in the neighboring putative sugar receptor clade [29].

To test whether these seven Grs can confer silkworm MP chemosensory neurons attomolar sensitivity to CGA and ISQ, we performed calcium imaging in HEK293T cells. Of the seven Grs-expressing cells, BmGr6-, BmGr8-, and BmGr9-expressing cells responded to 100 aM CGA, and BmGr6- and BmGr9-expressing cells responded to 100 aM ISQ (Fig. 2A-C). We recorded a dose-dependent response of BmGr6, BmGr8, BmGr9, and BmGr10 (Fig. 2D and E) and determined the EC_{50} values (Table S1). Although BmGr6- and BmGr9-expressing cells responded to <10 aM CGA and ISQ, BmGr8-expressing cells did not respond to 10 aM CGA (Fig. 2D). Based on these results, we focused on and further characterized BmGr6 and BmGr9 as candidates conferring MP neurons attomolar sensitivities toward CGA and ISQ. BmGr6- and BmGr9-expressing cells did not respond to β-sitosterol up to 2 μM (Fig. S1).

Since BmGr6 is regarded as a putative sugar receptor [29] and BmGr9 was reported as a fructose
receptor [16], we investigated their response to major sugars present in the mulberry leaves (D-
fructose, D-glucose, sucrose, and *myo*-inositol) [19]. Surprisingly, BmGr9-expressing cells responded
to D-fructose with aM sensitivity (Fig. 2C and D), whereas they exhibited relatively lower sensitivities
to D-glucose, sucrose, and *myo*-inositol (Fig. 2E and F). BmGr6-expressing cells showed pM to nM
sensitivity to four sugars (Fig. 2C-F). These results indicate that the two Grs are highly tuned to CGA
and ISQ, whereas BmGr9 also exhibits a high affinity with D-fructose.
119

BmGr6 and BmGr9 are expressed in subsets of MP chemosensory neurons
According to the RNA-seq data from a previous study, BmGr6 and BmGr9 are highly expressed in
the MP [30]. To confirm this at the protein level, we performed immunohistochemistry using anti-
BmGr6 and anti-BmGr9 antisera. Each sensillum in MP contains three to five chemosensory neurons
whose somas are usually clustered in the second segment (Fig. 3A and 3B) [31]. We observed three
BmGr6-antiserum-positive MP neurons (Fig. 3C). In the two representative cryosections, two of the
three neurons are close to each other (Fig. 3C), suggesting that the two neurons are contained in the
same sensillum. We observed seven BmGr9-antiserum-positive MP neurons, three of which are close
to each other (Fig. 3D). Thus, BmGr6 and BmGr9 are expressed in at least three and seven neurons
in the MP of silkworm larvae.
130

BmGr6-KO and BmGr9-KO larvae lose their ability to recognize mulberry leaves specifically
To test whether the two Grs are involved in host recognition *in vivo*, we generated BmGr6-KO and
BmGr9-KO silkworm strains via the CRISPR/Cas9 system. We designed single-guide RNAs
(sgRNAs) targeting the exons 5 and 3 of the *BmGr6* and *BmGr9* genes (Fig. 4A). By injecting the
sgRNAs and the Cas9 protein into eggs, we successfully established mutant strains carrying
homozygous null mutations of *BmGr6* and *BmGr9* (material and methods for details). In both strains,
premature stop codons were generated via indel mutations (Fig. 4A and S2).
The body mass of the fifth instar larvae of BmGr6-KO and BmGr9-KO strains, which have been fed
with artificial diets since hatching, were ~86% and ~91% of the body mass of wild-type (WT) larvae
(Fig. 4B). First, we conducted 1.5 h short-term feeding assay using artificial diets and mulberry leaves.
When feeding on leaves, the larvae must recognize host leaves by MP to initiate feeding (the first
step of the host acceptance system) [8]. This step can be skipped when feeding artificial diets with
abundant sugars on their surface are available. The feeding amount of artificial diets by BmGr6-KO
larvae was ~85% of the WT larvae (Fig. 4C), suggesting that the low body mass of BmGr6-KO larvae
is due to the low feeding amount of artificial diets. The feeding amount of BmGr9-KO larvae is
comparable with that of WT (Fig. 4C), suggesting that factors that affect long-term feeding, such as
the interval of feeding, are involved in the low body mass of BmGr9-KO larvae. The feeding amounts
of mulberry leaves by the two mutant larvae were significantly lower than the WT (Fig. 4D). To directly
observe the ability of host recognition, we next observed test bites toward mulberry leaves. Only 33%

and 35% of BmGr6-KO and BmGr9-KO larvae made test bites within 1 min, whereas 90% of WT
larvae made test bites (Fig. 4E). This low probability of test biting corresponds to those of some
nonhost leaves in WT larvae [8], suggesting that the two mutant strains lose their ability to recognize
mulberry leaves. Even when BmGr6-KO and BmGr9-KO larvae were making test bites within 1 min,
and their palpation time until making a first test bite was significantly longer than that of WT larvae
(Fig. 4F), suggesting that even if mutant larvae made a test bite, they were not able to recognize
mulberry leaves immediately. These results indicate that BmGr6 and BmGr9 are essential for host
recognition *in vivo*.

BmGr6 and BmGr9 are responsible for sensing ISQ and both CGA and ISQ, respectively: *in*
vivo
We performed electrophysiological recordings from MP to test whether the two Grs are involved in
CGA and ISQ sensing *in vivo*. The MP neurons of BmGr6-KO larvae started responding to CGA from
10 aM and showed almost the same sensitivity as WT larvae (Fig. 5A and B); they did not respond to
ISQ up to 1 μM (Fig. 5A and C). This suggests that the ISQ sensitivity of BmGr6-KO larvae was >10^{10}-
fold lower than the MP neurons of WT larvae. By contrast, both CGA and ISQ sensitivities of BmGr9-
KO larvae were ~1,000-fold lower than those of WT larvae (Fig. 5A-C), suggesting that BmGr9 is
required for attomolar sensitivities of both CGA and ISQ. We previously reported that the β-sitosterol
sensitivity of MP neurons is at the femtomolar level [8]. The number of spikes toward 10 fM β-sitosterol
was comparable between WT, BmGr6-KO, and BmGr9-KO strains (Fig. S3), suggesting that MP of
the two mutant strains preserved the sensitivity to β-sitosterol. BmGr6 and BmGr9 are required for
the attomolar sensitivity to ISQ, and BmGr9 is required for attomolar sensitivity to CGA; nevertheless,
BmGr6 is not involved in CGA sensing *in vivo*. Unlike *in vitro* study observations in Fig. 2, these results
indicate different roles of BmGr6 and BmGr9 *in vivo*.

Discussion
Functional unit of Grs: molecular basis of CGA and ISQ sensing
Ligand identification of insect Grs using heterologous expression systems (e.g., human cultural cells
and *Xenopus* oocytes) has often been unsuccessful and yielded little information. In the present study,
we identified some ligands of silkworm Grs. The EC_{50} value of BmGr9 to D-fructose from the present
study (aM level) is significantly lower than that previously reported by Sato *et al.* (mM level) [16]. They
did not observe BmGr9 response to D-glucose, sucrose, and *myo*-inositol up to 100 mM using calcium
imaging assay in the HEK293T cells [16]. Given that the detection sensitivity in our condition is much
higher because of the improvement of Gr expression levels and other unknown reasons, results from
Sato *et al.* and the present study were consistent. For instance, D-fructose exhibited the highest
affinity to BmGr9 among the tested sugars in our experiment (Fig. 2G), which was not inconsistent
with the negative results for D-glucose, sucrose, and *myo*-inositol in Sato *et al.* [16]. Meanwhile, the
CRISPR/Cas9-mediated KOs of BmGr6 and BmGr9 demonstrated that they could confer the MP

attomolar sensitivity to the mulberry compounds (Fig. 5) in partial agreement with the results from
calcium imaging using HEK293T cells (Fig. 2).
Our results confirm that single Gr can fully function as a receptor *in vitro*; nevertheless, it is evident
that heterologous expression of single Gr does not fully reconstitute its function *in vivo*. For example,
although BmGr6 served as a CGA receptor with attomolar sensitivity in HEK-293T cells (Fig. 2), the
KO of BmGr6 did not influence the MP sensitivity of CGA (Fig. 5). This is not due to functional
redundancy with other Grs, as the KO of BmGr9 resulted in a drastic decrease in CGA sensitivity (Fig.
5). It is possible that BmGr6 in MP neurons does not function as a homomeric complex but function
as a heteromeric complex by coupling with other Grs, as proposed in Ors in general and in some Grs
[32–35], resulting in different ligand spectra and sensitivity from the homomeric complex. Otherwise,
the difference in the expression levels and other factors between HEK293T cells and MP neurons
may affect the contribution of BmGr6 to the CGA sensitivity in MP neurons. Another example is that
both BmGr6 and BmGr9 are indispensable for ISQ sensing with attomolar sensitivity (Fig. 5),
suggesting a synergism between BmGr6 and BmGr9 to realize the attomolar sensitivity. In this case,
too, one possibility is that the two Grs couples in a functional heteromeric complex. Altogether, our
results suggest that BmGr6 and BmGr9 function as a heteromeric complex in MP neurons. Although
the heterologous expression of single Grs significantly contributed to the screening for Grs with
attomolar sensitivity to CGA and ISQ, its utility for dissecting *in vivo* molecular gustatory systems is
limited. We cannot rule out the possibility that Grs with lower sensitivities to CGA and ISQ than BmGr6
and BmGr9 *in vitro* are involved in attomolar sensing in MP by synergism with the two Grs. To
understand insect gustatory systems, it is required to decipher functional combinations of Grs and the
mechanism determining their ligand spectra.
210
Putative fructose and sugar Grs in the silkworm act as plant phenolics receptors
We demonstrate that BmGr6 and BmGr9 are responsible for ultrahigh-sensitive detection of CGA
and/or ISQ. The clades to which BmGr6 and BmGr9 belong are known as sugar receptor clades [29].
Nevertheless, we found that they are highly tuned to CGA and ISQ *in vitro* and responded to sugars
at relatively high concentrations (Fig. 2). As there was no drastic decrease in the feeding amount of
the artificial diet in BmGr6-KO and BmGr9-KO larvae in comparison with the WT (Fig. 4C), it is unlikely
that BmGr6 and BmGr9 also act as sucrose receptors in the MG. Unfortunately, we could not provide
direct evidence for that because we failed to record spikes from sugar-sensing neurons in the MG of
the N4 silkworm strains (both WT and mutants), probably because of their small amplitudes. Our data
suggest that the two Grs act as plant phenolics receptors rather than sugar receptors in silkworm
larvae's gustatory system. Phenolics are the most widely distributed plant secondary metabolites and
serve as stimulants and deterrents toward insects in feeding and oviposition [36]. Some Grs in the
putative sugar and fructose receptor clades may be receptors for plant phenolics which are
responsible for host plant selection in herbivorous insects.
225

The attomolar sensitivity is required for sensing leaf surface compounds in host plant selection

The experiments using BmGr6-KO and BmGr9-KO larvae provided further insights into the host recognition by MP. BmGr6-KO larvae lost their ability to sense ISQ but could still detect the remaining two essential compounds (CGA and β-sitosterol), inducing a test bite with the same ultrahigh sensitivity as the WT larvae (Fig. 5 and S3). Their low probability of test biting toward mulberry leaves (Fig. 4) confirmed the obligate authentication of the three compounds by MP. BmGr9-KO larvae have ~1,000-fold lower sensitivity toward CGA and ISQ. However, they exhibited high sensitivity even at the femtomolar level (Fig. 5), resulting in a lack of host recognition ability with high probability (Fig. 4E). This indicates that the femtomolar sensitivity is insufficient, and the attomolar sensitivity is required to sense leaf surface secondary compounds for host recognition. We recently showed silkworm MP's attomolar sensitivity toward feeding deterrents, such as coumarin and strychnine nitrate [37]. These ultrahigh sensitivities underlie feeding decisions by herbivorous larvae at the leaf surface.

In summary, we identified the two silkworms Grs responsible for the ultrahigh sensitivity in MP to sense leaf surface host phenolic compounds and recognize host leaves. Mechanistic insight into the gustatory system in herbivorous insects, especially on a functional unit of Grs, is needed to know how their Grs have adapted their tunings to plant secondary metabolites and their contribution to the evolution of feeding preferences.

Acknowledgments

The authors thank Shingo Kikuta (Ibaraki University) and Masataka G. Suzuki (The University of Tokyo) for technical assistance in calcium imaging at an early stage of this study and for establishing the CRISPR-Cas9 system. The Japan Society supported this research for the Promotion of Science (JSPS) KAKENHI (Grant Number 25660040 for RS; 18J00733 for HE).

Author contributions

H.E., K.T., and R.S. designed the research; H.E. and K.T. performed calcium imaging and electrophysiological recording studies; K.T. performed immunostaining and behavioral experiments; H.E., T.Y., and D.M. generated knockout mutants under the supervision of K.I. and S.N. H.E. wrote the manuscript with intellectual inputs from all authors.

Conflict of interest

The authors declare no conflicts of interest.

Material and methods

Insect rearing

The silkworm *Bombyx mori* larvae (N4 strain and Kinshu × Showa hybrid) were reared at 25°C on an artificial diet Silkmate 2M (Nihon-Nosan Co. Ltd.) under a 12 h light–12 h dark photoperiod. The N4

larvae were used for experiments in Fig. 2, 4, 5, S3, and S4. The Kinshu × Showa larvae were used for immunostaining in Fig. 3.

Phylogenetic analysis

The lepidopteran Gr phylogeny was generated using full-length amino acid sequences of Grs from *Bombyx mori* [30], *P. xylostella* [38], *H. melpomene* [39], PxutGr1 [23], and PrapGr28 [24]. The sequences were aligned with MAFFT v.7 (http://mafft.cbrc.jp/alignment/server/). The IQ-TREE web server (http://iqtree.cibiv.univie.ac.at/) generated the maximum-likelihood tree using the JTT+F+I+G4 model. The phylogenetic tree was visualized using iTOL (https://itol.embl.de/) [40].

cDNA cloning of *Bombyx* Grs

cDNA of BmGr6, BmGr7, BmGr9, and BmGr10 were cloned into the pcDNA3.1 vector (Thermo Fisher Scientific) as reported previously [19,41]. According to the nucleotide sequences reported by Guo *et al.* [30], cDNA of BmGr4, BmGr5, and BmGr8 were codon-optimized for mammalian cell expression, artificially synthesized (gBlocks®, Integrated DNA Technologies), and inserted into the EcoRV site of pcDNA3.1 vector using GeneArt® Seamless Cloning and Assembly Enzyme Mix (Thermo Fisher Scientific).

Cell culture and transfection

HEK293T cells were cultured in Dulbecco's modified Eagle medium (DMEM, Thermo fisher scientific) supplemented with 10% fetal bovine serum (BioWest), 4 mM GlutaMAX (Thermo Fisher Scientific), 100 units/ml penicillin (Meiji Seika Pharma Co., Ltd.), and 100 mg/ml streptomycin (Meiji Seika Pharma Co., Ltd.) with 5% CO_2 at 37°C. For preparing transfection solutions, 500 µL Opti-MEM (Thermo Fisher Scientific) were vortexed with 4–6 µg polyethylenimine (PEI Max, Polysciences, Inc., Warrington, PA) and incubated for 5 min. Afterward, 2 µg of an expression vector was added, vortexed vigorously, and incubated for 20 min. Approximately 50%–70% confluent HEK293T cells were seeded on a coverslip in a six-well plate and transfected with the transfection solution for 4 h with 5% CO_2 at 37°C. The media was replaced with DMEM, and the cells were incubated for 48 h.

Calcium imaging

For preparing the loading solution of Ca^{2+} indicator Fluo-4 AM (Thermo Fisher Scientific), 3 µL of 1 mM Fluo-4 AM (stock solution in DMSO) and 30 µL of PowerLoad Concentrate (100X, Thermo Fisher Scientific) were vortexed vigorously. The mixture (5 µL) was added into a six-well plate containing 500 µL Hank's buffered saline solution (HBSS, 137 mM NaCl, 5.4 mM KCl, 0.3 mM Na_2HPO_4, 0.4 mM KH_2PO_4, 4.2 mM $NaHCO_3$, 1.3 mM $CaCl_2$, 0.5 mM $MgCl_2$, and 0.4 mM $MgSO_4$, pH 7.4) incubated for 10 min at 37°C (final conc. ~1 µM Fluo4-AM and ~1X PowerLoad). After loading, the solution was replaced with HBSS containing 0.75 mM Probenecid (Thermo Fisher Scientific). After 1 h incubation, the cells were used for calcium imaging experiments.

Calcium imaging was performed with a flow system under the fluorescence microscope BX53
(Olympus) equipped with a MetaView imaging system (Molecular Devices). A coverslip on which
transfected cells were seeded was inverted and put on a self-manufactured chamber with a 5-mm-
wide flow channel. A ligand dissolved in HBSS was superfused at a proper flow rate with peristaltic
pumps (PERISTA pump, ATTO). CGA (Nacalai tesque, Kyoto, Japan), ISQ (Extrasynthese, Lyon,
France), D-glucose, D-fructose, sucrose, and *myo*-inositol (Fujifilm Wako Pure Chemical Corp.,
Osaka, Japan) were directly dissolved in HBSS. β-sitosterol (Abcam, Cambridge, UK) was dissolved
in a 45% solution of 2-hydroxypropyl-β-cyclodextrin at 37°C to prepare stable aqueous solutions. The
time-lapsed fluorescence images (1 Hz) were processed using Fiji [42], and the fluorescent ratio
($\Delta F/F$) was calculated as peak intensity divided by basal intensity (mean of 10 flames before ligand
perfusion). The EC_{50} value was calculated by fitting using the four-parameter logistic model (Prism
ver.8, Graph-Pad).

Immunostaining
The heads of the fifth instar larvae (Kinshu × Showa hybrid) were fixed with 4%
paraformaldehyde/phosphate-buffered saline (PBS, 137 mM NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4,
1.8 mM KH_2PO_4, pH 7.4) overnight at 4°C. The fixed heads were dissected, and the maxilla was
collected and immersed into a cryoprotection solution (10% sucrose/PBS at room temperature (RT)
for 3 h; 20% sucrose/PBS at 4°C overnight; 30% sucrose/PBS at 4°C overnight), embedded in Tissue-
Tek O.C.T. Compound (Sakura Finetek Japan Co., Ltd) and stored at −80°C. The embedded
specimens were sectioned at 14 μm thickness via cryostat (CM3050S, Leica, Wetzlar, Germany),
mounted on the coated slide glasses (MAS-03, Matsunami Glass, Osaka, Japan), and air dried at RT
for a day. After 5 min washes with Tris-NaCl-Tween buffer (TNT; 0.15 M Tris-HCl, 0.01 M NaCl, 0.05%
Tween20), the sections were permeabilized with 0.1% TritonX-100/TNT for 5–10 min, blocked with
2% bovine serum albumin (BSA)/TNT for 30 min, and incubated with primary antiserum raised
elsewhere [41,43] (anti-BmGr6 and anti-BmGr9 antisera in 2% BSA/TNT at a 1:100 dilution) for 12 h
at 4°C. After three 5 min washes with TNT, the sections were incubated in a secondary antiserum (an
antimouse IgG conjugated with Alexa Fluor 488, Molecular Probes, USA), at a 1:1,000 dilution in TNT
for 12 h at 4°C. After three TNT washes for 5 min, the sections were counterstained with 4, 6-
diamidino-2-phenylindole (l mg/mL, Sigma Aldrich, Schnelldorf, Germany) for 5 min at RT. After 5 min
of TNT wash, the sections were mounted in 90% glycerol/PBS containing 1, 4-diazabicyclo[2.2.2]
octane at RT and observed under fluorescence microscopy (BX53, Olympus, Tokyo, Japan).

Generation of Gr-KO silkworm strains using CRISPR/Cas9 genome editing
We used the Alt-R® CRISPR-Cas9 system (Integrated DNA Technologies) to generate silkworm
strains carrying null mutations in *BmGr6* or *BmGr9* genes. CRISPRdirect (Naito *et al.*, 2015) identified
unique target sequences for guide RNAs using *Bombyx mori* genome assembly as a reference. Alt-
R® crRNAs (for BmGr6: 5′-GTATTCGGCGGCGCTGGGAA<u>TGG</u>-3′; for BmGr9, 5′-

CTGGTGGCGGAGATAGGAGT<u>GGG</u>-3′), tracrRNA, and S.p. Cas9 Nuclease V3 were purchased
from Integrated DNA Technologies. CRISPR-Cas9 ribonucleoprotein (RNP) complexes containing 10
µM crRNA:tracrRNA duplex and 300 ng/µL Cas9 enzyme were prepared according to a user-
submitted protocol entitled "CRISPR-Cas9 RNP delivery—*B. tryoni* microinjection—A Choo" on the
Integrated DNA Technologies website. Newly laid eggs of the WT N4 strain were aligned and glued
onto a sliding glass, and RNP complexes were injected using IM 300 Microinjector (NARISHIGE)
within 5–8 h after oviposition. Generation 0 (G_0) moths were crossed with WT moths. To identify
Generation 1 (G_1) broods carrying a heterozygous Cas9-induced mutation, genomic DNA extracted
from ~10 G_1 eggs were used for the T7 Endonuclease I (T7EI) assay with Alt-R$^®$ Genome Editing
Detection Kit (Integrated DNA Technologies). T7EI assay-positive G_1 moths were subjected to sibling
mating or crossed with WT moths. Mutations were identified by DNA sequencing of genomic PCR
(gPCR) products. Mutant strains carrying a homozygous null mutation were generated by
intercrossing G_2 or G_3 moths.
353
Feeding behavior assays
The feeding behavior assays were performed as described previously [8]. For feeding assay, three to
five fifth instar larvae (Day 0) were put on individual Petri dishes, and the larval mass before and after
1.5 h feeding was measured. An increase in larval mass was regarded as the amount of food intake
because no feces were observed after feeding. We adopted an increase in body mass (%, w/w) as
the index of food intake because a significant correlation between the amount of food intake and the
original body mass was observed via Pearson's correlation analysis (Fig. S4). Mulberry leaves were
collected around the campus of the Tokyo University of Agriculture and Technology (Koganei, Tokyo,
Japan) and stored at 4°C. To observe test bites, videos of silkworm feeding on leaves were captured
by a D5100 camera (Nikon, Tokyo, Japan) equipped with a telephoto lens AF-S DX Micro Nikkor 85
mm (Nikon, Tokyo, Japan). The larvae were used 24–48 h after molting to the fifth instar. To prevent
nonspecific biting due to starvation, all larvae were fed on artificial diets for 2 min before use.
366
Tip recording
Tip recording was performed as described previously [8]. In brief, the heads of the fifth instar larvae
(Day 0) were excised and used for further experiments within ~20 min. The whole sensilla of MP were
capped with a glass capillary (Erma Inc., Tokyo, Japan) filled with a tastant solution dissolved in 5
mM NaCl using a micromanipulator (NARISHIGE, Tokyo, Japan). Electrical signals were amplified by
a TastePROBE amplifier (Syntech, Hilversum, The Netherlands), recorded via PowerLab 2/25 (AD
Instruments, Sydney, Australia), and analyzed by the LabChart software (AD Instruments). Spikes
were counted from 0.1 to 1.1 s after the onset of stimulation using custom codes written in MATLAB
(Mathworks). Thresholds for counting spikes were set from 3 to 10 mV depending on the baseline
amplitude.

Fig. 1. Phylogeny of the gustatory receptors in some lepidopteran insects. The combination of gustatory receptors and plant secondary metabolites as their ligands were highlighted with the same colors as reported in previous studies [23,24,26,27]. Bm: *B. mori* [30]; Pxyl: *P. xylostella* [38]; Hmel: *H. melpomene* [39].

Fig. 2. Calcium imaging of BmGrs-expressing HEK293T cells. (A) The response of BmGr-

expressing cells to 100 aM chlorogenic acid (CGA) and isoquercitrin (ISQ). The representative cell

images before (pre) and after (stim) stimulation. Fluorescent intensity is color-coded according to the

scale bar. (B and C) Time course of responses (ΔF/F) toward 100 aM CGA (B) and ISQ (C) (n = 30).

The arrowheads indicate the timing of the ligand application. (D and E) Dose-dependent responses

of BmGr6, BmGr8, BmGr9, and BmGr10 toward CGA (D) and ISQ (E). (F and G) Dose-dependent

responses of BmGr6 (F) and BmGr9 (G) toward CGA, ISQ, and sugars (B–G). Data are represented

as the mean ± SEM.

392
393

Fig. 3. Immunostaining using maxillary palp (MP) BmGr6 and BmGr9 antibodies. (A and B)
Image (A) and schematic diagram (B) of the maxilla of a silkworm larva. (C and D) Expression of
BmGr6 (C) and BmGr9 (D). Arrows indicate somas of immunoreactive sensory neurons. The areas
surrounded by broken white lines indicate the autofluorescence of the cuticle. Yellow broken lines
indicate the border between MP's second and third segments. Scale bar: 25 μm.

Fig. 4. Generation and characterization of BmGr6-KO and BmGr9-KO larvae. (A) Gene structures,

sgRNA target sites, including the adjacent protospacer motif (PAM), and obtained CRISPR/Cas9-

mediated mutations of *BmGr6* and *BmGr9*. Gene structures were drawn using Exon-Intron Graphic

Maker (http://wormweb.org/exonintron). (B) The body mass of freshly molted larvae (Day 0 of the fifth

instar) (n = 140–175). (C and D) The increase in larval body mass after 1.5 h feeding of artificial diets

(C) (n = 75–117) and mulberry leaves (D) (n = 60). (E) The proportion of larvae showing test biting

toward mulberry leaves. The biting of ten larvae was observed for one minute. Experiments were

repeated as independent biological replicates (n = 7–11). (F) Palpation time until the first bite toward

mulberry leaves (n = 23–99). Data are represented as the mean ± SEM. Statistical analysis was

performed by one-way ANOVA followed by the Tukey post hoc test. "ns" indicates no significant

difference; an asterisk indicates a significant difference (**$P < 0.01$; ****$P < 0.0001$).

Fig. 5. Electrophysiological recordings of the maxillary palp (MP) in BmGr6-KO and BmGr9-KO larvae. The representative traces (A) and frequencies of action potentials from the whole MP sensilla in response to chlorogenic acid (CGA) (B) and isoquercitrin (ISQ) (C). Spikes were counted from 0.1 to 1.1 s after the onset of stimulation. Error bars are represented as SEM (n = 6–8).

416
Fig. S1. Response of BmGr6- and BmGr9-expressing cells toward β-sitosterol. (A) The
representative cell images before (pre) and after (stim) stimulation by 2 μM β-sitosterol. Fluorescent
intensity is color-coded according to the scale bar. (B) Time course of responses (ΔF/F) toward 2 μM
β-sitosterol (n = 30). The arrowhead indicates the timing of the ligand application.

A

B

423 **Fig. S2. Deduced amino acid sequences of truncated BmGr6 (A) and BmGr9 (B) in knockout**

424 **larvae.** The sequences were aligned using CLC Sequence Viewer 8 (CLC Bio, Aarhus, Denmark).

Fig. S3. Response of the maxillary palp (MP) in BmGr6-KO and BmGr9-KO larvae toward β-sitosterol. The representative traces (A) and frequencies of action potentials from the whole MP sensilla in response to 10 fM β-sitosterol (B). Spikes were counted from 0.1 to 1.1 s after the onset of stimulation. Data are represented as the mean ± SEM (n = 3). Statistical analysis was performed via one-way ANOVA followed by the Tukey post hoc test. "ns" indicates no significant difference.

432 **Fig. S4. Correlation between original body mass and feeding amount.** Correlation coefficient (r)

433 using Pearson correlation analysis (n = 80).

434 **Table S1. EC$_{50}$ values of BmGrs in response to plant secondary metabolites (chlorogenic acid**
435 **(CGA) and isoquercitrin (ISQ)) and sugars.**

Gr ligand IC50 (aM)	BmGr6 CGA 1.329	BmGr8 CGA ~ 84.55	BmGr9 CGA ~ 3.226	BmGr10 CGA 4692479	BmGr6 ISQ ~ 0.4287	BmGr8 ISQ ~ 148.4	BmGr9 ISQ ~ 1.205	BmGr10 ISQ 4663

Gr ligand IC50 (M)	BmGr6 D-glucose 5.778E-13	BmGr6 D-fructose 6.993E-12	BmGr6 sucrose 3.909E-12	BmGr6 myo-inositol 8.378E-10	BmGr9 D-glucose ~ 1.059e-015	BmGr9 sucrose 4.198E-07	BmGr9 D-fructose ~ 3.903e-018	BmGr9 myo-inositol 4.051E-12

References

1. War AR, Paulraj MG, Ahmad T, Buhroo AA, Hussain B, Ignacimuthu S, et al. Mechanisms of plant defense against insect herbivores. Plant Signal Behav. 2012;7: 1306–1320.

2. Fraenkel GS. The Raison d'Être of Secondary Plant Substances. Science. 1959;129: 1466–1470.

3. Heidel-Fischer HM, Vogel H. Molecular mechanisms of insect adaptation to plant secondary compounds. Curr Opin Insect Sci. 2015;8: 8–14.

4. Simon J-C, d'Alençon E, Guy E, Jacquin-Joly E, Jaquiéry J, Nouhaud P, et al. Genomics of adaptation to host-plants in herbivorous insects. Brief Funct Genomics. 2015;14: 413–423.

5. Chapman RF. Contact chemoreception in feeding by phytophagous insects. Annu Rev Entomol. 2003;48: 455–484.

6. Chapman RF, Bernays EA. Insect behavior at the leaf surface and learning as aspects of host plant selection. Experientia. 1989;45: 215–222.

7. Renwick JAA. Chemical ecology of oviposition in phytophagous insects. Experientia. 1989;45: 223–228.

8. Tsuneto K, Endo H, Shii F, Sasaki K, Nagata S, Sato R. Diet choice: The two-factor host acceptance system of silkworm larvae. PLoS Biol. 2020;18: e3000828.

9. Cheng D, Mulder PPJ, van der Meijden E, Klinkhamer PGL, Vrieling K. The correlation between leaf-surface and leaf-tissue secondary metabolites: a case study with pyrrolizidine alkaloids in Jacobaea hybrid plants. Metabolomics. 2017;13: 47.

10. Chun MW. Dynamics of Feeding Responses in Pieris brassicae Linn. as a Function of Chemosensory Input: A Behavioural, Ultrastructural and Electrophysiological Study. de Wilde J, editor. Wageningen University and Research. 1972. Available: https://library.wur.nl/WebQuery/wurpubs/421851

11. Devitt BD, Smith JJB. ACTION OF MOUTHPARTS DURING FEEDING IN THE DARK-SIDED CUTWORM, EUXOA MESSORIA (LEPIDOPTERA: NOCTUIDAE). Can Entomol. 1985;117: 343–349.

12. Montell C. Drosophila sensory receptors-a set of molecular Swiss Army Knives. Genetics. 2021;217: 1–34.

13. Slone J, Daniels J, Amrein H. Sugar receptors in Drosophila. Curr Biol. 2007;17: 1809–1816.

14. Dahanukar A, Lei Y-T, Kwon JY, Carlson JR. Two Gr genes underlie sugar reception in Drosophila. Neuron. 2007;56: 503–516.

15. Chyb S, Dahanukar A, Wickens A, Carlson JR. Drosophila Gr5a encodes a taste receptor tuned to trehalose. Proc Natl Acad Sci U S A. 2003;100 Suppl 2: 14526–14530.

16. Sato K, Tanaka K, Touhara K. Sugar-regulated cation channel formed by an insect gustatory receptor. Proc Natl Acad Sci U S A. 2011;108: 11680–11685.

17. Miyamoto T, Slone J, Song X, Amrein H. A fructose receptor functions as a nutrient sensor in the Drosophila brain. Cell. 2012;151: 1113–1125.

18. Zhang H-J, Anderson AR, Trowell SC, Luo A-R, Xiang Z-H, Xia Q-Y. Topological and functional
characterization of an insect gustatory receptor. PLoS One. 2011;6: e24111.

19. Kikuta S, Endo H, Tomita N, Takada T, Morita C, Asaoka K, et al. Characterization of a ligand-
gated cation channel based on an inositol receptor in the silkworm, Bombyx mori. Insect
Biochem Mol Biol. 2016;74: 12–20.

20. Dweck HKM, Carlson JR. Molecular Logic and Evolution of Bitter Taste in Drosophila. Curr Biol.
2020;30: 17-30.e3.

21. Steck K, Walker SJ, Itskov PM, Baltazar C, Moreira J-M, Ribeiro C. Internal amino acid state
modulates yeast taste neurons to support protein homeostasis in Drosophila. Elife. 2018;7.
doi:10.7554/eLife.31625

22. Tauber JM, Brown EB, Li Y, Yurgel ME, Masek P, Keene AC. A subset of sweet-sensing
neurons identified by IR56d are necessary and sufficient for fatty acid taste. PLoS Genet.
2017;13: e1007059.

23. Ozaki K, Ryuda M, Yamada A, Utoguchi A, Ishimoto H, Calas D, et al. A gustatory receptor
involved in host plant recognition for oviposition of a swallowtail butterfly. Nat Commun. 2011;2:
542.

24. Yang J, Guo H, Jiang N-J, Tang R, Li G-C, Huang L-Q, et al. Identification of a gustatory
receptor tuned to sinigrin in the cabbage butterfly Pieris rapae. PLoS Genet. 2021;17:
e1009527.

25. Renwick JAA, Lopez K. Experience‐based food consumption by larvae of Pieris rapae :
addiction to glucosinolates? Entomol Exp Appl. 1999;91: 51−58.

26. Yang K, Gong X-L, Li G-C, Huang L-Q, Ning C, Wang C-Z. A gustatory receptor tuned to the
steroid plant hormone brassinolide in Plutella xylostella (Lepidoptera: Plutellidae). Elife. 2020;9.
doi:10.7554/eLife.64114

27. Kasubuchi M, Shii F, Tsuneto K, Yamagishi T, Adegawa S, Endo H, et al. Insect taste receptors
relevant to host identification by recognition of secondary metabolite patterns of non-host
plants. Biochem Biophys Res Commun. 2018;499: 901–906.

28. Liu X-L, Zhang J, Yan Q, Miao C-L, Han W-K, Hou W, et al. The Molecular Basis of Host
Selection in a Crucifer-Specialized Moth. Curr Biol. 2020;30: 4476-4482.e5.

29. Wanner KW, Robertson HM. The gustatory receptor family in the silkworm moth Bombyx mori
is characterized by a large expansion of a single lineage of putative bitter receptors. Insect Mol
Biol. 2008;17: 621–629.

30. Guo H, Cheng T, Chen Z, Jiang L, Guo Y, Liu J, et al. Expression map of a complete set of
gustatory receptor genes in chemosensory organs of Bombyx mori. Insect Biochem Mol Biol.
2017;82: 74–82.

31. Asaoka. Ultrastructure of maxillary sensilla in the silkworm, Bombyx mori: differences among
strains? J Insect Biotechnol Sericology. 2003. Available:
https://www.jstage.jst.go.jp/article/jibs/72/2/72_2_117/_article/-char/ja/

32. Vosshall LB, Amrein H, Morozov PS, Rzhetsky A, Axel R. A spatial map of olfactory receptor
expression in the Drosophila antenna. Cell. 1999;96: 725–736.
33. Del Mármol J, Yedlin MA, Ruta V. The structural basis of odorant recognition in insect olfactory
receptors. Nature. 2021. doi:10.1038/s41586-021-03794-8
34. Jiao Y, Moon SJ, Wang X, Ren Q, Montell C. Gr64f is required in combination with other
gustatory receptors for sugar detection in Drosophila. Curr Biol. 2008;18: 1797–1801.
35. Shim J, Lee Y, Jeong YT, Kim Y, Lee MG, Montell C, et al. The full repertoire of Drosophila
gustatory receptors for detecting an aversive compound. Nat Commun. 2015;6: 8867.
36. Singh S, Kaur I, Kariyat R. The Multifunctional Roles of Polyphenols in Plant-Herbivore
Interactions. Int J Mol Sci. 2021;22. doi:10.3390/ijms22031442
37. Shii F, Mang D, Kasubuchi M, Tsuneto K, Toyama T, Endo H, et al. Ultrasensitive detection by
maxillary palp neurons allows non-host recognition without consumption of harmful
allelochemicals. J Insect Physiol. 2021;132: 104263.
38. Engsontia P, Sangket U, Chotigeat W, Satasook C. Molecular evolution of the odorant and
gustatory receptor genes in lepidopteran insects: implications for their adaptation and
speciation. J Mol Evol. 2014;79: 21–39.
39. Briscoe AD, Macias-Muñoz A, Kozak KM, Walters JR, Yuan F, Jamie GA, et al. Female
behaviour drives expression and evolution of gustatory receptors in butterflies. PLoS Genet.
2013;9: e1003620.
40. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display
and annotation. Nucleic Acids Res. 2021;49: W293–W296.
41. Mang D, Shu M, Endo H, Yoshizawa Y, Nagata S, Kikuta S, et al. Expression of a sugar clade
gustatory receptor, BmGr6, in the oral sensory organs, midgut, and central nervous system of
larvae of the silkworm Bombyx mori. Insect Biochem Mol Biol. 2016;70: 85–98.
42. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-
source platform for biological-image analysis. Nat Methods. 2012;9: 676–682.
43. Mang D, Shu M, Tanaka S, Nagata S, Takada T, Endo H, et al. Expression of the fructose
receptor BmGr9 and its involvement in the promotion of feeding, suggested by its co-
expression with neuropeptide F1 in *Bombyx mori*. Insect Biochem Mol Biol. 2016;75: 58-69.