Title: Maize integrates light and volatile cues from neighboring plants into stronger

volatile emissions

Rocío Escobar-Bravo[1], Bernardus C.J. Schimmel[1#], Yaqin Zhang[1*], Lei Wang[1], Christelle A.M.

Robert[1], Gaétan Glauser[2], Carlos L. Ballaré[3,4], Matthias Erb[1]

[1] Institute of Plant Sciences, University of Bern, 3013, Bern, Switzerland.

[2] Neuchâtel Platform of Analytical Chemistry, University of Neuchâtel, CH2000, Neuchâtel,

Switzerland

[3] Facultad de Agronomía, Instituto de Investigaciones Fisiológicas y Ecológicas Vinculadas a la

Agricultura (IFEVA) and Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET),

Universidad de Buenos Aires, Av. San Martín 4453, Buenos Aires C1417DSE, Argentina

[4] 2IIBio, Consejo Nacional de Investigaciones Científicas y Técnicas, Universidad Nacional de San

Martín, Buenos Aires B1650HMP, Argentina

#Current address: Institute of Environmental Biology, Utrecht University, Padualaan 8, 3584 CH

Utrecht, The Netherlands

*Current address: Key Laboratory of Crop Ecophysiology and Farming System in Southwest China,

Ministry of Agriculture, College of Agronomy, Sichuan Agricultural University, Chengdu 611130, P.R.

China

Rocío Escobar-Bravo, Email: rocioebe@hotmail.com, ORCID iD: 0000-0003-1280-3241.

Bernardus CJ Schimmel, Email: bart.schimmel@ips.unibe.ch, ORCID iD: 0000-0001-9345-7883.

Yaqin Zhang, Email: yaqinzhang66@163.com

Lei Wang, Email: lei.wang@unibe.ch

Christelle AM Robert, Email: christelle.robert@ips.unibe.ch, ORCID iD: 0000-0003-3415-2371

Gaétan Glauser, Email: gaetan.glauser@unine.ch, ORCID iD: 0000-0002-0983-8614.

Carlos Ballaré, Email: ballare@ifeva.edu.ar, ORCID iD: 0000-0001-9129-4531

Matthias Erb, Email: matthias.erb@ips.unibe.ch, ORCID iD: 0000-0002-4446-9834.

Authors for correspondence: Rocío Escobar-Bravo (rocioebe@hotmail.com) and Matthias Erb,

(matthias.erb@ips.unibe.ch). Phone: +41 31 684 86 68

Supporting Information: Figs. S1, S2, S3, S4, S5, S6, S7, S8, S9, S10, S11 and S12, Table S1, and

Methods S1

Abstract

- Plants perceive the presence and defense status of their neighbors through light and volatile cues, but how plants integrate both stimuli is poorly understood.

- We investigated if and how low Red to Far red light (R:FR) ratios, indicative of shading or canopy closure, affects maize (*Zea mays*) responses to herbivore-induced plant volatiles (HIPVs), including the green leaf volatile (*Z*)-3-hexenyl acetate. We modulated light signaling and perception by using FR supplementation and a *phyB1phyB2* mutant and we determined volatile release as response readout. To gain mechanistic insights, we examined expression of volatile biosynthesis genes, hormone accumulation, and photosynthesis.

- Exposure to a full blend of HIPVs or (*Z*)-3-hexenyl acetate induced maize volatile release. Short-term FR supplementation increased this response. In contrast, prolonged FR supplementation or constitutive phytochrome B inactivation in *phyB1phyB2* plants showed the opposite response. Short-term FR supplementation enhanced photosynthesis and stomatal conductance, and (*Z*)-3-hexenyl acetate-induced JA-Ile levels.

- We conclude that FR-enriched light environments can prompt maize plants to respond more strongly to HIPVs emitted by neighbors, which might be explained by changes in photosynthetic processes and phytochrome B signaling. Our findings reveal interactive responses to light and volatiles cues with potentially important consequences for plant-plant and plant-herbivore interactions.

Introduction

The capacity to perceive and respond to fluctuating environments is essential to life on earth. As primary producers in terrestrial ecosystems, plants are constantly dealing with competitors, pests, and pathogens. Consequently, they have evolved systems to detect these stressors and respond to them appropriately. While recent work has advanced our knowledge of how plants perceive individual environmental cues to detect biotic stressors, much less is known about their capacity to simultaneouly integrate multiple ones (Nguyen *et al.*, 2016; Escobar-Bravo *et al.*, 2017).

Plants can perceive qualitative and quantitative changes in light. When plants grow in dense canopies there is a strong decline in photosynthetically active radiation (PAR) and a change in light quality due to the light absorption and reflection by leaves (Ballaré & Pierik, 2017). Plants typically absorb red (R) and blue (B) light whereas the far red (FR) component of the sunlight is mostly reflected and transmitted. Consequently, in dense or closing canopies, there is a decrease in the R to FR ratio (R:FR) resulting from the reduction in R light, which is strongly absorbed by chlorophylls, and the increase in FR radiation. A reduction in R:FR ratio is a signal of foliar shading and proximity of other plants (Ballaré *et al.*, 1990). Plants can respond to this FR-enriched light environment by upward bending of the leaves (hyponasty) and stem elongation, which allows them to reach more illuminated areas and outcompete their neighbors (Franklin, 2008; Ballaré & Pierik, 2017). Variations in R:FR ratios are perceived by the photoreceptor phytochrome B (phyB) (Fankhauser, 2001). PhyB is inactivated under low R:FR, which triggers the expression of growth-related pathways controlled by hormones such as auxins and gibberellins (Fiorucci & Fankhauser, 2017).

Besides light cues, plants can perceive and respond to volatile organic compounds emitted by herbivores (Helms *et al.*, 2017) and neighboring plants (Hu, 2022). When a plant is attacked, it releases distinct blends of herbivore induced plant volatiles (HIPVs). HIPVs can be perceived by non-attacked plants and induce or prime their defenses (Engelberth *et al.*, 2004). Plant responses to HIPVs includes the rapid release of volatiles (Engelberth *et al.*, 2004) and HIPV-induced priming, which allows plants to respond more rapidly and strongly once they are under attack (Ton *et al.*, 2007). In maize, HIPVs such as indole and (*Z*)-3-hexenyl acetate [(*Z*)-3-HAC] prime the jasmonic acid (JA) pathway (Erb *et al.*, 2015; Ye *et al.*, 2019), triggering the deployment of an array of defenses that enhance herbivore resistance and the attraction of parasitoids (Ton *et al.*, 2007). In addition, (*Z*)-3-HAC can directly induce defenses, volatile biosynthesis, and release (Engelberth *et al.*, 2004; Hu *et al.*, 2019).

Recent work has demonstrated that R:FR ratios influence defense deployment in plants, including volatiles. Reduced R:FR via FR supplementation to ambient light suppresses direct defenses against herbivores and pathogens (Moreno *et al.*, 2009; Cargnel *et al.*, 2014; Courbier *et al.*, 2021). This is predominantly mediated by crosstalk between phyB and the JA pathway (Moreno *et al.*, 2009; Campos *et al.*, 2016; Fernández-Milmanda *et al.*, 2020). When phyB is inactivated, JA signaling is

attenuated as a consequence of reduced levels of bioactive JA forms (Fernández-Milmanda *et al.*,
2020), decreased stability of MYC transcription factors (Chico *et al.*, 2014), and enhanced stability of
JAZ transcriptional suppressors (Chico *et al.*, 2014; Leone *et al.*, 2014). As volatile biosynthesis can
be regulated by JA, this might explain why phyB inactivation by low R:FR ratios modulates the
emission of some volatile compounds. Upon FR supplementation, for instance, barley (*Hordeum*
vulgare) plants show reduced constitutive emission of several terpenes but an increased emission of
the monoterpene linalool oxide (Kegge *et al.*, 2015). Similarly, reduced R:FR via FR supplementation
in combination with shading (i.e., low PAR) reduces constitutive and methyl jasmonate-induced
terpenes and green leaf volatiles emissions in *Arabidopsis thaliana*, but induces the emission of
ethylene, a volatile phytohormone commonly deployed upon low R:FR (Kegge *et al.*, 2013). The fact
that emission of specific volatiles is enhanced under low R:FR (Kegge *et al.*, 2015; Cortés *et al.*, 2016)
suggests that the response to competition signals may be highly specific.

How does the perception of light cues modulate the perception of volatiles emitted by
neighbors? In contrast to the well documented effects of light cues on volatile emissions, close to
nothing is known about this important question. That plants are capable of sensing and integrating
different types of information is illustrated, for instance, by seed germination initiation, which requires
a combination of environmental cues such as temperature, moisture, and light (Finch-Savage &
Leubner-Metzger, 2006). Furthermore, plants that are under attack by herbivores of different feeding
guilds often show unique response patterns due to the hormonal integration of different signals (Soler
et al., 2013; Zhang *et al.*, 2013; Kroes *et al.*, 2015; Chrétien *et al.*, 2018). Thus, plants might also
integrate light and volatiles cues into different responses. For instance, plants might reduce their
response to volatile cues when they perceive light cues that indicate the close proximity of neighboring
plants as it may reduce the risk of an individual plant being attacked (Barbosa *et al.*, 2009).

In this study, we employed a newly developed automated high-throughput volatile screening
platform with a programmable multi-channel LED lighting to investigate if and how maize (*Zea mays*)
plants integrate light and volatile cues into defense-associated responses. We exposed maize plants
to short-term FR supplementation as a light cue for vegetation shade and to HIPVs or (*Z*)-3-HAC as
volatile cues for the presence of herbivore-infested neighbors, and we determined direct induction and
priming responses of volatile emissions in time-series analyses. In addition, we used a maize
phyB₁phyB₂ double mutant and short- or long-term FR supplementation experiments to determine the
role of PhyB in maize integration of both environmental cues. To gain mechanistic insights, we
determined gene expression, phytohormone levels, and photosynthetic responses in response to
short-term FR treatments and (*Z*)-3-HAC. Our experiments demonstrate that short-term exposure to
FR-enriched light environments, characteristic of early neighbors' detection, enhances maize volatile
emissions in response to HIPVs and (*Z*)-3-HAC, illustrating how plants use different types of signals
from neighbors to reprogram their own phenotypes.

Materials and Methods

Plants

Experiments were conducted with the maize (*Zea mays*) inbred line B73, the *phyB$_1$phyB$_2$* maize double mutant, and its wild-type W22. The *phyB$_1$phyB$_2$* double mutant was provided by Prof. Michael Scanlon (School of Integrative Plant Science, Cornell University). This mutant is homozygous for the Mutator (Mu) transposon insertion alleles *phyb1-563* and *phyb2-12058* resulting in no detectable PHYB protein (Sheehan *et al.*, 2007; Dubois *et al.*, 2010). B73 responds to (*Z*)-3-HAC (Hu *et al.*, 2018) and FR (Dubois *et al.*, 2010) and releases moderate volatile emissions upon herbivory (Block *et al.*, 2018). For experiments with 11-day-old plants, seeds were sown in transparent cylindric plastic pots (11x4 cm) filled with commercial soil (Selmaterra, BiglerSamen, Switzerland) and wrapped with aluminum foil to prevent root exposure to light. For experiments with 30-day-old plants, seeds were sown in black plastic pots (9x9x10 cm) filled with the same commercial soil. Plants were grown in a ventilated greenhouse supplemented with artificial lighting under 50-70% relative humidity, 14/10 h light/dark photoperiod, and 14-22/10-14°C day/night temperatures.

Insects

Eggs of *Spodoptera littoralis* were provided by Oliver Kindler (Syngenta, Stein, CHE) and the larvae were reared on artificial diet as described in Maag *et al.*, 2014. *S. littoralis* oral secretions were collected from third- to fourth-instar larvae as described in Hu *et al.*, 2019 and stored at − 80°C until use.

Light treatments

Light treatments were conducted in a custom-made high-throughput phenotyping platform designed for volatile analyses. The platform consisted of 96 transparent cylindrical glass chambers (120 mm Ø, 450 mm height; abon life sciences, Switzerland) closed at the top with transparent glass lids and distributed over two tables. Each table was supplied with four light emitting diodes (LEDs) lamps placed above the glass chambers. Each lamp contained 120 LEDs with nine wavebands consisting of 380, 400, 420, 450, 520, 630, 660, 735 nm and 5700K white (RX30, Heliospectra). Individual glass chambers were supplied with an airflow of 0.8 L/min and an air outlet. Plants were placed inside these glass chambers and exposed to low (~0.5) or high (~2.65) R:FR ratios (R, λ = 600-700 nm; FR, λ = 700-800 nm) by providing supplemental FR to the same white light background (120 ± 15 μmol m^{-2} s^{-1}) (Supplemental Fig. S1) with a photoperiod of 16/8 h light/dark. This low R:FR corresponds to the ratios observed in high density maize canopies (Maddonni *et al.*, 2002), where low PAR levels as the ones used in our experimental set-up are commonly observed (Xue *et al.*, 2016). Light treatments were separated by white opaque curtains and started between 9:00 and 10:00 a.m. two days before the volatile exposure treatments.

(Z)-3-HAC treatment

Individual plants were exposed to one (Z)-3-HAC dispenser for 6 h within the transparent cylindrical glass chambers of the custom-made high-throughput phenotyping platform described above. (Z)-3-HAC dispensers were prepared as described in Hu *et al.*, 2019, each releasing approximately 70 ng h^{-1} , which has been previously described in herbivore-attacked maize plants (Mérey *et al.*, 2011). (Z)-3-HAC treatments started between 9:00-10:00 a.m.

Simulated herbivory

Two leaves (leaf 2 and 3 from the bottom) per plant were wounded over an area of ca. 0.5 cm^2 parallel to the central vein using a haemostat followed by the application of 10 µL of *S. littoralis* larval oral secretions (diluted 1:1 in autoclaved distilled water) in each wounding site. This treatment induces plant defense responses comparable to real herbivory (Erb *et al.*, 2009).

Linalool dispensers

Linalool dispensers were prepared as described in Hu *et al.*, 2019. The dispensers contained 100 µL of pure linalool (SIGMA) and were pierced through the rubber septum with a 1 µL micropipette (Drummond, Millan SA, Switzerland).

PTR-TOF-MS volatile analysis

Plant volatile emission was determined using the automated high-throughput phenotyping platform described above for light treatments. Volatiles emitted by a plant inside each transparent glass chamber were purged by a clean air flow (0.8 L/min) that carried the emitted volatiles directly to a Vocus proton transfer reaction time-of-flight mass spectrometer (PTR-TOF-MS) system (Tofwerk, Switzerland) for analysis. The Vocus PTR-TOF-MS was operated using H_3O^+ as the reagent ion and was integrated into an autosampler (Gonin *et al.*, 2022) that cycles between individual chambers/plants. Each plant was measured for 50 or 60 seconds. Between individual plants, a fast zero air measurement was performed for 5 seconds. Complete mass spectra (0-500 Th) were saved and plotted at 5 Hz (5 unique mass spectra per second). PTR-TOF-MS data processing consisted of mass calibration and calculation of peak intensities using the Tofware software package v3.2.0, followed by normalization of the signals to aboveground (stem and leaves) plant fresh biomass.

GC-MS volatile emission analysis

Dynamic volatile collection was conducted using adsorbent Super-Q traps followed by gas chromatography-mass spectrometry (GC-MS) analysis of volatile extracts**.** For this, air was pulled out from the PTR-TOF-MS transparent glass chambers containing the test plants through the air outlet ports (0.5 L/min) and across the Super-Q traps (Porapak-Q^{TM} 50 mg, Volatile collection Trap, USA). Volatiles were collected for periods of 2 h for 12 h. After each collection period, Super-Q traps were

exchanged for new ones and the volatiles collected on the trapping filters were extracted with 200 µl
of Dichloromethane (Roth) containing 200 ng of both N-octane and nonyl-acetate as internal
standards. Extracts were analyzed by GC-MS (Agilent 7820A GC interfaced with and Agilent 5977E
MSD, Palo Alto, CA, USA). Aliquots of 1 µL were injected into the column (HP5-MS, 30 m, 250 µm ID,
2.5-µm film, Agilent Technologies, Palo Alto, CA, USA) using a pulsed split-less mode and He as
carrier gas at a flow rate of 1 mL min^{-1}. A temperature gradient of 5°C min $^{-1}$ from 40°C to 200°C was
used. Compound identification was based on similarity to library matches (NIST search 2.2 Mass
Spectral Library. Gaithersburg, MD, USA), retention times, and spectral comparison with pure
compounds. Relative abundance was calculated by dividing the peak area of individual compounds
by the aboveground plant fresh biomass.

Phytohormone analysis

The concentrations of 12-oxo-phytodienoic acid (OPDA), JA, jasmonic acid-isoleucine (JA-Ile),
salicylic acid (SA), abscisic acid (ABA) and auxin (indole acetic acid, IAA) were determined in
aboveground tissues (stem and leaves) following the procedure described in Glauser *et al.*, 2014 with
minor modifications (Supplemental Methods S1).

Gene expression analysis

Total RNA was isolated using TRI Reagent (Sigma-Aldrich) following the manufacturer's instructions.
DNase treatment, reverse transcription, and first strand cDNA synthesis were performed using gDNA
Eraser (Perfect Real Time) and PrimeScript Reverse Transcriptase following manufacturer's
instructions (Takara Bio Inc. Kusatsu, Japan). For gene expression analysis, 2 µL of cDNA solution
(equivalent to 10 ng of total RNA) served as template in a 10 µL qRT-PCR reaction using ORA™ SEE
qPCR Mix (Axonlab) on an Applied Biosystems® QuantStudio® 5 Real-Time PCR System.
Normalized expression (NE) values were calculated as in Alba *et al.*, (2015). Gene identifiers, primer
sequences, and references are listed in Table S1.

Stomatal conductance, density, and net photosynthesis

Stomatal conductance was measured using an AP4 Porometer (Delta-T Devices Ltd) with cup
aperture dimension of 2.5 x 17.5 mm. Stomatal density was determined in epidermal imprints of the
abaxial surface using clear nail varnish. Epidermal imprints were imaged using a Leica DM2500 optical
microscope coupled to a Leika HD Digital microscope camera (Leica MC170 HD) (Leica
Microsystems, Wetzlar, Germany). The number of stomata was determined in 10 pictures per plant
and leaf using ImageJ software. Net photosynthesis was determined using a LI-6400 Portable
Photosynthesis system (LI-COR, Biosciences).

 Statistical analysis

Data analysis was conducted in R version 4.0.4 (R core Team, 2016). Effects of (1) sampling time, (2)

light, (3) genotype, (4) volatile exposure, and (5) their interactions on volatile emissions in the different

experiments were tested in linear mixed-effects models (LME) using nlme library. We included

individual plants as a random factor and a correlation structure when autocorrelation among residuals

was found to be significant ($p < 0.05$) in the models. Fitted models were subjected to type III analyses

of variance (ANOVAs) to produce a summary of the F- and p statistics (car package Fox & Weisberg,

2018). When needed, data from volatiles signatures were log10 or squared root transformed prior

analysis to correct for heteroscedasticity. In Fig. 4 differences among light treatments were tested for

all volatiles at each sampling time by estimated marginal means. Hormone concentrations and gene

expression were analyzed separately for each time point by two-way ANOVAs followed by Tukey HSD

post hoc test when light, (*Z*)-3-HAC, or their interaction were significant. Effects of light, (*Z*)-3-HAC,

herbivory, and their interactions on stomatal conductance, density, and net photosynthesis were tested

by two-way ANOVAs followed by Tukey HSD post hoc test.

Results

FR-enrichment boosts maize responses to herbivore-induced plant volatiles

HIPVs can induce the emission of volatiles in non-attacked plants and enhance their defense

responses when actual herbivory occurs (Farag & Paré, 2002; Engelberth *et al.*, 2004). To determine

whether different R:FR ratios alter growth and responses to HIPVs in maize, we exposed 11-day old

B73 seedlings to high or low R:FR for two days by modulating FR levels. Thereafter, we measured

growth-related parameters and compared their volatile emissions when exposed to volatiles emitted

by non-infested or *Spodoptera littoralis*-infested low R:FR treated maize plants. Low R:FR-treated

emitter plants infested with ten second-instar *S. littoralis* larvae or left uninfested were connected to

the glass chambers of receiver plants using Teflon tubing (Fig. 1a). The air flow moved from the

emitter's chamber towards the receiver's chamber. Volatiles simultaneously emitted by emitter and

receiver plants were collected from the glass chamber containing the receiver plant through an air

outlet connected to the PTR-TOF-MS system. Because all emitter plants were exposed to low R.FR

and volatile fluxes were unidirectional, differences in volatile emissions could only be explained by the

light treatment of the receivers (low or high R:FR).

Low R:FR increased the elongation of stem internodes and auxin levels (Supplemental Fig.

S2ab), confirming the effectiveness of our light treatment in triggering shade avoidance (Dubois &

Brutnell, 2009).

Emission of the green leaf volatiles (GLVs) hexenal, hexen-1-ol and hexenyl acetate, as well

as indole, monoterpenes, sesquiterpenes, and the homoterpenes DMNT and TMTT by *S. littoralis*-

infested emitter + intact receiver plants significantly increased over time in comparison to non-infested

emitter + intact receiver plants (Fig. 1b). Indole and sesquiterpenes emissions were significantly higher

when receiver plants were exposed to low R:FR and volatiles from *S. littoralis*-infested emitter plants

than to the same volatiles and high R:FR (Fig. 1b). A similar pattern was observed for monoterpenes

and DMNT. As emitter plants were all grown under low R:FR, showed similar levels of herbivore

damage (Supplemental Fig. S3), and did not receive any volatile cues from the receiver plants, these

differences must be the result of changes in the receivers' volatile emissions. Thus, FR-enriched light

environments enhance maize responses to HIPVs emitted by herbivore-infested neighbors.

FR-enrichment boosts maize responses to (Z)-3-HAC

GLVs such as (*Z*)-3-hexenal, (*Z*)-3-hexen-1-ol, and (*Z*)-3-HAC are HIPVs emitted by maize that can

induce the emission of volatiles in neighboring intact maize plants (Farag & Paré, 2002; Engelberth *et*

al., 2004). We next tested whether low R:FR modulates maize responses to the GLV (*Z*)-3-HAC. We

exposed 11-day old B73 maize seedlings to low or high R:FR for two days by modulating FR levels

and determined their volatile emissions by using the PTR-TOF-MS system during (*Z*)-3-HAC exposure

and subsequent simulated herbivory (Fig. 2a). These analyses focused on indole and terpenoid

emissions because these compounds responded to low R:FR and HIPVs in the previous experiment.

Exposure to (*Z*)-3-HAC induced volatile release in both high and low R:FR-treated maize

plants (Fig. 2b). Low R:FR, however, significantly enhanced (*Z*)-3-HAC-induced emissions of

monoterpenes and DMNT, while the opposite was observed for indole.

Upon simulated herbivory, pre-exposure to (*Z*)-3-HAC increased the emission of all the

volatiles signatures in both light treatments (Fig. 2b). Emission of DMNT was significantly higher in low

R:FR-treated plants than in high R:FR-treated plants pre-exposed to (*Z*)-3-HAC. Irrespective of (*Z*)-3-

HAC pre-treatment, herbivory-induced plants exposed to low R:FR emitted more monoterpenes,

sesquiterpenes, and DMNT than high R:FR-treated plants (Fig. 2b). An additional experiment

confirmed that low R:FR enhances herbivore-induced emissions of monoterpenes and DMNT, but it

reduced constitutive indole emissions and significantly increased them upon herbivory (Supplemental

Fig. S4).

The PTR-TOF-MS system only detects molecular mass features, while the structural identity

of isomers, e.g., different mono- or sesqui-terpenes, cannot be resolved. To determine the identity of

the induced volatile compounds and confirm the observed patterns with an established, orthogonal

method, we collected plant volatiles for periods of 2 h for 12 h using Super-Q traps followed by GC-

MS analysis of volatile extracts. (*Z*)-3-HAC exposure significantly increased the emissions of indole,

the monoterpene alcohol linalool, the sesquiterpenes (*E*)-β-farnesene and (*E*)-α-bergamotene, and

the homoterpene TMTT (Supplemental Fig. S5).

Figure 1. FR-enrichment boosts maize responses to herbivore-induced plant volatiles. (a) Schematic overview of the experimental set-up. Eleven-day old `B73´ maize (*Zea mays*) plants were exposed to low or high R:FR light conditions by modulating FR levels. After two days, receiver plants growing under low or high R:FR were exposed to constitutive volatiles (cVOC) or *S. littoralis* herbivory-induced plant volatiles (iVOC) from emitter plants growing under low R:FR. Emitter and Receiver plants were connected via Teflon tubing and their combined volatile emissions were measured by PTR-TOF-MS. Volatile measurements started at 12:00 a.m. **(b)** Time-series analysis of volatile emissions (mean ± SEM, *n* = 12) collected from paired plants. The effects of sampling time (T), light treatments of receivers (L), volatiles from emitters (VOC), and their interactions on volatile emissions were tested using linear mixed-effects models. Statistically significant effects are shown in each graph (*$p < 0.05$, ** $p < 0.001$, *** $p < 0.001$). DMNT and TMTT stand for the homoterpenes (*E*)-4,8-dimethyl-1,3,7-nonatriene and (*E*, *E*)-4,8,12-trimethyltrideca-1,3,7,11-tetraene, respectively. AU refers to arbitrary units.

Low R:FR alone had a positive effect on the emission of the (*E*)-β-farnesene, and it also increased
(*Z*)-3-HAC-induced emission of indole, linalool and (*E*)-α-bergamotene. We did not detect a significant
induction of the monoterpenes α-pinene and β-ocimene. Further analyses using dispensers with pure
linalool, which is a monoterpene alcohol, showed that the most abundant fragment ion of linalool has
an m/z of 137 ($C_{10}H_{17}^{+}$) (Supplemental Fig. S6) like other monoterpenes. Thus, the induced
monoterpene signal detected in the PTR-MS experiments upon (*Z*)-3-HAC and FR treatment is likely
due to enhanced linalool emissions. FR effects on indole emissions agreed with the enhanced
induction observed in Fig. 1b but differed from PTR-TOF-MS data displayed in Fig. 2b, indicating a
stronger context-dependent variation in the emission of this compound. Upon simulated herbivory,
linalool, (*E*)-β-farnesene, and (*E*)-α-bergamotene emissions were higher under low R:FR than under
high R:FR irrespective of (*Z*)-3-HAC pre-treatment. TMTT was below the limit of detection of the GC-
MS. During (*Z*)-3-HAC exposure and after simulated herbivory, DMNT emission was higher under low
R:FR than under high R:FR, albeit these differences were not statistically significant. This might be
explained by the lower number of sampling times compared to the PTR-TOF-MS experiments,
resulting in reduced statistical power.

In field-grown maize, heterogeneity in R/FR environments within maize canopies is detected
from early ontogenetic stages – i.e., 30 days from sowing - to flowering. This heterogeny is
characterized by an enriched FR environment that acts as a signal for leaf reorientation to escape
vegetation shade (Maddonni *et al.*, 2002). To put our results into a more realistic context, we tested if
the observed response to low R:FR and (*Z*)-3-HAC also occurs in 30-day-old B73 maize plants. Low
R:FR significantly enhanced monoterpenes, sesquiterpenes, DMNT, and TMTT emissions during (*Z*)-
3-HAC exposure, and it did not affect indole (Supplemental Fig. S7). Upon simulated herbivory, pre-
exposure to (*Z*)-3-HAC enhanced indole and sesquiterpenes emissions under both light treatments.
Irrespective of (*Z*)-3-HAC pre-treatment, low R:FR increased monoterpenes, sesquiterpenes, TMTT,
DMNT, and indole emissions upon herbivory. We also tested whether the slight variations in horizontal
PAR levels detected in this experimental set up could have influenced maize responses to (*Z*)-3-HAC.
There was no significant correlation between PAR and volatile emissions at the sampling time with the
strongest (*Z*)-3-HAC effect (Supplemental Fig. S8), confirming FR specific effects on maize responses
to (*Z*)-3-HAC.

Together, these results demonstrate that FR-enrichment increases maize responsiveness to
(*Z*)-3-HAC, resulting in higher direct and primed emission of terpenes in both 11- and 30-day-old maize
plants.

Figure 2. FR-enrichment boosts maize responses to (*Z*)-3-hexenyl acetate. (a) Schematic overview of the experimental set-up. Eleven-day old `B73´ maize (*Zea mays*) plants were exposed to low or high R:FR conditions for 2 d by modulating FR levels, and subsequently exposed to (*Z*)-3-hexenyl acetate (HAC) or left untreated for 6 h. After HAC treatment, plants were induced with simulated herbivory (wounding and application of *Spodoptera littoralis* oral secretions; W+OS). Volatile emissions were determined during HAC exposure and after simulated

herbivory by PTR-TOF-MS. **(b)** Time-series analysis of volatile emissions (mean ± SEM, *n* = 9) determined in
low and high R:FR-treated plants during exposure to HAC (+ HAC) or clean air (- HAC) and after induction with
simulated herbivory (priming phase). HAC exposure and volatile measurements started between 9:00 and 10:00
am. The effects of the sampling time (T), HAC, light treatment (L) and their interactions on volatile emissions
were tested separately for the HAC exposure and priming phase using linear mixed-effects models. Statistically
significant effects are shown in each graph (*$p < 0.05$, ** $p < 0.001$, *** $p < 0.001$). DMNT and TMTT stand for
the homoterpenes (*E*)-4,8-dimethyl-1,3,7-nonatriene and (*E*,*E*)-4,8,12-trimethyltrideca-1,3,7,11-tetraene,
respectively. AU refers to arbitrary units.

Maize volatile emissions in response to (Z)-3-HAC are reduced in the phyB1phyB2 mutant

Low R:FR is sensed by the phytochrome family of photoreceptors, of which phyB is the central
regulator in Arabidopsis (Ballaré & Pierik, 2017). In maize, gene duplication has resulted in two PhyB1
and PhyB2 genes, both involved in FR signaling (Sheehan *et al.*, 2007; Dubois *et al.*, 2010). We next
determined whether a non-functional phyB enhances maize responses to (*Z*)-3-HAC like low R:FR.

First, we exposed 11-day-old *phyB1phyB2* maize double mutant plants and its wild-type
(inbred line W22) to high R:FR for two days and determined volatile emissions during (*Z*)-3-HAC
exposure and after simulated herbivory using the PTR-TOF-MS system. Compared to the wild-type,
constitutive emission of indole, monoterpenes, sesquiterpenes, and DMNT was significantly lower in
phyB1phyB2 plants (Fig. 3). (*Z*)-3-HAC increased volatile emissions in wild-type and *phyB1phyB2*
plants (Fig. 3), but the induction of volatiles was significantly lower in *phyB1phyB2* plants compared
to the wild-type. Similar patterns were observed following simulated herbivory (Fig. 3).

To better understand the unexpected response of *phyB1phyB2* plants, we conducted
additional experiments. First, we analyzed photomorphogenic responses of *phyB1phyB2* and wild-
type W22 plants to low R:FR. As expected, *phyB1phyB2* plants grew longer stem internodes
irrespective of the light regime (Supplemental Fig. S9). The wild-type W22 did not show a strong
response to low R:FR, which might be explained by maize genotype-dependent responses to
supplemental FR (Dubois *et al.*, 2010).

Next, we compared the responsiveness of the wild-type W22 and *phyB1phyB2* plants to (*Z*)-
3-HAC when grown under high or low R:FR. Low R:FR significantly enhanced (*Z*)-3-HAC-induced
emission of monoterpenes, sesquiterpenes, and DMNT in wild-type W22 plants (Supplemental Fig.
S10) as observed before in B73 (Fig. 2b and Supplemental Fig. S5), whereas the effect on indole
differed along the time course. Light did not significantly affect volatile emissions upon simulated
herbivory. In *phyB1phyB2* plants, low R:FR slightly increased indole, monoterpenes, and
sesquiterpenes emissions at the start of the (*Z*)-3-HAC treatment (Supplemental Fig. S11), and it did
not significantly affect volatile emissions after simulated herbivory. These experiments indicate that
phyB1phyB2 plants retain partial, but not full, enhanced responsiveness to (*Z*)-3-HAC when
simultaneously exposed to low R:FR.

Figure 3. Maize volatile emissions in response to (Z)-3-HAC are reduced in the _phyB1phyB2_ mutant. Time-series analysis of volatile emissions (mean ± SEM, _n_ = 6) determined by PTR-TOF-MS in high R:FR-treated _PhyB₁B₂_ double mutant and wild-type (inbred line `W22´) maize (_Zea mays_) plants during exposure to (Z)-3-hexenyl acetate (+ HAC) or clean air (- HAC) and after induction with simulated herbivory (wounding and application of _Spodoptera littoralis_ oral secretions; W+OS) (priming phase). HAC exposure and volatile measurements started between 9:00 and 10:00 am. The effects of the sampling time (T), HAC, genotype (G), and their interactions on volatile emissions were tested separately for the HAC exposure and priming phase using linear mixed effects models. Statistically significant effects are shown in each graph (*_p_ < 0.05, ** _p_ < 0.001, *** _p_ < 0.001). DMNT and TMTT stand for the homoterpenes (E)-4,8-dimethyl-1,3,7-nonatriene and (E, E)-4,8,12-trimethyltrideca-1,3,7,11-tetraene, respectively. AU refers to arbitrary units.

Maize volatile responsiveness to (Z)-3-HAC is stronger under short-term exposure to FR than under long-term exposure

Why do *phyB1phyB2* mutant plants that behave as if continuously exposed to low R:FR show suppression of (*Z*)-3-HAC-induced volatile emissions? One possible explanation is that phyB is inactivated in *phyB1phyB2* mutant plants over the full course of their development, while in FR experiments plants have a deactivated phytochrome for only two days preceding volatile exposure. To test whether duration of phyB inactivation matters for volatile responsiveness, seven-day-old W22 plants were exposed to low or high R:FR for 6 days, or 4 days of high R:FR followed by 2 days of low R:FR (Fig. 4a). Then, we monitored volatile emissions during (*Z*)-3-HAC exposure and subsequent simulated herbivory using the PTR-TOF-MS system (Fig. 4b).

Wild-type W22 plants exposed to low R:FR for 2 days showed stronger volatile responses to (*Z*)-3-HAC than plants exposed to high R:FR (Fig. 4b) as in previous experiments (Supplemental Fig. S10). Short and long-term exposure to low R:FR induced higher indole, sesquiterpenes, and DMNT emissions, but the inverse pattern was observed for TMTT. Induced sesquiterpenes and DMNT emissions at early times of (*Z*)-3-HAC exposure were lower in plants exposed to low R:FR for 6 days than to shorter low R:FR exposure (2 days) and high R:FR only (Fig. 4b). Upon herbivory, short-term exposure (2 days) to low R:FR also enhanced monoterpenes and sesquiterpenes emissions, but this response was attenuated under long-term low R:FR exposure (6 days). Thus, long-term inactivation of phyB by low R:FR may partially explain the discrepancy between the *phyB1phyB2* double mutant phenotype and short-term FR supplementation experiments. These results also suggest that short- and long-term inactivation of phyB might result in different physiological responses.

Figure 4. Maize volatile responsiveness to (*Z*)-3-hexenyl acetate is stronger under short- than under long-term FR-enrichment. (a) Schematic overview of the experimental set-up. Seven-day-old W22 maize (*Zea mays*) seedlings were exposed to: (1) high R:FR for 6 days, (2) low R:FR for 6 days, or (3) high R:FR for 4 days and transferred to low R:FR conditions for 2 days. Thereafter, plants were exposed to (*Z*)-3-hexenyl acetate (HAC) for 6 h and subsequently induced with simulated herbivory (wounding and application of *Spodoptera littoralis* oral secretions; W+OS). **(b)** Time-series analysis of volatile emissions (mean ± SEM, *n* = 6) determined by PTR-TOF-MS during exposure to HAC (+ HAC) or clean air (-HAC) and after induction with simulated herbivory (priming phase). HAC exposure and volatile measurements started between 9:00 and 10:00 am. The effects of the sampling time (T), HAC, light treatment (L) and their interactions on volatile emissions were tested separately for the HAC exposure and priming phase using linear mixed effects models. Statistically significant effects are shown in each graph (*p < 0.05, ** *p* < 0.001, *** *p* < 0.001). Shaded grey squares containing different letters indicate

430 significant differences among treatments tested by estimated marginal means at $p < 0.05$. DMNT and TMTT
431 stand for the homoterpenes (E)-4,8-dimethyl-1,3,7-nonatriene and (E, E)-4,8,12-trimethyltrideca-1,3,7,11-
432 tetraene, respectively. AU refers to arbitrary units.

FR-enrichment enhances JA-Ile accumulation in response to (Z)-3-HAC

Volatile emissions in maize can be modulated by jasmonates and ABA (Bruinsma *et al.*, 2009; Seidl-
Adams *et al.*, 2015). To investigate the potential mechanisms by which short-term exposure to low
R:FR enhances volatile emissions, levels of these stress-related hormones were analyzed in 11-day
old B73 seedlings exposed to low or high R:FR for two days and subsequently to (Z)-3-HAC and
simulated herbivory (Fig. 5).

Exposure to (Z)-3-HAC increased OPDA, JA, JA-Ile, and ABA levels as previously described
(Hu *et al.*, 2019) in both light regimes (Fig. 5a). Light did not affect JA, JA-Ile, and SA levels, but low
R:FR slightly reduced OPDA concentrations and enhanced ABA levels. In addition, JA-Ile levels in low
R:FR-treated plants at 45 min after the start of (Z)-3-HAC exposure were significantly higher when
compared to high R:FR-treated plants.

Upon simulated herbivory, (Z)-3-HAC pre-treated plants displayed higher OPDA and JA-Ile
levels when compared to plants pre-exposed to clean air irrespective of light treatments (Fig. 5b). Low
R:FR enhanced JA, JA-Ile, and ABA levels at 180 min after simulated herbivory irrespective of (Z)-3-
HAC pre-treatment (Fig. 5b).

FR-enrichment modulates the expression of volatile biosynthetic genes in response to (Z)-3-HAC and herbivory

Next, we tested whether low R:FR-induced emission of volatiles in B73 seedlings correlated with
changes in the expression of terpene-related biosynthetic genes during (Z)-3-HAC exposure and
subsequent simulated herbivory in two separate time-series experiments (Fig. 6).

Exposure to (Z)-3-HAC significantly induced the expression of *FPSS3,* which provides
precursors for sesquiterpene biosynthesis (Richter *et al.*, 2015), *TPS10*, involved in the biosynthesis
(E)-α-bergamotene and (E)-β-farnesene (Schnee *et al.*, 2006), and *TPS2*, involved in the production
of linalool, DMNT and TMTT (Richter *et al.*, 2016) (Fig. 6a). *TPS1*, involved in the biosynthesis of (E)-
β-farnesene, linalool derivatives, and *ent*-kaurene, the latter serving as an intermediate in the
production of kauralexins (Schnee *et al.*, 2002; Fu *et al.*, 2016; Xu *et al.*, 2019), was also induced by
(Z)-3-HAC. Expression of *CYP92C5*, involved in the production of DMNT (Richter *et al.*, 2016), was
slightly reduced by (Z)-3-HAC at 180 min after the start of the treatment (Fig. 6a). (Z)-3-HAC effects
were not influenced by light. Low R:FR alone, however, reduced the expression of *CYP92C5* and
FPPS3 genes at 45 min and 90 min, respectively, and slightly enhanced *TPS1* expression at 45 min
after the start of the volatile treatment (Fig. 6a).

Figure 5. FR-enrichment enhances JA-Ile accumulation in response to (*Z*)-3-hexenyl acetate. Concentrations (mean ± SEM, *n* = 4-6) of 12-oxo-phytodienoic acid (OPDA), jasmonic acid (JA), jasmonic acid-isoleucine (JA-Ile), salicylic acid (SA), and abscisic acid (ABA) in high and low R:FR treated B73 maize (*Zea mays*) plants that were in **(a)** exposed to (*Z*)-3-hexenyl acetate **(**HAC) and in **(b)** induced with simulated herbivory (wounding and application of *Spodoptera littoralis* oral secretions) after exposure to HAC for 6 h. Effects of light treatment (L), HAC exposure and their interaction were tested by two-way ANOVA at each time point. Different letters denote significant differences among groups tested by Tukey-HSD post hoc test. Statistically significant effects are shown in each graph (*$p < 0.05$, ** $p < 0.001$, *** $p < 0.001$).

Upon simulated herbivory, (*Z*)-3-HAC pre-treatment enhanced the expression of *TPS2* and *TPS10*
and suppressed *CYP92C5* expression in low and high R:FR-treated plants, with no effect on *FPSS3*
and *TPS1* (Fig. 6b). Low R:FR enhanced *TPS2* expression in (*Z*)-3-HAC pre-treated plants and led to
a stronger (*Z*)-3-HAC-mediated CYP92C5 suppression upon simulated herbivory. Irrespective of (*Z*)-
3-HAC pre-treatment, low R:FR slightly enhanced *TPS2*, *TPS10*, and *CYP92C5* expressions after
simulated herbivory (Fig. 6b). These results show that FR-enriched environments and (*Z*)-3-HAC
interact to modulate the expression of some, but not all, terpene biosynthesis genes, and that these
slight changes are unlikely to fully explain the differences in volatile emissions.

Figure 6. FR-enrichment modulates the expression of volatile biosynthetic genes in response to (*Z*)-3-
hexenyl acetate and herbivory. Normalized expression (mean ± SEM, *n* = 6) of terpene metabolism-related

genes determined in aboveground (leaves and stem) tissues of low and high R: FR-irradiated B73 maize (*Zea*
mays) plants that were in **(a)** exposed to (*Z*)-3-hexenyl acetate (HAC) and in **(b)** induced with simulated herbivory
(wounding and application of *Spodoptera littoralis* oral secretions; W+OS) after exposure to HAC for 6 h. Effects
of light treatment (L), HAC and their interaction were tested by two-way ANOVA at each time point. Different
letters denote significant differences among groups tested by Tukey-HSD post hoc test. Statistically significant
effects are shown in each graph (*$p < 0.05$, ** $p < 0.001$, *** $p < 0.001$) Abbreviations for precursors and genes:
FPPS3, farnesyl pyrophosphate synthase 3; *TPS1, Terpene synthase 1; TPS10*, *Terpene synthase 10*; *TPS2*,
terpene synthase 2; *CYP92C5*, *P450 Monooxygenase CYP92C5.*

FR-enrichment increases stomatal conductance and photosynthesis

Emission of sesquiterpenes can be regulated by light-dependent stomata movements in maize (Seidl-
Adams *et al.*, 2015). We tested whether differences in stomata conductance accounted for the
increase in volatile emissions under low R:FR. For this, 11-day-old B73 seedlings were exposed to
low or high R:FR for two days by modulating FR levels, and leaf stomatal conductance was determined
at 3 h after (*Z*)-3-HAC exposure and at 4 h after subsequent simulated herbivory.

Exposure to (*Z*)-3-HAC significantly reduced stomatal conductance in maize (Fig. 7ab). Low
R:FR significantly increased stomatal conductance (Fig. 7ab), which was not explained by differences
in stomatal densities (Supplemental Fig. S12). We did not detect significant interactions between light
and (*Z*)-3-HAC on stomatal conductance. Simulated herbivory reduced stomatal conductance close
to the wounding sites (Fig. 7cd), a phenomenon driven by OPDA signaling in local leaves of
dicotyledonous plants (Savchenko et al., 2014; Meza-Canales et al., 2017). In leaf 2, stomatal closure
was slightly stronger in wounded plants exposed to low R:FR and (*Z*)-3-HAC pre-treatment (Fig. 7d).

Stomatal opening as well as volatile biosynthesis are tightly coupled to photosynthetic rates
(Arimura *et al.*, 2008; Way & Pearcy, 2012). We therefore evaluated if light and volatile cues interact
to regulate photosynthesis. While (*Z*)-3-HAC had no significant effect on photosynthesis,
photosynthetic rates were significantly increased under low R:FR in leaf 3 (Fig. 8ab).

Figure 7. FR-enrichment increases stomatal conductance in maize. Stomatal conductance (mean ± SEM, *n* = 20) determined in the abaxial side of **(a)** leaf #3 and **(b)** leaf #2 from the bottom of low and high R:FR-treated B73 maize (*Zea* mays) plants after 3 h of exposure to (*Z*)-3-hexenyl acetate (HAC). Effects of light treatment (L), HAC exposure and their interaction were tested by two-way ANOVAs. Stomatal conductance (mean ± SEM, *n* = 10) determined in the abaxial side of **(c)** leaf #3 and **(d)** leaf #2 of low and high R:FR-treated B73 plants that were induced with simulated herbivory (wounding and application of *Spodoptera littoralis* oral secretions; W+OS) after exposure to HAC for 6 h. Measurements were performed close to the wounding sites at 4 h after simulated herbivory. Effects of light treatment (L), HAC exposure, herbivory (H), and their interaction were tested by three-way ANOVAs. In all graphs, different letters denote significant differences among groups tested by Tukey-HSD post hoc test. Statistically significant effects are shown in each graph (*$p < 0.05$, ** $p < 0.001$, *** $p < 0.001$).

Figure 8. FR-enrichment increases photosynthesis in maize Photosynthesis rate (mean ± SEM, *n* = 10)
determined in the abaxial side of **(a)** leaf 3 and **(b)** leaf 2 from the bottom of high and low R:FR-treated B73 maize
plants at 3 h after the start of (*Z*)-3-hexenyl acetate (HAC) exposure. Effects of light treatment (L) and HAC on
stomatal density were tested by two-way ANOVAs. Statistically significant effects are shown in the graphs. ****p*
< 0.001. Different letters denote significant differences among groups tested by Tukey-HSD post hoc test at *p* <
0.05.

Discussion

In closing or dense canopies, plants can detect the proximity and physiological state of neighboring

plants through light and volatile cues. Our results demonstrate that maize plants integrate both stimuli

into distinct defense-related volatile signaling patterns. Below, we discuss the potential mechanisms

and ecological consequences of this phenomenon.

HIPVs can induce defense-related responses in undamaged plants including the biosynthesis

and emission of volatiles (Arimura *et al.*, 2002; Choh *et al.*, 2004; Engelberth *et al.*, 2004) and the

priming of defense responses when actual herbivory occurs (Ton *et al.*, 2007; Frost *et al.*, 2008; Erb

et al., 2015; Hu *et al.*, 2019; Paudel Timilsena *et al.*, 2020). Here, we report that FR-enriched light

environment prompts maize plants to respond more strongly to HIPVs emitted by their neighbors. We

found that direct and priming of volatile release in response to HIPVs and (*Z*)-3-HAC is enhanced in

plants exposed to FR-enriched conditions, the typical spectral signature of the light environment in

closing plant canopies and shaded environments. In line with our findings, Chautá and Kessler (2022)

also reported a higher emission of a sesquiterpene in tall goldenrod (*Solidago altissima*) plants

exposed to supplemental FR and HIPVs from neighbors, but the mechanism remained to be
determined.

Most of the volatile identities and PTR-TOF-MS emission patterns in the present work were
validated using a complementary dynamic volatile collection and GC-MS detection. The higher
sensitivity of the PTR-TOF-MS allowed the detection of less abundant volatiles such as TMTT, and it
also enhanced the statistical power of the analyses by increasing their temporal resolution. Strikingly,
(*Z*)-3-HAC-induced indole emissions showed varying patterns between experiments, being slightly
enhanced by FR in some of them and during the time-series analyses while strongly reduced in others
(Fig. 2b). Further experimentation is needed to determine the context-depended regulation of this
compound emissions.

Plant responses to volatiles are regulated by the jasmonate signaling pathway (Arimura *et al.*,
2000; Hu, 2022) as part of an integrated signaling network that controls plant growth and defense
responses (Schmelz *et al.*, 2003; Bosch *et al.*, 2014; Mujiono *et al.*, 2021). In maize, (*Z*)-3-HAC
induces the accumulation of jasmonates and expression of JA-related marker genes before and after
herbivory induction (Hu *et al.*, 2019). Here, we confirmed that (*Z*)-3-HAC strongly up-regulated JA
signaling in maize. Furthermore, we found that low R:FR or FR-enriched environment resulted in a
two-fold higher induction of JA-Ile levels. This is surprising considering the reported suppressing
effects of low R:FR on JA accumulation (Fernandez-Milmanda et al., 2020) and signaling in
Arabidopsis (Cerrudo *et al.*, 2012; de Wit *et al.*, 2013; Chico *et al.*, 2014; Leone *et al.*, 2014). Yet, low
R:FR induced characteristic shade-associated morphological responses in maize, such as the
elongation of stems. Furthermore, we observed strong inductions of auxin and ABA levels in low R:FR
treated plants, with both hormones involved in shade-induced suppression of branching (tillering in
grasses) (Küpers *et al.*, 2020; Fernández-Milmanda & Ballaré, 2021). PAR levels and R:FR ratios
used in our work were similar to conditions employed in previous work reporting the effects of
supplemental FR on hormonal signaling and defenses in Arabidopsis (Pierik *et al.*, 2009; Wit *et al.*,
2013) and tomato (Courbier *et al.*, 2020; Courbier *et al.*, 2021) Thus, although maize responses to
shade show similarities with those described for Arabidopsis and other plant species, there might be
significant points of divergence. Additional analyses using reverse genetics and pharmacological
approaches can reveal the importance of the enhanced jasmonate response and the potential role of
auxin and ABA in maize responses to (*Z*)-3-HAC.

Plant volatile emission is controlled by biochemical, physiological, and physicochemical
processes including the rate of volatile biosynthesis, the activities of enzymes, the volatility and
diffusion rates of distinct volatile compounds, and stomatal conductance (Niinemets *et al.*, 2004; Seidl-
Adams *et al.*, 2015). Our study showed that low R:FR changed the expression of a subset of volatile
biosynthesis genes in response to (*Z*)-3-HAC and simulated herbivory. However, the observed
interactions were weak and did not track the changes in volatile emissions. Thus, it is unlikely that light

and volatile cues interact at this level to boost volatile release in maize. Low R:FR effects on the
diffusion of volatiles through the cuticle is also an unlikely explanation, as the cuticular diffusion
coefficients of monoterpenes is lower than the diffusion coefficient of air and water, indicating that the
cuticle provides an efficient barrier for these plant volatiles (Niinemets, 2003). We therefore reasoned
that volatile emissions might be alternatively modulated by induced changes in stomatal conductance.
Our results showed that both exposure to (*Z*)-3-HAC and simulated herbivory reduced leaf stomatal
conductance, which contrasted with the strong increase in volatile emissions. This might be explained
by the fact that leaf stomata can only limit volatile emission if the decrease in stomatal conductance
(i.e., closure) is larger than the increase in the volatile diffusion gradient (Lin *et al.*, 2022). Most
probably, (*Z*)-3-HAC induction of volatile biosynthesis quickly increased their concentration in the
intercellular air spaces, enhancing volatile diffusion gradients and overruling stomatal limitations.

Interestingly, FR-enriched environment had a positive effect on leaf stomatal conductance.
This is consistent with previous findings in bean (*Phaseolus vulgaris*) where supplemental FR added
from above to sub-saturating levels of visible light increased stomatal conductance (Holmes *et al.*,
1986). More recently, FR has been shown to enhance photochemical efficiency and photosynthesis
in several dicotyledonous and C3 and C4 monocotyledonous species, including maize, when added
to a background of white light (Zhen & Bugbee, 2020). In maize, FR photons were recently shown to
account for a large fraction of the photosynthetic activity under vegetation shade (Zhen *et al.*, 2022).
This was explained by their enhancing linear electron transport efficiency and cyclic electron flow
around the photosystem I (Zhen *et al.*, 2022). In C4 species such as maize, this is especially relevant
due to the extra ATP requirement for the CO_2-concentrating mechanism in the bundle sheath cells
(Johnson, 2011). Hence, our results align with the expected stomatal and photosynthetic responses
of maize plants in dense canopies characterized by low irradiances in the visible light range (400-700
nm) and enrichment in the FR region of the spectrum. Zhen and Bugbee (2020) also found that the
addition of equal amounts of FR or white photons causes similar increases in net photosynthesis,
while the increase in stomatal conductance was comparatively smaller under FR. This suggested that
stomatal responses to FR were most likely a response to the increase in photosynthetic rate. Stomatal
conductance in Arabidopsis *phyB* mutants have been shown to be modulated by changes in ABA
sensitivity under drought stress (González *et al.*, 2012). Whether the increase in stomatal conductance
from adding FR result from increased photosynthetic rates and/or of alteration of ABA sensitivity needs
to be established. Yet, it is conceivable that the effects of FR-enriched environment on the emission
of volatiles in response to (*Z*)-3-HAC might be explained, at least in part, by the positive effects of
supplemental FR on stomatal conductance and/or higher photosynthetic rates. The latter might affect
the availability of precursors for volatile biosynthesis (Arimura *et al.*, 2008).

Unlike FR supplementation experiments, *phyB1phyB2* plants displayed a diminished response
to (*Z*)-3-HAC. These results align with the suppressive effect of phyB inactivation on plant defenses

(Ballaré, 2014; Pierik & Ballaré, 2021). We reasoned that long-term phyB inactivation (as present in

the *phyB1phyB2* mutant) may have different effects from those triggered by a short-term inactivation

via changes in R:FR. Indeed, longer exposure of maize seedlings to low R:FR resulted in a slightly

weaker effect of low R:FR on (*Z*)-3-HAC-induced volatile emissions. At the same time, the

phyB1phyB2 mutant retained some ability to respond to low R:FR with higher (*Z*)-3-HAC induced

volatile emissions, reinforcing the potential effects of FR on the photosynthetic machinery and stomatal

conductance. We therefore conclude that (i) the duration of phyB inactivation might explain FR effects

on maize responses to (*Z*)-3-HAC, and that (ii) an enhanced photosynthetic efficiency and/or (iii) other

FR photoreceptors may also play a role in this phenomenon. Under deep shade (R:FR < 0.8), phyA is

strongly activated and prevents the excessive elongation of seedlings in Arabidopsis, antagonizing

phyB FR-induced responses (Casal *et al.*, 2014; Martínez-García *et al.*, 2014; Lim *et al.*, 2018; Yang

et al., 2018). Furthermore, PhyA promotes photosynthetic processes in rice (Panda *et al.*, 2023). PhyA

therefore represents a candidate that may be involved in the responses to light and volatile cues in

maize.

From a biological perspective, the capacity to integrate light and volatile cues would seem

particularly beneficial to fend off an attacking herbivore. While induced volatiles can serve as reliable

cues for the presence of herbivores on neighboring plants (Moreira *et al.*, 2018; Meents & Mithöfer,

2020), light cues can provide information about the proximity of neighbors and, therefore, the risk of

being attacked next. Because the effects of herbivore attack are more severe under competition (Vries

et al., 2018), it might be advantageous for a plant to respond more strongly to induced volatiles in the

presence of neighboring plants and vegetative shade. The enhanced emission of volatiles might, for

instance, promote the attraction of natural enemies in a setting where their recruitment is a competitive

struggle (Kessler & Baldwin, 2001; Cortés et al., 2016; Aartsma *et al.*, 2017). Plant volatiles may also

decrease ovipositing of gravid female moths (Moraes *et al.*, 2001). However, these volatiles may also

attract herbivores themselves (Mérey *et al.*, 2013), in which case becoming "apparent" in a dense

stand would be the wrong strategy. Further studies on the adaptive benefits of volatile responses in

dense stands will help us to understand the evolutionary and adaptive character of plant responses to

light and volatile cues in agricultural and ecological settings. Clearly, the integration of light and volatile

cues deserves attention as a potentially major determinant of plant-plant, plant-herbivore, tri-trophic

interaction dynamics, and community structure.

Acknowledgments

We thank Prof. Dr. Tobias Züst for his help in the optimization of GC-MS protocols and the identification

of maize volatile compounds. We also thank Prof. Dr. Christian Fankhauser and Dr. Martina Legris for

their help with light spectral measurements. The work of BCJS was supported by the Marie

Sklodowska-Curie Action Individual Fellowship (European Union Horizon 2020, Grant Nr. 794,947).

The work of LW was supported by the Marie Sklodowska-Curie Action Individual Fellowship (European
Union Horizon 2020, Grant Nr. 886651). This work was supported by the Velux Foundation (Grant Nr.
1231), the Swiss National Science Foundation (Grant. Nr. 200355), the State Secretariat for
Education, Research, and Innovation SERI (Project CANWAS) and the University of Bern.

Author contributions

ME conceived the study. ME, RE-B, BCJS, CB, and CAMR designed experiments. RE-B conducted
the experiments. RE-B performed the PTR-TOF-MS measurements and analyses. LW assisted with
PTR-TOF-MS measurements and conducted the analysis of linalool emissions. RE-B and BCJS
conducted gene expression analyses. RE-B and GG performed hormone analyses. RE-B and YZ
conducted stomatal measurements. RE-B analyzed the data. RE-B, BCJS, CB and ME interpreted the
data. RE-B and ME wrote the manuscript and all the authors contributed to revisions.

Data availability

The data generated for this manuscript will be archived in Dryad and the data DOI will be included at
the end of the article.

References

Aartsma Y, Bianchi FJJA, van der Werf W, Poelman EH, Dicke M. 2017. Herbivore-induced plant
volatiles and tritrophic interactions across spatial scales. *New Phytologist* **216**: 1054–1063.
**Alba JM, Schimmel BCJ, Glas JJ, Ataide LMS, Pappas ML, Villarroel CA, Schuurink RC, Sabelis
MW, Kant MR. 2015.** Spider mites suppress tomato defenses downstream of jasmonate and
salicylate independently of hormonal crosstalk. *The New phytologist* **205**: 828–840.
**Arimura G, Köpke S, Kunert M, Volpe V, David A, Brand P, Dabrowska P, Maffei ME, Boland W.
2008.** Effects of feeding Spodoptera littoralis on lima bean leaves: IV. Diurnal and nocturnal damage
differentially initiate plant volatile emission. *Plant Physiology* **146**: 965–973.
Arimura G, Ozawa R, Nishioka T, Boland W, Koch T, Kühnemann F, Takabayashi J. 2002.
Herbivore-induced volatiles induce the emission of ethylene in neighboring lima bean plants. *The
Plant journal : for cell and molecular biology* **29**: 87–98.
Arimura G, Ozawa R, Shimoda T, Nishioka T, Boland W, Takabayashi J. 2000. Herbivory-induced
volatiles elicit defence genes in lima bean leaves. *Nature* **406**: 512–515.
Ballaré CL. 2009. Illuminated behaviour: phytochrome as a key regulator of light foraging and plant
anti-herbivore defence. *Plant, Cell & Environment* **32**: 713–725.
Ballaré CL. 2014. Light regulation of plant defense. *Annual review of plant biology* **65**: 335–363.
Ballaré CL, Pierik R. 2017. The shade-avoidance syndrome: multiple signals and ecological
consequences. *Plant, Cell & Environment* **40**: 2530–2543.
Ballaré CL, Scopel AL, Sánchez RA. 1990. Far-red radiation reflected from adjacent leaves: an early
signal of competition in plant canopies. *Science* **247**: 329–332.
Barbosa P, Hines J, Kaplan I, Martinson H, Szczepaniec A, Szendrei Z. 2009. Associational
Resistance and Associational Susceptibility: Having Right or Wrong Neighbors. *Annual Review of
Ecology, Evolution, and Systematics* **40**: 1–20.
Block AK, Hunter CT, Rering C, Christensen SA, Meagher RL. 2018. Contrasting insect attraction
and herbivore-induced plant volatile production in maize. *Planta* **248:** 105–116.

Bosch M, Wright LP, Gershenzon J, Wasternack C, Hause B, Schaller A, Stintzi A. 2014.
Jasmonic acid and its precursor 12-oxophytodienoic acid control different aspects of constitutive
and induced herbivore defenses in tomato. *Plant Physiology* **166**: 396–410.

Bruinsma M, Posthumus MA, Mumm R, Mueller MJ, van Loon JJA, Dicke M. 2009. Jasmonic
acid-induced volatiles of Brassica oleracea attract parasitoids: effects of time and dose, and
comparison with induction by herbivores. *Journal of Experimental Botany* **60**: 2575–2587.

Campos ML, Yoshida Y, Major IT, Oliveira Ferreira D de, **Weraduwage SM, Froehlich JE,**
Johnson BF, Kramer DM, Jander G, Sharkey TD *et al.* **2016.** Rewiring of jasmonate and
phytochrome B signalling uncouples plant growth-defense tradeoffs. *Nature Communications* **7**:
12570.

Cargnel MD, Demkura PV, Ballaré CL. 2014. Linking phytochrome to plant immunity: low red : far-
red ratios increase Arabidopsis susceptibility to Botrytis cinerea by reducing the biosynthesis of
indolic glucosinolates and camalexin. *New Phytologist* **204**: 342–354.

Casal JJ, Candia AN, Sellaro R. 2014. Light perception and signalling by phytochrome A. *Journal of*
Experimental Botany **65**: 2835–2845.

Cerrudo I, Keller MM, Cargnel MD, Demkura PV, de Wit M, Patitucci MS, Pierik R, Pieterse CMJ,
Ballaré CL. 2012. Low red/far-red ratios reduce Arabidopsis resistance to Botrytis cinerea and
jasmonate responses via a COI1-JAZ10-dependent, salicylic acid-independent mechanism. *Plant*
physiology **158:** 2042–2052.

Chautá A, Kessler A. Metabolic integration of spectral and chemical cues mediating plant responses
to competitors and herbivores. *Plants* **11**(20), 2768.

Chico J-M, Fernández-Barbero G, Chini A, Fernández-Calvo P, Díez-Díaz M, Solano R. 2014.
Repression of Jasmonate-Dependent Defenses by Shade Involves Differential Regulation of
Protein Stability of MYC Transcription Factors and Their JAZ Repressors in Arabidopsis. *The Plant*
Cell **26**: 1967–1980.

Choh Y, Shimoda T, Ozawa R, Dicke M, Takabayashi J. 2004. Exposure of lima bean leaves to
volatiles from herbivore-induced conspecific plants results in emission of carnivore attractants:
active or passive process? *Journal of Chemical Ecology* **30**: 1305–1317.

Chrétien LTS, David A, Daikou E, Boland W, Gershenzon J, Giron D, Dicke M, Lucas-Barbosa
D. 2018. Caterpillars induce jasmonates in flowers and alter plant responses to a second attacker.
New Phytologist **217**: 1279–1291.

Cortés LE, Weldegergis BT, Boccalandro HE, Dicke M, Ballaré CL. 2016. Trading direct for indirect
defense? Phytochrome B inactivation in tomato attenuates direct anti-herbivore defenses whilst
enhancing volatile-mediated attraction of predators. *New Phytologist* **212**: 1057–1071.

Courbier S, Grevink S, Sluijs E, Bonhomme P-O, Kajala K, van Wees SCM, Pierik R. 2020. Far-
red light promotes Botrytis cinerea disease development in tomato leaves via jasmonate-dependent
modulation of soluble sugars. *Plant, Cell & Environment* **43**: 2769–2781.

Courbier S, Snoek BL, Kajala K, Li L, van Wees SCM, Pierik R. 2021. Mechanisms of far-red light-
mediated dampening of defense against Botrytis cinerea in tomato leaves. *Plant Physiology* **187**:
1250–1266.

Dubois PG, Brutnell TP. 2009. Light Signal Transduction Networks in Maize. In: Bennetzen JL, Hake
SC, eds. *Handbook of maize. Its biology*. New York: Springer, 205–227.

Dubois PG, Olsefski GT, Flint-Garcia S, Setter TL, Hoekenga OA, Brutnell TP. 2010. Physiological
and genetic characterization of end-of-day far-red light response in maize seedlings. *Plant*
physiology **154**: 173–186.

Engelberth J, Alborn HT, Schmelz EA, Tumlinson JH. 2004. Airborne signals prime plants against
insect herbivore attack. *Proceedings of the National Academy of Sciences of the United States of*
America **101**: 1781–1785.

Erb M, Flors V, Karlen D, Lange E de, **Planchamp C, D'Alessandro M, Turlings TCJ, Ton J. 2009.** Signal signature of aboveground-induced resistance upon belowground herbivory in maize. *The Plant Journal* **59**: 292–302.

Erb M, Veyrat N, Robert CAM, Xu H, Frey M, Ton J, Turlings TCJ. 2015. Indole is an essential herbivore-induced volatile priming signal in maize. *Nature Communications* **6**: 6273.

Escobar-Bravo R, Klinkhamer PGL, Leiss KA. 2017. Interactive Effects of UV-B Light with Abiotic Factors on Plant Growth and Chemistry, and Their Consequences for Defense against Arthropod Herbivores. *Frontiers in Plant Science* **8**: 278.

Fankhauser C. 2001. The phytochromes, a family of red/far-red absorbing photoreceptors. *Journal of Biological Chemistry* **276**: 11453–11456.

Farag MA, Paré PW. 2002. C6-Green leaf volatiles trigger local and systemic VOC emissions in tomato. *Phytochemistry* **61**: 545–554.

Fernández-Milmanda GL, Ballaré CL. 2021. Shade Avoidance: Expanding the Color and Hormone Palette. *Trends in plant science* **26**: 509–523.

Fernández-Milmanda GL, Crocco CD, Reichelt M, Mazza CA, Köllner TG, Zhang T, Cargnel MD, Lichy MZ, Fiorucci A-S, Fankhauser C *et al.* **2020.** A light-dependent molecular link between competition cues and defence responses in plants. *Nature plants* **6**: 223–230.

Finch-Savage WE, Leubner-Metzger G. 2006. Seed dormancy and the control of germination. *The New phytologist* **171**: 501–523.

Fiorucci A-S, Fankhauser C. 2017. Plant Strategies for Enhancing Access to Sunlight. *Current biology : CB* **27**: R931-R940.

Franklin KA. 2008. Shade avoidance. *The New phytologist* **179**: 930–944.

Frost CJ, Mescher MC, Dervinis C, Davis JM, Carlson JE, de Moraes CM. **2008.** Priming defense genes and metabolites in hybrid poplar by the green leaf volatile cis-3-hexenyl acetate. *New Phytologist* **180**: 722–734.

Fu J, Ren F, Lu X, Mao H, Xu M, Degenhardt J, Peters RJ, Wang Q. 2016. A tandem array of ent-kaurene synthases in maize with roles in gibberellin and more specialized metabolism. *Plant Physiology* **170:** 742–751.

Glauser G, Vallat A, Balmer D. 2014. Hormone profiling. *Methods in molecular biology (Clifton, N.J.)* **1062**: 597–608.

Goning M, Lopez-hilfiker F, Hutterli M, Erb M, Pfander M. 2022. Autosampler. Patent number US11227755B2. https://worldwide.espacenet.com/patent/search/family/061157140/publication/US11227755B2?q=erb%20tofwerk

González CV, Ibarra SE, Piccoli PN, Botto JF, Boccalandro HE. 2012. Phytochrome B increases drought tolerance by enhancing ABA sensitivity in Arabidopsis thaliana. *Plant, Cell & Environment* **35**: 1958–1968.

Gouinguené SP, Turlings TCJ. 2002. The effects of abiotic factors on induced volatile emissions in corn plants. *Plant Physiology* **129**: 1296–1307.

Helms AM, Moraes CM de, **Tröger A, Alborn HT, Francke W, Tooker JF, Mescher MC. 2017.** Identification of an insect-produced olfactory cue that primes plant defenses. *Nature Communications* **8**: 337.

Holmes MG, Sager JC, Klein WH. 1986. Sensitivity to far-red radiation in stomata of Phaseolus vulgaris L.: Rhythmic effects on conductance and photosynthesis. *Planta* **168**: 516–522.

Hu L. 2022. Integration of multiple volatile cues into plant defense responses. *New Phytologist* **233**: 618–623.

Hu L, Ye M, Erb M. 2019. Integration of two herbivore-induced plant volatiles results in synergistic effects on plant defence and resistance. *Plant, Cell & Environment* **42**: 959–971.

Johnson GN. 2011. Physiology of PSI cyclic electron transport in higher plants. *Biochimica et biophysica acta* **1807**: 384–389.

Kegge W, Ninkovic V, Glinwood R, Welschen RAM, Voesenek LACJ, Pierik R. 2015. Red:far-red light conditions affect the emission of volatile organic compounds from barley (Hordeum vulgare), leading to altered biomass allocation in neighbouring plants. *Annals of botany* **115**: 961–970.

Kegge W, Weldegergis BT, Soler R, Eijk MV-V, Dicke M, Voesenek LACJ, Pierik R. 2013. Canopy light cues affect emission of constitutive and methyl jasmonate-induced volatile organic compounds in Arabidopsis thaliana. *New Phytologist* **200**: 861–874.

Kessler A, Baldwin IT. 2001. Defensive function of herbivore-induced plant volatile emissions in nature. *Science (New York, N.Y.)* **291**: 2141–2144.

Kroes A, van Loon JJA, Dicke M. 2015. Density-dependent interference of aphids with caterpillar-induced defenses in Arabidopsis: involvement of phytohormones and transcription factors. *Plant & cell physiology* **56**: 98–106.

Küpers JJ, Oskam L, Pierik R. 2020. Photoreceptors Regulate Plant Developmental Plasticity through Auxin. *Plants* **9**: 940.

Leone M, Keller MM, Cerrudo I, Ballaré CL. 2014. To grow or defend? Low red : far-red ratios reduce jasmonate sensitivity in Arabidopsis seedlings by promoting DELLA degradation and increasing JAZ10 stability. *New Phytologist* **204**: 355–367.

Lim J, Park J-H, Jung S, Hwang D, Nam HG, Hong S. 2018. Antagonistic Roles of PhyA and PhyB in Far-Red Light-Dependent Leaf Senescence in Arabidopsis thaliana. *Plant & cell physiology* **59**: 1753–1764.

Lin P-A, Chen Y, Ponce G, Acevedo FE, Lynch JP, Anderson CT, Ali JG, Felton GW. 2022. Stomata-mediated interactions between plants, herbivores, and the environment. *Trends in plant science* **27**: 287–300.

Maag D, Dalvit C, Thevenet D, Köhler A, Wouters FC, Vassão DG, Gershenzon J, Wolfender J-L, Turlings TCJ, Erb M *et al.* 2014. 3-β-D-Glucopyranosyl-6-methoxy-2-benzoxazolinone (MBOA-N-Glc) is an insect detoxification product of maize 1,4-benzoxazin-3-ones. *Phytochemistry* **102**: 97–105.

Maddonni GA, Otegui ME, Andrieu B, Chelle M, Casal JJ. 2002. Maize leaves turn away from neighbors. *Plant Physiology* **130**: 1181–1189.

Martínez-García JF, Gallemí M, Molina-Contreras MJ, Llorente B, Bevilaqua MRR, Quail PH. 2014. The shade avoidance syndrome in Arabidopsis: the antagonistic role of phytochrome a and B differentiates vegetation proximity and canopy shade. *PLOS ONE* **9**: e109275.

Meents AK, Mithöfer A. 2020. Plant-Plant Communication: Is There a Role for Volatile Damage-Associated Molecular Patterns? *Frontiers in plant science* **11**: 583275.

Mérey GE von, Veyrat N, D'Alessandro M, Turlings TCJ. 2013. Herbivore-induced maize leaf volatiles affect attraction and feeding behavior of Spodoptera littoralis caterpillars. *Frontiers in Plant Science* **4**: 209.

Mérey G von, Veyrat N, Mahuku G, Valdez RL, Turlings TCJ, D'Alessandro M. 2011. Dispensing synthetic green leaf volatiles in maize fields increases the release of sesquiterpenes by the plants, but has little effect on the attraction of pest and beneficial insects. *Phytochemistry* **72**: 1838–1847.

Meza-Canales ID, Meldau S, Zavala JA, Baldwin IT. 2017. Herbivore perception decreases photosynthetic carbon assimilation and reduces stomatal conductance by engaging 12-oxo-phytodienoic acid, mitogen-activated protein kinase 4 and cytokinin perception. *Plant, Cell & Environment* **40:** 1039– 1056.

Moraes CM de, Mescher MC, Tumlinson JH. 2001. Caterpillar-induced nocturnal plant volatiles repel conspecific females. *Nature* **410**: 577–580.

Moreira X, Nell CS, Katsanis A, Rasmann S, Mooney KA. 2018. Herbivore specificity and the
chemical basis of plant-plant communication in Baccharis salicifolia (Asteraceae). *New Phytologist*
220: 703–713.
Moreno JE, Tao Y, Chory J, Ballaré CL. 2009. Ecological modulation of plant defense via
phytochrome control of jasmonate sensitivity. *Proceedings of the National Academy of Sciences of*
the United States of America **106**: 4935–4940.
Mujiono K, Tohi T, Sobhy IS, Hojo Y, Shinya T, Galis I. 2021. Herbivore-induced and constitutive
volatiles are controlled by different oxylipin-dependent mechanisms in rice. *Plant, Cell &*
Environment **44**: 2687–2699.
Nguyen D, Rieu I, Mariani C, van Dam NM. 2016. How plants handle multiple stresses: hormonal
interactions underlying responses to abiotic stress and insect herbivory. *Plant Molecular Biology* **91**:
727–740.
Niinemets U, Loreto F, Reichstein M. 2004. Physiological and physicochemical controls on foliar
volatile organic compound emissions. *Trends in plant science* **9**: 180–186.
Niinemets Ü. 2003. Controls on the emission of plant volatiles through stomata: Differential sensitivity
of emission rates to stomatal closure explained. *Journal of Geophysical Research* **108**.
Panda D, Dash GK, Mohanty S, Sekhar S, Roy A, Tudu C, Behera L, Tripathy BC, Baig MJ, 2023.
Phytochrome A mediated modulation of photosynthesis, development and yield in rice (*Oryza sativa*
L.) in fluctuating light environment. *Environmental and Experimental Botany*, *206*, p.105183.
Paudel Timilsena B, Seidl-Adams I, Tumlinson JH. 2020. Herbivore-specific plant volatiles prime
neighboring plants for nonspecific defense responses. *Plant, Cell & Environment* **43**: 787–800.
Pierik R, Ballaré CL. 2021. Control of Plant Growth and Defense by Photoreceptors: From
Mechanisms to Opportunities in Agriculture. *Molecular Plant* **14**: 61–76.
Pierik R, Djakovic-Petrovic T, Keuskamp DH, Wit M de, **Voesenek LACJ. 2009.** Auxin and ethylene
regulate elongation responses to neighbor proximity signals independent of gibberellin and della
proteins in Arabidopsis. *Plant Physiology* **149**: 1701–1712.
Pierik R, Wit M de. **2014.** Shade avoidance: phytochrome signalling and other aboveground
neighbour detection cues. *Journal of experimental botany* **65**: 2815–2824.
**Richter A, Schaff C, Zhang Z, Lipka AE, Tian F, Köllner TG, Schnee C, Preiß S, Irmisch S, Jander
G** *et al.* **2016.** Characterization of Biosynthetic Pathways for the Production of the Volatile
Homoterpenes DMNT and TMTT in Zea mays. *The Plant cell* **28**: 2651–2665.
Richter A, Seidl-Adams I, Köllner TG, Schaff C, Tumlinson JH, Degenhardt J. 2015. A small,
differentially regulated family of farnesyl diphosphate synthases in maize (Zea mays) provides
farnesyl diphosphate for the biosynthesis of herbivore-induced sesquiterpenes. *Planta* **241**: 1351–
1361.
**Savchenko T, Kolla VA, Wang CQ, Nasafi Z, Hicks DR, Phadungchob B, Chehab WE, Brandizzi
F, Froehlich J, Dehesh K. 2014.** Functional convergence of oxylipin and abscisic acid pathways
controls stomatal closure in response to drought. *Plant Physiology* **164:** 1151–1160.
Schmelz EA, Alborn HT, Banchio E, Tumlinson JH. 2003. Quantitative relationships between
induced jasmonic acid levels and volatile emission in Zea mays during Spodoptera exigua herbivory.
Planta **216**: 665–673.
Schnee C, Köllner TG, Gershenzon J, Degenhardt J. 2002. The maize gene terpene synthase 1
encodes a sesquiterpene synthase catalyzing the formation of (E)-beta-farnesene, (E)-nerolidol,
and (E,E)-farnesol after herbivore damage. *Plant Physiology* **130**: 2049–2060.
Schnee C, Köllner TG, Held M, Turlings TCJ, Gershenzon J, Degenhardt J. 2006. The products
of a single maize sesquiterpene synthase form a volatile defense signal that attracts natural
enemies of maize herbivores. *Proceedings of the National Academy of Sciences of the United*
States of America **103**: 1129–1134.

Seidl-Adams I, Richter A, Boomer KB, Yoshinaga N, Degenhardt J, Tumlinson JH. 2015.
Emission of herbivore elicitor-induced sesquiterpenes is regulated by stomatal aperture in maize
(Zea mays) seedlings. *Plant, Cell & Environment* **38**: 23–34.
Sheehan MJ, Kennedy LM, Costich DE, Brutnell TP. 2007. Subfunctionalization of PhyB1 and
PhyB2 in the control of seedling and mature plant traits in maize. *The Plant journal : for cell and*
molecular biology **49**: 338–353.
Soler R, Erb M, Kaplan I. 2013. Long distance root-shoot signalling in plant-insect community
interactions. *Trends in plant science* **18**: 149–156.
Ton J, D'Alessandro M, Jourdie V, Jakab G, Karlen D, Held M, Mauch-Mani B, Turlings TCJ.
2007. Priming by airborne signals boosts direct and indirect resistance in maize. *The Plant journal*
: for cell and molecular biology **49**: 16–26.
Vries J de, **Poelman EH, Anten N, Evers JB. 2018.** Elucidating the interaction between light
competition and herbivore feeding patterns using functional-structural plant modelling. *Annals of*
botany **121**: 1019–1031.
Way DA, Pearcy RW. 2012. Sunflecks in trees and forests: from photosynthetic physiology to global
change biology. *Tree physiology* **32**: 1066–1081.
Wit M de, **Keuskamp DH, Bongers FJ, Hornitschek P, Gommers CMM, Reinen E, Martínez-Cerón**
C, Fankhauser C, Pierik R. 2016. Integration of Phytochrome and Cryptochrome Signals
Determines Plant Growth during Competition for Light. *Current biology : CB* **26**: 3320–3326.
Wit M de, **Spoel SH, Sanchez-Perez GF, Gommers CMM, Pieterse CMJ, Voesenek LACJ, Pierik**
R. 2013. Perception of low red:far-red ratio compromises both salicylic acid- and jasmonic acid-
dependent pathogen defences in Arabidopsis. *The Plant Journal* **75**: 90–103.
Xu G, Cao J, Wang X, Chen Q, Jin W, Li Z, Tian F. 2019. Evolutionary Metabolomics Identifies
Substantial Metabolic Divergence between Maize and Its Wild Ancestor, Teosinte. *The Plant Cell*
31: 1990–2009.
Xue J, Gou L, Zhao Y, Yao M, Yao H, Tian J, Zhang W. 2016. Effects of light intensity within the
canopy on maize lodging. *Field Crops Research* **188**: 133–141.
Yang C, Xie F, Jiang Y, Li Z, Huang X, Li L. 2018. Phytochrome A Negatively Regulates the Shade
Avoidance Response by Increasing Auxin/Indole Acidic Acid Protein Stability. *Developmental Cell*
44: 29-41.e4.
Ye M, Glauser G, Lou Y, Erb M, Hu L. 2019. Molecular Dissection of Early Defense Signaling
Underlying Volatile-Mediated Defense Regulation and Herbivore Resistance in Rice. *The Plant Cell*
31: 687–698.
Zhang P-J, Broekgaarden C, Zheng S-J, Snoeren TAL, van Loon JJA, Gols R, Dicke M. 2013.
Jasmonate and ethylene signaling mediate whitefly-induced interference with indirect plant defense
in Arabidopsis thaliana. *New Phytologist* **197**: 1291–1299.
Zhen S, Bugbee B. 2020. Far-red photons have equivalent efficiency to traditional photosynthetic
photons: Implications for redefining photosynthetically active radiation. *Plant, Cell & Environment*
43: 1259–1272.
Zhen S, van Iersel MW, Bugbee B. 2022. Photosynthesis in sun and shade: the surprising importance
of far-red photons. *New Phytologist.* doi: 10.1111/nph.18375.
918

919 **Supporting Information**

920 **Figure S1.** Light spectral composition determined in FR supplementation experiments.

921 **Figure S2.** Effects of FR-enrichment on maize growth and auxin levels.

922 **Figure S3.** Aboveground biomass in maize plants infested with *Spodoptera littoralis* larvae.

Figure S4. Effect of FR-enrichment on herbivory-induced plant volatile emission in maize

Figure S5. Effect of FR-enrichment on maize volatile emission during exposure to (*Z*)-3-hexenyl acetate and after simulated herbivory measured via GC-MS methods.

Figure S6. Measurement of pure linalool in the PTR-TOF-MS system.

Figure S7. Effect of FR-enrichment and (*Z*)-3-hexenyl acetate on volatile emissions in 30-day-old maize plants.

Figure S8 Spatial variations in photosynthetically active radiation (PAR) levels in the experimental set up did not influence 30-day-old maize responses to (*Z*)-3-hexenyl acetate.

Figure S9. Effect of FR-enrichment on the growth of *phyB1phyB2* maize double mutant and its wild-type.

Figure S10. Effect of FR-enrichment on volatiles emission during exposure to (*Z*)-3-hexenyl acetate and after simulated herbivory in the maize inbred line W22.

Figure S11. Effect of FR-enrichment on volatiles emission during exposure to (*Z*)-3-hexenyl acetate and after simulated herbivory in the *phyB1phyB2* maize double mutant.

Figure S12. Effect of FR-enrichment and (*Z*)-3-hexenyl acetate on stomatal density.

Supplementary Method S1. Detailed protocol for phytohormone extraction and analysis.

Supplementary Table S1. Primer's specifications for gene expression analyses