Immunological landscape of human lymphoid explants during measles virus infection

Joshua A Acklin*[1,2], Aum R Patel*[1,2], Andrew P Kurland[1], Shu Horiuchi[1], Arianna S Moss[1], Emma
J Degrace[1,2], Satoshi Ikegame[1], Jillian Carmichael[1], Shreyas Kowdle[1], Patricia Thibault[1], Naoko
Imai[1], Hideki Ueno[1], Benjamin Tweel[3], Jeffrey R Johnson[1], Brad R Rosenberg[1], Benhur Lee[1], and
Jean K Lim[1]

AFFILIATIONS:
[1] Department of Microbiology, Icahn School of Medicine at Mount Sinai, New York, NY, United
States,
[2] Graduate School of Biomedical Sciences, Icahn School of Medicine at Mount Sinai, New York,
NY, United States
[3] Department of Otolaryngology, Icahn School of Medicine at Mount Sinai, New York, NY, United
States
*indicates co-first authorship

Corresponding Author: One Gustave L. Levy Place, Box 1124, New York, NY. 10029;

jean.lim@mssm.edu; +1(212)-241-7811

ABSTRACT:

In humans, lymph nodes are the primary site of measles virus (MeV) replication. To understand

the immunological events that occur at this site, we infected human lymphoid tissue explants

using a pathogenic strain of MeV that expresses GFP. We found that MeV infected between 5-

15% of cells across donors. Using single cell RNA-Seq (scRNA-Seq) and flow cytometry, we

found that while most of the 29 cell populations identified in the lymphoid culture were susceptible

to MeV, there was a broad preferential infection of B cells and reduced infection of T cells. Further

subsetting of T cells revealed that this reduction may be driven by the decreased infection of naïve

T cells. Transcriptional changes in infected B cells were dominated by an interferon stimulated

gene (ISG) signature. To determine which of these ISGs were most substantial, we evaluated the

proteome of MeV-infected Raji cells by mass spectrometry. We found that IFIT1, IFIT2, IFIT3,

ISG15, CXCL10, MX2, and XAF1 proteins were the most highly induced, and positively correlated

with their expression in the transcriptome. These data provide insight into the immunological

events that occur in lymph nodes during infection and may lead to the development of therapeutic

interventions.

CONFLICTS OF INTEREST

The authors have declared that no conflict of interest exists.

INTRODUCTION:

Measles virus (MeV) is the most infectious human virus, with a reported R_0 value of 12-18 (1-7). MeV outbreaks have largely been controlled with the advent of the two-dose measles, mumps, and rubella (MMR) vaccine, yet MeV still causes ~200,000 deaths annually, primarily among unvaccinated children in developing countries (8-10). However, recent surges in vaccine-hesitancy have allowed MeV to re-emerge in countries like the United States and the United Kingdom, where the MMR vaccine coverage has historically been high (11-14). With many global vaccination campaigns for MeV stalled due to the COVID-19 pandemic, the risk of MeV outbreaks globally continues to grow (15). Compounding these crises is the lack of any licensed antiviral that targets MeV once individuals are infected (16, 17).

MeV is a morbillivirus of the family *Paramyxoviridae* that is transmitted through the respiratory tract, where alveolar macrophages and dendritic cells are the initial cellular targets of infection (18, 19). These infected immune cells then traffic to the draining lymph nodes where the virus replicates rapidly in lymphocytes that express the entry factor CD150/SLAMF1 (20-23), followed by egress through lung epithelium that is mediated by basolateral expression of the Nectin-4 receptor (23-27). MeV is also known for causing immune amnesia through the depletion of CD150[+] B and T cells in both primary and secondary lymphoid organs, increasing the morbidity and mortality rates from secondary infections with common childhood pathogens (28-35). Immunological amnesia following MeV infection has been shown to markedly reduce the antibody repertoire towards common childhood pathogens, such as the human parainfluenza viruses, respiratory syncytial virus, coronaviruses, and cytomegalovirus (36). Given that immune responses are primarily generated in the draining lymph node, and that this site is a critical launching pad for MeV infections, understanding virus: host interactions at this site is paramount for identifying factors that shape disease progression.

Modern reanalysis of early work on immunological amnesia caused by measles implicates T cells due to a delayed type-I hypersensitivity response to tuberculin antigen (37, 38). In vitro

characterization of MeV infection in primary lymphocytes revealed that B cells are the most

extensively infected population of lymphocytes, consistent with their high CD150 expression (39).

Characterization of MeV infection in PBMCs from both humans and macaques further

demonstrated the propensity of MeV for lymphocytes and that MeV infection is biased towards

naive B cells, memory B cells, and memory T lymphocytes, which was subsequently validated in

PBMCs derived from MeV-infected children (18, 34, 40, 41).

While these studies provide insights into MeV infection of lymphocytes, they do not

examine infection within the complex architecture of secondary lymphoid tissue. The draining

lymph nodes are organized with high-density B cell follicles surrounded by T cell zones (42), which

may be important to determine the in vivo cellular susceptibility as well as the kinetics of lymph

node infection. Studies in macaques have examined the geographic distribution of infected cells

within secondary lymphoid tissues and have identified that the majority of infection is established

within B-cell follicles (41). Further, these studies recapitulated the heightened susceptibility to

infection among memory but not naïve lymphocyte subsets within secondary lymphoid organs

(41). Studies using human tonsil explants have also been conducted, where lymphoid tissue

structures and native cell ratios are intact, which found that B cells are a preferential target of MeV

infection and that memory T cells were more extensively infected compared to naive T cell subsets

(43, 44). In this study, we revisited this human lymphoid explant model using a GFP-expressing

pathogenic isolate of MeV which has been shown to mimic human clinical outcomes in non-

human primates, commonly referred to as a "wild-type" isolate (45). Our findings both confirm and

extend our understanding of cellular susceptibility to MeV in humans. While a similar analysis of

MeV-infected airway epithelium has been conducted (46), we present transcriptional signatures

of MeV-infected lymphocytes with single cell resolution in human lymphoid tissue explants and

link these transcriptional signatures with translated products in the proteome.

RESULTS

MeV replicates efficiently in human lymphoid tissues ex vivo:

To evaluate how MeV infection proceeds within human lymphoid tissue, we infected human tonsil tissues ex vivo, the most accessible lymphoid tissue for laboratory use. Tissue samples from routine, non-inflamed tonsillectomies were infected with a pathogenic isolate of MeV (IC323) that expresses GFP (MeV-GFP) as previously described (44, 45, 47). While this model lacks a functional lymphatic system, and thus does not exactly mimic the way that MeV enters the draining lymph node during human infections, it benefits from retaining the 3-dimensional tissue architecture of human lymph tissue. To assess the extent to which MeV could replicate in human lymphoid tissues across eleven donors, we collected culture supernatants at days 3, 6, and 8 post-infection and measured virus production by fluorescence plaque assay. As shown in **Figure 1A**, viral titers increased for all donors, with ~2.5 log increase over the 8-day culture.

To assess the extent of infection within the tissues and to further confirm productive infection, we measured the frequency of GFP$^+$ cells over time by flow cytometry. As shown in **Figure 1B**, for one representative donor, the percentage of GFP$^+$ cells increased over time, while no GFP signal was detected in the uninfected condition. Quantification across 4 donors revealed a frequency of MeV-infected cells that ranged between 5-15% of live cells by day 8 post-infection (**Figure 1C**). Further, RNA transcripts for GFP were readily detected within the tissues through in situ hybridization (**Figure 1D**). Together, our data establish that human tonsils infected ex vivo are susceptible to MeV without any stimulation or infection enhancers, providing us with a robust model system to further characterize MeV-infected cells.

scRNA-Seq of MeV-infected tissues reveals broad cellular susceptibility:

Given the high percentage of GFP$^+$ cells detected, we sought to sort and analyze infected cells

using scRNA-Seq. To do this, we generated single cell suspensions from MeV-infected and

donor-matched uninfected tonsil tissue on day 8 post-infection, the time point at which the

maximum number of infected cells was observed. We sorted GFP$^+$ (infected) and GFP$^-$

(bystander) cells from the MeV-infected condition, as well as the GFP$^-$ cells (uninfected) from the

uninfected condition for scRNA-Seq using a workflow shown in **Figure 2A**. We validated the

quality of sequencing across the three groups by quantifying unique molecular identifier (UMI)

counts, unique genes captured, and the representation of mitochondrial transcripts

(Supplementary Figure 1A-C). As expected, only cells from the infected group had appreciable

MeV transcripts (**Figure 2B**). Among the infected cells, we examined the expression levels of viral

transcripts as a final confirmation of infection status. Infected cells showed a transcriptional

gradient of viral genes from 3' to 5', consistent with the phenomenon of run-off transcription that

occurs for paramyxoviruses (**Figure 2C**).

Following data integration, we annotated constituent cell clusters across conditions based

on immune cell reference data and supervised differential marker gene analysis (see *Methods*).

Using this strategy, we identified 29 distinct populations of cells (**Figure 2D**), with the vast majority

belonging to either T or B cell subsets (annotation strategy shown in **Supplementary Figure 1D**).

Interestingly, GFP$^+$ transcripts were identified in all 29 populations, albeit to different levels in each

population, spanning subpopulations of CD4 and CD8 T cells, B cells, tonsillar epithelium, and

tonsillar stromal cells (**Figure 2E**). We were surprised by this finding and assessed the expression

of *SLAMF1*, the gene encoding the canonical MeV receptor CD150, in each cluster. As shown in

Figure S1E, the detection of *SLAMF1* in our dataset was not robust. However, we do detect

higher levels among B cell populations, particularly among activated B cells, which is consistent

with previous reports (39, 41, 43). We next quantified the frequency of each of these populations

within the uninfected, bystander, and infected cell-sorted sample groups and found that, despite

T cells comprising the largest population of cells in human tonsils (as shown in uninfected and

bystander groups), they were not the majority among the MeV-infected cells. Instead, IgD$^+$ B cells

were overrepresented within the pool of GFP$^+$ cells (**Figure 2F**). To assess the extent to which

the transcriptome is co-opted for viral gene expression, we quantified the percentage of UMI

counts that mapped to MeV UMIs for each identified cell population. As shown in **Figure 2G**, MeV

transcripts constituted ~0.93% of all transcripts for each cluster on average (calculated using the

median % viral UMI per cluster). One possible explanation for differences in the frequency of cell

clusters among the GFP$^+$ condition would be that cells had either proliferated or died during or

because of preferential infection of various cell subsets found within the tissue. To assess this

possibility, we evaluated the B cell clusters for gene signatures associated with proliferation. As

shown in **Figure S1F-G**, while we observed high pathway scores for S phase and G2/M phase

among CD4$^+$ Cells (annotated as proliferating in **Figure 2**), we did not observe any elevation

within any B cell cluster, regardless of infection status. However, as this analysis was limited to a

single donor at a single time point, we can only conclude that MeV has a wide cellular tropism

within the lymphoid explants and suggest that IgD$^+$ B cells are the primary target.

B cells are preferential targets of MeV infection in lymphoid tissue explants:

While the scRNA-Seq analysis suggested that differences in susceptibility to MeV may exist within

the lymphoid tissue explants, these data evaluated only a single time point for a single donor. To

define MeV infection across donors and over time, we immunophenotyped major cell subsets

identified in the scRNA-Seq dataset by flow cytometry (n=3). We first quantified the frequency of

MeV-infected B and T cells and compared these frequencies to their frequency among GFP$^-$

bystander cells (from the infected condition) and donor-matched, uninfected cells (gating

schemata in **Figure 3A**). For ease of data interpretation, we also show the frequencies of GFP

over time within cell populations discussed in this section in **Figure S2** and present a complete

gating schema for all analyses in **Figure S3**. As shown in **Figure 3B**, there was a significantly

higher frequency of CD19$^+$ B cells among the GFP$^+$ cells compared to their frequency in the

uninfected or bystander populations. This difference was established by day 3 and maintained

across the 8-day culture. Conversely, we observed a decrease in the frequency of CD3$^+$ T cells

(**Figure 3C**). Evaluation of CD150 expression by flow cytometry showed a trend toward higher

levels of CD150 on CD19$^+$ B cells compared to CD3$^+$ T cells. While this did not reach statistical

significance, this trend is consistent with the higher CD150 transcript counts observed in B cell

populations compared to T cell populations by scRNAseq (**Figure 3D; Supplementary Figure**

1E).

We next asked whether the preference for B cell infection was driven by a specific B cell

subset, or if all B cells were more susceptible to infection. To test this, we subset B cells based

on CD38 and CD27 expression (gating schemata shown in **Supplementary Figure 4A**). To

discriminate between susceptibility differences shared among all B cells and those that are

subtype-specific, we calculated the frequency of each B cell subset as a percentage of the total

B cell pool among both GFP$^+$ B cells from infected tonsil explants, as well as among bystander

and uninfected B cells. We found essentially no differences in infection based on CD38/CD27

expression (**Figure S4B-E**). The lack of differences in infection between subsets is also consistent

with the relatively stable expression of CD150 among these populations **(Figure S4F)**. Given the

dramatic increase in the IgD$^+$ B cell cluster observed among GFP$^+$ cells in the scRNA-seq **(Figure**

2F), we next assessed if IgD status conferred heightened susceptibility to infection among B cells

across time. We recapitulated the finding that IgD$^+$ cells are more frequent among the GFP$^+$

population than among bystander and uninfected cells (**Figure S4G**). However, when examined

among CD19$^+$ cells in each group, we found no evidence of preferential infection based on IgD

status (**Figure S4H-I**). Likewise, CD150 expression was not different between IgD$^+$ and IgD$^-$ cells

at day 6 (**Figure S4J**). These data show that while B cells are more susceptible to MeV infection,

this is most likely not driven by any individual subset of B cells.

We also evaluated the susceptibility of T cell subpopulations by examining the frequency

of CD4$^+$ and CD8$^+$ cells among GFP$^+$, bystander, or uninfected cell subsets. As shown in **Figure**

3E-G, we identified no differences in susceptibility between helper (CD4$^+$) and cytotoxic (CD8$^+$)

designations, consistent with the lack of differences in CD150 expression, which was generally

low, between CD4$^+$ and CD8$^+$ T cells (**Figure 3H**).

Reduced susceptibility of naive T cell subsets in human lymphoid tissue:

Previous reports have shown that mature (CD45RA$^-$) T cell subsets are more susceptible to MeV

infection than naive (CD45RA$^+$) T cells (34, 40, 41). To assess this in our model, we evaluated

the frequency of naïve T cells (CD45RA$^+$CCR7$^+$), as well as non-naïve T central memory (TCM;

CD45RA$^-$CCR7$^+$), T effector memory (TEM; CD45RA$^-$CCR7$^-$) and T effector memory RA$^+$

(TEMRA; CD45RA$^+$CCR7$^-$) subsets among CD4$^+$ and CD8$^+$ T cells from the GFP$^+$, GFP$^-$

(bystander) and uninfected groups. As shown through **Figure 3I-T**, we observed reduced MeV

infection in CD4$^+$ and CD8$^+$ naïve T cells compared to the non-naïve subsets, which were less

frequent in the GFP$^+$ population than in the bystander or uninfected groups, particularly at earlier

time points. Evaluation of the non-naïve subsets did not reveal an increased frequency among

GFP$^+$ cells, suggesting they were no more likely to become infected than their proportion in the

culture. Assessment of CD150 expression on these T cell subsets shows that naïve T cells

trended toward less CD150 expression than mature subsets, these differences were not

significant, suggesting that CD150 expression alone does not explain these trends in

susceptibility. Lastly, we hypothesized that the proximity of follicular CD4$^+$ T cells to highly

susceptible B cells in the follicle could impact the susceptibility of these CD4$^+$ cells to infection.

To test this, we evaluated the frequency of MeV-infected cells among CD4$^+$ CD45RA$^-$ T cells

based on CXCR5 expression. As shown in **Figure S5**, we found no difference in susceptibility

based on CXCR5 status. Taken together, these data indicate a reduced susceptibility of naïve T

cells to MeV that is largely independent of CD4 or CD8 status, CD150 expression, and CXCR5

expression.

MeV induces a canonical ISG response in both B and T cell transcriptomes:

Given that the most striking susceptibility differences to MeV infection were observed between B and T cells, we next asked whether the host response to infection among these cell types could contribute to these differences in susceptibility. To test this, we randomly sampled an equal number of B cells or T cells from the uninfected, bystander, and GFP$^+$ groups from the scRNA-Seq data and conducted differential gene expression analysis. As expected, the most significantly induced genes among the infected cells were MeV genes and GFP transcripts (**Figure 4A-B**). Following the viral genes, the most significantly upregulated transcripts among GFP$^+$ B and T cells were associated with a canonical interferon (IFN) signature. This pattern of interferon induction was strikingly similar among bystander cells, which were GFP$^-$ (thus not containing viral transcripts). We detected expression of the edited MeV interferon antagonist transcript, *V*, in GFP$^+$ cells but were unable to make meaningful comparisons in the expression of these transcripts among infected cell clusters due to the low read coverage at the p-editing site, where non-templated nucleotide insertion distinguishes *V* transcripts from the more abundant *P* mRNAs (**Supplementary Figure 6**). To directly compare the host response in infected B and T cells, we constructed a Venn-diagram of significantly induced genes in each group. As shown in **Figure 4C**, we find a highly conserved response between both cell types, among which *IFIT1, IFIT2, IFIT3, MX1, MX2, XAF1,* and other canonical interferon stimulated genes (ISGs) were shared. While the only gene found to be uniquely induced in infected B cells as compared to infected T cells was the MeV *P* gene, T cells were found to have induced additional ISGs that were not significant in B cells, including *OAS1, OAS2, OAS3, OASL, USP18, HELZ2, SAMD9L,* and *HERC6*. This unique pattern of gene expression may be biologically meaningful, or instead a consequence of strict statistical thresholding. Careful and directed comparative analyses using protein-based approaches will need to be conducted to confirm the relevance of these differences. To further validate this IFN signature, we conducted qRT-PCR from the tissues of two additional tonsil donors over time. We evaluated the expression of *IFIT1*, *IFIT3,* and *MX1*, three of the highly

expressed type-I interferon stimulated genes (ISGs) from the scRNA-Seq analysis. As shown in

Figure 4D-F, we found potent induction of all three genes by day 8 post-infection, the time point

of scRNA-Seq. Taken together, these data suggest MeV induces a potent interferon response at

the transcriptional level in both infected and bystander B and T cells, with no notable differences

that account for the increased susceptibility of B cells.

MeV induces an interferon-driven response in B cells at the protein level:

Since we identified a potent interferon signature in response to MeV at the transcriptional level,

we next asked whether the corresponding proteins were expressed. As B cells were preferential

targets for MeV, we utilized Raji cells where we could carefully define the protein level response

to infection in a uniform cellular population. Raji cells were infected with MeV at an MOI of 0.1 for

72 hours, and infection was confirmed through GFP expression (**Figure 5A**). Infected and

uninfected Raji cells were lysed, trypsin digested and analyzed by quantitative mass

spectrometry. Protein abundance was quantified in each sample relative to uninfected samples.

We next conducted differential expression analysis to quantify altered protein expression during

MeV infection as compared to the uninfected condition. As shown in **Figure 5B**, MeV-infected B

cells have higher expression of ISGs, consistent with our scRNA-seq data.

To functionally annotate the significantly dysregulated proteins, we next conducted a gene

ontology (GO) analysis. Significant GO terms are shown in **Figure 5C**, where the directionality of

the response is artificially shown based on a positive (upregulated) or negative (downregulated)

transformation of the adjusted p-value for that term. We found that the most significantly

upregulated pathways in MeV-infected cells were involved in antiviral signaling or interferon

biology (colored in red; **Figure 5C**). To further validate these results, we selected IFIT3 and

ISG15, two of the most highly upregulated proteins identified in the proteomics data to examine

by western blot. Like the GFP signal, we found that both IFIT3 and ISG15 were expressed only

in MeV-infected Raji cells (**Figure 5D**). Given that the transcriptomic analysis was conducted in

primary human lymphoid tissue, while the proteomic analysis was conducted in a B cell line, we

wanted to identify the significantly upregulated hits that were identified in both systems. To do

this, we compared the MeV transcriptional signature among the pseudobulked B cell cluster

(containing stochastic sampling of each of the B cell clusters) from lymphoid explants to the

significant proteins identified in the proteome of infected Raji cells. As shown in **Figure 5E**, the

most highly upregulated hits from this correlation analysis were ISGs, dominated by *MX2*, *ISG15*,

IFIT1, *IFIT2*, *IFIT3*, *CXCL10* and *XAF1*. To further characterize this host response, we also

compared the transcriptome and proteome of infected Raji cells. Raji cells were infected as in the

MS experiment but collected for bulk RNA sequencing. As shown in **Figure S7**, we identified a

conserved set of interferon stimulated genes that were robustly and significantly upregulated at

both the RNA and protein level. In addition, we noted that the expression of the FC-epsilon

receptor (*FCER2*) was significantly downregulated. Taken together, our integrated approach of

assessing the host response to infection at both the transcriptional and protein level reveals a

potently induced interferon signature that is conserved between distinct infection systems.

277
278

DISCUSSION

Measles virus pathogenesis is dependent upon early replication within the draining lymph node, yet our understanding of how infection proceeds in this organ and its link to disease outcomes is incomplete (48). In this study, we sought to model MeV infection in primary human lymphoid tissue explants and comprehensively characterize the immunological events that occur following MeV infection. scRNA-Seq analysis of infected cells was made possible by the high infection rates achieved in this system coupled with a GFP-expressing pathogenic strain of MeV. Our study also contributes an analysis of the MeV-induced proteome within infected B cells. Our unbiased approaches suggest that MeV has a remarkably wide lymphoid tropism, as we found MeV transcripts in most of the 29 cell types identified within the lymphoid tissue explants. Our data confirm previous findings of lymphocyte susceptibility and demonstrate a strong IFN signature associated with infection at both the RNA and protein levels.

Previous studies on MeV pathogenesis have identified a bias in infection towards B cells and away from T cells, with susceptibility differences driven by the expression of CD150 (18, 20, 27, 34, 40, 48, 49). Our data are consistent with this notion, as we found that B cells were the largest targets of infection, having heightened susceptibility and trending higher CD150 expression compared to T cells. Our analysis also extends these findings, revealing that while all B cells are highly susceptible to infection, accounting for the majority of infected cells, there were no observed differences in susceptibility based on B cell phenotype. Since CD150 expression was generally high among these B cell subsets, we interpret these findings to indicate that a baseline level of CD150 expression is sufficient to confer susceptibility, but differences beyond this threshold do not alter susceptibility. Of note, we found that germinal center B cells (GCBs) were by far the largest population of B cells found in this tissue, and thus, also comprised the greatest number of MeV-infected cells. These findings may suggest that immunological amnesia may extend beyond existing immunological to hamper future germinal center responses.

An interesting observation that we uncovered was that all cell types that we identified in

the tonsil cultures were found to have some level of MeV transcripts. Several cell types had

extremely low levels of MeV transcripts, including stromal and epithelial cells. While we took many

measures to ensure that these were bona fide infected cells (such as dead cell removal, flow

sorting for GFP, and a careful hashing strategy), we cannot eliminate the possibility that these

cells are a byproduct of exosomes or ambient viral RNA co-encapsulated with the GFP⁻

populations rather than truly infected cells. Our findings in larger populations, such as subsets of

CD4 and CD8 cells, were recapitulated with our flow cytometry approach, however, future work

should assess the possibility of MeV infection in rare tonsillar populations, such as stroma and

epithelium.

Many early studies on MeV pathogenesis focused on infection of T cells within secondary

lymphoid tissue (21, 34, 37, 43, 49, 50). Indeed, immunological amnesia was originally described

as a T cell phenotype, where children who had previously tested positive for a hypersensitivity

test to tuberculin antigen began to test negative following MeV infection (51). Subsequent work in

thymus and macaques revealed that MeV preferentially infects and depletes memory T cells over

naive T cell subsets, consistent with CD150 expression (32, 41). Our approach of assessing the

relative susceptibility of both T cells broadly, as well as within individual subsets both confirmed

and extended these findings. We found that antigen experience (CD45RA negativity) influenced

susceptibility to infection, while CXCR5 expression, used here as a proxy for localization within

the lymphoid explants, as well as CD150 expression, did not. Future work assessing the

susceptibility of these antigen-experienced subsets should focus on directly testing if factors other

than CD150 expression, such as spatial localization, promote susceptibility. Indeed, one

parameter that may be interesting to evaluate would be the extent to which directed cell migration

occurs within the tissue, and if infection influences immune cell trafficking.

Previous groups have shown that MeV does not induce a potent interferon (IFN) response,

as the viral V and C proteins can inhibit the induction of IFN (52-60). However, some groups have

observed the opposite, where MeV induces potent IFN expression (46, 61, 62). In general, this

discrepancy has been attributed to the presence of defective interfering (DI) RNAs, which can be

enriched as a byproduct of in vitro replication (52, 59, 63, 64). While these would not be captured

by our scRNA-Seq modality, we can conclude that the presence of viral V transcripts at day 8 was

not sufficient to shut down the IFN response, whether induced by DI RNAs or viral replication.

One hypothesis that might explain how MeV replicates in the presence of an IFN response would

be that the IFN response is induced to the benefit of MeV, not the detriment. The idea that viruses

may utilize IFN responses to promote infection has recently been demonstrated for influenza

virus, whereby the virus utilizes the host ISG *IFIT2* to enhance the translational efficiency of viral

RNAs (65). Alternatively, the addition of GFP into the viral genome may be indirectly involved as

placement of GFP in the first transcriptional unit of our MeV-GFP may decrease the relative

amounts of P-derived V and C proteins that antagonize type-I IFN responses. MeV-C is known to

reduce the production of DIs by enhancing the processivity of the viral polymerase (66) with C-

deficient MeV generating ~10-fold more DIs than the parental virus (64). Future studies should

assess the impact of this transcriptional shift on DI production as well as the downstream ability

to antagonize the type-I IFN response.

One major limitation of our study is that we do not know the impact of MeV infection

disease outcomes, such as immunological amnesia. Our results suggest that MeV infection of

GBCs may impact the germinal center responses, an outcome that would amplify the impact of

MeV on immunological amnesia. A second limitation of our study is that the transcriptomic and

proteomic analyses were conducted in entirely different systems due to the heterogeneous nature

of the lymphoid tissue explants. In the absence of single-cell proteomics, we limited our approach

to a correlative analysis between the two methodologies and systems. Therefore, we have high

confidence that these molecules are indeed a conserved response to MeV infection. A third

limitation in our analysis is that we are unable to differentiate between cell death by MeV versus

cell susceptibility to infection. While our data suggest that the broad susceptibility of T and B cells

is positively associated with the expression of the entry receptor CD150, this does not exclude

the possibility that some cell subpopulations have a greater capacity to survive while infected with

MeV. Indeed, it has previously been established that MeV is capable of depleting CD150$^+$ cells in

tonsil explants (43). However, given the high similarity in the frequency of bystander cells and

uninfected cells across all cell populations identified, we can presume that an enhanced frequency

among GFP$^+$ cells is indicative of enhanced susceptibility to infection. Understanding the impact

of MeV on cell death and proliferation may prove critical to understanding the complete pathology

of measles disease. Another limitation of our study is that we do not know the impact of the GFP

produced by the MeV-GFP on the induction of the innate immune response in tonsil tissues and/or

cell lines. The use of the GFP-expressing MeV enabled us to distinguish between infected and

bystander cells in our culture system. This was unavoidable to sort and conduct scRNAseq for

this study. However, a study utilizing this MeV-GFP showed that this strain is fully pathogenic in

macaques, suggesting that the introduction of GFP, and its possible ISG induction, does not affect

viral pathogenesis (67).

Our findings here represent a thorough analysis of the immunological events following

MeV infection of human lymphoid tissue explants. The finding that MeV has a broad tropism within

B cell, T cell, myeloid, and non-hematopoietic compartments may unlock new aspects of viral

pathogenesis in humans. While we do not know the role of each of these cell types in the collective

immunological response to infection, future studies should investigate how these cell types shape

the progression of measles disease. Our findings also represent a model system for the testing

of MeV antivirals, for which there are no current intervention strategies. One possibility would be

that by targeting specific aspects of the induced IFN response, MeV pathogenesis could be

ameliorated.

In toto, we present a thorough kinetic examination of the process of MeV infection in

human lymphoid explants, confirming previous groups' findings and broadening our

understanding of the key players in MeV infection within its natural target organ architecture.

Further, our integrated transcriptional and proteomic approach in this model establishes tonsil

explants as a unique platform for the identification of host factors important for MeV replication

and screening of targeted antivirals. Future work in this model should focus on understanding

how MeV replicates in the face of this potently induced interferon response to identify junctions at

which viral replication can be inhibited.

MATERIALS AND METHODS:

Sex as a biological variable

Our study received human tonsil tissue from both males and females. We did not observe any clear difference in MeV replication, so these data were analyzed together.

Cells and plasmids:

Vero-hCD150 cells were provided by Dr. Yanagi at Kyushu University and maintained in DMEM with 10% FBS (Biowest). RAJI-DCSIGNR cells were gifted by Ted Pierson (NIH/VRC, Bethesda USA) and cultured in RPMI with 10% FBS (68). The genome coding plasmid for MeV (p(+) MV323-AcGFP) was kindly gifted from Dr. Makoto Takeda (University of Tokyo, Tokyo Japan) (47). The MeV genome sequence was transferred into a pEMC vector, adding an optimal T7 promoter, a hammerhead ribozyme, and an eGFP transcriptional unit at the 3' end of the genome (pEMC-IC323-eGFP) as previously described (47).

MeV rescue and amplification:

MeV (IC323-eGFP) rescue was performed in BSR-T7 cells, seeded in 6-well format. Upon confluency, pEMC-IC323eGFP (5µg), T7-MeV-N (1.2µg), T7-MeV-P (1.2µg), T7-MeV-L (0.4µg), a plasmid encoding a codon-optimized T7 polymerase (3µg), PLUS reagent (5.8µL, Invitrogen), and Lipofectamine LTX (9.3µL, Invitrogen) were combined in Opti-MEM (200µl; Invitrogen). After a 30 min incubation at RT, the transfection mixture was added dropwise onto cells and incubated for 24hrs at 37°C. Following, rescued virus was amplified once on Vero-hCD150 cells for 72hrs to generate a P1 virus, in infection media (made in DMEM+2% FBS). This virus was then titered (see plaque assay method below) and used at an MOI = 0.01 on Vero-hCD150 cells to generate a P2 virus (amplified as above). Supernatants were collected, clarified of cell debris,

ultracentrifuged through a 20% sucrose gradient at 24,000 RPM for 3hrs, reconstituted in fresh

infection media, and frozen at -80°C.

MeV quantification by plaque assay:

Vero-hCD150 were plated in 12 well format until ~90-95% confluence. 10-fold dilutions of samples

(made in DMEM +2% FBS) were applied to these monolayers in a total volume of 250μL, and

infections were allowed to incubate for 2hrs at 37°C. Viral inoculum was replaced with 500μL/well

of methylcellulose (in DMEM + 2% FBS + 7.5% $NaHCO_3$). At 72hrs, wells were imaged for GFP$^+$

plaques on the Celigo S platform.

Processing and infection of human lymphoid tissue:

Human tonsils from routine tonsillectomies performed at the Mount Sinai Hospital and the New

York Eye and Ear Infirmary of Mount Sinai were collected under IRB-approved protocols within a

few hours after surgery. Tonsils were cut into 2mm^3 blocks, and 9 tissue blocks per well were

placed on top of collagen gelfoams (Cardinal Health) in a 6-well plate as previously described,

utilizing three wells per condition (a total of 27 blocks per experimental sample) (43). In all

experiments, triplicate wells were harvested as a single sample to reduce variability (44). After

overnight incubation, individual tissue blocks were individually inoculated with 5μl containing

1,666 PFU MeV-GFP (for a final concentration of 5,000 PFU/ml) or left uninfected. Media was

collected and replaced at days 3, 6, and 8 post-infection. Tonsil donors consisted of three male

and eight female donors. The median age of donors was 23 years old, with a range of 4 to 54

years old. The reasons for tonsillectomy included sleep apnea, breathing disorders, and chronic

tonsillitis.

Visualization of MeV-infected cells in tonsillar explants by in situ hybridization:

Tissues were fixed in 10% neutral buffered formalin and paraffin-embedded. In situ hybridization

using RNAscope® (ACDBio) was performed on 5µm sections to detect RNA encoding GFP.

Deparaffinization was performed by baking slides at 55°C for 20 min. Slides were washed twice

with xylene, twice in 100% ethanol, and were dried for 5 min at 60°C. Slides were then incubated

with hydrogen peroxide for 10 min at RT and were subsequently washed in diH2O. Slides were

placed in Target Retrieval solution at 100°C for 15 min, washed with water, and transferred into

100% ethanol for 3min, before drying. Sections were treated with RNAscope® Protease Plus and

fluorescence in situ hybridization was subsequently performed according to the manufacturer's

protocol (ACD# 323110) with RNAscope® Probe EGFP (ACD #400281; binds eGFP RNA) as

previously described (69). Slides were then mounted with Vectashield hard-set mounting medium

with DAPI (Vector Laboratories) and analyzed using an AxioImager Z2 microscope (Zeiss) and

Zen 2012 software (Zeiss).

Generating single cell suspensions from tonsil histocultures:

Single cell suspensions were generated by dissociating tissue (merged from the three technical

triplicate wells) using Collagenase IV (Worthington Biochemical) incubated for 30 min at 37°C

with gentle shaking as previously described (44). Samples were homogenized with mortar and

pestle before filtration over a 100µm cell filter and washed once with cold PBS before downstream

application.

Single cell RNA sequencing (scRNA-Seq):

Samples for scRNA-Seq were pooled for multiplex processing and analysis with a cell hashing

antibody strategy (70). Hash antibodies were generated by conjugating IDT synthesized oligos

(barcode sequences from 10x Genomics Chromium index SI-GA-F11; HBC21-29 for hash #1-8)

to antibodies utilizing Thunder-Link® PLUS oligo Antibody Conjugation Kit. Single cell

suspensions were generated from one donor-matched infected and uninfected culture at day 8

and dead cells were depleted from samples using the EasySep™ Dead Cell Removal (Annexin

V) Kit (Stemcell Technologies, #17899). Cells were blocked with Human TruStain FcX™

(Biolegend, #422302). Cells from the uninfected tonsil were split into 4 hashing groups (hash #1-

4), and cells from the infected tonsil were split into two infected hashing groups (hash #5, 6) and

two bystander hashing groups (hash #7, 8). Samples were stained with corresponding hashing

antibodies for 30 min at 4°C and washed 3x in FACS buffer (PBS + 1mM EDTA + 2% BSA). Cell

suspensions were filtered over a 70µm filter, stained with propidium iodide (PI) for viability, and

sorted as follows: live/GFP$^-$ cells from the uninfected condition, live/GFP$^+$ cells from the infected

condition, and live/GFP$^-$ cells from the bystander condition on a BD FACS Aria III. Sorted cells

were counted, and 10,000 uninfected cells, 5,000 infected cells, and 5,000 bystander cells were

pooled and processed for scRNA-Seq on the 10x Genomics Chromium platform, utilizing the 10x

3' v3 kit. A scRNA-Seq library was generated as per the manufacturer's protocol and sequenced

on an Illumina NextSeq500 instrument. A corresponding library of barcoded hash antibody

oligonucleotides was indexed with a standard Illumina D701 index and sequenced as above.

Processing of scRNA-Seq data:

Raw sequencing data output (BCL files) was converted to fastq files with CellRanger mkfastq

v3.0.2 (10X Genomics). Per cell gene count and hashtag antibody count matrices were generated

with CellRanger count v3.0.2 (10X Genomics), using a human genome reference (GRCh38,

Ensembl v96 transcript annotations) appended with the MeV-eGFP reference and corresponding

transcript annotations (MeV-IC323-eGFP, Genbank: MW401770). Data were read into the R

statistical framework (v4.0.3) for additional analysis with Seurat(71, 72) (v4.0.1). Hashtag

antibody data were center log ratio normalized by feature, and individual samples were

demultiplexed with the Seurat HTODemux function with the positive.quantile parameter set to
0.99.

Quality control and filtering of scRNA-Seq:

Data exploration and HTODemux classifications were used to set QC thresholds on per cell
transcript unique molecular identifiers (UMI) counts, detected gene counts, and the percentage
of detected mitochondrial transcripts. Cells with fewer than 2,500 transcript UMIs, fewer than 800
detected genes, and greater than 15% mitochondrial transcripts were excluded from downstream
analyses. After filtering, these data included 5,737 cells in the uninfected group, 2,736 cells in the
infected group, and 2,944 in the bystander group.

scRNA-Seq data analysis:

Datasets were normalized with SCTransform (73), with the per-cell mitochondrial transcript
percentage included as a regression variable. Data from all groups were integrated in Seurat
using 3,000 anchor features; MeV genes were excluded from all integration and clustering steps
to avoid group-specific artifacts. Dimensionality reduction was performed by principal component
analysis on integrated data, and the first 20 components were selected for graph-based clustering
by smart local moving algorithm (74) at a resolution of 1.4 (determined by Clustering Tree
assessment (75)).

General cell types were annotated by SingleR (76) from human immune cell reference
data (77). Clusters were assigned to one of each major cell group: T/NK, B, plasma, myeloid,
stromal, and epithelial. Those major cell groups with multiple component clusters were subset
and re-analyzed (normalization, principal components analysis dimensionality reduction, and
clustering at ClusterTree determined optimal resolution) for further annotation. For each major
cell group subset analysis, "marker genes" distinguishing component clusters were identified with
the FindAllMarkers (on the uninfected group) and/or FindConservedMarkers (on all groups)

functions. Intergroup differential gene expression analysis was performed with edgeR (78, 79)

(v3.32.1), including modifications of scRNA-Seq data (80). The edgeR linear model incorporated

factors for cellular gene detection rate (to account for scRNA-Seq "dropout") and experimental

group and included only those genes detected in at least 20% of cells in any contrast condition.

Statistical thresholds were set at Benjamini-Hochberg adjusted *p*-value less than 0.0001 and

absolute log fold-change greater than 1.58 for differential expression.

Immunophenotyping by flow cytometry:

Cells were stained with the Zombie Red fixable viability kit (Biolegend, #423109) for 10min at RT,

washed once with FACS buffer, and then blocked with Human TruStain FcX^{TM} (Biolegend,

#422302). Samples were incubated for 30min on ice with a cocktail of antibodies against: **(B Cell**

Panel) CD150 (Biolegend; Clone: A127d4; PE), CD38 (eBioscience; Clone: HB7; PE-Cy7), CD27

(Biolegend; Clone: O323; APC), CD45 (BD Horizon; Clone: HI30; BV605), CD19 (Biolegend;

Clone: HIB19; BV750) and CD3 (Biolegend; Clone: OKT3; BV785); **(T Cell Panel)** CD4

(eBioscience; Clone: OKT4; PerCP-Cy5.5), CD150 (Biolegend; Clone: A127d4; PE), CD45RA

(Biolegend; Clone: HI100; AlexaFluor-700), CXCR5 (Biolegend; Clone: J252D4; BV421), CD8

(Biolegend; Clone: RPA-T8; BV570), CD45 (BD Horizon; Clone: HI30; BV605), CD19 (Biolegend;

Clone: HIB19; BV750) and CD3 (Biolegend; Clone: OKT3; BV785). The antibody cocktail was

supplemented with Brilliant Stain buffer (BD Horizon^{TM,} #563794). All antibodies were used at a

concentration of 1µg/ml, except for CD27 (4µg/ml). Cells were washed 3 times with FACS buffer

before fixation with BD Cytofix^{TM} (BD Biosciences; #554655). Single color controls were

generated on UltraComp eBeads™ (Invitrogen™, #01-2222-42), except for GFP and Live/Dead

controls, which were generated using cells. All samples were analyzed on an Aurora Cytek®, and

unmixed samples were analyzed in FlowJo v10.8.1.

qRT-PCR:

Resuspended single tonsil cell suspensions were placed in 1ml of TriZol, and RNA was isolated

using Direct-zol RNA MiniPrep Plus kit (Zymo). 1µg of RNA was reverse transcribed with random

hexamer primers (Applied Biosystems). 1µl of cDNA was utilized per reaction, and primer/probes

for *IFIT1* (HS03027069_S1), *IFIT3* (HS01922752_S1), and *MX1* (Hs00895608_m) were utilized

to amplify ISG transcripts. Fold induction was calculated using the ΔΔCT method by normalizing

expression to *GAPDH* expression (NC_000012.11).

Sample preparation for Mass Spectrometry:

Uninfected or MeV-infected Raji-DCSIGNR cells were lysed in an 8M urea lysis buffer (with 100

mM ammonium bicarbonate, 150 mM NaCl, and 1x protease/phosphatase inhibitor cocktail HALT

(Thermo Fisher Scientific)). Lysates were sonicated, and protein concentrations were quantified

by micro-BCA assay (Thermo Fisher Scientific). 50µg of protein for each sample were treated

with Tris-(2-carboxyethyl)phosphine (TCEP) at a 4mM final concentration and incubated for 30

min at RT. Iodoacetamide (IAA) was added to a 10mM final concentration and samples were

incubated for 30 min at RT. Free IAA was quenched with the addition of Dithiothreitol (DTT) at a

10mM final concentration for 30min. Samples were diluted with 5 sample volumes of 100 mM

ammonium bicarbonate. Lysates were next digested with Trypsin Gold (Promega) at a 1:100

(enzyme: protein) ratio and lysates were rotated for 16hrs at RT. Trypsin activity was quenched

by adding 10% v/v trifluoroacetic acid (TFA) to a final concentration of 0.1%. Samples were

desalted on BioPure™ SPN MIDI C18 Spin columns. Samples were eluted from these columns

with 200µL 40% ACN/0.1% TFA, dried by vacuum centrifugation, and stored at -80°C.

Protein abundance Mass Spectrometry.

Samples were analyzed on an Orbitrap Eclipse mass spectrometry system (Thermo Fisher
Scientific) equipped with an Easy nLC 1200 ultra-high pressure liquid chromatography system
(Thermo Fisher Scientific) interfaced via a Nanospray Flex nanoelectrospray source. Immediately
before spectrometry, lyophilized samples were resuspended in 0.1% formic acid. Samples were
injected on a C18 reverse phase column (30 cm x 75 µm (ID)) packed with ReprosilPur 1.9 µm
particles). Mobile phase A consisted of 0.1% FA, and mobile phase B consisted of 0.1% FA/80%
ACN. Peptides were separated by an organic gradient from 5% to 35% mobile phase B over 120
min followed by an increase to 100% B over 10 min at a flow rate of 300 nL/min. Analytical
columns were equilibrated with 3 µL of mobile phase A. To build a spectral library, samples from
each set of biological replicates were pooled and acquired in a data-dependent manner. Data-
dependent analysis (DDA) was performed by acquiring a full scan over a m/z range of 375-1025
in the Orbitrap at 120,000 resolution(@200 m/z) with a normalized AGC target of 100%, an RF
lens setting of 30%, and an instrument-controlled ion injection time. Dynamic exclusion was set
to 30 seconds, with a 10ppm exclusion width setting. Peptides with charge states 2-6 were
selected for MS/MS interrogation using higher energy collisional dissociation (HCD) with a
normalized HCD collision energy of 28%, with three seconds of MS/MS scans per cycle. Data-
independent analysis (DIA) was performed on all individual samples. An MS scan was performed
at 60,000 resolution (@200m/z) over a scan range of 390-1010 m/z, an instrument-controlled
AGC target, an RF lens setting of 30%, and an instrument-controlled maximum injection time,
followed by DIA scans using 8 m/z isolation windows over 400-1000 m/z at a normalized HCD
collision energy of 28%.

MS Data analysis:

Peptides/Proteins were first identified with Spectronaut (81). False discovery rates (FDR) were

estimated using a decoy database strategy. All data were filtered to achieve an FDR of 0.01 for

peptide-spectrum matches, peptide identifications, and protein identifications. Search parameters

included a fixed modification for carbamidomethyl cysteine and variable modifications for N-

terminal protein acetylation and methionine oxidation. All other search parameters were defaults

for the respective algorithms. Analysis of protein expression utilized the MSstats statistical

package in R. Output data from Spectronaut was annotated based on a publicly available *Homo*

sapiens proteome (Proteome ID: UP000005640) and the reference sequence for IC323-eGFP

(Genbank: MW401770.1). Technical and biological replicates were integrated to estimate log2

fold changes, p values, and adjusted p values. All data were normalized by equalizing median

intensities, the summary method was Tukey's median polish, and the maximum quantile for

deciding censored missing values was 0.999. Significantly dysregulated proteins were defined as

those that had a fold change value >2 or <-2, with a p-value of <0.05. The mass indices for the

most significantly dysregulated proteins were transformed with the Quantile function in R and then

visualized using the pheatmap package.

Gene Ontology (GO) analysis

Gene Ontology (GO) analysis was conducted in R by quantifying the number of significantly

upregulated or downregulated proteins for each GO term identified from the gene ontology

resource, downloaded on 2.18.21. Significance was determined by a hypergeometric test. Only

terms with >2 proteins detected in the dataset were included in this analysis. This approach was

similarly taken to identify significantly downregulated terms. GO terms were graphed in GraphPad

PRISM based on their p-value, and terms related to the antiviral response were colored in red.

Western blot analysis of RAJI cell lysates.

Raji cells were infected with MeV at an MOI= 0.1 for 72 hours and compared to uninfected controls

(n=3). Cells were lysed as described above, and 10ug samples of whole cell lysate were mixed

1:1 with Lamelli Buffer (containing b-mercaptoethanol) and were heated at 95°C for 10min.

Samples were then electrophoresed on a 4-20% gradient SDS-PAGE gel (BioRad) and

transferred onto a methanol-activated polyvinylidene difluoride (PVDF) membrane (Bio-Rad). The

membrane was blocked with 5% Milk in PBST (0.1% Tween-20) for 1hr. The following antibody

staining protocols were run sequentially: 1) anti-IFIT3 (ThermoFisher; Clone: OTI1G1; 1:1000)

developed with goat-anti-mouse IgG-HRP (Catalog: G-21040; 1:10,000); 2) anti-ISG15 (Clone:

7H29L24; 1:5000) developed with goat-anti-rabbit IgG-HRP antibody (Thermofisher; Catalog: 65-

6120; 1:10,000); and 3) anti-beta Actin (Thermofisher; Clone: 15G5A11/E2; 1:1000) and anti-GFP

(Thermofisher; Clone: GF28R; 1:1000) simultaneously, developed with anti-mouse Alexa 647

antibody (Thermofisher; clone A-21235; 1:2000). HRP signals were detected between each

incubation with SuperSignal West Pico™ PLUS reagent (Thermofisher; 1:1 luminol/enhancer),

and images were acquired on a Chemidoc™ MP. Western images were merged for presentation

in FIJI.

Correlation Analysis of RNA and protein response to infection

Data from scRNAseq and mass spectrometry were further processed in RStudio to correlate RNA

and protein levels. All B cells from the scRNAseq dataset were rebulked, and total read counts in

this new "B cell" cluster were calculated, normalized to counts per kb million, and log2

transformed. To evaluate the fold change (FC) between mock-treated and MeV-infected samples,

the log2 values from the mock condition were subtracted from the MeV condition. A correlation

scatter plot was created using ggplot, and protein labels were added only if the log2FC values

were greater than 2 in both RNA and protein.

Bulk-RNA sequencing of infected RAJI cells

RAJI-DCSIGNR cells were infected with MeV as during the preparation of MS samples above.

Cells were pelleted and resuspended in 500uL of Trizol. RNA was extracted using the Direct-zol

RNA miniprep kit (Zymo Research), and frozen RNA was shipped to Azenta Life Sciences for

library preparation and sequencing. ERCC RNA Spike-in Max kit was added to normalize total

RNA prior to library preparation following the manufacturer's protocol (cat # 4456740). RNA

sequencing libraries were prepared using the NEBNext Ultra II RNA Library Prep Kit for Illumina.

Libraries were quantified with Agilent TapeStation, Qubit 2.0 and by quantitative PCR prior to

sequencing on an Illumina NovaSeq XPlus 25B. Samples were sequenced using a standard

2x150bp paired-end configuration. Raw sequence data was converted into fastq files and de-

multiplex using Illumina's bcl2fastq 2.2.0 software. The quality of sequencing was assessed with

FastQC, and sequencing reads were aligned to indexed reference genomes using the STAR

aligner. Expression matrices were calculated using featureCounts, and data were exported into

R for data analysis and visualization. Transcripts where fewer than 10 transcripts were collected

across all samples were excluded from further analysis. Data normalization and differential gene

expression analysis was conducted using the DESEQ2 package (version 1.42.1).

Statistics:

For scRNA-Seq and mass spectrometry analysis, statistical analysis methodology is detailed in

the above methods section. For all comparisons of infection susceptibility over time, significance

was determined by two-way ANOVA using the Geisser-Greenhouse correction with Tukey's

multiple comparison test. For all comparisons of CD150 expression among multiple (>2) groups,

significance was determined by one-way ANOVA using Friedman's test with Dunnett's multiple

comparison test. For pairwise comparisons, a non-parametric 2-tailed paired T-test was utilized

(Wilcoxon matched-pairs signed rank test). For experiments with three replicates, the median with

the 95% confidence intervals is shown instead of p-value. For all other statistical analyses, *

indicates $p<0.05$, ** indicates $p<0.01$, *** indicates $p<0.001$ and **** indicates $p<0.0001$

Study Approval:

All tonsil tissues were obtained with informed consent under IRB# 16-01425-CR002 at the New

York Eye and Ear Infirmary of Mount Sinai, or the Mount Sinai Hospital under IRB HS#12-0045.

Data Availability:

Large data sets will be made available on NCBI GEO and SRA. BulkSeq (Accession

#GSE272426); scRNASeq (Accession #GSE272481). Proteomics data are available on PRIDE.

Raw data values for data shown in this manuscript can be accessed in the "Supporting data

values" XLS file. Information about human subjects is limited by IRB, however anonymized

information will be made available upon request to the corresponding author where possible.

AUTHOR CONTRIBUTIONS: JAA, BRR, BL and JKL conceptualized the project. JA, ARP, SH,

APK, PT, SI, JC, HU, BT, JRJ, BRR, BL, and JKL contributed to the work intellectually. JAA, ARP,

SH, ASM, PT, and NI conducted the experiments. SI generated the virus used in this study. BT

conducted tonsillectomies included in this work. JAA, ARP, SH, EJD, and BRR conducted data

analysis. JAA, ARP, and JKL prepared the manuscript. All authors edited the manuscript. The

order of the co-first authors was determined based on contributions to the conceptualization and

execution of this work.

ACKNOWLEDGMENTS:

Research was supported by NIH applications R21AI149033 (JKL, BRR, and BL), R01AI071002

(BL), and CDMRP PR192188 (JKL and BL). JAA was supported by an NIH fellowship

F31HL149295. ARP and EJD were supported in part by the NIH training grant T32AI007647-23.

P.A.T. was funded by CIHR-IRSC: 0041001056. The authors thank Dr. Rachel Brody and

colleagues at the ISMMS Biorepository and Pathology CoRE, the ISMMS Human Immune

Monitoring Core (HIMC), and Dr. Emilia Bagiella, for her expertise and guidance on the statistical

methodology.

REFERENCES

1. Olsen LF, et al. Oscillations and chaos in epidemics: A nonlinear dynamic study of six childhood diseases in Copenhagen, Denmark. *Theoretical Population Biology.* 1988;33(3):344-70.

2. Merler S, and Ajelli M. Deciphering the relative weights of demographic transition and vaccination in the decrease of measles incidence in Italy. *Proc Biol Sci.* 2014;281(1777):20132676.

3. Edmunds WJ, et al. The pre-vaccination epidemiology of measles, mumps and rubella in Europe: implications for modelling studies. *Epidemiol Infect.* 2000;125(3):635-50.

4. Metcalf CJ, et al. Seasonality and comparative dynamics of six childhood infections in pre-vaccination Copenhagen. *Proc Biol Sci.* 2009;276(1676):4111-8.

5. Wallinga J, et al. Estimation of measles reproduction ratios and prospects for elimination of measles by vaccination in some Western European countries. *Epidemiol Infect.* 2001;127(2):281-95.

6. Mossong J, and Muller CP. Estimation of the basic reproduction number of measles during an outbreak in a partially vaccinated population. *Epidemiol Infect.* 2000;124(2):273-8.

7. Guerra FM, et al. The basic reproduction number (R(0)) of measles: a systematic review. *Lancet Infect Dis.* 2017;17(12):e420-e8.

8. Chovatiya R, and Silverberg JI. Inpatient morbidity and mortality of measles in the United States. *PLoS One.* 2020;15(4):e0231329.

9. Tanne JH. Measles cases and deaths are increasing worldwide, warn health agencies. *Bmj.* 2020;371:m4450.

10.   López-Perea N, et al. Measles in Vaccinated People: Epidemiology and

Challenges in Surveillance and Diagnosis in the Post-Elimination Phase. Spain,

2014-2020. *Viruses.* 2021;13(10).

11.   Hussain A, et al. The Anti-vaccination Movement: A Regression in Modern

Medicine. *Cureus.* 2018;10(7):e2919.

12.   Gallegos M, et al. Anti-vax: the history of a scientific problem. *J Public Health (Oxf).*

2022.

13.   Kuylen E, et al. Clustering of susceptible individuals within households can drive

measles outbreaks: an individual-based model exploration. *Sci Rep.*

2020;10(1):19645.

14.   Zucker JR, et al. Consequences of Undervaccination - Measles Outbreak, New

York City, 2018-2019. *N Engl J Med.* 2020;382(11):1009-17.

15.   Gaythorpe KA, et al. Impact of COVID-19-related disruptions to measles,

meningococcal A, and yellow fever vaccination in 10 countries. *Elife.* 2021;10.

16.   Rota PA, et al. Measles. *Nat Rev Dis Primers.* 2016;2:16049.

17.   Plemper RK, and Snyder JP. Measles control--can measles virus inhibitors make

a difference? *Curr Opin Investig Drugs.* 2009;10(8):811-20.

18.   Lemon K, et al. Early target cells of measles virus after aerosol infection of non-

human primates. *PLoS Pathog.* 2011;7(1):e1001263.

19.   Ludlow M, et al. Measles virus infection of epithelial cells in the macaque upper

respiratory tract is mediated by subepithelial immune cells. *J Virol.*

2013;87(7):4033-42.

20.   Welstead GG, et al. Measles virus replication in lymphatic cells and organs of

737   CD150 (SLAM) transgenic mice. *Proc Natl Acad Sci U S A.* 2005;102(45):16415-

738   20.

739   21.   Dhiman N, et al. Measles virus receptors: SLAM and CD46. *Rev Med Virol.*

740   2004;14(4):217-29.

741   22.   Ono N, et al. Measles viruses on throat swabs from measles patients use signaling

742   lymphocytic activation molecule (CDw150) but not CD46 as a cellular receptor. *J*

743   *Virol.* 2001;75(9):4399-401.

744   23.   Takeda M, et al. Wild-Type Measles Virus is Intrinsically Dual-Tropic. *Front*

745   *Microbiol.* 2011;2:279.

746   24.   Lin LT, and Richardson CD. The Host Cell Receptors for Measles Virus and Their

747   Interaction with the Viral Hemagglutinin (H) Protein. *Viruses.* 2016;8(9).

748   25.   Noyce RS, et al. Tumor cell marker PVRL4 (nectin 4) is an epithelial cell receptor

749   for measles virus. *PLoS Pathog.* 2011;7(8):e1002240.

750   26.   Mühlebach MD, et al. Adherens junction protein nectin-4 is the epithelial receptor

751   for measles virus. *Nature.* 2011;480(7378):530-3.

752   27.   Laksono BM, et al. Measles Virus Host Invasion and Pathogenesis. *Viruses.*

753   2016;8(8).

754   28.   Ashbaugh HR, et al. Association of Previous Measles Infection With Markers of

755   Acute Infectious Disease Among 9- to 59-Month-Old Children in the Democratic

756   Republic of the Congo. *J Pediatric Infect Dis Soc.* 2019;8(6):531-8.

757   29.   Le NTH, et al. Biphasic pattern in the effect of severe measles infection; the

758   difference between additive and multiplicative scale. *BMC Infect Dis.*

759   2021;21(1):1249.

760   30.   Sato R, and Haraguchi M. Effect of measles prevalence and vaccination coverage

761   on other disease burden: evidence of measles immune amnesia in 46 African

762   countries. *Hum Vaccin Immunother.* 2021;17(12):5361-6.

763   31.   Morales GB, and Muñoz MA. Immune amnesia induced by measles and its effects

764   on concurrent epidemics. *J R Soc Interface.* 2021;18(179):20210153.

765   32.   de Swart RL. [The measles paradox explained: how measles virus both

766   suppresses and activates the immune system]. *Ned Tijdschr Geneeskd.* 2020;164.

767   33.   Galassi FM, et al. Measles-induced immune amnesia likely recorded in the 18th

768   century. *J Clin Virol.* 2021;141:104899.

769   34.   Laksono BM, et al. Studies into the mechanism of measles-associated immune

770   suppression during a measles outbreak in the Netherlands. *Nat Commun.*

771   2018;9(1):4944.

772   35.   Mina MJ, et al. Long-term measles-induced immunomodulation increases overall

773   childhood infectious disease mortality. *Science.* 2015;348(6235):694-9.

774   36.   Mina MJ, et al. Measles virus infection diminishes preexisting antibodies that offer

775   protection from other pathogens. *Science.* 2019;366(6465):599-606.

776   37.   Valdimarsson H, et al. Measles virus receptor on human T lymphocytes. *Nature.*

777   1975;255(5509):554-6.

778   38.   VON PIRQUET C. Verhalten der kutanen tuberkulin-reaktion wahrend der Masern.

779   *Deutsch Med Wochenschr.* 1908;34: 1297-1300.

39. Laksono BM, et al. In Vitro Measles Virus Infection of Human Lymphocyte Subsets

Demonstrates High Susceptibility and Permissiveness of both Naive and Memory

B Cells. *J Virol.* 2018;92(8).

40. de Swart RL, et al. Predominant infection of CD150+ lymphocytes and dendritic

cells during measles virus infection of macaques. *PLoS Pathog.* 2007;3(11):e178.

41. de Vries RD, et al. Measles immune suppression: lessons from the macaque

model. *PLoS Pathog.* 2012;8(8):e1002885.

42. Victora GD, and Nussenzweig MC. Germinal Centers. *Annu Rev Immunol.*

2022;40:413-42.

43. Condack C, et al. Measles virus vaccine attenuation: suboptimal infection of

lymphatic tissue and tropism alteration. *J Infect Dis.* 2007;196(4):541-9.

44. Grivel JC, and Margolis L. Use of human tissue explants to study human infectious

agents. *Nat Protoc.* 2009;4(2):256-69.

45. Takeda M, et al. Recovery of pathogenic measles virus from cloned cDNA. *J Virol.*

2000;74(14):6643-7.

46. Hippee CE, et al. Measles virus exits human airway epithelia within dislodged

metabolically active infectious centers. *PLoS Pathog.* 2021;17(8):e1009458.

47. Ikegame S, et al. Fitness selection of hyperfusogenic measles virus F proteins

associated with neuropathogenic phenotypes. *Proc Natl Acad Sci U S A.*

2021;118(18).

48. Griffin DE. Measles immunity and immunosuppression. *Curr Opin Virol.* 2021;46:9-

14.

49. Okamoto Y, et al. Immature CD4+CD8+ thymocytes are preferentially infected by

measles virus in human thymic organ cultures. *PLoS One.* 2012;7(9):e45999.

50. Welstead GG, et al. Mechanism of CD150 (SLAM) down regulation from the host

cell surface by measles virus hemagglutinin protein. *J Virol.* 2004;78(18):9666-74.

51. VON PIRQUET C. FREQUENCY OF TUBERCULOSIS IN CHILDHOOD. *Journal*

of the American Medical Association. 1909;LII(9):675-8.

52. Nakatsu Y, et al. Measles virus circumvents the host interferon response by

different actions of the C and V proteins. *J Virol.* 2008;82(17):8296-306.

53. Naniche D, et al. Evasion of host defenses by measles virus: wild-type measles

virus infection interferes with induction of Alpha/Beta interferon production. *J Virol.*

2000;74(16):7478-84.

54. Shivakoti R, et al. Limited in vivo production of type I or type III interferon after

infection of macaques with vaccine or wild-type strains of measles virus. *J*

Interferon Cytokine Res. 2015;35(4):292-301.

55. Rabaan AA, et al. Updates on Measles Incidence and Eradication: Emphasis on

the Immunological Aspects of Measles Infection. *Medicina (Kaunas).* 2022;58(5).

56. Fontana JM, et al. Regulation of interferon signaling by the C and V proteins from

attenuated and wild-type strains of measles virus. *Virology.* 2008;374(1):71-81.

57. Komune N, et al. Measles virus V protein inhibits NLRP3 inflammasome-mediated

interleukin-1β secretion. *J Virol.* 2011;85(24):13019-26.

58. Sparrer KM, et al. Measles virus C protein interferes with Beta interferon

transcription in the nucleus. *J Virol.* 2012;86(2):796-805.

59.   Runge S, et al. In vivo ligands of MDA5 and RIG-I in measles virus-infected cells.

PLoS Pathog. 2014;10(4):e1004081.

60.   Shaffer JA, et al. The C protein of measles virus inhibits the type I interferon

response. *Virology.* 2003;315(2):389-97.

61.   Herschke F, et al. Cell-cell fusion induced by measles virus amplifies the type I

interferon response. *J Virol.* 2007;81(23):12859-71.

62.   Plumet S, et al. Cytosolic 5'-triphosphate ended viral leader transcript of measles

virus as activator of the RIG I-mediated interferon response. *PLoS One.*

2007;2(3):e279.

63.   Calain P, et al. Molecular cloning of natural paramyxovirus copy-back defective

interfering RNAs and their expression from DNA. *Virology.* 1992;191(1):62-71.

64.   Pfaller CK, et al. Measles Virus Defective Interfering RNAs Are Generated

Frequently and Early in the Absence of C Protein and Can Be Destabilized by

Adenosine Deaminase Acting on RNA-1-Like Hypermutations. *J Virol.*

2015;89(15):7735-47.

65.   Tran V, et al. Influenza virus repurposes the antiviral protein IFIT2 to promote

translation of viral mRNAs. *Nat Microbiol.* 2020;5(12):1490-503.

66.   Siering O, et al. C Proteins: Controllers of Orderly Paramyxovirus Replication and

of the Innate Immune Response. *Viruses.* 2022;14(1).

67.   de Vries RD, et al. In vivo tropism of attenuated and pathogenic measles virus

expressing green fluorescent protein in macaques. *J Virol.* 2010;84(9):4714-24.

68.   Davis CW, et al. West Nile virus discriminates between DC-SIGN and DC-SIGNR

for cellular attachment and infection. *J Virol.* 2006;80(3):1290-301.

69.   Cao P, et al. Fluorescence in situ hybridization is superior for monitoring Epstein

848   Barr viral load in infectious mononucleosis patients. *BMC Infectious Diseases.*

849   2017;17(1):323.

850   70.   Stoeckius M, et al. Cell Hashing with barcoded antibodies enables multiplexing

851   and doublet detection for single cell genomics. *Genome Biol.* 2018;19(1):224.

852   71.   Stuart T, et al. Comprehensive Integration of Single-Cell Data. *Cell.*

853   2019;177(7):1888-902.e21.

854   72.   Butler A, et al. Integrating single-cell transcriptomic data across different

855   conditions, technologies, and species. *Nat Biotechnol.* 2018;36(5):411-20.

856   73.   Hafemeister C, and Satija R. Normalization and variance stabilization of single-cell

857   RNA-seq data using regularized negative binomial regression. *Genome Biol.*

858   2019;20(1):296.

859   74.   Waltman L, and van Eck NJ. A smart local moving algorithm for large-scale

860   modularity-based community detection. *The European Physical Journal B.*

861   2013;86(11):471.

862   75.   Zappia L, and Oshlack A. Clustering trees: a visualization for evaluating clusterings

863   at multiple resolutions. *Gigascience.* 2018;7(7).

864   76.   Aran D, et al. Reference-based analysis of lung single-cell sequencing reveals a

865   transitional profibrotic macrophage. *Nat Immunol.* 2019;20(2):163-72.

866   77.   Monaco G, et al. RNA-Seq Signatures Normalized by mRNA Abundance Allow

867   Absolute Deconvolution of Human Immune Cell Types. *Cell Rep.*

868   2019;26(6):1627-40.e7.

869   78.   McCarthy DJ, et al. Differential expression analysis of multifactor RNA-Seq

870   experiments with respect to biological variation. *Nucleic Acids Res.*

871   2012;40(10):4288-97.

872   79.   Robinson MD, et al. edgeR: a Bioconductor package for differential expression

873   analysis of digital gene expression data. *Bioinformatics.* 2010;26(1):139-40.

874   80.   Soneson C, and Robinson MD. Bias, robustness and scalability in single-cell

875   differential expression analysis. *Nat Methods.* 2018;15(4):255-61.

876   81.   Bruderer R, et al. Extending the limits of quantitative proteome profiling with data-

877   independent acquisition and application to acetaminophen-treated three-

878   dimensional liver microtissues. *Mol Cell Proteomics.* 2015;14(5):1400-10.

881
Figure 1: MeV productively infects human lymphoid tissue explants. Tonsil tissues (n=11)
were infected with MeV-GFP. Cumulative viral plaque forming units (PFU/ml) was quantified from
supernatants collected on days 3, 6, and 8 post-infection by plaque assay **(A)**. Representative
flow plots quantifying infection (GFP) are shown for one donor **(B)** and quantified across 4 donors
over time **(C)**. In situ hybridization for GFP RNA (green) on paraffin-embedded tissues collected
on day 8 post-infection compared to a donor-matched, uninfected control **(D)**. Nuclei were
counterstained with DAPI (blue). Scale bars represent 50µm. Significance was determined by
one-way ANOVA using Friedman's test with Dunnett's multiple comparison test. *indicates
$p<0.05$, ** indicates $p<0.01$, *** indicates $p<0.001$ and **** indicates $p<0.0001$.
891
892

893
Figure 2: scRNA-Seq identifies 29 unique cell populations in tonsils susceptible to MeV.
Tonsil tissue from MeV-GFP-infected and uninfected explants on day 8 from one donor were
sorted for scRNA-Seq. Schemata of scRNA-Seq workflow are shown in **(A)**. Cells from the
infected condition were sorted and hash-labeled into GFP$^+$ and GFP$^-$ groups. Uninfected GFP$^-$
cells were sorted from a donor-matched uninfected control. 5,000 GFP$^+$ cells, 5,000 GFP$^-$ cells,
and 10,000 uninfected cells were encapsulated for sequencing. MeV RNA unique molecular
identifiers (UMIs) were quantified for quality control and filtering **(B)**. Normalized expression of
each MeV transcript in infected cells was quantified and shown as violin plots ordered from 3'to
5'in the MeV genome **(C)**. Canonical correlation analysis (CCA) was conducted on all groups
(combined) and individual clusters were functionally annotated (see also **Figure S1**). Clusters
were visualized by UMAP **(D)** and then split into conditions based on captured hashing
oligonucleotide sequences for further analysis **(E)**. The frequencies of each cell cluster identified
in **(E)** were calculated for each group, and quantification is shown in **(F)**. The percentage of the
transcriptome that is MeV RNA is shown in **(G)**, with a +1 pseudocount artificially added to the
values for display on a $\log_{10}$ axis. All cluster annotations are labeled by the color legend shown.

Figure 3: MeV preferentially infects B cells and is restricted among naïve T cells. Cells from
MeV-GFP-infected and donor-matched uninfected tissues were collected at days 3, 6, and 8 post-
infection, and were then immunophenotyped by flow cytometry (n=3; gating schemata in **A, E, I,**
and **O**). The frequency of CD19+ B cells **(B)** and CD3+ T cells **(C)** among all CD45+ cells are
quantified and compared over time for GFP+ cells, GFP- bystander cells, and donor-matched
uninfected controls. The mean fluorescent intensity (MFI) of surface CD150 among CD19+ and
CD3+ cells is compared **(D)**. Susceptibility to infection among CD4 **(F)** and CD8 **(G)** populations
is shown, with CD150 expression compared between populations **(H)**. Naive (CD45RA+/CCR7+),
TCM (CD45+/CCR7-), TEM (CD45RA-/CCR7-), and TEMRA (CD45RA+/CCR7-) populations are
quantified and compared among CD4+ cells **(I-M)** and CD8+ cells **(O-S)**. CD150 expression is
compared among CD4+ **(N)** and CD8+ **(T)** subpopulations. For all immunophenotyping panels,
significance was determined by two-way ANOVA using the Geisser-Greenhouse correction with
Tukey's multiple comparison test. For panels **(D)** and **(H)**, significance was determined by the
Wilcoxon matched-pairs signed rank test. For panels **(N)** and **(T)**, Significance was determined
by one-way ANOVA using Friedman's test with Dunnett's multiple comparison test. For all plots,
the median with the 95% confidence interval are shown.

926
Figure 4: Host response to MeV in lymphoid tissue is dominated by a type I interferon
response. All B and T cell clusters from the scRNA-Seq analysis were combined with stochastic
downsampling. Differential gene expression analysis was conducted with EdgeR, and expression
of the most significant genes is shown for B cells **(A)** and T cells **(B)**. Statistical thresholds for
significant differential gene expression were set at Benjamini-Hochberg adjusted *p* value <0.0001
and absolute LogFC>1.58. Genes that were significant in either the infected:uninfected or the
bystander:uninfected comparison were plotted. Genes significantly induced during infection were
compared between B and T cells **(C)**. MeV-infected tonsil explants were collected for RNA
extraction and analysis by qRT-PCR. RNA was examined for the expression of *IFIT1* **(D)**, *IFIT3*
(E), and *MX1* **(F)**. The fold change in expression levels relative to uninfected controls is shown,
with the expression of each ISG normalized to the expression of *GAPDH* (ΔΔCT method).
938

939
Figure 5: Proteins involved in a type I Interferon response are potently upregulated in
response to MeV. Raji-DCSIGNR cells were infected with MeV-GFP (MOI=0.1) or left uninfected
for 72 hours before processing for protein abundance mass spectrometry (n=4). Infection was
confirmed by quantifying GFP expression by flow cytometry **(A)**. The most significantly
dysregulated proteins (columns) for each sample (rows) were visualized, with clustering based
on protein expression **(B)**. GO analysis was conducted, and the most significant functional terms
were visualized. Terms were given a positive value if the term was upregulated during infection
or a negative value if downregulated. Red bars indicate involvement in antiviral responses **(C)**.
Validation of IFIT3 and ISG15 upregulation in infected Raji cells was conducted by western blot
(D). Correlation of the transcriptome (pseudobulked B cell supercluster from Figure 4) and
proteome (Raji cells, Figure 5) was conducted. Values that had a log2FC>2 in both the proteome
and transcriptome were labeled **(E)**. For the box and whisker plot, the mean with SEM is shown.

952
Figure S1 (related to Figure 2): Cluster annotation strategy for scRNA-Seq. Total counts of
unique molecular identifiers (UMI; **A**), individual genes **(B)**, and mitochondrial transcripts **(C)** were
quantified for quality control and filtering. Dot plot showing the expression of cluster-defining
features across each identified cell cluster. Average expression (Z scaled) is shown for each
feature by color, while the percentage of cells in each cluster expressing that feature is shown by
the size of the dot **(D)**. Normalized SLAMF1 expression across all cell types was determined **(E)**.
Pathway scores for S phase **(F)** and G2M **(G)** were calculated and compared across B cell
subsets and proliferating CD4 T cells.

961
962

Figure S2 (Related to Figure 3): Quantification of infection kinetics in different cell populations. Secondary analysis of donors from Figure 3. The percentage of GFP$^+$ cells within each cell type identified in the explants is quantified for B and T cells **(A-B)**, CD4$^+$ and CD8$^+$ cells **(C-D)**, B cell subsets **(E-H)**, CD4$^+$ T cell subsets **(I-L)** and CD8$^+$ T cell subsets **(M-P)**. Each dot/line represents a single donor.

 Figure S3 (Related to Figure 3). Gating schemata for flow cytometry-based
immunophenotyping experiments. Representative flow plots are shown for one uninfected
donor. Single cells are selected, followed by dead cell exclusion, gating upon CD45[+] cells, cell
gating, and then binning cells based on GFP status **(A)**. Within GFP[+] or GFP- gates, cells are
further phenotyped based on CD3/CD19 (B vs T cells), CD4 vs CD8 lineages within the CD3[+]
gate, and then memory subsets within these lineages. Histograms demonstrating CD150
expression are compared between CD3[+] and CD19[+] cells **(B)**, between CD4[+] and CD8[+] cells **(C)**,
or between CD4[+] **(D)** or CD8[+] **(E)** memory subsets for one representative donor in the day 6
uninfected condition.

Figure S4 (related to Figure 3): MeV infects B cell subsets proportionally. Donors (n=3) from Figure 3 were immunophenotyped to identify B cell subsets using CD38 and CD27 within the CD19$^+$ cells, with the gating strategy utilized shown in **(A)**. Naive (CD27$^-$CD38$^-$; **B**), germinal center (CD27$^-$CD38$^-$; **C**), memory (CD27$^+$CD38$^-$; **D**), and activated memory (CD27$^+$CD38$^+$; **E**) B cells were quantified by comparing their frequency among both uninfected and GFP$^+$ cells over time. CD150 expression was calculated for each population among uninfected cells at 6 days post-infection and the mean fluorescent intensity (MFI) of CD150 expression is shown **(F)**. Representative flow plots are shown demonstrating the enrichment of IgD$^+$ B cells among all infected cells at 8DPI **(G)**. Susceptibility to infection was assessed by quantifying the frequency of IgD- **(H)** or IgD$^+$ **(I)** cells among uninfected, bystander, or infected cells. CD150 expression on IGD- and IgD$^+$ populations at day 6 are shown in **(J)**. For all immunophenotyping panels, significance was determined by two-way ANOVA using the Geisser-Greenhouse correction with Tukey's multiple comparison test. For panel **(F)** significance was determined by one-way ANOVA using Friedman's test with Dunnett's multiple comparison test. Significance in **(J)** was determined using the Wilcoxon matched-pairs signed rank test. For all plots, the median with the 95% confidence interval is shown.

1000
Figure S5 (related to Figure 3): CXCR5 status has no impact on the susceptibility of CD4^{+}
cells. Non-Naïve CD4^{+} cells (n=3) were subset based on CXCR5 status as shown in (**A**). Shown
are the frequency of non-follicular (CXCR5^{-}; **B**) and follicular (CXCR5^{+}; **C**) cells among uninfected,
GFP^{-} (bystander), or GFP^{+} non-naïve CD4^{+} cells. CD150 expression was compared between non-
follicular and follicular cells (**D**). Significance for **B-C** was determined by two-way ANOVA using
the Geisser-Greenhouse correction with Tukey's multiple comparison test, and CD150 expression
significance was determined by the Wilcoxon matched-pairs signed rank test (**D**). For all plots,
the median with the 95% confidence interval are shown.
1009

A.

B.

1010
Figure S6 (related to Figures 2 and 4): Detection of P-editing by scRNA-Seq. (A) Read
coverage for all libraries along the length of the MeV P transcript. The red box denotes the
region of the MeV linear genome that is shown in the histogram box, and the boxes under the
histogram are a selection of reads that map to these positions. Gray bars indicate that the read
matches the reference sequence, with colored letters representing mismatches to the reference.
(B) Zoomed-in coverage of the p-edited region, showing representative reads that map to this
region of the gene in gray. "I" represents indel mappings, which may indicate P-edited
transcripts. There were 235 total reads covering the putative edit site.

Figure S7 (Related to Figure 5). Comparison of the transcriptome and proteome in infected Raji cells. Raji cells were infected with MeV and collected for bulk RNA sequencing (n=3). **(A)** Detection of MeV transcripts in infected cells is shown as normalized counts for each MeV gene compared to uninfected controls. Median and 95% confidence intervals are shown. **(B)** Correlation plot showing the relationship between differentially expressed proteins (y-axis) identified by MS in Figure 5 with differentially expressed transcripts identified by bulk RNA sequencing (x-axis). Log2FC values for transcripts and proteins that were detected in both RNA and protein assays are shown. Simple linear regression was conducted, and the Pearson correlation value was reported on the plot, along with the significance of the correlation. **(C)** Correlation plot demonstrating the relationship between the list of significantly altered proteins (y-axis) and transcripts (x-axis). Hits that were significant in both assays are denoted in the table shown in **(D)**.