Alternatively Spliced Landscape of *PPARγ* mRNA in Podocytes is Distinct from Adipose Tissue

Claire Bryant[1], Amy Webb[2], Alexander S Banks[3], Dawn Chandler[4,5], Rajgopal Govindarajan[6,7]
Shipra Agrawal*[1,5,8]

[1]*Center for Clinical and Translational Research, Abigail Wexner Research Institute at Nationwide Children's Hospital, Columbus, OH*
[2]*Department of Bioinformatics, The Ohio State University, Columbus, OH*
[3]*Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA*
[4]*Center for Childhood Cancer and Blood Disease, Abigail Wexner Research Institute at Nationwide Children's Hospital, Columbus, OH*
[5]*Department of Pediatrics, College of Medicine, The Ohio State University*
[6]*Translational Therapeutics, The Ohio State University Comprehensive Cancer Center*
[7]*Division of Pharmaceutics and Pharmacology, College of Pharmacy, The Ohio State University*
[8]*Division of Nephrology and Hypertension, Department of Medicine, Renaissance School of Medicine, Stony Brook University, Stony Brook, NY*

Short Title: Alternative Splicing of *PPARγ* mRNA in Podocytes

*Correspondence:

Shipra Agrawal, Ph.D.

Division of Nephrology and Hypertension

Department of Medicine

Health Sciences Center (HSC), T-15, Room 114, 101 Nicolls Rd.

Renaissance School of Medicine, Stony Brook University

Stony Brook, NY 11794

E-mail: Shipra.Agrawal@stonybrookmedicine.edu

ABSTRACT

Podocytes are highly differentiated epithelial cells, and their structural and functional integrity is compromised in a majority of glomerular and renal diseases, leading to proteinuria, chronic kidney disease, and kidney failure. Traditional agonists (e.g. pioglitazone) and selective modulators (e.g. GQ-16) of peroxisome-proliferator-activated-receptor-γ (PPARγ) reduce proteinuria in animal models of glomerular disease, and protect podocytes from injury via PPARγ activation. This indicates a pivotal role for PPARγ in maintaining glomerular function through preservation of podocytes distinct from its well-understood role in driving insulin sensitivity and adipogenesis. While its transcriptional role in activating adipokines and adipogenic genes is well-established in adipose tissue, liver and muscle, understanding of podocyte PPARγ signaling remains limited. We performed a comprehensive analysis of *PPARγ* mRNA variants due to alternative splicing, in human podocytes and compared with adipose tissue. We found that podocytes exclusively express the ubiquitous *PPARγ* Var 1 (encoding γ1) and not Var2 (encoding γ2), which is mostly restricted to adipose tissue and liver. Additionally, we detected expression at very low level of Var4, and barely detectable levels of other variants, Var3, Var11, VartORF4 and Var9, in podocytes. Podocyte vs adipose tissue expression of genes containing PPAR-promoter-response-elements suggested differential regulation by podocyte-specific PPARγ1 variant, distinct from the adipocyte-specific γ2 variant. This is also evident from distinct ingenuity-pathway-profile of RNAseq data of the PPARγ agonist-treated podocytes/glomeruli vs adipocytes/adipose tissue. In summary, podocytes and glomeruli express several PPARγ variants, including Var1 (γ1) and excluding adipocyte-specific Var2 (γ2), which may have implications in podocyte-specific signaling and pathophysiology.

INTRODUCTION

Thiazolidinediones (TZDs) or peroxisome proliferator activated receptor γ (PPARγ) agonists such as pioglitazone, directly protect podocytes from injury as demonstrated in podocyte culture *in vitro* studies [1-5], and reduce proteinuria and glomerular injury in various animal models of glomerular disease, as reported in preclinical *in vivo* studies [5-13]. Moreover, these beneficial effects of PPARγ have been shown to be mediated by activation of podocyte PPARγ as demonstrated by elegant studies using podocyte-specific *Pparg* knock out (KO) mouse model [7,10]. This indicates a pivotal role for PPARγ in maintaining glomerular function through preservation of podocytes [1,5-7,10]. Podocytes are highly differentiated epithelial cells in the kidneys, whose structural and functional integrity is critical for the maintenance of glomerular filtration barrier [14-16]. Accordingly, their dysfunction or loss is the initiating and progressing characteristic factor in a vast majority of renal diseases, leading to chronic kidney disease and kidney failure. Glomerular diseases characterized by high proteinuria manifest as nephrotic syndrome (NS) which is often associated with co-morbidities such as hypoalbuminemia, hypercholesterolemia, edema, and hyper-coagulopathy [17-22]. Recently, our group has further demonstrated that selective modulation of PPARγ with a partial agonist, GQ-16, provides high efficacy in reducing proteinuria, as well as NS-associated comorbidities in an experimental model of nephrotic syndrome [23]. Thus, a multitude of evidence shows that targeting the podocyte PPARγ pathway offers an attractive therapeutic strategy in NS. While PPARγ has an established role in driving many adipogenic and lipid metabolizing genes mainly in adipose tissue, liver, and muscle, the understanding of molecular pathways regulated by PPARγ in podocytes and glomeruli remains limited [5-7,9,10,24-28]. Furthermore, it is well-known that PPARγ exists in two major isoforms, γ1 and γ2, which are a result of different promoter usage as well as alternative splicing (AS) [29-31]. As a result, γ2 contains 30 additional amino acids at its N terminal, despite a longer mRNA sequence towards the 5'UTR of γ1. While γ1 is more widely expressed, γ2 is mostly restricted to adipose tissue and liver. Few other forms have been discovered, including some with dominant negative function [32-36] [37]. The specific variant or isoform(s) expression of *PPARγ* and the role of its AS in podocytes and glomerular disease is unexplored. In this study, we sought out to answer this critical gap in literature by performing a comprehensive analysis of different AS variants of PPARγ in human podocytes.

EXPERIMENTAL PROCEDURES

Podocyte Cell Culture and Treatments

Immortalized human podocytes were cultured in RMPI 1640 (Corning, Tewksbury, MA) supplemented with 10% fetal bovine serum (FBS) (Corning, Tewksbury, MA), 1% 100X Penicillin Streptomycin L-Glutamine (PSG) (Corning, Tewksbury, MA), and 1% Insulin-Transferrin-Selenium-Ethanolamine (ITS-X) (Gibco, Gaithersburg, MD) [38]. Proliferating podocytes were cultured in a humidified atmosphere with 5% CO_2 at 33ºC, and the media was changed twice per week and cells were passaged at ~70%-80% confluence. Differentiation was induced by placing the cells at 37ºC under the same atmospheric conditions, for 14 days. Differentiated cells were treated with puromycin aminonucleoside (PAN) (Sigma-Aldrich, St. Louis, MO) at 25 ug/ml or with pioglitazone (Alfa Aesar, Tewksbury, MA) at 10 µM. Control cells received treatment of vehicle dimethyl sulfoxide (DMSO) (Sigma-Aldrich, St. Louis, MO). Cells were harvested after 24 hours and total RNA isolated.

Kidney, Liver and Adipose Tissues

This study was approved by the Institution Animal Care and Use Committee at Nationwide Children's Hospital and the experiments were performed according to their guidelines. Male Wistar rats weighing ~150-200 g (Envigo, Indianapolis, IN) were purchased, acclimated for 3 days, and housed under standard conditions. They were sacrificed to collect kidneys, liver, and white adipose tissue (WAT) epididymal fat. Tissues were flash frozen in liquid nitrogen prior to RNA isolation. Glomeruli were isolated from harvested kidneys using the sequential sieving method as previously described [23]and total RNA was isolated. Commercially available human and rat kidneys total RNA was purchased from Zyagen (San Diego, CA).

RNA Isolation

Rat epididymal fat and liver tissue samples were lysed in RNA extraction buffer (RLT) from the RNeasy kit (Qiagen, Germantown, MD) with stainless-steel disruption beads for 4 minutes at 30.0 Hz using the Qiagen TissueLyser (Germantown, MD). Total RNA was immediately isolated from the lysate using the RNeasy kit (Qiagen, Germantown, MD), following the manufacturer's instructions. Total RNA was extracted from rat glomeruli tissue and human podocyte samples using the mirVanaTM Isolation Kit (Invitrogen, Carlsbad, CA), according to the manufacturer's instructions. RNA yield and purity was checked prior to downstream applications by measuring the absorbance at 260, 280, and 230 nm with a nanodrop spectrophotometer (ThermoFisher, Waltham, MA) and by calculating appropriate ratios (260/280, 260/230).

DNase Treatment, cDNA Synthesis, and PCR

Total RNA was DNase treated according to manufacturer's instructions (Zymo Research, Irvine, CA). 500 ng - 1µg DNase-treated RNA was reverse transcribed using the iScript cDNA Synthesis Kit (Bio-Rad, Hercules, CA), according to the manufacturer's instruction and the resulting cDNA was used for reverse transcription-polymerase chain reaction (RT-PCR) using HotStarTaq Plus Master Mix Kit (Qiagen, Germantown, MD). The PCR conditions used were as follows: 95°C for 5 minutes, 40 cycles [95°C for 30 seconds, 50-60°C (depending on the primer pair) for 30 seconds, 72°C for 30 seconds], hold at 4°C. Housekeeping gene *RPL6* was amplified only for 28 cycles to prevent over-saturation. The PCR-products were separated on a 1.5-2.5% agarose gel along with molecular weight ladder (ThermoFisher, Waltham,

MA), stained with 0.5µg/mL ethidium bromide (Research Products International, Mount Prospect, IL), and images were captured using the Chemidoc (Bio-Rad, Hercules, CA) equipment. Densitometry was performed using ImageJ Software (National Institutes of Health, Bethesda, MD) and band density was calculated by subtracting background and normalizing to *RPL6*. Quantitative RT-PCR (qRT-PCR) was performed on samples using SYBR green (Bio-Rad, Hercules, CA) on the Applied Biosystems 7500 Real-Time PCR System (Waltham, MA). The PCR conditions were 95ºC for 10 minutes, 40 X (95ºC for 15 seconds, 50-60ºC for 1 minute) followed by a melt curve to ensure specific products. Analysis was performed using the ΔΔCt method [39] with normalization to *RPL6*.

Primer Design and Synthesis

Primers were custom designed to detect various PPARγ variants for alternatively spliced mRNA products using the Reference Sequence Database (National Center for Biotechnology Information, Bethesda, MD) and confirmed for the potential binding transcripts in the annotated database using the Primer Blast program (National Center for Biotechnology Information, Bethesda, MD) (Table 1, Figure 1A). For variants with exon skipping, primers were designed to span the exon-exon junction to create a specific primer pair that would exclusively amplify the desired mRNA variant.

PPAR-Promoter Responsive Element Prediction

Key podocyte and adipocyte specific genes measured in this study were queried through the PPARgene database (ppargene.org) to predict PPAR-responsive elements (PPRE) on their promoters [40]. Upon query submission, p-value and confidence level were generated if the gene was predicted as a PPAR target gene and contained putative PPREs in the 5 kb transcription start site flanking region. They were assigned into categories of confidence as: high ($p > 0.8$), medium ($0.8 \geq p > 0.6$), and low ($0.6 \geq p > 0.45$). A p value ≤ 0.45 was predicted as negative.

Statistical Analysis

Statistical analysis was performed using GraphPad Prism Software version 8.2.0 for Windows (GraphPad Software, San Diego, CA). Data are expressed as mean ± standard error of mean and compared using unpaired Student's t-test. P value significance is depicted as *p<0.05, **p<0.01, ****p<0.0001.

RESULTS

Podocyte Specific PPARγ Splice Variants

To identify the PPARγ variants expressed in podocytes and to elucidate differences in podocytes vs adipocytes, we examined the known variants of PPARγ by reviewing the literature, scoping out the annotated versions (RefSeq database, NCBI), and by performing variant specific PCRs using custom-designed primers. A schematic of different variant forms (Figure 1A) and their exon/intron usage information and annotation (Figure 1B) is presented. Conducting PCR utilizing variant-specific primer pairs (Figure 1A and Table 1), we could identify total PPARγ expression, specific variants corresponding to γ1 (Var1 and 4) and other forms in differentiated podocytes, but not the γ2 form (Var 2) (Figure 1C), which is highly expressed in differentiated adipocytes [41]. Variant 2 (encodes γ2, a protein of 505 amino acids) includes exon B which contains a start codon. Variant 1 (encodes γ1, a protein of 475 amino acids) utilizes exons A1 and A2 instead of B in the 5'UTR of the gene, resulting in the usage of a start codon ATG located on exon 1. Variants 3 and 4 both encode γ1 as well, but they have differing 5'UTRs. Variant 3 contains an alternate exon A1.2, and variant 4 skips exon A2. Variant 11 has a similar 5'UTR to variant 1, but it skips exon 5, resulting in a frameshift and a premature stop codon in exon 6, encoding a potential product of 248 amino acids (γ3). Variant 9 utilizes a start codon located in exon 3 and skips exon 4, encoding γ5 (266 amino acids). Variant 10 or tORF4 uses a partially retained intron after exon 4 which contains a stop codon, making a product of 266 amino acids (γ6). Variant 10 version of tORF4 is annotated and contains exon A1.2, but there is no annotated variant that uses exon A1. The possibility of expression of other variants was considered using primer pairs that would amplify specific products (Figure 1C). The doublet primer pair was able to discriminate between the cumulative expression of variants 1,9 and 11 vs variant 4. Cumulative expression of variants 3, 5, 7, 10 and 12 was ruled out using primer pair for Var3 (dF and aR), which would also amplify these other mentioned variants.

Figure 1. Alternatively Spliced Variants and Isoforms of PPARγ in Podocytes. (A) The schematic depicts the exon and intron usage and the transcription and translation start and stop sites for various PPARγ variants. The major isoforms of PPARγ are isoforms γ1 and γ2, which encode for proteins containing 475 and 505 amino acids, respectively. The most common human annotated variants 1 (var1: NM_138712) and 2 (var2: NM_015869) encode isoforms γ1 and γ2 (depicted in the topmost PPARγ illustration). These two variants are transcribed by different promoter usage by RNA Polymerase II (depicted in tall arrows) and alternative splicing (depicted by exon skipping). PPARγ1 variant 1 includes exons A1 and A2, a start codon (ATG) on exon 1, and γ2 variant 2 excludes exons A1 and A2, but includes exon B and a start codon in the same exon B. Thus, γ1 protein product is 30 amino acids shorter than γ2. The less known and understood variants are represented below, which include Variants 3, 4, 11, 9 and tORF4/Var 10. These are depicted with A1 and A2 usage in the schematic to emphasize those likely to be expressed in podocytes, although their adipocyte counterparts with exon B usage at the 5' UTR also exist. A1 and A2 exons are shown in light blue, exon B in dark blue, internal exons are in green boxes, introns are black lines, spliced out exons are marked with

blue lines. Primer pairs used to amplify major variants and other variants are depicted in grey arrows and circles (A1, Ex1, a-j, Table 1). For the variants that skip an exon, the primer spans the exon-exon junctions and is shown as a split grey arrow underneath the introns (c, g i). Protein isoforms are shown below their corresponding RNA in purple and amino acid (aa) length is listed. **(B)** Summary of the protein isoforms (γ) and RNA variants (var) with detailed exon/intron usage and annotations. γ2 is found primarily in adipose tissue, while γ1 is found in podocytes. Variants 1, 3, and 4 are variants of γ1 that are likely to be found in podocytes, while variants 9, 10, and 11 are additional less explored variants that would encode different isoforms. The variant 10 version of tORF4 is annotated that contains exon A1.2, but there isn't an annotated version of tORf4 that would contain exon A1. **(C)** A representative gel showing the expression of several variants of PPARγ in differentiated podocytes. PCR products were generated using specific primer pairs and run on an agarose gel and stained with ethidium bromide. All PPARγ [primer pair Ex1 F, aR; 315bp]; γ2 [primer pair bF, aR; expected product 476bp]; variants 1, 9, 11 (432 bp PCR product) and Var 4 (358 bp product) [doublet primer pair, A1F, aR]; variants 3,5, and 10 (415bp product); variant 7 (341bp product); and variant 12 (503bp product) [primer pair dF, aR]. Primers are listed in Table 1.

Table 1. Primers Used in the Current Study

Name	Species	Forward	Reverse
PPARγ			
Total (All)	Homo Sapiens	ATGACCATGGTTGACACAG Ex1	GGAGTTGGAAGGCTCTTCAT a
Doublet	Homo Sapiens	CGAGGACACCGGAGAGGG A1	GGAGTTGGAAGGCTCTTCAT a
Var2 (γ2)	Homo Sapiens	TTTTAACGGATTGATCTTTTG b	GGAGTTGGAAGGCTCTTCAT a
Var1 (γ1)	Homo Sapiens	GAAAGAAGCCAACACTAAACCAC c	GGAGTTGGAAGGCTCTTCAT a
Var3 (γ1)	Homo Sapiens	TTCTGCTTAATTCCCTTTCC d	GGAGTTGGAAGGCTCTTCAT a
Var4 (γ1)	Homo Sapiens	GCCGCCGTGGCCGCAGAAA e	GGAGTTGGAAGGCTCTTCAT a
Var11 (γ3)	Homo Sapiens	CTTGCAGTGGGGATGTCTCA f	CAGCAAACCTGGGCGGTTGA g
Var9 (γ5)	Homo Sapiens	AGAGCCTTCCAACTCCCTCA h	GTCATAGATAACGAATGGCAT i
tORF4	Homo Sapiens	AGAGCCTTCCAACTCCCTCA h	AAACCCAAAACAACTTCCCG j
Var1 (γ1)*	Rattus Norvegicus	GAAAGAAGCTGTGAACCACT Rat c	GGAGTTTGAAGGCTCTTCAT Rat a
(γ2)*	Rattus Norvegicus	GCACTTCACAAGAAATTACC Rat b	GGAGTTTGAAGGCTCTTCAT Rat a
Total (All)*	Rattus Norvegicus	ATTACCATGGTTGACACAG Rat Ex1	GGAGTTTGAAGGCTCTTCAT Rat a
Podocyte Specific Genes			
NPHS1	Homo Sapiens	CGCAGGAGGAGGTGTCTTATTC	CGGGTTCCAGAGTGTCCAAG
Nphs1	Rattus Norvegicus	TGCTCTTTGCAGTTGGTGGT	TCCTGATCCTGTCCTCCGAC
SYNPO	Homo Sapiens	AGGTAGGCGTGGAGGAGG	GAGGTTCTGGTTGGGTTTGG
Synpo	Rattus Norvegicus	CCGCTTGGGTCCCCTTC	TGAACTCGTTCACCCTCTGC
WT1	Rattus Norvegicus/ Homo Sapiens	CCCTACAGCAGTGACAATTTATAC	TGCCCTTCTGTCCATTTC
Adipose Tissue Specific Genes			
CD36	Homo Sapiens	GGACTGCAGTGTAGGACTTTCC	TTCCGGTCACAGCCCATTTT
Cd36	Rattus Norvegicus	AAGTTATTGCGACATGATT	GATCCAAACACAGCATAGAT
ADIPOQ	Homo Sapiens	ACTGCAGTCTGTGGTTCTGA	ACTCCGGTTTCACCGATGTC
Adipoq	Rattus Norvegicus	TGTTCCTCTTAATCCTGCCCA	CCAACCTGCACAAGTTTCCTT
ADIPSIN	Homo Sapiens	CGCCCCGTGGTCGGAT	GACAGCTGTAGCAGCAGGAG
Adipsin	Rattus Norvegicus	AAGCTCTCCCACAATGCCTC	CATGGTACGTGCGCAGATTG
AP2	Homo Sapiens	AAAGTCAAGAGCACCATAACCTT	TGACGCCTTTCATGACGCATTCC
Ap2	Rattus Norvegicus	AAAGTGAAGAGCATCATAACCCT	TCACGCCTTTCATGACACATTCC
Housekeeping Genes			
GAPDH	Homo Sapiens	AGACACCATGGGGAAGGTGA	GACGGTGCCATGGAATTTGC
RPL6	Homo Sapiens	GCAACCCTGTCCTTGTCAGA	GCCAGCTGCTTCAGGAAAAC
Rpl6	Rattus Norvegicus	AGGCATCAGGAGGGTGAGAT	TGTGAGGGTACATGCCGTTC

*Rat nomenclature is annotated differently than human γ1 and γ2 variants.
 Primers a-j, Ex1, A1 are depicted in Figure 1A.
 Rat primers correspond to locations listed for human primers in Figure 1A.

PPARγ Splice Variants in Podocytes vs Adipose Tissue

PPARγ is a transcription factor and an established master regulator of fat cell physiology and differentiation in adipose tissue and liver [27,42]. While podocyte-specific *Pparg* KO mice illustrated its role in podocytopathies [7,10], to understand the cell- and tissue-specific determinants of PPARγ function in glomeruli and in podocytes, we performed specific PCRs to detect the expression of total (all) variants and exclusively of variant 1 (encoding γ1) and variant 2 (encoding γ2) in human podocytes and whole kidney and rat tissues of glomeruli, whole kidney, liver and WAT. In accordance with the hypothesis of cell specific action of PPARγ, while expression of variant encoding the PPARγ2 isoform was observed in WAT, liver, and whole kidneys, γ2 variant was undetected in human podocytes and in the rat glomeruli (Figure 2A). As shown in Figure 1A, γ1 and γ2 use different, promoter sites and undergo AS. In accordance, we observed a marked disparity in the expression of key functional genes in podocytes vs white adipose tissue (WAT) containing identified or putative PPREs (Figure 2B). While the podocyte markers (*NPHS1, SYNPO* and *WT1*) are expressed at a higher level in podocytes vs adipose tissue, expression of adipokines (*ADIPOQ* and *ADIPSIN*) and adipogenic genes (*CD36* and *AP2*) was found to be high in WAT and absent or modestly expressed in podocytes (Figure 2B). PPARγ binds to its promoter response elements (PPRE) along with other transcription factors such as retinoid X receptor (RXR) or to other DNA elements in association with factors such as nuclear factor kappa B (NFκB), activator protein 1 (AP1), and in proximity to CCAAT enhancer-binding proteins (C/EBPα) [5,24,43]. We carried out promoter element prediction using a PPARgene database (ppargene.org) [40] on genes known to play critical roles in podocyte pathophysiology, such as *NPHS1* (nephrin), *SYNPO* (synpatopodin), and *WT1* (wilms tumor 1) or those that have established roles in adipogenesis, such as *CD36, ADIPOQ, ADIPSIN*, and *AP2* [28,42,44-48] (Table 2). All the adipogenic genes depicted high level of PPRE prediction, and *NPHS1*, *SYNPO,* and *WT1* showed low to high levels of PPRE prediction. In summary, we propose that the podocyte genes are driven by the γ1 form of PPARγ, while the adipogenic genes are driven by the γ2 form. Expression analysis of several uncommon variants described in Figure 1 demonstrated that they are present at low and varying levels in the whole human kidney (Figure 2C). Most of these (variants 4, 3, 11, 9, and tORF4) were found to be expressed at very low to undetectable levels in podocytes (Figure 2C). Moreover, analysis of the variants encoding γ1 (Var 1 and 4) showed that their levels remained unchanged with PAN-induced injury of podocytes or treatment with PPARγ agonist, pioglitazone (Figure 2D).

Figure 2. Expression of PPARγ Splice Variants in Podocytes vs Adipose Tissue. (A) Total RNA was isolated from glomeruli, liver, and WAT samples harvested from rats, and from cultured differentiated human podocytes. Expression of the RNA variants encoding PPARγ1 and γ2 (All variants), γ1 only (Var1), and γ2 only (Var 2) isoforms was measured by RT-PCR in isolated samples as well as commercially available human and rat whole kidney RNA. Rat total (all) γ [primer pair ex1 F, ex 2 R; 315bp], rat γ1 [primer pair exA2 F, ex2 R; 380bp], rat γ2 [primer pair exB F, ex2 R; 329bp], rat *Rpl6* [178bp], human total (all) γ [primer pair Ex1 F, aR; 315bp], human γ1 [primer pair cF, aR; 387bp], human γ2 [primer pair bF, aR; 476bp], Human *RPL6* [370bp]. **(B)** Podocyte marker genes (*NPHS1* [Hu: 234bp, Rat: 138bp]], *SYNPO* [Hu: 182bp, Rat: 325bp], and *WT1* [Hu/Rat: 134bp]) and adipose tissue genes (*CD36* [Hu: 304bp, Rat: 123bp], *ADIPOQ* [Hu: 347bp, Rat: 95bp], *ADIPSIN* [Hu: 301bp, Rat: 202bp], and *AP2* [Hu: 133bp, Rat: 133bp]), and housekeeping gene *RPL6* [Hu: 370bp, Rat: 178bp] were analyzed. **(i)** Representative gels depicting the PCR products of key genes in podocytes and adipose tissue. **(ii-iii)** Quantitative mRNA fold - changes measured and graphically presented using RT-PCR and real time qRT-PCR quantification of key genes in podocytes and adipose tissue. **(C)** Several PPARγ mRNA AS variants were analyzed by RT-PCR in cultured differentiated human podocytes and commercially available human whole kidney RNA using the primer pairs drawn in Figure 1A and detailed in Supplementary Table 1. Var1 [primer pair cF, aR; 387bp], Var3 [primer pair dF, aR; 415bp], Var4 [primer pair eF, aR; 333bp], Var11 [primer pair fF, gR; 242bp], Var9 [primer pair hF, iR; 254bp], tORF4 [primer pair hF, jR; 490bp], Total [primer pair Ex1 F, aR; 315bp], *RPL6* [370bp] **(D)** Expression of PPARγ mRNA variants 1 [primer pair cF, aR] and 4 [primer pair eF, aR], as well as total (All) PPARγ [primer pair Ex1 F, aR] and GAPDH in puromycin aminonucleoside (PAN)-injured and pioglitazone-treated podocytes. Primers are listed in Table 1.

Table 2. PPAR-Responsive Element (PPRE) Prediction in Target Genes

Gene	Tissue Tested in the Current Study	p Value*	Confidence Level	PPREs Predicted#
Podocyte Prominent Genes				
Nphs1	Glomerular/WAT	0.63696	Medium	4
Synpo	Glomerular/WAT	0.8715	High	5
Wt1	Glomerular/WAT	0.55556	Low	5
Adipose Tissue Prominent Genes				
Cd36	Glomerular/WAT	0.97886	High	3
Adipoq	Glomerular/WAT	0.97153	High	12
Adipsin	Glomerular/WAT	0.96216	High	6
Ap2/Fabp4	Glomerular/WAT	0.99997	High	10
Housekeeping				
Rpl6	Glomerular/WAT	–	–	–

*Probability of being a PPAR target gene, higher value means a higher confidence
High-confidence ($p > 0.8$), median-confidence ($0.8 \geq p > 0.6$), low-confidence category ($0.6 \geq p > 0.45$).
Genes with p value ≤ 0.45 were predicted as negative.
#Putative PPREs in the 5Kb transcription start site (TSS) flanking region
WAT = white adipose tissue

DISCUSSION

PPARγ agonism has a well-established beneficial role in the setting of diabetes, which has led to the generation of TZDs for the treatment of type II diabetes (Figure 3) [49,50]. Its beneficial role in kidney cells beyond its favorable systemic metabolic effects in diabetes originated from numerous preclinical studies and meta-analyses [51-55]. Earlier these effects of PPARγ in influencing non-diabetic glomerular disease was thought to be through its anti-inflammatory actions on endothelial and myeloid cells [56-58]. However, recent discoveries and advances suggest the role of podocytes in mediating these effects. Seminal studies from our group and other research teams suggested that TZDs such as pioglitazone, directly protect podocytes from injury [1-5] and reduce proteinuria and glomerular injury in various animal models of glomerular disease [5-13] (Figure 3). Moreover, studies using podocyte-specific *Pparg* KO mouse demonstrated that the beneficial effects of PPARγ are mediated by activation of podocyte PPARγ, thus indicating a pivotal role for PPARγ in maintaining glomerular function through preservation of podocytes [1,5-7,10].

Figure 3. Illustration Describing the Roles of PPARγ 1 and 2 Isoforms. Schematic of the suggested roles of PPARγ 1 and 2 forms in regulating podocyte/glomerular disease specific effects (right panel) vs metabolic/systemic and adipogenic effects (left panel). PPARγ agonism plays a well-established role in the setting of treatment of Type II Diabetes, and treatment with its agonists, thiazolidinediones (TZDs), leads to increased ('a' pathway) insulin sensitivity and glycemic control [49,50]. However, these beneficial effects are also accompanied

with adverse effects ('a' pathway), such as increased cholesterol, adipogenesis and decreased osteoblastogenesis [24]. TZDs or PPARγ agonists have now been demonstrated to reduce podocyte injury ('b' pathway) by enhancing their viability and cytoskeletal integrity and decreasing apoptosis [1-5]. A few pathways involved in this process include cross-talk with the glucocorticoid receptor (GR), decreased caspase-3 activity and decreased TRPC6 expression. Gain of PPARγ activity or its activation by TZDs ('c' pathway) has been shown to reduce proteinuria and glomerulosclerosis in various animal models of glomerular disease, such as 5/6 nephrectomy (Nx), minimal change disease (MCD), focal segmental glomerulosclerosis (FSGS) and glomerulonephritis (GN) [5-13] [23]. On the other hand, loss of PPARγ in podocyte-specific *Pparg* KO mouse ('d' pathway) has demonstrated exacerbation of glomerular injury and proteinuria in animal disease models of FSGS and GN [1,5-7,10]. Analysis of the podocyte vs adipose tissue specific expression of genes downstream of PPARγ containing putative or identified PPREs (Figure 2B) informs us that these downstream effects in podocytes/glomeruli are likely directed by the PPARγ1 splice variant, distinct from the adipocyte-regulatory γ2 variant.

Since their identification in 1990, PPARγ has been recognized as a nuclear receptor superfamily member, a ligand-dependent transcription factor and a master regulator of adipogenesis and metabolism, which accounts for the insulin sensitizing effects of its agonists or anti-diabetic drugs such as TZDs [59]. While its role in regulating adipogenesis and lipid metabolism in adipose tissue, liver and skeletal muscle is well-characterized, the knowledge of PPARγ-regulated signaling pathways in podocytes is scarce [5-7,9,10,24-28]. In our previous studies we speculated a cross-talk model between the PPARγ and glucocorticoid receptor and further demonstrated that while pioglitazone could provide proteinuria reducing benefits, it could also enhance the ability of low-dose steroids to reduce proteinuria [6]. This suggested the potential clinical use of these FDA-approved drugs in enabling the reduction of steroid dose, toxicity, and side effects to treat NS. We further translated these findings and demonstrated the ability of pioglitazone administration to improve clinical outcome in a child with steroid-refractory NS, which has now been extended in other NS patients as well [6,60]. Moreover, we have recently found that targeting PPARγ with a selective modulator, GQ-16, offers an even better therapeutic strategy in NS, by reducing proteinuria and NS-associated side effects with a greater efficacy while providing reduced adipogenic effects than traditional TZDs [23]. However, to understand the proteinuria-reducing podocyte protective effects of PPARγ activation, it is critical to build an understanding of podocyte-specific transcriptional activity of PPARγ as the lack of such knowledge is a barrier to modulating PPARγ activity effectively. The existence of two isoforms γ1 and γ2 as a result of different promoter usage and AS further complicates the PPARγ biology [29-31]. As a result, γ2 contains 30 additional amino acids at its N terminal, despite a longer γ1 mRNA sequence at its 5'UTR. We found that podocytes as well as glomeruli, exclusively express the more ubiquitous form of PPARγ, the γ1 form and not the γ2 form, which is mostly restricted to adipose tissue and liver. Interestingly a few other spliced forms of PPARγ have been discovered in the past few years, which may confer varied functions [32-36]. Approximately 95% of mammalian genes undergo AS that can result in different protein products as well as microRNA sensitivity, mRNA stability, localization, and translation efficiency [61,62]. Moreover, AS is associated with many cellular processes and diseases such as differentiation, cancer, and immunity to name a few [61,63]. Moreover, expression of PPARγ1 and 2 mRNAs and isoforms in adipose tissue, liver, and skeletal muscle is known to be regulated in animal models of obesity, diabetes, and nutrition [64]. As the expression and role of the AS of PPARγ in podocyte biology is unexplored, we performed a landscape analysis of other PPARγ forms, which have been identified mostly in adipose tissues in recent years, including some with dominant negative function [32-36] [37]. We could

detect low expression of variants 4, 3, 11, and tORF4, and variant 9 was undetectable in the whole kidney. In podocytes, Var 9 was detectable at very low levels, and these additional variants were barely detectable consistently. We have recently also observed the presence of variant 11 form due to skipping of exon 5 (delta 5) in the WAT, which was particularly reduced with both pioglitazone and GQ-16 [23]. Ligand-mediated PPARγ activation has been shown to induce skipping of exon 5 which by itself lacks ligand-dependent transactivation ability, and thus acts as a dominant negative form of PPARγ [32]. Analysis of the podocyte vs adipose tissue specific expression of genes downstream of PPARγ containing putative or identified PPREs informs us that these downstream effects in podocytes are likely directed by the PPARγ1 splice variant, distinct from the adipocyte-regulatory γ2 variant (Figure 3). This is also evident from differential gene expression, ingenuity pathway profile, and signaling pathways in the podocytes/glomeruli vs adipocytes/adipose tissues derived from cells/animals treated with PPARγ agonists [23,65]. At the same time, we acknowledge the limitations of *in silico* prediction of PPREs, species-specific differences that may exist in the location and functionality of these PPREs [66], and the involvement of other co-transcription factors that determine cell specificity of PPARγ [43]. Overall, previous evidence indicates that the: 1) activation of PPARγ results in reduction in podocyte injury and proteinuria [1,4-6,12,13], 2) podocyte-specific PPARγ KO results in resistance to protection of proteinuria by PPARγ agonists [7,10], and 3) selective modulation of PPARγ enhances proteinuria reducing efficacy with reduced adipogenic potential [23]]. Taken together with the findings in the current study suggesting that the podocytes do not express the γ2 form, yet they are responsive to treatments by PPARγ agonists, implicate that those new selective modulators can be potentially developed that will be able to distinguish between the two forms, γ1 and γ2, thus forming a basis of novel targeted therapeutic avenues.

PPARγ also undergoes post-translational modifications and its differential phosphorylation at serine (Ser) 273 is an important determinant of its effect on adipogenesis and insulin sensitivity [42,65,67]. PPARγ phosphorylation at Ser273 was demonstrated in obesity models and treatment with the PPARγ agonist dephosphorylates this residue in the adipose tissue. GQ-16, a selective modulator of PPARγ has been shown to dephosphorylate PPARγ in an *in vitro* assay like traditional TZDs [68,69]. Since this site of phosphorylation on Ser273 is encoded by the last codon on exon 4, it remains conserved in major variants, 1 and 2, and several other variants in the current study, except for Var9, which was only detectable in podocytes at very low levels. It should be noted that the relative position is Ser243 in the variants including A1 and A2 exons instead of Ser273 in the B exon. Moreover, our recent findings suggested that relative phosphorylated PPARγ levels (at Ser273) remained unaltered during glomerular injury and with treatments with pioglitazone and GQ-16 treatments in the glomeruli [23]. These data suggest that while PPARγ-Ser273 is critical in determining the insulin sensitizing effects of PPARγ in adipose tissue, it is unlikely to be a major mechanism of regulation in podocytes or glomeruli.

In the present study, we have observed a clear indication of the disparate roles of γ1 vs γ2 in podocyte vs adipocyte function and physiology and demonstrated the presence of other AS variants of PPARγ mRNA in podocytes that would share the 5'UTR with variant 1 (encoding γ1 isoform), although the roles of these other variants in podocytopathies and podocyte differentiation is still unclear. Their detection of differential expression is a challenge as they are observed to be present at too low levels in podocytes (unlike the adipocyte variants). It might be worthwhile to pursue studies in future wherein these variants are overexpressed using reporter plasmids followed by the analysis of their effects on podocyte physiology,

actin cytoskeleton and function in the presence of injury causing agents, such as PAN. The demonstration of presence of these variants also warrants their detection in kidney biopsies from patients treated with PPARγ agonists for diabetic nephropathy or type II diabetes.

In summary, we have shown that podocytes and glomeruli express the variants encoding the major form of PPARγ1 isoform and do not express the variants encoding the other major form γ2. Moreover, expression of other AS forms of PPARγ has been detected at very low levels in podocytes which may have implications in its pathophysiology. Future studies directed at developing new selective modulators that can distinguish between the two forms, γ1 and γ2, to enhance beneficial effects and reduce harmful effects of PPARγ activation has the potential to provide novel targeted therapeutic avenues.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article material, further inquiries can be directed to the corresponding author.

ETHICS STATEMENT

The animal study was reviewed and approved by the Abigail Wexner Research Institute, Nationwide Children's Hospital (NCH) Institution Animal Care and Use Committee (IACUC).

AUTHOR CONTRIBUTIONS

Conceptualization, S.A.; Methodology, S.A., R.G., A.W., D.C., A.B. and C.B.; Formal Analysis, C.B., A.W. and S.A.; Investigation, C.B. and S.A.; Resources, S.A.; Data Curation, C.B., A.W. and S.A.; Writing – Original Draft Preparation, C.B. and S.A.; Writing – Review & Editing, S.A., R.G., A.W., D.C., A.B. and C.B.; Visualization, C.B. and S.A.; Supervision, S.A.; Project Administration, S.A.; Funding Acquisition, S.A.

FUNDING

This study was supported by the American Heart Association Career Development Award (CDA34110287) to SA, funds from Nationwide Children's Hospital (NCH) and Stony Brook University Renaissance School of Medicine to SA.

ACKNOWLEDGEMENTS

The authors thank Dr. Jeffrey Kopp at the National Institute of Health and Dr. Moin Saleem at the University of Bristol for sharing the immortalized human podocyte cell line, and members of the animal resource core at the Abigail Wexner Research Institute, Nationwide Children's Hospital (NCH).

REFERENCES

1. Agrawal, S.; Guess, A.J.; Benndorf, R.; Smoyer, W.E. Comparison of direct action of thiazolidinediones and glucocorticoids on renal podocytes: protection from injury and molecular effects. *Molecular pharmacology* **2011**, *80*, 389-399, doi:10.1124/mol.111.071654.

2. Miglio, G.; Rosa, A.C.; Rattazzi, L.; Grange, C.; Camussi, G.; Fantozzi, R. Protective effects of peroxisome proliferator-activated receptor agonists on human podocytes: proposed mechanisms of action. *British journal of pharmacology* **2012**, *167*, 641-653, doi:10.1111/j.1476-5381.2012.02026.x.

3. Miglio, G.; Rosa, A.C.; Rattazzi, L.; Grange, C.; Collino, M.; Camussi, G.; Fantozzi, R. The subtypes of peroxisome proliferator-activated receptors expressed by human podocytes and their role in decreasing podocyte injury. *British journal of pharmacology* **2011**, *162*, 111-125, doi:10.1111/j.1476-5381.2010.01032.x.

4. Kanjanabuch, T.; Ma, L.J.; Chen, J.; Pozzi, A.; Guan, Y.; Mundel, P.; Fogo, A.B. PPAR-gamma agonist protects podocytes from injury. *Kidney Int* **2007**, *71*, 1232-1239, doi:5002248 [pii] 10.1038/sj.ki.5002248.

5. Agrawal, S.; He, J.C.; Tharaux, P.L. Nuclear receptors in podocyte biology and glomerular disease. *Nat Rev Nephrol* **2021**, *17*, 185-204, doi:10.1038/s41581-020-00339-6.

6. Agrawal, S.; Chanley, M.A.; Westbrook, D.; Nie, X.; Kitao, T.; Guess, A.J.; Benndorf, R.; Hidalgo, G.; Smoyer, W.E. Pioglitazone Enhances the Beneficial Effects of Glucocorticoids in Experimental Nephrotic Syndrome. *Sci Rep* **2016**, *6*, 24392, doi:10.1038/srep24392.

7. Henique, C.; Bollee, G.; Lenoir, O.; Dhaun, N.; Camus, M.; Chipont, A.; Flosseau, K.; Mandet, C.; Yamamoto, M.; Karras, A.; et al. Nuclear Factor Erythroid 2-Related Factor 2 Drives Podocyte-Specific Expression of Peroxisome Proliferator-Activated Receptor gamma Essential for Resistance to Crescentic GN. *Journal of the American Society of Nephrology : JASN* **2016**, *27*, 172-188, doi:10.1681/ASN.2014111080.

8. Ma, L.J.; Marcantoni, C.; Linton, M.F.; Fazio, S.; Fogo, A.B. Peroxisome proliferator-activated receptor-gamma agonist troglitazone protects against nondiabetic glomerulosclerosis in rats. *Kidney Int* **2001**, *59*, 1899-1910, doi:kid695 [pii]10.1046/j.1523-1755.2001.0590051899.x.

9. Platt, C.; Coward, R.J. Peroxisome proliferator activating receptor-gamma and the podocyte. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association* **2016**, doi:10.1093/ndt/gfw320.

10. Sonneveld, R.; Hoenderop, J.G.; Isidori, A.M.; Henique, C.; Dijkman, H.B.; Berden, J.H.; Tharaux, P.L.; van der Vlag, J.; Nijenhuis, T. Sildenafil Prevents Podocyte Injury via PPAR-gamma-Mediated TRPC6 Inhibition. *Journal of the American Society of Nephrology : JASN* **2017**, *28*, 1491-1505, doi:10.1681/ASN.2015080885.

11. Yang, H.C.; Deleuze, S.; Zuo, Y.; Potthoff, S.A.; Ma, L.J.; Fogo, A.B. The PPARgamma agonist pioglitazone ameliorates aging-related progressive renal injury. *Journal of the American Society of Nephrology : JASN* **2009**, *20*, 2380-2388, doi:ASN.2008111138 [pii]10.1681/ASN.2008111138.

12. Yang, H.C.; Ma, L.J.; Ma, J.; Fogo, A.B. Peroxisome proliferator-activated receptor-gamma agonist is protective in podocyte injury-associated sclerosis. *Kidney Int* **2006**, *69*, 1756-1764, doi:5000336 [pii]10.1038/sj.ki.5000336.

13. Zuo, Y.; Yang, H.C.; Potthoff, S.A.; Najafian, B.; Kon, V.; Ma, L.J.; Fogo, A.B. Protective effects of PPARgamma agonist in acute nephrotic syndrome. *Nephrology, dialysis, transplantation :*

official publication of the European Dialysis and Transplant Association - European Renal Association **2012**, *27*, 174-181, doi:10.1093/ndt/gfr240.

14. Benzing, T. Molecular Design of the Kidney Filtration Barrier. *Trans Am Clin Climatol Assoc* **2020**, *131*, 125-139.

15. Butt, L.; Unnersjo-Jess, D.; Hohne, M.; Edwards, A.; Binz-Lotter, J.; Reilly, D.; Hahnfeldt, R.; Ziegler, V.; Fremter, K.; Rinschen, M.M.; et al. A molecular mechanism explaining albuminuria in kidney disease. *Nat Metab* **2020**, *2*, 461-474, doi:10.1038/s42255-020-0204-y.

16. Assady, S.; Wanner, N.; Skorecki, K.L.; Huber, T.B. New Insights into Podocyte Biology in Glomerular Health and Disease. *Journal of the American Society of Nephrology : JASN* **2017**, *28*, 1707-1715, doi:10.1681/ASN.2017010027.

17. Kerjaschki, D. 2015 Homer W. Smith Award: The Podocyte from Periphery to Center Stage. *Journal of the American Society of Nephrology : JASN* **2016**, *27*, 3266-3270, doi:10.1681/ASN.2016040490.

18. Siddall, E.C.; Radhakrishnan, J. The pathophysiology of edema formation in the nephrotic syndrome. *Kidney Int* **2012**, *82*, 635-642, doi:10.1038/ki.2012.180.

19. Radhakrishnan, J. Hypercoagulopathy in nephrotic syndrome. Available online: https://www.uptodate.com/contents/hypercoagulability-in-nephrotic-syndrome#H22

20. Araya, C.E.; Wasserfall, C.H.; Brusko, T.M.; Mu, W.; Segal, M.S.; Johnson, R.J.; Garin, E.H. A case of unfulfilled expectations. Cytokines in idiopathic minimal lesion nephrotic syndrome. *Pediatr Nephrol* **2006**, *21*, 603-610, doi:10.1007/s00467-006-0026-5.

21. Araya, C.; Diaz, L.; Wasserfall, C.; Atkinson, M.; Mu, W.; Johnson, R.; Garin, E. T regulatory cell function in idiopathic minimal lesion nephrotic syndrome. *Pediatr Nephrol* **2009**, *24*, 1691-1698, doi:10.1007/s00467-009-1214-x.

22. Agrawal, S.; Zaritsky, J.J.; Fornoni, A.; Smoyer, W.E. Dyslipidaemia in nephrotic syndrome: mechanisms and treatment. *Nat Rev Nephrol* **2018**, *14*, 57-70, doi:10.1038/nrneph.2017.155.

23. Bryant, C.; Rask, G.; Waller, A.P.; Webb, A.; Galdino-Pitta, M.R.; Amato, A.A.; Cianciolo, R.; Govindarajan, R.; Becknell, B.; Kerlin, B.A.; et al. Selective modulator of nuclear receptor PPARgamma with reduced adipogenic potential ameliorates experimental nephrotic syndrome. *iScience* **2022**, *25*, 104001, doi:10.1016/j.isci.2022.104001.

24. Lefterova, M.I.; Haakonsson, A.K.; Lazar, M.A.; Mandrup, S. PPARgamma and the global map of adipogenesis and beyond. *Trends in endocrinology and metabolism: TEM* **2014**, *25*, 293-302, doi:10.1016/j.tem.2014.04.001.

25. Berger, J.P.; Akiyama, T.E.; Meinke, P.T. PPARs: therapeutic targets for metabolic disease. *Trends in pharmacological sciences* **2005**, *26*, 244-251, doi:10.1016/j.tips.2005.03.003.

26. Ahmadian, M.; Suh, J.M.; Hah, N.; Liddle, C.; Atkins, A.R.; Downes, M.; Evans, R.M. PPARgamma signaling and metabolism: the good, the bad and the future. *Nat Med* **2013**, *19*, 557-566, doi:10.1038/nm.3159.

27. Evans, R.M.; Barish, G.D.; Wang, Y.X. PPARs and the complex journey to obesity. *Nat Med* **2004**, *10*, 355-361, doi:10.1038/nm1025.

28. Agrawal, S.; Guess, A.J.; Chanley, M.A.; Smoyer, W.E. Albumin-induced podocyte injury and protection are associated with regulation of COX-2. *Kidney Int* **2014**, *86*, 1150-1160, doi:10.1038/ki.2014.196.

29. Fajas, L.; Auboeuf, D.; Raspe, E.; Schoonjans, K.; Lefebvre, A.M.; Saladin, R.; Najib, J.; Laville, M.; Fruchart, J.C.; Deeb, S.; et al. The organization, promoter analysis, and expression of the

human PPARgamma gene. *The Journal of biological chemistry* **1997**, *272*, 18779-18789, doi:10.1074/jbc.272.30.18779.

30.   Mukherjee, R.; Jow, L.; Croston, G.E.; Paterniti, J.R., Jr. Identification, characterization, and tissue distribution of human peroxisome proliferator-activated receptor (PPAR) isoforms PPARgamma2 versus PPARgamma1 and activation with retinoid X receptor agonists and antagonists. *The Journal of biological chemistry* **1997**, *272*, 8071-8076, doi:10.1074/jbc.272.12.8071.

31.   Agrawal, S.; He, J.C.; Tharaux, P.L. Nuclear Receptors in Podocyte Biology and Glomerular Disease. *Nature Reviews Nephrology, Review. Accepted (In Press).* **2020**.

32.   Aprile, M.; Cataldi, S.; Ambrosio, M.R.; D'Esposito, V.; Lim, K.; Dietrich, A.; Bluher, M.; Savage, D.B.; Formisano, P.; Ciccodicola, A.; et al. PPARgammaDelta5, a Naturally Occurring Dominant-Negative Splice Isoform, Impairs PPARgamma Function and Adipocyte Differentiation. *Cell reports* **2018**, *25*, 1577-1592 e1576, doi:10.1016/j.celrep.2018.10.035.

33.   Aprile, M.; Ambrosio, M.R.; D'Esposito, V.; Beguinot, F.; Formisano, P.; Costa, V.; Ciccodicola, A. PPARG in Human Adipogenesis: Differential Contribution of Canonical Transcripts and Dominant Negative Isoforms. *PPAR research* **2014**, *2014*, 537865, doi:10.1155/2014/537865.

34.   Chen, Y.; Jimenez, A.R.; Medh, J.D. Identification and regulation of novel PPAR-gamma splice variants in human THP-1 macrophages. *Biochimica et biophysica acta* **2006**, *1759*, 32-43, doi:10.1016/j.bbaexp.2006.01.005.

35.   Sabatino, L.; Casamassimi, A.; Peluso, G.; Barone, M.V.; Capaccio, D.; Migliore, C.; Bonelli, P.; Pedicini, A.; Febbraro, A.; Ciccodicola, A.; et al. A novel peroxisome proliferator-activated receptor gamma isoform with dominant negative activity generated by alternative splicing. *The Journal of biological chemistry* **2005**, *280*, 26517-26525, doi:10.1074/jbc.M502716200.

36.   Zhou, J.; Wilson, K.M.; Medh, J.D. Genetic analysis of four novel peroxisome proliferator activated receptor-gamma splice variants in monkey macrophages. *Biochemical and biophysical research communications* **2002**, *293*, 274-283, doi:10.1016/S0006-291X(02)00138-9.

37.   Mukha, A.; Kalkhoven, E.; van Mil, S.W.C. Splice variants of metabolic nuclear receptors: Relevance for metabolic disease and therapeutic targeting. *Biochim Biophys Acta Mol Basis Dis* **2021**, *1867*, 166183, doi:10.1016/j.bbadis.2021.166183.

38.   Saleem, M.A.; O'Hare, M.J.; Reiser, J.; Coward, R.J.; Inward, C.D.; Farren, T.; Xing, C.Y.; Ni, L.; Mathieson, P.W.; Mundel, P. A conditionally immortalized human podocyte cell line demonstrating nephrin and podocin expression. *Journal of the American Society of Nephrology : JASN* **2002**, *13*, 630-638.

39.   Pfaffl, M.W. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic acids research* **2001**, *29*, e45, doi:10.1093/nar/29.9.e45.

40.   Fang, L.; Zhang, M.; Li, Y.; Liu, Y.; Cui, Q.; Wang, N. PPARgene: A Database of Experimentally Verified and Computationally Predicted PPAR Target Genes. *PPAR research* **2016**, *2016*, 6042162, doi:10.1155/2016/6042162.

41.   Tontonoz, P.; Hu, E.; Spiegelman, B.M. Stimulation of adipogenesis in fibroblasts by PPAR gamma 2, a lipid-activated transcription factor. *Cell* **1994**, *79*, 1147-1156, doi:10.1016/0092-8674(94)90006-x.

42.   Choi, J.H.; Banks, A.S.; Estall, J.L.; Kajimura, S.; Bostrom, P.; Laznik, D.; Ruas, J.L.; Chalmers, M.J.; Kamenecka, T.M.; Bluher, M.; et al. Anti-diabetic drugs inhibit obesity-linked phosphorylation of PPARgamma by Cdk5. *Nature* **2010**, *466*, 451-456, doi:10.1038/nature09291.

43. Lefterova, M.I.; Steger, D.J.; Zhuo, D.; Qatanani, M.; Mullican, S.E.; Tuteja, G.; Manduchi, E.; Grant, G.R.; Lazar, M.A. Cell-specific determinants of peroxisome proliferator-activated receptor gamma function in adipocytes and macrophages. *Mol Cell Biol* **2010**, *30*, 2078-2089, doi:10.1128/MCB.01651-09.

44. Ettou, S.; Jung, Y.L.; Miyoshi, T.; Jain, D.; Hiratsuka, K.; Schumacher, V.; Taglienti, M.E.; Morizane, R.; Park, P.J.; Kreidberg, J.A. Epigenetic transcriptional reprogramming by WT1 mediates a repair response during podocyte injury. *Sci Adv* **2020**, *6*, eabb5460, doi:10.1126/sciadv.abb5460.

45. Eddy, A.A.; Fogo, A.B. Plasminogen activator inhibitor-1 in chronic kidney disease: evidence and mechanisms of action. *Journal of the American Society of Nephrology : JASN* **2006**, *17*, 2999-3012, doi:10.1681/ASN.2006050503.

46. Kobayashi, N.; Ueno, T.; Ohashi, K.; Yamashita, H.; Takahashi, Y.; Sakamoto, K.; Manabe, S.; Hara, S.; Takashima, Y.; Dan, T.; et al. Podocyte injury-driven intracapillary plasminogen activator inhibitor type 1 accelerates podocyte loss via uPAR-mediated beta1-integrin endocytosis. *American journal of physiology. Renal physiology* **2015**, *308*, F614-626, doi:10.1152/ajprenal.00616.2014.

47. Cheng, H.; Fan, X.; Guan, Y.; Moeckel, G.W.; Zent, R.; Harris, R.C. Distinct roles for basal and induced COX-2 in podocyte injury. *Journal of the American Society of Nephrology : JASN* **2009**, *20*, 1953-1962, doi:10.1681/ASN.2009010039.

48. Asanuma, K.; Kim, K.; Oh, J.; Giardino, L.; Chabanis, S.; Faul, C.; Reiser, J.; Mundel, P. Synaptopodin regulates the actin-bundling activity of alpha-actinin in an isoform-specific manner. *J Clin Invest* **2005**, *115*, 1188-1198, doi:10.1172/JCI23371.

49. Ahsan, W. The Journey of Thiazolidinediones as Modulators of PPARs for the Management of Diabetes: A Current Perspective. *Curr Pharm Des* **2019**, *25*, 2540-2554, doi:10.2174/1381612825666190716094852.

50. Hofmann, C.A.; Colca, J.R. New oral thiazolidinedione antidiabetic agents act as insulin sensitizers. *Diabetes care* **1992**, *15*, 1075-1078, doi:10.2337/diacare.15.8.1075.

51. Sarafidis, P.A.; Stafylas, P.C.; Georgianos, P.I.; Saratzis, A.N.; Lasaridis, A.N. Effect of thiazolidinediones on albuminuria and proteinuria in diabetes: a meta-analysis. *Am J Kidney Dis* **2010**, *55*, 835-847, doi:S0272-6386(09)01583-2 [pii]10.1053/j.ajkd.2009.11.013.

52. Buckingham, R.E.; Al-Barazanji, K.A.; Toseland, C.D.; Slaughter, M.; Connor, S.C.; West, A.; Bond, B.; Turner, N.C.; Clapham, J.C. Peroxisome proliferator-activated receptor-gamma agonist, rosiglitazone, protects against nephropathy and pancreatic islet abnormalities in Zucker fatty rats. *Diabetes* **1998**, *47*, 1326-1334, doi:10.2337/diab.47.8.1326.

53. Cha, D.R.; Zhang, X.; Zhang, Y.; Wu, J.; Su, D.; Han, J.Y.; Fang, X.; Yu, B.; Breyer, M.D.; Guan, Y. Peroxisome proliferator activated receptor alpha/gamma dual agonist tesaglitazar attenuates diabetic nephropathy in db/db mice. *Diabetes* **2007**, *56*, 2036-2045, doi:10.2337/db06-1134.

54. Tanimoto, M.; Fan, Q.; Gohda, T.; Shike, T.; Makita, Y.; Tomino, Y. Effect of pioglitazone on the early stage of type 2 diabetic nephropathy in KK/Ta mice. *Metabolism* **2004**, *53*, 1473-1479.

55. Calkin, A.C.; Giunti, S.; Jandeleit-Dahm, K.A.; Allen, T.J.; Cooper, M.E.; Thomas, M.C. PPAR-alpha and -gamma agonists attenuate diabetic kidney disease in the apolipoprotein E knockout mouse. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association* **2006**, *21*, 2399-2405, doi:10.1093/ndt/gfl212.

56. Chinetti, G.; Griglio, S.; Antonucci, M.; Torra, I.P.; Delerive, P.; Majd, Z.; Fruchart, J.C.; Chapman, J.; Najib, J.; Staels, B. Activation of proliferator-activated receptors alpha and gamma induces apoptosis of human monocyte-derived macrophages. *The Journal of biological chemistry* **1998**, *273*, 25573-25580.

57. Bouhlel, M.A.; Derudas, B.; Rigamonti, E.; Dievart, R.; Brozek, J.; Haulon, S.; Zawadzki, C.; Jude, B.; Torpier, G.; Marx, N.; et al. PPARgamma activation primes human monocytes into alternative M2 macrophages with anti-inflammatory properties. *Cell metabolism* **2007**, *6*, 137-143, doi:S1550-4131(07)00166-0 [pii]10.1016/j.cmet.2007.06.010.

58. Straus, D.S.; Glass, C.K. Anti-inflammatory actions of PPAR ligands: new insights on cellular and molecular mechanisms. *Trends in immunology* **2007**, *28*, 551-558, doi:10.1016/j.it.2007.09.003.

59. Heikkinen, S.; Auwerx, J.; Argmann, C.A. PPARgamma in human and mouse physiology. *Biochimica et biophysica acta* **2007**, *1771*, 999-1013, doi:10.1016/j.bbalip.2007.03.006.

60. Hunley, T.E.; Ng, K.H.; Smoyer, W.E.; Hidalgo, G. Pioglitazone (Pio) in Pediatric Nephrotic Syndrome (NS). *American Society of Nephrology (Kidney Week) Abstract* **2019**, *SA-PO679*, 939.

61. Scotti, M.M.; Swanson, M.S. RNA mis-splicing in disease. *Nature reviews. Genetics* **2016**, *17*, 19-32, doi:10.1038/nrg.2015.3.

62. Blijlevens, M.; Li, J.; van Beusechem, V.W. Biology of the mRNA Splicing Machinery and Its Dysregulation in Cancer Providing Therapeutic Opportunities. *Int J Mol Sci* **2021**, *22*, doi:10.3390/ijms22105110.

63. Montes, M.; Sanford, B.L.; Comiskey, D.F.; **Chandler, D.S.** <u>RNA Splicing and Disease: Animal Models to Therapies</u>. *Trends Genet* **2019**, *35*, 68-87, doi:10.1016/j.tig.2018.10.002.

64. Vidal-Puig, A.; Jimenez-Linan, M.; Lowell, B.B.; Hamann, A.; Hu, E.; Spiegelman, B.; Flier, J.S.; Moller, D.E. Regulation of PPAR gamma gene expression by nutrition and obesity in rodents. *J Clin Invest* **1996**, *97*, 2553-2561, doi:10.1172/JCI118703.

65. Hall, J.A.; Ramachandran, D.; Roh, H.C.; DiSpirito, J.R.; Belchior, T.; Zushin, P.H.; Palmer, C.; Hong, S.; Mina, A.I.; Liu, B.; et al. Obesity-Linked PPARgamma S273 Phosphorylation Promotes Insulin Resistance through Growth Differentiation Factor 3. *Cell metabolism* **2020**, *32*, 665-675 e666, doi:10.1016/j.cmet.2020.08.016.

66. Schmidt, S.F.; Jorgensen, M.; Chen, Y.; Nielsen, R.; Sandelin, A.; Mandrup, S. Cross species comparison of C/EBPalpha and PPARgamma profiles in mouse and human adipocytes reveals interdependent retention of binding sites. *BMC Genomics* **2011**, *12*, 152, doi:10.1186/1471-2164-12-152.

67. Choi, J.H.; Banks, A.S.; Kamenecka, T.M.; Busby, S.A.; Chalmers, M.J.; Kumar, N.; Kuruvilla, D.S.; Shin, Y.; He, Y.; Bruning, J.B.; et al. Antidiabetic actions of a non-agonist PPARgamma ligand blocking Cdk5-mediated phosphorylation. *Nature* **2011**, *477*, 477-481, doi:10.1038/nature10383.

68. Amato, A.A.; Rajagopalan, S.; Lin, J.Z.; Carvalho, B.M.; Figueira, A.C.; Lu, J.; Ayers, S.D.; Mottin, M.; Silveira, R.L.; Souza, P.C.; et al. GQ-16, a novel peroxisome proliferator-activated receptor gamma (PPARgamma) ligand, promotes insulin sensitization without weight gain. *The Journal of biological chemistry* **2012**, *287*, 28169-28179, doi:10.1074/jbc.M111.332106.

69. Coelho, M.S.; de Lima, C.L.; Royer, C.; Silva, J.B.; Oliveira, F.C.; Christ, C.G.; Pereira, S.A.; Bao, S.N.; Lima, M.C.; Pitta, M.G.; et al. GQ-16, a TZD-Derived Partial PPARgamma Agonist, Induces the Expression of Thermogenesis-Related Genes in Brown Fat and Visceral White Fat and

Decreases Visceral Adiposity in Obese and Hyperglycemic Mice. *PloS one* **2016**, *11*, e0154310, doi:10.1371/journal.pone.0154310.