Hourglass, a tool to mine bioimaging data, uncovers sex-disparities in the IL-6-associated T cell response in pancreatic tumors

Authors and affiliations

Kazeera Aliar[1], Henry R. Waterhouse[1], Foram Vyas[1,2], Niklas Krebs[1,3], Emily Poulton[1,2], Bowen Zhang[1,2], Nathan Chan[1], Peter Bronsert[4], Sandra E. Fischer[5,6], Steven Gallinger[7,8,9,10], Barbara T. Grünwald[1] *, Rama Khokha[1,2,11] *

Authors and Affiliations
Princess Margaret Cancer Centre University Health Network, Toronto, ON M5G 1L7, Canada
Department of Medical Biophysics, University of Toronto, Toronto, ON M5G 1L7, Canada
Department of Gastroenterology, Gastrointestinal Oncology & Endocrinology, University Medical Center Göttingen, 37075 Göttingen, Germany
Core Facility for Histopathology and Digital Pathology, Medical Center-University of Freiburg, 79106 Freiburg, Germany
Department of Laboratory Medicine and Pathobiology, University of Toronto, University Health Network, Toronto, ON M5G 2M9, Canada
Division of Anatomic Pathology, Laboratory Medicine Program, University Health Network, Toronto, ON M5G 2C4, Canada
PanCuRx Translational Research Initiative, Ontario Institute for Cancer Research, Toronto, ON M5G 0A3, Canada
Department of Surgery, University of Toronto, Toronto, ON M5T 1P5, Canada
Lunenfeld Tanenbaum Research Institute, Mount Sinai Hospital, Toronto, ON M5G 1X5, Canada
Hepatobiliary/Pancreatic Surgical Oncology Program, University Health Network, Toronto, ON M5G 2M9, Canada
Lead Contact
*   These authors contributed equally: Barbara T. Grünwald, Rama Khokha

Contact info
Correspondence: barbara.grunwald@uhnresearch.ca, rama.khokha@uhnresearch.ca

Summary

        Recent advances in digital pathology have led to an explosion in high-content multidimensional imaging approaches. Yet, our ability to gainfully process, visualize, integrate and mine the resulting mass of bioimaging data remains a challenge. We have developed Hourglass, an open access user-friendly software that streamlines complex biology-driven post-processing and visualization of multiparametric data. Directed at datasets derived from tissue microarrays or imaging methods that analyze multiple regions of interest per patient specimen, Hourglass systematically organizes observations across spatial and global levels

as well as within patient subgroups. Application of Hourglass to our large and complex
pancreatic cancer bioimaging dataset (540,617 datapoints derived from 26 bioimaging
analyses applied to 596 specimens from 165 patients) consolidated a breadth of known IL-6
functions in a well-annotated human pancreatic cancer cohort and uncovered new
unprecedented insights into a sex-linked Interleukin-6 (IL-6) association with immune
phenotypes. Specifically, regional effects of IL-6 on the intratumoral T cell response were
restricted to male patients only. In conclusion, Hourglass facilitates multi-layered knowledge
extraction from complex multiparametric bioimaging datasets and provides tailored analytical
means to productively harness heterogeneity at the sample and patient level.

Keywords
bioimage analysis, data visualization, digital pathology, interleukin-6, pancreatic cancer

Introduction

The digital pathology space is rapidly expanding and producing an immense amount
of data with the potential for insights in the magnitude of OMICs profiling approaches (1).
Typical sources of complex bioimaging datasets include imaging mass cytometry (IMC) and
multiplexed immunohistochemistry (IHC) staining, often applied to large numbers of samples
and/or tissue microarrays (TMA). In bioimaging data, one typically encounters multiple
quantitative parameters per molecular target (*e.g.* percentage of positive pixels, number of
positively stained cells, H scores, etc. for each stain; Table 1), as opposed to a single value
per target such as obtained from OMICs experiments (*e.g.* a single protein or mRNA
abundance value). The inherent multidimensionality of bioimaging data thereby poses
significant challenges for targeted analysis and visualization. As a result, manual iterative
hypothesis generation becomes a lengthy, labor-intensive task and a major challenge.
Moreover, the ability to robustly integrate multiple parameters is equally challenging without
computational knowledge yet remains indispensable for exposing meaningful biological
patterns in such complex, high-dimensional data. Thus, the wealth of information that lies
within these datasets is often barely utilized. In addition, bioimaging datasets and the
corresponding analyses often stem from contexts with significant heterogeneity at the
specimen and clinical patient levels, posing further challenges to their analysis.

Heterogeneity is a fundamental property of all tissues and oftentimes becomes
exacerbated in disease conditions (2–5). Intratumoral heterogeneity is increasingly recognized
as a major cancer driver (6–9), with key implications for treatment design and patient selection
and a rapidly increasing appreciation of the critical disease contributions stemming from
regional cellular communities (10,11). Imaging-based approaches, and thus the digital
pathology field, are uniquely positioned to resolve such spatial differences down to a cellular
level, enabling a more comprehensive biological understanding of the highly heterogeneous
tumor ecosystem (1,12). Yet, even though standard TMA designs and typical IMC experiments
incorporate multiple regions of interest in order to minimize the impact of sampling bias, the
inherent opportunity of such experimental design to systematically compare and dissect
regional from global biological patterns by analyzing multiple regions of the same tissue is
rarely harnessed.

In order to support effective analysis of such multi-patient multi-sample bioimaging
datasets, we developed Hourglass. Built as a computational toolkit, Hourglass was designed

to facilitate streamlined and reproducible processing, integration, visualization and mining of multiparametric datasets, and specifically tailored to tissue-based bioimaging data. The software is available as a stand-alone R package with an optional user-friendly desktop application (at https://kazeera.github.io/Hourglass). To assess the utility of Hourglass, we queried the effects of Interleukin-6 (IL-6) in pancreatic ductal adenocarcinoma (PDAC) using an extensive dataset comprising 34 read-outs from 26 bioimaging analyses, each performed on 596 samples from 165 PDAC patients. IL-6 is a powerful regulator of tumor biology, with complex functions in both the tumor and tumor microenvironment (TME)(13,14). Several clinical trials on targeting IL-6 along with chemotherapy in pancreatic cancer are currently ongoing (e.g., NCT02767557, NCT04191421, NCT00841191). Moreover, a recent study exposed concrete effects of IL-6 on functional tumor-associated fibroblast diversity in PDAC, which has major and topical immunotherapeutic implications (15). In addition to the versatile immunomodulatory and proinflammatory effects of IL-6 in pancreatic cancer are well-established (16–18), altogether forming a strong rationale for the pursuit of combination immunotherapy/IL-6 targeting strategies in PDAC and other cancers (19–23). Notably, IL-6 levels are known to be regionally heterogeneous (19,24) with varying clinical associations, which could hint at hidden regional biological patterns evolving around this influential cytokine and this heterogeneity has potential implications for the selection of patients for future IL-6-based treatments.

We here applied Hourglass for the systematic dissection of IL-6-associated effects across multiple cellular compartments and biological processes in a well-annotated human PDAC cohort, under consideration of regional heterogeneity in IL-6 expression and clinically relevant patient subsets (sex, molecular subtypes). Our findings not only consolidate known functions of IL-6 in regulating both tumoral and stromal biology in the context human PDAC but identify a substantial involvement of regional IL-6 expression levels in distinct spatial immune phenotypes. Moreover, we discovered that the regional association of IL-6 levels with T cell communities and FAP+ fibroblasts was nearly exclusive to the male subset of PDAC patients.

Results

Hourglass workflow for systematic mining of annotated multiparametric bioimaging datasets

The Hourglass workflow follows three steps: dataset aggregation and import (Fig. 1A); data processing (defining comparisons, data subsets and analysis settings, selecting statistics; Fig. 1B); output mining and quality control based on visualized analysis results (Fig. 1C). For this study, we have chosen a PDAC cohort (n = 165) with detailed clinical and molecular annotations (Fig. 1A, *left*), such as transcriptional subtype, patient sex, overall survival time, tumor differentiation, etc. This cohort was used to construct a TMA, from which we generated a typical bioimaging dataset (Fig. 1A, *right*) consisting of 34 readouts from 26 stains that covered a variety of IHC markers for malignant behavior (e.g., proliferation, EMT) and TME composition (e.g., different immune cell subpopulations, fibroblasts) along with second harmonic generation microscopy and histochemical collagen stains. All stains were digitally quantified via either or both pixel- and cell-based detection metrics (Fig. S1). Where appropriate, CK19-based estimation of TME and tumor area aided compartment-specific

132 normalization or, alternatively, quantification was restricted to manual annotations of tumoral
133 or stromal compartments by a pathologist. Aggregation of this bioimaging data with the
134 corresponding cohort annotations led to a final input dataset comprising 540,617 unique
135 entries.
136
To enable the systematic and reproducible analysis of such heterogeneous
multiparametric datasets, Hourglass facilitates the import of large numeric data files with
extensive annotation, transforms and subsets the data based on user specifications, and
produces publication-ready visualizations including pre-selected statistical tests (Fig. 1B). A
variety of analysis and corresponding plot types is supported and can be selected for each
analysis run; this includes comparisons between discrete groups (e.g., box plots, bar plots,
slopegraphs), correlation analyses (e.g., correlation matrices, scatter plots) and Kaplan Meier
survival analysis (see Table 2). The output (Fig. 1C) is stored in a structured file hierarchy with
user-defined biological comparison folders and additional subfolders containing user-defined
feature subsets. To enables visual data quality control and for identification of potential hidden
confounders, Hourglass furthermore routinely plots all available parameters for each feature
(in this context: stain) as a function of the user-defined comparison. Finally, analysis settings
are logged and stored in a separate file for each run to enable full cross-run reproducibility.
150
151 **Graphical user interface (GUI) supports navigation through Hourglass workflow**
152 **without prior computational expertise**
153
The software application for Hourglass was made accessible to non-expert users,
while also supporting customizable analysis at the experienced user level. Therefore, a
graphical user interface (GUI) was built to take users through the Hourglass algorithm via step-
by-step tabs (Fig. 2A-C). In brief, the Hourglass application allows manual selection of input
data and defining biological comparisons of interest based on values present in the input data
files (Fig. 2A), selection of feature subsets and customization of colors (Fig. 2B), along with
advanced options to iteratively refine analyses and a final selection of parameters for the
current run, including manual selection from supported plot types and quality control settings
(Fig. 2C). The application then saves the user analysis specifications (Fig. 2D) as an Excel file
and interfaces to the R statistical environment. Finally, the corresponding Hourglass R
package produces the output as a sequential background process (Fig. 2E). Computational
expert users can work directly within the Hourglass R package, without the GUI.
166
167 **Inter- and intra- tumoral heterogeneity in IL-6 expression levels**
168
To test the utility of Hourglass, we chose to profile the breadth of IL-6-related biology
in human PDAC, aiming at a systematic dissection of the powerful multipronged effects under
consideration of the known intratumoral heterogeneity of IL-6. An IL-6 IHC stain was optimized
using a set of control tissues (Fig. S2A) and then applied to a TMA from well-annotated PDAC
patients (Fig. 1A). IHC showed varying IL-6 expression levels across PDAC patients with
staining in malignant epithelial cells, fibroblasts and immune cells (Fig. 3A, *left*). Furthermore,
IL-6 levels varied intratumorally within both malignant epithelial and stromal regions (Fig. 3A,
right). As a result, when we divided all samples based on 33% quantiles (IL-6^{lo}, IL-6^{int} and IL-
6^{hi} Fig. S2B), we observed that different TMA cores from the same patient exhibited varying
IL-6 expression categories (Fig. 3B) in 33.6% of PDAC cases. While our analysis here focuses
on IL-6, we observed similar levels intratumoral heterogeneity (i.e., multiple cores from the

same tumor being classified as IL-6^{lo}, IL-6^{int} and/or IL-6^{hi}) for a range of markers in our staining
panel (e.g., CD3: 46.5%; CD68: 42.2%; αSMA: 52.2% of cases). Altogether, IL-6 was
commonly expressed in human PDAC and exhibited a spatially heterogeneous expression
pattern.
184
185 **IL-6^{high} PDAC patients show distinct myeloid, endothelial and ECM properties but little**
186 **involvement of lymphoid cells**
187
Bioimaging data from TMAs or other multiple regions of interest (ROI)-based methods
are typically collapsed to the patient level, i.e., expression levels from multiple sampled regions
are averaged to achieve robust patient stratification (Fig. 4A). This common analysis standard
was implemented in Hourglass and served as entry point for investigation of the functions of
IL-6 in PDAC. First, we compared PDAC patients by IL-6 expression quantiles (Fig. S3A)
across our full stain panel (Fig. 4B) to get an overview of the involvement of IL-6 in human
PDAC disease biology.
195
At the patient level (Fig. 4C), IL-6 showed no significant associations with malignant
cell behavior yet was strongly linked to changes in the TME. Specifically, high IL-6 levels were
associated with increased vascularization, lower ECM content, and increased presence of the
neutrophil marker CD15, monocyte marker CD11b and macrophage marker CD68 but not the
M2 phenotype marker CD206. In contrast, CAF markers such as αSMA and FAPα were not
significantly altered and neither were T cell markers (CD3, CD4, CD8, FoxP3) nor checkpoint
molecules (PDL1, IDO1). Accordingly, composite analysis of all myeloid and all lymphoid
markers confirmed an overall enrichment of myeloid but not lymphoid cells in human PDAC
tumors along with increasing IL-6 expression (Fig. 4D). Inspection of the individual datapoints
further showed a clean dose-dependent associations with IL-6 for the following markers:
CD15, CD11b and CD68 (higher along with IL-6; Fig. 4E), total collagen content and number
of collagen fibers (lower with high IL-6; Fig. 4F), and vascularization (CD31, HIF1; higher along
with IL-6; Fig. 4G). The B cell marker CD20 showed a weak, non-dose-dependent association
as it was lower in IL-6^{hi} compared to IL-6^{Int} patients (Fig. S3B). Altogether, IL-6^{hi} PDAC tumors
exhibited increased myeloid cell population and vascularization along with reduced ECM
content.
212
Intriguingly, none of the T cell marker or checkpoint molecule stains showed any
association with IL-6 levels and none except PDL1 even trended along with IL-6^{hi} levels (Fig.
4H). This was unexpected given the well-established and central role of IL-6 in T cell
functionality, expansion and survival in a variety of pathological contexts and its shown
benefits in combination with immunotherapy (16,19,25–28). We thus performed a targeted
manual analysis to count tumor infiltrating CD8+ lymphocytes (CD8 TILs) in direct physical
contact with malignant epithelial cells, which is indicative of the presence of functional CD8
effector T cells. This indeed revealed a small but significant increase in CD8+ TILs in IL-6^{hi}
PDAC patients (Fig. 4I). In light of the known regional heterogeneity in the intratumoral T cell
response in PDAC (10) we decided to explore this observation further in the regional analysis
formats that are supported by Hourglass.
224
225 **Intratumoral heterogeneity-informed Hourglass workflow links IL-6 expression to**
226 **regional T cell enrichment**
227

To complement the patient-level analysis (Fig. 4A), Hourglass was designed to support an otherwise identical analysis in parallel at the individual sample level (*i.e.,* by-sample (Fig. 5A). Notably, in this regional analysis, IL-6^{hi} samples showed a tendency to spontaneously group together across all stains (Fig. 5B), suggesting IL-6 is overall involved in tissue composition. As before at the patient level (Fig. 4C), high regional IL-6 was strongly linked to differences in TME composition, associating with increased myeloid cells and endothelium, and decreased ECM content (Fig. 5C). Only the B cell marker CD20 did not associate with regional IL-6 levels (Fig. S4). Beyond these differences, this analysis level revealed higher numbers of FAP+ CAFs, as well as an increase in CD3+, CD8+ cells and checkpoint molecules IDO-1 and PD-L1 in IL-6^{hi} regions in human PDAC. Accordingly, composite analysis of the myeloid and lymphoid cells showed an overall enrichment of both immune compartments in regions with high IL-6 expression in human PDAC (Fig. 5D), which had not been apparent in the global analysis (Fig. 4D). Inspection of the individual datapoints also showed the same clean expression-dependent associations with myeloid markers (CD11b, CD15, CD68; Fig. 5E), ECM content (Fig. 5F) and vascularization (CD31; Fig. 5G) as before. In addition, there was a clear association of regional IL-6 levels with CD3+ and CD8+ T cells as well as PDL-1 and IDO1 (Fig. 5H) and a prominent increase in CD8 TIL counts (Fig. 5I).

To test if these additional and/or stronger trends were merely a reflection of increased sample numbers, we compared paired IL-6^{hi} vs IL-6^{lo} regions within the same tumors, leveraging the presence of IL-6^{hi} and IL-6^{lo} cores from the same patients in our TMA (Fig. 3B, Fig. 5J *left*). This showed that regional differences in intratumoral IL-6 expression trend to be directly accompanied by regional enrichments in myeloid cells, endothelial cells, FAPα+ (but not αSMA+) CAFs, tumor-infiltrating CD8+ T cells, and a decrease in ECM (Fig. 5J). Altogether, regional variation in IL-6 expression appears to have distinct consequences for local TME composition, which can be masked when averaging regions of various expression within patients.

Systematic comparison of IL-6 effects between patient subsets identifies sex-specific immune phenotypes in PDAC

PDAC is clinically heterogeneous and biological patterns might vary between patient subgroups. Iterative refinement of analyses to identify confounder subsets can thus be instrumental to understand underlying biology. For instance, molecular subtypes of PDAC (29) differ in their baseline immune microenvironments (30,31) while IL-6 itself has been implicated in sex-differences in the context of hepatocellular carcinoma (32). To test whether the involvement of IL-6 in key biological patterns (immune phenotypes, TME composition) was confounded within specific PDAC patient subsets, we applied parallel subgroup comparison under otherwise identical analytical conditions across global and regional analysis levels, leveraging all steps in the Hourglass workflow (Fig. 6A).

While IL-6 expression levels were the same in male and female patients (Fig. S5A), we noticed that male patients trended towards worse survival outcomes as a function of high IL-6 expression (Fig. 6B). At the global level (all samples averaged by patient), this was accompanied by similar myeloid cell involvement in male vs female IL-6^{hi} patients (Fig. 6C, *heatmaps*) but a significant increase in both CD3 TILs and CD8 TILs that was absent in female patients (Fig. 6C, *boxplots*), despite a trend towards higher CD8 TILs that remained non-

significant. Strikingly, at the regional level (samples divided by IL-6 expression levels), we again observed a consistent increase of T cell markers (CD3, CD8, CD4, FoxP3) and checkpoint molecules (PDL1, IDO1) as well as CD3 TILs and CD8 TILs in IL-6^{hi} PDAC regions yet this was entirely restricted to male patients (Fig 5H-I vs. Fig. 6D). This indicates the presence of a male-specific regional T cell response linked to IL-6 expression levels in human PDAC. Moreover, when we compared the TME composition of paired intratumoral IL-6^{hi} and IL-6^{lo} regions between sexes (Fig. 6E), we observed larger scale differences: in male patients, IL-6^{hi} compared to IL-6^{lo} regions exhibited more CD3 T cells, more CD11b myeloid cells, more FAPα+ CAFs, and less ECM content. In female patients, on the other hand, we observed unaltered CD3 T cells, and unaltered FAP+ CAFs, while the reduced ECM content and the increase in CD11b myeloid cells were consistent between both sexes. Altogether, this suggests that regional variations in IL-6 levels within human PDAC lead to spatial effects on TME composition that are in part sex-specific, namely with male TMEs being overall more inflamed and more CAF-activated in the presence of IL-6.

IL-6 effects vary only slightly between molecular subtypes of PDAC

We then proceeded to test whether there are differences in the IL-6 response between PDAC basal-like and classical molecular subtypes (29), since we had noticed that IL-6 expression was higher in classical than in basal-like PDAC tumors (Fig. S6B). Patients with basal-like tumors exhibited significantly worse survival outcomes as a function of high IL-6 expression (Fig. S6C). At the global level (all samples averaged by patient), this was accompanied with a much less pronounced myeloid involvement in IL-6^{hi} regions of basal-like than of classical PDAC tumors (Fig. S6D, *heatmaps*). We in turn observed higher CD3 TILs and CD8 TILs in IL-6^{hi} regions of basal PDAC tumors (Fig. S6D, *boxplots*) even though classical PDACs trended in the same direction. Yet, at the regional level (samples by IL-6 expression levels), both basal and classical tumors exhibited similar differences in response to IL-6 across lymphoid and myeloid markers as well as CD3 TILs and CD8 TILs (Fig. S6E). Likewise, when we compared the TME composition of paired IL-6^{hi} and IL-6^{lo} regions within tumors of different subtypes (Fig. S6F), we observed very similar effects across both subtypes: unaltered T cells (CD3), increase FAP+ CAFs and decreased ECM; only myeloid cells (CD11b) were more strongly associated with IL-6 levels in the classical subset of PDAC patients.

Altogether, this systematic analysis of IL-6 in human PDAC through the Hourglass computational toolkit has revealed that a) T cell responses in PDAC were linked to IL-6 expression levels but largely occur as a regional phenomenon, b) regional IL-6 dependent T cell responses in human PDAC were largely restricted to male patients and c) higher IL-6 expression was associated with worse survival outcomes in basal-like subset of PDAC patients , which was however not directly linked to major differences covered by our tumor biology staining panel.

Discussion

Technological advances in biomedical research are producing immense amounts of data and the resultant downstream analysis can become a formidable undertaking. To address this obstacle, we constructed a framework for systematic interrogation of multiparametric datasets. The software package, called Hourglass, is informed by tissue heterogeneity and

able to strategically process multiparametric datasets with a focus on functionalities that are
frequently required for bioimaging datasets, such as assembly of feature subsets and regional
versus global analysis levels. We showcase its utility for knowledge extraction from a PDAC
TMA dataset, where it derives novel insights into IL-6-related regional biology and the
identification of clinically relevant patient subsets in the context of future IL-6 directed
immunotherapeutic strategies for PDAC.
328
Digital image analysis, as opposed to semi-quantitative scoring, is becoming a
common standard and has significant benefits in terms of larger scale experiments such as a
wide staining panel applied to a TMA, normalization and implementation on analysis standards
across labs and institutions, and cost-effectiveness. (33–36) Hourglass was designed as a
user-friendly and robust analysis tool to systematically and reproducibly mine large bioimaging
datasets in the rapidly expanding digital pathology space. Naturally, alongside the boom of big
data in biomedical and clinical spaces, tools are constantly created to query public and private
datasets. However, there are very limited tools that facilitate complex computational bioimage
analysis in a streamlined approach. This is due to several reasons, but most notably, the
majority of intuitive and/or interactive tools are geared towards omics analysis and
consequently provide little to no support for complex multiparametric data structures. At the
same time, computational experts in the bioimaging space are mostly engaged in rather high-
end artificial intelligence-based analytical approaches (36–38), which are undoubtedly
extremely powerful but not well-suited to generate software solutions suitable for systematic
data mining of standard bioimaging applications. Consequently, we hope that Hourglass will
particularly appeal to non-expert users and encourage more iterative and heterogeneity-
informed mining of large bioimaging dataset with extensive annotations, as it supports
customizable, iterative, and reproducible data exploration via a streamlined and semi-
automated workflow.
348
Careful analysis of multiparametric bioimaging datasets bears specific demands and
challenges, such as a high need for quality control across sample due to sample dropouts
and/or staining artifacts. In addition, with multiple options for how to analyze different stains
(cell-based versus pixel-based metrics, weighted scoring such as allred or H-scores), the
ability to systematically compare and benchmark multiple parameters for each stain and select
the appropriate one is essential for optimal analyses. Yet, this becomes laborious and error-
prone if done manually across multiple stains. Hourglass has thus been tailored to address
specific needs of both multiparametric data (manual parameter selection, parallel comparison
under otherwise identical conditions even within larger stain panels) and bioimaging datasets
(integration and subsetting of large stain panels, heterogeneity-informed navigation of regional
and global levels, quality control across all parameters per stain). In addition, Hourglass
supports various required data post-processing steps that might be particularly challenging for
computational non-expert users. For example, to account for dropouts (*e.g.,* insufficient
staining quality, cores lost at washing steps, tissue folds), Hourglass can perform imputation
of missing data to then facilitate downstream visualization per unsupervised clustering in
heatmaps and correlation analyses. Based on user specifications, Hourglass then aggregates
and visualizes data subsets and applies statistical testing. Outputs are structured in a coherent
file hierarchy and analytical specifications are logged, to facilitate iterative hypothesis
generation and testing. We are to the best of our knowledge not aware of any other software
applications that provide such a comprehensive toolkit tailored towards relevant functionalities
for bioimaging and multiparametric datasets to date.

370
Tissue heterogeneity and regional cell communities are increasingly understood as a relevant
disease-driving features rather than noise (10,11). Nevertheless, the most common approach
to preparing bioimaging data from TMAs is to average over multiple samples or ROIs per
patient while a whole wealth of information may be contained within those multiple individual
regions per tumor. Hourglass was designed to access all levels of this unique data structure.
Specifically, we harness this standard redundant sampling approach taken in TMAs and other
multi-ROI based bioimaging approaches to systematically dissect regional from inter-patient
heterogeneity and generate a new level of insight into tissue biology from the otherwise
unused or discarded information. In essence, Hourglass incorporates different analytical levels
in its heterogeneity-informed approach by supporting parallel analysis of patients, individual
samples, and patient-paired samples compared as a function of a parameter of interest (here,
IL-6 expression). We showcase how such systematic dissection of intra- and intertumoral
heterogeneity can lead to new insights into disease-relevant traits and clinically relevant IL-6
biology. Notably, these different levels of analysis can be used alone or in combination, *i.e.*
users can focus on the standard by-patient analysis only or perform the analysis under
identical settings at the global, regional and patient-paired levels to systematically identify
commonalities and differences. Altogether, we strongly recommend exploring all levels first
and then decide based on the obtained results. For instance, if regional differences are driving
global trends, as we have observed for IL-6 in male vs female PDAC case, then heterogeneity
or sampling bias is highly relevant in this context. On the other hand, when regional analyses
do not provide additional insight such as in the case IL-6 in the context of basal-like vs classical
PDAC, subsetting may simply not be warranted for this specific research question.
393
IL-6 is a key therapeutic target, not only in PDAC but many inflammation-related diseases (39)
and affects essentially all aspects of malignant biology. Yet, there are numerous conflicting
reports as to whether IL-6 levels are a prognostic indicator in PDAC and other cancers, and
these varying outcomes suggest significant clinical heterogeneity in the effects of IL-6. In
addition, IL-6 expression is known to be regionally heterogeneous (19,24), posing challenges
for sampling and due to spatial confinement of key biological associations that might be
missed in current standard approaches. By systematically navigating through all these layers
of complexity, Hourglass revealed novel, largely sex-specific, spatial effects of IL-6 expression
on TME composition in human PDAC. In specific, we found that T cells, as opposed to the
major myeloid cell populations, responded to regional variation in IL-6 expression, and these
were masked when averaging multiple regions of various expression within all patients.
Moreover, this regional effect of IL-6 on the T cell compartment was almost entirely restricted
to male patients. Notably, we also observed that FAP+ fibroblasts, another key stromal cell
population with presumed immune-modulatory functions (40) were also restricted to these IL-
6^{hi} regions in male patients only. In turn, there was a consistent regional and global
anticorrelation of IL-6 with the amount of ECM across sexes and subtypes. Notably, CD8 TILs
were the only T cell subset that fairly consistently aligned with IL-6 levels across multiple
patient subsets, hinting at a more robust link between IL-6 and CTL functionality. Altogether,
this exposes novel associations of IL-6 with TME biology in human PDAC, sex-disparities
within the lymphoid arm of the intratumoral immune response and provides a systematic
roadmap for navigating such complex phenomena across spatial levels and patient subsets.
The here-discovered sex disparities in the IL-6 related intratumoral T cell response in PDAC
have furthermore major implications for the design of immunotherapeutic strategies and the

selection of patients for such new treatments. Moreover, such disparities in patient subgroups might in part be responsible for the conflicting clinical correlates of IL-6.

References

1. Barsoum I, Tawedrous E, Faragalla H, Yousef GM. Histo-genomics: digital pathology at the forefront of precision medicine. Diagn Berl Ger. 2019 Aug 27;6(3):203–12.

2. Li X, Ma Z, Zhu YZ. Regional Heterogeneity of Perivascular Adipose Tissue: Morphology, Origin, and Secretome. Front Pharmacol [Internet]. 2021 [cited 2022 Jul 30];12. Available from: https://www.frontiersin.org/articles/10.3389/fphar.2021.697720

3. Goldstein NS, Hastah F, Galan MV, Gordon SC. Fibrosis heterogeneity in nonalcoholic steatohepatitis and hepatitis C virus needle core biopsy specimens. Am J Clin Pathol. 2005 Mar;123(3):382–7.

4. Böttcher C, Schlickeiser S, Sneeboer MAM, Kunkel D, Knop A, Paza E, et al. Human microglia regional heterogeneity and phenotypes determined by multiplexed single-cell mass cytometry. Nat Neurosci. 2019 Jan;22(1):78–90.

5. Süß P, Hoffmann A, Rothe T, Ouyang Z, Baum W, Staszewski O, et al. Chronic Peripheral Inflammation Causes a Region-Specific Myeloid Response in the Central Nervous System. Cell Rep. 2020 Mar 24;30(12):4082-4095.e6.

6. Manini I, Caponnetto F, Dalla E, Ius T, Della Pepa GM, Pegolo E, et al. Heterogeneity Matters: Different Regions of Glioblastoma Are Characterized by Distinctive Tumor-Supporting Pathways. Cancers. 2020 Oct 13;12(10):E2960.

7. Wang Y, Li X, Peng S, Hu H, Wang Y, Shao M, et al. Single-Cell Analysis Reveals Spatial Heterogeneity of Immune Cells in Lung Adenocarcinoma. Front Cell Dev Biol [Internet]. 2021 [cited 2022 Jul 30];9. Available from: https://www.frontiersin.org/articles/10.3389/fcell.2021.638374

8. Vitale I, Shema E, Loi S, Galluzzi L. Intratumoral heterogeneity in cancer progression and response to immunotherapy. Nat Med. 2021 Feb;27(2):212–24.

9. Iacobuzio-Donahue CA, Litchfield K, Swanton C. Intratumor heterogeneity reflects clinical disease course. Nat Cancer. 2020 Jan;1(1):3–6.

10. Grünwald BT, Devisme A, Andrieux G, Vyas F, Aliar K, McCloskey CW, et al. Spatially confined sub-tumor microenvironments in pancreatic cancer. Cell. 2021 Oct 28;184(22):5577-5592.e18.

11. Jackson HW, Fischer JR, Zanotelli VRT, Ali HR, Mechera R, Soysal SD, et al. The single-cell pathology landscape of breast cancer. Nature. 2020;578(7796):615–20.

12. Heindl A, Nawaz S, Yuan Y. Mapping spatial heterogeneity in the tumor microenvironment: a new era for digital pathology. Lab Invest. 2015 Apr;95(4):377–84.

13. Kumari N, Dwarakanath BS, Das A, Bhatt AN. Role of interleukin-6 in cancer progression and therapeutic resistance. Tumor Biol. 2016 Sep 1;37(9):11553–72.

14. Jones SA, Jenkins BJ. Recent insights into targeting the IL-6 cytokine family in inflammatory diseases and cancer. Nat Rev Immunol. 2018 Dec;18(12):773–89.

15. McAndrews KM, Chen Y, Darpolor JK, Zheng X, Yang S, Carstens JL, et al. Identification of Functional Heterogeneity of Carcinoma-Associated Fibroblasts with Distinct IL6-Mediated Therapy Resistance in Pancreatic Cancer. Cancer Discov. 2022 Jun 2;12(6):1580–97.

16. Bent EH, Millán-Barea LR, Zhuang I, Goulet DR, Fröse J, Hemann MT. Microenvironmental IL-6 inhibits anti-cancer immune responses generated by cytotoxic chemotherapy. Nat Commun. 2021 Oct 28;12(1):6218.

17. Lesina M, Wörmann SM, Neuhöfer P, Song L, Algül H. Interleukin-6 in inflammatory and malignant diseases of the pancreas. Semin Immunol. 2014 Feb;26(1):80–7.

18. Wörmann SM, Diakopoulos KN, Lesina M, Algül H. The immune network in pancreatic cancer development and progression. Oncogene. 2014 Jun;33(23):2956–67.

19. Mace TA, Shakya R, Pitarresi JR, Swanson B, McQuinn CW, Loftus S, et al. IL-6 and PD-L1 antibody blockade combination therapy reduces tumour progression in murine models of pancreatic cancer. Gut. 2018 Feb 1;67(2):320–32.

20. Ware MB, McQuinn C, Zaidi MY, Knochelmann H, Mace TA, Chen Z, et al. Dual blockade of IL-6 and CTLA-4 regresses pancreatic tumors in a CD4+ T cell-dependent manner [Internet]. bioRxiv; 2020 [cited 2022 Aug 2]. p. 2020.02.07.939199. Available from: https://www.biorxiv.org/content/10.1101/2020.02.07.939199v1

21. Lin JH, Huffman AP, Wattenberg MM, Walter DM, Carpenter EL, Feldser DM, et al. Type 1 conventional dendritic cells are systemically dysregulated early in pancreatic carcinogenesis. J Exp Med. 2020 Aug 3;217(8):e20190673.

22. Yang F, He Z, Duan H, Zhang D, Li J, Yang H, et al. Synergistic immunotherapy of glioblastoma by dual targeting of IL-6 and CD40. Nat Commun. 2021 Jun 8;12(1):3424.

23. Liu C, Yang L, Xu H, Zheng S, Wang Z, Wang S, et al. Systematic analysis of IL-6 as a predictive biomarker and desensitizer of immunotherapy responses in patients with non-small cell lung cancer. BMC Med. 2022 May 13;20(1):187.

24. Soares-Lima SC, Gonzaga IM, Camuzi D, Nicolau-Neto P, Vieira da Silva R, Guaraldi S, et al. IL6 and BCL3 Expression Are Potential Biomarkers in Esophageal Squamous Cell Carcinoma. Front Oncol. 2021;11:3102.

25. Yan Y, Ramanan D, Rozenberg M, McGovern K, Rastelli D, Vijaykumar B, et al. Interleukin-6 produced by enteric neurons regulates the number and phenotype of microbe-responsive regulatory T cells in the gut. Immunity. 2021 Mar 9;54(3):499-513.e5.

26. Jiang Z, Liao R, Lv J, Li S, Zheng D, Qin L, et al. IL-6 trans-signaling promotes the expansion and anti-tumor activity of CAR T cells. Leukemia. 2021 May;35(5):1380–91.

27. Weber R, Groth C, Lasser S, Arkhypov I, Petrova V, Altevogt P, et al. IL-6 as a major regulator of MDSC activity and possible target for cancer immunotherapy. Cell Immunol. 2021 Jan;359:104254.

28. Long KB, Tooker G, Tooker E, Luque SL, Lee JW, Pan X, et al. IL6 Receptor Blockade Enhances Chemotherapy Efficacy in Pancreatic Ductal Adenocarcinoma. Mol Cancer Ther. 2017 Sep 1;16(9):1898–908.

29. Moffitt RA, Marayati R, Flate EL, Volmar KE, Loeza SGH, Hoadley KA, et al. Virtual
microdissection identifies distinct tumor- and stroma-specific subtypes of pancreatic
ductal adenocarcinoma. Nat Genet. 2015 Oct;47(10):1168–78.

30. Raghavan S, Winter PS, Navia AW, Williams HL, DenAdel A, Lowder KE, et al.
Microenvironment drives cell state, plasticity, and drug response in pancreatic cancer.
Cell. 2021 Dec 9;184(25):6119-6137.e26.

31. Storrs EP, Usmani A, Chati P, Sloan I, Krasnick BA, Babbra R, et al. High-dimensional
deconstruction of pancreatic ductal adenocarcinoma identifies tumor
microenvironmental communities associated with survival [Internet]. medRxiv; 2022
[cited 2022 Aug 2]. p. 2022.04.29.22274376. Available from:
https://www.medrxiv.org/content/10.1101/2022.04.29.22274376v2

32. Prieto J. Inflammation, HCC and sex: IL-6 in the centre of the triangle. J Hepatol. 2008
Feb 1;48(2):380–1.

33. Aeffner F, Zarella MD, Buchbinder N, Bui MM, Goodman MR, Hartman DJ, et al.
Introduction to Digital Image Analysis in Whole-slide Imaging: A White Paper from the
Digital Pathology Association. J Pathol Inform. 2019;10:9.

34. Bankhead P. Developing image analysis methods for digital pathology. J Pathol. 2022
Jul;257(4):391–402.

35. Hamilton PW, Bankhead P, Wang Y, Hutchinson R, Kieran D, McArt DG, et al. Digital
pathology and image analysis in tissue biomarker research. Methods San Diego Calif.
2014 Nov;70(1):59–73.

36. Jahn SW, Plass M, Moinfar F. Digital Pathology: Advantages, Limitations and Emerging
Perspectives. J Clin Med. 2020 Nov 18;9(11):E3697.

37. Madabhushi A, Lee G. Image analysis and machine learning in digital pathology:
Challenges and opportunities. Med Image Anal. 2016 Oct;33:170–5.

38. Wang S, Yang DM, Rong R, Zhan X, Xiao G. Pathology Image Analysis Using
Segmentation Deep Learning Algorithms. Am J Pathol. 2019 Sep;189(9):1686–98.

39. Hirano T. IL-6 in inflammation, autoimmunity and cancer. Int Immunol. 2021 Mar
1;33(3):127–48.

40. Mhaidly R, Mechta-Grigoriou F. Role of cancer-associated fibroblast subpopulations in
immune infiltration, as a new means of treatment in cancer. Immunol Rev. 2021
Jul;302(1):259–72.

Figure Legends

<u>Main Figures</u>

534 **Figure 1.** **Hourglass workflow for systematic mining of annotated multiparametric**
535 **bioimaging datasets**

Schematic: **(A)** Multiparameteric dataset acquisition from digital bioimaging of TMA from well-annotated patient cohort, **(B)** data processing via Hourglass, and **(C)** user-selected output.

Figure 2. Graphical user interface supports navigation through Hourglass workflow
GUI tab layout supporting **(A)** data import **(B)** customization of colors for graphs, and **(C)** user-selected options.
(D) Key steps in analysis pipeline corresponding to tabs in GUI.
(E) Run button logs user-defined data file paths and choices to Excel file and submits to R package to create output folder via R functions and dependencies.

Figure 3. Inter- and intra-tumoral heterogeneity of IL-6 expression levels in PDAC
(A) Representative images of human PDAC tumors showing varying intertumoral (left) and intratumoral (right) IL-6 levels; arrowheads indicate IL-6 low (*blue*) vs. high (*red*) regions.
(B) Varying IL-6 expression within different TMA cores from four representative PDAC tumors.

Figure 4. IL-6^{hi} PDAC patients show distinct myeloid, endothelial and ECM properties but little involvement of lymphoid cells
(A) Schematic: Hourglass can average and compare IL-6 expression over multiple samples/regions of interest (ROIs) per patient.
(B) Staining panel used in this study.
(C) Heatmap of fold-change expression and significance levels for all stains in (**B**) across IL-6^{hi} (hi), IL-6^{int} (int) and IL-6^{lo} (lo) expressing human PDAC tumors. SHG1: Mean Col Width, SHG2: Mean Col Straightness, SHG3: Mean Col Length, SHG4: Col Alignment, SHG5: Number Col (where Col is Collagen Fiber).
(D) Composite plot of all myeloid (*top*) and lymphoid (*bottom*) marker quantification results per patient grouped by IL-6 levels (hi/left, int/middle, lo/right).
(E) Individual digital quantification results of IHC stains for CD11b, CD15, CD68, CD206 (myeloid markers), **(F)** SHG1, MT (ECM), **(G)** CD31 (vascularization), **(H)** CD3, CD8, CD4, FOXP3, PD-L1 (lymphoid), as well as manual counts of **(I)** tumor infiltrating CD8+ lymphocytes (CD8+ TILs). **(F)** Number of collagen fibers as assessed through second harmonic generation microscopy. **(E-I)** Boxplots compare IL-6^{high} (hig), IL-6^{int} (int) and IL-6^{low} (low) patients. **(C, E-I)** n = 143; Wilcoxon test. *p < 0.05, **p < 0.01, ***p < 0.001,****p < 0.0001.

Figure 5. Intratumoral heterogeneity-informed Hourglass workflow links IL-6 expression to regional T cell enrichment
(A) Schematic: Hourglass can compare IL-6 expression in individual samples/ROIs.
(B) Heatmap visualizing hierarchal clustering of markers in staining panel in all samples (n= 516), grouped by IL-6 level.
(C) Heatmap of fold-change expression and significance levels for all stains across IL-6^{hi} (hi), IL-6^{int} (int) and IL-6^{lo} (lo) samples. SHG1: Mean Col Width, SHG2: Mean Col Straight, SHG3: Mean Col Length, SHG4: Col Alignment, SHG5: Number Col (where Col is Collagen Fiber).
(D) Composite plot of all myeloid (*top*) and lymphoid (*bottom*) marker quantification results per sample grouped by IL-6 levels (hi/left, int/middle, lo/right).

(E) Individual digital quantification results of IHC stains for CD11b, CD15, CD68, CD206 (myeloid markers), **(F)** SHG1, MT (ECM), **(G)** CD31 (vascularization), **(H)** CD3, CD8, CD4, FOXP3, PD-L1 (lymphoid), as well as manual counts of **(I)** tumor infiltrating CD8+ lymphocytes (CD8+ TILs). **(F)** Number of collagen fibers as assessed through second harmonic generation microscopy. **(E-I)** Boxplots compare IL-6^{high} (hig), IL-6^{int} (int) and IL-6^{low} (low) samples. **(J)** Schematic: Hourglass can compare difference IL-6 levels within the same patient (left); Patient-paired slope-graphs show CD11b, CD68 (myeloid), FAPα (CAF), MT (ECM) CD31 (vascularization), and CD8+ TILS levels in IL-6^{high}+ IL-6^{int} (hig) and IL-6^{low} (low) samples within the same patient (right).
(C, E-J) n = 516.Wilcoxon test. *p < 0.05, **p < 0.01, ***p < 0.001,****p < 0.0001.

Figure 6. **Systematic comparison of IL-6 effects between patient subsets identifies sex-specific immune phenotypes in PDAC**
(A) Schematic: Hourglass systematically dissects observations across patient subgroups within user-defined biological sets of features/stains.
(B) Kaplan-Meier analysis of overall survival time in all patients (left), and basal-like (middle), classical (right) sub-cohorts; strata: IL-6 level; log-rank tests. M, male; F, female;
(C-E) Comparison of IL-6 effects in male (top panels) versus female (bottom panels) PDAC patients across global (**C**), regional (**D**) and patient-paired (**E**) levels. (**C, D**) Heatmaps depicting foldchange expression and significance levels for selected immune markers; boxplots show manual TIL counts. **(E)** Patient-paired slope-graphs depicting CD3 (T cells), CD11b, (myeloid), FAPα (CAF), MT (ECM) and CD8+ TILS levels. **(C-E)** All levels are stratified by IL-6 expression quantiles into IL-6^{hi} (hi), IL-6^{int} (int) and IL-6^{lo} (lo) groups. Wilcoxon test. *p < 0.05, **p < 0.01, ***p < 0.001,****p < 0.0001.

Supplemental Figures

Figure S1. **Quantification metrics for digital IHC images, related to Figure 1**
Where applicable, stains were digitally quantified using QuPath via either or both cell- (*left*) and pixel- (*right*) based detection methods.

Figure S2. **Intratumoral heterogeneity of IL-6 and other stains within patients, related to Figure 3**
(A) IL-6 staining controls (kidney, liver, pancreas); representative images.
(B) Binning of human PDAC samples based on tertiles of IL-6 expression.

Figure S3. **Expression of IL-6 and CD20 across patients, related to Figure 4**
(A) Binning of human PDAC tumors based on tertiles of IL-6 expression.
(B) Digitally quantified CD20 IHC stain results plotted as function of IL-6 expression per tumor.

Figure S4. **Expression of CD20 across samples, related to Figure 5**
Digitally quantified CD20 IHC stain results plotted as function of IL-6 expression per sample.

Figure S5. **IL-6 effects vary only slightly between molecular subtypes of PDAC, related to Figure 6**

Boxplots show expression of IL-6 in patients across **(A)** sexes and **(B)** Moffitt molecular
subtypes.
(C) Kaplan-Meier analysis of overall survival time in all patients (*left*), and basal-like (*middle*),
classical (*right*) sub-cohorts; strata: IL-6 level; log-rank tests.
(D-F) Comparison of IL-6 effects in PDAC patients with basal-like (top panels) versus
classical (bottom panels) transcriptomic subtype across global (**D**), regional (**E**) and patient-
paired (**F**) levels. (**D, E**) Heatmaps depicting foldchange expression and significance levels
for selected immune markers; boxplots show manual TIL counts. **(F)** Patient-paired slope-
graphs depicting CD3 (T cells), CD11b, (myeloid), FAPα (CAF), MT (ECM) and CD8+ TILS
levels. (**D-F**) All levels are stratified by IL-6 expression quantiles into IL-6^{hi} (hi), IL-6^{int} (int)
and IL-6^{lo} (lo) groups. Wilcoxon test. *p < 0.05, **p < 0.01, ***p < 0.001,****p < 0.0001.
641
642

643 # Methods

645 ## Resource Availability

646 **Lead contact**
647 Further information and requests for resources and reagents should be directed to and will be
648 fulfilled by the lead contact, Rama Khokha (rama.khokha@uhnresearch.ca).
649
650 **Materials availability**
651 This study did not generate new materials.

652 # Data and code availability

653 Hourglass source code is available on Github, see https://kazeera.github.io/Hourglass for user
654 manual.
655

Method details

Acquisition of pancreatic cancer tissue microarray with immunohistochemistry
Hourglass was built around the PDAC IHC TMA dataset, presented in (10). Briefly,
pancreatic cancer tissues were obtained from patients, samples were placed as cores
(diameter 1.2 mm) into TMAs, 5 μm lateral sections, and stained for different genes or
features. Pixel-based or cell-based detection parameters were manually set and optimized for
each stain. Quantification results were quality controlled by visual inspection of the 10 cores
with the highest detected staining levels and the 10 cores with the lowest detected staining
levels for each slide for each stain. Next, images of slides were quantified by Numeric readouts
or parameters acquired for each stain for each core, for example the area, number of positive
pixels, etc. Next, a merging script was used to merge all the stain files into one table and
integrated with patient/sample annotations information including age, sex, cancer subtype,
treatment history, mutation statuses. Each region of interest had a unique identifier containing
the core location on the TMA grid, slide, links to other dataset acquisition (e.g. sequencing)
and links to clinical and histopathology. Each patient has a unique case ID and 2-8
samples/regions of interest.
672
Description of bioimaging data

The dataset used in this study consists of 596 cores or tissue samples from 165
patients (i.e. 1 to 15 samples per patient; average 3.6), first published in (10). Each sample
had 26 stains applied, including histochemical stains, immunohistochemical stains, and
second harmonic generation microscopy. Digital quantification of these stains using the
bioimage analysis softwares QuPath and CT-FIRE recorded 22 unique parameters (Table 1).
The final *sample x feature-parameter* matrix dimensions are 648 x 696, resulting in
451,008 data points. In addition, patient data was used to define multiple recorded subgroups,
such as age, cancer subtypes, and treatment history that could be compared (represented as
a clipboard in Fig. 1A). In our dataset, 50 and 59 annotations were labeled for patients and for
samples respectively.

Statistical Analyses
Where appropriate, the quantitative analyses are described in the relevant sections of
the Method details. Statistical and visual analyses were performed using Hourglass and R
(v4.1.0). Quantitative variables were compared by Wilcoxon rank sum test, pairwise in
slopegraphs. Fold-changes were calculated by dividing means of respective groups. Survival
curves were plotted by the Kaplan–Meier method and p values were assessed based on
logrank test. The specific statistical tests used are indicated in the figure legends. All tests
were two-sided. Statistical significance was set at $p<0.05$. Correlation coefficients were
determined by the Spearman method.

Development of R package
The standalone Hourglass R package was built using R v4.1.0. We compiled a number
of existing data filtering functions (such as subset retrieval, imputation and outlier removal)
and plotting functions and wrote our own.

Development of graphical user interface
The Kivy design language (v2.0.0) and pyinstaller (v4.8) in python3 was used to create
a standalone executable.

Details of IL-6 Hourglass run
For this study, we customized an Hourglass analysis to interrogate the role of IL-6 in
PDAC with 8 main comparisons (comparing subgroups in samples and patients; e.g. looking
at IL-6 in basal vs classical tumors), 22 feature sets (grouping stains; e.g. vascularization,
immune, epithelial behavior) and 10 plot types. In parallel, Hourglass imputed and ran the
same comparisons on this version of the dataset. Hourglass ran as a background sequential
process for 7.393 hours on a Windows System (32GB RAM, Intel (Alt-0174) $^®$ Core™(Alt-
0153) i7 CPU). The result was an output folder with a total of 17,386 PDF files contained in
subfolders with relevant biological analysis, amounting to 3.40 GB (3,401,516,890 bytes) of
publication-ready plots.

Acknowledgments

717 This work was supported by grants to RK from CCS and CIHR. BTG held fellowships from the
718 Princess Margaret Foundation, EMBO (ALTF-116-2018), Alexander von Humboldt

719 Foundation (DEU-1199182-FLFP), and CRS (Next Generation of Scientists Scholarship). We
720 thank Parinaz Nasr Esfahani for supporting software development.
721
722 The authors declare no competing financial interests.
723

Author contributions

725 K.A. and B.T.G. conceptualized the study. K.A., H.R.W., and B.Z. designed and implemented
726 the software tool. F.V., E.P., N.C., N.K., and B.T.G. performed data analysis. K.A. and B.T.G.
727 prepared the final visualizations. B.T.G. and K.A. wrote the manuscript. All authors proofread
728 the manuscript. B.T.G. and R.K. supervised the project.
729

Declaration of interests

731 The authors declare no competing interests.
732

A Dataset aggregation

Clinical & Molecular Annotations + Staining and Imaging Samples → Digital Image Analysis → Bioimaging Data

Multiparametric data + annotations

B Data processing

Input

Hourglass

Defining analysis setting
Defining comparisons and subgroups
Selecting visualization options
Selecting Statistics

Graphs

— Sample
— Patient
— Comparison1
— Comparison2
— QC
— Set1
— Set2

Output

Visualizations + log files

C Output mining & Quality Control

Visualization Portfolio

Figure 1

Figure 2

Human PDAC

Intertumoral Heterogeneity

Intratumoral Heterogeneity

IL-6 IHC

B

	Patient #1	Patient #2	Patient #3	Patient #4

IL-6 expression

hi

lo

Figure 3

Figure 4

Figure 5

Figure 6