Collective protection against the type VI secretion system in bacteria

Elisa T. Granato*[1,2], William P. J. Smith* [1,2,3], Kevin R. Foster[1,2]

*Equal contribution

[1] Department of Biology, University of Oxford, Oxford, UK

[2] Department of Biochemistry, University of Oxford, Oxford, UK

[3] Division of Genomics, Infection and Evolution, University of Manchester, UK

ABSTRACT

Bacteria commonly face attacks from other strains via the type VI secretion system (T6SS), a
molecular speargun that stabs and intoxicates competitors. Here we show how bacteria can work
together to collectively defend themselves against these attacks. This project began with an
outreach activity: while developing an online computer game of bacterial warfare, we noticed that
one strategist ("Slimy") that made extracellular polymeric substances (EPS) was able to resist
attacks from another strategist that employed the T6SS ("Stabby"). This observation motivated
us to model this scenario more formally, using dedicated agent-based simulations. The model
predicts that EPS production can serve as a collective defence mechanism, which protects both
producing cells and neighbouring cells that do not make EPS. We then tested our model with a
synthetic community that contains a T6SS-wielding attacker (*Acinetobacter baylyi*), and two
T6SS-sensitive target strains (*Escherichia coli*) that either secrete EPS, or not. As predicted by
our modelling, we find that the production of EPS leads to collective protection against T6SS
attacks, where EPS producers can shield each other and nearby non-producers. Our work shows
how EPS-producing bacteria can work together to defend themselves from attackers.

INTRODUCTION

Bacteria commonly live in densely-populated, multispecies communities where they must compete for limited space and nutrients (1–3). Life in these environments has driven the evolution of an array of bacterial weapon systems, used to inhibit or kill competing microbes (4, 5). One of the most widespread and sophisticated of these weapons is the type VI secretion system (T6SS), a contractile nanomachine that fires a needle into neighbouring bacteria to deliver toxic effector proteins (6).

Possession of the T6SS can confer a strong ecological advantage to bacteria cells (7–10). Conversely, being on the receiving end of a T6SS attack can be lethal. Defending against the T6SS, however, presents a unique set of problems for targeted cells, as—unlike many other weapon systems—the T6SS does not rely on the target cell's surface receptors or transport systems. Instead, it uses mechanical force to pierce the target cell's membrane and deliver its toxic cargo, which limits the potential range of mechanisms that can be employed in defence (11).

In the face of these challenges, here we show how bacteria can work together to protect themselves against T6SS attacks. Individual-level defence mechanisms are known, including immunity proteins, capsule formation, target modification, envelope stress responses, and the removal of susceptibility factors (12–16). However, some of the most powerful defences in bacteria come about when bacteria act as a collective, allowing groups of cells to take advantage of their large numbers (17–22).

This project began when we serendipitously identified a potential collective defence mechanism against the T6SS while developing a video game, as part of an outreach activity. The game—*Gut Wars*—uses an agent-based model developed by our group for research (17, 23–25) to illustrate different bacterial warfare strategies in the human gut microbiome. Running the game repeatedly, we noticed that "Slimy" strategists, which secrete extracellular polymeric substances (EPS), were particularly resistant against "Stabby" competitors armed with the T6SS. Secretion of EPS is widespread among bacteria and plays a key role in the formation of biofilms (26, 27). Unlike typical anti-T6SS defence mechanisms that act at the level of the individual cell, we saw that EPS released into the environment has the potential to protect both the cell that produces it and others around it, thereby functioning as a powerful collective defence mechanism. There is evidence that EPS has the potential to be physically resistant to T6SSs (12, 13, 15). Moreover, biofilm formation—involving EPS secretion—is known to reduce T6SS susceptibility (14), and is a

response to bacterial competition via T6SS (28). We, therefore, decided to investigate whether
EPS allows bacteria to work collectively in the face of T6SS attacks.

Here, we develop agent-based models of the competitions between T6SS attackers and EPS-
producing target cells. As suggested by the computer game, these analyses predict that secreting
EPS is a collective defence mechanism in which producer cells both protect each other, and also
cross-protect nearby non-producing cells. We test our predictions using a synthetic community,
in which T6SS-wielding *Acinetobacter baylyi* are competing with *Escherichia coli* strains secreting
free (i.e. non-membrane-bound) EPS in amounts comparable to clinical *E. coli* isolates. These
experiments revealed that EPS can indeed serve as a powerful collective defence against T6SS
attacks.

RESULTS

Agent-based modelling predicts that EPS can protect against T6SS
Bacteria have evolved a wide repertoire of traits and strategies to increase their survival chances
in densely-populated communities. We recently devised a simple browser-based computer
game—*Gut Wars*—to showcase some of these competition strategies in a format accessible to
children as well as adults. The game asks the user to select different characters representing
distinct bacterial competition strategies, which are then pitted against each other in a simulated
gut microbiome environment using agent-based modelling (Fig. 1A). Two of the available
strategies are EPS production ("Slimy") and T6SS firing ("Stabby"), both common traits in
bacteria. While not originally implemented to fulfil this function, we noticed that i) "Slimy" bacteria
were surprisingly resistant to T6SS attacks, and ii) the simulated EPS particles covered a large
area around the producing cells. This led us to ask whether the secretion of loose (i.e. non-
membrane-bound) EPS could protect cells from T6SS firing events, and so help EPS-producing
bacteria survive when faced with a T6SS attacker.

To explore these questions, we implemented EPS secretion within an existing individual-based
model of T6SS competition (24, 25). Following previous models (29–31), EPS production is
modelled as cells secreting loose, spherical particles into the environment, at a rate P_{EPS} (see
Materials and Methods). As they grow, divide, and secrete EPS, producing cells create small
pockets of EPS particles around themselves, which helps to separate cells from T6SS-wielding
competitors (Fig. 1B). Note, we do not assume here that EPS can physically block T6SS needles,

as this effect may not be universal (15). However, were this to occur, the expectation is that it will

only increase the protection conferred to cells.

We performed simulation parameter sweeps to identify conditions where EPS production was

most protective against T6SS attacks. Comparing simulations with and without EPS (respectively,

P_{EPS} = 0.1, 0), we used a simple metric to quantify the benefit of EPS protection:

$$EPS\ benefit = (N_{susc}(t_{end})\ /\ N_{susc}(t_{start}))_{EPS+} - (N_{susc}(t_{end})\ /\ N_{susc}(t_{start}))_{EPS-}\ ,$$

with $N_{susc}(t)$ the number of surviving susceptible cells present at time t. Varying the starting number

of attacker and susceptible cells shows that EPS production always offers at least some T6SS

protection (*EPS benefit > 0*), and that production is most beneficial for intermediate attacker- and

susceptible cell densities (Fig. 1C).

We performed additional simulations to test the robustness of EPS-mediated T6SS protection

with respect to other simulation parameters (Fig. S1). Repeats of competition simulations with

different EPS production rates showed that, as might be expected, susceptible survival decreases

as the per-cell rate of EPS secretion is reduced (Fig. S1A); below a minimal secretion rate,

survival is indistinguishable from that of a non-producing susceptible strain. We also tested a

model in which EPS production is costly, such that susceptible cells suffer a growth penalty

proportional to their EPS production rate,

$$Susceptible\ growth\ rate = k_{grow}(1 - c_{EPS}\ P_{EPS}),$$

with k_{grow} the maximum cell growth rate, P_{EPS} the probability of EPS secretion per simulation

timestep, and c_{EPS} the unit cost of EPS production. For various $c_{EPS} > 0$, costly EPS production

remains net-beneficial to susceptible cells (Fig. S1B), such that the optimal secretion rate $P_{EPS}^* >$

0. Predictably, EPS investment collapses for excessive costs (Fig. S1B), and is contingent on

attacker presence (Fig. S1C). Overall, our simulations predict that production of loose, non-

membrane-bound EPS is an effective means for cells to protect themselves against the T6SS,

even when costly.

Fig. 1: Agent-based modelling of EPS secretion demonstrates collective defence against the T6SS. **(A)** Snapshots of our agent-based video game 'GutWars', showing EPS-producing 'Slimy' bacteria outcompeting T6SS-armed 'Stabby' rivals. **(B)** Agent-based simulations compete T6SS attackers (green) with EPS non-producing (EPS-, P_{EPS} = 0.0) and EPS-producing (EPS+, P_{EPS} = 0.1) susceptible cells (orange and magenta, right and left columns, respectively). Snapshots show i) initial cell seeding (0h), with 400 cells of each type scattered within a circular homeland, and ii) cell configuration after 3 hours' competition (3h). Scale bars: 50 μm. Zoomed sections highlight dead (black) susceptible cells and distribution of EPS particles. Inset scale bars: 10 μm. **(C)** Contour plot showing measurements of mean EPS benefit (see main text) as a function of initial attacker and susceptible cell density. N=20 simulation replicates per parameter combination. **(D)** Plot showing susceptible final:initial cell counts against initial susceptible density, exemplifying density-dependent benefit of EPS production (fixed attacker density of 400 cells). Individual data points are shown; lines and error bars correspond to data means and standard deviations. Dashed line denotes Final:Initial susceptible ratio = 1, i.e. no net loss or gain of susceptible strain. N=20 simulation replicates per parameter combination; data replotted from (C).

EPS protection is predicted to function as a collective trait

We next used the simulations to more formally evaluate whether EPS protection is a collective trait. A key feature of many collective traits is a positive relationship between the number of individuals with the trait, and the benefit conferred (32). To look for this, we plotted the benefit of EPS production against the starting number of susceptible (EPS producer) cells, which revealed that the relationship is positive for many levels of EPS secretion (Fig. 1D, S2A, S2B). Note that our metric controls for any density-dependent survival effects emerging solely from changes to cell group size (33). That is, while survival is expected to generally increase with susceptible cell number, this effect becomes much stronger when susceptible cells also produce EPS (Fig. 1D).

Next, we examined the extent to which the benefits from EPS production are shared among cells, which is a second key feature of collective traits. Using the same competition protocol as before, we ran simulations pitting T6SS attackers against susceptible EPS- cells, susceptible EPS+ cells, or both together (Fig. 2A), for a range of initial cell densities. We then quantified the survival of both the EPS+ cells and the EPS- cells when they grew either i) alone with the attacker or ii) paired with the other susceptible cell type (Fig. 2B). As expected when the protective effect of EPS is shared between cells, EPS- survival increased in the presence of EPS+ cells (Fig. 2B, 'rescue'), and, conversely, EPS+ survival decreased in the presence of EPS- cells (Fig. 2B, 'exploitation').

A synthetic community demonstrates collective protection via EPS

To test the model's predictions empirically, we turned to an established experimental system for studying T6SS killing: competitions between a T6SS+ *A. baylyi* attacker strain and a T6SS-susceptible *E. coli* strain (MG1655; (34)). Using their T6SS, *A. baylyi* cells can inject a cocktail of different effector proteins into neighbouring cells which rapidly induce lysis in susceptible *E. coli* (24). To compare the relative T6SS susceptibility of EPS-producing and non-producing *E. coli*, we created *E. coli* MG1655 pRcsA2 (EPS+), a strain capable of undergoing IPTG-inducible production of the exopolysaccharide colanic acid, which is a major component of natural *E. coli* biofilms (35, 36). Colanic acid production is induced via plasmid-based expression of RcsA, the major positive regulator of colanic acid biosynthesis in *E. coli* (37). Its plasmid-free parent MG1655 serves as the non-EPS-producing control strain (EPS-), as MG1655 does not produce significant amounts of colanic acid under the conditions used in this study (Fig. S3).

Fig. 2: Modelling and experiments show that EPS secretion protects both producer and non-producer *E. coli* from T6SS attacks. (A) End-state snapshots of agent-based simulations competing T6SS attackers (green) with EPS- or EPS+ (orange, magenta) strains, or both strains at once. **(B)** Survival of each susceptible strain (EPS+, magenta; EPS-, orange) growing alone against T6SS attackers (dashed lines), or together with the other susceptible strain type (solid lines). * denotes the scenario depicted in (A). **(C+D)** We conducted analogous competition assays *in vitro* between three bacterial strains: *A. baylyi* ADP1 (T6SS attacker), *E. coli* MG1655 Gen[R] (EPS-), and *E. coli* MG1655 pRcsA2 (EPS+; colanic acid overproducer). For each strain, cell recovery (CFU) after 6h of co-culturing at different starting cell densities (optical density; OD) is shown. Individual data points (n=3) and their means are shown as dots and lines, respectively. Arrows highlight key comparisons. **(C)** Survival of *E. coli* EPS- (left), EPS+ (centre) or a 1:1 mix of EPS- and EPS+ (right), when competed against T6SS+ *A.*

baylyi. At high seeding density (OD = 10^{-1}), EPS+ cells survive pairwise competitions against a T6SS attacker better than EPS- cells (middle and left; p < 0.0001). When co-cultured with EPS+ and competing against the attacker (right), survival of EPS- cells is strongly increased relative to when they face the attacker alone (left; p < 0.0001). When EPS+ cells are co-cultured with EPS- and compete against the attacker (right), EPS+ survival is reduced compared to when they face the attacker alone (left; p < 0.0001). **(D)** Repeat of (C) with *A. baylyi* lacking a functional T6SS. When competed against a T6SS-attacker in pairwise competitions, no difference in cell recovery between EPS+ and EPS- cells was detected (p = 0.66). For both (C) and (D), competitions were analysed per strain combination and seeding density, with statistical analyses on log-transformed data via two-way ANOVA (α = 0.01) with Tukey post-hoc tests (****: p < 0.0001).

We first sought to measure the amount of colanic acid secreted by the EPS+ strain, as well as

test where it is localised with respect to the cell membrane. Across *E. coli* strains, colanic acid

can be secreted both as a capsule tethered to the producing cell membrane, as well as in

untethered, loose form (38). The difference between these two states is likely to strongly influence

the role of colanic acid in collective protection from external assaults. To benchmark our colanic

acid quantification and localization assays, we compared the EPS+ and EPS- strains to six clinical

E. coli isolates (Table S2). As for the majority of the clinical isolates, phase-contrast microscopy

showed that our EPS+ strain has no visible membrane-bound capsule. This observation is

consistent with colanic acid being untethered to producing cells and released into the extracellular

environment (Fig. S3A). This conclusion was further supported by colanic acid quantification

assays where we measured the amount of colanic acid found in total cell extracts vs. supernatant

fractions of cultures harvested from agar plates. For the EPS+ strain, the entirety of the detected

colanic acid was recovered in the supernatant (Fig. S3B). This same assay also revealed that the

amount of loose colanic acid secreted by EPS+ is comparable to that secreted by two clinical

isolates (#0172 and #0993, Fig. S3B). The EPS+ *E. coli* strain, therefore, secretes loose colanic

acid at rates comparable to naturally occurring *E. coli,* allowing for a direct test of the role of EPS

in T6SS-mediated competitions.

Next, we compared the EPS+ and EPS- *E. coli* strains in competition against a T6SS-wielding *A.*

baylyi (ADP1; T6SS+) and a T6SS-deficient mutant (ADP1 Δ*tssM*; T6SS-). We seeded a 1:1

mixture of *A. baylyi* with either EPS+ or EPS- *E. coli* at various densities on agar plates and

measured strain recovery after 6h of co-culture using selective plating (Fig. 2C). At low seeding

densities, the T6SS is less efficient because it relies on cell-cell-contact for killing (24, 33). At high

seeding densities, we expect the T6SS to be maximally effective; this was confirmed in our

competition assays, as the EPS- control strain suffered maximal killing at high densities (Fig. 2C,

left). In contrast, the EPS+ strain survived much better at high densities (Fig. 2C, middle), with a

10^5-fold increase in survival against T6SS+ *A. baylyi* compared to EPS-. This advantage was

dependent on T6SS-activity, as there was no detectable difference in recovery between EPS+

and EPS- after pairwise competition with the T6SS- *A. baylyi* (Δ*tssM*) mutant (Fig. 2D). To test

whether the presence of EPS in the environment alters T6SS activity, we measured firing rates in

hundreds of T6SS+ *A. baylyi* cells exposed to cell-free supernatants with and without EPS, and

were unable to detect a difference (Fig. S4). These results strongly support our model's prediction

that non-membrane-bound EPS can protect susceptible cells from killing by the T6SS.

We then examined the potential for a producer strain to cross-protect a non-producer, as expected

for a collective defence mechanism. We conducted 3-way competitions between T6SS+ *A. baylyi*

and both the EPS+ and EPS- *E. coli* (2:1:1 ratio of *A. baylyi* : EPS+ : EPS-) and quantified strain

recovery as above. At high seeding densities, as predicted by the model, the survival of EPS-

increased 1000-fold in the presence of EPS+ (Fig. 2C, right) compared to when they faced the

T6SS attacker alone (Fig. 2C, left), indicating that the presence of secreted EPS can rescue

otherwise susceptible cells from T6SS attacks even if they do not produce EPS themselves.

Further consistent with the modelling, we observed that relative survival of the EPS+ strain

decreased in presence of EPS- (Fig. 2C, right) compared to when grown alone with the T6SS

attacker (Fig. 2C, left) suggesting that the presence of non-producing cells partially compromises

the protective effects of EPS production.

Cross-protection can emerge from flank protection

Our modelling and experiments show that EPS can function as a collective defence, whose

benefit is shared among cells in a colony or biofilm. We next sought to understand the

mechanism(s) underpinning this collective property. We initially reasoned that loose EPS would

be shared between cells, resulting in the protection of non-producing cells seen in both modelling

and experiments. To test this hypothesis, we modified our model so that EPS+ cells no longer

secreted loose EPS particles, but instead became intrinsically resistant to T6SS attacks (requiring

2 hits to kill instead of 1), as has been demonstrated in capsule-forming bacteria (15). The goal

of this model was to remove the possibility of EPS sharing between cells, so that we could test

whether sharing was necessary for the cross-protection of non-producer cells (Fig. 3).

Fig. 3. Modelling shows that cross protection can occur without EPS sharing (A) End-state snapshots of agent-based simulations competing T6SS attackers (green) with an EPS- strain (orange), a T6SS-resistant strain (purple) that secretes no EPS particles, or both strains at once. Box corresponds to zoomed section in (C). **(B)** Survival of Resistant strain (purple) and the EPS- susceptible strain (orange) growing alone against T6SS attackers (dashed lines), or together with the other strain type (solid lines). N=5 simulations per parameter combination; lines denote mean survival. **(C)** Magnified views showing Resistant strain shielding EPS- from the attacker (top), similar to where protection occurs via secretion of loose EPS (bottom).

We then competed the resistant cell type against T6SS attackers as before, either in the presence

or absence of the EPS- susceptible cell type (Fig. 3A). To our surprise, this again lead to collective

T6SS defence, which displayed both cross-protection and the same density-dependence seen

for loose EPS (Fig. 3B). Close inspection of simulations revealed that resistant cells act as

barriers, protecting the flanks of susceptible cell groups by physically separating them from T6SS

attackers (Fig. 3C). We therefore term this mechanism 'flank protection'. Because we see cross

protection, flank protection is also a collective effect, but one that can be provided by a purely

private resistance mechanism, as well as by loose EPS.

Time-lapse microscopy demonstrates cross-protection involves EPS sharing

Our modelling shows that EPS-producing bacteria can collectively defend themselves against

T6SS attacks via flank protection, without sharing of EPS between cells. However, an important

simplification of the model is that it represents EPS as small solid spheres. This limits the ability

of EPS to flow past and around cells in a manner that may not always reflect the behaviour of real

EPS (39). Given this potential limitation of the model, we decided to investigate whether our

original hypothesis of EPS sharing was important in the experimental data, even if it was not

necessary for collective defence in the model. To do this, we turned to time-lapse confocal

microscopy, which enables direct observation of cells over time (Fig 4). We competed fluorescent

protein-expressing strains of *A. baylyi* (*vipA-sfGFP*, green) with EPS+ *E. coli* (*mScarlet*, magenta)

and EPS- *E. coli* (*BFP*, orange) and performed competitions on agar pads as before, except we

imaged different locations in the competition area hourly, using confocal microscopy. End-state

images of the competition area (Fig. 4A) provided visual confirmation of i) improved survival of

EPS+ relative to EPS-, and ii) improved survival of EPS- in the presence of EPS+. Surviving EPS-

cells also tended to co-localise with EPS+ cells (bottom right), similar to behaviour observed in

our model.

To look for signatures of EPS-mediated cross-protection in our time-lapse data, we examined

T6SS-mediated killing at boundaries between *A. baylyi* and *E. coli* cell groups (Fig. 4B) using

automated analysis of confocal time-lapse images. We reasoned that, if cross-protection occurred

via the association of loose EPS with EPS- cells, then an EPS- cell group should be able to

withstand the T6SS attacker better when EPS+ producers were also present. Specifically, the

boundaries between the T6SS attacker and EPS- cells should shift less in the favour of the T6SS

attacker, if EPS+ cells are also present nearby. Accordingly, we measured the rate of *A. baylyi*

encroachment into EPS+ and EPS- cell groups (Fig. 4B, "Invasion"), normalising this by the total

length of interface between *A. baylyi* and each individual susceptible strain (Fig. 4B, "Boundary").

The resulting metric, which we term the normalised invasion rate, provides an estimate of how

susceptible each *E. coli* strain is to T6SS attack. Crucially, since the normalised invasion rate

examines *A. baylyi*–EPS+ and *A. baylyi*–EPS- strain interactions separately, it excludes any

possible effects of flank protection.

Our measurements of normalised invasion rates over time showed that the EPS- strain was

indeed invaded less rapidly when grown together with the EPS+ strain, vs. when grown alone

(Fig. 4C, left). This improvement was most pronounced between hours 3 and 4 of competition.

Meanwhile, the EPS+ strain was invaded at a consistently lower rate (Fig. 4C right) than the EPS-

strain, with normalised invasion rates largely unchanged by the presence / absence of the EPS-

strain (i.e., no significant exploitation effect detected). These effects are summarised in Fig. 4D:

integrating over all time points, the EPS- cell groups suffered significantly more invasion when

the EPS+ strain was absent. Importantly, this variation was contingent on T6SS activity: control

experiments showed minimal variation in EPS- strain invasion with EPS+ strain presence when

attackers lacked a functional T6SS (Fig. 4E, right). Note that these data do not exclude a role for

flank protection in the experiments, but they do provide evidence that EPS+ cells increase survival

of EPS- cells *at a distance*. This result is consistent with collective protection being mediated—at

least in part—by shared EPS.

Fig. 4: ***A. baylyi* invades EPS-** *E. coli* **cell groups less rapidly in the presence of EPS+ strain, supporting ranged protection via EPS sharing. (A)** Attacker (green), EPS- (orange) and EPS+ (magenta) strains expressing fluorescent proteins were mixed and co-cultured as before (Fig. 2C,D), this time analysing competition dynamics using confocal microscopy. Colonies were imaged for 7h at 1h intervals, across five 470 x 470 µm frames spanning the colony edge (position 1) to its centre (position 5). Example images shown are zoomed-in areas of interest from position 4 (t = 6h); scale bars: 50 µm. **(B)** Automated image analysis for measuring rate of *A. baylyi* invasion into *E. coli* cell groups. We used segmented time-lapse images to identify i) boundaries between *A. baylyi* and EPS-, EPS+ *E. coli* cell groups (grey pixels), and ii) the rates at which *A. baylyi* expanded into EPS-, EPS+ *E. coli* groups

(orange- and magenta-edged pixels). **(C)** Measurements of *A. baylyi* invasion rates, normalised by boundary length, comparing invasion of EPS- (left) and EPS+ (right) cell groups, grown alone (dashed lines) or together (solid lines). Lines connect data means, and error bars denote standard deviations. Time-lapse data aggregated over colony positions 2-5; Position 1 was excluded for analysis since cells quickly grew beyond a monolayer, confounding our 2-D analysis. N=3 biological replicates; 12 data points per time point. **(D)** Comparison of total normalised invasion (area under curves in (D)) across all strain pairings, including T6SS- controls. Red cross denotes outlier data point (> Q3 + 1.5 * IQR). Significance assessed using one-sample Kruskal-Wallis tests; *** denotes p = 0.0004; N.S., non-significant.

DISCUSSION

Our modelling shows how EPS can serve as a highly protective, collective defence strategy

against T6SS attacks, and one that can also mediate cross-protection of neighbouring, non-

producing cells. Moreover, an experimental test of our model revealed that EPS dramatically

increases *E. coli* survival in competition with T6SS-wielding *A. baylyi*. Here, colanic acid is

secreted into the extracellular environment and we find evidence that it has the potential to protect

cells other than the producer. Secretion of colanic acid is widespread in *Enterobacteriaceae* such

as *E. coli*, *Klebsiella*, *Salmonella* and *Enterobacter* (40, 41), where it plays an important role in

biofilm formation (35, 36, 42), and confers protection against various environmental stressors

such as desiccation, osmotic stress, and oxidative stress (43, 44). Moreover, elevated EPS

production (mucoidy) readily evolves in *E. coli* and other species in response to predation by

phages (45–47), and membrane-anchored capsules can protect pathogenic bacteria from attacks

by the immune system (48–50). EPS-mediated protection from T6SS attacks could therefore also

evolve as a by-product of adaptation to other types of stressors, as well as a direct response to

T6SS attacks itself.

With both modelling and experiments, we also found that EPS non-producing cells are able to

survive T6SS attacks better when co-cultured with EPS producers. Additionally, our simulations

identified a mechanism of cross-protection that we call flank protection, whereby groups of T6SS-

resistant producers reduce the exposure of non-producers to attackers. However, in the

experiments, we find evidence for a second, perhaps more intuitive, mechanism whereby loose

EPS covers non-producers and directly shields them. Specifically, our microscopy experiments

support the presence of direct protection via EPS sharing, as evidenced by a reduction in attacker

invasion at the edges of non-producer cell groups. This implies that secreted EPS can shield non-

producing cells, at a range spanning multiple cell lengths.

As discussed, our analyses do not exclude flank protection as a contributor to cross-protection in

our experiments; we expect it to occur in parallel with EPS sharing. Indeed, flank protection has

the potential to be highly general: it is predicted to emerge whenever sensitive cell groups

interface with resistant cell groups, without assuming anything about the underlying resistance

mechanism. Collective resistance, therefore, may emerge widely, including in cases where EPS

is not shared between cells (15). More generally, flank protection has the potential to be mediated

by other mechanisms of T6SS resistance, such as envelope stress responses, or the production

of immunity proteins. In principle, it could generate unexpected collective protection against any

contact-dependent weapon, including via the T4SS, T7SS, and Cdi systems (4), adding further

complexity to interbacterial interactions.

T6SS-wielding bacteria are increasingly being considered as potential biotherapeutics, capable

of delivering toxins into pathogenic bacteria (51, 52). Our work cautions that many species have

the potential to resist such attacks due to the widespread ability of bacteria to work as collectives.

Consistent with this, bacteria are already known to display collective defences against antibiotics

(17, 19, 22), predators (53) and phage (20), and biofilm formation continues to be a major issue

in industry and medicine (54–57). Strategies that seek to remove problem strains, therefore,

needs to consider the ability of bacteria to work both as individuals and as groups.

MATERIALS AND METHODS

Agent-based modelling

We ran agent-based model simulations using the open-source modelling software Cellmodeller

(58). To simulate T6SS-mediated interactions and EPS particle secretion, we devised additional

code modules, available for download at https://github.com/WilliamPJSmith/CellModeller. The

key processes incorporated in our model are summarised here.

<u>Model description</u>

Cell growth and division. Bacterial cells are modelled as exponentially-growing rigid elastic rods

with fixed radius R, which push on one another as they elongate and divide (after doubling their

initial volume V_0, plus a small random noise term η_{div}). All cells are assumed to have equal access

to growth-limiting nutrients. Generally, cells pay no growth penalty for using either T6SS needles

or EPS secretion, and so a cell's volume V increases over time as $\Delta V = k_{grow} V \Delta t,$ with k_{grow} the

maximum specific growth rate, and Δt the simulation timestep. However, we also explored

scenarios in which EPS secretion carried a range of growth costs proportional to the EPS

secretion rate (see Fig. S2B,C). Immediately after cell growth, an energy penalty method is

applied each timestep to compute cell movements necessary to minimise total cell-cell overlap,

subject to viscous drag forces acting on each cell. This process, described previously in detail

(23, 58), approximates the elastic repulsion forces acting between cells in physical contact.

EPS secretion. Similar to previous work, we model agglomerations of EPS polymers as inert,

pseudospherical particles (29–31). EPS+ cells secrete these particles from randomly chosen

points on their surface. These particles have radius 0.4 µm, rod segment length 0.01 µm, and are

assumed mechanically identical to bacterial cells. The secretion probability per time step, P_{EPS}, is

at most 0.1, such that the max secretion rate is capped at 1 particle per timestep. EPS particles

persist indefinitely, and completely ignore the effects of T6SS toxin injection (see below).

T6SS interactions. Each time step, T6SS+ cells can fire needles of length *R/2,* projecting

orthogonally from randomly chosen sites on their cell surface. The number of firings per cell per

unit time is drawn from a Poisson distribution, whose mean is k_{fire}. After firing, each needle is

checked to determine if it comes into contact with any other cell in the population (line-segment

method). Subsequent to this contact finding, each cell's state is updated to indicate the number

of T6SS needle hits it sustained. Cells die (cease growing) after being struck N_{hits} times in this

way, and lyse (disappear) following a fixed delay of *1 / k_{lysis}*. Each T6SS needle can deliver its

toxin payload to at most one cell. T6SS+ cells are assumed completely immune to their own T6SS

toxins, such that they effectively ignore all collateral fire from clone-mates. EPS particles do not

obstruct T6SS needles, and so EPS particles can only protect bacterial cells from T6SS attacks

by enforcing spatial separation of attacker and susceptible cells.

<u>Computation and post-processing</u>

All agent-based simulations were run using a 2017 Apple MacBook Pro laptop computer, with

concurrent simulations distributed between an Intel 3.1 GHz quadcore i7-7920HQ CPU, an Intel

HD 630 Graphics card, and an AMD Radeon Pro 560 Compute Engine. Simulation data were

visualised using Paraview software, and analysed using custom Matlab and R scripts.

<u>Simulation protocols</u>

All simulations were initialised by randomly scattering different combinations of cells on a flat

surface, within a circular 'homeland' of radius 50 µm. At simulation time *t* = 0, all cells lie flat on

the surface, but have randomised orientations. All simulations were run for 130 time steps (*Δt* =

0.025h), totalling 3.25h of simulated time. Depending on conditions (most crucially, the number

of EPS particles present), most simulations had a runtime of between 10 and 30 minutes

(approximately 10 simulations per hour after concurrency).

Bacterial growth conditions

All strains were routinely grown overnight in 5 ml LB medium (per L: 10g Tryptone, 10g NaCl, 5g Yeast Extract) in 15 mL polypropylene tubes with agitation (220 rpm). Where necessary, the medium was supplemented with streptomycin (50 µg/mL), ampicillin (100 µg/mL), carbenicillin (100 µg/mL) or gentamycin (15 µg/mL). Routine culturing was carried out on 1.5% w/v LB Agar at 37°C (*E. coli*) or 30°C (*A. baylyi*). All pre-cultures and experiments were carried out at 30°C on 0.8% w/v LB Agar, unless indicated otherwise. Optical density (OD) of liquid cultures was measured at 600 nm. For time-lapse microscopy experiments, samples were kept at 30°C throughout the duration of the experiment using a custom-built incubation chamber.

Strain construction

For the generation of MG1655 *mScarlet-I*::Tn7 and MG1655 *mTagBFP2*::Tn7, MG1655 cells were transformed with either pGRG25-P*max*::*mScarlet-I* or pGRG25-P*max*::*mTagBFP2* via electroporation and selected on ampicillin at 30°C. The resulting transformants were cultured for 16 h at 30°C in liquid LB medium supplemented with ampicillin and arabinose (0.5% w/v). A small volume (appr. 5 µL) of these cultures was streaked onto 1.5% w/v LB agar supplemented with arabinose (0.5% w/v) and incubated at 42°C for 12 h. For each construct, three fluorescent colonies were then selected, streaked onto 1.5% w/v LB agar and incubated at 42°C for 12 h. Resulting colonies were tested for ampicillin sensitivity. For the generation of *E. coli* MG1655 *mScarlet-I*::Tn7 pRcsA2, MG1655 *mScarlet-I*::Tn7 cells were transformed with pRcsA2 (37) via electroporation and selected on 100 µg/mL carbenicillin at 37°C. All plasmids and strains used in this study are listed in Table S2.

Competition assays

To monitor survival of different *E. coli* strains in competition with *A. baylyi* strains, overnight cultures of *A. baylyi* and *E. coli* were diluted (*A. baylyi* 1:10; *E. coli* 1:100) into 5 mL of fresh LB medium supplemented with appropriate antibiotics (Table S2) and 0.5 mM IPTG to induce production of colanic acid. We used *vipA-sfGFP*-labelled *A. baylyi* ADP1 as the T6SS-positive competitor, whereas an isogenic *tssM* deletion mutant was used as T6SS-negative control competitor. Cultures were then grown for 3 h with agitation, washed twice with LB medium, and resuspended and normalized in LB medium to an optical density at 600 nm (OD600) of 0.1. An *A. baylyi* strain was then mixed with an *E. coli* strain at a ratio of 1:1 (100 µL : 100 µL), and for

each strain combination 5 µL were spotted in triplicates in 24-well plates pre-filled with LB agar

supplemented with 0.5 mM IPTG (1 mL per well). For experiments at different seeding densities

(Fig. 2), the initial 1:1 mix was further diluted 10-fold (for OD = 0.01) or 100-fold (for OD = 0.001)

before being spotted on LB agar. For experiments with three competitors (Fig. 2), one *A. baylyi*

strain and two *E. coli* strains were normalized to OD = 0.1, mixed at a ratio of 2:1:1 (*A. baylyi* : *E.*

coli : *E. coli*), and further diluted if required before being spotted on LB agar. Plates were then

dried in a laminar flow at room temperature for approximately 20 min to allow droplet drying,

before being incubated at 30°C for 6 h. After incubation, competition spots were washed off with

500 µL NaCl (0.8%) each and subjected to seven rounds of 10-fold serial dilution. Colony-forming

units (CFUs) of *A. baylyi* and *E. coli* strains were determined by plating on streptomycin (100

µg/mL; selecting for *A. baylyi*), ampicillin/carbenicillin (100 µg/mL; selecting for *E. coli* MG1655

pRcsA2) or gentamycin (15 µg/mL; selecting for *E. coli* MG1655 Gen^R) media, incubated at 30°C

(*A. baylyi*) or 37°C (*E. coli*).

Determination of T6SS firing rate

To determine T6SS firing rate in *A. baylyi* cells exposed to EPS produced by *E. coli* (Fig. S4A),

cells of *E. coli* EPS+ and EPS- were grown overnight for 16 hours at 37°C, and 1 mL of bacterial

suspension was centrifuged for 5 min at 17000 g to sediment cells. The resulting supernatants

were sterile-filtered using 0.2 µm syringe filters (Sartorius), and 5 µL were each pipetted onto

separate cut-outs (ca. 5 mm x 5 mm) of 1% w/v LB agarose and left to dry for 20 min at room

temperature. In parallel, cells of the focal strain (*A. baylyi* T6SS+ *clpV-mCherry2*) were grown to

exponential phase in liquid culture (OD600 ~ 0.8-1.0) before spotting 1 µL of culture onto the

agarose pads treated with supernatant, and leaving them to dry for 5 min at room temperature.

Pads were then inverted onto a 5-cm diameter glass bottom Petri dish with a 3-cm diameter

uncoated n° 1.5 glass window (MatTek Corporation), so that the cells were sandwiched between

the agarose and the glass. The sample was then moved to the microscope and imaged

immediately. Time-lapse fluorescence microscopy was performed using a Zeiss Axio Observer

inverted microscope with a Zeiss Plan-Apochromat 63x oil immersion objective (NA = 1.4) and

ZEN Blue software (version 1.1.2.0). Time-lapse images were recorded for 3 minutes at a 10-

second interval. Exposure times were 1.05 ms for brightfield and 500 ms for mCherry2 (ex:

589 nm | em: 610 nm). The rate of T6SS firing was measured in a total of n = 916 *A. baylyi* cells

via detection of ClpV (*mCherry*) fluorescent foci (24). Each new ClpV focus was assumed to mark

one T6SS contraction event in a focal cell. Three separate fields of view at different locations on

the same agarose pad were monitored for each treatment. Image analysis was performed using

FIJI 2.1.0/1.53c (59). For each field of view, total number of cells and ClpV foci were determined

each minute via thresholding, segmentation and object counting in the brightfield channel, and

the FIJI built-in "Find maxima" function in the mCherry fluorescence channel, respectively

Colanic acid quantification

To quantify the amount of colanic acid (CA) produced by different *E. coli* strains (Fig. S2), 1 mL

of overnight cultures were washed once with fresh LB medium and density-adjusted to an OD of

1.0. For each strain, 50 μL were then spotted in triplicates in 24-well plates pre-filled with LB agar

supplemented with 0.5 mM IPTG (1 mL per well). Plates were then dried in a laminar flow at room

temperature for approximately 20 min to allow droplet evaporation, before being incubated at

30°C for 6h. Cell material was then harvested by adding 1 mL of sterile ddH$_2$O to the well and

pipetting up and down until all material was suspended, and then transferred to a sterile 1.5 mL

tube. Tubes were vortexed for 5-7 seconds, and the OD600 was determined for all replicates. Of

these volumes, twice 400 μL were then transferred to two separate reaction tubes pre-filled with

600 μL ddH$_2$O: one for quantification of total CA, and one for quantification of CA secreted into

the extracellular environment. For quantification of total CA (Fig. S3), samples were boiled for 20

min at 95°C, centrifuged at 17'000 g for 20 min, and 1 mL of supernatant was used for further

processing. For quantification of CA in the extracellular environment (Fig. S3), samples were

centrifuged at 17'000 g for 20 min, and 1 mL of supernatant was used for further processing. To

test whether sterile-filtering removes EPS from supernatants (Fig. S4B), *E. coli* MG1655 pRcsA2

(EPS+) and MG1655 Gen^R (EPS-) strains were grown overnight for 16 hours at 37°C, and 1 mL

of bacterial suspension was centrifuged for 5 min at 17000 g to sediment cells. The resulting

supernatants were either sterile-filtered using 0.2 μm syringe filters (Sartorius) or left unfiltered,

and 900uL of supernatant was used for further processing. Relative CA production was then

determined through the quantification of L-Fucose (60). Briefly, samples were boiled with 4.5 mL

H$_2$SO$_4$·H$_2$O (6:1 H$_2$SO$_4$ and H$_2$O) for 20 min and cooled to room temperature. Then, 900 μL of

each sample was transferred to a spectrophotometer cuvette and absorbance was measured at

396 nm (A$_{396}$) and 427 nm (A$_{427}$). To each cuvette was then added 100 μL of freshly prepared L-

cysteine hydrochloride (1M), mixed by pipetting, and absorbance was again measured at 396 nm

(A-cy$_{396}$) and 427 nm (A-cy$_{427}$). The concentration of L-Fucose was then determined by calculating

$(A\text{-}cy_{396} - A_{396}) - (A\text{-}cy_{427} - (A_{427})$ and applying a previously generated L-Fucose standard
curve (0-100 µg/mL).

Confocal microscopy

To image competition dynamics, fluorescently tagged versions of all relevant strains (Table S2)
were pre-cultured and density-adjusted to OD600 = 0.1 as described in "Competition assays".
For each strain combination, 5 µL were then spotted onto a glass bottom Petri dish filled with 8
mL 0.8% w/v LB agar supplemented with 0.5 mM IPTG. After drying for ten minutes at room
temperature, the plate was inverted to allow for immediate imaging of the competition area
through air. Time-lapse confocal imaging was then carried out using a Zeiss Plan-Apochromat
10x objective (NA = 0.45) on a Zeiss LSM880 confocal laser scanning unit using ZEN Black
software (version 14.0.18.201). Images (each covering an area of 472.33 µm x 472.33 µm) were
acquired hourly for 7 h, over 5 different colony positions, spanning in total the entire radius of the
competition area (edge to centre). Three biological replicates were carried out for each strain
combination.

Confocal image analysis

Image segmentation and stabilization. Following acquisition, raw confocal image data were
analyzed using FIJI (59). Multichannel images were split into individual channels, with each
channel subjected to separate background subtraction (rolling ball radius 50 px) and conversion
to binary image data via user-defined thresholds. To maximise the accuracy of cell group
boundary tracking, we stabilised images using the FIJI 'StackReg' plugin (61), applied to the GFP
(*A. baylyi*) channel. This stabilisation made it necessary to exclude later time points (t=6,7h) as
these were typically confluent, confounding accurate registration of images. We also excluded
colony position 1 (colony edge); here, coffee-ring effects (62) consistently led to the formation of
3-D cell groups as early as 2h into the competition. As our image analysis is limited to 2-D, we
were unable to accurately track inter-strain boundaries at this position. For visualization purposes,
mtagBFP2 fluorescence is false-coloured orange in all images throughout the manuscript.

Normalised invasion rates. To estimate T6SS killing rates at interstrain boundaries (Fig. 4), we
tracked changes in strain occupancy at each pixel across consecutive time-lapse images, using

custom MATLAB scripts. We compared consecutive time points (t-1, t; t=1,2,3,4,5h) in each
stablised time lapse. Any pixel where EPS+ channel signal was lost, and replaced with new *A.*
baylyi signal, was marked as being 'invaded' by *A. baylyi*, giving a measure of EPS+ cell killing
by the attacker. EPS+ invasion pixel counts were normalised by the amount of Ab-EPS+ contact
found in the t-1 image. We measured Ab-EPS+ contact by dilating *Ab* cell group images (MATLAB
imdilate method; 3 pixel spherical structuring element) and computing their overlap (in pixels) with
EPS+ cell groups. Performing this procedure in reverse (dilate EPS+ onto *Ab*) yielded similar
boundary pixel counts. EPS- invasion was calculated and normalised in the same way.

Statistical Analysis

Statistical analysis and data visualization were performed using RStudio version 1.2.5033 (63)
and packages dplyr (64), Rmisc (65), ggplot2 (66), and cowplot (67); as well as Matlab version
2017b (9.3.0.713579). For all statistical tests, the significance level α was set to 0.05. To test
whether cell recovery differed between competitions (Fig. 2), two-way ANOVA of log-transformed
CFU values with a post-hoc Tukey's HSD test was performed. To test whether colanic acid
production differed between strains (Fig. S3B), and whether firing rate differed when cells were
exposed to EPS (Fig. S4A), a two-sided Wilcoxon rank sum test was performed. To test whether
filtering removed EPS from supernatants (Fig. S4B), a two-sided, paired samples Wilcoxon signed
rank test was performed. To test whether total normalised invasion of the EPS- strain changed
when grown alongside the EPS+ strain (or vice versa; Fig. 4), we used one-sample Kruskal-Wallis
tests. Statistical details (sample size, test values, p-values) for each experiment are provided in
the respective Figure legends.

ACKNOWLEDGEMENTS

We thank Marek Basler, Colin Kleanthous and Frances Spragge for strains, Chris French and
Thomas Meiller-Legrand for plasmids, and Erik Bakkeren, Ravinash Krishna Kumar, Jake Palmer,
and Rachel Wheatley for comments on the manuscript. ETG is funded by a Postdoc Mobility
Fellowship from the Swiss National Science Foundation (P400PB_183878) and a BBSRC
Discovery Fellowship (BB/V004328/1). WPJS and KRF are supported by the National Institutes
of Health (project numbers R01AI093771 and R01AI120633), by European Research Council
Grant 787932, and by Wellcome Trust Investigator award 209397/Z/17/Z. WPJS is also funded
by a Sir Henry Wellcome Postdoctoral fellowship award, 222795/Z/21/Z.

REFERENCES

1. M. Ghoul, S. Mitri, The Ecology and Evolution of Microbial Competition. *Trends Microbiol.* **24**, 833–845 (2016).

2. M. E. Hibbing, C. Fuqua, M. R. Parsek, S. B. Peterson, Bacterial competition: Surviving and thriving in the microbial jungle. *Nat. Rev. Microbiol.* **8**, 15–25 (2010).

3. J. D. Palmer, K. R. Foster, Bacterial species rarely work together. *Science* **376**, 581–582 (2022).

4. E. T. Granato, T. A. Meiller-Legrand, K. R. Foster, The Evolution and Ecology of Bacterial Warfare. *Curr. Biol.* **29**, R521–R537 (2019).

5. S. B. Peterson, S. K. Bertolli, J. D. Mougous, The Central Role of Interbacterial Antagonism in Bacterial Life. *Curr. Biol.* **30**, R1203–R1214 (2020).

6. F. R. Cianfanelli, L. Monlezun, S. J. Coulthurst, Aim, Load, Fire: The Type VI Secretion System, a Bacterial Nanoweapon. *Trends Microbiol.* **24**, 51–62 (2016).

7. L. P. Allsopp, P. Bernal, L. M. Nolan, A. Filloux, Causalities of War: The connection between T6SS and microbiota. *Cell. Microbiol.*, cmi.13153 (2019).

8. P. Bernal, L. P. Allsopp, A. Filloux, M. A. Llamas, The Pseudomonas putida T6SS is a plant warden against phytopathogens. *ISME J.* **11**, 972–987 (2017).

9. C. Chen, X. Yang, X. Shen, Confirmed and Potential Roles of Bacterial T6SSs in the Intestinal Ecosystem. *Front. Microbiol.* **0** (2019).

10. R. Gallegos-Monterrosa, S. J. Coulthurst, The ecological impact of a bacterial weapon: microbial interactions and the Type VI secretion system. *FEMS Microbiol. Rev.* (2021).

11. B. Chassaing, E. Cascales, Antibacterial Weapons: Targeted Destruction in the Microbiota. *Trends Microbiol.* **26**, 329–338 (2018).

12. N. Flaugnatti, *et al.*, Human commensal gut Proteobacteria withstand type VI secretion attacks through immunity protein-independent mechanisms. *Nat. Commun.* **12**, 5751 (2021).

13. S. J. Hersch, *et al.*, Envelope stress responses defend against type six secretion system attacks independently of immunity proteins. *Nat. Microbiol.* **5**, 706–714 (2020).

14. S. Robitaille, E. Trus, B. D. Ross, Bacterial Defense against the Type VI Secretion System. *Trends Microbiol.* (2020) https:/doi.org/10.1016/j.tim.2020.09.001 (November 22, 2020).

15. J. Toska, B. T. Ho, J. J. Mekalanos, Exopolysaccharide protects Vibrio cholerae from exogenous attacks by the type 6 secretion system. *Proc. Natl. Acad. Sci.*, 201808469 (2018).

16. S. J. Hersch, K. Manera, T. G. Dong, Defending against the Type Six Secretion System: beyond Immunity Genes. *Cell Rep.* **33**, 108259 (2020).

17. I. Frost, *et al.*, Cooperation, competition and antibiotic resistance in bacterial colonies.
ISME J. **12**, 1582–1593 (2018).

18. E. T. Granato, K. R. Foster, The Evolution of Mass Cell Suicide in Bacterial Warfare. *Curr.*
Biol. (2020) https:/doi.org/10.1016/j.cub.2020.05.007 (June 5, 2020).

19. R. Krishna Kumar, *et al.*, Droplet printing reveals the importance of micron-scale structure
for bacterial ecology. *Nat. Commun.* **12**, 857 (2021).

20. A. Lopatina, N. Tal, R. Sorek, Abortive Infection: Bacterial Suicide as an Antiviral Immune
Strategy. *Annu. Rev. Virol.* **7**, 371–384 (2020).

21. D. A. I. Mavridou, D. Gonzalez, W. Kim, S. A. West, K. R. Foster, Bacteria Use Collective
Behavior to Generate Diverse Combat Strategies. *Curr. Biol.* **28**, 345-355.e4 (2018).

22. D. A. C. Stapels, *et al.*, Salmonella persisters undermine host immune defenses during
antibiotic treatment. *Science* **362**, 1156–1160 (2018).

23. W. P. J. Smith, *et al.*, Cell morphology drives spatial patterning in microbial communities.
Proc. Natl. Acad. Sci. U. S. A. **114**, E280–E286 (2017).

24. W. P. J. Smith, *et al.*, The evolution of the type VI secretion system as a disintegration
weapon. *PLOS Biol.* **18**, e3000720 (2020).

25. W. P. J. Smith, *et al.*, The evolution of tit-for-tat in bacteria via the type VI secretion
system. *Nat. Commun.* **11**, 5395 (2020).

26. H.-C. Flemming, J. Wingender, The biofilm matrix. *Nat. Rev. Microbiol.* **8**, 623–633 (2010).

27. H. C. Flemming, S. Wuertz, Bacteria and archaea on Earth and their abundance in
biofilms. *Nat. Rev. Microbiol.* **17**, 247–260 (2019).

28. B. Lories, *et al.*, Biofilm Bacteria Use Stress Responses to Detect and Respond to
Competitors. *Curr. Biol.* **30**, 1231-1244.e4 (2020).

29. P. Ghosh, J. Mondal, E. Ben-Jacob, H. Levine, Mechanically-driven phase separation in a
growing bacterial colony. *Proc. Natl. Acad. Sci.* **112**, E2166–E2173 (2015).

30. J. U. Kreft, J. W. Wimpenny, Effect of EPS on biofilm structure and function as revealed by
an individual-based model of biofilm growth. *Water Sci. Technol. J. Int. Assoc. Water*
Pollut. Res. **43**, 135–141 (2001).

31. J. B. Xavier, K. R. Foster, Cooperation and conflict in microbial biofilms. *Proc. Natl. Acad.*
Sci. **104**, 876–881 (2007).

32. A. Berdahl, C. J. Torney, C. C. Ioannou, J. J. Faria, I. D. Couzin, Emergent Sensing of
Complex Environments by Mobile Animal Groups. *Science* **339**, 574–576 (2013).

33. D. B. Borenstein, P. Ringel, M. Basler, N. S. Wingreen, Established Microbial Colonies
Can Survive Type VI Secretion Assault. *PLOS Comput. Biol.* **11**, e1004520 (2015).

34. P. D. Ringel, D. Hu, M. Basler, The Role of Type VI Secretion System Effectors in Target
Cell Lysis and Subsequent Horizontal Gene Transfer. *Cell Rep.* **21**, 3927–3940 (2017).

35. P. N. Danese, L. A. Pratt, R. Kolter, Exopolysaccharide Production Is Required for
Development of Escherichia coli K-12 Biofilm Architecture. *J. Bacteriol.* (2000)
https:/doi.org/10.1128/JB.182.12.3593-3596.2000 (December 28, 2021).

36. M. Horvat, A. Pannuri, T. Romeo, I. Dogsa, D. Stopar, Viscoelastic response of
Escherichia coli biofilms to genetically altered expression of extracellular matrix
components. *Soft Matter* **15**, 5042–5051 (2019).

37. N. Joshi, B. T. Ngwenya, C. E. French, Enhanced resistance to nanoparticle toxicity is
conferred by overproduction of extracellular polymeric substances. *J. Hazard. Mater.* **241–**
242, 363–370 (2012).

38. C. Whitfield, Biosynthesis and Assembly of Capsular Polysaccharides in Escherichia coli.
Annu. Rev. Biochem. **75**, 39–68 (2006).

39. J. van Gestel, F. J. Weissing, O. P. Kuipers, Á. T. Kovács, Density of founder cells affects
spatial pattern formation and cooperation in Bacillus subtilis biofilms. *ISME J.* **8**, 2069–
2079 (2014).

40. M. Rättö, *et al.*, Colanic acid is an exopolysaccharide common to many enterobacteria
isolated from paper-machine slimes. *J. Ind. Microbiol. Biotechnol.* **33**, 359–367 (2006).

41. C. Sande, C. Whitfield, Capsules and Extracellular Polysaccharides in Escherichia coli and
Salmonella. *EcoSal Plus* **9**, eESP-0033-2020 (2021).

42. M. M. Hahn, J. S. Gunn, Salmonella Extracellular Polymeric Substances Modulate Innate
Phagocyte Activity and Enhance Tolerance of Biofilm-Associated Bacteria to Oxidative
Stress. *Microorganisms* **8**, 253 (2020).

43. J. Chen, S. M. Lee, Y. Mao, Protective effect of exopolysaccharide colanic acid of
Escherichia coli O157:H7 to osmotic and oxidative stress. *Int. J. Food Microbiol.* **93**, 281–
286 (2004).

44. T. Ophir, D. L. Gutnick, A role for exopolysaccharides in the protection of microorganisms
from desiccation. *Appl. Environ. Microbiol.* **60**, 740–745 (1994).

45. K. Mizoguchi, *et al.*, Coevolution of Bacteriophage PP01 and Escherichia coli O157:H7 in
Continuous Culture. *Appl. Environ. Microbiol.* **69**, 170–176 (2003).

46. P. D. Scanlan, A. Buckling, Co-evolution with lytic phage selects for the mucoid phenotype
of Pseudomonas fluorescens SBW25. *ISME J.* **6**, 1148–1158 (2012).

47. S. Wielgoss, T. Bergmiller, A. M. Bischofberger, A. R. Hall, Adaptation to Parasites and
Costs of Parasite Resistance in Mutator and Nonmutator Bacteria. *Mol. Biol. Evol.* **33**,
770–782 (2016).

48.   P. Domenico, R. J. Salo, A. S. Cross, B. A. Cunha, Polysaccharide capsule-mediated
resistance to opsonophagocytosis in Klebsiella pneumoniae. *Infect. Immun.* **62**, 4495–
4499 (1994).

49.   T. A. Russo, *et al.*, The K1 capsular polysaccharide of Acinetobacter baumannii strain 307-
0294 is a major virulence factor. *Infect. Immun.* **78**, 3993–4000 (2010).

50.   M. R. Wessels, A. E. Moses, J. B. Goldberg, T. J. DiCesare, Hyaluronic acid capsule is a
virulence factor for mucoid group A streptococci. *Proc. Natl. Acad. Sci.* **88**, 8317–8321
(1991).

51.   M. Raffatellu, Learning from bacterial competition in the host to develop antimicrobials.
Nat. Med. **24**, 1097–1103 (2018).

52.   C. Rüter, P. R. Hardwidge, 'Drugs from Bugs': bacterial effector proteins as promising
biological (immune-) therapeutics. *FEMS Microbiol. Lett.* **351**, 126–132 (2014).

53.   G. Corno, K. Jürgens, Direct and Indirect Effects of Protist Predation on Population Size
Structure of a Bacterial Strain with High Phenotypic Plasticity. *Appl. Environ. Microbiol.* **72**,
78–86 (2006).

54.   R. M. Donlan, J. W. Costerton, Biofilms: Survival Mechanisms of Clinically Relevant
Microorganisms. *Clin. Microbiol. Rev.* **15**, 167–193 (2002).

55.   A. Schulze, F. Mitterer, J. P. Pombo, S. Schild, Biofilms by bacterial human pathogens:
Clinical relevance – development, composition and regulation – therapeutical strategies.
Microb. Cell **8**, 28–56 (2021).

56.   M. Jamal, *et al.*, Bacterial biofilm and associated infections. *J. Chin. Med. Assoc. JCMA*
81, 7–11 (2018).

57.   S. Galié, C. García-Gutiérrez, E. M. Miguélez, C. J. Villar, F. Lombó, Biofilms in the Food
Industry: Health Aspects and Control Methods. *Front. Microbiol.* **9** (2018).

58.   T. J. Rudge, P. J. Steiner, A. Phillips, J. Haseloff, Computational modeling of synthetic
microbial biofilms. *ACS Synth. Biol.* **1**, 345–352 (2012).

59.   J. Schindelin, *et al.*, Fiji: An open-source platform for biological-image analysis. *Nat.*
Methods **9**, 676–682 (2012).

60.   B. Obadia, *et al.*, Influence of Tyrosine-Kinase Wzc Activity on Colanic Acid Production in
Escherichia coli K12 Cells. *J. Mol. Biol.* **367**, 42–53 (2007).

61.   P. Thevenaz, U. E. Ruttimann, M. Unser, A pyramid approach to subpixel registration
based on intensity. *IEEE Trans. Image Process.* **7**, 27–41 (1998).

62.   A. Agrawal, S. Sinha, R. Mukherjee, D. Mampallil, Dynamics of bacterial deposition in
evaporating drops. *Phys. Fluids* **32**, 093308 (2020).

63.   R Development Core Team, *R: A language and environment for statistical computing.* (R
Foundation for Statistical Computing, 2018).

64. L. H. and K. M. Hadley Wickham, Romain François, dplyr: A Grammar of Data
Manipulation. R package version 0.7.6 (2018).

65. R. M. Hope, Rmisc: Ryan Miscellaneous. R package version 1.5. (2013).

66. P. M. Valero-Mora, *ggplot2: Elegant Graphics for Data Analysis* (Springer, 2010).

67. C. O. Wilke, cowplot: Streamlined Plot Theme and Plot Annotations for "ggplot2". R
package version 0.9.4. (2019).

661