ARHGAP18-Ezrin functions as a RhoA autoregulatory module in the organization of apical microvilli

Authors: Andrew T. Lombardo[1]*, Cameron A. R. Mitchell[1], Riasat Zaman[1], David J. McDermitt[1], Anthony Bretscher[1]

Affiliations:

[1]Department of Molecular Biology and Genetics, Weill Institute for Cell and Molecular Biology, Cornell University; Ithaca, N.Y., USA

*Corresponding author. Email: ATL73@cornell.edu

Abstract: The assembly of a three-dimensional actin cytoskeleton is largely mediated by local Rho GTPases. Restriction of microvilli to the epithelial apical surface requires local regulation of actin structure. On epithelial cells, the presence of microvilli depends on RhoA activation of its effector kinases LOK and SLK that phosphorylate ezrin to function as a membrane-actin linker. Further, previous work has revealed that active ezrin negatively regulates RhoA, but the mechanism was unknown. Here, we show this regulation is mediated by recruitment and activation of ARHGAP18 by active ezrin, which stimulates its GTPase activity towards RhoA. Loss of ARHGAP18 enhances ezrin phosphorylation and results in abundant stubby microvilli that turn over rapidly. This phenotype can be rescued by re-expression of ARHAGP18, and by a construct that targets just the GTPase domain to active ezrin, but not a GAP-dead version. Loss of ARHGAP18 also results in enhanced phosphorylation of myosin light chain, the classic response to elevated RhoA levels. This leads to inappropriate assembly of non-muscle myosin-II into microvilli and the disruption of the normal apical actomyosin organization. Thus, the localization and activation of ARGHAP18 by active ezrin constitutes a local auto-regulatory module to fine tune RhoA to ensure appropriate microvilli organization.

Introduction:

Nearly all vertebrate cells are polarized with regions of morphologically distinct plasma membrane structures supported by directional cytoskeletal filaments. A canonical example is the polarized epithelial cell found along villi of the gut, where microvilli are exclusively found on the apical membrane surface to form a brush border maximizing nutrient absorption. A requirement for this process is the attachment of the plasma membrane to the underlying actin, which depends on members of the Ezrin, Radixin and Moesin (ERM) family of proteins (Bretscher, 1983; Casaletto et al., 2011; Fehon et al., 2010). The human ERMs share a similar domain structure with an N-terminal membrane binding band 4.1 ERM (FERM) domain linked through an α-helical region to a C-terminal ~80 residue domain that binds tenaciously to the FERM domain to mask membrane protein binding sites on the FERM domain and an actin-binding site in the C-terminal domain (Gary & Bretscher, 1995; Turunen et al., 1994) (Fig. 1A). ERMs interact with the apical surface of cells by binding of the FERM domain to phosphatidylinositol 4,5-bisphosphate $(PI(4,5)P_2)$(Fievet et al., 2004). Once there, specific microvilli-targeted kinases LOK and SLK phosphorylate a conserved ERM threonine in the C-terminal domain (T567 in ezrin) (Belkina et al., 2009; Pelaseyed et al., 2017). Phosphorylation then opens ERMs for binding to actin, linking them with the plasma membrane (Matsui et al., 1998; Nakamura et al., 1995; Turunen et al., 1994). This linkage is a requirement for microvilli formation, where the plasma membrane envelops a core bundle of polarity-aligned actin filaments. The actin bundle within microvilli is anchored into a terminal web of actin filaments and non-muscle myosin-2, where force, tension and microvillus length are regulated in a carefully balanced system (Chinowsky et al., 2020; Zaman et al., 2021). A long-standing question in the field has centered around the nucleation and maintenance of these actin networks and the microvilli core bundle. While the microvillus actin nucleator remains elusive, a proposed model suggests that non-muscle myosin-2 regulates microvillus length by depolymerizing the actin core bundle where it embeds into the

terminal actin web (Meenderink et al., 2019). How non-muscle myosin-2 is activated within the terminal web but excluded from the microvillus only a few hundred nanometers away has remained an open question.

Activation of non-muscle myosin-2 occurs via phosphorylation of the regulatory light chains bound to non-muscle myosin-2, allowing for assembly into bipolar filaments of 14-30 individual myosin molecules (Sellers & Heissler, 2019). Phosphorylation occurs predominantly, but not exclusively, through the activity of the RhoA effector Rho-Associated Kinase (ROCK) (Heissler & Manstein, 2013). An intimate signaling connection between Rho-GTPases and ERMs was first suggested in early work using rodent model systems (Tsukita et al., 1994). Understanding of this system was expanded by reports that in the fly moesin acts antagonistically to RhoA (Speck et al., 2003) and in humans RhoA acts upstream of ERM phosphorylation (Hirao et al., 1996). However, unraveling the connections between RhoA signaling and ERMs in humans has been complicated by two factors: first, a controversy to define the relevant endogenous kinases of ERMs (see (Sauvanet et al., 2015)for review) and second, genetic redundancy of these proteins in humans. Humans have three genes specifying the ERM proteins, two activating kinases (LOK and SLK), 20 Rho family of GTPases and at least 49 Rho GTPase activating proteins (ARHGAPs). The functional redundancy of ERMs is most clearly observed in contrasting the relatively mild defects found in moesin knockout mice, where ezrin and radixin can compensate (Doi et al., 1999), with the removal the single ERM homolog moesin in *Drosophila*, which is lethal (Speck et al., 2003).

In humans, both the Rho-GTPase family and ERMs are direct participants in cell proliferation and cancer progression and thus have generated significant clinical attention (Hoskin et al., 2019; Martin et al., 2003). Despite this importance, the relationship between these signaling pathways at the plasma membrane has remained unknown. Two reports have suggested a possible involvement of Rho GTPase-

Activating Protein 18, also called ARHGAP18, in regulating ERM proteins. First, in a proteomic screen in human cells, ARGHAP18 was identified as a candidate interactor of active ezrin (Viswanatha et al., 2013). Second, a two-hybrid study in the fly identified a potential homolog of ARHGAP18, known as Conundrum, by its ability to bind the FERM domain of active moesin and to bind to fly RhoA. Further, loss of Conundrum was shown to affect epithelial morphogenesis, but genetic experiments were unable to define a specific role in moesin regulation (Neisch et al., 2013). Additionally, ARHGAP18 has been shown biochemically to be a specific GTPase Activating Protein (GAP) for RhoA (Maeda et al., 2011). In this study, we show how ARHGAP18, through its recruitment and activation by ezrin, fine-tunes the local level of RhoA to contribute to the appropriate morphogenesis of the apical aspect of epithelial cells.

Results:

ARHGAP18 binds active, open ezrin through its FERM domain:

ARHGAP18 is a specific RhoA GTPase Activating Protein (GAP) (Maeda et al., 2011) that was also identified as a potential ezrin interactor in a proteomic screen by our group (Viswanatha et al., 2013). To characterize the proteomics finding, we expressed a full-length flag-tagged ezrin in the human placental epithelial cell line Jeg3 and performed an anti-flag pulldown experiment (Fig. 1A&B). Endogenous ARHGAP18 co-precipitated with flag-ezrin (Fig. 1B). To explore this interaction further, cell lysates from Jeg3 cells expressing various flag-tagged ezrin constructs were passed over a resin to which bacterially expressed recombinant human-ARHGAP18 was bound. Truncated ezrin 1-479 (FERM) lacking the C-terminal regulatory domain bound at approximately 10-fold the level compared to the full-length protein (Fig.1 C, E). These results indicate that ARHGAP18 binds to a region of ezrin within the FERM domain that is unavailable in the full-length closed protein.

To determine if the interaction between the FERM domain of ezrin and ARHGAP18 was direct, we purified both proteins using bacterial recombinant expression (Fig. S1A). Either purified FERM or full-length ezrin was passed over a SUMO-ARHGAP18 column and the bound and flow through fractions analyzed (Fig. S1A). Purified ezrin FERM domain bound preferentially to the SUMO-ARHGAP18 column compared to full-length ezrin. Using Isothermal Calorimetry (ITC), the *in vitro* binding affinity of purified ARHGAP18 to full-length ezrin or the FERM domain was determined (Fig. S1B). The FERM bound with a dissociation constant of K_d 19.29 ± 3.53μM (mean ± SEM) (Fig. S1B). Therefore, ezrin's C-terminal region inhibits the FERM domain from binding ARHGAP18. To test if activation and opening of full-length ezrin enhanced ARHGAP18's ability to access the FERM domain, both phosphomimetic ezrin-T567E and constitutively open ezrin(1-583) mutants were expressed in Jeg3 cells (Fig. 1C). Ezrin's T567 is the target of the activating kinases LOK and SLK, while the ezrin (1-583) construct is a four amino acid truncation previously shown to disrupt the interaction between ezrin's C-terminal and FERM domains (Viswanatha et al., 2012; Zaman et al., 2021). Both these mutations greatly enhanced ezrin's ability to bind ARHGAP18 compared to the wildtype-ezrin or the non-phoshorylatable ezrinT567A mutant (Fig. 1C, E).

Phosphorylation at ezrin-T567 is abolished in the absence of the kinases LOK and SLK (Zaman et al., 2021). Strikingly, replication of the experiment in cells genetically lacking LOK and SLK showed no binding of full-length wildtype ezrin to ARHGAP18 (Fig. 1D, E). However, phosphomimetics or truncation-activated ezrin constructs were still capable of binding ARHGAP18, albeit at a somewhat diminished level (Fig. 1E). Using the DLD-1 colorectal cell line, the interaction of ezrin and ARHGAP18 was similarly phosphorylation site-dependent (Fig. S2), indicating this interaction was not limited to Jeg3 cells. We conclude that ARHGAP18 interacts with ezrin directly through a region on the FERM domain masked in inactive ezrin but available when phosphorylated at T567 by LOK or SLK.

ARHGAP18 localizes to microvilli and reduces ezrin T567 phosphorylation to regulate apical microvilli organization and dynamics

We next looked at the localization of ARHGAP18-flag in fixed Jeg3 cells using confocal microscopy. Cells expressing the construct had reduced numbers of microvilli depending on the expression level compared to wildtype Jeg3 cells (Fig. 2A). Additionally, the ARHGAP18-flag localized with actin and ezrin within the apical microvilli of the cells (Fig. 2A inset). We were interested if ARHGAP18 localized to just the base of microvilli where its downstream target, RhoA, is believed to be active (Shaw et al., 1998). Increased resolution afforded by Structured Illumination Microscopy (SIM) revealed that the ARHGAP18 signal was throughout the microvilli, overlapping with the ezrin signal (Fig. 2B). These data indicate that ARHGAP18 was targeted specifically to microvilli through its ability to bind active ezrin.

To investigate the effect of loss of ARHGAP18 on the presence of microvilli and ezrin regulation, CRISPR/CAS9 was used to genetically knock out endogenous ARHGAP18 within Jeg3 cells (Fig. 2C, D). Imaging of ezrin in the ARHGAP18$^{-/-}$ cells revealed an increase in the number of microvilli compared to wildtype cells (Fig. 2E, F). Scanning electron microscopy (SEM) of the apical surface of the cells (Fig. 2G) showed numerous smaller but variable-sized microvilli in the ARHGAP18$^{-/-}$ cells compared to the wildtype Jeg3 cells. Since phosphocycling of ERMs at the conserved ezrin T567 site is required for microvilli assembly and maintenance (Viswanatha et al., 2012), western blotting was performed to test if loss of ARHGAP18 influenced levels of ezrin phosphorylation. Knocking out ARHGAP18 resulted in approximately 30% increase in the level of ezrin T567 phosphorylation, while overexpressing ARHGAP18 resulted in the opposite effect (Fig. 2C, D). It should be noted that these are significant effects as in unperturbed cells, the steady state level of phosphorylation is about 50% of total cellular ezrin (Viswanatha et al., 2012).

To see what effect the enhanced level of phospho-ezrin has on microvillar dynamics, we transfected wildtype and ARHGAP18$^{-/-}$ cells with GFP-EBP50, a construct we have used for its excellent live cell imaging of microvilli (Garbett & Bretscher, 2012). In both conditions, microvilli moved across the apical surface with an overall bias for retrograde trajectories from the apical periphery toward the cell center (Movie 1). In the ARHGAP18$^{-/-}$ cells, we observed not only the expected increased numbers of shorter microvilli but also increased turnover of individual microvilli in the knockout cells compared to wildtype cells (Fig. 3A Movie 1). Tracking of the individual microvilli (Fig. 3A blue arrows) showed that in contrast to microvilli in wildtype cells, those in the ARHGAP18$^{-/-}$ cells present at the beginning of a time series regularly disappeared and were replaced by new microvilli (Fig. 3A magenta arrows) over a period of 10 minutes.   We conclude that ARHGAP18 is involved in a negative feedback loop impacting ezrin phosphorylation that regulates microvilli abundance, morphology, and dynamics.

ARHGAP18 function is dependent on ezrin binding and GAP-activity

An attractive model is that ARHGAP18's binding to ezrin targets ARHGAP18 to locally reduce RhoA activity within microvilli. To investigate this model, we expressed various ARHGAP18-flag constructs in the ARHGAP18$^{-/-}$ cells and explored how well they could function to restore normal microvillar organization (Fig. 3B). Expression of full length tagged ARHGAP18 rescued the increased microvilli phenotype back to wildtype levels (Fig. 3C, 2F). On the other hand, expression of a tagged, truncated ARHGAP18 with only the structurally predicted GAP domain (GAP-flag), was more diffuse in its cellular localization and lacked the enrichment to the apical surface as seen with expression of the wild-type protein (Fig. 3C,D side projections). Additionally, ezrin targeting and microvilli formation were severely disrupted in the GAP-flag condition (Fig. 3D). Expression of full-length ARHGAP18 with a point mutation ablating the catalytic 'arginine finger' (ARHGAP18(R356A)-flag) (Fig. 3B) was targeted correctly to microvilli (Fig. 3D inset). However, it had no observable influence on the increased microvilli

phenotype characteristic of ARHGAP18[-/-] cells, indicating that GAP activity is required for rescue (Fig. 3C). These results suggest that ARHGAP18 must be correctly targeted to active ezrin in microvilli to mediate RhoA regulation for the cell to maintain normal microvilli. To test this model, we expressed a chimeric protein with ARHGAP18's GAP domain fused to the 37 amino acid ezrin binding motif of EBP50 (GAP-flag-EBP50(tail)) (Fig. 3B,D) This construct was targeted to microvilli (Fig. 3D inset) and reduced the number of microvilli on ARHGAP18[-/-] cells closer to wild type levels (Fig. 3D).

Ezrin binding to ARHGAP18 enhances its GAP activity to locally reduce active RhoA inside microvilli

We next examined whether binding of ezrin to ARHGAP18 regulated its RhoA GAP activity. Using an *in vitro* RhoA GAP colorimetric assay that measures free phosphate (Pi) released by RhoA-dependent hydrolysis of GTP to GDP, the intrinsic GAP activity of recombinant RhoA was found to be enhanced about four-fold in the presence of purified ARHGAP18 (Fig. 4A, B). This activity was about half the rate obtained with the purified control Rho GTPase-activating protein 1(p50RhoA GAP) (Fig. 4A, B). When the ezrin FERM domain was included with ARHGAP18, the GAP activity was enhanced about two-fold to the level of the control p50RhoA GAP. We conclude that the binding of ARHGAP18 to ezrin's FERM domain activates its catalytic activity thereby enhancing its activity to microvilli where active ezrin exists. Activation in this way would limit the potency of its GAP activity to areas of the cell where active, open ezrin exists.

Loss of ARHGAP18 should increase the levels of active RhoA at the apical surface of the cells, but due to the presence of numerous potentially redundant GAPs it was not clear what effect this might have. To visualize active RhoA, we expressed a biosensor, AHPH-GFP, that preferentially binds active RhoA-GTP (Piekny & Glotzer, 2008). In wildtype cells, the GFP signal was localized to cell-cell junctions as reported (Priya et al., 2015) (Fig. 4C, yellow arrowheads). Expression of the biosensor in ARHGAP18[-]

/- cells showed a reorganization of RhoA-GTP to the apical surface of the cell, though some signal remained at junctions (Fig. 4C). Next, western blotting on cell lysates was used to test the expression and activation of RhoA downstream targets. Expression of RhoA itself remained unchanged in the knockout cells (Fig. 4D). Interestingly, one target of RhoA activation, phosphorylation of myosin light chain (pMLC), was increased in ARHGAP18[-/-] cells (Fig. 4D, E).

Immunofluorescence detection of pMLC is typically observed in contractile actomyosin bundles or networks where Nm2 is active. When we imaged the apical surface of wildtype jeg3 cells, pMLC was found along the cell-cell junctions and in punctate structures near the terminal actin web, just below the apical membrane (Fig. 4F inset). This observation agrees with reports of active non-muscle myosin-2 localizing to the region directly below the microvilli within the terminal web of actin, where it depolymerizes the core bundle and regulates microvilli length (Meenderink et al., 2019). However, pMLC in the ARHGAP18[-/-] cells was inappropriately localized inside microvilli where it overlapped with the actin core bundles (Fig. 4F inset). SIM images for non-muscle myosin-2b similarly showed a rearrangement in the ARHGAP18[-/-] cells compared to wildtype cells which could be rescued by the expression of ARHGAP18-flag (Fig. S3). Thus, ARHGAP18 is necessary to modulate local RhoA activity in microvilli to ensure that myosins are not inappropriately assembled there, while ensuring sufficient RhoA levels to locally activate LOK and SLK to phosphorylate ezrin.

Another notable feature of ARHGAP18[-/-] cells is that the actin architecture of the terminal web appears to be disrupted (Fig. 4F inset). Characterization of this network is limited by the resolution of conventional imaging techniques. We, therefore, employed Stochastic Optical Reconstruction Microscopy (STORM) on regions where the cells expanded out away from the monolayer, becoming thinner in height. In these regions, the apical surface could be imaged using total internal fluorescence microscopy, allowing for STORM reconstructions with approximately 40nm resolution in the x-y dimensions. Under these

conditions, we observed a dramatic reorganization of the actin architecture in favor of smaller branched or individual actin filaments in the ARHGAP18$^{-/-}$ cells compared to wildtype cells (Fig. 4H). Loss of ARHGAP18 resulted in a near total depletion of large, bundled actin in this region which are dependent on RhoA regulation of non-muscle myosin-2 activation. However, smaller bundles and branched networks remained. These results collectively promote a model where ARHGAP18 acts as a negative regulator of RhoA locally within regions where ERMs are active. This activity regulates non-muscle myosin-2 activation and the actin at the apical surface and within microvilli. ARHGAP18 is regulated downstream of LOK/SLK, where activation of ERMs both localizes ARHGAP18 and enhances its GAP activity (Fig. 5).

Discussion:

ARHGAP18 limits the activity of RhoA in microvilli through binding to active ezrin.

This study reveals that active ezrin directly recruits and activates ARHGAP18 to microvilli to locally regulate RhoA activity for appropriate ezrin phosphorylation and to limit apical actomyosin organization. In earlier work we characterized the Jeg3 cells used here lacking expression of ERMs or LOK/SLK and demonstrated that LOK/SLK are the major kinases that phosphorylate ERM proteins in humans. We also showed that a constitutively active form of RhoA directly binds LOK (Zaman et al., 2021). An independent proteomic screen identifying Rho GTPase family protein interactors showed that RhoA directly binds SLK leading to dimerization and autophosphorylation (Bagci et al., 2020). These results define an activation pathway where RhoA activates LOK/SLK that then activates ERMs. Additionally, we found that in the absence of LOK/SLK or ERMs, the level of active RhoA was elevated, indicating that active ERMs in some way down-regulates active RhoA (Zaman et al., 2021). These findings indicated a missing negative regulatory feedback loop through an unknown effector, an idea supported by studies in the genetically simpler fly model (Speck et al., 2003). As introduced earlier,

ARHGAP18 was identified both in our proteomic screen for active ezrin interactors (Viswanatha et al., 2013), and its probable fly homolog Conundrum in a two-hybrid screen with fly moesin (Neisch et al., 2013). We find that ARHGAP18 binding to active ezrin activates the RhoA-GAP activity of ARHGAP18, and this activity is necessary for appropriate regulation of the apical domain (Fig. 5).

Potential mechanism of apical regulation through RhoA, LOK/SLK, ERMs and ARHGAP18

Cells lacking either LOK/SLK or all ERMs have no apical microvilli, underscoring the important roles of these two classes of proteins (Zaman et al., 2021). ERMs interact through their FERM domain with PI(4,5)P2 which is preferentially enriched at the plasma membrane. Biochemical reconstitution studies show that activation of ERMs by LOK or SLK then occurs through a co-incidence detection mechanism whereby ERMs are activated through phosphorylation only when bound simultaneously to LOK/SLK and PI(4,5)P2 (Pelaseyed et al., 2017). Once active the ERMs serve a required function of linking the plasma membrane to the actin core bundle formed within the microvilli. Our data show that ARHGAP18 is recruited to the microvilli through its preferential binding to the open active ERM's FERM domain (Fig. 1C). The localization of ARHGAP18 in microvilli is particularly interesting as we are unaware of any other GAP or GEF that localizes specifically to microvilli. An independent screen of brush border enterocytes proteins also did not identify one; however, the study did identify Rho precursors to isoforms A, B, C & G (McConnell et al., 2011). These results suggest RhoA is not excluded from microvilli but instead kept modulated locally by ARHGAP18. RhoA is active just below the microvilli within the terminal web of actin (Shaw et al., 1998), and ARHGAP18's ability to enhance its GAP activity when bound exclusively to active, open ERMs within microvilli indicates that the microvilli are regulated as a separate RhoA microdomain on the individual microvilli scale (200-1000nm). This model may represent a broader mechanism employed to regulate spatiotemporal coordination of Rho GTPases across multiple cell surface protrusions through various GEFs and GAPs. For example, the ARHGAP35 gene

product p190A has been reported to localize to cilia where Rho activity is required for proper cilia formation (Stewart et al., 2016), RhoA acts specifically at sites of protrusion initiation in lamellipodia (MacHacek et al., 2009) and SRGAP1 controls podocyte morphology through localized regulation of Rho activity (Rogg et al., 2021).

Cells lacking ARHGAP18 have enhanced RhoA activity that increases ezrin phosphorylation and increases microvilli, showing that ARHGAP18 negatively regulates the RhoA/LOK/SLK pathway. RhoA also regulates actin and non-muscle myosin-2 through the ROCK pathway. Loss of LOK/SLK, ERMs, and ARHGAP18 all result in increased active RhoA at the apical surface and downstream increased activation of non-muscle myosin-2(Fig. 4C-E) (Maeda et al., 2011; Porazinski et al., 2015; Zaman et al., 2021). Increased activation of non-muscle myosin-2 results in depolymerization of the microvilli core bundle within the terminal web of actin (Meenderink et al., 2019). In cells lacking ARHGAP18, this results in the increased turnover of microvilli (Fig. 4G) as the actin core bundles are depolymerized more quickly by increased non-muscle myosin-2 activity. Thus, by combining these two effects in ARHGAP18$^{-/-}$ cells, the number of microvilli is enhanced as a result of increased active ERMs, yet microvilli length is decreased as a result of increased non-muscle myosin-2 activity within the terminal web.

Although there are no microvilli in cells lacking LOK/SLK or ERMs, cells lacking ARHGAP18 are similar in that the terminal web's actin organization is disrupted; all have the increased non-muscle myosin-2 activity, and the stiffness of the cells is similarly increased as a result (Zaman et al., 2021) (Fig. 4F, H) (Fig. S4). However, the disorganization of the actin in this region differs between the ARHGAP18$^{-/-}$ cells, which show no large stress-fiber-like bundles (Fig. 4H), and the cells lacking active ERMs which express excessive large contractile actin bundles at the apical surface (Zaman et al., 2021). Our result of reduced contractile bundles in ARHGAP18$^{-/-}$ cells agrees with previous reports using siRNA knockdown of ARHGAP18 (Porazinski et al., 2015). One explanation for this apparent discrepancy may relate to the

availability of G-actin. Removal of all microvilli in the LOK[-/-]SLK[-/-] cells would leave an increased abundance of G-actin that would otherwise be sequestered into microvilli bundles. On the other hand, the increased number of microvilli in ARHGAP18[-/-] cells may deplete this pool resulting in a change in the F-actin assembly and organization.

It is of particular interest to note that the overall mechanism proposed here was correctly speculated previously. To explain the phenotypes observed in *Drosophila* moesin mutants, Neisch *et al.* suggested that it is the joined effects of increased RhoA dependent apical actomyosin contractility and loss of moesin mediated membrane-cytoskeleton cross-linking activity (Neisch et al., 2013). Our data support this model within the human system, and additionally provide characterization and localization of the specific interactors facilitating these signaling interactions at the apical surface.

Implications for the local regulation of Rho proteins.

A major gap in our understanding of active Rho GTPases, that are free to diffuse in the plasma membrane, is how they select and discriminate between their many effectors (reviewed in (Denk-Lobnig & Martin, 2019)). This discrimination could be mediated by effectors with different affinities for active Rho, or their abundance, or location, or any combination of these. In the simple case investigated here, where active RhoA must discriminate between the LOK/SLK pathway to activate ERM proteins and the ROCK pathway that leads to myosin activation, we now have some insights into how this may occur. In the case of LOK/SLK, these effector kinases are specifically targeted to the apical membrane by the C-terminal half of the molecule (Pelaseyed et al., 2017; Viswanatha et al., 2012). However, as revealed by the current study, this selective targeting of the kinase is not sufficient for functional discrimination by active RhoA, as loss of ARHGAP18 results in the over-activation of ezrin and inappropriate assembly of non-muscle myosin-2. Thus, targeting of ARHGAP18 to microvilli by ezrin is necessary to locally tune the level of active RhoA to ensure that sufficient ezrin is phosphorylated, while ensuring non-muscle myosin-2 is not

activated. An attractive mechanism could be to locally reduce the level of active RhoA to a level commensurate with the local microvillar abundance of LOK/SLK, thereby driving selectivity by mass action. As part of this regulation, the level of active ezrin is itself determined by the activity of RhoA through LOK/SLK, and that in turn determines the recruitment and activity of ARHGAP18, which then negatively regulates RhoA (Fig. 5). This auto-regulatory loop would ensure that excessive RhoA activity is immediately modulated down to the appropriate level, and too little activity would be modulated up to the appropriate level. This highly localized auto-regulatory loop thereby fine-tunes RhoA activity for the appropriate morphogenesis of the apical membrane. We expect that similar fine-tuning of the activity of other small GTPases by down-stream targets will emerge as more is learned about the GAP proteins that vastly out-number the small number of GTPases they regulate.

Figure 1: ARHGAP18 binds active, open ezrin through the FERM domain. A) Protein domain schematics of human ARHGAP18 and ezrin. B) Western blot from pulldown experiment from passing WT-Jeg3 placental epithelial cell lysate over an anti-flag resin then blotting against flag-tagged ezrin and endogenous ARHGAP18. Ezrin tagged internally (iflag) within a flexible linker region at residue 479. C) Western blot of ARHGAP18 bound resin pulldown experiment blotting against ezrin constructs expressed in WT-Jeg3 cells. D) Western blot of ARHGAP18 bound resin pulldown experiment blotting against ezrin-constructs expressed in LOK[-/-]SLK[-/-]-Jeg3 cells. E) Quantification of normalized band intensity from experiments presented in C & D. Bars represents mean ± SEM; n=4. WT and LOK[-/-]SLK[-/-] conditions significantly different (Ratio paired t-test; p=0.0235).

Figure 2: ARHGAP18 localizes to microvilli and alters ezrin T567 phosphorylation. A) Confocal immunofluorescence images of WT-Jeg3 cells expressing ARHGAP18-flag. Scale Bar 10μm; inset 2μm B) Structured Illumination Microscopy (SIM) reconstructions of WT-Jeg3 cells expressing ARHGAP18-flag with zoom in on boxed area. Scale bar 10μm; inset 2μm. C) Western blot showing CRISPR KO of endogenous ARHGAP18 and ezrin pT567 phosphorylation relative to total ezrin. Antibody against ARHGAP18 was confirmed in part by identification of endogenous ARHGAP18, GFP-ARHGAP18 and purified ARHGAP18 while showing no detection in KO cells. D) Quantification of normalized band intensity from experiments presented in C. Bars represents mean ± SEM; n=4 paired t-test; *, P ≤ 0.05). E) SIM images of WT vs. ARHGAP18$^{-/-}$ Jeg3 cells with ezrin as a marker for the abundance of microvilli at the apical surface. F) Quantification of the density of microvilli on the surface of Jeg3 cells in WT, ARHGAP18$^{-/-}$, and ARHGAP18 overexpression conditions. Microvilli counts per cell were normalized to the area of each cell to account for cell size variability. Bars represents mean ± SEM; n=3 replicates; t-test; **, P ≤ 0.001). G) Scanning Electron Microscopy (SEM) micrographs of WT-Jeg3 cells' apical surface compared to those of ARHGAP18$^{-/-}$ cells. Left panels imaged at 4000x; scale 5μm. Right panels imaged at 10000x within of the boxed areas; scale 1μm.

Figure 3: ARHGAP18 KO rescue is dependent on ezrin binding and GAP-activity A) First frame of Movie 1 with inset zoom in of the magenta box. Each frame at shows the tip of the microvilli present in the inset image at t=0 (blue arrowhead). In the ARHGAP18$^{-/-}$ condition the original microvilli disappear and are replaced by new microvilli not present in the first image (magenta arrowheads). Scale bars 10μm; inset 1μm. B) Schematic of constructs used in this figure. C) Immunofluorescence confocal images of ARHGAP18$^{-/-}$ Jeg3 cells expressing flag-tagged WT-ARHGAP18-flag (rescue) or the control flag vector. XY images above; side projections showing apical/basolateral localization below. Scale bars 10μm D) Immunofluorescence confocal images of ARHGAP18$^{-/-}$ Jeg3 cells expressing ARHGAP18(319-523) GAP domain only, a point mutant GAP-dead variant or a chimera variant of the GAP domain of ARHGAP18(319-523) linked to the ezrin binding motif of EBP50(486-523). Expressing the GAP domain alone does not target to microvilli, disrupts microvilli formation and ezrin targeting to the apical surface. The GAP-dead variant is localized to microvilli and apical ezrin but does not rescue the knockout microvilli phenotype (zoomed inset). GAP-flag-EBP50(tail) targets to microvilli (inset), is sufficient to reduce microvilli abundance in the ARHGAP18$^{-/-}$ cells and ensures ezrin localization to the apical surface. Scale bars 10μm, insets 2μm. E) Quantification of the density of microvilli on the surface of ARHGAP18$^{-/-}$ cells expressing the conditions shown in C&D. Microvilli counts per cell were normalized to the area of each cell to account for cell size variability. ARHGAP18$^{-/-}$+ARHGAP18-flag condition duplicated from Fig. 2F for comparison between conditions. Bars represents mean ± SEM; n≥3; t-test; *, P ≤ 0.05). Side projections expanded in Z-dimension for clarity.

A

ARHGAP18   1   319   R365 523 663
                                     RhoA GAP

p50RhoAGAP   1   195
Catylitic domain   RhoA GAP

Ezrin   1   296
FERM   FERM

RhoA   1   193
            RhoA

B

Relitive Absorbance 550nm (A.U)

RhoA   - + - - - - + + + +
ARHGAP18   - - + - - + + + + -
FERM   - - - + - + - - + -
p50RhoAGAP   - - - - + - - - - +

C

WT Jeg3   ARHGAP18 −/−

AHPH-GFP
(Active RhoA BioSensor)

Apical-
Basal-

D

                WT Jeg3   ARHGAP18 −/−

15   pMLC
15   MLC
      RhoA
15
55   Actin
35
55   Tubulin

E

pMLC/MLC band Intensity(A.U.)

2.0
1.5
1.0
0.5
0.0
        WT Jeg3   ARHGAP18 −/−

F

        Actin   pMLC   Merge/DAPI   Inset

WT Jeg3

ARHGAP18 −/−

G

STORM Actin Reconstruction

WT Jeg3   ARHGAP18 −/−

2 µm   2 µm

Figure 4: Ezrin binding to ARHGAP18 enhances GAP activity excluding active RhoA inside microvilli A) Cartoon schematic of purified proteins used in (B) B) Representative image of RhoA GAP assay and graph of combined results from multiple assays. Each assay normalized to the buffer only condition. Bars represents mean $\pm$ SEM; n=4; t-test; (*, P $\leq$ 0.05).C) Z-projection confocal fluorescent images of active RhoA biosensor. Side projections of magenta area shown below each image. Yellow arrowhead indicates active-RhoA signal at cell-cell junctions. Scale bars 10μm. Side projections expanded in Z-dimension for clarity. D) Western blotting indicating changes in pMLC in response to altered expression of ARHGAP18 in Jeg3 cells. E) Quantification of pMLC over total MLC from replicates of the data in (D) F) Bars represents mean $\pm$ SEM; n=3; t-test;(*, P $\leq$ 0.05). F) immunofluorescence staining of pMLC in WT Jeg3 cells vs ARHGAP18$^{-/-}$ cells inset showing localization to the microvilli in the KO cells. Scale bars 10μm; inset 2μm. G) Super resolution TIRF/STORM reconstructions of actin in single Jeg3 cells. Scale 2μm.

Figure 5: Model of Ezrin and RhoA feedback loops in microvilli. (**1**) RhoA is activated and binds to and activates LOK and SLK kinases. These kinases are localized specifically to the microvilli of epithelial cells; (**2**) In cooperation with PIP2, the kinases phosphorylate and activate ezrin. Once active, ezrin links the plasma membrane to the actin core bundle within microvilli and also makes available a binding site for ARHGAP18 (**3**) ARHGAP18 binds the FERM domain of ezrin and this activates its RhoA-GAP activity (**4**) This activity locally reduces the concentration of active-RhoA in microvilli, thereby modulating the ability of RhoA to activate LOK to phosphorylate ezrin. (**5**) Inactive Rho-GDP cannot bind or activate LOK kinase thus completing a negative feedback system (top center of figure). (**6**) In the terminal web, RhoA-GTP activates the assembly of non-muscle myosin-II through the ROCK pathway (**7**) regulating microvilli length and apical actin organization. Loss of ARHGAP18 results in increased RhoA activation inside the microvilli. The consequences are locally increased active ERMs (**7**) and increased non-muscle myosin-2 assembly (**8**) inside microvilli. These changes result in increased microvilli formation (through ERMs) and disruption of the F-actin core bundle within microvilli (through non-muscle myosin-2).

Methods

Western blotting:

Western blotting was done using 12-8% split SDS-PAGE gels except for proteins less than 30kDa that used 14% SDS-PAGE gels for better resolution. The gels were transferred at 16V to 0.45µm pore size polyvinylidene difluoride (PVDF) membranes (EMD Millipore Immobilon-FL; catalog no. IPFL85R), or 0.2µm pore size PVDF membranes (EMD Millipore Immobilon-P^{SQ}; catalog no. ISEQ08130) for smaller proteins less than 30kDa. Cell lysates were washed and harvested with warm Laemmli sample buffer (70°C) by scraping then boiling. Membranes were blocked with 5% nonfat milk in Tris-buffered-saline with 0.1%tween (TBS-T), except for phosphorylation-specific antibodies which used Immobilon Block – PO phosphoprotein blocking buffer (EMD Millipore; catalog no. WBAVDP001). Primary antibodies diluted in either 5% BSA in TBS-T or in Immobilon Block – PO were incubated with the membrane at room temperature for 1 hour, or at 4°C overnight. Signal was detected using fluorescent secondary antibodies, anti-mouse Alexa Fluor IRDye 680RD, and anti-rabbit Alexa Fluor IRDye 800CW (LI-COR Biosciences; catalog nos. 926-68070 and 926-32211) at 1:10,000 in 5% nonfat milk. Blots were imaged using a Bio-Rad ChemiDoc, and band intensities were determined with ImageJ's gel analysis package by normalizing band intensity and calculating relative intensities. Gel band intensities were then input to GraphPad Prism for quantification and statistical analysis.

ARHGAP18 and ezrin iFlag pulldown assays:

In order to determine the interaction between ARHGAP18 and ezrin, pulldown assays were performed using cell lysate from Jeg3 wild-type and Jeg3 LOK$^{-/-}$SLK$^{-/-}$ cells. Transfections were performed using PEI MAX® polyethylenimine reagent (Polysciences; catalog no. 24765). After transfection, cells were washed with PBS and harvested with lysis buffer (25mM Tris, 5% glycerol, 150mM NaCl, 50mM NaF, 0.1mM Na3VO$_4$,10mM βGP, 0.2% Triton X-100,250mM Calyculin A, 1mM DTT, 1X complete protease inhibitor cocktail (Roche Cat no. 11836153001) by scraping. Lysates were then sonicated, and any insoluble material was centrifuged at 8000xg for 10 min at 4°C. Before incubating with the cell lysates, SUMO-ARHGAP18 NiNTA or M2-Flag resin beads were equilibrated and washed with wash into lysis buffer. The a sample of the supernatant was taken for input then the rest added to the SUMO-ARHGAP18 NiNTA or M2-Flag beads and nutated for 3h at 4°C. After incubation, the beads were pelleted and washed 4x before boiling in 40µL 2X Laemmli sample buffer.

Cell culture:

Jeg3 cells originally obtained from ATCC.org (catalog no. Htb-36) were maintained in a humidified incubator at 37°C and 5% CO$_2$. Jeg3 cells were maintained in 1X MEM (Thermo Fisher Scientific; catalog no. 10370088) with penicillin/streptomycin (Thermo Fisher Scientific; catalog no. 15070063), 10% fetal bovine serum (Thermo Fisher Scientific; catalog no. 26140079) and GlutaMAX (Thermo Fisher Scientific; catalog no. 35050061). Cultures were maintained on Corning 100 x 20mm style culture dishes (catalog no. 430167). Transient transfections were accomplished using a PEI MAX polyethylenimine reagent (Polysciences; catalog no. 24765). DLD-1 cells originally obtained from ATCC.org (catalog no. CCL-221) were handled identically with the exception of using DMEM media.

Crispr knockout lines:

CRISPR analysis tools from Benchling Inc. (San Francisco, CA) were used to design Single-guide RNAs (sgRNAs). Guides were cloned into puromycin-resistant pLenti-CRISPRV2 (Addgene; catalog no. 49535) as described in (Sanjana et al., 2014).The sgRNA sequence used against the first exon of human ARHGAP18 was 5′- CAGCGGCAAGGACCAGACCG -3′ and the construct was transfected into 293TN cells with psPAX3 and pCMV-VSV-G (a gift from Jan Lammerding, Weill Institute for Cell and Molecular Biology, Cornell University, Ithaca, NY) for 48–72 h before virus collection. Jeg3 cells were then transduced with viral media supplemented with Polybrene to 8µg/mL twice a day for 2 days prior to puromycin selection at 2µg/mL. Single cell sorting was performed then expanded in puromycin selection before screening by western blotting.

Microvilli quantification:

To determine if ARHGAP18's presence influences the abundance of microvilli in Jeg3 cells, cells were transfected with wild-type ezrin as a marker for microvillar abundance. The abundance of microvilli was quantified using ImageJ's cell counter toolkit and total microvilli counts for each cell were normalized to their respective cell areas to account for variability in cell sizes. Values were then input to GraphPad Prism for quantification and statistical analysis.

Fixed cell preparation:

Jeg3 cells were grown on glass coverslips prior to fixing cells for immunofluorescence. Cells were fixed in 3.7% formaldehyde for 10 minutes at room temperature, then washed in PBS three times. Cell membranes were permeated using 0.2% Triton X-100 diluted in phosphate buffered saline (PBS). Blocking was performed using 2% FBS in PBS for 10 minutes. Primary and secondary antibodies were incubated on the cells for 1 hr. at room temperature each and were diluted in the 2% FBS blocking solution. To visualize actin or the nucleus, Alexa Fluor 647 Phalloidin at 1:25 dilution in methanol and DAPI at 1:10000 dilution in H20 were included with the secondary antibody incubation. The cells were mounted with either SlowFade Diamond Antifade (Thermo Fisher Scientific; catalog no. S36967), or Prolong Diamond Antifade (Thermo Fisher Scientific; P36965). For AHPH-GFP (Addgene; catalog no. 68026) active RhoA biosensor cells were fixed using 10% trichloroacetic acid (TCA) instead of formaldehyde.

SEM imaging:

SEM imaging was performed at the University of Vermont Imaging Facility (Burlington, VT). Cells were grown on Thermonox coverslips fixed using Karnovski's fixative (2.5% glutaraldehyde, 2% Paraformaldehyde 0.1M cacodylate buffer, pH 7.2) for 60 min at 4°C then rinsed 4 times in 0.1M cacodylate buffer, pH 7.2. Cells were then post-fixed with 1% osmium tetroxide in 0.1M cacodylate buffer, pH 7.2 for 1 hr. at 4°C followed by 3 rinses. Cells were then immersed in 1% tannic acid in 0.05M cacodylate buffer, 1hr at RT before rinsing into 0.05M cacodylate buffer then into water. Cells were then incubated with 0.5% uranyl acetate in MilliQ water for 1 hr. at room temperature, rinsed into water then stored overnight in 0.05M cacodylate buffer at 4° C. Cells were then dehydrated using a series of increasing ethanol concentration washes until equilibrated with 100% anhydrous ethanol where the cells were then critical point dried with liquid CO_2. The samples were then mounted onto aluminum specimen mounts with a conductive carbon pain and dried overnight using desiccation. The desiccated cells were

then sputter coated with gold/palladium in a polaron sputter coater (Model 5100) and stored under dissection conditions. Prepared cells were then imaged with a JSM-6060 scanning electron microscope from JEOL USA, Inc. (Peabody, MA).

Confocal imaging:

Fixed cell confocal imaging was done at room temperature (23°C) and was accomplished using a spinning-disk (Yokogawa CSU-X1; Intelligent Imaging Innovations) Leica DMi600B microscope with a spherical aberration correction device and either a 100×/1.46 NA or 63×/1.40 NA Leica objective. Images were captured using a Hamamatsu ORCA-Flash 4.0 camera metal-oxide semiconductor device, and Z-slices of acquired images were assembled using SlideBook 6 software (Intelligent Imaging Innovations). Maximum- or summed-intensity projections were then assembled in SlideBook 6 or ImageJ and exported to Adobe Illustrator for editing. Side projections were vertically expanded using Illustrator to increase visual clarity of apical/basal localization.

Live imaging:

Live-cell imaging of cells expressing GFP-EBP50 (Garbett & Bretscher, 2012) was done using an inverted Leica DMi8 wide field microscope equipped with a Leica 100× NA air objective, a Leica DFC 9000 GTC camera, Leica Application Suite X THUNDER deconvolution software, and Leica adaptive focus control. Jeg3 cells were transiently transfected 24h prior to imaging, and cell medium was changed to FluoroBrite DMEM (Thermo Fischer Scientific; catalog no. A1896701) with 10% FBS immediately before imaging. The cells were then kept in a 5% CO_2 humidified environmental chamber at 37°C throughout the duration of the imaging as cells were imaged every 1 minute for 15 minutes. Deconvolution images of GFP-EBP50 were performed on the Leica DMi8 wide field microscope, Leica Application Suite X THUNDER deconvolution software, and Leica adaptive focus control listed above. The default small sample Leica THUNDER deconvolution settings were used for this, apart from adjusting structure size to 1,000 nm and reducing deconvolution strength to 30%.

Statistical methods:

Statistical comparisons were performed in GraphPad Prism. The type of statistical tests utilized, as well as the number of independent data points (*n*) and the configuration of error bars are detailed in the figure legends respective to the data tested. Choice technical or biological replicas decided on an experiment-by-experiment basis. Non-parametric or parametric testing was justified through assumption of the tested data to have distribution or not. Any additional analyses related to specific techniques are described in the materials and methods sections.

Super resolution SIM and STORM:

Structured Illumination Microscopy and STORM imaging were performed by using a Zeiss Elyra super-resolution inverted Axio Observer.Z1 microscope provided by Cornell University's Institute of Biotechnology Imaging Facility. Illumination was performed using lasers emitting 405-, 488-, 561-, and 640-nm wavelengths through a 63×/1.4 NA oil objective and capturing images on a pco.edge 5.5m camera. Exposure time ranged from 50 to 400 milliseconds and was dependent on sample and channel to optimize SIM reconstruction. As calculated through the ZEN black software, the number of Z-slice steps was set to the optimized minimum for each illumination channel. SIM grating was set to five rotations for all

conditions, and processing was completed using default settings from the automatic SIM processing toolset provided by the ZEN software. By applying the built-in color alignment, ZEN software to a Z-stack of images containing 100-nm multicolor beads from an Invitrogen Molecular Probes TetraSpeck Fluorescent Microspheres Size Kit (catalog no. T14792), color channels were able to be corrected for chromatic aberration. Preparation of cells for STORM imaging of actin in Jeg3 cells was performed as described in (Heaslip et al., 2014) and reconstructions of STORM imaging were performed using the Zeiss ZEN analysis software.

Reagents and cDNA:

Ezrin antibodies were a mouse anti-ezrin antibody (Developmental Studies Hybridoma Bank; catalog no. CPTC-ezrin-1; Research Resource Identification AB_2100318) used at 1:1,250 (western blot) or 1:100 (immunofluorescence), or a previously characterized polyclonal rabbit antibody raised against full-length human ezrin (APB B64) used at 1:1,000 (western blot) or 1:200 (immunofluorescence) (Zaman et al., 2021). A previously characterized home-made rabbit anti-phospho-ERM, raised against recombinant phosphopeptide, was used at 1:1,000 (western blot) (Zaman et al., 2021). Flag antibodies were rabbit anti-Flag (Cell Signaling Technology; catalog no. 14793) used at 1:500 (western blot) or 1:500 (immunofluorescence), and mouse anti-flag M2 (Sigma-Aldrich; catalog no. F1804) which was used at 1:1,000 (western blot) or 1:200 (immunofluorescence). Mouse anti-GFP (B-2) (Santa Cruz Biotechnology, Inc.; catalog no. SC-9996) was used at 1:1,000 (western blot). Mouse anti-actin C4 (EMD Millipore; catalog no. MAB1501) was used at 1:1,000). Rabbit anti-non muscle myosin-IIB (BioLegend; catalog no. 909902) was used at 1:100 (western blot and immunofluorescence). Anti-myosin light chain antibodies were rabbit anti-MLC2 (catalog no. 3672) used at 1:500 (western blot) and 1:50 (immunofluorescence) and rabbit anti–phospho-MLC2 raised against phospho-Thr18/Ser19 (Cell Signaling Technology; catalog no. 3674) used at 1:500 (western blot) and 1:50 (immunofluorescence). Anti-cofilin antibodies were rabbit anti-phospho-cofilin (Abcam; catalog no. ab12866) used at 1:1,000 (western blot). Rabbit anti-RhoA (Cell Signaling Technology; catalog no. 2117) was used at 1:500 (western blot). Mouse anti-tubulin (Sigma-Aldrich; catalog no. T5168) was used at 1:5,000 (western blot). To stain actin, Alexa Fluor anti-actin 647 phalloidin (Invitrogen; catalog no. A30107) was used at 1:25 (immunofluorescence). For nuclear staining, DAPI (Invitrogen; catalog no. D1306) was used at 1: 250 (immunofluorescence). Secondary antibodies used for immunofluorescence were anti-mouse Alexa Fluor 488 (LI-COR Biosciences; catalog no. A21202) and anti-rabbit Alexa Fluor 568 (LI-COR Biosciences; catalog no. A11036), both used at 1:250. Calyculin A was purchased from Enzo Life Sciences (catalog no. BML-El92-0100).

Human ARHGAP18 constructs were created using Polymerase Chain Reaction (PCR) using New England Biolabs (NEB) Phusion High-Fidelity PCR Kit (catalog no. E0553L) off a construct originally derived from the Harvard Plasma Database (ID # HsCD00379004). PCR and Gibson Assembly products were purified using NEB Monarch PCR & DNA Cleanup Kit (catalog no. T1030S) and Thermo Scientific GeneJET Gel Extraction Kit (Catalog no. K0692). These products were cloned into the mammalian expression vector PQCXIP using NEB Gibson Assembly cloning Kit (catalog no. E5510S), then transformed into OneShot TOP10 bacteria (ThermoFisher Catalog no.C404010). Successfully cloned constructs were selected from using ampicillin resistance then sequenced for verification. Single sight mutagenesis for the gap-dead construct was performed using New England Biolab Q5 Site-Directed

Mutagenesis Kit (Catalog no. E0554S). EBP50 tail sequence was generated by PCR off the GFP-EBP50 construct described below and inserted into the GAP-flag-EBP50(tail) vector using Gibson assembly. GFP tagged ARHGAP18 used as a control in antibody quality control was a generous gift from Jennifer Gamble at the University of Sydney. Transfections were performed using PEI MAX® polyethylenimine reagent (Polysciences; catalog no. 24765).

ARHGAP18 antibody:

Purified human ARHGAP18 was produced by bacterial expression of an N-terminal-SUMO-HIS tagged protein purified using a NiNTA resin. The SUMO tag was cleaved using purified ULP1 and then the ARHGAP18 was further purified using a HiLoad 16/600 superdex 200 size exclusion chromatography installed on a General Electric AKTA pure FPLC. For antibody production this purified protein was exchanged into PBS and denatured by adding 1mM DTT and then boiled for 5 min at a final concentration of 2.1 mg/ml before flash freezing in liquid nitrogen. The frozen ARHGAP18 antigen was then shipped to Pocono Rabbit Farm & Laboratory, Inc (Canadensis, PA) for antibody production in rabbits. ARHGAP18 antibody specificity was confirmed by western blot against the antigen, the ARHGAP18[-/-] cells and observation of a band shift when expressing a GFP tagged variant of ARHGAP18 in Jeg3 cells (Fig. 2C). The used protocol was approved by Cornell University IACUC number 2014-0109 to A. Bretscher.

Isothermal calorimetry (ITC):

Purified ARHGAP18 was produced as described in the antibody production except that the SUMO tag was not cleaved as it was found that the protein was insoluble above 2mg/ml without it. Purified ezrin and FERM were produced as described in (Viswanatha et al., 2012). Ezrin or FERM were diluted to 75uM-80uM depending on the purified volume and number of injections needed for ITC. The proteins were then injected into 30uM SUMO-ARHGAP18 using automated injections on a TA Instruments-Waters LLC (New Castle, DE) Affinity ITC-LV. Injections were performed using ITC run software (TA Instruments) with a minimum pulse time of 200s and 2-5ul injections and a stirring rate of 175 rotations per minute. The system was calibrated using the injection of water into water as a negative control and the injection of 0.95mM $CaCl_2$ into 0.150mM EDTA as appositive control. Injections were continued until saturation of the binding was observed through a plateau of the released energy per injection or the max volume of injection syringe was depleted.. Collected ITC data was then imported into Nano analyze software (TA Instruments) and a baseline linear curve representing to energy from injecting water into water was subtracted leaving a normalized energy injection dataset. Injection points from either the first injection or injections after the binding affinity had plateaued were excluded and the remaining data points were then fit to a single independent binding curve. FL-Ezrin or FERM injections that did not show an absorbance curve were not fit to one in the Nano analyze program as part of a pre exclusion criteria.

Force indentation:

Cells were plated at 50% confluence to form uniform monolayers. Four hours prior to indentation media was exchanged with fresh media containing 10uM Hepes buffer. Cells were then mounted on an inverted epifluorescence microscope (Observer Z1, Zeiss) and force indentation curves were collected using nanoindentation device (Chiaro, Optics11, Netherlands). Measurements were performed at room temperature with a 0.030 N/m cantilever with a spherical tip of 3μm radius. Young's elastic modulus was

calculated using the Optics11 DataViewer software using by applying a Hertzian model (equation below). Any indentation with a fit for the Hertzian model below an R2 value of 0.8 was excluded from further analysis. Visualization of each indentation fit was performed by graphing the force as a function of displacement using the equation $F = \frac{4}{3} * E * \sqrt{PR} * h^{\frac{3}{2}}$, where F is the force in Newton's, E is the measured Young's modulus, PR is the probe radius, and h is displacement in meters starting at the point of contact. Graphing and analysis of Young's modulus was performed using GraphPad Prism.

RhoA GAP assay:

RhoA GAP assays were performed using a kit provided by Cytoskeleton Inc (Denver Colorado Cat no. BK105) and by following the provided kit protocol summarized below. FERM ezrin and ARHGAP18 were purified as described above and were dialyzed into fresh GAP assay buffer provided in the kit. RhoA and p50 RhoGAP proteins were provided in the kit. For each reaction, +/- 4µg of RhoA, 4ug FERM, 4ug ARHGAP18 or, 6ug p50RhoA GAP, depending on the experimental condition, was brought to a final concentration of 30ul in GAP buffer. All reactions for the various conditions were prepared and run simultaneously in a single 96 well dish for each replicate. The reactions were started simultaneously by adding 200uM GTP to the wells using a multichannel pipette. The reaction was incubated at 37°C for 20 minutes before adding CytoPhos reagent which increased absorbance at 550nm as free phosphate is released. Absorbance signal was measured at 550nm using a Molecular Dynamics SpectraMax i3X and an average normalized signal was calculated then plotted in GraphPad Prism.

Data and materials availability statement:

Data, newly synthesized materials, coding programs, and analysis from the current study are available from the corresponding author upon reasonable request.

Acknowledgments

We thank A. Martin, N. Buffard for their helpful discussions and J. Lammerding, N. Zuela-Sopilniak and A. Varlet for expertise with force indentation. This work was supported by National Institutes of Health grant R35GM131751 to A. Bretscher and the Sam and Nancy Fleming Research Fellowship to A. Lombardo. GFP tagged ARHGAP18 used as a control in antibody quality control was a generous gift from J. Gamble. Scanning electron microscopy was performed at the Microscopy Imaging Center at the University of Vermont (RRID# SCR_018821), and the Cornell Institute of Biotechnology Imaging Facility is supported by National Science Foundation funding (1428922).

Citations

Bagci, H., Sriskandarajah, N., Robert, A., Boulais, J., Elkholi, I. E., Tran, V., Lin, Z. Y., Thibault, M. P., Dubé, N., Faubert, D., Hipfner, D. R., Gingras, A. C., & Côté, J. F. (2020). Mapping the proximity interaction network of the Rho-family GTPases reveals signalling pathways and regulatory mechanisms. *Nature Cell Biology*, *22*(1), 120–134. https://doi.org/10.1038/s41556-019-0438-7

Belkina, N. v., Liu, Y., Hao, J. J., Karasuyama, H., & Shaw, S. (2009). LOK is a major ERM kinase in resting lymphocytes and regulates cytoskeletal rearrangement through ERM phosphorylation. *Proceedings of the National Academy of Sciences of the United States of America*, *106*(12), 4707–4712. https://doi.org/10.1073/pnas.0805963106

Bretscher, A. (1983). Purification of an 80,000-dalton protein that is a component of the isolated microvillus cytoskeleton, and its localization in nonmuscle cells. *The Journal of Cell Biology*, *97*(2), 425–432. https://doi.org/10.1083/jcb.97.2.425

Casaletto, J. B., Saotome, I., Curto, M., & McClatchey, A. I. (2011). Ezrin-mediated apical integrity is required for intestinal homeostasis. *Proceedings of the National Academy of Sciences of the United States of America*, *108*(29), 11924–11929. https://doi.org/10.1073/pnas.1103418108

Chinowsky, C. R., Pinette, J. A., Meenderink, L. M., Lau, K. S., & Tyska, M. J. (2020). Nonmuscle myosin-2 contractility-dependent actin turnover limits the length of epithelial microvilli. *Molecular Biology of the Cell*, *31*(25), 2803–2815. https://doi.org/10.1091/mbc.E20-09-0582

Denk-Lobnig, M., & Martin, A. C. (2019). Modular regulation of Rho family GTPases in development. In *Small GTPases* (Vol. 10, Issue 2, pp. 122–129). Taylor and Francis Inc. https://doi.org/10.1080/21541248.2017.1294234

Doi, Y., Itoh, M., Yonemura, S., Ishihara, S., Takano, H., Noda, T., Tsukita, S., & Tsukita, S. (1999). Normal development of mice and unimpaired cell adhesion/cell motility/actin-based cytoskeleton without compensatory up-regulation of ezrin or radixin in moesin gene knockout. *Journal of Biological Chemistry*, *274*(4), 2315–2321. https://doi.org/10.1074/jbc.274.4.2315

Fehon, R. G., McClatchey, A. I., & Bretscher, A. (2010). Organizing the cell cortex: The role of ERM proteins. In *Nature Reviews Molecular Cell Biology* (Vol. 11, Issue 4, pp. 276–287). Nature Publishing Group. https://doi.org/10.1038/nrm2866

Fievet, B. T., Gautreau, A., Roy, C., del Maestro, L., Mangeat, P., Louvard, D., & Arpin, M. (2004). Phosphoinositide binding and phosphorylation act sequentially in the activation mechanism of ezrin. *Journal of Cell Biology*, *164*(5), 653–659. https://doi.org/10.1083/jcb.200307032

Garbett, D., & Bretscher, A. (2012). PDZ interactions regulate rapid turnover of the scaffolding protein EBP50 in microvilli. *Journal of Cell Biology*, *198*(2), 195–203. https://doi.org/10.1083/jcb.201204008

Gary, R., & Bretscher, A. (1995). Ezrin self-association involves binding of an N-terminal domain to a normally masked C-terminal domain that includes the F-actin binding site. *Molecular Biology of the Cell*, *6*(8), 1061–1075. https://doi.org/10.1091/mbc.6.8.1061

Heaslip, A. T., Nelson, S. R., Lombardo, A. T., Beck Previs, S., Armstrong, J., & Warshaw, D. M. (2014). Cytoskeletal Dependence of Insulin Granule Movement Dynamics in INS-1 Beta-Cells in Response to Glucose. *PLoS ONE*, *9*(10), e109082. https://doi.org/10.1371/journal.pone.0109082

Heissler, S. M., & Manstein, D. J. (2013). Nonmuscle myosin-2: Mix and match. In *Cellular and Molecular Life Sciences* (Vol. 70, Issue 1, pp. 1–21). Springer. https://doi.org/10.1007/s00018-012-1002-9

Hirao, M., Sato, N., Kondo, T., Yonemura, S., Monden, M., Sasaki, T., Takai, Y., Tsukita, S., & Tsukita, S. (1996). Regulation mechanism of ERM (ezrin/radixin/moesin) protein/plasma membrane association: Possible

involvement of phosphatidylinositol turnover and rho-dependent signaling pathway. *Journal of Cell Biology*, *135*(1), 37–51. https://doi.org/10.1083/jcb.135.1.37

Hoskin, V., Ghaffari, A., & Elliott, B. E. (2019). Ezrin, more than a metastatic determinant? In *Oncotarget* (Vol. 10, Issue 63, pp. 6755–6757). Impact Journals LLC. https://doi.org/10.18632/oncotarget.27227

MacHacek, M., Hodgson, L., Welch, C., Elliott, H., Pertz, O., Nalbant, P., Abell, A., Johnson, G. L., Hahn, K. M., & Danuser, G. (2009). Coordination of Rho GTPase activities during cell protrusion. *Nature*, *461*(7260), 99–103. https://doi.org/10.1038/nature08242

Maeda, M., Hasegawa, H., Hyodo, T., Ito, S., Asano, E., Yuang, H., Funasaka, K., Shimokata, K., Hasegawa, Y., Hamaguchi, M., & Senga, T. (2011). ARHGAP18, a GTPase-activating protein for RhoA, controls cell shape, spreading, and motility. *Molecular Biology of the Cell*, *22*(20), 3840–3852. https://doi.org/10.1091/mbc.E11-04-0364

Martin, T. A., Harrison, G., Mansel, R. E., & Jiang, W. G. (2003). The role of the CD44/ezrin complex in cancer metastasis. In *Critical Reviews in Oncology/Hematology* (Vol. 46, Issue 2, pp. 165–186). Elsevier Ireland Ltd. https://doi.org/10.1016/S1040-8428(02)00172-5

Matsui, T., Maeda, M., Doi, Y., Yonemura, S., Amano, M., Kaibuchi, K., Tsukita, S., & Tsukita, S. (1998). Rho-kinase phosphorylates COOH-terminal threonines of ezrin/radixin/moesin (ERM) proteins and regulates their head-to-tail association. *Journal of Cell Biology*, *140*(3), 647–657. https://doi.org/10.1083/jcb.140.3.647

McConnell, R. E., Benesh, A. E., Mao, S., Tabb, D. L., & Tyska, M. J. (2011). Proteomic analysis of the enterocyte brush border. *American Journal of Physiology - Gastrointestinal and Liver Physiology*, *300*(5), G914–G926. https://doi.org/10.1152/ajpgi.00005.2011

Meenderink, L. M., Gaeta, I. M., Postema, M. M., Cencer, C. S., Chinowsky, C. R., Krystofiak, E. S., Millis, B. A., & Tyska, M. J. (2019). Actin Dynamics Drive Microvillar Motility and Clustering during Brush Border Assembly. *Developmental Cell*, *50*(5), 545-556.e4. https://doi.org/10.1016/j.devcel.2019.07.008

Nakamura, F., Amieva, M. R., & Furthmayr, H. (1995). Phosphorylation of threonine 558 in the carboxyl-terminal actin-binding domain of moesin by thrombin activation of human platelets. *Journal of Biological Chemistry*, *270*(52), 31377–31385. https://doi.org/10.1074/jbc.270.52.31377

Neisch, A. L., Formstecher, E., & Fehon, R. G. (2013). Conundrum, an ARHGAP18 orthologue, regulates RhoA and proliferation through interactions with Moesin. *Molecular Biology of the Cell*, *24*(9), 1420–1433. https://doi.org/10.1091/mbc.E12-11-0800

Pelaseyed, T., Sauvanet, C., Viswanatha, R., Filter, J. J., Goldberg, M. L., & Bretscher, A. (2017). Ezrin activation by LOK phosphorylation involves a PIP2-dependent wedge mechanism. *ELife*, *6*. https://doi.org/10.7554/eLife.22759

Piekny, A. J., & Glotzer, M. (2008). Anillin Is a Scaffold Protein That Links RhoA, Actin, and Myosin during Cytokinesis. *Current Biology*, *18*(1), 30–36. https://doi.org/10.1016/j.cub.2007.11.068

Porazinski, S., Wang, H., Asaoka, Y., Behrndt, M., Miyamoto, T., Morita, H., Hata, S., Sasaki, T., Krens, S. F. G., Osada, Y., Asaka, S., Momoi, A., Linton, S., Miesfeld, J. B., Link, B. A., Senga, T., Castillo-Morales, A., Urrutia,

A. O., Shimizu, N., … Furutani-Seiki, M. (2015). YAP is essential for tissue tension to ensure vertebrate 3D body shape. *Nature*, *521*(7551), 217–221. https://doi.org/10.1038/nature14215

Priya, R., Gomez, G. A., Budnar, S., Verma, S., Cox, H. L., Hamilton, N. A., & Yap, A. S. (2015). Feedback regulation through myosin II confers robustness on RhoA signalling at E-cadherin junctions. *Nature Cell Biology*, *17*(10), 1282–1293. https://doi.org/10.1038/ncb3239

Rogg, M., Maier, J. I., Dotzauer, R., Artelt, N., Kretz, O., Helmstädter, M., Abed, A., Sammarco, A., Sigle, A., Sellung, D., Dinse, P., Reiche, K., Yasuda-Yamahara, M., Biniossek, M. L., Walz, G., Werner, M., Endlich, N., Schilling, O., Huber, T. B., & Schell, C. (2021). SRGAP1 controls small rho GTPases to regulate podocyte foot process maintenance. *Journal of the American Society of Nephrology*, *32*(3), 563–579. https://doi.org/10.1681/ASN.2020081126

Sanjana, N. E., Shalem, O., & Zhang, F. (2014). Improved vectors and genome-wide libraries for CRISPR screening. In *Nature Methods* (Vol. 11, Issue 8, pp. 783–784). Nature Publishing Group. https://doi.org/10.1038/nmeth.3047

Sauvanet, C., Wayt, J., Pelaseyed, T., & Bretscher, A. (2015). Structure, Regulation, and Functional Diversity of Microvilli on the Apical Domain of Epithelial Cells. *Annual Review of Cell and Developmental Biology*, *31*(1), 593–621. https://doi.org/10.1146/annurev-cellbio-100814-125234

Sellers, J. R., & Heissler, S. M. (2019). Nonmuscle myosin-2 isoforms. In *Current Biology* (Vol. 29, Issue 8, pp. R275–R278). Cell Press. https://doi.org/10.1016/j.cub.2019.03.022

Shaw, R. J., Henry, M., Solomon, F., & Jacks, T. (1998). RhoA-dependent phosphorylation and relocalization of ERM proteins into apical membrane/actin protrusions in fibroblasts. *Molecular Biology of the Cell*, *9*(2), 403–419. https://doi.org/10.1091/mbc.9.2.403

Speck, O., Hughes, S. C., Noren, N. K., Kulikauskas, R. M., & Fehon, R. G. (2003). Moesin functions antagonistically to the Rho pathway to maintain epithelial integrity. *Nature*, *421*(6918), 83–87. https://doi.org/10.1038/nature01295

Stewart, K., Gaitan, Y., Shafer, M. E. R., Aoudjit, L., Hu, D., Sharma, R., Tremblay, M., Ishii, H., Marcotte, M., Stanga, D., Tang, Y. C., Boualia, S. K., Nguyen, A. H. T., Takano, T., Lamarche-Vane, N., Vidal, S., & Bouchard, M. (2016). A Point Mutation in p190A RhoGAP Affects Ciliogenesis and Leads to Glomerulocystic Kidney Defects. *PLOS Genetics*, *12*(2), e1005785. https://doi.org/10.1371/journal.pgen.1005785

Tsukita, S., Oishi, K., Sato, N., Sagara, J., Kawai, A., & Tsukita, S. (1994). ERM family members as molecular linkers between the cell surface glycoprotein CD44 and actin-based cytoskeletons. *Journal of Cell Biology*, *126*(2), 391–401. https://doi.org/10.1083/jcb.126.2.391

Turunen, O., Wahlström, T., & Vaheri, A. (1994). Ezrin has a COOH-terminal actin-binding site that is conserved in the ezrin protein family. *Journal of Cell Biology*, *126*(6), 1445–1453. https://doi.org/10.1083/jcb.126.6.1445

Viswanatha, R., Ohouo, P. Y., Smolka, M. B., & Bretscher, A. (2012). Local phosphocycling mediated by LOK/SLK restricts ezrin function to the apical aspect of epithelial cells. *Journal of Cell Biology*, *199*(6), 969–984. https://doi.org/10.1083/jcb.201207047

Viswanatha, R., Wayt, J., Ohouo, P. Y., Smolka, M. B., & Bretscher, A. (2013). Interactome analysis reveals ezrin can adopt multiple conformational states. *Journal of Biological Chemistry*, *288*(49), 35500–35510. https://doi.org/10.1074/jbc.M113.505669

Zaman, R., Lombardo, A., Sauvanet, C., Viswanatha, R., Awad, V., Bonomo, L. E. R., McDermitt, D., & Bretscher, A. (2021). Effector-mediated ERM activation locally inhibits rhoa activity to shape the apical cell domain. *Journal of Cell Biology*, *220*(6). https://doi.org/10.1083/jcb.202007146