ATF4 Dependent Increase in Mitochondrial-Endoplasmic Reticulum Tethering
Following OPA1 Deletion in Skeletal Muscle

Antentor Hinton, Jr.,[#1,2,3], Prasanna Katti[#4], Margaret Mungai[1,2], Duane D. Hall[1], Olha Koval[1,2],
Jianqiang Shao[5], Zer Vue[3], Edgar Garza Lopez[1], Rahmati Rostami[6], Kit Neikirk[3], Jessica
Ponce[1,2], Jennifer Streeter[1,2], Brandon Schickling[1,7], Serif Bacevac[1,2], Chad Grueter[1,2], Andrea
Marshall[3], Heather K. Beasley[3], Young Do Koo[1,2], Sue C. Bodine[1,2,8], Nayeli G. Reyes Nava[9],
Anita M. Quintana[9], Long-Sheng Song[1,2], Isabella Grumbach[1,2], Renata O. Pereira[1,2*], Brian
Glancy[4,10*], and E. Dale Abel[1,2,11*]

Affiliations:
1. Department of Internal Medicine, University of Iowa - Carver College of Medicine, Iowa City,
IA, USA 52242
2. Fraternal Order of Eagles Diabetes Research Center, Iowa City, IA, USA 52242
3. Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville, TN
37232
4. National Heart, Lung, and Blood Institute, National Institutes of Health, Bethesda, MD, USA,
20892
5. Central Microscopy Research Facility, Iowa City, IA USA 52242
6. Department of Genetic Medicine, Joan & Sanford I. Weill Medical College of Cornell
University, New York, NY, USA, 10065
7. Department of Medicine, Duke University, Durham, NC, USA 27708
8. Oklahoma Medical Research Foundation, Oklahoma City, OK, USA 73104
9. Department of Biological Sciences and Border Biomedical Research Center, The University of
Texas at El Paso, El Paso, TX, USA 79968
10. National Institute of Arthritis and Musculoskeletal and Skin Diseases, National Institutes of
Health, Bethesda, MD, USA 20892
11. Department of Medicine, UCLA School of Medicine, Los Angeles, CA, USA 90095
27
28 # Authors Contributed Equally to Manuscript, co-first
29
30 * co-Senior Authors
31
32 *Co-corresponding authors:
33
E. Dale Abel, M.D., Ph.D.

Department of Medicine

David Geffen School of Medicine and UCLA Health

757 Westwood Plaza, Suite 7236

Los Angeles, CA 90095

Phone: 310-267-3144 | Fax: 310-206-7291

Email: DOMChair_DaleAbel@mednet.ucla.edu

Brian Glancy, Ph.D.
National Heart, Lung, and Blood Institute
National Institute of Arthritis and Musculoskeletal and Skin Diseases
National Institutes of Health
10 Center Dr, Room B1D416
Bethesda MD 20892
Phone: 301-496-2679
Brian.glancy@nih.gov

50
51

Summary:

Mitochondria and endoplasmic reticulum (ER) contact sites (MERCs) are protein- and lipid-

enriched hubs that mediate interorganellar communication by contributing to the dynamic

transfer of Ca^{2+}, lipid, and other metabolites between these organelles. Defective MERCs are

associated with cellular oxidative stress, neurodegenerative disease, and cardiac and skeletal

muscle pathology via mechanisms that are poorly understood. We previously demonstrated that

skeletal muscle-specific knockdown (KD) of the mitochondrial fusion mediator optic atrophy 1

(OPA1) induced ER stress and correlated with an induction of Mitofusin-2, a known MERC

protein. In the present study, we tested the hypothesis that *Opa1* downregulation in skeletal

muscle cells alters MERC formation by evaluating multiple myocyte systems, including from

mice and *Drosophila*, and in primary myotubes. Our results revealed that OPA1 deficiency

induced tighter and more frequent MERCs in concert with a greater abundance of MERC

proteins involved in calcium exchange. Additionally, loss of OPA1 increased the expression of

activating transcription factor 4 (ATF4), an integrated stress response (ISR) pathway effector.

Reducing *Atf4* expression prevented the OPA1-loss-induced tightening of MERC structures.

OPA1 reduction was associated with decreased mitochondrial and sarcoplasmic reticulum, a

specialized form of ER, calcium, which was reversed following ATF4 repression. These data

suggest that mitochondrial stress, induced by OPA1 deficiency, regulates skeletal muscle MERC

formation in an ATF4-dependent manner.

Keywords: Optic atrophy 1, Mitochondria Endoplasmic Reticulum Contact, Activating

Transcription Factor 4, Integrated Stress Response, Serial Block Face-Scanning Electron

Microscopy, Transmission Electron Microscopy, Focus Ion Beam-Scanning Electron

Microscopy,

<u>Introduction</u>

The endoplasmic reticulum (ER) and mitochondria continuously exchange molecules to

sustain intracellular homeostasis. This is facilitated, in part, by physical contacts between the two

organelles known as mitochondria-associated membranes (MAMs) or mitochondrial-ER contact

sites (MERCs) (Rieusset, 2018). MERC dysfunction, including alterations in MERC formation

and morphology, has been associated with disease states characterized by altered mitochondrial

oxidative metabolism and insulin resistance including obesity, type II diabetes (T2D),

cardiovascular disease and aging (Rieusset, 2018).

MERC microdomains are enriched in proteins involved in mitochondrial Ca^{2+} flux, lipid

transfer, and the regulation of mitochondrial function (Tubbs et al., 2014, 2018). The size,

density, and thickness of MERCs regulate their function, and changes in MERC distances can

modulate Ca^{2+} exchange, lipid homeostasis, and cell death (Csordás et al., 2018; Delprat et al.,

2019; Giacomello & Pellegrini, 2016; Hirabayashi et al., 2017; Rieusset, 2018).

Several important MERC-tethering proteins are expressed in the MERC microdomain,

including Mitofusin-2 (MFN2), the inositol 1,4,5-triphosphate receptor (IP$_3$R3), glucose-related

protein 75 (GRP75), voltage-dependent anion channel (VDAC) complex, the vesicle-associated

membrane proteins B and C, and others. Most known MERC proteins are located on the outer

mitochondrial membrane (OMM). A relationship between the inner mitochondria membrane

protein optic atrophy 1 (OPA1) and regulation of MERCs, was recently described (Cartes-

Saavedra et al., 2022). However, a complete understanding of mechanisms linking inner

mitochondrial membrane proteins, such as OPA1, to the regulation of MERC formation remains

to be achieved.

We previously showed that *OPA1* knockdown (KD) in skeletal muscle (OPA1smko/KD)

impairs mitochondrial fusion and bioenergetics, induces ER stress, and correlates with the

induction of MFN2 (Pereira et al., 2017; Tezze et al., 2017). *OPA1* KD in skeletal muscle also

increases activating transcription factor (ATF4) levels (Pereira et al., 2017). Interestingly, ATF4

activation is involved in several stress processes, including having a key role in the integrated

stress response (ISR) (Costa-Mattioli & Walter, 2020; Hartwick Bjorkman & Oliveira Pereira,

2021; Kasai et al., 2019; Sasaki et al., 2020). ATF4 is also linked to muscle atrophy (Pereira et

al., 2017). Skeletal muscle atrophy and reduced exercise activity are associated with MERC

changes (Romanello & Sandri, 2015). Age-related physical inactivity is also associated with

muscle loss and reduced OPA1 levels (Tezze et al., 2017). However, the mechanisms by which

ATF4 may influence the morphology of MERC microdomains in these and other contexts are

incompletely understood.

We hypothesized that OPA1-induced mitochondrial stress increases MERC abundance and

narrows MERC distance, potentially as a response to ER stress. To test this hypothesis, we used

OPA1 loss of function in myotubes, mouse skeletal muscle, and *Drosophila* flight muscle to

assess the effects of OPA1 deficiency on MERC structure. Across models, we demonstrate that

OPA1 deficiency in muscle cells induces the expression of MERC proteins and increases MERC

tethering. Additionally, we find that the MERC structural changes in OPA1-deficient muscle

cells are mediated, at least in part, by ATF4. Indeed, *Atf4* loss reverses the narrowing of MERC

distances driven by *Opa1* loss, while *Atf4* overexpression alone is sufficient to increase MERC

length and narrow MERC distances. This increase in MERC formation may represent an initially

compensatory, but ultimately maladaptive mechanism that attempts to restore calcium

homeostasis in response to OPA1-deficiency-induced mitochondrial stress.

Materials and Methods

Animal Studies

The University of Iowa Institutional Animal Care and Use Committee (IACUC) approved all

protocols for animal work. For primary satellite cells isolation, 16 male mice on the C57BL/6J

genetic background were used. We previously reported that male and female mice demonstrate

similar effects to loss of *OPA1* (Pereira et al., 2017). For all gastrocnemius samples, mice were

aged to 9 months.

Generation of OPA1 KO animals and TUDCA treatment was achieved using protocols

previously described by us (Pereira et al., 2017). The genotypes of OPA1 deficient and control

mice are as described for satellite cell experiments. Standard chow was fed to mice during their

4-week recovery period following tamoxifen injections at 4 weeks of age. A subgroup of OPA1

deficient and control mice were treated with TUDCA (Millipore: Billerica, MA, USA;

250 mg/kg), administered via intraperitoneal injection. vehicle treated mice received equivalent

volumes of phosphate-buffered saline. All animals had free access to standard chow and water in

a 22 °C environment with a 12h/12h light/dark cycle.

Atf4$^{fl/fl}$ mice were generated as previously described (Ebert et al., 2012). We also generated *Opa1*

and *Atf4* double floxed mice from which satellite cells were also isolated.

Primary Cell Culture

Satellite cell isolation was performed as previously described (Pereira et al., 2017). Eight mice of

each experimental condition [control (*OPA1*$^{flox/flox}$) and skeletal muscle OPA1-deficient mice

(*OPA1$^{flox/flox}$ has-CreER^{T2}*)] were used to isolate primary satellite cells. Satellite cells from

OPA1$^{fl/fl}$, *Atf4$^{fl/fl}$*, and *OPA1$^{fl/fl}$ + Atf4*$^{fl/fl}$ double floxed mice were plated on BD Matrigel-coated

dishes and differentiated into myoblasts in Dulbecco's modified Eagle medium (DMEM)-F12

containing 20% fetal bovine serum (FBS), 40 ng/ml basic fibroblast growth factor, 1× non-

essential amino acids, 0.14 mM β-mercaptoethanol, 1× penicillin/streptomycin, and Fungizone.

Myoblasts were maintained with 10 ng/ml basic fibroblast growth factor and differentiated in

DMEM-F12 containing 2% FBS and 1× insulin–transferrin–selenium when 90% confluency was

reached. Three days after differentiation, myotubes were infected with an adenovirus expressing

GFP-Cre to achieve OPA1 deletion, ATF4 deletion, or double knockout of OPA1 and ATF4.

Additionally, we obtained an ATF4 adenovirus (Ad5CMVATF4/RSVeGFP) from the University

of Iowa Viral Vector Core facility to induce ATF4 overexpression in Atf4$^{fl/fl}$ myotubes and in

OPA1 ablated myotubes. . Experiments were performed between 3 and 7 days after infection.

Human Cell Culture

Human and primary mouse myotubes were grown and maintained in DMEM-F12 (Gibco:

Waltham, MA, USA) supplemented with 20% fetal bovine serum (Gibco), 10 ng/ml basic

fibroblast growth factor, 1% penicillin/streptomycin, 300 µl/100 ml Fungizone, 1% non-essential

amino acids, and 1 mM β-mercaptoethanol. Before plating, dishes were coated with Matrigel to

enhance cell adherence. On alternate days, the cells were washed with phosphate-buffered saline,

and the media replaced.

RNA Sample Preparation and Quantitative RT-PCR in Mouse Samples

RNA was isolated from the gastrocnemius of 40-week-old mice using TRIzol and tissue grinder

(Sigma-Aldrich: St. Louis, MO, USA). Reverse polymerase chain reaction (PCR) was performed

to generate cDNA using SuperScript III (Invitrogen). For quantitative reverse transcriptase-PCR,

50 ng of cDNA was used for each reaction with iTAQ Universal SYBR green (Biorad: Hercules,

CA, USA), using the QuantiStudio 6 Flex system (Applied Biosystems: Waltham, MA, USA).

Gene expression was analyzed using the $\Delta\Delta C_T$ method, and relative expression was normalized

to *Gapdh*. Sequences for primers are listed in supplementary materials (See Table 1).

RNA Extraction and Quantitative RT-PCR for Cells

RNA extraction and quantification were performed as previously described (Pereira et al., 2017).

Briefly, RNA was extracted from samples using TRIzol reagent and purified with the RNeasy kit

(Qiagen Inc: Hilden, Germany). The RNA was reverse transcribed, followed by quantitative

PCR (qPCR) reactions using SYBR Green. Samples were then loaded onto a 384-well plate, and

a qPCR was performed using an ABI Prism 7900HT instrument. The primers used are listed in

Supplemental Table 1.

Western Blotting

To obtain total protein extracts from differentiated myotubes, cells were washed with ice-cold

PBS before adding cold lysis buffer (25 mM Tris HCl, pH 7.9, 5 mM MgCl$_2$, 10% glycerol,

100 mM KCl, 1% NP40, 0.3 mM dithiothreitol, 5 mM sodium pyrophosphate, 1 mM

sodium orthovanadate, 50 mM sodium fluoride, and protease inhibitor cocktail (Roche Applied

Science: Penzberg, Germany). Cells were scraped, homogenized with a 25-gauge needle, and

centrifuged at 14,000 rpm for 10 min at 4 °C. Supernatants were collected and mixed with

Laemmli sample buffer to get a final concentration of 1X. Cell lysates were subjected to sodium

dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), and proteins were transferred

to nitrocellulose membranes (BioRad: Berkeley, California, USA). Membranes were blocked

with 5% bovine serum albumin (BSA)-Tris-buffered saline with Tween-20 (TBST) and

incubated with antibodies as indicated. Quantification was performed using Image Studio Lite

Ver 5.2 with three biological replicates for each protein of interest.

Primary antibodies used for Western blotting and their working dilutions include: OPA1

(1:1,000; BD Biosciences: San Jose, CA, USA; #612606), glyceraldehyde 3-phosphate

dehydrogenase (GAPDH; 1:5,000; Cell Signaling Technology: Danvers, MA, USA; #2118),

MFN1 (1:1,000; Abcam, Cambridge, UK; #ab57602), MFN2 (1:1,000; Abcam; #ab101055),

GRP75 (1:1000; Cell Signaling Technology; #D13H4), PACS2 (1:1,000; Abcam; #ab129402),

BIP (1:5,000; BD Biosciences; #610978), ATF4 (1:1000; Proteintech: Rosemont, IL, USA;

10835-1-AP), VDAC (1:1000; Cell Signaling Technology; #4866), SERCA (1:1000; Cell

Signaling Technology; #4388), MCU (1:1000; ThermoFisher Scientific: Waltham, MA, USA;

#PA5-68153), Alpha Tubulin (1:1000; Cell Signaling Technology; #2144), and IP$_3$R3 (1:1000;

BD Biological Sciences; #610312). Secondary antibodies include: IRDye 800CW anti-mouse

(1:10,000; LI-COR: Lincoln, NE, USA; #925- 32212) and Alexa Fluor anti-rabbit 680 (1:10,000;

Invitrogen; #A27042). Fluorescence was quantified using the LiCor Odyssey imager.

MERC Confocal Analysis

Cells were grown in culture media on a #1.5 cover glass for optimized microscope optics and

either embedded into a petri dish or divided with plastic-walled growth chambers. Cells were

incubated for 30 minutes with MitoTracker Orange FM (Invitrogen, 400 nmol/L) and

subsequently fixed in 4% paraformaldehyde to prepare for colocalization staining with GRP78.

Confocal image stacks were captured with a Zeiss LSM-5, Pascal 5 Axiovert 200 microscope

using LSM 5 version 3.2 image capture and analysis software and a Plan-APOCHROMAT

40x/1.4 Oil DIC objective. Images were deconvoluted with NIH (National Institutes of Health)

ImageJ software and BITPLANE Imaris software (Oxford Instruments: Abingdon, United

Kingdom). Subsequently, images were subjected to analysis using Imaris software for Mander's

and Pearson's coefficient analyses. Each experiment was performed at least three times, and

quantified from 10–20 cells per condition.

Sample Preparation for Transmission Electron Microscope (TEM) Analysis

Mice were anesthetized with isoflurane according to prior protocols (Garza-Lopez et al., 2022;

Lam et al., 2021; Neikirk, Vue, et al., 2023; Vue et al., 2022). Soleus muscle was freshly excised

and immediately washed in ice-cold saline. Samples were processed as previously described by

us (Pereira et al., 2017). Soleus samples were fixed with Trump's TEM fixative as previously

described(Hinton et al., 2023) and stained. Random images of intermyofibrillar (IMF) sections

were obtained. TEM was used to view the ultrastructure of the samples to quantify MERCs.

Cells were fixed in 2.5% glutaraldehyde in sodium cacodylate buffer for 1 hour at 37 °C,

embedded in 2% agarose, postfixed in buffered 1% osmium tetroxide, stained in 2% uranyl

acetate, and dehydrated with an ethanol-graded series. After embedding in EMbed-812 resin, 80

nm sections were cut on an ultramicrotome and stained with 2% uranyl acetate and lead citrate.

Images were acquired on a JEOL JEM-1230 Transmission Electron Microscope, operating at

120 kV.

TEM Analysis

Measurements of mitochondrial area, circularity, and length were performed using the Multi-

Measure region of interest (ROI) tool in ImageJ (Garza-Lopez et al., 2022; Lam et al., 2021;

Neikirk, Vue, et al., 2023; Parra et al., 2013; Vue et al., 2022), in accordance with prior

recommendations (Neikirk, Lopez, et al., 2023). To measure cristae morphology, we used three

distinct ROIs of all the same magnification in ImageJ to determine the cristae area, circulatory

index, number, and cristae score. Cristae volume was determined by the sum of all cristae area

divided by total area of mitochondria (Patra et al., 2016). The NIH ImageJ software was used to

manually trace and analyze all mitochondria or cristae using the freehand told in the ImageJ

application (Parra et al., 2013). We also measured MERC distance, length, and percent coverage.

MERC length was defined as the horizontal distance between the mitochondria and ER interface.

MERC distance was defined as the vertical distance between the mitochondria and ER interface.

MERC percentage coverage was calculated as the total ER or mitochondrion surface area

normalized to the distance between the mitochondria and the ER.

Serial Block Face-Scanning Electron Microscopy (SBF-SEM) Processing for *Drosophila*

Muscle Fibers

Protocols follow previously established methods (Crabtree et al., 2023; Garza-Lopez et al., 2022;

Vue et al., 2022, 2023). Briefly, tissue was fixed in 2% glutaraldehyde in 0.1 M cacodylate

buffer and processed using a heavy metal protocol adapted from a previously published protocol

(Courson et al., 2021; Mustafi et al., 2014). Tissue was immersed in 3% potassium ferrocyanide

and 2% osmium tetroxide (1 hour at 4 °C), followed by filtered 0.1% thiocarbohydrazide (20

min), 2% osmium tetroxide (30 min), and left overnight in 1% uranyl acetate at 4 °C (several de-

ionized H_2O washes were performed between each step). The next day, samples were immersed

in a 0.6% lead aspartate solution (30 min at 60 °C and dehydrated in graded acetone (as

described for TEM). The tissues were impregnated with epoxy TAAB 812 hard resin, embedded

in fresh resin, and polymerized at 60 °C for 36–48 hours. After polymerization, the block was

sectioned for TEM to identify the area of interest, then trimmed to 0.5 mm × 0.5 mm and glued

to an aluminum pin. The pin was placed into an FEI/Thermo Scientific Volumescope 2 scanning

electron microscope (SEM), a state-of-the-art SBF imaging system. For 3D EM reconstruction,

300 to 400 thin (0.09 µm) serial sections were obtained from single blocks that were processed

for conventional TEM. Serial sections were collected onto formvar coated slot grids (Pella,

Redding CA), stained, and imaged, as described above. The segmentation of SBF-SEM

reconstructions was performed by manually tracing structural features through sequential slices

of the micrograph block. Images were collected from 30–50 serial sections, which were then

stacked, aligned, and assembled into videos using Amira (Thermo Fisher Scientific, Waltham,

Massachusetts, USA) to quantify and visualize volumetric structures (Garza-Lopez et al., 2022).

Proximity Ligation Assay in Cells

We used Duolink In Situ Detection Reagents Red (Sigma-Aldrich; DUO92008) to verify the

proximity of two antibodies of interest in cell culture. Previously validated antibodies were used.

The first antibody was mouse monoclonal antibody against MFN1 (Abcam #ab57602), the

second antibody was a rabbit polyclonal antibody against MFN2 (Abcam #ab50843), produced

using a synthetic peptide corresponding to human Mitofusin 2, aa 557–576, conjugated to

keyhole limpet hemocyanin (KLH). Both antibodies were determined to work at a 1:200 dilution.

Three controls were included in these experiments. Each primary antibody was omitted from

separate cells to determine the non-specific binding of the primary antibodies and to confirm the

optimal antibody titer. One well was devoid of all primary antibodies to detect any non-specific

binding of the Duolink[©] PLA probes.

Cells were fixed in 4% paraformaldehyde in PBS at 37 °C for 10 minutes. After three rinses with

PBS, a permeabilization solution containing 0.02% Triton X-100 in PBS was applied for 10

minutes. A ready-to-use blocking solution from the kit was applied for 1 hour at 37 °C. A

cocktail containing both antibodies, or control solution, diluted in the provided diluent, was

applied to the cells overnight at 4 °C. The next day, cells were washed and incubated for 1 hour

at 37 °C with a pair of oligonucleotide-labeled secondary antibodies, one against mouse and one

against rabbit. Cells were washed again and hybridized with the ligase solution for 30 minutes at

37 °C, connecting the oligos in close proximity to each other and forming a closed circle DNA

template. Cells were then washed, rolling circle amplification (RCA) was performed with the

template, with the probe acting as a primer for the DNA polymerase. RCA was allowed to

proceed in a dark hybridization chamber at 37 °C for 1 hour and 40 minutes. Then, cells were

washed and mounted in Duolink mounting media and stored at 4 °C overnight and imaged with

an SP-8 confocal inverted microscope using a white laser light set to an excitation wavelength of

594 nm and an emission wavelength of 624 nm ± 10 nm.

Proximity Ligation Assay in Tissue

Duolink In Situ Detection Reagents Red (Sigma-Aldrich; DUO92008) was used to verify the

proximity of two antibody locations in tissue samples. We used the two antibodies described for

cells. Both antibodies were determined to work at a 1:50 dilution. One tissue slide was devoid of

both primary antibodies to detect any non-specific binding of the Duolink PLA probes. The

tissues were fixed in a manner that described forlike PLA in cells above except , tissue fixation

occurred in warm 4% paraformaldehyde in PBS at 37 °C for 24 hoursinstead of . After fixation,

tissues were mounted in optimal cutting temperature (OCT) medium.

Fly Strains and Genetics

Genetic crosses in *Drosophila melanogaster* were performed on yeast corn medium at 22 °C

unless otherwise noted. The W^{1118} strain was used as a control for the respective genetic

backgrounds. *Mef2-Gal4* (III) was used to drive muscle-specific *Opa1-like* (Opa1)(Rai et al.,

2014; Vue et al., 2022). *Tub-Gal80^{ts}*; *Mef2 Gal4* (BS# 27390) was used for the conditional

Opa1-like muscle-specific gene KD. Genetic crosses were set up at 18 °C and then shifted to 29

°C at the larval stage. *UAS-mito-GFP* (II chromosome) was used to visualizemitochondria.

Stocks were requested or obtained from the Bloomington *Drosophila* stock center. All

chromosomes and gene symbols are as described in Flybase (http://flybase.org).

Mitochondrial Staining

Indirect flight muscles (IFMs) from 2–3-day-old adult flies were dissected and fixed in 4%

paraformaldehyde for 1.5–2 hours on a rotor with a minimum rpm. Samples were washed in

PBSTx (PBS +0.03% TritonX-100) three times each for 15 min and incubated in PBS with 500–

1000 nM Mitotracker (M22425, Molecular Probes: Eugene, OR, USA; 1 mM stock in dimethyl

sulfoxide, DMSO) at 37 °C for 30 min. *DMef2-Gal4* driven *UAS-mito-GFP* was also used to

image mitochondria. Muscle samples were washed and counterstained with Phalloidin-TRITC or

FITC (P1951-TRITC, Sigma-Aldrich, 50 µg/ml stock; 46950, Sigma-Aldrich) and mounted

using Prolong Glass Antifade Mountant with NucBlue (P36985, ThermoFisher, USA). Samples

were imaged using a ZEISS 780 confocal microscope and processed using ZEN software

(version 3.2.0.115).

Immunohistochemistry

Two-day-old adult fly thoraces were dissected to obtain IFMs in 4% paraformaldehyde

(Sigma), with muscle fixed in 4% paraformaldehyde (Sigma-Aldrich) for 1.5–2 hours at 25 °C

on a rotor. Samples were washed three times with PBSTx (Phosphate Saline Buffer with Tween)

for 15 min and blocked for 2 hours at 25 °C or overnight at 4 °C using 2% BSA (Sigma-

Aldrich). Samples were incubated with respective primary antibodies at 4 °C overnight. Later,

samples were washed three times for 10 min with PBSTx and incubated for 2.5 hours in

respective secondary antibodies at 25 °C or overnight at 4 °C. Samples were then incubated for

40 min with Phalloidin-TRITC (2.5 µg/ml; P1951, Sigma-Aldrich, USA) to counterstain

samples. After incubation for 20 min at 25 °C in Prolong Glass Antifade Mountant with NucBlue

stain (P36985, ThermoFisher, USA), samples were mounted. Images were acquired using the

ZEISS 780 confocal microscope and processed using ZEN software (version 3.2.0.115).

Antibodies used for the staining were rabbit anti-PAC (1:20), rabbit anti-VDAC1 (1:200), rabbit

anti-IP3 (1:200), and Alexa Fluor-488-labeled anti-rabbit IgG (1:500, ThermoFisher, USA).

Electron Microscopy (EM) Sample Preparation in Flies

Flies (2–3 days old) were dissected in standard fixative (2.5% glutaraldehyde, 1%

paraformaldehyde, and 0.12 M sodium cacodylate buffer). Individual *Drosophila* muscle fibers

(IFMs) were isolated and transferred to a fresh fixative tube. Fixation and imaging were

performed as described above.

RNA Sample Preparation in Flies

RNA was isolated from the IFMs of 1–2-day-old flies. IFMs were removed from the bisected

thoraces at 4 °C and immersed in TRIzol (Sigma-Aldrich). Total RNA was extracted in TRIzol

reagent (Invitrogen) using a tissue grinder. Reverse polymerase chain reaction was performed to

generate cDNA using SuperScript III (Invitrogen). For quantitative reverse transcriptase-PCR, 50

ng of cDNA were used for each reaction with iTAQ Universal Sybergreen (Biorad), using the

QuantiStudio 6 Flex system (Applied Biosystems). Gene expression was analyzed using the

$\Delta\Delta C_T$ method, and relative expression was normalized to GAPDH. Sequences for the primers are

listed in Supplementary Table 2.

ChIP-seq

ChIP-Seq data was analyzed from data sets generated previously (J. Han et al., 2013; Tameire et

al., 2019; Zou et al., 2022).

Statistical Analysis

Results are presented as the mean ± standard error of the mean. Data were analyzed using an

unpaired T-test. If more than two groups were compared, one-way analysis of variance

(ANOVA) was performed, and significance was assessed using Fisher's protected least

significance difference test. GraphPad PRISM and Statplus software package was used for t-tests

and ANOVA analyses (SAS Institute: Cary, NC, USA). For all statistical analyses, a significant

difference was accepted when $P < 0.05$.

Results

***Opa1-like* KD in *Drosophila* Flight Muscle Alters Mitochondria, Cristae, and MERC Morphology**

Mammalian OPA1 plays an essential role in mitochondrial fusion, and *OPA1* KD increases the number of fragmented mitochondria and the average number of mitochondria per cell (Cipolat et al., 2004; Frezza et al., 2006; Parra et al., 2013; Patten et al., 2014; Scorrano et al., 2002; Tezze et al., 2017; Varanita et al., 2015). We previously reported that *Opa1* downregulation in skeletal muscle of mice induces the mitochondrial fusion and MERC protein MFN2, suggesting that *Opa1* deletion may play a role in regulating MERCs (Pereira et al., 2017). To determine the role of OPA1 in MERC formation, maintenance, and function, we analyzed the effects of *Opa1-like* KD on MERC gene mRNA transcripts in fly muscle tissue (Fig. 1). Using quantitative polymerase chain reaction (qPCR), we find the expression of genes encoding ER and mitochondrial proteins, such as the mitochondrial fission protein dynamin-related protein 1 (*DRP1*) and the ER stress-related markers *ATF4*, *BIP,* and inositol-requiring enzyme 1 (*IRE1*) were increased in *Opa1-like* KD compared with wild-type (WT) *Drosophila* muscles (Fig. 1A) despite the reduced size and activity of the *Opa1-like* KD flies (Supplementary Figure [SF] 1A-B). However, no significant difference in the ER stress marker *ATF6* expression was observed between WT and *Opa1-like* KD muscles (Fig. 1A). Transcripts encoding Ca^{2+}-related MERC components, IP_3R3 (*Itpr3*) and GRP75 (*Hspa9),* were elevated in *Opa1-like* KD fly muscle (Fig. 1A). Interestingly, the gene expression levels of the OMM protein VDAC (Fig. 1A) were decreased. These results indicate that *Opa1-like* KD induces the transcription of MERC components, and ER stress genes.

To examine the effects of *Opa1-like* KD on mitochondrial morphology in *Drosophila*

skeletal muscle, we used confocal microscopy and transmission electron microscopy (TEM).

Confocal microscopy revealed that *Opa1-like* KD increased mitochondrial fragmentation (Fig.

1B-G), as assessed by mitochondrial number normalized to the distance of three sarcomeres (Fig.

1H). Mitochondrial area and aspect ratio (length:width ratio) were also reduced in the *Opa1-like*

KD muscle (Fig. 1I-J). Similar results were obtained by TEM (Fig. 1K-N), showing that *Opa1-*

like KD increased the mean circularity index, decreased the mean mitochondrial area, and

increased the number of mitochondria (Fig. 1O-Q). Additionally, we assessed the abundance and

quality of cristae using the cristae score, cristae area, cristae number, and cristae volume, which

were all decreased in *Opa1-like* KD mitochondria relative to control mitochondria (Fig. 1R–U).

In addition, *Opa1-like* KD was associated with loss of lamellar cristae and increased tubular

cristae. The increased mitochondrial fragmentation and alterations in cristae shape are consistent

with the many previous reports that OPA1 plays a critical role in remodeling mitochondrial and

cristae morphology (Cipolat et al., 2004; Frezza et al., 2006; Parra et al., 2013; Patten et al.,

2014; Scorrano et al., 2002; Tezze et al., 2017; Varanita et al., 2015).

After confirming the expected *Opa1-like* KD phenotype [Supplemental Video (SV)1;

SV2-3; Figure 1)., TEM was then used to visualize and quantify MERC morphology in *Opa1-*

like KD *Drosophila* muscle (Fig. 1V-Y). Relative to WT, *Opa1-like KD* trended toward an

increased MERC contact length (Fig. 1Z), decreased the mean MERC distance (Fig. 1AA),

increased the percentage of mitochondrial coverage by ER (Fig. 1AB), and increased the

percentage of ER coverage by mitochondria (Fig. 1AC), all suggesting that *Opa1-like* deficiency

increased MERC tethering. Because TEM only captures structural MERC changes in two

dimensions (2D), serial block face-scanning electron microscopy (SBF-SEM), which offers

numerous advantages in volumetric renderings (Marshall et al., 2023), was used to validate

MERC changes in a three-dimensional (3D) context. The percentage of mitochondrial surface

area covered by the ER and the percentage of the ER surface area covered by mitochondria in

Drosophila muscle both significantly increased in the *Opa1-like* KD relative to the WT muscle

(Fig. 1AD-AI, SF 2A-B, SV 4-5). Together, these results demonstrate that OPA1 deficiency

increases MERC formation in *Drosophila* skeletal muscle.

OPA1 Deficiency in Myotubes and Mouse Skeletal Muscle Increases MERC Sites and the

Expression of MERC-Tethering Proteins

To examine whether our *Drosophila* findings translated to mammalian muscle cells, we

transduced primary myoblast and myotube cell cultures derived from *Opa1* floxed mice with

recombinant adenovirus containing the green fluorescent protein gene (Ad-GFP) or Cre

recombinase (Ad-CRE) and evaluated MERC protein abundance and morphology. KO of *OPA1*

in primary myoblasts (SF 3A) induced ER stress, as indicated by increased BIP and ATF4

protein levels and increased levels of the MERC proteins MFN2 and GRP75 (SF 3B), similar to

what was observed in fly muscle. Structurally, increased ER and mitochondrial colocalization in

OPA1 KO myoblasts and myotubes compared to controls was first confirmed by the overlapping

fluorescent signals from Mitotracker and GRP78 (SF 3C-S). Additionally, *OPA1* KO cells

showed increased MFN2–MFN1 interactions in both primary myoblasts (SV 6-7) and myotubes

(SF 3T-Z) as assessed by *in situ* proximity ligation assays (PLA). At the ultrastructural level (SF

3AA-AD, AM-AP; SV   8-9), *OPA1* KO again decreased MERC distance (SF 3AE) and

increased the percentage of mitochondria surface covered by ER (SF 3AF, AQ) and the

percentage of the ER surface covered by mitochondria in two and three dimensions (2D/3D),

respectively (SF 3AG, AR), despite a decrease in mitochondrial size (SF 3AH). Also, similar to

results observed in *Drosophila*, OPA1 deficiency in myotubes decreased the number, area, score,

and volume of cristae (SF 3AI–AL). These data suggest that OPA1 loss increases MERC protein

expression and MERC tethering similarly in mammalian primary myoblasts and myotubes and

adult fly muscles.

To determine whether OPA1 regulation of MERCs extends to mature mammalian

skeletal muscle, we next investigated the effects of *OPA1* deletion on MERC structure and

expression in mice with skeletal muscle-specific inducible knockdown of OPAl (*OPA1* smKO)

(Figure 2). Consistent with the fly and cell culture data, OPA1-deficient skeletal muscles from

40-week-old mice, which represent adult mice, showed increased levels of MERC proteins BIP,

PACS2, GRP75, MFN2 and MFN1 (Fig. 2A-B). Similarly, mRNA levels of MERC genes as

assessed by qPCR were also increased in *OPA1* smKO muscles (Fig. 2C). Additionally, *in situ*

PLA (Fig. 2D-I) revealed increased MFN2–MFN1 (Fig. 2F: SV 10-11) and IP$_3$R3–VDAC

interactions in *OPA1* smKO muscle fibers (Fig. 2I; SV 12–13). TEM analysis (Fig. 2 J-M)

showed that MERC distance was decreased (Fig. 2N) and contact between mitochondrial and ER

membranes was increased in 2D and 3D images (Fig. 2O-V; SV 14-15) in *OPA1* smKO mice

compared to controls. These data demonstrate that MERC formation also increases in mouse

skeletal muscle following adult-onset OPA1 deficiency.

We previously reported that *OPA1* smKO animals exhibited ER stress activation and

increased levels of the MERC protein MFN2 and its interacting partner MFN1 (Pereira et al.,

2017). To evaluate the role of ER stress in OPA1-mediated MERC formation, four weeks after

tamoxifen injections, *OPA1* smKO mice were treated with either PBS or the chemical ER

chaperone tauroursodeoxycholic acid (TUDCA), which prevents unfolded protein response

dysfunction (Yoon et al., 2016), for 4 weeks to attenuate ER stress activation (Pereira et al.,

2017; Tezze et al., 2017). Following TUDCA treatment, we observed decreased expression of

Ca^{2+}-related MERC genes *VDAC3*, IP$_3$R*3, ATF4*, and *GRP75* in 12-week-old *OPA1* smKO mice

relative to untreated *OPA1* smKO mice (Figure 2W–Z). These results support the hypothesis that

ER stress may mediate the MERC response to OPA1 deficiency.

ATF4 Mediates MERC Formation and Spacing in *OPA1*-deficient Skeletal Muscle

To better understand the relationship between ER stress and MERC formation in OPA1-

deficient muscles, we hypothesized that the integrated stress response transcription factor ATF4,

which is induced in OPA1-deficient muscles and can be activated downstream of the unfolded

protein response (UPR), binds to the promoters of MERC genes to induce expression. By

performing an *in-silico* study using publicly available ChIP-Seq data, we identified ATF4

binding to the promoters of *Hspa9* (GRP75) and *Vdac3* (SF 4A-B), and potentially two sites in

Itpr3 that encodes IP$_3$R3 (SF 4C) in WT samples that are reduced or absent in the ATF4 KO

samples. For *Vdac3*, the major peak may be within an intron, suggesting activation downstream

of the promoter. These observations support a potential role for stress response factor ATF4 in

the observed changes in MERC formation and protein expression under OPA1-deficient

conditions.

To further investigate the relationships between ATF4, OPA1, and MERC formation

across several models, we evaluated MERC structure and protein abundance in primary mouse

myotubes with knockdown or overexpression of ATF4 in concert with OPA1 deficiency (Fig. 3).

ATF4 overexpression alone increased the abundance of Ca^{2+}-related MERC proteins, IP3R3 and

VDAC (Fig. 3A-D), suggesting involvement in MERC formation through ER stress. When we

knocked out both OPA1 and ATF4 in muscle cells, the mean mitochondrial area was

significantly decreased, whereas the mean mitochondrial area was significantly increased in

ATF4 OE muscle cells (Fig. 3E-K) suggesting that ATF4 functions to also modulate

mitochondrial fusion. Double OPA1 and ATF4 KO (DKO) and OPA1 KO/ATF4 OE resulted in

non-significant reductions in the mitochondrial area relative to control skeletal muscle cells (Fig.

3K), suggesting the pluralistic effects of these proteins serve moderating effects. In contrast to

OPA1 KO, ATF4 KO significantly increased MERC distance (Fig. 3L). Conversely, ATF4 OE

significantly decreased MERC distance compared to the control (Fig. 3L), indicating that ATF4

regulates the spacing between mitochondrial and ER membranes in myotubes. Moreover, in

DKO cells, a significant increase in MERC distance was observed, relative to both control and

OPA1 KO cells. These data suggest that OPA1 regulation of MERC spacing is dependent on

ATF4 expression.

We next sought to recapitulate these findings in *Drosophila* flight muscle, and SBF-SEM

was used to further quantify ER and MERC lengths in *ATF4* KD and *ATF4* OE muscle (Fig. 3M-

R, SV 16–18). We found a significant increase in ER length, or length in contact with ER, in

ATF4 OE muscle cells compared with control cells (Fig. 3S), while *ATF4* loss did not alter ER

length. The ER volume did not significantly change in *ATF4* OE, but significantly increased in

ATF4 KD cells, likely a byproduct of both increased MERC length and distance (Fig. 3T).

Together, these data support the hypothesis that ATF4 expression regulates MERC formation by

inducing the expression of MERC proteins, decreasing the width of the MERC space and

increasing MERC length and modulating ER morphology. Specifically, a change of both *ATF4*

through OE or KD can attenuate OPA1-dependent changes in mitochondrial size, while *ATF4*

expression is necessitated for significant MERC tethering.

OPA1 Deficiency Alters Mitochondrial Ca^{2+} Uptake

Given the alterations in mitochondrial and ER structure observed in relation to OPA1 and ATF4 expression, we sought to explore potential functional consequences of these changes. Prior work suggests that mitochondrial dynamics could play a role in intracellular calcium oscillations in part by regulating MERC formation (Casellas☐Díaz et al., 2021; De Brito & Scorrano, 2008; S. Han et al., 2021). We, therefore, sought to determine changes in Ca^{2+} organellar partitioning in OPA1-deficient myotubes. In cells treated with caffeine (20 mM) to release Ca^{2+} from the ER we estimated mitochondrial Ca^{2+} concentration using mitoPericam fluorescence from control (OPA1$^{fl/fl}$) and OPA1 KO (OPA1$^{fl/fl}$-Cre) primary myotubes (Fig. 4A). The area under the curve was decreased in OPA1 deficient cells, indicating that OPA1deficiency reduces mitochondrial Ca^{2+} uptake (Fig. 4B). The peak amplitude for mitoPericam was also decreased in OPA1 KO, which could implicate reduced mitochondrial calcium uniporter (MCU) activity (Fig. 4C). To further evaluate this possibility, we measured MCU levels in myoblasts, myotubes, and mouse skeletal muscle (SF 5). 4). In all three models, loss of OPA1 increased MCU expression (SF 5B, E, H). To assess ER/SR calcium regulation, levels of the sarcoplasmic/endoplasmic reticulum Ca^{2+}-ATPase were measured and noted to be increased upon loss of OPA1 (SF4 C, F, I). Using Fura-2 to assess total cellular calcium, we observed that OPA1 deficiency reduced the area under curve and peak amplitude following caffeine stimulation (Figure 4D -F). When Ca^{2+} re-uptake by the ER was inhibited by thapsigargin (1 µM), both area under the curve and peak amplitude of cytosolic Ca^{2+} were decreased in OPA1-deficient cells following caffeine (20 mM) (Fig. 4G-I). Together these findingsindicate reduced ER Ca^{2+} stores. Thus, OPA1 deficiencyreduces SR calcium, which could contribute to impaired mitochondrial Ca^{2+} uptake, despitecompensatory induction of MCU and SERCA2.

Given the ATF4 dependence of MERC modulation in OPA1 deficient cells described

above, we hypothesized that ATF4-dependent MERC formation contributed to altered Ca^{2+}

homeostasis and reducing ATF4 could restore Ca^{2+} homeostasis in cells lacking OPA1. We

compared cytosolic Ca^{2+} measurements using Fura-2 from control, OPA1 KO and OPA1/ATF4

DKO primary myotubes following caffeine exposure (Fig. 4J). Loss of OPA1 reduced the area

under the curve (Fig. 6K) and the peak amplitude, consistent with reduced Ca^{2+} release from the

ER into the cytosol (Fig. 4L). However, the loss of ATF4 alone or the simultaneous loss of

ATF4 and OPA1 restored area under the curve and peak amplitude to levels similar to those seen

in control mice (Fig. 4J-L). This shows that in DKO, ATF4 can rescue impaired calcium

homeostasis caused by OPA1 deficiency. These data suggest that ATF4-induced MERCs

remodeling contributes to dysregulated Ca^{2+} homeostasis in OPA1-deficient muscle cells (Fig.

4J-L).

<u>Discussion</u>

Previous studies have used 3D reconstruction to better elucidate diverse phenotypes of

mitochondria and ER communication modalities, including the description of curved sheets of

rough ER around mitochondria referred to as wrappER (Anastasia et al., 2021). However, to our

knowledge, no studies have yet investigated the changes incurred by the loss of OPA1 on MERC

formations using a mixed-technique approach including 3D reconstruction. Numerous studies

have shown that Protein kinase RNA-like ER Kinase (also known as PERK) is a facilitator of

MERC tethering during ER-stress (van Vliet & Agostinis, 2016; Verfaillie et al., 2012). OPA1

function has been linked to the ER in numerous ways, including facilitating calcium transfer and

homeostasis (Cartes-Saavedra et al., 2022). Thus, our study supports earlier work demonstrating

a link between OPA1 function and the regulation of MERCs. This study advances the field by

demonstrating the central role of ATF4 in regulating MERC formation in the context of altered

OPA1 expression.

MERCs mediate important intracellular communication processes between the ER and

the mitochondria, including Ca^{2+} and lipid transfer that could influence mitochondrial

morphology and apoptotic signaling (Giacomello & Pellegrini, 2016; Moltedo et al., 2019;

Sassano et al., 2022). MERC defects are associated with the pathophysiology of various

disorders including neurodegenerative illnesses, obesity, and diabetes (Kim & Roy, 2020; Liao et

al., 2017; Pereira et al., 2017; Tubbs et al., 2018; Verma et al., 2016). Evaluating mechanisms

that regulate MERC formation and maintenance has begun to elucidate the role of MERCs in

cellular physiology and metabolism. However, the relationship between mitochondrial dynamics,

mitochondrial stress, and MERCs remained incompletely understood. Here, we demonstrate that

OPA1 deficiency increased key MERC proteins, including MFN2, MFN1, PACS2, and BIP that

paralleled bioenergetics defects. OPA1-deficiency also altered MERC tethering at the

ultrastructural level and increased MERC volume and percent coverage in skeletal muscle, as

determined by SBF-SEM. Importantly, we also elucidated that ATF4 might regulate MERC gene

expression in response to mitochondrial stress, which could represent an initial compensatory

mechanism that becomes persistent and ultimately maladaptive.

Reduced OPA1 expression results in increased expression of the ER stress protein ATF4,

and of MFN2, a tethering MERC protein that regulates Ca^{2+} levels in the MERC space

(Ainbinder et al., 2015)·(De Brito & Scorrano, 2008; Naon et al., 2016; Pereira et al., 2017; Sood

et al., 2014). MERC thickness is thought to be precisely maintained to ensure normal Ca^{2+}

transport, as an insufficient distance can result in steric hindrance among various components of

the Ca^{2+} transporter machinery (Giacomello & Pellegrini, 2016). Ca^{2+} uptake between the ER

and mitochondria is more likely to occur when the organelles are in close proximity, with an

ideal distance ranging from 15–30 nm (Csordás et al., 2018; Giacomello & Pellegrini, 2016;

Thoudam et al., 2019). Reduced OPA1 expression decreased MERC distance, induced structural

changes in MERCs, and were correlated with an increase in the MERC proteins IP3R, GRP75,

and VDAC3 and are crucial for Ca^{2+} transfer between the two organelles. Interestingly, despite

this reduction in MERCs distance and increased MERC protein levels, we confirmed prior

results (Cartes-Saavedra et al., 2020) showing mitochondrial Ca^{2+} uptake was reduced in OPA1-

deficient myotubes. The discrepancy between the expression of proteins that promote

mitochondrial Ca^{2+} uptake (including MCU), versus the measured reductions in SR and

mitochondrial calcium content, underscore additional mechanisms through which OPA1 loss

impairs cellular Ca^{2+} homeostasis. Examining decreased store-operated Ca^{2+} entry during OPA1

loss may offer greater insight into Ca^{2+} availability. The induction of the ATF4-dependent

pathways identified, could represent an adaptive response. We validated previous results (Cretin

et al., 2021; Parra et al., 2013) showing that loss of OPA1 increases mitochondrial fragmentation

in concert with the induction of *Drp1,* a regulator of mitochondrial fission (Favaro et al., 2019;

Möpert et al., 2009). Mitochondrial size is also known to influence calcium uptake (Golic et al.,

2014; Handran et al., 1997; Kowaltowski et al., 2019), and represents an additional mechanism

by which OPA1 deficiency could impair intracellular calcium homeostasis.

Changes in MERC properties may also modulate ER to mitochondrial Ca^{2+} transfer,

which serves an essential role in the regulation of mitochondrial energy metabolism and

oxidative phosphorylation (Bravo et al., 2011; Glancy & Balaban, 2012; Rieusset, 2018).

Mitochondrial Ca^{2+} concentration regulates the activation of the TCA cycle, and increased Ca^{2+}

levels in the mitochondrial matrix activate multiple enzymes involved in energy production

(Carreras-Sureda et al., 2019; Denton et al., 1975; Glancy et al., 2013; Ivannikov & Macleod,

2013; Traaseth et al., 2004). Thus, the reduced intramitochondrial Ca^{2+} levels could contribute

to the bioenergetics impairment and altered crista morphology we found to be associated with

OPA1 deficiency. Altered MERC morphology in skeletal muscle following OPA1 loss could

represent a compensatory mechanism to restore Ca^{2+} homeostasis, given MERCs are understood

to serve as regulatory hubs in ER and mitochondrial Ca^{2+} homeostasis (Cartes-Saavedra et al.,

2020, 2021; Fülöp et al., 2011; Kushnareva et al., 2013). Similarly, the induction of IP_3R and

SERCA could also represent an adaptive responses. The inhibition of IP_3R mimics sustained ER

stress by impairing mitochondrial function, and inducing autophagy (Jia et al., 2019; Rashid et

al., 2015). Moreover, SERCA induction may improve mitochondrial quality control (Tan et al.,

2020), offering clues as to the pluralistic roles ATF4 expression has on mitochondrial structure

in addition to only MERC formation. However, we cannot exclude the possibility that the

associated ATF4-dependent MERCs changes observed in response to OPA1 deficiency could be

linked to the impairment of mitochondrial and SR calcium homeostasis via additional

mechanisms.

Changes in mitochondria–ER cross-talk accompany the early stages of ER stress,

suggesting that exchange in metabolites between the mitochondria and ER are altered in

response to various cellular insults (Bravo et al., 2011). The ER-stress-associated protein IRE1α,

which was induced in *Opa1-like* KD flies in our experiments, has non-canonical functions in

MERC regulation, Ca^{2+} flux between the ER and mitochondria, and in regulating mitochondrial

respiration(Carreras-Sureda et al., 2019), indicating a cloe association between ER stress, MERC

formation, and the regulation of mitochondrial metabolism.

OPA1 deficiency in murine gastrocnemius skeletal muscle induced ER stress-related

proteins, including the unfolded protein response (UPR) transcription factor ATF4, ATF6, and

CHOP(Cawley et al., 2011). Recent studies have described that the UPR can also modulate and

restore mitochondrial proteostasis which includes reducing unfolded protein load (Hetz, 2012),

representing an adaptive response to relieve ER stress (Rainbolt et al., 2014). ATF6-deficiency

has been linked to dysfunction in mitochondrial potential, homeostasis, and biogenesis (Wang et

al., 2018). Similarly, in neuroblastoma models, IRE1α-XBP1 inhibition decreases MERC spaces

(Chu et al., 2021). NRF2, another transcription factor induced by OPA1 deletion, plays

pleiotropic roles in mitochondrial biology (Buttari et al., 2022). CHOP which is induced during

ER stress might trigger mitochondria-dependent apoptosis (Hu et al., 2019). Thus, many lines of

evidence support a link between the UPR transcription factors and mitochondrial function. Our

in-silico study confirmed that ATF4 binds to downstream factors of MERC genes encoding

GRP75 and VDAC3. ATF4 induction following OPA1 deficiency may be an adaptive

mechanism for directly regulating MERC gene expression as a means of modifying MERC

formation. It is therefore possible that ATF6, XBP1, CHOP, and NRF2 also regulate MERC

gene expression by a similar mechanism. Previously the regulation of MERCs by the UPR has

been suggested (Amodio et al., 2021; Wilson & Metzakopian, 2021), yet other studies have

shown that MERCs can suppress proteins associated with the UPR (Li et al., 2023), underscoring

the need for additional studies into this interplay.

In summary, using a model of OPA1-deficiency-induced mitochondrial and ER stress we

identified a novel role for ATF4 induction in regulating MERC formation. OPA1 deficiency

632 induces mitochondrial dysfunction that leads to complex compensatory mechanisms, including

633 altered MERC structure and function and perturbed Ca^{2+} homeostasis. Our data suggest that

634 these mechanisms are, at least in part, mediated by ATF4, which regulates ER-mitochondria

635 coupling and calcium homeostasis in skeletal muscle cells likely via transcriptional regulation of

636 core MERC proteins and additional undetermined mechanisms.

640 **Collaboration Acknowledgements:**

641 We thank Dr. Christopher M. Adams for providing the Ad5CMVATF4/RSVeGFP and ATF4

642 floxed mice. We would like to also thank George R. Marcotte for his assistance in optimizing the

643 Ad5CMVATF4/RSVeGFP for myotubes.

644 **Funding Acknowledgements:**

645 This work was supported by NIH grants R01HL108379 and R01DK092065 to E.D.A; Division

646 of Intramural Research of the National Heart, Lung, and Blood Institute (NHLBI,

647 1ZIAHL006221 and the National Institute of Arthritis and Musculoskeletal and Skin Diseases

648 (NIAMS) to B.G.; The United Negro College Fund/Bristol-Myers Squibb E.E. Just Faculty

649 Fund, Burroughs Wellcome Fund Career Awards at the Scientific Interface Award, Burroughs

650 Wellcome Fund Ad-hoc Award, National Institutes of Health Small Research Pilot Subaward to

651 5R25HL106365-12 from the National Institutes of Health PRIDE Program, DK020593,

652 Vanderbilt Diabetes and Research Training Center for DRTC Alzheimer's Disease Pilot &

653 Feasibility Program to A.H.J.; National Institute of Health (NIH) NIDDK T-32, number

654 DK007563 entitled Multidisciplinary Training in Molecular Endocrinology to Z.V.; NIH NIA

655 RF1AG55549, NIH NINDS R01NS107265, RO1AG062135, AG59093, AG072899 to E.T.;

656 United Negro College Fund/Bristol-Myers Squibb E.E. Just Postgraduate Fellowship in the Life

657 Sciences Fellowship to H.K.B.; and NSF grant MCB #2011577I and NIH T32 5T32GM133353

658 to S.A.M. Its contents are solely the responsibility of the authors and do not necessarily represent

659 the official view of the NIH. The funders had no role in study design, data collection and

660 analysis, decision to publish, or preparation of the manuscript.

662 Competing interests:

663 The authors have no disclosures to report.

665 References:

666 Ainbinder, A., Boncompagni, S., Protasi, F., & Dirksen, R. T. (2015). Role of Mitofusin-2 in

667   mitochondrial localization and calcium uptake in skeletal muscle. *Cell Calcium*, *57*(1),

668   14–24. https://doi.org/10.1016/j.ceca.2014.11.002

669 Amodio, G., Pagliara, V., Moltedo, O., & Remondelli, P. (2021). Structural and Functional

670   Significance of the Endoplasmic Reticulum Unfolded Protein Response Transducers and

671   Chaperones at the Mitochondria–ER Contacts: A Cancer Perspective. *Frontiers in Cell*

672   *and Developmental Biology*, *9*.

673   https://www.frontiersin.org/articles/10.3389/fcell.2021.641194

674 Anastasia, I., Ilacqua, N., Raimondi, A., Lemieux, P., Ghandehari-Alavijeh, R., Faure, G.,

675   Mekhedov, S. L., Williams, K. J., Caicci, F., Valle, G., Giacomello, M., Quiroga, A. D.,

676   Lehner, R., Miksis, M. J., Toth, K., de Aguiar Vallim, T. Q., Koonin, E. V., Scorrano, L.,

677   & Pellegrini, L. (2021). Mitochondria-rough-ER contacts in the liver regulate systemic

678   lipid homeostasis. *Cell Reports*, *34*(11), 108873.

679   https://doi.org/10.1016/j.celrep.2021.108873

Bravo, R., Vicencio, J. M., Parra, V., Troncoso, R., Munoz, J. P., Bui, M., Quiroga, C.,

Rodriguez, A. E., Verdejo, H. E., Ferreira, J., Iglewski, M., Chiong, M., Simmen, T.,

Zorzano, A., Hill, J. A., Rothermel, B. A., Szabadkai, G., & Lavandero, S. (2011).

Increased ER-mitochondrial coupling promotes mitochondrial respiration and

bioenergetics during early phases of ER stress. *Journal of Cell Science*, *124*(Pt 13),

2143–2152. https://doi.org/10.1242/jcs.080762

Buttari, B., Arese, M., Oberley-Deegan, R. E., Saso, L., & Chatterjee, A. (2022). NRF2: A

crucial regulator for mitochondrial metabolic shift and prostate cancer progression.

Frontiers in Physiology, *13*.

https://www.frontiersin.org/articles/10.3389/fphys.2022.989793

Carreras-Sureda, A., Jaña, F., Urra, H., Durand, S., Mortenson, D. E., Sagredo, A., Bustos, G.,

Hazari, Y., Ramos-Fernández, E., Sassano, M. L., Pihán, P., van Vliet, A. R., González-

Quiroz, M., Torres, A. K., Tapia-Rojas, C., Kerkhofs, M., Vicente, R., Kaufman, R. J.,

Inestrosa, N. C., … Hetz, C. (2019). Non-canonical function of IRE1α determines

mitochondria-associated endoplasmic reticulum composition to control calcium transfer

and bioenergetics. *Nature Cell Biology*, *21*(6), 755–767. https://doi.org/10.1038/s41556-

019-0329-y

Cartes-Saavedra, B., Arancibia, D., Burté, F., Sjoberg, M., Estela Andres, M., Yu-Wai-Man, P.,

Hajnoczky, G., & Eisner, V. (2020). OPA1 GTPase and GE Domain-Specific Mutations

Differentially Alter Mitochondrial Fusion Dynamics and Calcium Homeostasis.

Biophysical Journal, *118*(3), 184a. https://doi.org/10.1016/j.bpj.2019.11.1125

Cartes-Saavedra, B., Macuada, J., Lagos, D., Arancibia, D., Andrés, M. E., Yu-Wai-Man, P.,

Hajnóczky, G., & Eisner, V. (2022). OPA1 Modulates Mitochondrial Ca2+ Uptake

Through ER-Mitochondria Coupling. *Frontiers in Cell and Developmental Biology*, *9*.

https://www.frontiersin.org/article/10.3389/fcell.2021.774108

Cartes-Saavedra, B., Yu-Wai-Man, P., Hajnoczky, G., & Eisner, V. (2021). Mitochondrial Ca2+

Uptake in OPA1 Cells is Upregulated through Functional ER-Mitochondria Coupling.

Biophysical Journal, *120*(3), 348a–349a. https://doi.org/10.1016/j.bpj.2020.11.2167

Casellas☐Díaz, S., Larramona☐Arcas, R., Riqué☐Pujol, G., Tena☐Morraja, P.,

Müller☐Sánchez, C., Segarra☐Mondejar, M., Gavaldà☐Navarro, A., Villarroya, F.,

Reina, M., & Martínez☐Estrada, O. M. (2021). Mfn2 localization in the ER is necessary

for its bioenergetic function and neuritic development. *EMBO Reports*, e51954.

Cawley, K., Deegan, S., Samali, A., & Gupta, S. (2011). Chapter Two—Assays for Detecting the

Unfolded Protein Response. In P. M. Conn (Ed.), *Methods in Enzymology* (Vol. 490, pp.

31–51). Academic Press. https://doi.org/10.1016/B978-0-12-385114-7.00002-7

Chu, B., Li, M., Cao, X., Li, R., Jin, S., Yang, H., Xu, L., Wang, P., & Bi, J. (2021). IRE1α-

XBP1 Affects the Mitochondrial Function of Aβ25–35-Treated SH-SY5Y Cells by

Regulating Mitochondria-Associated Endoplasmic Reticulum Membranes. *Frontiers in*

Cellular Neuroscience, *15*.

https://www.frontiersin.org/articles/10.3389/fncel.2021.614556

Cipolat, S., de Brito, O. M., Dal Zilio, B., & Scorrano, L. (2004). OPA1 requires mitofusin 1 to

promote mitochondrial fusion. *Proceedings of the National Academy of Sciences*,

101(45), 15927–15932. https://doi.org/10.1073/pnas.0407043101

Costa-Mattioli, M., & Walter, P. (2020). The integrated stress response: From mechanism to

disease. *Science*, *368*(6489), eaat5314. https://doi.org/10.1126/science.aat5314

Courson, J. A., Landry, P. T., Do, T., Spehlmann, E., Lafontant, P. J., Patel, N., Rumbaut, R. E.,

& Burns, A. R. (2021). Serial Block-Face Scanning Electron Microscopy (SBF-SEM) of

Biological Tissue Samples. *Journal of Visualized Experiments : JoVE*, *169*,

10.3791/62045. https://doi.org/10.3791/62045

Crabtree, A., Neikirk, K., Marshall, A. G., Vang, L., Whiteside, A. J., Williams, Q., Altamura, C.

T., Owens, T. C., Stephens, D., Shao, B., Koh, A., Killion, M., Lopez, E. G., Lam, J.,

Rodriguez, B., Mungai, M., Stanley, J., Dean, E. D., Koh, H.-J., … Hinton, A. (2023).

Defining Mitochondrial Cristae Morphology Changes Induced by Aging in Brown

Adipose Tissue. *Advanced Biology*, e2300186. https://doi.org/10.1002/adbi.202300186

Cretin, E., Lopes, P., Vimont, E., Tatsuta, T., Langer, T., Gazi, A., Sachse, M., Yu-Wai-Man, P.,

Reynier, P., & Wai, T. (2021). High-throughput screening identifies suppressors of

mitochondrial fragmentation in OPA1 fibroblasts. *EMBO Molecular Medicine*, *13*(6),

e13579. https://doi.org/10.15252/emmm.202013579

Csordás, G., Weaver, D., & Hajnóczky, G. (2018). Endoplasmic Reticulum–Mitochondrial

Contactology: Structure and Signaling Functions. *Trends in Cell Biology*, *28*(7), 523–

540. https://doi.org/10.1016/j.tcb.2018.02.009

De Brito, O. M., & Scorrano, L. (2008). Mitofusin 2 tethers endoplasmic reticulum to

mitochondria. *Nature*, *456*(7222), 605–610.

Delprat, B., Rieusset, J., & Delettre, C. (2019). Defective Endoplasmic Reticulum–Mitochondria

Connection Is a Hallmark of Wolfram Syndrome. *Contact*, *2*, 251525641984740.

https://doi.org/10.1177/2515256419847407

Denton, R. M., Randle, P. J., Bridges, B. J., Cooper, R. H., Kerbey, A. L., Pask, H. T., Severson,

D. L., Stansbie, D., & Whitehouse, S. (1975). Regulation of mammalian pyruvate

748  dehydrogenase. *Molecular and Cellular Biochemistry*, *9*(1), 27–53.

749  https://doi.org/10.1007/BF01731731

750 Ebert, S. M., Dyle, M. C., Kunkel, S. D., Bullard, S. A., Bongers, K. S., Fox, D. K., Dierdorff, J.

751  M., Foster, E. D., & Adams, C. M. (2012). Stress-induced skeletal muscle Gadd45a

752  expression reprograms myonuclei and causes muscle atrophy. *The Journal of Biological*

753  *Chemistry*, *287*(33), 27290–27301. https://doi.org/10.1074/jbc.M112.374777

754 Favaro, G., Romanello, V., Varanita, T., Desbats, M. A., Morbidoni, V., Tezze, C., Albiero, M.,

755  Canato, M., Gherardi, G., & De Stefani, D. (2019). DRP1-mediated mitochondrial shape

756  controls calcium homeostasis and muscle mass. *Nature Communications*, *10*(1), 1–17.

757 Frezza, C., Cipolat, S., De Brito, O. M., Micaroni, M., Beznoussenko, G. V., Rudka, T., Bartoli,

758  D., Polishuck, R. S., Danial, N. N., & De Strooper, B. (2006). OPA1 controls apoptotic

759  cristae remodeling independently from mitochondrial fusion. *Cell*, *126*(1), 177–189.

760 Fülöp, L., Szanda, G., Enyedi, B., Várnai, P., & Spät, A. (2011). The Effect of OPA1 on

761  Mitochondrial Ca2+ Signaling. *PLoS ONE*, *6*(9), e25199.

762  https://doi.org/10.1371/journal.pone.0025199

763 Garza-Lopez, E., Vue, Z., Katti, P., Neikirk, K., Biete, M., Lam, J., Beasley, H. K., Marshall, A.

764  G., Rodman, T. A., Christensen, T. A., Salisbury, J. L., Vang, L., Mungai, M.,

765  AshShareef, S., Murray, S. A., Shao, J., Streeter, J., Glancy, B., Pereira, R. O., … Hinton,

766  A. (2022). Protocols for Generating Surfaces and Measuring 3D Organelle Morphology

767  Using Amira. *Cells*, *11*(1), Article 1. https://doi.org/10.3390/cells11010065

768 Giacomello, M., & Pellegrini, L. (2016). The coming of age of the mitochondria–ER contact: A

769  matter of thickness. *Cell Death & Differentiation*, *23*(9), 1417–1427.

770  https://doi.org/10.1038/cdd.2016.52

Glancy, B., & Balaban, R. S. (2012). Role of mitochondrial Ca2+ in the regulation of cellular

energetics. *Biochemistry*, *51*(14), 2959–2973. https://doi.org/10.1021/bi2018909

Glancy, B., Willis, W. T., Chess, D. J., & Balaban, R. S. (2013). Effect of calcium on the

oxidative phosphorylation cascade in skeletal muscle mitochondria. *Biochemistry*,

52(16), 2793–2809. https://doi.org/10.1021/bi3015983

Golic, I., Velickovic, K., Markelic, M., Stancic, A., Jankovic, A., Vucetic, M., Otasevic, V.,

Buzadzic, B., Korac, B., & Korac, A. (2014). Calcium-induced alteration of

mitochondrial morphology and mitochondrial-endoplasmic reticulum contacts in rat

brown adipocytes. *European Journal of Histochemistry: EJH*, *58*(3), 2377.

https://doi.org/10.4081/ejh.2014.2377

Han, J., Back, S. H., Hur, J., Lin, Y.-H., Gildersleeve, R., Shan, J., Yuan, C. L., Krokowski, D.,

Wang, S., & Hatzoglou, M. (2013). ER-stress-induced transcriptional regulation

increases protein synthesis leading to cell death. *Nature Cell Biology*, *15*(5), 481–490.

Han, S., Zhao, F., Hsia, J., Ma, X., Liu, Y., Torres, S., Fujioka, H., & Zhu, X. (2021). The role of

Mfn2 in the structure and function of endoplasmic reticulum-mitochondrial tethering in

vivo. *Journal of Cell Science*, *134*(13), jcs253443. https://doi.org/10.1242/jcs.253443

Handran, S. D., Werth, J. L., DeVivo, D. C., & Rothman, S. M. (1997). Mitochondrial

morphology and intracellular calcium homeostasis in cytochrome oxidase-deficient

human fibroblasts. *Neurobiology of Disease*, *3*(4), 287–298.

https://doi.org/10.1006/nbdi.1996.0125

Hartwick Bjorkman, S., & Oliveira Pereira, R. (2021). The Interplay Between Mitochondrial

Reactive Oxygen Species, Endoplasmic Reticulum Stress, and Nrf2 Signaling in

Cardiometabolic Health. *Antioxidants & Redox Signaling*, ars.2020.8220.

https://doi.org/10.1089/ars.2020.8220

Hetz, C. (2012). The unfolded protein response: Controlling cell fate decisions under ER stress

and beyond. *Nature Reviews Molecular Cell Biology*, *13*(2), Article 2.

https://doi.org/10.1038/nrm3270

Hinton, A., Katti, P., Christensen, T. A., Mungai, M., Shao, J., Zhang, L., Trushin, S., Alghanem,

A., Jaspersen, A., Geroux, R. E., Neikirk, K., Biete, M., Lopez, E. G., Shao, B., Vue, Z.,

Vang, L., Beasley, H. K., Marshall, A. G., Stephens, D., … Abel, E. D. (2023). A

Comprehensive Approach to Sample Preparation for Electron Microscopy and the

Assessment of Mitochondrial Morphology in Tissue and Cultured Cells. *Advanced*

Biology, e2200202. https://doi.org/10.1002/adbi.202200202

Hirabayashi, Y., Kwon, S.-K., Paek, H., Pernice, W. M., Paul, M. A., Lee, J., Erfani, P.,

Raczkowski, A., Petrey, D. S., Pon, L. A., & Polleux, F. (2017). ER-mitochondria

tethering by PDZD8 regulates Ca^{2+} dynamics in mammalian neurons. *Science*,

358(6363), 623–630. https://doi.org/10.1126/science.aan6009

Hu, H., Tian, M., Ding, C., & Yu, S. (2019). The C/EBP Homologous Protein (CHOP)

Transcription Factor Functions in Endoplasmic Reticulum Stress-Induced Apoptosis and

Microbial Infection. *Frontiers in Immunology*, *9*.

https://www.frontiersin.org/articles/10.3389/fimmu.2018.03083

Ivannikov, M. V., & Macleod, G. T. (2013). Mitochondrial free Ca^{2+} levels and their effects on

energy metabolism in Drosophila motor nerve terminals. *Biophysical Journal*, *104*(11),

2353–2361. https://doi.org/10.1016/j.bpj.2013.03.064

Jia, S., Xu, X., Zhou, S., Chen, Y., Ding, G., & Cao, L. (2019). Fisetin induces autophagy in

pancreatic cancer cells via endoplasmic reticulum stress- and mitochondrial stress-

dependent pathways. *Cell Death & Disease*, *10*(2), Article 2.

https://doi.org/10.1038/s41419-019-1366-y

Kasai, S., Yamazaki, H., Tanji, K., Engler, M. J., Matsumiya, T., & Itoh, K. (2019). Role of the

ISR-ATF4 pathway and its cross talk with Nrf2 in mitochondrial quality control. *Journal*

of Clinical Biochemistry and Nutrition, *64*(1), 1–12. https://doi.org/10.3164/jcbn.18-37

Kim, D., & Roy, S. (2020). Effects of Diabetes on Mitochondrial Morphology and Its

Implications in Diabetic Retinopathy. *Investigative Ophthalmology & Visual Science*,

61(10), 10. https://doi.org/10.1167/iovs.61.10.10

Kowaltowski, A. J., Menezes-Filho, S. L., Assali, E. A., Gonçalves, I. G., Cabral-Costa, J. V.,

Abreu, P., Miller, N., Nolasco, P., Laurindo, F. R. M., Bruni-Cardoso, A., & Shirihai, O.

S. (2019). Mitochondrial morphology regulates organellar Ca2+ uptake and changes

cellular Ca2+ homeostasis. *FASEB Journal: Official Publication of the Federation of*

American Societies for Experimental Biology, *33*(12), 13176–13188.

https://doi.org/10.1096/fj.201901136R

Kushnareva, Y. E., Gerencser, A. A., Bossy, B., Ju, W.-K., White, A. D., Waggoner, J.,

Ellisman, M. H., Perkins, G., & Bossy-Wetzel, E. (2013). Loss of OPA1 disturbs cellular

calcium homeostasis and sensitizes for excitotoxicity. *Cell Death & Differentiation*,

20(2), 353–365. https://doi.org/10.1038/cdd.2012.128

Lam, J., Katti, P., Biete, M., Mungai, M., AshShareef, S., Neikirk, K., Garza Lopez, E., Vue, Z.,

Christensen, T. A., Beasley, H. K., Rodman, T. A., Murray, S. A., Salisbury, J. L.,

Glancy, B., Shao, J., Pereira, R. O., Abel, E. D., & Hinton, A. (2021). A Universal

Approach to Analyzing Transmission Electron Microscopy with ImageJ. *Cells*, *10*(9),

2177. https://doi.org/10.3390/cells10092177

Li, T., Zhao, H., Guo, G., Xia, S., & Wang, L. (2023). VMP1 affects endoplasmic reticulum

stress sensitivity via differential modulation of the three unfolded protein response arms.

Cell Reports, *42*(3), 112209. https://doi.org/10.1016/j.celrep.2023.112209

Liao, C., Ashley, N., Diot, A., Morten, K., Phadwal, K., Williams, A., Fearnley, I., Rosser, L.,

Lowndes, J., Fratter, C., Ferguson, D. J. P., Vay, L., Quaghebeur, G., Moroni, I., Bianchi,

S., Lamperti, C., Downes, S. M., Sitarz, K. S., Flannery, P. J., … Poulton, J. (2017).

Dysregulated mitophagy and mitochondrial organization in optic atrophy due to OPA1

mutations. *Neurology*, *88*(2), 131–142. https://doi.org/10.1212/WNL.0000000000003491

Marshall, A. G., Neikirk, K., Stephens, D. C., Vang, L., Vue, Z., Beasley, H. K., Crabtree, A.,

Scudese, E., Lopez, E. G., Shao, B., Krystofiak, E., Rutledge, S., Davis, J., Murray, S. A.,

Damo, S. M., Katti, P., & Hinton, A. (2023). Serial Block Face-Scanning Electron

Microscopy as a Burgeoning Technology. *Advanced Biology*, e2300139.

https://doi.org/10.1002/adbi.202300139

Moltedo, O., Remondelli, P., & Amodio, G. (2019). The mitochondria–endoplasmic reticulum

contacts and their critical role in aging and age-associated diseases. *Frontiers in Cell and*

Developmental Biology, *7*, 172.

Möpert, K., Hajek, P., Frank, S., Chen, C., Kaufmann, J., & Santel, A. (2009). Loss of Drp1

function alters OPA1 processing and changes mitochondrial membrane organization.

Experimental Cell Research, *315*(13), 2165–2180.

https://doi.org/10.1016/j.yexcr.2009.04.016

Mustafi, D., Kikano, S., & Palczewski, K. (2014). Serial Block Face☐Scanning Electron

Microscopy: A Method to Study Retinal Degenerative Phenotypes. *Current Protocols in*

Mouse Biology, *4*(4), 197–204.

Naon, D., Zaninello, M., Giacomello, M., Varanita, T., Grespi, F., Lakshminaranayan, S.,

Serafini, A., Semenzato, M., Herkenne, S., Hernández-Alvarez, M. I., Zorzano, A., De

Stefani, D., Dorn, G. W., & Scorrano, L. (2016). Critical reappraisal confirms that

Mitofusin 2 is an endoplasmic reticulum–mitochondria tether. *Proceedings of the*

National Academy of Sciences, *113*(40), 11249–11254.

https://doi.org/10.1073/pnas.1606786113

Neikirk, K., Lopez, E.-G., Marshall, A. G., Alghanem, A., Krystofiak, E., Kula, B., Smith, N.,

Shao, J., Katti, P., & Hinton, A. O. (2023). Call to Action to Properly Utilize Electron

Microscopy to Measure Organelles to Monitor Disease. *European Journal of Cell*

Biology, 151365. https://doi.org/10.1016/j.ejcb.2023.151365

Neikirk, K., Vue, Z., Katti, P., Rodriguez, B. I., Omer, S., Shao, J., Christensen, T., Garza Lopez,

E., Marshall, A., Palavicino-Maggio, C. B., Ponce, J., Alghanem, A. F., Vang, L.,

Barongan, T., Beasley, H. K., Rodman, T., Stephens, D., Mungai, M., Correia, M., …

Hinton Jr., A. O. (2023). Systematic Transmission Electron Microscopy-Based

Identification and 3D Reconstruction of Cellular Degradation Machinery. *Advanced*

Biology, *7*(6), 2200221. https://doi.org/10.1002/adbi.202200221

Parra, V., Verdejo, H. E., Iglewski, M., del Campo, A., Troncoso, R., Jones, D., Zhu, Y.,

Kuzmicic, J., Pennanen, C., Lopez☐Crisosto, C., Jaña, F., Ferreira, J., Noguera, E.,

Chiong, M., Bernlohr, D. A., Klip, A., Hill, J. A., Rothermel, B. A., Abel, E. D., …

Lavandero, S. (2013). Insulin Stimulates Mitochondrial Fusion and Function in

Cardiomyocytes via the Akt-mTOR-NFκB-Opa-1 Signaling Pathway. *Diabetes*, *63*(1),

75–88. https://doi.org/10.2337/db13-0340

Patra, M., Mahata, S. K., Padhan, D. K., & Sen, M. (2016). CCN6 regulates mitochondrial

function. *Journal of Cell Science*, *129*(14), 2841–2851.

https://doi.org/10.1242/jcs.186247

Patten, D. A., Wong, J., Khacho, M., Soubannier, V., Mailloux, R. J., Pilon-Larose, K.,

MacLaurin, J. G., Park, D. S., McBride, H. M., Trinkle-Mulcahy, L., Harper, M.-E.,

Germain, M., & Slack, R. S. (2014). OPA1-dependent cristae modulation is essential for

cellular adaptation to metabolic demand. *The EMBO Journal*, *33*(22), 2676–2691.

https://doi.org/10.15252/embj.201488349

Pereira, R. O., Tadinada, S. M., Zasadny, F. M., Oliveira, K. J., Pires, K. M. P., Olvera, A.,

Jeffers, J., Souvenir, R., Mcglauflin, R., & Seei, A. (2017). OPA 1 deficiency promotes

secretion of FGF 21 from muscle that prevents obesity and insulin resistance. *The EMBO*

Journal, *36*(14), 2126–2145.

Rai, M., Katti, P., & Nongthomba, U. (2014). Drosophila Erect wing (Ewg) controls

mitochondrial fusion during muscle growth and maintenance by regulation of the Opa1-

like gene. *Journal of Cell Science*, *127*(Pt 1), 191–203.

https://doi.org/10.1242/jcs.135525

Rainbolt, T. K., Saunders, J. M., & Wiseman, R. L. (2014). Stress-responsive regulation of

mitochondria through the ER unfolded protein response. *Trends in Endocrinology &*

Metabolism, *25*(10), 528–537. https://doi.org/10.1016/j.tem.2014.06.007

Rashid, H.-O., Yadav, R. K., Kim, H.-R., & Chae, H.-J. (2015). ER stress: Autophagy induction,

inhibition and selection. *Autophagy*, *11*(11), 1956–1977.

https://doi.org/10.1080/15548627.2015.1091141

Rieusset, J. (2018). The role of endoplasmic reticulum-mitochondria contact sites in the control

of glucose homeostasis: An update. *Cell Death & Disease*, *9*(3), 388.

https://doi.org/10.1038/s41419-018-0416-1

Romanello, V., & Sandri, M. (2015). Mitochondrial Quality Control and Muscle Mass

Maintenance. *Frontiers in Physiology*, *6*, 422. https://doi.org/10.3389/fphys.2015.00422

Sasaki, K., Uchiumi, T., Toshima, T., Yagi, M., Do, Y., Hirai, H., Igami, K., Gotoh, K., & Kang,

D. (2020). Mitochondrial translation inhibition triggers ATF4 activation, leading to

integrated stress response but not to mitochondrial unfolded protein response. *Bioscience*

Reports, *40*(11), BSR20201289. https://doi.org/10.1042/BSR20201289

Sassano, M. L., Felipe-Abrio, B., & Agostinis, P. (2022). ER-mitochondria contact sites; a

multifaceted factory for Ca2+ signaling and lipid transport. *Frontiers in Cell and*

Developmental Biology, *10*.

https://www.frontiersin.org/articles/10.3389/fcell.2022.988014

Scorrano, L., Ashiya, M., Buttle, K., Weiler, S., Oakes, S. A., Mannella, C. A., & Korsmeyer, S.

J. (2002). A distinct pathway remodels mitochondrial cristae and mobilizes cytochrome c

during apoptosis. *Developmental Cell*, *2*(1), 55–67. https://doi.org/10.1016/s1534-

5807(01)00116-2

Sood, A., Jeyaraju, D. V., Prudent, J., Caron, A., Lemieux, P., McBride, H. M., Laplante, M.,

Tóth, K., & Pellegrini, L. (2014). A Mitofusin-2-dependent inactivating cleavage of Opa1

links changes in mitochondria cristae and ER contacts in the postprandial liver.

Proceedings of the National Academy of Sciences of the United States of America,

111(45), 16017–16022. https://doi.org/10.1073/pnas.1408061111

Tameire, F., Verginadis, I. I., Leli, N. M., Polte, C., Conn, C. S., Ojha, R., Salas Salinas, C.,

Chinga, F., Monroy, A. M., Fu, W., Wang, P., Kossenkov, A., Ye, J., Amaravadi, R. K.,

Ignatova, Z., Fuchs, S. Y., Diehl, J. A., Ruggero, D., & Koumenis, C. (2019). ATF4

couples MYC-dependent translational activity to bioenergetic demands during tumour

progression. *Nature Cell Biology*, *21*(7), Article 7. https://doi.org/10.1038/s41556-019-

0347-9

Tan, Y., Mui, D., Toan, S., Zhu, P., Li, R., & Zhou, H. (2020). SERCA Overexpression

Improves Mitochondrial Quality Control and Attenuates Cardiac Microvascular

Ischemia-Reperfusion Injury. *Molecular Therapy. Nucleic Acids*, *22*, 696–707.

https://doi.org/10.1016/j.omtn.2020.09.013

Tezze, C., Romanello, V., Desbats, M. A., Fadini, G. P., Albiero, M., Favaro, G., Ciciliot, S.,

Soriano, M. E., Morbidoni, V., Cerqua, C., Loefler, S., Kern, H., Franceschi, C., Salvioli,

S., Conte, M., Blaauw, B., Zampieri, S., Salviati, L., Scorrano, L., & Sandri, M. (2017).

Age-Associated Loss of OPA1 in Muscle Impacts Muscle Mass, Metabolic Homeostasis,

Systemic Inflammation, and Epithelial Senescence. *Cell Metabolism*, *25*(6), 1374-

1389.e6. https://doi.org/10.1016/j.cmet.2017.04.021

Thoudam, T., Ha, C.-M., Leem, J., Chanda, D., Park, J.-S., Kim, H.-J., Jeon, J.-H., Choi, Y.-K.,

Liangpunsakul, S., Huh, Y. H., Kwon, T.-H., Park, K.-G., Harris, R. A., Park, K.-S.,

Rhee, H.-W., & Lee, I.-K. (2019). PDK4 Augments ER-Mitochondria Contact to

Dampen Skeletal Muscle Insulin Signaling During Obesity. *Diabetes*, *68*(3), 571–586.

https://doi.org/10.2337/db18-0363

Traaseth, N., Elfering, S., Solien, J., Haynes, V., & Giulivi, C. (2004). Role of calcium signaling

in the activation of mitochondrial nitric oxide synthase and citric acid cycle. *Biochimica*

Et Biophysica Acta, *1658*(1–2), 64–71. https://doi.org/10.1016/j.bbabio.2004.04.015

Tubbs, E., Chanon, S., Robert, M., Bendridi, N., Bidaux, G., Chauvin, M.-A., Ji-Cao, J., Durand,

C., Gauvrit-Ramette, D., Vidal, H., Lefai, E., & Rieusset, J. (2018). Disruption of

Mitochondria-Associated Endoplasmic Reticulum Membrane (MAM) Integrity

Contributes to Muscle Insulin Resistance in Mice and Humans. *Diabetes*, *67*(4), 636–

650. https://doi.org/10.2337/db17-0316

Tubbs, E., Theurey, P., Vial, G., Bendridi, N., Bravard, A., Chauvin, M.-A., Ji-Cao, J., Zoulim,

F., Bartosch, B., & Ovize, M. (2014). Mitochondria-associated endoplasmic reticulum

membrane (MAM) integrity is required for insulin signaling and is implicated in hepatic

insulin resistance. *Diabetes*, *63*(10), 3279–3294.

van Vliet, A. R., & Agostinis, P. (2016). When under pressure, get closer: PERKing up

membrane contact sites during ER stress. *Biochemical Society Transactions*, *44*(2), 499–

504. https://doi.org/10.1042/BST20150272

Varanita, T., Soriano, M. E., Romanello, V., Zaglia, T., Quintana-Cabrera, R., Semenzato, M.,

Menabò, R., Costa, V., Civiletto, G., & Pesce, P. (2015). The OPA1-dependent

mitochondrial cristae remodeling pathway controls atrophic, apoptotic, and ischemic

tissue damage. *Cell Metabolism*, *21*(6), 834–844.

Verfaillie, T., Rubio, N., Garg, A. D., Bultynck, G., Rizzuto, R., Decuypere, J.-P., Piette, J.,

Linehan, C., Gupta, S., Samali, A., & Agostinis, P. (2012). PERK is required at the ER-

mitochondrial contact sites to convey apoptosis after ROS-based ER stress. *Cell Death &*

Differentiation, *19*(11), Article 11. https://doi.org/10.1038/cdd.2012.74

Verma, A., Zhu, P., Prasad, T., Liang, Z., & Li, Q. (2016). Reduced Optic Atrophy 1 (OPA1)

gene expression in diabetic retina- implication of mitochondrial dynamics in pathogenesis

of diabetic retinopathy. *Investigative Ophthalmology & Visual Science*, *57*(12), 5446.

Vue, Z., Garza-Lopez, E., Neikirk, K., Vang, L., Beasley, H., Marshall, A. G., Murray, S. A., &

Hinton Jr, A. (2022). Mouse Skeletal Muscle Decrease in the MICOS Complex and

Altered Mitochondrial Networks with age. *bioRxiv*, 2022–03.

Vue, Z., Neikirk, K., Vang, L., Garza-Lopez, E., Christensen, T. A., Shao, J., Lam, J., Beasley,

H. K., Marshall, A. G., Crabtree, A., Anudokem, J., Rodriguez, B., Kirk, B., Bacevac, S.,

Barongan, T., Shao, B., Stephens, D. C., Kabugi, K., Koh, H.-J., … Hinton, A. (2023).

Three-Dimensional Mitochondria Reconstructions of Murine Cardiac Muscle Changes in

Size Across Aging. *American Journal of Physiology-Heart and Circulatory Physiology*.

https://doi.org/10.1152/ajpheart.00202.2023

Wang, S., Hu, B., Ding, Z., Dang, Y., Wu, J., Li, D., Liu, X., Xiao, B., Zhang, W., Ren, R., Lei,

J., Hu, H., Chen, C., Chan, P., Li, D., Qu, J., Tang, F., & Liu, G.-H. (2018). ATF6

safeguards organelle homeostasis and cellular aging in human mesenchymal stem cells.

Cell Discovery, *4*(1), Article 1. https://doi.org/10.1038/s41421-017-0003-0

Wilson, E. L., & Metzakopian, E. (2021). ER-mitochondria contact sites in neurodegeneration:

Genetic screening approaches to investigate novel disease mechanisms. *Cell Death &*

Differentiation, *28*(6), Article 6. https://doi.org/10.1038/s41418-020-00705-8

Yoon, Y. M., Lee, J. H., Yun, S. P., Han, Y.-S., Yun, C. W., Lee, H. J., Noh, H., Lee, S.-J., Han,

H. J., & Lee, S. H. (2016). Tauroursodeoxycholic acid reduces ER stress by regulating of

Akt-dependent cellular prion protein. *Scientific Reports*, *6*(1), Article 1.

https://doi.org/10.1038/srep39838

Zou, Z., Ohta, T., Miura, F., & Oki, S. (2022). ChIP-Atlas 2021 update: A data-mining suite for

exploring epigenomic landscapes by fully integrating ChIP-seq, ATAC-seq and Bisulfite-

seq data. *Nucleic Acids Research*, gkac199. https://doi.org/10.1093/nar/gkac199

1000

Figure Legends:

Figure 1. *Opa1-Like* knockdown (KD) in skeletal muscle alters mitochondrial morphology and

mitochondrial-ER contact sites in *Drosophila*. (A) mRNA expression levels of genes encoding

mitochondria–endoplasmic reticulum (ER) contact (MERC) proteins, *IP$_3$R3*, *GRP75* and *VDAC*

and ER stress induced proteins in *Opa1-like* KD compared with WT (Control) IFMs. (B-G)

Adult myofibrils were stained with Phalloidin-FITC (F-actin) and mito-GFP (mitochondria).

IFMs from WT flies revealed mitochondria with tubular morphology, whereas the *Opa1-like* KD

flies showed clusters of spherical mitochondria. (H) Mitochondria number (per three sarcomeres)

in *Opa1-like* KD (n = 34) and WT (n = 32) IFMs. (I) Quantification of mitochondrial area. (J)

Mitochondrial aspect ratio (major axis/minor axis) in *Opa1-like KD* (n = 141) and WT (n = 105)

IFMs. (K-N) Transmission electron microscopy (TEM) images showing cristae and

mitochondrial morphology in WT and *Opa1-like* KD IFMs (n = 20). (O) Circularity index in

Opa1-like KD IFMs relative to WT IFMs. (P). Quantification of mitochondrial area in *Opa1-like*

KD and WT. (Q) Mitochondria number in *Opa1-like* KD and WT. (R) Cristae quality and

abundance was assessed using the cristae score, on a scale from 0 (no sharply defined cristae) to

4 (many regular cristae). The average cristae score was significantly lower in *Opa1-like* KD than

in WT. (S) Quantification of cristae area in *Opa1-like* KD and WT. (T) Cristae number in *Opa1-*

like KO and WT. (U) Cristae volume in *Opa1-like* KD and WT. (V-Y) TEM images displaying

MERCS in WT and *Opa1-like* KD IFMs (n = 20). (Z) Quantification of MERC lengths between

Opa1-like KD and WT IFMs. (AA) Quantification of MERC distance in *Opa1-like* KD and WT

IFMs. (AB) Percentage of mitochondrial surface area in direct contact with the ER in *Opa1-like*

KD and WT IFMs. (AC) Percentage of ER surface area in direct contact with mitochondria in

Opa1-like KD and WT IFMs. (AD-AG) The 3D distribution of single continuous and stationary

mitochondria (purple, purple arrows) and ER (blue, blue arrows), reconstructed from serial block

face-scanning electron microscopy (SBF-SEM) image stacks of Drosophila indirect flight muscle

(IFM) fibers. (AH, AI) Percentage of ER surface area in contact with mitochondria (AH) and

percentage mitochondrial surface area in direct contact with ER (AI) in *Opa1-like* KD compared

with wild-type (WT) *Drosophila* skeletal muscle (SV1 [WT] SV2-3 [*Opa1-like* KD]). SBF-SEM

reconstructions from 7 to 23 fully constructed mitochondria, ER, or MERCs. *Significance was*

*determined by two-tailed Student's t-test. * P<0.05, ** P<0.01, *** P<0.001, **** P < 0.0001.*

1031

Figure 2. OPA1 deficiency promotes mitochondria–endoplasmic reticulum (ER) contact

(MERC) tethering in murine gastrocnemius skeletal muscle. (A) Representative immunoblots

showing MERC protein levels (normalized to GAPDH) in 40-week-old WT or *OPA1* smKO

muscle (n = 6). (B) Densitometric quantification showing a significant increase in protein levels

of BIP, GRP-75, MFN-2, and MFN-1 in *OPA1* smKO compared with WT mice. (C)

Quantification of mRNA expression of *OPA1*, ER stress genes *ATF4*, *BIP*, *CHOP*, and *ATF6*,

MERC-tethering gene *MFN-2*, and calcium-related MERC genes IP$_3$R*3*, *Grp75*, and *VDAC3* in

40-week-old *OPA1* smKO compared with WT mice (n = 5). Data are expressed as fold changes

vs. WT mice. (D-I). *In situ* proximal ligation assay (PLA) visualization of MFN1–MNF2 (D-E,

red punctae) and IP$_3$R3–VDAC interactions (G-H, red punctae) from 40-week-old WT and *Opa1*

smKO mice. Quantification demonstrating (F) increased MFN1–MFN2 and (I) IP$_3$R3–VDAC

interactions in *Opa1* smKO mice compared with WT (n = 3). (J-M). Transmission electron

microscopy (TEM) images showing MERCs in 20-week-old wild-type (WT) and *Opa1* skeletal

muscle specific knockout (*Opa1* smKO) mice (n = 3). (N). MERC distance in *Opa1* smkO and

WT controls. (O, P). Percentage coverage of (O) mitochondrial surface area and (P) ER surface

area in *Opa1* smKO relative to WT. (Q-T) The 3D distribution of single continuous and

stationary mitochondria (purple, purple arrows) and ER (blue, blue arrows), reconstructed from

serial block face-scanning electron microscopy (SBF-SEM) image stacks of *Opa1* skeletal

muscle specific knockout (*Opa1* smKO) mouse muscle. (U, V). Percentage of ER surface area in

contact with mitochondria (U) and the percentage of mitochondria surface area in direct contact

with ER (V) in *OPA1* KO compared with WT skeletal muscle. (W–Z). mRNA levels in

12☐week☐old *OPA1* smKO and WT mice 4 weeks after TUDCA treatment (n = 3). *IP$_3$R3*, (W)

ATF4, (X) *GRP75 (Y)*, and *VDAC3* (Z) gene expression were significantly reduced in *OPA1*-

smKO mice treated with TUDCA compared to those without TUDCA. Data are presented as fold

changes vs. WT mice. *SBF-SEM reconstructions from 7 to 23 fully constructed mitochondria,*

ER, or MERCs. Data are presented as the mean ± SEM. Significance was determined by

Student's t-test. *P < 0.05, **P < 0.01, ***P < 0.001, **** P < 0.0001.

1059

Figure 3. ATF4 regulates mitochondria–endoplasmic reticulum (ER) contact (MERC) formation

in *Opa1*-deficient muscle. (A) Representative immunoblot assessing ATF4, IP$_3$R3, and VDAC

levels from control (with no genetic modifications) or Ad-*Atf4* overexpression (OE) mouse

skeletal muscle myotubes (n=6) (B-D). Densitometric quantification demonstrating an increase

in (B) ATF4, (C) IP$_3$R3 and (D) VDAC protein levels in *Atf4* OE compared with control

myotubes. (E-J) TEM panel comparing MERCs area in control, *Opa1* KO, *Atf4* KO, *Atf4* OE,

double *Opa1/Atf4* KO (DKO), and *Opa1* KO/*Atf4* OE myotubes. Representative mitochondria

are identified with purple arrows. (K) Mitochondrial area in *Opa1* KO, *Atf4* KO, *Atf4* OE,

Opa1/Atf4 KO (DKO), and *Opa1* KO/*Atf4* OE relative to control myotubes. (L) MERC distance

in *Opa1* KO, *Atf4* KO, *Atf4* OE, *Opa1/Atf4* KO (DKO), and *Opa1* KO/*Atf4* OE relative to control

myotubes. (M-R). Individual electron micrograph of MERCs in control (M), *Atf4* KD (N), and

Atf4 OE myotubes and their serial block face-scanning electron microscopy (SBF-SEM) 3D

reconstructions (P-R), showing continuous, and stationary mitochondria (blue), nuclei (yellow),

and ER (pink) in *Drosophila* flight muscle. Quantification of ER length (S) and ER volume (T)

in control, *Atf4* KD and *Atf4* OE in *Drosophila* flight muscle. *Significance was determined by*

Student's t-test. *P < 0.05, **P < 0.01, ***P < 0.001, **** P < 0.0001.

1076
1077

Figure 4. ATF4 regulates calcium homeostasis in OPA1 deficient skeletal myotubes. (A).

Mitochondria Ca^{2+} measurements after caffeine administration (20mM) using $_{mt}$pericam in

control and *Opa1* KO primary myotubes. (n=3). (B). Area under the curve for pericam showing

decreased mitochondrial Ca^{2+} in *Opa1* KO. (C). Quantification of peak amplitude pericam signal

showing reduction in OPA1 primary myotubes. (D). Caffeine-induced (20mM) cytosolic Ca^{2+}

tracing measured by Fura-2 fluorescence in control and *Opa1* KO primary myotubes(n=5). (E).

Decreased area under the curve for cytosolic Ca^{2+}in *Opa1* KO myotubes. (F). Quantification of

peak Fura-2 amplitude signal in *Opa1* primary myotubes compared to control. (G). Cytosolic

Ca^{2+} levels estimated by Fura-2 in control and *Opa1*-KO primary myotubes following treatment

with Thapsigargin (Thaps) (1uM) and Caffeine (20mM) (n=5). (H). Area under the curve for

Fura-2 fluorescence in *Opa1* KO myotubes compared to control. (I). Quantification of Fura-2

peak amplitude signal in *Opa1* primary myotubes compared to control. (J). Tracing of cytosolic

calcium by Fura-2 in primary myotubes from *Opa1* KO, *Atf4* KO, and double KO (DKO) of

Opa1 and *Atf4*. (K). Area under the curve is lower in *Opa1* KO compared control, but

normalized in DKO, which is not significantly different relative to control. (L). Peak amplitude

quantification significantly decreases in KO of *Opa1* compared to control and is normalized in

DKO.

1095

1096

1097
1098   **Supplementary Materials:**

1099
1100   **Supplemental Figure 1.** Reduced pupal size and reduced mobility in *Opa1-like* knockdown

1101   (KD) (A). Representative images of pupa showing reduced size of *Opa1-like* KD compared with

1102   WT (control) fly. (B). The number of steps per second were significantly decreased in adult

1103   *Opa1-like* KD flies compared with WT. *Data are presented as the mean ± SEM. Significance*

1104   *was determined using Student's t-test. ****P < 0.0001.*

1105

1106   **Supplemental Figure 2.** Serial block face-scanning electron microscopy (SBF-SEM)

1107   demonstrates that the loss of *Opa1* increases mitochondria–endoplasmic reticulum (ER) contacts

1108   (MERCs). MERCs are shown in white and mitochondria are purple in wild type (left) and *Opa1*

1109   deficient (right) in (A-B) *Drosophila* muscle, (C-D) primary skeletal muscle myotubes, and (E-

1110   F) murine muscle.

1111

Supplemental Figure 3. OPA1 knockout (*Opa1* KO) increases mitochondria–endoplasmic

reticulum (ER) contact (MERC) formation in primary skeletal myoblasts and myotubes. (A)

Representative immunoblots of ER stress proteins, ATF4 and BIP, and MERC-tethering

proteins, MFN-2 and GRP-75, after infecting *Opa1$^{fl/fl}$* myotubes with Ad-Cre (*Opa1$^{fl/fl}$*-Cre). (B)

Densitometric quantification demonstrating an increase in BIP, ATF4, MFN2, and GRP75 levels

in *Opa1$^{fl/fl}$*-Cre myotubes compared with WT, normalized to alpha tubulin. (C-J). Representative

confocal images of *Opa1$^{fl/fl}$* and *Opa1$^{fl/fl}$*-Cre myoblasts (C-J) and myotubes (K-P), with DAPI-

labeled nuclei (blue) (C,G), mitotracker-labeled mitochondria (red) (D,H,K,N), GRP78 (E, I, L,

O) showing the ER (green), and a composite image showing all channels (F, J, M, P; n = 10

cells). (Q) Co-localization (Pearson's correlation coefficient) is increased for ER and

mitochondria in *Opa1* KO compared with WT myoblasts (R, S). Mander's overlap coefficient,

demonstrating increased co-occurrence of ER and mitochondria in *Opa1$^{fl/fl}$*-Cre (R) myoblasts

(S) and myotubes compared with WT. (T-Y) *In situ* proximity ligation assay (PLA) showing

MFN1–MFN2 interactions (red punctae) in *Opa1$^{fl/fl}$* and *Opa1$^{fl/fl}$*-Cre myoblasts (n = 10 cells, 3

independent experiments). Nuclei are labeled with 4′,6-diamidino-2-phenylindole (DAPI, blue).

(X) Quantification showing increased MFN1–MFN2 puncta in *Opa1* KO compared with WT

myoblasts. (AA-AD) Transmission electron microscopy (TEM) images of MERCs in primary

Opa1$^{fl/fl}$ and *Opa1$^{fl/fl}$*-Cre myotubes (Red arrows indicate mitochondria–ER interactions; n= 3 of

10 cells/experiment). Scale bar = 2 µm (2.5K magnification) and 500 nm (15K magnification).

(AE) Quantification of MERC distance in *Opa1$^{fl/fl}$* and *Opa1$^{fl/fl}$*-Cre myotubes. (AF, AG). (AF)

Percentage of mitochondrial surface area in direct contact with ER (AG) and percentage of ER

surface area in direct contact with mitochondria in *Opa1$^{fl/fl}$*-Cre compared with *Opa1$^{fl/fl}$*

myotubes. (AH) MERC area in *Opa1$^{fl/fl}$*-Cre compared with *Opa1$^{fl/fl}$* myoblasts and myotubes.

(AI-AL). Cristae number (AI), area (AJ), score (AK) and volume (AL) in *Opa1$^{fl/fl}$*-Cre compared

with *Opa1$^{fl/fl}$*. (AM-AP) The 3D distribution of single continuous and stationary mitochondria

(purple, purple arrows) and ER (blue, blue arrows), reconstructed from serial block face-

scanning electron microscopy (SBF-SEM) image stacks of *Opa1fl/fl*-Cre primary myotubes. (AQ-

AR). Percentage of ER surface area in contact with mitochondria (AQ) and the percentage of

mitochondria surface area in direct contact with ER (AR) in *Opa1fl/fl*-Cre compared with *Opa1fl/fl*

myotubes. SBF-SEM reconstructions from 7 to 23 fully constructed mitochondria, ER, or

MERCs. *Data are presented as the mean ± SEM. Significance was determined by Student's t-*

*test. *P < 0.05, **P < 0.01, ***P < 0.001, **** P < 0.0001.*

1144

Supplemental Figure 4. ATF4 binds to the promoters of genes encoding the IP_3R-GRP75-

VDAC MERC Complex. ChIP-seq analysis (represented in blue for the reverse strand and red

for the forward strand) from WT and *ATF4* deficient skeletal-muscle derived cells at (A) *Hspa9*

(GRP75), (B) *VDAC3*, and (C) *Itpr3* (IP_3R3) gene loci. Data obtained from a publicly available

database: https://chip-atlas.org. Data displays wild-type fibroblast and ATF4 knockout

fibroblasts, with ATF4 target genes binding peaks within 3kb from the Transcription Start Site of

the annotated gene. The mRNA-Seq data (yellow) reveals the gene expression levels, allowing

for gene expression relative to binding to be considered.

1153

Supplemental Figure 5. Loss of *Opa1* in myotubes and skeletal muscle increases Mitochondrial

Calcium Uniporter (MCU) and Sarcoplasmic/Endoplasmic Reticulum Ca^{2+}-ATPase (SERCA)

levels. (A). Western blot showing MCU and SERCA levels following OPA1 KO in myoblasts.

(B,C) Densitometry for MCU and SERCA respectively, normalized to alpha-tubulin. (D)

Western blot showing MCU and SERCA levels following OPA1 KO in myotubes. (E,F)

Densitometry for MCU and SERCA respectively. (G) Western blot showing MCU and SERCA

levels following OPA1 KO in mouse skeletal gastrocnemius muscle. (H,I) Densitometry for

MCU and SERCA respectively.

1162

1163   **Supplemental Table 1.** qPCR primers used for all studies involving mouse samples.

1164

1165   **Supplemental Table 2.** qPCR primers used for all studies involving *Drosophila*.

1166

1167   **Supplemental Videos 1–3.** Loss of *Opa1-like* in *Drosophila* skeletal muscle decreases muscle

1168   movement. Live videos of fly movement for WT (Video 1) and *Opa1-like* KD (Video 2 and 3)

1169   flies.

1170

1171   **Supplemental Videos 4-5.** Representative 360° rotational view of mitochondria–endoplasmic

1172   reticulum (ER) contacts (MERCs) in WT (Video 4) *Drosophila* and after *Opa1-like* knockdown

1173   (Video 5), captured with serial block face-scanning electron microscopy (SBF-SEM).

1174   Mitochondria are labeled blue, and ER are labeled pink.

1175

1176   **Supplementary Videos 6–7.** Representative 360° rotational view of confocal images in primary

1177   myoblasts after *OPA1* knockdown. Nuclei are labeled blue and MFN2–MFN1 interactions are

1178   labeled red. Supplemental Video 4 shows OPA1$^{fl/fl}$ while Supplemental Video 5 shows OPA-

1179   1$^{fl/fl}$Cre.

1180

1181   **Supplemental Videos 8–9.** Representative 360° rotational view of mitochondria–endoplasmic

1182   reticulum (ER) contacts (MERCs) in primary myotubes as WT (Video 8) and after *OPA1*

1183 knockdown (Video 9), captured with serial block face-scanning electron microscopy (SBF-

1184 SEM). Mitochondria are labeled blue, and ER are labeled pink.

1185

1186 **Supplementary Videos 10-13.** Representative 360° rotational views of confocal images of

1187 mouse skeletal muscle. Muscle fibers are labeled gray, and nuclei are labeled blue. Protein

1188 antibody interactions, MFN1–MFN2 and IP_3R3–VDAC interactions, are labeled in red. MFN1–

1189 MFN2 interactions are shown in WT (Video 10) and OPA1 smKO (Video 11). IP_3R3–VDAC

1190 interactions are shown in WT (Video 12) and OPA1 smKO (Video 13).

1191

1192 **Supplemental Videos 14-15.** Representative 360° rotational view of mitochondrial–endoplasmic

1193 reticulum (ER) contacts (MERCs) in mouse skeletal tissue as WT (Video 14) and after *OPA1*

1194 knockdown (Video 15), captured using serial block face-scanning electron microscopy (SBF-

1195 SEM). Mitochondria are labeled blue, and ER is labeled pink.

1196

1197 **Supplemental Videos 16–18.** Representative 360° rotational view of mitochondria–endoplasmic

1198 reticulum (ER) contacts (MERCs) in *Drosophila* flight muscle from control (Video 16), *ATF4*

1199 KO (Video 17), and *ATF4* overexpression (Video 18), captured with serial block face-scanning

1200 electron microscopy (SBF-SEM). Mitochondria are labeled blue, ER is labeled pink, and the

1201 nucleus is labeled yellow.

Figure 1

Figure 2

Figure 3

Figure 4