The *Aphelenchoides* genomes reveal major events of horizontal gene transfers in clade IV nematodes

Cheng-Kuo Lai[1,2], YiChien Lee[1,3,4], Huei-Mien Ke[1], Min R Lu[1], Wei-An Liu[1], Hsin-Han Lee[1], Yu-Ching Liu[1], Toyoshi Yoshiga[5], Taisei Kikuchi[6], Peichen J. Chen[7]* and Isheng Jason Tsai[1,2]*

Correspondence: janetchen@nchu.edu.tw and ijtsai@gate.sinica.edu.tw

[1]Biodiversity Research Center, Academia Sinica, Taipei 11529, Taiwan.
[2]Genome and Systems Biology Degree Program, National Taiwan University and Academia Sinica, Taipei, Taiwan.
[3]Biodiversity Program, Taiwan International Graduate Program, Academia Sinica and National Taiwan Normal University, Taipei, Taiwan
[4]Department of Life Science, National Taiwan Normal University, 116 Wenshan, Taipei, Taiwan
[5]Faculty of Agriculture, Saga University, Saga 840-8502 Japan
[6]Department of Integrated Biosciences, Graduate School of Frontier Sciences, The University of Tokyo, Chiba, 277-8562, Japan
[7]Department of Plant Pathology, National Chung Hsing University, Taichung, Taiwan

Abstract

Aphelenchoides besseyi is a plant-parasitic nematode (PPN) in the Aphelenchoididae family capable of infecting more than 200 plants. *A. besseyi* is also a species complex with strains exhibiting varying pathogenicity to plants. We present the genome and annotations of six *Aphelenchoides* species, four of which belonged to the *A. besseyi* species complex. Most *Aphelenchoides* have a genome size of 44.7-47.4 Mb and are amongst the smallest in the clade IV, with the exception of *A. fujianensis*, which has a size of 143.8 Mb and is the largest. Phylogenomic analysis successfully delimited the species complex into *A. oryzae* and *A. pseudobesseyi* and revealed a reduction of transposon elements in the last common ancestor of *Aphelenchoides*. Synteny analyses between reference genomes indicated that three chromosomes in *A. besseyi* were derived from fission and fusion events. A systematic identification of horizontal gene transfer (HGT) genes across 27 representative nematodes allowed us to identify two major episodes of acquisition corresponding to the last common ancestor of clade IV or major PPNs, respectively. These genes were mostly lost and differentially retained between clades or strains. Most HGT events were acquired from bacteria, followed by fungi, and also from plants which was especially prevalent in *Bursaphelenchus mucronatus*. Our results establish a comprehensive understanding on new origins of horizontal gene transfer in nematodes.

Introduction

The ability to parasitise plants has evolved in the phylum Nematoda on at least four occasions[1,2]. The major plant parasites belonged to the Aphelenchodidae and Parasitaphelenchidae families making up the Aphelenchoidea superfamily and the Tylenchida order of clade IV nematodes[3]; these plant parasitic nematodes (PPNs) collectively cause worldwide agriculture damages of over US$80 billion each year[4]. Root-knot nematodes in *Meloidogyne* genus cause the majority of these losses and were the first of PPNs to have their genomes sequenced[5], followed by pinewood nematode *Bursaphelenchus xylophilus*[6,7], potato cyst nematode *Globodera pallida*[8], soybean cyst nematode *Heterodera glycines*[9] and others[10,11]. Comparing these genomes yield insight into several adaptions that allow PPNs to parasitize plants. Examples include effectors such as carbohydrate active enzymes (CAZyme), which are known to be secreted by PPNs and are hypothesized to be involved in degrading or modifying the composition of different plant structural tissues[12,13]. Some of these PPN-specific genes are known to be acquired from bacteria or fungi through horizontal gene transfer (HGT)[14], giving nematodes the ability to adapt to different environments[14]. Although numerous HGT genes have been identified and documented in different nematodes, research on the timing and subsequent maintenance of these genes, and why their copy numbers differ, has been restricted to a few PPN clades[15].

Currently, the only major groups containing plant parasitic nematodes that lack a reference genome are Trichodoridae and Aphelenchoididae. Of particular interest is the *Aphelenchoides besseyi*, which is a foliar nematode that infects almost 200 plants in 35 genera[16]. This nematode is 10mm in body size, has a life cycle of this nematode is around 10 to 12 days and it can reproduce in extreme environments, making it hard to eliminate. Better known as the rice white tip, *A. besseyi* infects important agronomic crops such as rice, soybeans and strawberries[17,18], causing necrosis and distortion of its host's leaves[16,18,19]. The nematode has reportedly been responsible for up to a 60% crop loss in some cases[20,21] and was listed among the top ten plant parasitic nematodes in a recent review[22]. Despite the economic damage these parasitic

nematodes inflict, particularly in the Asian region, little is known about the basic biology,

genetic diversity or evolution of *A. besseyi* and other Aphelenchoididae members. It has

been reported that *A. besseyi* isolated from different hosts have different levels of

pathogenicity. For instance, the populations of *A. besseyi* isolated from strawberries

were unable to parasitise rice[21]. However, the populations of this species from bird's-

nest fern can reproduce in both rice and strawberries[19]. Despite their almost overlapping

morphological features, we previously identified copy number variations of genes

encoding cell-wall-degrading enzymes including glycosyl hydrolase family 5 (GH5) and

GH45 cellulases between *A. besseyi* of different host origins[23]. An 18S phylogeny

separated the strains isolated from rice and fern unambiguously, suggesting that *A.*

besseyi might be a species complex, with literatures also identifying variations in

different molecular markers in different hosts that are original to *A. besseyi*[18,24].

Subbotin *et al*. recently used a combination of molecule makers (28S, ITS and

mitochondria *COI* gene)[17] to reclassify foliar nematodes into three separated clades: the

A. besseyi isolated mainly from strawberries, the *A. oryzae* mainly isolated from rice and

A. pseudobesseyi from wood fern suggesting this species complex may be well

differentiated at the genome level. From an evolutionary perspective, *A. besseyi* is also

interesting because its primitive plant parasitism was a relatively recent evolutionary

adaptation[25].

In this study, we sequenced and annotated the genomes of four *A. besseyi*

species complex strains isolated from different plants which we later designated as *A.*

pseudobesseyi and *A. oryzae,* and another two species in the Aphelenchoididae family

(*Aphelenchoides bicaudatus* and *Aphelenchoides fujianensis*). We compared the

proteomes of six Aphelenchoididae members with 21 other representative nematodes to

delimit species relationships and investigated their gene family dynamics. We identified

synteny with representative nematodes and inferred rearrangement events to determine

how the three chromosomes of *A. besseyi* was evolved. The availability of the

Aphelenchoides assemblies allowed us to systematically determine the horizontal gene

transfer-acquired genes in nematode genomes. By inferring the evolutionary origins of

these HGT genes we found historical HGT events that shaped nematode evolution.

The major event occurred in the last common ancestor of clade IV nematodes and may
have contributed to the early adaptation of these nematodes.

Results

Genome assemblies and annotations of six *Aphelenchoides* species

We sequenced and assembled the genomes of six nematodes in the

Aphelenchoides genus (four *A. besseyi*, one *A. bicaudatus* and one *A. fujianensis*).

These species were chosen to represent the Aphelenchoididae family and *A. besseyi*

strains isolated from three plant hosts (**supplementary table S1**) to delimit their

relationship within the species complex. For each species, an initial assembly was

produced from either 70-148X Oxford Nanopore or 113-422X Pacbio reads using Flye

assembler[26] and further polished using Illumina reads (**supplementary table S2**).

Among *A. besseyi* assemblies, the VT strain isolated from tape grass *Vallisneria spiralis*

had the highest genome quality with N50 5.4 Mb (hereafter denoted as APVT). The

contigs of this strain were further scaffolded with 150X Hi-C reads using the Juicer

program[27] (**supplementary fig. S1**) yielding a final assembly of 44.7 Mb (N50 = 16.9

Mb). More than 99% of this assembly was in three scaffolds, presumably corresponding

to three chromosomes[28] (2n=6). Five *Aphelenchoides* assemblies ranged from 44.7 to

47.4 Mb (N50 = 12.2-17.8 Mb; **supplementary table S3**), and a sixth assembly (*A.

fujianensis*) of 143.8 Mb (N50 = 553 kb; **supplementary table S3**) was estimated to be

triploid[29] (**and supplementary fig. S2**). Although not present in the assemblies, the

telomere motif TTAGGC was identified in the reads of *A. peseudobesseyi* at low

coverage (**Supplementary Info**), which is consistent to the sister group species of *B.

xylophilus* (**supplementary table S4**) indicating the presence of telomeres in these

species.

Using the proteomes of *Bursaphelenchus xylophilus* and *Caenorhabditis

elegans*, and the transcriptomes of pooled worms in each species as evidences, we

predicted 11,701 to 12,948 protein coding genes in six *Aphelenchoides* species with

Maker2 pipeline[30] (**supplementary table S3**). With the exception of *A. fujianensis*,

these were fewer protein coding genes in these species than in Tylenchida nematodes

(12,762 to 19,212) and free-living nematodes (20,184 to 20,992). The completeness of annotated genes was estimated to be 76.4–81.3% based on a BUSCO assessment, lower than that of *Bursaphelenchus* species (83.0–89.4%), but are higher than that of Tylenchida (59.8–73.8%) nematodes. The lower BUSCO completeness in *Aphelenchoides* was likely clade specificm as re-annotation of the APVT strain with trained models based on manual curation of 975 genes also gave a similar score of 78.2%. Among them, 66.5% to 71.0% of genes in *Aphelenchoides* could be assigned at least a domain from the Protein family (Pfam) database[31]. In addition, orthologous groups were inferred with the proteomes of six *Aphelenchoides* with 21 other nematodes using Orthofinder[32]. With the exception of *A. fujianensis*, 78.5–85.4% (*A. fujianensis* = 48.7%), 69.4–76.9% (*A. fujianensis* = 42.8%) and 87.5–98.7% of *Aphelenchoides* genes were orthologous to *B. xylophilus*, *C. elegan*s and at least one other nematode species, respectively, suggesting that the reduced proteome in most *Aphelenchoides* was mainly comprised of conserved genes among nematodes.

Phylogenomics delimit species complex of *Aphelenchoides besseyi*

To investigate the evolution of plant-parasitic nematodes and the relationships among members in the *A. besseyi* species complex, a maximum-likelihood phylogenetic tree was constructed based on 74 low-copy orthologues. The phylogeny is consistent with a the previous study[33]: the major plant parasitic nematodes were divided into Aphelenchoidea and Tylenchida, and six *Aphelenchoides* species were grouped as sister to *Bursaphelenchus* (**fig. 1a**). The *A. besseyi* strains were clustered into two groups based on their hosts, suggesting that relationships in these species within the *A. besseyi* species complex can be unambiguously resolved based on their different lifestyles and host preferences. Combined with the previous 28S phylogeny of the *A. besseyi* species complex[17] (**supplementary fig. S3**), we further designated these two groups as *A. oryzae* and *A. pseudobesseyi* groups isolated from rice or other plants (land grass and bird's-nest fern). The median nucleotide and amino acid identity was 86.6% and 90% between these two groups, respectively (**supplementary fig. S4**). Strains in each group also differed in heterozygosity (0.017-0.019% in *A. oryzae* vs 0.071-0.075% in *A. pseudobesseyi*) and changes in recent effective population sizes

inferred using pairwise sequentially Markovian coalescent (PSMC) analysis[34]

(**supplementary fig. 5**). Together these results emphasised that relationships among

species in the *A. besseyi* species complex were highly diversified at the genome level

despite being challenging to differentiate based solely on morphology[17].

Genome reduction as a result of transposable element loss

The *Aphelenchoides* genomes were smaller than those of other plant-parasitic

nematodes (**fig. 1b** and **supplementary table S3**), indicating that genome reduction

took place in the last common ancestor of the *Aphelenchoides* genus. Much of the

reduction can be explained by the reduced markup of repeat content compared to other

nematodes (**fig. 1b**). The dominant transposable elements of *Aphelenchoides* were

DNA transposons—which were reduced in content (0.14–1.36 Mb vs. 4.2–22.1 Mb)—

and number of families (1–7 in *Aphelenchoides* compared to 9 and 26 in *B. xylophilus*

and *H. glycines*, respectively) compared to other nematodes. Fewer LTR (0.07–0.8 Mb

vs. 0.24–9.3 Mb) and LINE (0.0006–0.66 Mb vs 0.02–4.5 Mb) retrotransposons were

also observed in this genus. These results suggest that the reduced genome sizes in

Aphelenchoides might have been caused by the rapid loss of transposable elements

and led to the eventual loss of entire families in some cases (**fig. 2a**). Within the *A.*

besseyi species complex, *A. peseudobesseyi* contained significantly fewer DNA

transposons, LTR and LINE retrotransposons than *A. oryzae* (**fig. 2a and**

supplementary fig. 6).

Figure 1 Phylogenomic analysis of plant-parasitic nematodes. a. The phylogeny of 27
representative nematodes was inferred based on the concatenation of protein sequences from
74 low-copy orthologues. The blue dots in branches denote a bootstrap value of 100. **b.** The
size of the genome features in nematodes, repeats containing DNA transposons (DNA), long
interspersed nuclear elements (LINE), long terminal repeats (LTR), rolling-circular (RC) and the
unclassified repeats are labelled as "other repeats". Numbers in brackets denote genome
assembly sizes in megabases.

Figure 2 Repeat and proteome contents in nematodes. a. The genome proportions of DNA,

LINE and LTR transposable elements in nematodes shown by genome percentage **b-c.** Protein

families and CAZyme gene copy numbers vary significantly among nematodes. The dot size

represents the copy number of each domain and the different colour represents the copy

number of domains larger or lower than average copies shown in brackets.

Gene family specialization in the *Aphelenchoides* species

We observed 66 enriched and 31 reduced protein domains in the four member of

the *A. besseyi* species complex compared to 21 other nematodes. (**fig. 2b** and

supplementary table S5). Domain reduction included collagen (90–109 copies in the

A. besseyi species complex vs. 72–407 in others), Somatomedin B and BTG. Genes

containing collagen domains were reportedly associated with capsule formation; the

reduced copy of collagen domains in *Trichinella spiralis* were thought to contribute its

lower host-specificity than other nematodes[35], and may be related to the wide host

range of *A. besseyi*. In contrast, Aphelenchoidea members possess on average four-

fold (91–314 vs. 4–555 copies) more aspartic proteases (ASP) than other nematodes

(**supplementary table S5**). ASPs were reported to associate with the digestion of host

haemoglobin in *Haemonchus controtus*[36], and also skin penetration in hookworms[37],

and may play an important role in the *Aphelenchoides*'s parasitism process. Other expansions included LIM and peptidase C13 domains, which participate in participating in the regulation of cell motility and cell growth[38] or degradation of protein tissues in a host[39], emphasizing that these domain dynamics are associated with adaptations to plant parasitism.

The plant cell wall acts as a primary defensive barrier and the production of carbohydrate-active enzyme (CAZyme) families are important for PPNs to infect plants. A total of 132 CAZyme families were identified in the representative the 27 nematodes. Of these, 59–67% of the CAZyme families were observed in Aphelenchoidea which is similar to the 55–66% and 58-68% of the families in Tylenchida and free-living nematodes (**supplementary table S6**), respectively. A total of 13 families were significantly expanded or lost in the *Aphelenchoides* genus (**fig. 2c**), including GH16, GH27 and GH45. GH16 serves as the putative β-glycanases involved in the degradation or remodelling of cell wall polysaccharides[40], GH16 had one to six copies in Aphelenchoididae nematodes and was not identified outside this clade except in *D. destructor*, in which there were three copies. There are three to 11 copies of GH27— which are reportedly involved in the function of hemicellulose and associated with α-galactosidase activity in both bacteria and fungi[41]— in Aphelenchoidea, but fewer in the Tylenchida nematodes. The previously identified GH45 present in Aphelenchoidea nematodes[23]—involved in the degradation of beta-1,4-glucans in the plant cell wall[19]— possess different copy numbers between *A. pseudobesseyi* and *A. oryzae* and were absent in *A. fujianensis* and *A. bicaudatus,* suggesting differential maintenance of these genes in the same genus may have contributed to variations of pathogenicity to plants.

Chromosome evolution of PPNs

To investigate the extent of the karyotype rearrangements in *Aphelenchoides*, we inferred the synteny relationships among *A. peseudobesseyi* (chromosome n=3)[28], *B. xylophilus* (n=6) and *C. elegans* (n=6) using single copy orthologs. Within the three *A. peseudobesseyi* chromosomes, orthologs belonging to all *C. elegans* chromosomes were clustered into distinctive blocks (**fig. 3a**) suggesting a fusion of ancestral chromosomes. These regions remained contiguous and contained 148-801 orthologous

genes that could be assigned from individual chromosomes presumably not yet broken down yet by recombination, allowing us to pinpoint the fusion points and infer the order of rearrangement events based the constitution of chromosomes (**fig. 3b**). We encountered instances of where an ancestral chromosome was found in different parts of the *A. peseudobesseyi* chromosomes, suggesting fission also took place. In the case of chr IV—which remained homologous in *C. elegans* and *B. xylophilus*—corresponding synteny blocks in *A. peseudobesseyi* were identified in the arm of chr 2 and chr 1 separated by regions of chr III origin (**fig. 3b**). The majority of the ancestral sex chromosomes were unambiguously assigned to chr 2, and remapping of male sequences showed equal coverage along the chromosomes (**supplementary figure S7**), suggesting that the Aphelenchoidea superfamily including *A. besseyi* exhibited a stochastic sex determination system that was recently characterized in *B. xylophilus*[42]. Within the *A. besseyi* species complex, a total of 91% and 88% of genomes were in synteny between APVT and AORT, respectively. Intra-chromosomal inversions were common at chromosome arms. In addition, we identified a major inversion of length 3.4 Mb long located in the centre of chr 2 (**fig. 3c**) suggesting rearrangement is still ongoing. Both the LTR and LINE retrotransposons were enriched in the chromosome arms of the *A. oryzae* strain (AORT) (**fig. 3c and supplementary fig. S6**), which is consistent with the hallmark of nematode chromosome evolution[43]. In contrast, only the LTR retrotransposons were found in the two chromosome arms of *A. peseudobesseyi*, suggesting that these repeats were differentially maintained after speciation.

Figure 3. Chromosome evolution of plant-parasitic nematodes. a. The density of pairwise

single-copy orthologs between *A. pseudobesseyi* VT and *B. xylophilus*. Colours denote *B.*

xylophilus chromosomes and the putative chromosome fusion sites in APVT are labelled with

red triangles. **b.** Colours denote corresponding *C. elegans* chromosomes, and dashed lines

indicate linkage groups corresponding to ancestral chromosomes. **c.** The synteny relationship

and the distribution of transposable between *A. pseudobesseyi* VT and *A. oryzae* RT. Blocks

indicate synteny links between the two strains, and the line colours correspond to inversion

(blue) and inter-chromosomal rearrangement (grey). Distribution of DNA transposons (red), long

interspersed elements (green) and long terminal repeats (blue) between two stains are shown.

Major episode of HGT in clade IV nematodes

In plant parasitic nematodes, the GH5 cellulase was found present in Tylenchida and only *A. peseudobesseyi* and *A. bicaudatus* within the Aphelenchoidea clade[23,44], raising the possibility that many of the horizontal gene transferred genes were acquired in the last common ancestor of major PPNs but were differentially lost. To identify such events, a total of 27 proteomes from representative nematodes including the *Aphelenchoides* genomes were searched for evidence of HGT by calculating the Alien Index (AI) score using Alienness[45]. We identified a total of 1,675 HGT orthogroups in 21 nematodes. Placing these orthologs designated as events onto the species phylogeny assuming a parsimonious scenario[46], indicated that HGT started in the last common ancestor of clade IV nematodes (**fig 4a**). Examples include GH16, GH32, GH43 and the aforementioned GH5 cellulases. We inferred a total of 161 orthogroups were acquired in this episode, and most of their origins were inferred to be bacteria (78.3%) (**supplementary table S7**) belonging to different genera, suggesting multiple acquisitions took place. Of these, we found 36 Pfam terms such as ABC transporter that were identified in multiple orthogroups suggesting some convergence in the acquired functions (**supplementary table S8**).

The revised GH5 cellulase phylogeny indicated an ancient duplication took place before the divergence of PPNs (**fig 5a**). One clade contains orthologs of the three *Panagrolaimid* (*P.* sp. PS1159, *P. superbus* and *P. davidi*), and Tylenchida, and the other clade contains members of Aphelenchoidea and Tylenchida nematodes, which emphasises that the fate of the HGT genes was governed by duplications and loss. Interestingly, the closest GH5 bacterial orthologs were *Salinimicrobium xinjiangense* and *Leeuwenhoekiella sp.*, which belonged to Flavobacteriaceae family and were from marine environments. We observed two GH16 subfamilies in nematodes. GH16_3 in Tylenchida and *Bursaphelenchida* nematodes were clustered with bacterial origin sequences, whereas GH16_1 of *Aphelenchoides* and *Panagrolaimus* nematodes were clustered with fungal origin (**fig 5b**), suggesting that the two GH16 groups arose independently. GH32 in *G. pallida*[13] is believed to play a role in the function of fructose hydrolysis and was found in one *Panagrolaimus* in addition to several Tylenchida nematodes (**supplementary fig. S8**). GH43 was identified at two distinct clusters of

bacterial origin in Tylenchida and *Panagrolaimid* nematodes which have been proposed to be involved in degradation of the hemicellulose in plants[47] (**supplementary fig. S9**).

The next major episode of acquisition took place in the common ancestor of PPNs, with 47 orthogroups (**fig. 4a**). These families included pectate lyases 3 (PL3) which is associated with cell wall degradation[48]. The orthologs of PL3 in *Aphelenchus avenae* and two *Bursaphelenchus* nematodes were grouped together with distinct clusters of *Meloidogyne* species (**fig. 5c**) is consistent with previous phylogeny finding in PPNs[44]. The closest bacterial ortholog in the *Meloidogyne* clade was from *Curtobacterium flaccmfaciens* which is also known to cause bacterial wilt in the Fabaceae family[49]. Together, these results suggested some genes that were thought to play important roles in plant parasitism were in fact acquired earlier than the common ancestor of plant parasitic nematodes.

The majority of HGT gene families were of bacterial followed by fungal origin (**fig 4b**). We also identified genes that were acquired from non-bacterial donors in the last common ancestors of clade IV, as well as in more recent, different PPN lineages (**fig 4a**). This included the previously characterised fungal origin of GH45[23,50], This cellulase family is present in most Aphelenchoidea nematodes except *A. fujinensis* and *A. bicaudatus*. The GH16 family was independently acquired from a bacterial and fungal donor in the last common ancestor of clade IV nematodes and the *Aphelenchoides* genus, respectively (**fig 5b**). Notably, we identified 40 orthogroups among PPNs that were transferred from the plant phylum Streptophyta, which is consistent with the finding of several sequences that are highly similar to plants in *H. glycine*[51] (**fig. 4b**). The closest plant orthologs included rice, maple and oak (**fig. 5d**) which are common hosts to many PPNs. Strikingly, of these orthogroups, 27 were present in *B. mucronatus* and enriched in the detoxification of cadmium and copper ion function (**supplementary table S9**), suggesting these genes may help *Bursaphelenchus* nematodes to degrade the toxin in pine wood hosts.

Figure 4 The evolution of genes acquired from horizontal gene transfer (HGT) in clade IV nematodes. a. HGT orthogroups were inferred by the AI score[52] > 0 across 27 representative nematodes; the HGT families gained or lost are shown in the branches. Horizontally acquired CAZymes are annotated. The proportions of donor origins in each HGT orthogroup belonging to different kingdoms of donors are shown as pie charts. The size of the pie chart corresponds to the total number of HGT orthologues in branches; the chart was normalized using: log_5(total number). **b.** The distribution of HGT families transferred from different kingdoms, with the same denoted color scheme as same as figure a. **c.** Number of HGT events among different inferred time points corresponding with the branches which are triangle marked in the phylogeny.

Figure 5 Phylogenies of nematode HGT orthogroups and their donor origin. a. GH5_2 **b.**
GH16 **c.** PL3 **d.** OG0003777 and OG0004537. Different colours denote different kingdoms and
species as shown in legend. Nematode gene copies with negative AI values were marked with
an asterisk. Additional Pfam domains are labelled when available. Nodes with iqtree UFBoot
and SH-aLRT bootstrap support > 80% are labelled as grey circles.

We identified 0.3-2.4%, 0.6-2.1% and 0.1-5.4% proteomes among

Aphelenchoidea, Tylenchida and *Panagrolaimomorpha* nematodes that were predicted

to be HGT (**fig. 4c**). The majority of these differences were the result of clade-specific

evolution after speciation. The high copy number of HGT genes observed in *M.*

incognita was a result of duplication[53], indicated by the fact that the number of HGT

orthologs of bacteria origin were over two times higher than any other species

(**supplementary fig. S10**). The high number of HGT genes in *P. superbus* was

consistent with a previous study[54] and likely to be species specific.

To independently assess the accuracy of our approach and interrogate the fate

of HGT genes, we constructed a phylogeny for every orthogroup containing identified

HGT candidates. Members of Aphelenchoididae and Tylenchida orthologs in the

majority of these orthogroups were predicted to be all HGT genes (with AI > 0; 54.6-

76.5% vs. 77.3-89.4%). Genes from a species were typically grouped together in the

orthogroup phylogeny regardless of being identified as HGT candidates, suggesting the

genes that were not detected using our threshold shared common ancestries with those

that were. Presumably, this was a result of accumulating substitutions over time.

Consistent with this observation, the more ancient acquired HGT orthogroups in PPNs

contained higher copy number of these genes compared to recently acquired families

(**supplementary fig. S11**). The instances included GH5 families with 12.5-70.6% of

copies in Tylenchida were failed to identify as HGT candidates, suggesting duplication

and possibly neo-functionalisation of the GH family in PPNs after being acquired from

bacterial origin (**fig. 5a**). The differentiation was ongoing and observed in the *A. besseyi*

species complex, which included the GH45 orthogroup with negative AI in two *A. oryzae*

strains (**supplementary fig. S12**).

Discussion

Characterising the diversity and comparing the genomes of plant parasitic

nematodes has been of fundamental importance in understanding how such lifestyles

arise and of practical importance in identifying candidate effectors and control methods.

The latter has been addressed in several studies, focusing mainly on *Meloidogyne*[55].

The *Aphelenchoides* genome assemblies presented in this study allowed us to gain a

holistic view of the evolution of clade IV nematodes, which appeared to gain and lose

many adaptations, including plant parasitism[56]. In their evolution, HGT genes have

played important roles in functions related to these adaptations. The most recent

comprehensive analyses of HGT in nematodes focused on plant parasitic nematodes[15]

and found many of these genes were PPN specific. Of these, donors of gene families

involved in plant cell-wall modifications were previously found to be associated with

plants which was appealing that were sympatric with plant parasitic nematodes which

made HGT possible[44]. Additional HGT events were identified in other clade IV

nematodes[13,54,57–59] but were part of analysing new *de novo* genomes.

Our systematic investigation of HGT has instead shown that many of the

aforementioned families were acquired much earlier in the last common ancestor of

clade IV. Many clade IV nematodes are known to survive extreme desiccation[54,59,60] and

the acquired HGT genes may be central to their resistance in harsh environment and

subsequently catalysis to successful plant parasitism[61]. Sources of these donors may

be the symbionts like the case of insects[62], but currently nematode endosymbionts are

restricted to *Wolbachia* and *Cardinium*[63] and were not identified in our analyses.

Interestingly, many of the closest bacterial donors were from marine environments,

raising the possibility that the last common ancestor of clade IV may have lived in a

marine environment that underwent habitat transition[64]. However, we also identified

donors of non-bacterial origin that were usually found in the environments that fit

nematodes' present day lifestyle. Now that more genome sequences are available,

historical HGT events were detected in the most recent common ancestor of major

organism groups such as land plants[65] , of moths and butterflies[66], which contributed

the hosts' developmental roles and adaptations. These acquisitions were found to be

episodic and likely took place in a time when either the host development or genome

defence was vulnerable. We speculate that the gain and absence of gene families in

clade IV nematodes may have played a role in retention of HGT genes.

The successful delimitation of the *A. besseyi* species complex unambiguously

into *A. oryzae* and the recently proposed *A. pseudobesseyi* has important implications

in nematode management. Congruent delimitation was observed between genomes

and 28S phylogenies confirming the utility of species identification with existing

molecular markers[18]. *A. besseyi* is generally believed to have limited mobility in natural

habitats, so its lack of population structure in China[24] was suggested as a consequence

of human-mediated dispersal. Our results also supported that *A. oryzae* appears to be

more rice plant specific compared to *A. pseudobesseyi* which was isolated more

frequently in ornamental plants and other agronomic crops[18]. A comprehensive

collection across a wider geographical range and resequencing of strains previously

designated as *A. besseyi* could confirm whether *A. oryzae* was responsible for all the

white tip disease in rice plants and may lead to better characterisations of the

biogeography and evolution of different cryptic species.

The reduction of genome size and reduced chromosome numbers of *A. besseyi*

represent an interesting outcome for the typical six nematode ancestral chromosomes

around hundred megabases in length. Genome rearrangement and reduction are

common across the tree of life including plants[67], butterflies[68] and nematodes[69]. We

show that *A. besseyi* underwent multiple chromosome fission and fusion events, and a

possible explanation together with genome reduction may be the missing of meiosis

genes and the telomeric repeat maintenance genes, which resulted in truncated meiosis

(**supplementary table S10**); this was observed in an extreme case of *Diploscapter*

pachys[70] possessing a single chromosome. Alterations in meiosis may lead to genome

shrinkage due to a loss of transposable elements as a result of imbalanced chromatin

as observed in *Caenorhabditis nigoni*[71]. It is likely that *Aphelenchoides* underwent a

similar scenario. However, members of *Bursaphelenchus* with six chromosomes also

failed to identify these aforementioned orthologs (**supplementary table S10**),

suggesting their divergence has taken place since the last common ancestor of

Aphelenchoidea. Further cell and molecular evidence are needed to confirm the

integrity of meiosis in *A. besseyi*.

To conclude, the availability of the *Aphelenchoides* genome and our comparative

analyses allowed us to pinpoint the major events of horizontal gene transfer in clade IV

nematodes. The results have reinforced the importance of horizontal gene transfers

contributing to multiple adaptations of these nematodes including plant parasitism. In

addition, the various *A. besseyi* genomes will assist in developing molecular diagnostic

tools to distinguish the specific diseases caused by the species complex.

Methods

DNA, RNA extraction and sequencing

Nematodes were cultured with *Alternaria citri* on PDA (potato dextrose agar) medium. All stages of nematodes were collected from the medium, washed with sterile distilled water, and purified by sucrose gradients. Genomic DNA was extracted using Qiagen Genomic-tip 100/G according to the manufacturer's instructions, RNA extraction was conducted using Trizol, and then purified using a lithium chloride purification method. The DNA paired-end libraries were constructed using either a Nextera DNA Flex or KAPA hyper library prep kit (Illumina, San Diego, USA); the RNA paired-end libraries were constructed using a TruSeq Stranded mRNA library prep kit (Illumina, San Diego, USA). Both DNA and RNA pair-end followed with standard protocol and were sequenced by Illumina HiSeq 2500 (Illumina, USA) to produce 150-bp paired-end reads. The HiC library preparation was performed by Phase Genomics (Seattle, WA, USA) proximo HiC animal protocol with some modification in tissue processing. The enriched worms were finely chopped by microtube pellet pestle rods for about 2 minutes. The tissues were crosslinked by adding 1 ml crosslinking solution and incubate for 25 minutes with occasional mixing by rotation. 100 ul quenching solution was added to the crosslinked tissue and mixed for 20 minutes by rotation. The rest of the preparation steps follow the protocol. The library was sequenced by Illumina HiSeq 2500 (Illumina, USA) to produce 150-bp paired-end reads. APFT and AORT were using Pacbio sequencing system to produce long-read, and the rest of 4 *Aphelenchoides* strains (APVT, AORJ, *A. bicaudatus*, *A. fujianensis*) were sequenced using the Oxford Nanopore sequencing platform. The raw nanopore signals were basecalled by Guppy[72] (ver 0.5.1) producing a total of 5.0-28.4 Gb sequences at least 1 kb in length.

Assemblies of six *Aphelenchoides* spp.

Raw reads of each species were assembled using Flye (ver 2.8.2)[26] assembler. The assemblies from Nanopore reads were corrected using Nanopore reads using Racon[73] (ver 1.4.6) and Medaka[74] (ver 0.10.0). All assemblies were further corrected using Illumina reads using Pilon[75] (ver 1.22) with five iterations. The *A. pesudobesseyi* VT assembly was scaffolded using HiC reads and subsequently curated in Juice-box[27]

tools. The other five *Aphelenchoides* genomes were reference scaffolded based on this

assembly using Ragtag[76] (ver 1.1).

Gene prediction and functional annotation

The identification of repetitive elements were computed by RepeatModeler[77] (ver

1.0.8), TransposonPSI (ver 1.0.0; https://github.com/NBISweden/TransposonPSI) and

USEARCH[78] (ver 8.1) based on the protocol by Berriman *et al*.[79]. Repeat locations were

then identified by Repeatmasking[80] (ver 4.0.9). RNA-seq reads of six *Aphelenchoides*

strains were trimmed by Trimmomatic[81] (ver 0.36), and aligned to corresponding

assemblies using STAR[82] (ver 2.7.1a). From these mappings, transcripts were inferred

using three approaches: i) assembled based on the mappings as guides using Trinity[83]

(ver 2.84; option: default setting), reconstructed using ii) Stringtie[84] (ver 1.3.4; option:

default setting) and iii) CLASS2[85] (ver 2.17; option: default setting). Transcripts

generated from Trinity were realigned to the reference using GMAP[86] (ver 2017-11-15).

The RNAseq mappings were also used in BRAKER[87] to train species parameter and

generate an initial set of annotations. Proteomes of *Bursaphelenchus xylophilus* and

Caenorhabditis elegans were downloaded from Wormbase (WBPS14;

https://wormbase.org) and used as homology guides to pick the best transcripts for

each putative locus using MIKADO[88] (ver 1.2.4; option: three Mikado steps, containing

"prepare", "serialize" and "pick" procedures), and were also used to train MAKER2.

Finally, MAKER2 was invoked to generate a final set of gene annotations using picked

EST evidence and protein evidence from MIKADO transcript and proteomes from

closely related species (*Bursaphelenchus xylophilus* and *Caenorhabditis elegans*), and

used gene models (BUSCO[89], BRAKER, SNAP[90] and Augustus[91]) as EST hints to train

predicted data with three iterations.

Comparative analyses

Proteomes of five plant-parasitic nematodes (*Bursaphelenchus xylophilus,*

Meloidogyne hapla, Meloidogyne incognita, Globodera pallida, Ditylenchus destructor),

two free-living nematodes (*Caenorhabditis elegans, Caenorhabditis briggsae*), six

Panagrolaimomorpha (*Propanagrolaimus* sp. JU765, *Panagrellus revidius,*

Panagrolaimus superbus, Panagrolaimus sp. PS1159*, Panagrolaimus davidi* and

Halicephalobus mephisto) and one animal parasitic nematode (*Brugia malayi*) were

downloaded from Wormbase[92,93](ver 14). Orthogroups were determined by

Orthofinder[32] (ver 2.2.7; options: -S diamond). Sequence alignments of each of the

single-copy orthogroups were generated by MAFFT (ver 7.310; options: --maxterate

1000). Then, the concatenated alignment of all single-copy orthogroups was used to

compute a maximum likelihood phylogeny using RAxML[94] (ver 8.2.3; options: -s -T 32 -

N 100 -f a -m PROTGAMMILGF) with 100 bootstrap replicates. Pfam copy numbers of

all 27 nematodes were identified from the results of nematode proteomes blast against

the database of Pfam website (ver 31; https://pfam.xfam.org/) using HMMER engine

with e-value smaller than 0.001. Effector enzymes were identified by searching the

nematode proteomes against the CAZyme[95] database (http://www.cazy.org) using

HMMER engine with sequence length larger than 80bp. The identified sequence was at

least larger than 0.35 proportion of conserved domain from database and had an e-

value smaller than 1e-15.

Identification of the HGT genes

The probability of genes having been acquired via HGT was estimated by using

Alienness Index (AI)[45]. Our donor group were generated by non-Metazoans from NCBI

nr database, and the recipient were Metazoans excluding the following species to

prevent self-alignment: Aphelenchoidea, Tylenchida, Rhabditina, Spirurina and

Cephaloboidea. The Alien Index (AI) was estimated by calculating the e-value of

diamond[96] (ver 2.0.14; option: blastx --evalue 0.001) best hits between the donor and

recipient database. Orthogroups having at least one gene with an AI value over 30 were

selected for further analysis. Gains and losses at each node were inferred using Phylip-

Dollop[46] (ver 3.69; options: fdollop -method d -ancseq). Some of the HGT family

acquired branches were manually curated by their evolutionary place of gene phylogeny

due to the fact that nematode genes with AI < 0 were clustered with other HGT genes.

The highest AI value of nematode genes with classified taxonomy hit were chosen to

represent the HGT origin in each orthogroup. Orthogroups with the same CAZyme

annotated and nematode orthology gene AI higher than -50 in those Orthogroups were

selected. AI < 0 genes were labelled with "*". The orthologs were further combined with the HGT identified donor sequence from nr database and the specific cellulase sequence from CAZyme database. To reduce contamination, orthologs of Pfam domain were annotated and filtered by having at least one major domain (cellulase or pectate lyase). Sequences of each HGT orthogroup were aligned using MAFFT (options: --maxiterate 1000 --genafpair) and trimmed by using trimAl[97] (ver 1.4; options: -gappyout). The ortholog phylogeny were computed by using IQtree[98] (ver 1.6.6; options: -bb 1000 -alrt 1000). For the CAZyme unclassified HGT orthogroups, the top 2 blast hits sequences from separated Uniprot (bacteria, fungi, land plants and insect) were used to confirm the HGT origin.

AUTHORS CONTRIBUTION

IJT and PJC conceived the study. IJT led the study. YCL, TY and PJC sampled the *Aphelenchoides* nematodes. YiCL, HMK, WAL conducted the experiments. CKL analysed the data with input from HHL, YuCL and MRL. IJT and CKL wrote the manuscript with input from TY, TK, PJC

DATA AVAILABILITY

The sequencing data and annotation of six *Aphelenchoides* nematodes are publicly available in NCBI with the BioProject accession PRJNA834627 and scheduled in the next release WBPS18 of WormBase Parasite. The accession numbers of individual samples are listed in **supplementary table S2**. The information of Clade IV acquired HGT orthogroups could be found in github (https://github.com/lihowfun/CladeIV_HGT.git).

Funding

This work was supported by Academia Sinica grant (AS-CDA-107-L01) to IJT.

References

1.   Blaxter, M. L. *et al.* A molecular evolutionary framework for the phylum Nematoda. *Nat. 1998 3926671* **392**, 71–75 (1998).

2.   Van Megen, H. *et al.* A phylogenetic tree of nematodes based on about 1200 full-length

small subunit ribosomal DNA sequences. *Nematology* **11**, 927–950 (2009).

3.   Bird, D. M. K., Jones, J. T., Opperman, C. H., Kikuchi, T. & Danchin, E. G. J. Signatures

of adaptation to plant parasitism in nematode genomes. *Parasitology* **142**, S71 (2015).

4.   Nicol, J. M. *et al.* Current Nematode Threats to World Agriculture. *Genomics Mol. Genet.*

Plant-Nematode Interact. 21–43 (2011). doi:10.1007/978-94-007-0434-3_2

5.   Abad, P. *et al.* Genome sequence of the metazoan plant-parasitic nematode Meloidogyne

incognita. *Nat. Biotechnol. 2008 268* **26**, 909–915 (2008).

6.   Kikuchi, T., Cotton, J. A., Dalzell, J. J., Hasegawa, K. & Kanzaki, N. Genomic Insights

into the Origin of Parasitism in the Emerging Plant Pathogen Bursaphelenchus xylophilus

Genomic Insights into the Origin of Parasitism in the Emerging Plant Pathogen. (2011).

doi:10.1371/journal.ppat.1002219

7.   Dayi, M. *et al.* Nearly Complete Genome Assembly of the Pinewood Nematode

Bursaphelenchus xylophilus Strain Ka4C1 . *Microbiol. Resour. Announc.* **9**, (2020).

8.   Cotton, J. A. *et al.* The genome and life-stage specific transcriptomes of Globodera

pallidaelucidate key aspects of plant parasitism by a cyst nematode. *Genome Biol. 2014*

153 **15**, 1–17 (2014).

9.   Masonbrink, R. *et al.* The genome of the soybean cyst nematode (Heterodera glycines)

reveals complex patterns of duplications involved in the evolution of parasitism genes.

BMC Genomics **20**, 119 (2019).

10.   Wu, S. *et al.* A reference genome of bursaphelenchus mucronatus provides new

resources for revealing its displacement by pinewood nematode. *Genes (Basel).* **11**,

(2020).

11.   Koutsovoulos, G. D. *et al.* Genome assembly and annotation of Meloidogyne enterolobii,

an emerging parthenogenetic root-knot nematode. *Sci. Data 2020 71* **7**, 1–13 (2020).

12.   Ali, M. A., Azeem, F., Li, H. & Bohlmann, H. Smart parasitic nematodes use multifaceted

strategies to parasitize plants. *Frontiers in Plant Science* (2017).

doi:10.3389/fpls.2017.01699

13.   Danchin, E. G. J., Guzeeva, E. A., Mantelin, S., Berepiki, A. & Jones, J. T. Horizontal

Gene Transfer from Bacteria Has Enabled the Plant-Parasitic Nematode Globodera

pallida to Feed on Host-Derived Sucrose. *Mol. Biol. Evol.* **33**, 1571–1579 (2016).

14.   Jones, J. T., Furlanetto, C. & Kikuchi, T. Horizontal gene transfer from bacteria and fungi

as a driving force in the evolution of plant parasitism in nematodes. *Nematology* (2005).

doi:10.1163/156854105775142919

15.   Grynberg, P. *et al.* Comparative genomics reveals novel target genes towards specific

control of plant-parasitic nematodes. (2020). doi:10.20944/preprints202010.0449.v1

16. Jen, F. Y., Tsay, T. T. & Chen, P. Aphelenchoides bicaudatus from ornamental nurseries in Taiwan and its relationship with some agricultural crops. *Plant Dis.* **96**, 1763–1766 (2012).

17. Subbotin, S. A. *et al.* The taxonomic status of Aphelenchoides besseyi Christie, 1942 (Nematoda: Aphelenchoididae) populations from the southeastern USA, and description of Aphelenchoides pseudobesseyi sp. N. *Nematology* (2020). doi:10.1163/15685411-bja10048

18. Oliveira, C. J. *et al.* Morphological and Molecular Identification of Two Florida Populations of Foliar Nematodes (Aphelenchoides spp.) Isolated from Strawberry with the Description of Aphelenchoides pseudogoodeyi sp. N. (Nematoda: Aphelenchoididae) and notes on their bionomics. *Plant Dis.* **103**, 2825–2842 (2019).

19. Wang, F. *et al.* Transcriptomic analysis of the rice white tip nematode, phelenchoides besseyi (Nematoda: Aphelenchoididae). *PLoS One* **9**, (2014).

20. Lilley, C. J., Kyndt, T. & Gheysen, G. Nematode Resistant GM Crops in Industrialised and Developing Countries. in *Genomics and Molecular Genetics of Plant-Nematode Interactions* 517–541 (Springer Netherlands, 2011). doi:10.1007/978-94-007-0434-3_24

21. Koenning, S. R. *et al.* Survey of crop losses in response to phytoparasitic nematodes in the United States for 1994. *Journal of Nematology* **31**, 587–618 (1999).

22. Jones, J. T. *et al.* Top 10 plant-parasitic nematodes in molecular plant pathology. (2013). doi:10.1111/mpp.12057

23. Wu, G. L., Kuo, T. H., Tsay, T. T., Tsai, I. J. & Chen, P. J. Glycoside hydrolase (gh) 45 and 5 candidate cellulases in aphelenchoides besseyi isolated from bird's-nest fern. *PLoS One* **11**, (2016).

24. Xu, X. *et al.* Population structure and species delimitation of rice white tip nematode, Aphelenchoides besseyi (Nematoda: Aphelenchoididae), in China. *Plant Pathol.* **69**, 159–167 (2020).

25. Rybarczyk-Mydłowska, K. *et al.* Small subunit ribosomal DNA-based phylogenetic analysis of foliar nematodes (Aphelenchoides spp.) and their quantitative detection in complex DNA backgrounds. *Phytopathology* (2012). doi:10.1094/PHYTO-05-12-0114-R

26. Kolmogorov, M., Yuan, J., Lin, Y. & Pevzner, P. A. Assembly of long, error-prone reads using repeat graphs. *Nat. Biotechnol.* (2019). doi:10.1038/s41587-019-0072-8

27. Durand, N. C. *et al.* Juicebox Provides a Visualization System for Hi-C Contact Maps with Unlimited Zoom. *Cell Syst.* (2016). doi:10.1016/j.cels.2015.07.012

28.　Yoshida, K., Hasegawa, K., Mochiji, N. & Miwa, J. Early Embryogenesis and Anterior-
Posterior Axis Formation in the White-Tip Nematode Aphelenchoides besseyi
(Nematoda: Aphelenchoididae). *J. Nematol.* **41**, 17–22 (2009).

29.　Ranallo-Benavidez, T. R., Jaron, K. S. & Schatz, M. C. GenomeScope 2.0 and
Smudgeplots: Reference-free profiling of polyploid genomes. *bioRxiv* 747568 (2019).
doi:10.1101/747568

30.　Cantarel, B. L. *et al.* MAKER: An easy-to-use annotation pipeline designed for emerging
model organism genomes. *Genome Res.* (2008). doi:10.1101/gr.6743907

31.　Finn, R. D. *et al.* Pfam: The protein families database. *Nucleic Acids Research* (2014).
doi:10.1093/nar/gkt1223

32.　Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biol.* **20**, 238 (2019).

33.　Kikuchi, T., Eves-Van Den Akker, S. & Jones, J. T. Genome Evolution of Plant-Parasitic
Nematodes. (2017). doi:10.1146/annurev-phyto-080516

34.　Nadachowska-Brzyska, K., Burri, R., Smeds, L. & Ellegren, H. PSMC analysis of effective
population sizes in molecular ecology and its application to black-and-white Ficedula
flycatchers. *Mol. Ecol.* **25**, 1058–1072 (2016).

35.　Mitreva, M. *et al.* The draft genome of the parasitic nematode Trichinella spiralis. *Nat.*
Genet. (2011). doi:10.1038/ng.769

36.　Tcherepanova, I., Bhattacharyya, L., Rubin, C. S. & Freedman, J. H. Aspartic proteases
from the nematode Caenorhabditis elegans: Structural organization and developmental
and cell-specific expression of asp-1. *J. Biol. Chem.* **275**, 26359–26369 (2000).

37.　McKerrow, J. H. *et al.* Strongyloides stercoralis: Identification of a protease that facilitates
penetration of skin by the infective larvae. *Exp. Parasitol.* **70**, 134–143 (1990).

38.　Koch, B. J., Ryan, J. F. & Baxevanis, A. D. The Diversification of the LIM Superclass at
the Base of the Metazoa Increased Subcellular Complexity and Promoted Multicellular
Specialization. doi:10.1371/journal.pone.0033261

39.　Dall, E. & Brandstetter, H. Structure and function of legumain in health and disease.
Biochimie **122**, 126–150 (2016).

40.　Holm Viborg, A. *et al.* A subfamily roadmap for functional glycogenomics of the
evolutionarily diverse Glycoside Hydrolase Family 16 (GH16).
doi:10.1074/jbc.RA119.010619

41.　Coconi Linares, N., Dilokpimol, A., Stålbrand, H., Mäkelä, M. R. & de Vries, R. P.
Recombinant production and characterization of six novel GH27 and GH36 α-

galactosidases from Penicillium subrubescens and their synergism with a commercial

mannanase during the hydrolysis of lignocellulosic biomass. *Bioresour. Technol.* (2020).

doi:10.1016/j.biortech.2019.122258

42. Shinya, R. *et al.* Possible stochastic sex determination in Bursaphelenchus nematodes.

Nat. Commun. 2022 131 **13**, 1–14 (2022).

43. Woodruff, G. C. Patterns of putative gene loss suggest rampant developmental system

drift in nematodes. *bioRxiv* 627620 (2019). doi:10.1101/627620

44. Danchin, E. G. J. *et al.* Multiple lateral gene transfers and duplications have promoted

plant parasitism ability in nematodes. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 17651–17656

(2010).

45. Rancurel, C., Legrand, L. & Danchin, E. G. J. Alienness: Rapid detection of candidate

horizontal gene transfers across the tree of life. *Genes (Basel).* **8**, (2017).

46. Campoy, E. & González-Martín, A. The Geography as a Regulator of Genetic Flow and

Genetic Structure in Andorra. *Adv. Anthropol.* (2017). doi:10.4236/aa.2017.72008

47. Morais, M. A. B. *et al.* Two distinct catalytic pathways for GH43 xylanolytic enzymes

unveiled by X-ray and QM/MM simulations. *Nat. Commun. 2021 121* **12**, 1–13 (2021).

48. Atanasova, L. *et al.* Evolution and functional characterization of pectate lyase PEL12, a

member of a highly expanded Clonostachys rosea polysaccharide lyase 1 family. *BMC*

Microbiol. **18**, 1–19 (2018).

49. PATHOGENICITY OF CURTOBACTERIUM FLACCUMFACIENS pv.

FLACCUMFACIENS TO SEVERAL PLANT SPECIES on JSTOR. Available at:

https://www.jstor.org/stable/45156052#metadata_info_tab_contents. (Accessed: 26th

August 2022)

50. Kikuchi, T., Jones, J. T., Aikawa, T., Kosaka, H. & Ogura, N. A family of glycosyl

hydrolase family 45 cellulases from the pine wood nematode Bursaphelenchus

xylophilus. *FEBS Lett.* **572**, 201–205 (2004).

51. Elling, A. A. *et al.* Sequence mining and transcript profiling to explore cyst nematode

parasitism. *BMC Genomics* **10**, (2009).

52. Rancurel, C., Legrand, L. & Danchin, E. G. J. Alienness: Rapid detection of candidate

horizontal gene transfers across the tree of life. *Genes (Basel).* **8**, (2017).

53. Szitenberg, A. *et al.* Comparative genomics of apomictic root-knot nematodes:

Hybridization, ploidy, and dynamic genome change. *Genome Biol. Evol.* **9**, 2844–2861

(2017).

54. Schiffer, P. H. *et al.* Signatures of the Evolution of Parthenogenesis and Cryptobiosis in

the Genomes of Panagrolaimid Nematodes. *iScience* **21**, 587–602 (2019).

55. Grynberg, P. *et al.* Comparative genomics reveals novel target genes towards specific

control of plant-parasitic nematodes. (2020). doi:10.20944/preprints202010.0449.v1

56. Holterman, M. *et al.* Disparate gain and loss of parasitic abilities among nematode

lineages. *PLoS One* (2017). doi:10.1371/journal.pone.0185445

57. Han, Z. *et al.* Horizontally Acquired Cellulases Assist the Expansion of Dietary Range in

Pristionchus Nematodes. *Mol. Biol. Evol.* **39**, (2022).

58. Zheng, J. *et al.* The ditylenchus destructor genome provides new insights into the

evolution of plant parasitic nematodes. *Proc. R. Soc. B Biol. Sci.* (2016).

doi:10.1098/rspb.2016.0942

59. Wan, X. *et al.* The Aphelenchus avenae genome highlights evolutionary adaptation to

desiccation. *Commun. Biol. 2021 41* **4**, 1–8 (2021).

60. Jagdale, G. B. & Grewal, P. S. Infection behavior and overwintering survival of foliar

nematodes, Aphelenchoides fragariae, on Hosta. *J. Nematol.* (2006).

61. Haegeman, A., Jones, J. T. & Danchin, E. G. J. Horizontal Gene Transfer in Nematodes:

A Catalyst for Plant Parasitism? *http://dx.doi.org/10.1094/MPMI-03-11-0055* **24**, 879–887

(2011).

62. Xia, J. *et al.* Whitefly hijacks a plant detoxification gene that neutralizes plant toxins. *Cell*

(2021). doi:10.1016/j.cell.2021.02.014

63. Brown, A. M. V. *et al.* Comparative genomics of wolbachia-cardinium dual endosymbiosis

in a plant-parasitic nematode. *Front. Microbiol.* **9**, 2482 (2018).

64. Holterman, M., Schratzberger, M. & Helder, J. Nematodes as evolutionary commuters

between marine, freshwater and terrestrial habitats. *Biol. J. Linn. Soc.* **128**, 756–767

(2019).

65. Ma, J. *et al.* Major episodes of horizontal gene transfer drove the evolution of land plants.

Mol. Plant **15**, 857–871 (2022).

66. Li, Y. *et al.* HGT is widespread in insects and contributes to male courtship in

lepidopterans. *Cell* **185**, 2975-2987.e10 (2022).

67. Ren, L., Huang, W., Cannon, E. K. S., Bertioli, D. J. & Cannon, S. B. A mechanism for

genome size reduction following genomic rearrangements. *Front. Genet.* (2018).

doi:10.3389/fgene.2018.00454

68. Cicconardi, F. *et al.* Chromosome Fusion Affects Genetic Diversity and Evolutionary

Turnover of Functional Loci but Consistently Depends on Chromosome Size. *Mol. Biol.*

Evol. **38**, 4449–4462 (2021).

69.   Wang, J. *et al.* Comparative genome analysis of programmed DNA elimination in
781   nematodes. *Genome Res.* **27**, 2001–2014 (2017).

782   70.   Fradin, H. *et al.* Genome Architecture and Evolution of a Unichromosomal Asexual
783   Nematode. *Curr. Biol.* **27**, 2928-2939.e6 (2017).

784   71.   Yin, D. *et al.* Rapid genome shrinkage in a self-fertile nematode reveals sperm
785   competition proteins. *Science (80-.).* **359**, 55–61 (2018).

786   72.   Wick, R. R., Judd, L. M. & Holt, K. E. Performance of neural network basecalling tools for
787   Oxford Nanopore sequencing. *Genome Biol.* (2019). doi:10.1186/s13059-019-1727-y

788   73.   Vaser, R., Sović, I., Nagarajan, N. & Šikić, M. Fast and accurate de novo genome
789   assembly from long uncorrected reads. *Genome Res.* (2017). doi:10.1101/gr.214270.116

790   74.   Medaka: Sequence correction by using ONT research.
791   https://github.com/nanoporetech/medaka

792   75.   Walker, B. J. *et al.* Pilon: An integrated tool for comprehensive microbial variant detection
793   and genome assembly improvement. *PLoS One* (2014).
794   doi:10.1371/journal.pone.0112963

795   76.   Alonge, M. *et al.* RaGOO: Fast and accurate reference-guided scaffolding of draft
796   genomes. *Genome Biol.* **20**, 224 (2019).

797   77.   Flynn, J. M. *et al.* RepeatModeler2 for automated genomic discovery of transposable
798   element families. *Proc. Natl. Acad. Sci. U. S. A.* (2020). doi:10.1073/pnas.1921046117

799   78.   Edgar, R. C. & Bateman, A. Search and clustering orders of magnitude faster than
800   BLAST. *Bioinformatics* **26**, 2460–2461 (2010).

801   79.   Coghlan, A., Coghlan, A., Tsai, I. J. & Berriman, M. Creation of a comprehensive repeat
802   library for a newly sequenced parasitic worm genome. *Protoc. Exch.* (2018).
803   doi:10.1038/protex.2018.054

804   80.   Tarailo-Graovac, M. & Chen, N. Using RepeatMasker to identify repetitive elements in
805   genomic sequences. *Current Protocols in Bioinformatics* (2009).
806   doi:10.1002/0471250953.bi0410s25

807   81.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: A flexible trimmer for Illumina
808   sequence data. *Bioinformatics* (2014). doi:10.1093/bioinformatics/btu170

809   82.   Dobin, A. & Gingeras, T. R. Mapping RNA-seq Reads with STAR. *Curr. Protoc.*
810   *Bioinforma.* (2015). doi:10.1002/0471250953.bi1114s51

811   83.   Grabherr, M. G. *et al.* Full-length transcriptome assembly from RNA-Seq data without a
812   reference genome. *Nat. Biotechnol.* (2011). doi:10.1038/nbt.1883

84.   Pertea, M. *et al.* StringTie enables improved reconstruction of a transcriptome from RNA-

seq reads. *Nat. Biotechnol.* (2015). doi:10.1038/nbt.3122

85.   Song, L., Sabunciyan, S. & Florea, L. CLASS2: Accurate and efficient splice variant

annotation from RNA-seq reads. *Nucleic Acids Res.* (2016). doi:10.1093/nar/gkw158

86.   Wu, T. D. & Watanabe, C. K. GMAP: A genomic mapping and alignment program for

mRNA and EST sequences. *Bioinformatics* (2005). doi:10.1093/bioinformatics/bti310

87.   Hoff, K. J., Lange, S., Lomsadze, A., Borodovsky, M. & Stanke, M. BRAKER1:

Unsupervised RNA-Seq-based genome annotation with GeneMark-ET and AUGUSTUS.

Bioinformatics (2016). doi:10.1093/bioinformatics/btv661

88.   Venturini, L., Caim, S., Kaithakottil, G. G., Mapleson, D. L. & Swarbreck, D. Leveraging

multiple transcriptome assembly methods for improved gene structure annotation.

Gigascience **7**, (2018).

89.   Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M.

BUSCO: Assessing genome assembly and annotation completeness with single-copy

orthologs. *Bioinformatics* (2015). doi:10.1093/bioinformatics/btv351

90.   Korf, I. Gene finding in novel genomes. *BMC Bioinformatics* **5**, 1–9 (2004).

91.   Stanke, M. *et al.* AUGUSTUS: A b initio prediction of alternative transcripts. *Nucleic Acids*

Res. (2006). doi:10.1093/nar/gkl200

92.   Howe, K. L., Bolt, B. J., Shafie, M., Kersey, P. & Berriman, M. WormBase ParaSite − a

comprehensive resource for helminth genomics. *Mol. Biochem. Parasitol.* **215**, 2 (2017).

93.   Davis, P. *et al.* WormBase in 2022—data, processes, and tools for analyzing

Caenorhabditis elegans. *Genetics* **220**, (2022).

94.   Stamatakis, A. RAxML version 8: A tool for phylogenetic analysis and post-analysis of

large phylogenies. *Bioinformatics* (2014). doi:10.1093/bioinformatics/btu033

95.   Xu, J. -H. Carbohydrate Active Enzyme database. in *Catalysis from A to Z* (2020).

doi:10.1002/9783527809080.cataz02801

96.   Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using

DIAMOND. *Nature Methods* (2014). doi:10.1038/nmeth.3176

97.   Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldón, T. trimAl: A tool for automated

alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* (2009).

doi:10.1093/bioinformatics/btp348

98.   Nguyen, L. T., Schmidt, H. A., Von Haeseler, A. & Minh, B. Q. IQ-TREE: A fast and

effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol. Biol.*

Evol. (2015). doi:10.1093/molbev/msu300