**The *Aphelenchoides* genomes reveal substantial horizontal gene transfers in the last common ancestor of free-living and major plant parasitic nematodes**

Cheng-Kuo Lai[1,2], Yi-Chien Lee[1,3,4], Huei-Mien Ke[5], Min R Lu[1], Wei-An Liu[1], Hsin-Han Lee[1], Yu-Ching Liu[1], Toyoshi Yoshiga[6], Taisei Kikuchi[7], Peichen J. Chen[8]* and Isheng Jason Tsai[1,2,3,4]*

Correspondence: janetchen@nchu.edu.tw and ijtsai@gate.sinica.edu.tw

[1]Biodiversity Research Center, Academia Sinica, Taipei 11529, Taiwan.
[2]Genome and Systems Biology Degree Program, National Taiwan University and Academia Sinica, Taipei, Taiwan.
[3]Biodiversity Program, Taiwan International Graduate Program, Academia Sinica and National Taiwan Normal University, Taipei, Taiwan
[4]Department of Life Science, National Taiwan Normal University, 116 Wenshan, Taipei, Taiwan
[5] Department of Microbiology, Soochow University, Taipei, 111, Taiwan
[6]Faculty of Agriculture, Saga University, Saga 840-8502 Japan
[7]Department of Integrated Biosciences, Graduate School of Frontier Sciences, The University of Tokyo, Chiba, 277-8562, Japan
[8]Department of Plant Pathology, National Chung Hsing University, Taichung, Taiwan

**Abstract**

*Aphelenchoides besseyi* is a plant-parasitic nematode (PPN) in the Aphelenchoididae family capable of infecting more than 200 plant species. *A. besseyi* is also a species complex with strains exhibiting varying pathogenicity to plants. We present the genome and annotations of six *Aphelenchoides* species, four of which belonged to the *A. besseyi* species complex. Most *Aphelenchoides* genomes have a size of 44.7-47.4 Mb and are amongst the smallest in clade IV, with the exception of *A. fujianensis*, which has a size of 143.8 Mb and is the largest. Phylogenomic analysis successfully delimited the species complex into *A. oryzae* and *A. pseudobesseyi* and revealed a reduction of transposon elements in the last common ancestor of *Aphelenchoides*. Synteny analyses between reference genomes indicated that three chromosomes in *A. besseyi* were derived from fission and fusion events. A systematic identification of horizontal gene transfer (HGT) genes across 27 representative nematodes allowed us to identify two major episodes of acquisition corresponding to the last common ancestor of clade IV or major PPNs, respectively. These genes were mostly lost and differentially retained between clades or strains. Most HGT events were acquired from bacteria, followed by fungi, and also from plants; plant HGT was especially prevalent in *Bursaphelenchus mucronatus*. Our results comprehensively improve the understanding of horizontal gene transfer in nematodes.

**Introduction**

The ability to parasitise plants has evolved in the phylum Nematoda on at least four occasions (Blaxter et al., 1998; Van Megen et al., 2009). The major plant parasites belonged to the Aphelenchodidae and Parasitaphelenchidae families making up the Aphelenchoidea superfamily and the Tylenchida order of clade IV nematodes (Bird, Jones, Opperman, Kikuchi, & Danchin, 2015); these plant parasitic nematodes (PPNs) collectively cause worldwide agriculture damages of over US$80 billion each year (Nicol et al., 2011). Root-knot nematodes in *Meloidogyne* genus cause the majority of these losses and were the first of PPNs to have their genomes sequenced (Abad et al., 2008; Opperman et al., 2008), followed by pinewood nematode *Bursaphelenchus xylophilus* (Dayi et al., 2020; Kikuchi, Cotton, Dalzell, Hasegawa, & Kanzaki, 2011), potato cyst nematode *Globodera pallida* (Cotton et al., 2014), soybean cyst nematode *Heterodera glycines* (Masonbrink et al., 2019) and others (Eves-van den Akker et al., 2016; Koutsovoulos et al., 2020; Wu et al., 2020). Comparing these genomes yield insight into several adaptions that allow PPNs to parasitize plants. Examples include effectors such as carbohydrate active enzymes (CAZyme), which are known to be secreted by PPNs and are hypothesized to be involved in degrading or modifying the composition of different plant structural tissues (Ali, Azeem, Li, & Bohlmann, 2017; Danchin, Guzeeva, Mantelin, Berepiki, & Jones, 2016). Some of these PPN-specific genes are known to be acquired from bacteria or fungi through horizontal gene transfer (HGT) (Jones, Furlanetto, & Kikuchi, 2005), giving nematodes the ability to adapt to different environments (Jones et al., 2005). Although numerous HGT genes have been identified and documented in different nematodes, research on the timing and subsequent maintenance of these genes, and why their copy numbers differ, has been restricted to a few PPN clades (Grynberg, Coiti Togawa, et al., 2020).

Currently, the only major groups containing plant parasitic nematodes that lack a reference genome are Trichodoridae and Aphelenchoididae. Of particular interest is *Aphelenchoides besseyi***, a foliar nematode that infects almost 200 plant species in 35 genera (Jen, Tsay, & Chen, 2012). This nematode is 0.4-0.8 mm in body size, has a life

cycle around 10 to 12 days and it can reproduce in extreme environments, making it hard to eliminate. Better known as the rice white tip, *A. besseyi* infects important agronomic crops such as rice, soybeans and strawberries (Oliveira et al., 2019; Subbotin et al., 2020), causing necrosis and distortion of its host's leaves (Jen et al., 2012; Oliveira et al., 2019; Wang et al., 2014). The nematode has reportedly been responsible for up to a 60% crop loss in some cases (Koenning et al., 1999; Lilley, Kyndt, & Gheysen, 2011) and was listed among the top ten plant parasitic nematodes in a 2013 review (Jones et al., 2013). Despite the economic damage these parasitic nematodes inflict, particularly in Asia, little is known about the basic biology, genetic diversity or evolution of *A. besseyi* and other Aphelenchoididae members.

        The taxonomic status of *A. besseyi* was reevaluated several times before molecular data were available. After its first description by Christie (1942), Yokoo (1949) identified another *Aphelenchoides oryzae* species, but this was later considered a synonym of *A. besseyi* due to the overlaps of several morphological characters (Allen, 1952). Several characteristics such as stylet length (Fortuner, 1970) have been added as standards to identify *A. besseyi* globally (Bulletin, 2017). More recently, molecular analyses indicated that *A. besseyi* may consist of several genetically diverged clades. We previously identified copy number variations of genes encoding cell-wall-degrading enzymes including glycosyl hydrolase family 5 (GH5) and GH45 cellulases between *A. besseyi* of different host origins (Wu, Kuo, Tsay, Tsai, & Chen, 2016). An 18S phylogeny separated these strains isolated from rice and fern unambiguously, and more recently new literature identified variations in different molecular markers in different hosts that are original to *A. besseyi* (Oliveira et al., 2019; Xu et al., 2020). Subbotin *et al*. used a combination of molecular makers (28S, ITS and mitochondria *COI* gene) (Subbotin et al., 2020) to reclassify foliar nematodes into three distinct clades: *A. besseyi* isolated mainly from strawberries, *A. oryzae* isolated mainly from rice and *A. pseudobesseyi* from wood fern. This study proposed this *A. besseyi* is a species complex, with each cryptic species associated with, but not restricted to, particular plant hosts. It has been reported that *A. besseyi* isolated from different hosts have varying levels of pathogenicity. For instance, *A. besseyi* populations isolated from strawberries

were unable to parasitise rice (Koenning et al., 1999), while populations of this species

from bird's-nest fern were able to reproduce in both rice and strawberries (Wang et al.,

2014). From an evolutionary perspective, *A. besseyi* is also interesting because its

primitive plant parasitism was a relatively recent evolutionary adaptation (Rybarczyk-

Mydłowska et al., 2012).

In this study, we sequenced and annotated the genomes of four *A. besseyi*

strains isolated from different plants which we later designated as *A. pseudobesseyi*

and *A. oryzae,* and another two species in the Aphelenchoididae family

(*Aphelenchoides bicaudatus* and *Aphelenchoides fujianensis*). We compared the

proteomes of six Aphelenchoididae members with 21 other representative nematodes to

delimit species relationships and investigated their gene family dynamics. We identified

synteny with representative nematodes and inferred rearrangement events to determine

how the three chromosomes of *A. besseyi* was evolved. The availability of the

*Aphelenchoides* assemblies allowed us to systematically determine the horizontal gene

transfer-acquired genes in nematode genomes. By inferring the evolutionary origins of

these HGT genes we found historical HGT events that shaped nematode evolution.

The major event occurred in the last common ancestor of clade IV nematodes and may

have contributed to the early adaptation of these nematodes.

144
145
**Results**

**Genome assemblies and annotations of six *Aphelenchoides* species**

We sequenced and assembled the genomes of six nematodes in the

*Aphelenchoides* genus (four *A. besseyi*, one *A. bicaudatus* and one *A. fujianensis*).

These species were chosen to represent the Aphelenchoididae family and *A. besseyi*

strains isolated from three plant hosts (**table S1**) to delimit their relationship within the

species complex. For each species, an initial assembly was produced from either 70-

148X Oxford Nanopore or 113-422X Pacbio reads using Flye assembler (Kolmogorov,

Yuan, Lin, & Pevzner, 2019) and further polished using Illumina reads (**table S2**).

Among *A. besseyi* assemblies, the VT strain isolated from tape grass *Vallisneria spiralis*

had the highest genome quality with N50 5.4 Mb (hereafter denoted as APVT). The contigs of this strain were further scaffolded with 150X Hi-C reads using the Juicer program (Durand et al., 2016) (**fig. S1**) yielding a final assembly of 44.7 Mb (N50 = 16.9 Mb). More than 99% of this assembly was in three scaffolds, presumably corresponding to three chromosomes (Yoshida, Hasegawa, Mochiji, & Miwa, 2009) (2n=6). Five *Aphelenchoides* assemblies ranged from 44.7 to 47.4 Mb (N50 = 12.2-17.8 Mb; **table S3**), and a sixth assembly (*A. fujianensis*) of 143.8 Mb (N50 = 553 kb; **table S3**) was estimated to be triploid (Ranallo-Benavidez, Jaron, & Schatz, 2020) (**fig. S2**). Although not present in the assemblies, the telomere motif TTAGGC was identified in the reads of *A. pseudobesseyi* at low coverage (**Supplementary Info**), which is consistent with the sister group species of *B. xylophilus* (**table S4**) and indicates the presence of telomeres in these species.

Using the proteomes of *Bursaphelenchus xylophilus* and *Caenorhabditis elegans*, and the transcriptomes of pooled worms in each species as evidences, we predicted 11,701 to 12,948 protein-coding genes in six *Aphelenchoides* species with Maker2 (Cantarel et al., 2008) (**table S3**). With the exception of *A. fujianensis*, these were fewer protein coding genes in these species than in Tylenchida nematodes (12,762 to 19,212) and free-living nematodes (20,184 to 20,992). The completeness of annotated genes was estimated to be 76.4–81.3% based on a BUSCO assessment, lower than that of *Bursaphelenchus* species (83.0–89.4%), but higher than that of Tylenchida (59.8–75.4%) nematodes. 66.5% to 71.0% of genes in *Aphelenchoides* could be assigned at least a domain from the Protein family (Pfam) database (Finn et al., 2014). In addition, orthologous groups were inferred with the proteomes of six *Aphelenchoides* with 21 other nematodes using Orthofinder (Emms & Kelly, 2019). With the exception of *A. fujianensis*, 78.5–85.4% (*A. fujianensis* = 48.7%), 69.4–76.9% (*A. fujianensis* = 42.8%) and 87.5–98.7% of *Aphelenchoides* genes were orthologous to *B. xylophilus*, *C. elegans* and at least one other nematode species, respectively, suggesting that the reduced proteome in most *Aphelenchoides* was mainly comprised of conserved genes among nematodes.

**Phylogenomics delimit species complex of *Aphelenchoides besseyi***

To investigate the evolution of plant-parasitic nematodes and the relationships

among members in the *A. besseyi* species complex, a maximum-likelihood phylogenetic

tree was constructed based on 75 low-copy orthologues. The phylogeny is consistent

with the previous study (Kikuchi, Eves-Van Den Akker, & Jones, 2017): the major plant

parasitic nematodes were divided into Aphelenchoidea and Tylenchida, and six

*Aphelenchoides* species were grouped as sister to *Bursaphelenchus* (**fig. 1a**). The *A.*

*besseyi* strains were clustered into two groups based on their hosts, suggesting that

relationships in these species within the *A. besseyi* species complex were associated

with host preferences. Combined with the previous 28S phylogeny of the *A. besseyi*

species complex (Subbotin et al., 2020) (**fig. S3**), we further designated these two

groups as *A. oryzae* and *A. pseudobesseyi* groups isolated from rice or other plants

(land grass and bird's-nest fern). The median nucleotide and amino acid identity was

86.6% and 90% between these two groups, respectively (**fig.S4**). Such levels of

divergence were comparable to species comparisons of other clades (87.2% and 92.8%

amino acid identity in *B. xylophilus–B. mucronatus* and *M. incognita–M. floridensis,*

respectively). Strains in each group also differed in heterozygosity (0.017-0.019% in *A.*

*oryzae* vs 0.071-0.075% in *A. pseudobesseyi*) and changes in recent effective

population sizes inferred using pairwise sequentially Markovian coalescent (PSMC)

analysis (Nadachowska-Brzyska, Burri, Smeds, & Ellegren, 2016) (**fig.5**). Together

these results demonstrated that relationships among the two designated clades in the

*A. besseyi* species complex were distinct species, well-differentiated at the genome

level, despite being challenging to differentiate based solely on morphology (Subbotin et

al., 2020).

**Most *Aphelenchoides* species have reduced genomes and repetitive elements**

The *Aphelenchoides* genomes were smaller than those of other plant-parasitic

nematodes (**fig. 1b** and **table S3**). Much of the reduction can be explained by the

reduced markup of repeat content compared to other nematodes (**fig. 1b**). The

dominant transposable elements of *Aphelenchoides* were DNA transposons—which

were reduced in content (0.14–1.36 Mb vs. 4.2–22.1 Mb in other nematodes)—and

number of families (1–7 in *Aphelenchoides* compared to 9 and 26 in *B. xylophilus* and

*H. glycines,* respectively) compared to other nematodes. Fewer LTR (0.07–0.8 Mb vs.

0.24–9.3 Mb) and LINE (0.0006–0.66 Mb vs 0.02–4.5 Mb) retrotransposons were also

observed in this genus. These results suggest that the reduced genome sizes in

*Aphelenchoides* might have been caused by the loss of transposable elements and led

to the eventual loss of entire families in some cases (**fig. 2a**). Within the *A. besseyi*

species complex, *A. pseudobesseyi* contained significantly fewer DNA transposons,

LTR and LINE retrotransposons than *A. oryzae* (**fig. 2a and fig.6**).

**Figure 1 Phylogenomic analysis of plant-parasitic nematodes. a.** The phylogeny of 27

representative nematodes was inferred based on the concatenation of protein sequences from

75 low-copy orthologues. The blue dots in branches denote a bootstrap value of 100. **b.** The

size of the genome features in nematodes, repeats containing DNA transposons (DNA), long

interspersed nuclear elements (LINE), long terminal repeats (LTR), rolling-circular (RC) and the

unclassified repeats are labelled as "other repeats". Numbers in brackets denote genome

assembly sizes in megabases.

**Figure 2 Repeat and proteome contents in nematodes. a.** The genome proportions of DNA,
LINE and LTR transposable elements in nematodes shown by genome percentage **b-c.** Protein
families and CAZyme gene copy numbers vary significantly among nematodes. The dot size
represents the copy number of each domain and the different colour represents the copy
number of domains larger or lower than average copies shown in brackets. Pfam and CAZyme
families that were significantly different between *Aphelenchoides* species and other nematodes
are denoted by the "+" symbol. CAZyme families that was identified as having been acquired
horizontally are denoted by the "*" symbol.

**Gene family specialization in the *Aphelenchoides* species**

To gain insight into the genomic features associated with the biology and plant
parasitism of *Aphelenchoides*, we aimed to identify expansion or contraction of protein
domains and carbohydrate-active enzymes (CAZymes) specific to this clade. We
observed 66 enriched and 31 reduced protein domains in the four members of the *A.
besseyi* species complex compared to 21 other nematodes. (Wilcoxon rank-sum test q
value < 0.05; **fig. 2b** and **table S5**). Reduced domains included collagen (90–109
copies in the *A. besseyi* species complex vs. 72–407 in others), Somatomedin B and
BTG/Tob (Winkler, 2010). Genes containing collagen domains were reportedly

associated with capsule formation; the reduced copy of collagen domains in *Trichinella spiralis* were thought to contribute its lower host-specificity than other nematodes (Mitreva et al., 2011), and may be related to the wide host range of *A. besseyi*. In contrast, Aphelenchoidea members possess on average four-fold (91–314 vs. 4–555 copies) more aspartic proteases (ASP) than other nematodes (**table S5**). Other expansions included LIM and peptidase C13 domains, which participate in participating in the regulation of cell motility and cell growth (Koch, Ryan, & Baxevanis, 2021) or degradation of protein tissues in a host (Dall & Brandstetter, 2016), emphasizing that these domain dynamics are associated with adaptations to plant parasitism.

The plant cell wall acts as a primary defensive barrier and the production of CAZyme families are important for PPNs to infect plants. A total of 132 CAZyme families were identified in the representative 27 nematode species. Of these, 59–67% of the CAZyme families were observed in Aphelenchoidea which is similar to the 55–66% and 58-68% of the families in Tylenchida and free-living nematodes (**table S6**), respectively. A total of 13 families were significantly expanded or lost in the *Aphelenchoides* genus (**fig. 2c**), including GH16, GH27 and GH45. GH16 serves as the putative β-glycanases involved in the degradation or remodelling of cell wall polysaccharides (Holm Viborg et al., 2019), GH16 had one to six copies in Aphelenchoididae nematodes and was not identified outside this clade except in *D. destructor*, in which there were three copies. There are three to 11 copies of GH27— which are reportedly involved in the function of hemicellulose and associated with α-galactosidase activity in both bacteria and fungi (Coconi Linares, Dilokpimol, Stålbrand, Mäkelä, & de Vries, 2020)— in Aphelenchoidea, but fewer in the Tylenchida nematodes. The previously identified GH45 present in Aphelenchoidea nematodes (Wu et al., 2016)—involved in the degradation of beta-1,4-glucans in the plant cell wall (Wang et al., 2014)—possess different copy numbers between *A. pseudobesseyi* and *A. oryzae* and were absent in *A. fujianensis* and *A. bicaudatus,* suggesting differential maintenance of these genes in the same genus may have contributed to variations of pathogenicity to plants.

**Chromosomal evolution of PPNs**

To investigate the extent of the karyotype rearrangements in *Aphelenchoides*, we inferred the synteny relationships among *A. pseudobesseyi* (chromosome n=3) (Yoshida et al., 2009), *B. xylophilus* (n=6) and *C. elegans* (n=6) using single copy orthologs. Within the three *A. pseudobesseyi* chromosomes, orthologs belonging to all *C. elegans* chromosomes were clustered into distinctive blocks (**fig. 3a**) suggesting a fusion of ancestral chromosomes. These regions remained contiguous and contained 148-801 orthologous genes that could be assigned from individual chromosomes presumably not yet broken down yet by recombination, allowing us to pinpoint the fusion points and infer the order of rearrangement events based the constitution of chromosomes (**fig. 3b**). We encountered instances of where an ancestral chromosome was found in different parts of the *A. pseudobesseyi* chromosomes, suggesting fission also took place. In the case of chromosome IV—which remained homologous in *C. elegans* and *B. xylophilus*—corresponding synteny blocks in *A. pseudobesseyi* were identified in the arm of chromosome 2 and chromosome 1 separated by regions of chromosome III origin (**fig. 3b**). The majority of the ancestral sex chromosomes were unambiguously assigned to chromosome 2, and remapping of male sequences showed equal coverage along the chromosomes (**fig.S7**), suggesting that the Aphelenchoidea superfamily including *A. besseyi* exhibited a stochastic sex determination system that was recently characterized in *B. xylophilus* (Shinya et al., 2022). Within the *A. besseyi* species complex, a total of 91% and 88% of genomes were in synteny between APVT and AORT, respectively. Intra-chromosomal inversions were common at chromosome arms. In addition, we identified a major inversion of length 3.4 Mb long located in the centre of chr 2 (**fig. 3c**) suggesting rearrangement is still ongoing. Both the LTR and LINE retrotransposons were enriched in the chromosome arms of the *A. oryzae* strain (AORT) (**fig. 3c and fig.S6**), which is consistent with repetitive element distributions in other nematode chromosomes (Woodruff, 2019). In contrast, only the LTR retrotransposons were found in the two chromosome arms of *A. pseudobesseyi*, suggesting that these repeats were differentially maintained after speciation.

**Figure 3. Chromosome evolution of plant-parasitic nematodes. a.** The density of pairwise single-copy orthologs between *A. pseudobesseyi* VT and *B. xylophilus*. Colours denote the Nigon elements inferred from 15 rhabditid nematodes (de la Rosa et al., 2021) and the putative chromosome fusion sites in APVT are labelled with red triangles. **b.** Dashed lines indicate rearrangement events. **c.** The synteny relationship and the distribution of transposable between *A. pseudobesseyi* VT and *A. oryzae* RT. Blocks indicate synteny links between the two strains, and the line colours correspond to inversion (blue) and inter-chromosomal rearrangement (grey). Distribution of DNA transposons (red), long interspersed elements (green) and long terminal repeats (blue) between two stains are shown.

**Major episode of HGT in clade IV nematodes**

In plant parasitic nematodes, the GH5 cellulase was found present in Tylenchida and only *A. pseudobesseyi* and *A. bicaudatus* within the Aphelenchoidea clade (Danchin et al., 2010; Wu et al., 2016), raising the possibility that many of the horizontal gene transferred (HGT) genes were acquired in the last common ancestor of major PPNs but were differentially lost. To identify such events, a total of 27 proteomes from representative nematodes including the *Aphelenchoides* genomes were searched for evidence of HGT by calculating the Alien Index (AI) score using Alienness (Rancurel, Legrand, & Danchin, 2017), which has been applied to infer HGT candidates in various nematode species (Grynberg, Togawa, et al., 2020; Schiffer et al., 2019; Siddique et al., 2021). We identified a total of 1,675 HGT orthogroups in 22 nematodes. Placing these orthologs designated as events onto the species phylogeny assuming a parsimonious scenario (Campoy & González-Martín, 2017), indicated that HGT started in the last common ancestor of clade IV nematodes (**fig 4a**). Examples include GH16, GH32, GH43 and the aforementioned GH5 cellulases. We inferred a total of 161 orthogroups were acquired in this episode, and most of their origins were inferred to be bacteria (78.3%) (**table S7**) belonging to different genera, suggesting multiple acquisitions took place. Of these, we found 36 Pfam terms such as ABC transporter that were identified in multiple orthogroups suggesting some convergence in the acquired functions (**table S8**).

The revised GH5 cellulase phylogeny indicated an ancient duplication took place before the divergence of PPNs (**fig 5a**). One clade contains orthologs of the three *Panagrolaimid* (*P.* sp. PS1159, *P. superbus* and *P. davidi*), and Tylenchida, and the other clade contains members of Aphelenchoidea and Tylenchida nematodes, which emphasises that the fate of the HGT genes was governed by duplications and loss. Interestingly, the closest GH5 bacterial orthologs were *Salinimicrobium xinjiangense* and *Leeuwenhoekiella sp.*, which belonged to Flavobacteriaceae family and were from marine environments. We observed two GH16 subfamilies in nematodes. GH16_3 in Tylenchida and *Bursaphelenchida* nematodes were clustered with bacterial origin sequences, whereas GH16_1 of *Aphelenchoides* and *Panagrolaimus* nematodes were clustered with fungal origin (**fig 5b**), suggesting that the two GH16 groups arose

independently. GH32 in *G. pallida* (Danchin et al., 2016) is believed to play a role in the function of fructose hydrolysis and was found in one *Panagrolaimus* in addition to several Tylenchida nematodes (**fig.S8**). GH43 was identified at two distinct clusters of bacterial origin in Tylenchida and *Panagrolaimid* nematodes which have been proposed to be involved in degradation of the hemicellulose in plants (Morais et al., 2021) (**fig.S9**).

The next major episode of acquisition took place in the common ancestor of PPNs, with 47 orthogroups (**fig. 4a**). These families included pectate lyases 3 (PL3) which is associated with cell wall degradation (Atanasova et al., 2018). The orthologs of PL3 in *Aphelenchus avenae* and two *Bursaphelenchus* nematodes were grouped together with distinct clusters of *Meloidogyne* species (**fig. 5c**) is consistent with previous phylogenetic findings in PPNs (Danchin et al., 2010). The closest bacterial ortholog in the *Meloidogyne* clade was from *Curtobacterium flaccmfaciens* which is also known to cause bacterial wilt in the Fabaceae family (Júnior et al., 2012). Together, these results suggested some genes that were thought to play important roles in plant parasitism were in fact acquired earlier than the common ancestor of plant parasitic nematodes.

The majority of HGT gene families were of bacterial followed by fungal origin (**fig 4b**). We also identified genes that were acquired from non-bacterial donors in the last common ancestors of clade IV, as well as in more recent, different PPN lineages (**fig 4a**). This included the previously characterised fungal origin of GH45 (Kikuchi, Jones, Aikawa, Kosaka, & Ogura, 2004; Wu et al., 2016), This cellulase family is present in most Aphelenchoidea nematodes except *A. fujinensis* and *A. bicaudatus*. The GH16 family was independently acquired from a bacterial and fungal donor in the last common ancestor of clade IV nematodes and the *Aphelenchoides* genus, respectively (**fig 5b**). Notably, we identified 40 orthogroups among PPNs that were transferred from the plant phylum Streptophyta, which is consistent with the finding of several sequences that are highly similar to plants in *H. glycine* (Elling et al., 2009) (**fig. 4b**). The closest plant orthologs included rice, maple and oak (**fig. 5d**) which are common hosts to many PPNs. Strikingly, of these orthogroups, 27 were present in *B. mucronatus* and enriched

in the detoxification of cadmium and copper ion function **(table S9)**, suggesting these

genes may help *Bursaphelenchus* nematodes to degrade the toxin in pine wood hosts.

**Figure 4 The evolution of genes acquired from horizontal gene transfer (HGT) in clade IV**

**nematodes. a.** HGT orthogroups were inferred by the AI score (Rancurel, Legrand, & Danchin,

2017) > 0 across 27 representative nematodes; the HGT families gained or lost are shown in

the branches. Horizontally acquired CAZymes are annotated. The proportions of donor origins

in each HGT orthogroup belonging to different kingdoms of donors are shown as pie charts. The

size of the pie chart corresponds to the total number of HGT orthologues in branches; the chart

was normalized using: $\log_5$(total number). **b.** The distribution of HGT families transferred from

different kingdoms, with the same denoted color scheme as same as figure a. **c.** Number of

HGT events among different inferred time points corresponding with the branches which are

marked with triangles in the phylogeny.

**Figure 5 Phylogenies of nematode HGT orthogroups and their donor origin. a.** GH5_2 **b.**
GH16 **c.** PL3 **d.** OG0003777 and OG0004537. Different colours denote different kingdoms and
species as shown in legend. Nematode gene copies with negative AI values were marked with
an asterisk. Additional Pfam domains are labelled when available. Nodes with iqtree UFBoot
and SH-aLRT bootstrap support > 80% are labelled as grey circles.

We identified 0.3-2.4%, 0.6-2.1% and 0.1-5.4% proteomes among

Aphelenchoidea, Tylenchida and *Panagrolaimomorpha* nematodes that were predicted

to be HGT (**fig. 4c**). The majority of these differences were the result of clade-specific

evolution after speciation. The high copy number of HGT genes observed in *M.*

*incognita* was a result of duplication (Szitenberg et al., 2017), indicated by the fact that

the number of HGT orthologs of bacteria origin were over two times higher than any

other species (**fig. S10**). The high number of HGT genes in *P. superbus* was consistent

with a previous study (Schiffer et al., 2019) and likely to be species-specific.

To independently assess the accuracy of our approach and interrogate the fate

of HGT genes, we constructed a phylogeny for every orthogroup containing identified

HGT candidates. Members of Aphelenchoididae and Tylenchida orthologs in the

majority of these orthogroups were predicted to be all HGT genes (with AI > 0; 54.6-

76.5% vs. 77.3-89.4%). Genes from a species were typically grouped together in the

orthogroup phylogeny regardless of being identified as HGT candidates, suggesting the

genes that were not detected using our threshold shared common ancestries with those

that were. Presumably, this was a result of accumulating substitutions over time.

Consistent with this observation, the more ancient acquired HGT orthogroups in PPNs

contained higher copy numbers of these genes compared to recently acquired families

(**fig.S11**). The instances included GH5 families with 12.5-70.6% of copies in Tylenchida

that could not be identified as HGT candidates, suggesting duplication and possibly

neo-functionalisation of the GH family in PPNs after being acquired from bacteria (**fig.**

**5a**). The differentiation was ongoing and observed in the *A. besseyi* species complex,

which included the GH45 orthogroup with negative AI in two *A. oryzae* strains (**fig.S12**).

## Discussion

441  Characterising the diversity and comparing the genomes of plant parasitic
nematodes has been of fundamental importance in understanding how such lifestyles
arise and of practical importance in identifying candidate effectors and control methods.
The latter has been addressed in several studies, focusing mainly on *Meloidogyne*
(Grynberg et al., 2020). The *Aphelenchoides* genome assemblies presented in this
study allowed us to gain a holistic view of the evolution of clade IV nematodes, which
appeared to gain and lose many adaptations, including plant parasitism (Holterman et
al., 2017). In their evolution, HGT genes have played important roles in functions related
to these adaptations. The most recent comprehensive analyses of HGT in nematodes
focused on plant parasitic nematodes (Grynberg et al., 2020) and found many of these
genes were PPN-specific. The donors of these PPN-specific gene families were
sympatric plant bacteria which may have facilitated the possibility of HGT (Danchin et
al., 2010). Additional HGT events were identified in other clade IV nematodes (Danchin
et al., 2016; Han et al., 2022; Schiffer et al., 2019; Wan et al., 2021; Zheng et al., 2016)

456  Our systematic investigation of HGT has shown that many of these families were
acquired much earlier in the last common ancestor of clade IV. Sources of these donors
may be symbionts like the case of insects (Xia et al., 2021), but currently nematode
endosymbionts are restricted to *Wolbachia* and *Cardinium* (Brown et al., 2018) and
were not identified in our analyses. Interestingly, many of the closest bacterial donors
were from marine environments, raising the possibility that the last common ancestor of
clade IV may have lived in a marine environment that underwent habitat transition
(Holterman, Schratzberger, & Helder, 2019). However, we also identified donors of non-
bacterial origin that were usually found in the environments that fit nematodes' present
day lifestyle. Now that more genome sequences are available, historical HGT events
were detected in other nematode clades (Mayer et al., 2011; Zarlenga et al., 2019) as
well as in the most recent common ancestor of major organism groups such as land
plants (Ma et al., 2022) , of moths and butterflies (Li et al., 2022), which contributed the
hosts' developmental roles and adaptations. These acquisitions were found to be
episodic and likely took place in a time when either the host development or genome
defence was vulnerable.

The successful delimitation of the *A. besseyi* species complex unambiguously

into *A. oryzae* and the recently proposed *A. pseudobesseyi* has important implications

in nematode management. Congruent delimitation was observed between genomes

and 28S phylogenies, confirming the utility of species identification with existing

molecular markers (Oliveira et al., 2019). We thus recommend that the taxonomic status

of all *A. besseyi* strains be reclassified into either species. *A. besseyi* is generally

believed to have limited mobility in natural habitats, so its lack of population structure in

China (Xu et al., 2020) was suggested as a consequence of human-mediated dispersal.

Our results also supported that *A. oryzae* appears to be more rice plant-specific

compared to *A. pseudobesseyi* which was isolated more frequently in ornamental plants

and other agronomic crops (Oliveira et al., 2019). A comprehensive collection across a

wider geographical range and resequencing of strains previously designated as *A.*

*besseyi* could confirm whether *A. oryzae* was responsible for all the white tip disease in

rice plants and may lead to better characterisations of the biogeography and evolution

of different cryptic species.

To conclude, the availability of the *Aphelenchoides* genome and our comparative

analyses allowed us to pinpoint the major events of horizontal gene transfer in clade IV

nematodes. The results have reinforced the importance of horizontal gene transfers

contributing to multiple adaptations of these nematodes including plant parasitism. In

addition, the various *A. besseyi* genomes will assist in developing molecular diagnostic

tools to distinguish the specific diseases caused by the species complex.

**Methods**

**DNA, RNA extraction and sequencing**

Nematodes were cultured with *Alternaria citri* on PDA (potato dextrose agar)

medium. All stages of nematodes were collected from the medium, washed with sterile

distilled water, and purified by sucrose gradients. Genomic DNA was extracted using

Qiagen Genomic-tip 100/G according to the manufacturer's instructions, RNA extraction

was conducted using Trizol, and then purified using a lithium chloride purification

method. The DNA paired-end libraries were constructed using either a Nextera DNA

Flex or KAPA hyper library prep kit (Illumina, San Diego, USA); the RNA paired-end

libraries were constructed using a TruSeq Stranded mRNA library prep kit (Illumina, San

Diego, USA). Both DNA and RNA pair-end followed with standard protocol and were

sequenced by Illumina HiSeq 2500 (Illumina, USA) to produce 150-bp paired-end reads.

The HiC library preparation was performed by Phase Genomics (Seattle, WA, USA)

proximo HiC animal protocol with some modification in tissue processing. The enriched

worms were finely chopped by microtube pellet pestle rods for about 2 minutes. The

tissues were crosslinked by adding 1 ml crosslinking solution and incubate for 25

minutes with occasional mixing by rotation. 100 ul quenching solution was added to the

crosslinked tissue and mixed for 20 minutes by rotation. The rest of the preparation

steps follow the protocol. The library was sequenced by Illumina HiSeq 2500 (Illumina,

USA) to produce 150-bp paired-end reads. APFT and AORT were using Pacbio

sequencing system to produce long-read, and the rest of 4 *Aphelenchoides* strains

(APVT, AORJ, *A. bicaudatus*, *A. fujianensis*) were sequenced using the Oxford

Nanopore sequencing platform. The raw nanopore signals were basecalled by Guppy

(Wick, Judd, & Holt, 2019) (ver 0.5.1) producing a total of 5.0-28.4 Gb sequences at

least 1 kb in length.

**Assemblies of six *Aphelenchoides* spp*.**

Raw reads of each species were assembled using Flye (ver 2.8.2)(Kolmogorov

et al., 2019) assembler. The assemblies from Nanopore reads were corrected using

Nanopore reads using Racon (Vaser, Sović, Nagarajan, & Šikić, 2017) (ver 1.4.6) and

Medaka (ver 0.10.0; https://github.com/nanoporetech/medaka). All assemblies were further

corrected using Illumina reads using Pilon (Walker et al., 2014) (ver 1.22) with five

iterations. The *A. pesudobesseyi* VT  assembly was scaffolded using HiC reads and

subsequently curated in Juice-box (Durand et al., 2016) tools. The other five

*Aphelenchoides* genomes were reference scaffolded based on this assembly using

Ragtag (Alonge et al., 2019) (ver 1.1).

**Gene prediction and functional annotation**

The identification of repetitive elements were computed by RepeatModeler (Flynn

et al., 2020) (ver 1.0.8), TransposonPSI (ver 1.0.0;

https://github.com/NBISweden/TransposonPSI) and USEARCH (Edgar & Bateman,

2010) (ver 8.1) based on the protocol by Berriman *et al*. (Coghlan, Coghlan, Tsai, &

Berriman, 2018). Repeat locations were then identified by Repeatmasker (Tarailo-

Graovac & Chen, 2009) (ver 4.0.9). RNA-seq reads of six *Aphelenchoides* strains were

trimmed by Trimmomatic (Bolger, Lohse, & Usadel, 2014) (ver 0.36), and aligned to

corresponding assemblies using STAR (Dobin & Gingeras, 2015) (ver 2.7.1a). From

these mappings, transcripts were inferred using three approaches: i) assembled based

on the mappings as guides using Trinity (Grabherr et al., 2011) (ver 2.84; option: default

setting), reconstructed using ii) Stringtie (Pertea et al., 2015) (ver 1.3.4; option: default

setting) and iii) CLASS2 (Song, Sabunciyan, & Florea, 2016) (ver 2.17; option: default

setting). Transcripts generated from Trinity were realigned to the reference using GMAP

(Wu & Watanabe, 2005) (ver 2017-11-15). The RNA-seq mappings were also used in

BRAKER (Hoff, Lange, Lomsadze, Borodovsky, & Stanke, 2016) to train species

parameter and generate an initial set of annotations. Proteomes of *Bursaphelenchus*

*xylophilus* and *Caenorhabditis elegans* were downloaded from Wormbase ParaSite

(WBPS14; Howe, Bolt, Shafie, Kersey, & Berriman, 2017) and used as homology

guides to pick the best transcripts for each putative locus using MIKADO (Venturini,

Caim, Kaithakottil, Mapleson, & Swarbreck, 2018) (ver 1.2.4; option: three Mikado

steps, containing "prepare", "serialize" and "pick" procedures), and were also used to

train MAKER2. Finally, MAKER2 was invoked to generate a final set of gene

annotations using picked EST evidence and protein evidence from MIKADO transcript

and proteomes from closely related species (*Bursaphelenchus xylophilus* and

*Caenorhabditis elegans*), and used gene models (BUSCO (Simão, Waterhouse,

Ioannidis, Kriventseva, & Zdobnov, 2015), BRAKER, SNAP (Korf, 2004) and Augustus

(Stanke et al., 2006)) as EST hints to train predicted data with three iterations. The

APVT strain was predicted using trained models based on the manual curation of 975

genes.

**Comparative analyses**

Proteomes of five plant-parasitic nematodes (*Bursaphelenchus xylophilus,*

*Meloidogyne hapla, Meloidogyne incognita, Globodera pallida, Ditylenchus destructor*),

two free-living nematodes (*Caenorhabditis elegans, Caenorhabditis briggsae*), six

*Panagrolaimomorpha* (*Propanagrolaimus* sp. JU765, *Panagrellus revidius,*

*Panagrolaimus superbus, Panagrolaimus* sp. PS1159*, Panagrolaimus davidi* and

*Halicephalobus mephisto*) and one animal parasitic nematode (*Brugia malayi*) were

downloaded from Wormbase WBP17 (Howe, Bolt, Shafie, Kersey, & Berriman, 2017)

and the longest isoforms were selected for the further analyses. Only the longest

isoform per gene was considered for subsequent analyses. Orthogroups were

determined by Orthofinder (Emms & Kelly, 2019) (ver 2.2.7; options: -S diamond). Low-

copy orthogroups were applied due to multiple genome duplication events in *M.*

*incognita*, *A. avenae* and two *Panagrolaimus* species. Sequence alignments of each of

the single-copy orthogroups were generated by MAFFT (ver 7.310; options: --maxterate

1000). Then, the concatenated alignment of all single-copy orthogroups was used to

compute a maximum likelihood phylogeny using RAxML (Stamatakis, 2014) (ver 8.2.3;

options: -s -T 32 -N 100 -f a -m PROTGAMMILGF) with 100 bootstrap replicates. Pfam

copy numbers of all 27 nematodes were identified from the results of nematode

proteomes blast against the database of Pfam website (ver 31; https://pfam.xfam.org/)

using hmmscan (e value < 0.001; Finn, Clements, & Eddy, 2011). To identify putative

effector enzymes, we searched the nematode proteomes against the CAZyme database

(http://www.cazy.org; Drula et al., 2022) using hmmscan (Finn et al., 2011). We

considered only sequences that were at least 80bp in length, had a conserved domain

proportion of at least 0.35 of its length, and an e-value of less than 1e-15. Nigon

elements of 15 rhabditid nematodes in *A. pseudobesseyi*, *B. xylophilus* and *C. elegans*

were inferred based on (https://github.com/pgonzale60/vis_ALG; de la Rosa et al.,

2021) which assigned Nigon units using BUSCO output (ver 4.8.4, options: genome;

dataset: nematoda_odb10).

**Identification of the HGT genes**

The probability of genes having been acquired via HGT was estimated by using

Alienness Index (AI) (Rancurel et al., 2017). Our donor group were generated by non-

Metazoans from NCBI nr database, and the recipient were Metazoans excluding the

following species to prevent self-alignment: Aphelenchoidea, Tylenchida, Rhabditina,

Spirurina and Cephaloboidea. The Alien Index (AI) was estimated by calculating the e-value of diamond (Buchfink, Xie, & Huson, 2014) (ver 2.0.14; option: blastx --evalue 0.001) best hits between the donor and recipient database. Orthogroups having at least one gene with an AI value over 30 were selected for further analysis. Gains and losses at each node were inferred using Phylip-Dollop (Campoy & González-Martín, 2017) (ver 3.69; options: fdollop -method d -ancseq). Some of the HGT family acquired branches were manually curated by their evolutionary place of gene phylogeny due to the fact that nematode genes with AI < 0 were clustered with other HGT genes. The highest AI value of nematode genes with classified taxonomy hit were chosen to represent the HGT origin in each orthogroup. Orthogroups with the same CAZyme annotated and nematode orthology gene AI higher than -50 in those Orthogroups were selected. AI < 0 genes were labelled with "*". The orthologs were further combined with the HGT identified donor sequence from nr database and the specific cellulase sequence from CAZyme database. To reduce contamination, orthologs of Pfam domain were annotated and filtered by having at least one major domain (cellulase or pectate lyase). Sequences of each HGT orthogroup were aligned using MAFFT (options: --maxiterate 1000 --genafpair) and trimmed by using trimAl (Capella-Gutiérrez, Silla-Martínez, & Gabaldón, 2009) (ver 1.4; options: -gappyout). The ortholog phylogenies were computed by using IQtree (Nguyen, Schmidt, Von Haeseler, & Minh, 2015) (ver 1.6.6; options: -bb 1000 -alrt 1000). For the CAZyme unclassified HGT orthogroups, the top 2 blast hits sequences from separated Uniprot (bacteria, fungi, land plants and insect) were used to confirm the HGT origin.

**AUTHORS CONTRIBUTION**

IJT and PJC conceived the study. IJT led the study. YCL, TY and PJC sampled the *Aphelenchoides* nematodes. YiCL, HMK, WAL conducted the experiments. CKL analysed the data with input from HHL, IJT and CKL wrote the manuscript with input from YuCL, MRL, TY, TK and PJC.

**DATA AVAILABILITY**

The sequencing data and annotation of six *Aphelenchoides* nematodes are publicly available in NCBI under the BioProject accession PRJNA834627. The accession numbers of individual assemblies are listed in **table S1** and scheduled in the WBPS18 of WormBase Parasite (https://parasite.wormbase.org/index.html). The accession numbers of individual samples are listed in **table S2**. The raw sequences, alignments and phylogenies of the HGT orthologs are available at https://github.com/lihowfun/CladeIV_HGT. The code used to perform this study is deposited at https://github.com/lihowfun/Aphelenchoides.git.

**Acknowledgement**

We thank Pablo Gonzalez for his comments on the preprint version. We thank Christian Rödelsperger and the two reviewers for their comments. We would like to thank National Core Facility for Biopharmaceuticals (NCFB, 111-2740-B-492-001) and National Center for High-performance Computing (NCHC) of National Applied Research Laboratories (NARLabs) of Taiwan for providing computational resources and storage resources. This work was supported by Academia Sinica grant (AS-CDA-107-L01) to IJT.

**References**

Abad, P., Gouzy, J., Aury, J.-M., Castagnone-Sereno, P., Danchin, E. G. J., Deleury, E., … Wincker, P. (2008). Genome sequence of the metazoan plant-parasitic nematode Meloidogyne incognita. *Nature Biotechnology 2008 26:8*, *26*(8), 909–915. doi: 10.1038/nbt.1482

Ali, M. A., Azeem, F., Li, H., & Bohlmann, H. (2017). Smart parasitic nematodes use multifaceted strategies to parasitize plants. *Frontiers in Plant Science*. doi:

661   10.3389/fpls.2017.01699

662 Allen, M. W. (1952). Taxonomic status of the bud and leaf nematodes related to

663   Alphelenchoides fragariae (Ritzema Bos, 1891). *Proceedings of the Helminthological*

664   *Society of Washington*, *19*(2), 108–120.

665 Alonge, M., Soyk, S., Ramakrishnan, S., Wang, X., Goodwin, S., Sedlazeck, F. J., … Schatz, M.

666   C. (2019). RaGOO: Fast and accurate reference-guided scaffolding of draft genomes.

667   *Genome Biology*, *20*(1), 224. doi: 10.1186/s13059-019-1829-6

668 Atanasova, L., Dubey, M., Grujić, M., Gudmundsson, M., Lorenz, C., Sandgren, M., … Karlsson,

669   M. (2018). Evolution and functional characterization of pectate lyase PEL12, a member of a

670   highly expanded Clonostachys rosea polysaccharide lyase 1 family. *BMC Microbiology*,

671   *18*(1), 1–19. doi: 10.1186/S12866-018-1310-9/METRICS

672 Bird, D. M. K., Jones, J. T., Opperman, C. H., Kikuchi, T., & Danchin, E. G. J. (2015).

673   Signatures of adaptation to plant parasitism in nematode genomes. *Parasitology*,

674   *142*(Suppl 1), S71. doi: 10.1017/S0031182013002163

675 Blaxter, M. L., De Ley, P., Garey, J. R., Liu, L. X., Scheldeman, P., Vierstraete, A., … Thomas,

676   W. K. (1998). A molecular evolutionary framework for the phylum Nematoda. *Nature 1998*

677   *392:6671*, *392*(6671), 71–75. doi: 10.1038/32160

678 Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina

679   sequence data. *Bioinformatics*. doi: 10.1093/bioinformatics/btu170

680 Brown, A. M. V., Wasala, S. K., Howe, D. K., Peetz, A. B., Zasada, I. A., & Denver, D. R. (2018).

681   Comparative genomics of wolbachia-cardinium dual endosymbiosis in a plant-parasitic

682   nematode. *Frontiers in Microbiology*, *9*(OCT), 2482. doi: 10.3389/FMICB.2018.02482

683 Buchfink, B., Xie, C., & Huson, D. H. (2014). Fast and sensitive protein alignment using

684   DIAMOND. *Nature Methods*. doi: 10.1038/nmeth.3176

685 Bulletin (2017). PM 7/39 (2) Aphelenchoides besseyi. *47*(3), 384–400. doi: 10.1111/EPP.12432

686 Campoy, E., & González-Martín, A. (2017). The Geography as a Regulator of Genetic Flow and

687   Genetic Structure in Andorra. *Advances in Anthropology*. doi: 10.4236/aa.2017.72008

688 Cantarel, B. L., Korf, I., Robb, S. M. C., Parra, G., Ross, E., Moore, B., … Yandell, M. (2008).

689   MAKER: An easy-to-use annotation pipeline designed for emerging model organism

690   genomes. *Genome Research*. doi: 10.1101/gr.6743907

691 Capella-Gutiérrez, S., Silla-Martínez, J. M., & Gabaldón, T. (2009). trimAl: A tool for automated

692   alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*. doi:

693   10.1093/bioinformatics/btp348

694 Christie, J. R. (1942). A description of Aphelenchoides besseyi n. sp., the summer-dwarf

nematode of strawberries, with comments on the identity of Aphelenchoides subtenuis

(Cobb, 1926) and Aphelen-choides hodsoni Goodey, 1935. *Proceedings of the*

*Helminthological Society of Washington*, *9*, 82–84.

Coconi Linares, N., Dilokpimol, A., Stålbrand, H., Mäkelä, M. R., & de Vries, R. P. (2020).

Recombinant production and characterization of six novel GH27 and GH36 α-

galactosidases from Penicillium subrubescens and their synergism with a commercial

mannanase during the hydrolysis of lignocellulosic biomass. *Bioresource Technology*. doi:

10.1016/j.biortech.2019.122258

Coghlan, A., Tsai, I. J., & Berriman, M. (2018). Creation of a comprehensive repeat library for a

newly sequenced parasitic worm genome. *Protocol Exchange*. doi:

10.1038/protex.2018.054

Cotton, J. A., Lilley, C. J., Jones, L. M., Kikuchi, T., Reid, A. J., Thorpe, P., … Urwin, P. E.

(2014). The genome and life-stage specific transcriptomes of Globodera pallidaelucidate

key aspects of plant parasitism by a cyst nematode. *Genome Biology 2014 15:3*, *15*(3), 1–

17. doi: 10.1186/GB-2014-15-3-R43

Dall, E., & Brandstetter, H. (2016). Structure and function of legumain in health and disease.

*Biochimie*, Vol. 122, pp. 126–150. Elsevier B.V. doi: 10.1016/j.biochi.2015.09.022

Danchin, E. G. J., Guzeeva, E. A., Mantelin, S., Berepiki, A., & Jones, J. T. (2016). Horizontal

Gene Transfer from Bacteria Has Enabled the Plant-Parasitic Nematode Globodera pallida

to Feed on Host-Derived Sucrose. *Molecular Biology and Evolution*, *33*(6), 1571–1579. doi:

10.1093/molbev/msw041

Danchin, E. G. J., Rosso, M. N., Vieira, P., De Almeida-Engler, J., Coutinho, P. M., Henrissat,

B., & Abad, P. (2010). Multiple lateral gene transfers and duplications have promoted plant

parasitism ability in nematodes. *Proceedings of the National Academy of Sciences of the*

*United States of America*, *107*(41), 17651–17656. doi: 10.1073/PNAS.1008486107/

Dayi, M., Sun, S., Maeda, Y., Tanaka, R., Yoshida, A., Tsai, I. J., & Kikuchi, T. (2020).  Nearly

Complete Genome Assembly of the Pinewood Nematode Bursaphelenchus xylophilus

Strain Ka4C1 . *Microbiology Resource Announcements*, *9*(42). doi: 10.1128/mra.01002-20

de la Rosa, P. M. G., Thomson, M., Trivedi, U., Tracey, A., Tandonnet, S., & Blaxter, M. (2021).

A telomere-to-telomere assembly of Oscheius tipulae and the evolution of rhabditid

nematode chromosomes. *G3: Genes, Genomes, Genetics*. doi:

10.1093/G3JOURNAL/JKAA020

Dobin, A., & Gingeras, T. R. (2015). Mapping RNA-seq Reads with STAR. *Current Protocols in*

*Bioinformatics*. doi: 10.1002/0471250953.bi1114s51

Drula, E., Garron, M. L., Dogan, S., Lombard, V., Henrissat, B., & Terrapon, N. (2022). The
carbohydrate-active enzyme database: functions and literature. *Nucleic Acids Research*,
*50*(D1), D571–D577. doi: 10.1093/NAR/GKAB1045

Durand, N. C., Robinson, J. T., Shamim, M. S., Machol, I., Mesirov, J. P., Lander, E. S., &
Aiden, E. L. (2016). Juicebox Provides a Visualization System for Hi-C Contact Maps with
Unlimited Zoom. *Cell Systems*. doi: 10.1016/j.cels.2015.07.012

Edgar, R. C., & Bateman, A. (2010). Search and clustering orders of magnitude faster than
BLAST. *Bioinformatics*, *26*(19), 2460–2461. doi: 10.1093/BIOINFORMATICS/BTQ461

Elling, A. A., Mitreva, M., Gai, X., Martin, J., Recknor, J., Davis, E. L., … Baum, T. J. (2009).
Sequence mining and transcript profiling to explore cyst nematode parasitism. *BMC
Genomics*, *10*. doi: 10.1186/1471-2164-10-58

Emms, D. M., & Kelly, S. (2019). OrthoFinder: phylogenetic orthology inference for comparative
genomics. *Genome Biology*, *20*(1), 238. doi: 10.1186/s13059-019-1832-y

Eves-van den Akker, S., Laetsch, D. R., Thorpe, P., Lilley, C. J., Danchin, E. G. J., Da Rocha,
M., … Jones, J. T. (2016). The genome of the yellow potato cyst nematode, Globodera
rostochiensis, reveals insights into the basis of parasitism and virulence. *Genome Biology
2016 17:1*, *17*(1), 1–23. doi: 10.1186/S13059-016-0985-1

Finn, R. D., Bateman, A., Clements, J., Coggill, P., Eberhardt, R. Y., Eddy, S. R., … Punta, M.
(2014). Pfam: The protein families database. *Nucleic Acids Research*. doi:
10.1093/nar/gkt1223

Finn, R. D., Clements, J., & Eddy, S. R. (2011). HMMER web server: Interactive sequence
similarity searching. *Nucleic Acids Research*. doi: 10.1093/nar/gkr367

Flynn, J. M., Hubley, R., Goubert, C., Rosen, J., Clark, A. G., Feschotte, C., & Smit, A. F.
(2020). RepeatModeler2 for automated genomic discovery of transposable element
families. *Proceedings of the National Academy of Sciences of the United States of
America*. doi: 10.1073/pnas.1921046117

Fortuner, R. (1970). On the Morphology of Aphelenchoides besseyi Christie, 1942 and A.
siddiqii n.sp. (Nematoda, Aphelenchoidea). *Journal of Helminthology*, *44*(2), 141–152. doi:
10.1017/S0022149X00021702

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., … Regev, A.
(2011). Full-length transcriptome assembly from RNA-Seq data without a reference
genome. *Nature Biotechnology*. doi: 10.1038/nbt.1883

Grynberg, P., Togawa, R. C., Freitas, L. D. de, Antonino, J. D., Rancurel, C., Costa, M. M. do
C., … Danchin, E. G. J. (2020). Comparative Genomics Reveals Novel Target Genes

towards Specific Control of Plant-Parasitic Nematodes. *Genes 2020, Vol. 11, Page 1347*, *11*(11), 1347. doi: 10.3390/GENES11111347

Han, Z., Sieriebriennikov, B., Susoy, V., Lo, W.-S., Igreja, C., Dong, C., … Sommer, R. J. (2022). Horizontally Acquired Cellulases Assist the Expansion of Dietary Range in Pristionchus Nematodes. *Molecular Biology and Evolution*, *39*(2). doi: 10.1093/molbev/msab370

Hoff, K. J., Lange, S., Lomsadze, A., Borodovsky, M., & Stanke, M. (2016). BRAKER1: Unsupervised RNA-Seq-based genome annotation with GeneMark-ET and AUGUSTUS. *Bioinformatics*. doi: 10.1093/bioinformatics/btv661

Holm Viborg, A., Terrapon, N., Lombard, V., Michel, G., Czjzek, M., Henrissat, B., & Brumer, H. (2019). *A subfamily roadmap for functional glycogenomics of the evolutionarily diverse Glycoside Hydrolase Family 16 (GH16)*. doi: 10.1074/jbc.RA119.010619

Holterman, M., Karegar, A., Mooijman, P., Van Megen, H., Van Den Elsen, S., Vervoort, M. T. W., … Helder, J. (2017). Disparate gain and loss of parasitic abilities among nematode lineages. *PLoS ONE*. doi: 10.1371/journal.pone.0185445

Holterman, M., Schratzberger, M., & Helder, J. (2019). Nematodes as evolutionary commuters between marine, freshwater and terrestrial habitats. *Biological Journal of the Linnean Society*, *128*(3), 756–767. doi: 10.1093/BIOLINNEAN/BLZ107

Howe, K. L., Bolt, B. J., Shafie, M., Kersey, P., & Berriman, M. (2017). WormBase ParaSite − a comprehensive resource for helminth genomics. *Molecular and Biochemical Parasitology*, *215*, 2. doi: 10.1016/J.MOLBIOPARA.2016.11.005

Jen, F. Y., Tsay, T. T., & Chen, P. (2012). Aphelenchoides bicaudatus from ornamental nurseries in Taiwan and its relationship with some agricultural crops. *Plant Disease*, *96*(12), 1763–1766. doi: 10.1094/PDIS-03-12-0229-RE

Jones, J. T., Furlanetto, C., & Kikuchi, T. (2005). Horizontal gene transfer from bacteria and fungi as a driving force in the evolution of plant parasitism in nematodes. *Nematology*. doi: 10.1163/156854105775142919

Jones, J. T., Haegeman, A., Danchin, E. G. J., Gaur, H. S., Helder, J., Jones, M. G. K., … Perry, R. N. (2013). *Top 10 plant-parasitic nematodes in molecular plant pathology*. doi: 10.1111/mpp.12057

Kikuchi, T, Cotton, J. A., Dalzell, J. J., Hasegawa, K., & Kanzaki, N. (2011). *Genomic Insights into the Origin of Parasitism in the Emerging Plant Pathogen Bursaphelenchus xylophilus Genomic Insights into the Origin of Parasitism in the Emerging Plant Pathogen*. doi: 10.1371/journal.ppat.1002219

Kikuchi, T, Eves-Van Den Akker, S., & Jones, J. T. (2017). *Genome Evolution of Plant-Parasitic*
*Nematodes*. doi: 10.1146/annurev-phyto-080516

Kikuchi, T, Jones, J. T., Aikawa, T., Kosaka, H., & Ogura, N. (2004). A family of glycosyl
hydrolase family 45 cellulases from the pine wood nematode Bursaphelenchus xylophilus.
*FEBS Letters*, *572*(1–3), 201–205. doi: 10.1016/j.febslet.2004.07.039

Koch, B. J., Ryan, J. F., & Baxevanis, A. D. (2012). *The Diversification of the LIM Superclass at*
*the Base of the Metazoa Increased Subcellular Complexity and Promoted Multicellular*
*Specialization*. doi: 10.1371/journal.pone.0033261

Koenning, S. R., Overstreet, C., Noling, J. W., Donald, P. A., Becker, J. O., & Fortnum, B. A.
(1999). Survey of crop losses in response to phytoparasitic nematodes in the United States
for 1994. *Journal of Nematology*, Vol. 31, pp. 587–618. Society of Nematologists.

Kolmogorov, M., Yuan, J., Lin, Y., & Pevzner, P. A. (2019). Assembly of long, error-prone reads
using repeat graphs. *Nature Biotechnology*. doi: 10.1038/s41587-019-0072-8

Korf, I. (2004). Gene finding in novel genomes. *BMC Bioinformatics*, *5*(1), 1–9. doi:
10.1186/1471-2105-5-59/TABLES/4

Koutsovoulos, G. D., Poullet, M., Elashry, A., Kozlowski, D. K. L., Sallet, E., Da Rocha, M., …
Danchin, E. G. J. (2020). Genome assembly and annotation of Meloidogyne enterolobii, an
emerging parthenogenetic root-knot nematode. *Scientific Data 2020 7:1*, *7*(1), 1–13. doi:
10.1038/s41597-020-00666-0

Li, Y., Liu, Z., Liu, C., Shi, Z., Pang, L., Chen, C., … Shen, X.-X. (2022). HGT is widespread in
insects and contributes to male courtship in lepidopterans. *Cell*, *185*(16), 2975-2987.e10.
doi: 10.1016/J.CELL.2022.06.014

Lilley, C. J., Kyndt, T., & Gheysen, G. (2011). Nematode Resistant GM Crops in Industrialised
and Developing Countries. In *Genomics and Molecular Genetics of Plant-Nematode*
*Interactions* (pp. 517–541). Springer Netherlands. doi: 10.1007/978-94-007-0434-3_24

Júnior, T.A.F. Silva, D.R. Negrão, A.T. Itako, and A.C. Maringoni. (2012). Pathogenicity of
Curtobacterium curtobacterium flaccumfaciens pv. flaccumfaciens to several plant species.
*Journal of Plant Pathology* 94, no. 2: 427–30. http://www.jstor.org/stable/45156052.

Ma, J., Wang, S., Zhu, X., Sun, G., Chang, G., Li, L., … Huang, J. (2022). Major episodes of
horizontal gene transfer drove the evolution of land plants. *Molecular Plant*, *15*(5), 857–
871. doi: 10.1016/J.MOLP.2022.02.001

Masonbrink, R., Maier, T. R., Muppirala, U., Seetharam, A. S., Lord, E., Juvale, P. S., … Baum,
T. J. (2019). The genome of the soybean cyst nematode (Heterodera glycines) reveals
complex patterns of duplications involved in the evolution of parasitism genes. *BMC*

*Genomics, 20*(1), 119. doi: 10.1186/s12864-019-5485-8

Mayer, W. E., Schuster, L. N., Bartelmes, G., Dieterich, C., & Sommer, R. J. (2011). Horizontal

gene transfer of microbial cellulases into nematode genomes is associated with functional

assimilation and gene turnover. *BMC Evolutionary Biology, 11*(1), 1–10. doi: 10.1186/1471-

2148-11-13

Medaka: Sequence correction by using ONT research.

https://github.com/nanoporetech/medaka. (2019).

Mitreva, M., Jasmer, D. P., Zarlenga, D. S., Wang, Z., Abubucker, S., Martin, J., … Wilson, R.

K. (2011). The draft genome of the parasitic nematode Trichinella spiralis. *Nature Genetics*.

doi: 10.1038/ng.769

Morais, M. A. B., Coines, J., Domingues, M. N., Pirolla, R. A. S., Tonoli, C. C. C., Santos, C.

R., … Murakami, M. T. (2021). Two distinct catalytic pathways for GH43 xylanolytic

enzymes unveiled by X-ray and QM/MM simulations. *Nature Communications 2021 12:1,*

*12*(1), 1–13. doi: 10.1038/s41467-020-20620-3

Nadachowska-Brzyska, K., Burri, R., Smeds, L., & Ellegren, H. (2016). PSMC analysis of

effective population sizes in molecular ecology and its application to black-and-white

Ficedula flycatchers. *Molecular Ecology, 25*(5), 1058–1072. doi: 10.1111/MEC.13540

Nguyen, L. T., Schmidt, H. A., Von Haeseler, A., & Minh, B. Q. (2015). IQ-TREE: A fast and

effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular*

*Biology and Evolution*. doi: 10.1093/molbev/msu300

Nicol, J. M., Turner, S. J., Coyne, D. L., Nijs, L. den, Hockland, S., & Maafi, Z. T. (2011). Current

Nematode Threats to World Agriculture. *Genomics and Molecular Genetics of Plant-*

*Nematode Interactions*, 21–43. doi: 10.1007/978-94-007-0434-3_2

Oliveira, C. J., Subbotin, S. A., Álvarez-Ortega, S., Desaeger, J., Brito, J. A., Xavier, K. V., …

Inserra, R. N. (2019). Morphological and Molecular Identification of Two Florida

Populations of Foliar Nematodes (Aphelenchoides spp.) Isolated from Strawberry with the

Description of Aphelenchoides pseudogoodeyi sp. N. (Nematoda: Aphelenchoididae) and

notes on their bionomics. *Plant Disease, 103*(11), 2825–2842. doi: 10.1094/PDIS-04-19-

0752-RE

Opperman, C. H., Bird, D. M., Williamson, V. M., Rokhsar, D. S., Burke, M., Cohn, J., …

Windham, E. (2008). Sequence and genetic map of Meloidogyne hapla: A compact

nematode genome for plant parasitism. *Proceedings of the National Academy of Sciences,*

*105*(39), 14802–14807. doi: 10.1073/PNAS.0805946105

Pertea, M., Pertea, G. M., Antonescu, C. M., Chang, T. C., Mendell, J. T., & Salzberg, S. L.

(2015). StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nature Biotechnology*. doi: 10.1038/nbt.3122

Ranallo-Benavidez, T. R., Jaron, K. S., & Schatz, M. C. (2020). GenomeScope 2.0 and Smudgeplot for reference-free profiling of polyploid genomes. *Nature Communications 2020 11:1*, *11*(1), 1–10. doi: 10.1038/s41467-020-14998-3

Rancurel, C., Legrand, L., & Danchin, E. G. J. (2017). Alienness: Rapid detection of candidate horizontal gene transfers across the tree of life. *Genes*, *8*(10). doi: 10.3390/genes8100248

Rybarczyk-Mydłowska, K., Mooyman, P., Van Megen, H., Van Den Elsen, S., Vervoort, M., Veenhuizen, P., … Helder, J. (2012). Small subunit ribosomal DNA-based phylogenetic analysis of foliar nematodes (Aphelenchoides spp.) and their quantitative detection in complex DNA backgrounds. *Phytopathology*. doi: 10.1094/PHYTO-05-12-0114-R

Schiffer, P. H., Danchin, E. G. J., Burnell, A. M., Creevey, C. J., Wong, S., Dix, I., … Blaxter, M. (2019). Signatures of the Evolution of Parthenogenesis and Cryptobiosis in the Genomes of Panagrolaimid Nematodes. *IScience*, *21*, 587–602. doi: 10.1016/J.ISCI.2019.10.039

Shinya, R., Sun, S., Dayi, M., Tsai, I. J., Miyama, A., Chen, A. F., … Sternberg, P. W. (2022). Possible stochastic sex determination in Bursaphelenchus nematodes. *Nature Communications 2022 13:1*, *13*(1), 1–14. doi: 10.1038/s41467-022-30173-2

Siddique, S., Radakovic, Z. S., Hiltl, C., Pellegrin, C., Baum, T. J., Beasley, H., … Akker, S. E. den. (2021). The genome and lifestage-specific transcriptomes of a plant-parasitic nematode and its host reveal susceptibility genes involved in trans-kingdom synthesis of vitamin B5. *BioRxiv*.

Siddique, S., Radakovic, Z. S., Hiltl, C., Pellegrin, C., Baum, T. J., Beasley, H., … Akker, S. E. den. (2022). The genome and lifestage-specific transcriptomes of a plant-parasitic nematode and its host reveal susceptibility genes involved in trans-kingdom synthesis of vitamin B5. *Nature Communications*, *13*(1), 6190. doi: 10.1038/S41467-022-33769-W

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015). BUSCO: Assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*. doi: 10.1093/bioinformatics/btv351

Song, L., Sabunciyan, S., & Florea, L. (2016). CLASS2: Accurate and efficient splice variant annotation from RNA-seq reads. *Nucleic Acids Research*. doi: 10.1093/nar/gkw158

Stamatakis, A. (2014). RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics*. doi: 10.1093/bioinformatics/btu033

Stanke, M., Keller, O., Gunduz, I., Hayes, A., Waack, S., & Morgenstern, B. (2006). AUGUSTUS: A b initio prediction of alternative transcripts. *Nucleic Acids Research*. doi:

10.1093/nar/gkl200

Subbotin, S. A., Oliveira, C. J., Álvarez-Ortega, S., Desaeger, J. A., Crow, W., Overstreet, C., …

Inserra, R. N. (2020). The taxonomic status of Aphelenchoides besseyi Christie, 1942

(Nematoda: Aphelenchoididae) populations from the southeastern USA, and description of

Aphelenchoides pseudobesseyi sp. N. *Nematology*. doi: 10.1163/15685411-bja10048

Szitenberg, A., Salazar-Jaramillo, L., Blok, V. C., Laetsch, D. R., Joseph, S., Williamson, V.

M., … Lunt, D. H. (2017). Comparative genomics of apomictic root-knot nematodes:

Hybridization, ploidy, and dynamic genome change. *Genome Biology and Evolution*, *9*(10),

2844–2861. doi: 10.1093/gbe/evx201

Tarailo-Graovac, M., & Chen, N. (2009). Using RepeatMasker to identify repetitive elements in

genomic sequences. *Current Protocols in Bioinformatics*. doi:

10.1002/0471250953.bi0410s25

Van Megen, H., Van Den Elsen, S., Holterman, M., Karssen, G., Mooyman, P., Bongers, T., …

Helder, J. (2009). A phylogenetic tree of nematodes based on about 1200 full-length small

subunit ribosomal DNA sequences. *Nematology*, *11*(6), 927–950. doi:

10.1163/156854109X456862

Vaser, R., Sović, I., Nagarajan, N., & Šikić, M. (2017). Fast and accurate de novo genome

assembly from long uncorrected reads. *Genome Research*. doi: 10.1101/gr.214270.116

Venturini, L., Caim, S., Kaithakottil, G. G., Mapleson, D. L., & Swarbreck, D. (2018). Leveraging

multiple transcriptome assembly methods for improved gene structure annotation.

*GigaScience*, *7*(8). doi: 10.1093/gigascience/giy093

Walker, B. J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S., … Earl, A. M.

(2014). Pilon: An integrated tool for comprehensive microbial variant detection and genome

assembly improvement. *PLoS ONE*. doi: 10.1371/journal.pone.0112963

Wan, X., Saito, J. A., Hou, S., Geib, S. M., Yuryev, A., Higa, L. M., … Alam, M. (2021). The

Aphelenchus avenae genome highlights evolutionary adaptation to desiccation.

*Communications Biology 2021 4:1*, *4*(1), 1–8. doi: 10.1038/s42003-021-02778-8

Wang, F., Li, D., Wang, Z., Dong, A., Liu, L., Wang, B., … Liu, X. (2014). Transcriptomic

analysis of the rice white tip nematode, phelenchoides besseyi (Nematoda:

Aphelenchoididae). *PLoS ONE*, *9*(3). doi: 10.1371/journal.pone.0091591

Wick, R. R., Judd, L. M., & Holt, K. E. (2019). Performance of neural network basecalling tools

for Oxford Nanopore sequencing. *Genome Biology*. doi: 10.1186/s13059-019-1727-y

Winkler, G. S. (2010). The mammalian anti-proliferative BTG/Tob protein family. *Journal of*

*Cellular Physiology*. doi: 10.1002/jcp.21919

Woodruff, G. C. (2019). Patterns of putative gene loss suggest rampant developmental system
drift in nematodes. *BioRxiv*, p. 627620. bioRxiv. doi: 10.1101/627620

Wu, G. L., Kuo, T. H., Tsay, T. T., Tsai, I. J., & Chen, P. J. (2016). Glycoside hydrolase (gh) 45
and 5 candidate cellulases in aphelenchoides besseyi isolated from bird's-nest fern. *PLoS*
*ONE*, *11*(7). doi: 10.1371/journal.pone.0158663

Wu, S., Gao, S., Wang, S., Meng, J., Wickham, J., Luo, S., … Sun, J. (2020). A Reference
Genome of Bursaphelenchus mucronatus Provides New Resources for Revealing Its
Displacement by Pinewood Nematode. *Genes*, *11*(5), 570. doi: 10.3390/genes11050570

Wu, T. D., & Watanabe, C. K. (2005). GMAP: A genomic mapping and alignment program for
mRNA and EST sequences. *Bioinformatics*. doi: 10.1093/bioinformatics/bti310

Xia, J., Guo, Z., Yang, Z., Han, H., Wang, S., Xu, H., … Zhang, Y. (2021). Whitefly hijacks a
plant detoxification gene that neutralizes plant toxins. *Cell*. doi: 10.1016/j.cell.2021.02.014

Xu, X., Qing, X., Xie, J. L., Yang, F., Peng, Y. L., & Ji, H. L. (2020). Population structure and
species delimitation of rice white tip nematode, Aphelenchoides besseyi (Nematoda:
Aphelenchoididae), in China. *Plant Pathology*, *69*(1), 159–167. doi: 10.1111/ppa.13113

Yokoo, T. (1949). Aphelenchoides oryzae Yokoo n. sp., a Nematode parasitic to Rice Plant.
*Japanese Journal of Phytopathology*, *13*(1–2), 40–43. doi: 10.3186/JJPHYTOPATH.13.40

Yoshida, K., Hasegawa, K., Mochiji, N., & Miwa, J. (2009). Early Embryogenesis and Anterior-
Posterior Axis Formation in the White-Tip Nematode Aphelenchoides besseyi (Nematoda:
Aphelenchoididae). *Journal of Nematology*, *41*(1), 17–22. Retrieved from
http://www.ncbi.nlm.nih.gov/pubmed/22661773

Zarlenga, D. S., Mitreva, M., Thompson, P., Tyagi, R., Tuo, W., & Hoberg, E. P. (2019). A tale of
three kingdoms: members of the Phylum Nematoda independently acquired the detoxifying
enzyme cyanase through horizontal gene transfer from plants and bacteria. *Parasitology*,
*146*(4), 445–452. doi: 10.1017/S0031182018001701

Zheng, J., Peng, D., Chen, L., Liu, H., Chen, F., Xu, M., … Sun, M. (2016). The ditylenchus
destructor genome provides new insights into the evolution of plant parasitic nematodes.
*Proceedings of the Royal Society B: Biological Sciences*. doi: 10.1098/rspb.2016.0942