Association of Whole-Person Eigen-Polygenic Risk Scores with Alzheimer's Disease

Short title: Whole-Person PRS and Alzheimer's Disease

Amin Kharaghani[1,2], Earvin Tio[1,3], Milos Milic[1], David A. Bennett[4], Philip L. De Jager[5], Julie A. Schneider[4], Lei Sun[2,6,*], Daniel Felsky[1,2,3,7,*]

1) Krembil Centre for Neuroinformatics, Centre for Addiction and Mental Health, Toronto, ON, Canada

2) Division of Biostatistics, Dalla Lana School of Public Health, University of Toronto, Toronto, ON, Canada

3) Institute of Medical Science, University of Toronto, Toronto, ON, Canada

4) Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago, IL, USA

5) Centre for Translational and Computational Neuroimmunology, Columbia University Medical Center, New York, NY, USA

6) Department of Statistical Sciences, University of Toronto, Toronto, ON, Canada

7) Department of Psychiatry, University of Toronto, Toronto, ON, Canada

 *Corresponding Authors

Daniel Felsky PhD

Independent Scientist, Krembil Centre for Neuroinformatics, Centre for Addiction and Mental Health

Assistant Professor, Department of Psychiatry and Dalla Lana School of Public Health, University of Toronto

12th Floor, 250 College Street, Toronto ON, M5T 1R8, Canada

Daniel.felsky@camh.ca; dfelsky@gmail.com

www.felskylab.com

(416) 535 8501 x33587

Lei Sun PhD

Professor, Department of Statistical Sciences and Dalla Lana School of Public Health, University of Toronto

700 University Avenue, Toronto ON, M5G 1Z5, Canada

Lei.sun@utoronto.ca; sun@utsat.toronto.edu

https://www.utstat.toronto.edu/sun/#

(416) 978 3452

Keywords: Polygenic risk scores, Alzheimer's disease, WGCNA, eigen-PRS, whole-person

Abstract

Late-Onset Alzheimer's Disease (LOAD) is a heterogeneous neurodegenerative disorder with complex etiology and high heritability. Its multifactorial risk profile and large portions of unexplained heritability suggest the involvement of yet unidentified genetic risk factors. Here we describe the "whole person" genetic risk landscape of polygenic risk scores for 2,218 traits in 2,044 elderly individuals and test if novel eigen-PRSs derived from clustered subnetworks of single-trait PRSs can improve prediction of LOAD diagnosis, rates of cognitive decline, and canonical LOAD neuropathology. Principal component analyses of thousands of PRSs found generally poor global correlation among traits. However, component loadings confirmed covariance of clinically and biologically related traits and diagnoses, with the top PCs representing autoimmune traits, cardiovascular traits, and general pain medication prescriptions, depending on the PRS variant inclusion threshold. Network analyses revealed distinct clusters of PRSs with clinical and biological interpretability. Novel eigen-PRSs (ePRS) derived from these clusters were significantly associated with LOAD-related phenotypes and improved predictive model performance over the state-of-the-art LOAD PRS alone. Notably, an ePRS representing clusters of traits related to cholesterol levels was able to improve variance explained in a model of brain-wide beta-amyloid burden by 1.7% (likelihood ratio test $p=9.02 \times 10^{-7}$). While many associations of ePRS with LOAD phenotypes were eliminated by the removal of *APOE*-proximal loci, some modules (e.g. retinal defects, acidosis, colon health, ischaemic heart disease) showed associations at an unadjusted type I error rate. Our approach reveals new relationships between genetic risk for vascular, inflammatory, and other age-related traits and offers improvements over the existing single-trait PRS approach to capturing heritable risk for cognitive decline and beta-amyloid accumulation. Our results are catalogued for the scientific community, to aid in the generation of new hypotheses based on our maps of clustered PRSs and associations with LOAD-related phenotypes.

Introduction

Late-Onset Alzheimer's disease (LOAD) is the most common cause of dementia in older adults [1]. It is a highly polygenic and multifactorial disease [2] with heritability estimated to be between 58% to 79% [3]. Despite the identification of its canonical pathologies decades ago, no effective disease-modifying treatments are available. This is likely a result of the etiological complexity of the illness and challenges associated with studying causality for a cascade of events that precede the emergence of symptoms by decades. High-throughput genetic approaches and the ongoing era of genome-wide association study (GWAS) discovery have provided clues into the complex, potentially causal mechanisms of LOAD [4] and highlighted disturbances across multiple processes and systems. The most recent genetic evidence points toward inflammatory, neurotrophic, metabolic, and cellular senescence mechanisms [5–9], and genetic correlations have been observed between LOAD and a multitude of physical, psychiatric, and other neurological illnesses [8,10].

From an epidemiological perspective, factors affecting risk for and progression of LOAD are highly varied. A systematic review [11] on LOAD risk factors highlighted statin use, alcohol consumption, diet, educational attainment, physical and cognitive activity, head injury (in males only), age, diabetes, estrogen therapy, smoking, and social engagement as major risk factors (in addition to genetic risk and protective factors, such as *APOE* ε4 and ε2, respectively). A recent review [12] of factors specifically affecting the progression of LOAD identified six with cohesive evidence (malnutrition, genetic variants, altered gene regulation, baseline cognitive level, neuropsychiatric symptoms, and extrapyramidal signs), though nine more showed conflicting associations in the literature. Additional factors with less clear, potentially bidirectional, associations with LOAD include a large host of often comorbid conditions including sleep disturbances, depression, and gastrointestinal, cardiovascular, and other inflammatory diseases [13,14]. Adding to the complexity of contributing risk factors, the age at which some emerge can affect an individual's LOAD risk profile. For example, obesity [15], hypertension [16], high cholesterol [17] and statin use [18] in mid-life are associated with an increased risk of dementia later in life. Importantly, these lifestyle, environmental, clinical, and behavioural factors are themselves heritable, and exert their neuropathological effects via multiple biochemical cascades which may be unique to each person [19]. This calls for a more sophisticated multivariate, multi-trait analyses of the

genetic component of risk for predicting LOAD, incorporating the whole-person scope of

potential phenotypic contributions.

To our knowledge, there have only been a small handful of studies attempting to unravel the

phenome-wide genetic risk landscape of LOAD using polygenic approaches. One group used

results from the UK Biobank PheWAS [20] to identify patterns of correlated phenotypes and

individual genetic variants across thousands of human traits, including LOAD, using singular

value decomposition applied to summary statistics [21]. Similar work, focusing specifically

on LOAD, proposed another summary statistics-focused method [22], identifying 48 PRSs

each with significant association with LOAD genetic risk. Other analyses from the

Alzheimer's Disease Genomics Consortium (ADGC) have examined multifactorial risk for

LOAD by calculating PRSs for 22 candidate modifiable risk factors, finding only an

educational attainment PRS significantly associated with LOAD [23]. Here, we build on

previous efforts in four important ways: 1) by performing dimensionality reduction on

phenome-wide PRSs calculated on a population of elderly, rather than operating on summary

statistics alone, 2) by examining how broad patterns of phenotypic overlap across PRSs are

affected by the adjustment of SNP inclusion thresholds (including sensitivity analyses with

the inclusion and exclusion of the *APOE* and the MHC genetic regions), 3) by deriving and

testing novel composite eigen-PRSs (ePRSs) through the application of the weighted gene

correlation network analysis (WGCNA), and 4) by evaluating the added value of these novel

ePRSs explicitly against the standard LOAD-specific PRS [24] for association with

longitudinal cognitive decline and post-mortem neuropathological burden.

We hypothesize that a set of phenome-wide PRSs will yield discrete intra-correlated clusters

of related traits and that patterns of correlation and clustering will change as a function of

SNP inclusion criteria, possibly reflecting between-trait differences in polygenicity. We also

hypothesize that the proposed composite ePRSs, derived from the WGCNA algorithm, will

capture genetic risk from proximal, possibly overlapping etiological risk factors for LOAD

thus improving the prediction of LOAD and LOAD-related phenotypes over the single-PRS

method.

Materials and Methods

The Religious Orders Study (ROS) and Memory Aging Project (MAP) Participants

ROS and MAP are two longitudinal cohort studies based out of the Rush Alzheimer's Disease Centre (RADC) in Chicago (IL) [25] designed with harmonized protocols to be analyzed together. Participants are older, recruited free of known dementia, and agree to annual detailed clinical evaluation, cognitive testing, and organ donation at the time of death. A total of 2,044 participants were determined to be of European ancestry and included in our analyses (see **Supplementary Methods**). All phenotypic data are available via controlled access through the RADC Resource Sharing Hub (www.radc.rush.edu) and genomics data are available through Synapse (https://www.synapse.org/#!Synapse:syn3219045). The Institutional Review Board of Rush University Medical Center approved each study, and all subjects provided informed, written consent and signed an Anatomical Gift Act. All participants signed a repository consent for resource sharing.

Table 1 Demographic summary of Religious Orders Study (ROS) and Memory Aging Project (MAP) participants included in analysis

Total (n=2,044)	‡Autopsied (n=1,443)				Not deceased (n=601)			
	CN (n=473)	MCI (n=345)	LOAD (n=599)	Other (n=26)	CN (n=442)	MCI (n=111)	LOAD (n=47)	Other (n=1)
Sex (M/F)	167/306	128/217	184/415	9/17	98/344	27/84	7/40	0/1
Age at study entry (mean (SD))	78.77 (6.96)	81.16 (6.36)	81.69 (6.74)	78.48 (7.82)	73.73 (6.92)	76.69 (7.33)	78.17 (8.48)	87.24
Age at death (mean (SD))	87.26 (6.84)	89.23 (6.31)	91.22 (5.89)	88.01 (6.65)	--	--	--	--
§Postmortem interval (mean (SD))	9.46 (8.93)	9.24 (7.56)	8.52 (6.55)	10.5 (10.2)	--	--	--	--
Education, years (mean (SD))	16.39 (3.70)	16.12 (3.45)	16.19 (3.63)	17.04 (3.90)	16.70 (3.39)	15.68 (3.32)	16.23 (3.10)	16.00
†Mean LOAD PRS (mean (SD))	-0.15 (0.79)	-0.03 (0.78)	0.12 (0.80)	0.00 (0.86)	-0.03 (0.80)	0.16 (0.88)	0.13 (0.80)	0.13

‡191 deceased participants with genotype information did not have quantitative neuropathological data available at time of analysis. §Postmortem interval statistics were calculated on 1,252 people with complete data. †LOAD PRS values shown are averaged over all 15 α-value thresholds within each subgroup (calculated on the whole genome, without removing any regions *a priori*). CN = no cognitive

146 impairment; LOAD = late-onset Alzheimer's disease; MCI = mild cognitive impairment; Other = cause of
147 dementia not likely Alzheimer's disease (e.g. vascular dementia); SD = standard deviation.

Clinical diagnosis of LOAD

For all subjects, available clinical data were reviewed at the time of death by a neurologist with
expertise in dementia. Summary diagnoses were made blinded to all post-mortem data and case
conferences with one or more neurologists and a neuropsychologist were used to achieve
consensus when required. Diagnoses fell into one of six possible categories: no cognitive
impairment (NCI), mild cognitive impairment (MCI) with or without secondary cause of
impairment, probable late onset Alzheimer's disease (LOAD; NINCDS criteria [26]), possible
LOAD (with additional cause of impairment), and other primary cause of dementia [27]. For
the purposes of our case-control modelling of consensus diagnosis at time of death, we
considered only probable LOAD and NCI participants; all other analyses of neuropathological
and cognitive outcomes included all participants with available data.

Assessment of longitudinal cognitive decline

At each annual evaluation, all subjects were administered 21 cognitive tests of which 19 were
used in both ROS and MAP [28]. The raw scores from these 19 tests were standardized to z-
scores across the cohort and averaged to yield the global cognitive function summary score at
each study visit. Slopes of change in global cognitive performance over time (cognitive
decline) were calculated per subject using linear mixed modelling, with a random effect for
study visit, controlling for age at baseline, as previously described [29].

Assessment of postmortem beta-amyloid and tau neuropathology

Board-certified neuropathologists examined all brains without prior knowledge of clinical or
demographic information. Brain removal and fixing procedures have been detailed extensively
[30]. Complete data from neuropathological examinations were available for up to 1,252
participants at time of study. Immunohistochemistry was used to quantify total beta-amyloid
and paired helical filament tau (PHFtau) across eight brain regions (hippocampus, entorhinal
cortex, midfrontal cortex, inferior temporal, angular gyrus, calcarine cortex, cingulate cortex,
frontal cortex), as described previously [31]. The square root transformed brain-wide averages
of beta-amyloid and tau pathologies were used as in previous analyses of these data [32].
Figure S1 illustrates the distribution of the pathological measurements alongside global
cognitive decline of the ROS/MAP participants.

Genotyping and imputation

Genotype data were available for 2,067 ROS/MAP participants, between two batches: n_{batch1}=1,686 genotyped using the Affymetrix GeneChip 6.0 and n_{batch2}=381 genotyped using the Illumina OmniQuad Express platform; batch was included as a covariate in our analyses. Details of raw genotype quality control (QC) have been previously described [33]. Briefly, prior to submission for imputation, genotypes were preprocessed using the TOPMed Imputation Server-recommended data preparation pipeline (https://topmedimpute.readthedocs.io/en/latest/prepare-your-data.html). Each batch was imputed separately using the TOPMed Imputation Server (TOPMed reference r^2)[34], including Eagle (v2.4) for allelic phasing and Minimac4 (v1.5.7) for imputation. Imputed output data from the TOPMed server for each batch were filtered for imputation quality (removing SNPs with r^2<0.8) before merging and mapping to rsIDs (dbSNP build 155). This resulted in a set of 9,329,439 high-quality, bi-allelic autosomal SNPs. Additionally, we verified the European ancestral background of genotyped participants by mapping the multidimensional scaling (MDS) plots of the ROS/MAP SNP genotypes against the 1000 Genome project reference populations (phase 3) [35] (details in **Supplementary Methods** and **Figure S2**). Prior to our PRS calculation, additional sample and variant quality controls were performed, removing SNPs and subjects according to the following criteria:

- SNPs with minor allele frequency (MAF) < 0.01 (1,549,210 variants)
- SNPs with Hardy-Weinberg Equilibrium (HWE) *p*-value < 1×10^{-6} (132 variants)
- SNPs with missingness rate > 0.01 (978 variants)
- Individuals with no cognitive or neuropathological phenotype data available were removed (7 individuals)
- Individuals with higher-than-expected rates of SNP heterozygosity within 3 standard deviations of the mean (15 individuals)
- Individuals with non-white European ancestry were removed (1 individual)

Following QC, a total of 2,044 individuals and 7,723,915 variants remained for analysis.

Selection of GWAS summary statistics from the UK Biobank

For phenome-wide PRS calculations, we used GWAS summary statistics derived from the UK Biobank, a large-scale population-based study including approximately 500,000 individuals [36]. The Pan-UK Biobank Consortium (https://pan.ukbb.broadinstitute.org/) has

recently conducted GWAS for all available study phenotypes with sufficient statistical power.

Out of a total of 7,221 phenotypes with GWAS performed in the European ancestry sample

(which matches our sample of genetically verified European-ancestry ROS/MAP

participants), we selected 2,218 phenotypes that had heritability estimates greater than or

equal to 5% and were sex non-specific. The heritability scale of each phenotype was provided

by Pan-UKBB consortium (calculated using SAIGE [37]), and selected sex non-specific

phenotypes using their provided meta-data ("pheno_sex" column indicating "both").

Additional manual selection was performed to remove phenotypes with no cases for either

sex (indicated by columns "n_cases_full_cohort_females" or "n_cases_full_cohort_males"),

indicating geographical or administrative information, or duplicate indications or codings of

the same underlying measurement. The full list of selected phenotypes for our study are

provided in **Table S1**, and further details of selection criteria are outlined in **Supplementary**

Methods. For calculating a benchmark PRS for LOAD, we also downloaded summary

statistics from the latest LOAD GWAS, published by Bellenguez et al. [38]

Calculation and compilation of whole-person PRS matrices

PRSice (v2.3.3) [39] was used to generate PRSs across all traits at 15 different SNP inclusion

p-value thresholds (α-values) including 1 (all SNPs), 0.1, 0.05, 0.01, 5×10^{-3}, 1×10^{-3},

5×10^{-4}, 1×10^{-4}, 5×10^{-5}, 1×10^{-5}, 5×10^{-6}, 1×10^{-6}, 5×10^{-7}, 1×10^{-7}, and

5×10^{-8} (only genome-wide significant SNPs).

For each α-value threshold, we obtained a set of PRSs for up to 2,218 traits (fewer traits are

included at more stringent thresholds, as they may not have identified any SNPs reaching that

threshold). For PRS calculations, we used the default "average" method of PRSice [40]. To

account for patterns of linkage disequilibrium (LD), we applied a 500Kb sliding window and

r^2 threshold of 0.2 during variant clumping, as in previous work [41]. To control for fine

population structure [42], the first ten principal components of genotype data and a variable

representing genotype batch were regressed out of each score independently. We then

conducted further QC, removing any PRS with fewer than five contributing SNPs. We also

identified any pairs of PRSs with extremely highly correlation (Pearson correlation > 0.95)

and retained only one (phenotypes with ICD10 code were preferred otherwise selected

randomly). Finally, each PRS was standardized (mean subtracted and then divided by

standard deviation) to obtain z-scores.

This resulted in 15 sets (for the 15 different α-value thresholds) of PRS "matrices" for

downstream analysis. Each PRS "matrix" is a $n \times k$ matrix of phenome-wide PRSs

calculated across traits, where $n = 2,044$ is the number of participants after QC, and $k \leq$

2,218 is the number of traits. The value of k varies across the 15 sets (outlined in **Table S2**),

because at a given α-value threshold, some PRSs may be constructed from fewer than five

SNPs thus removed as part of the QC.

Notably, previous work has observed a profound influence of variants in the *APOE* region of

chromosome 19 on LOAD phenotypes in PRS analyses [43]. Additionally, many psychiatric

and neurological disorders are strongly associated with genetic variants in the major

histocompatibility complex (MHC) region of chromosome 6 [44], which is also notoriously

challenging to impute due to complex LD [45]. To assess the influence of these regions on

the clustering of PRSs and resulting associations with LOAD-related phenotypes, we

additionally calculated all the 15 sets of PRSs excluding a) the MHC region (GRCh37:

chr6:25477797-36448354) [21] (55,204 variants), and b) the *APOE* region (GRCh37:

chr19:45300000-45500000) [46] (622 variants).

This resulted in a total of $15 \times 4 = 60$ sets of phenome-wide PRS matrices, for the 15

different α-value thresholds and four different *APOE* and/or *MHC* inclusion and exclusion

criteria: **1)** including all genotyped and imputed variants genome-wide (primary analysis;

denoted "ALL"), **2)** removing only *APOE* SNPs (denoted "APOE-"), **3)** removing only MHC

SNPs (denoted "MHC-"), and **4)** removing both *APOE* and MHC SNPs (denoted

"MHC/APOE-"). For clarity, we denote these PRS matrices as $\Pi_{subscript}$, where the

subscript indicates whether the matrix was calculated on the whole genome or with indicated

regions excluded (i.e. ALL, MHC-, APOE-, MHC/APOE-) and the α-value threshold used

for SNP inclusion (e.g. 1×10^{-4}, 5×10^{-8}). For example, $\Pi_{MHC-,10^{-8}}$ is for the matrix

when a PRS was calculated without SNPs in the MHC region and at an α-value threshold of

1×10^{-8}. **Table S2** outlines the dimension of each PRS matrix after QC.

Principal component analysis (PCA) of whole-person PRS matrices

270   PCA is a multivariate dimension-deduction technique that can describe the covariance among

271   a set of variables (in our case, PRSs) [47]. All PCA analyses were performed using R (v4.2.0)

(https://www.r-project.org/). PCA was first applied to each of the 60 PRS matrices separately to identify the major variance components of phenome-wide genetic risk and describe the contributors to each top principal component. This exploratory PCA analysis served three purposes: 1) as a cursory test of accuracy for the large sets of PRS calculations and post-processing (do PCs load onto known epidemiologically and biologically related traits?), 2) to inform expectations for cohesion in downstream clustering analyses, and 3) to provide confirmation of observations made previously based on GWAS summary statistics that the majority of phenome-wide variance in genetic risk for human traits is explained by genetic risk for metabolic and cardiovascular health-related measures [21].

Clustering of PRS matrices and calculation of eigen-PRSs (ePRS) using WGCNA

To identify meaningful, discrete clusters of correlated PRSs across the phenome for ePRS construction, we used the weighted gene co-expression network analysis (WGCNA) framework [48],which leverages the network properties of biological phenomena to cluster input measures. Although WGCNA was originally developed for use in gene expression experiments to identify biologically coherent co-expressed gene modules, the principles on which it operates are not unique to such measures; WGCNA has been used to build epigenetic networks [49] and to subtype patient populations [50]. Notably, the original authors of WGCNA have also adapted the method for network-based analyses of GWAS summary statistics (the so-called weighted SNP correlation network analysis, or WSCNA [51]), which aims to identify clusters of SNPs with similar effects on a given phenotype after controlling for LD. In contrast, we used the WGCNA framework for clustering PRSs across a large number of phenotypes, rather than clustering SNP effects across the genome for a single phenotype.

Extensive details on the WGCNA/WSCNA (v1.71) pipeline have been published previously [48,51]. In our analysis, for each PRS matrix, pairwise Pearson correlations were calculated among all PRSs. For the construction of a weighted network, these PRS-PRS correlations were raised to a soft-thresholding power (β), determined empirically using criteria that encourage approximate scale-free topology of the resulting network (via the WGCNA package function *pickSoftThreshold*). We chose to calculate an *unsigned* network, meaning

that adjacency was based off of the absolute value of correlation, rather than discarding

negative correlations. This was important, as we calculated PRS across thousands of

phenotypes, among which both negative and positive correlations would be of interest.

To cluster this network, a topological overlap matrix (TOM) was calculated and used to

define dissimilarity for hierarchical clustering. The resulting dendrogram was then clustered

using dynamic tree cutting [52] which applies a variable-height tree cut algorithm to

maximize the capture of meaningful branch structures (i.e. PRS modules). WGCNA was run

using the following additional parameters for detection of PRS modules: module detection

cut-height of 0.999, minimum module size of 10, and high branch split sensitivity

(deepSplit=4).

Our soft-thresholding power selection varied across PRS matrices (refer to **Figure S3**),

depending mostly on the SNP inclusion α-value threshold, and therefore the number of PRSs

included in a given matrix. For each matrix, the lowest value of β was chosen at which the

scale-free topology model fit (R^2) reached above 0.8 (resulting β values ranged from 1 to 6)

to satisfy a scale-free topological network. We also chose a low minKMEtoStay parameter

(0.01), which governs the behaviour of the dendrogram clustering algorithm and results in the

inclusion of nearly all input PRS into some module (rather than excluding the many lowly

correlated PRSs from membership in any module). We did this to allow the ePRS pipeline to

capitalize on even small genetic influences of weakly-correlated PRSs within clusters, rather

than eliminating these subtle contributions entirely. Following the identification of PRS

modules, PCA is applied to each module separately, with the first principal component

representing that module's ePRS. These ePRSs were used for downstream modelling of

LOAD phenotypes.

Association analysis of ePRSs with LOAD phenotypes

For each iteration of our PRS analysis (i.e. for each matrix calculated using a different SNP

inclusion α-value threshold and including/excluding *APOE* and/or *MHC*), we used

multivariate regression models to identify the association of derived ePRSs with the

following LOAD-related variables: longitudinal global cognitive decline slope, consensus

clinical diagnosis of LOAD at time of death, and brain-wide post-mortem levels of β-amyloid

(Aβ) and PHFtau neuropathology. Among the four outcomes, LOAD diagnosis is a binary

outcome and was analyzed using logistic regression, with the remaining three continuous traits analyzed using linear models.

For LOAD diagnosis and cognitive decline, regression covariates included biological sex, age at death (age at baseline for decline), and educational attainment. For neuropathological outcomes, educational attainment was replaced with postmortem interval (PMI). False discovery rate (FDR) correction (i.e. q-value [53]) was used to mitigate multiple testing concerns among all ePRS derived from a given matrix; this matrix-specific FDR correction can be thought as a stratified FDR control [54]. For direct comparison with the benchmark PRS from the latest and most powerful GWAS [38] of LOAD, ePRSs demonstrating significant association with at least one of the LOAD phenotypes (q-value<0.05) were carried forward for further analysis.

Evaluating performance of ePRSs against LOAD PRS benchmark

First, we optimized a single-trait PRS for LOAD (PRS_{LOAD}) in the ROS/MAP sample. This was achieved using the same PRSice-based method described previously, calculating PRS_{LOAD} at the 15 different SNP inclusion α-value thresholds, and selecting the top associated (q-value $\leq$ 0.05) score for each AD-related phenotype. We note that this model selection approach is expected to result in a degree of overfitting for our benchmark PRS_{LOAD}, rendering subsequent estimates of ePRS performance gains naturally conservative.

To compare the performance of selected LOAD phenotype-associated ePRSs with our benchmark PRS_{LOAD}, we first built two sets of baseline regression models (linear for continuous and logistic for binary outcomes) to establish the performance of covariates alone (Model 1) and covariates + PRS_{LOAD} (Model 2) on our clinical and neuropathological outcomes (Y):

Model 1. Covariates only:
(Clinical Y) $Y \sim covariates\ (sex + age\ at\ baseline +\ education\ years)$
(Neuropathology Y) $Y \sim covariates\ (sex + age\ at\ death + PMI)$

Model 2. Covariates and PRS_{LOAD}:

368 $$Y \sim covariates + PRS_{LOAD}$$
369
370 Then, for each ePRS significantly associated with at least one LOAD-related outcome in our
371 preceding analyses, we built a third regression model (Model 3) including covariates,
372 PRS_{LOAD}, and the selected ePRS:
373
374 **Model 3.** Covariates, PRS_{LOAD}, and ePRS:
375 $$Y \sim covariates + PRS_{LOAD} + ePRS$$
376
The likelihood ratio test was then used, comparing model 2 with model 3, to assess whether
the ePRS significantly improved model fit over PRS_{LOAD} for a given outcome. The difference
in model performance was reported as a ΔAUC for logistic and ΔR^2 for linear models, again
comparing models 2 and 3. As each significant ePRS was selected based on association with
at least one LOAD-related outcome, we mitigated concerns of overfitting by performing
0.632+ bootstrapping [55] to derive model performance estimates.
383
384

Results

Principal components analysis of whole-person PRSs in elderly

Table 1 summarizes demographic information for the individuals included in our study. We first sought to explore the covariance structure of PRSs calculated in our sample for over 2,000 diverse phenotypes when no genomic regions were excluded from our calculations ($\Pi_{ALL,\alpha}$). **Figure 1** illustrates the overall variation explained by each PC for $\Pi_{ALL,\alpha}$ (refer to **Figures S4-S6** for the other PRS matrices), and the number of PCs explaining 80% of the total variation of the corresponding matrix. When considering PRSs calculated using a genome-wide significant α-value SNP inclusion threshold of 5×10^{-8} (349 PRSs), the first principal component (PC) explained 7.38% of the total variance in all PRSs. In evaluating variable contributions to this PC, we found that the strongest contributing PRSs were those for autoimmune traits, such as chronic hepatitis (total contribution: 3.16%), diffuse diseases of connective tissue (total contribution: 2.98%), Sicca syndrome (total contribution: 2.68%) (**Figure S7**). The LOAD PRS (PRS_{LOAD}) was loaded onto PC28 (total contribution of 2.1% to this PC) at this threshold along with arthrosis-related traits, family history of AD, delirium, dementia, and education, shown in **Figure S7**.

Across all the 15 α-value thresholds, this pattern was fairly consistent, with autoimmune conditions representing the majority of variability in the top PCs. In addition, there was a consistency in the top contributing PRSs to PC1 across all 15 α-value thresholds. At the highest α-value threshold (1.0), PC1 (explaining 1.90% of the variance) was dominated by opioid and generic over-the-counter pain relief medications. Further, PC1 of matrices calculated with somewhat lower α-value threshold values of 0.1, 0.05, and 0.01 (explaining 1.73%, 1.58% and 1.26% of the variance respectively) loaded most prominently onto body mass index (BMI) and weight, along with hypertension and vascular problems. At even lower thresholds of 0.005, 0.001, and 5×10^{-4} (explaining 1.25%, 1.63% and 1.80% of the variance respectively) PC1 was primarily represented by PRSs for intestinal malabsorption and hypertension. The PRS for thyrotoxicosis hyperthyroidism was the strongest contributor to PC1 at thresholds of 1×10^{-4}, 5×10^{-5}, 1×10^{-5} and 5×10^{-6} (explaining 2.08%, 2.19%, 2.22% and 2.37% of the variance respectively). PRS of Chronic hepatitis was heavily loaded on to PC1 at α-value threshold of 1×10^{-6}, 5×10^{-7}, 1×10^{-5} (explaining 5.28% and 6.11% of variance respectively), and lastly usage of insulin loaded onto PC1 at α-value

threshold 1×10^{-7} explaining 6.70% of the total variance. Detailed breakdowns of PC1 and

PC2 at α-value threshold of 0.001 are shown in **Figure S8**.

When the MHC region was removed prior to PRS calculations ($\Pi_{MHC-,\alpha}$), we found that the

top three contributors to PC1 of PRS matrices at lower α-value thresholds (i.e. 5×10^{-8},

1×10^{-7}, 5×10^{-7}, 1×10^{-6}, 5×10^{-6} and 1×10^{-5} explaining 5.17%, 5.03%, 4.78%,

3.81%, 1.21% and 1.02% of the total variance respectively) were mainly influenced by PRSs

such as presence of cardiac and vascular implants and grafts, coronary artery bypass grafts,

heart attacks, and other heart related problems. PRSs of mean arterial pressure, systolic blood

pressure, high blood pressure and hypertension were loaded as top contributors to PC1 at α-

value threshold of 5×10^{-5}, 1×10^{-4}, 5×10^{-4}, 0.001 and 0.005 (explaining 0.99%,

0.98%, 1.07%, 1.07% and 1.15% of the total variance respectively). At higher α-value

thresholds (α-value of 0.01, 0.05, 0.1), the top PRS contributors to PC1 (explaining 1.25%,

1.57% and 1.71% of the variance respectively) were BMI along with hypertension and other

vascular problems. Lastly, PC1 at α-value threshold of 1 was similar to the PC1 calculated

with the whole genome, and it consisted of PRSs of opioid use, and other over the counter

pain killers. More detailed description of PC1 and PC2 at α-value threshold of 0.001 are

shown in **Figure S9**.

A few of the PCs were heavily influenced by LOAD, dementia, and family history of AD

with other known AD-related traits such as high cholesterol, usage of statin, apolipoprotein

biomarker, directly assessed levels of LDL cholesterol (LDL) and delirium. Though these

PCs were mainly present in $\Pi_{ALL,\alpha}$ and $\Pi_{MHC-,\alpha}$. Additionally, the contribution of PRS of

LOAD was low compared to the top contributing PRSs to these PCs. Interestingly, PRS of

LOAD was not present in AD-related PRSs PC when the MHC region was not included.

Figures S7, S10 and S11 illustrate a few selected LOAD related PCs from $\Pi_{ALL,\alpha}$ and

$\Pi_{MHC-,\alpha}$. In the absence of the *APOE* region ($\Pi_{APOE-,\alpha}$), there was little notable change in

the variability explained by PCs or in the structure of those PCs (when compared to $\Pi_{ALL,\alpha}$).

This was true across all α-value thresholds, with a few exceptions. PC1 at α-value thresholds

of 5×10^{-8} (350 PRSs, explaining 7.32% of the total variations) influenced autoimmune

traits like chronic hepatitis (contribution of 3.16%), Thyrotoxicosis hyperthyroidism

(contribution of 2.99%), and Sicca syndrome (contribution of 2.78%). In the absence of both

influential genomic loci ($\Pi_{MHC/APOE-,\alpha}$) PC1 at α-value thresholds of 5×10^{-8} (315 PRSs,

explaining 5.12% of the total variance) was mostly influenced by coronary artery bypass

grafts (contribution of 4.96%). Lastly, the role of PRS_{LOAD} in contributing to the PCs from

both $\Pi_{APOE-,\alpha}$ and $\Pi_{MHC/APOE-,\alpha}$ dropped notably.

Figure 1. Principal component analysis of whole-person PRS matrices.

Beeswarm plot summarizing the results from PCA on Π_{ALL}. Each point represents a PC derived from each PRS matrix, and the colour of each point represents the PC number. The majority of PCs explain less than 1% of the total variation of their corresponding matrices (y-axis). Additionally, a large number of PCs are required to explain 80% of the total variation of the original matrix. However, at lower α-value thresholds (x-axis), we notice more PCs explain more than 5% of the total variation of their original matrix, largely due to the smaller number of PRSs at lower α-value thresholds. The number one contributing PRS of each PC, with their contributing percentage to the component, is labelled for the first PC of each matrix. **Figures S4-S6** show result summaries for the PCA results on other PRS matrices ($\Pi_{MHC-}, \Pi_{APOE}, \Pi_{MHC/APOE-}$).

Clustering of PRS matrices using WGCNA

Following descriptive PCA and the assessment of phenome-wide variance in PRSs at different α-value thresholds, we sought to group PRSs into discrete clusters based on their network-based similarity using WGCNA. Unlike PCA, WGCNA provides discrete, mutually exclusive groupings of PRS subnetworks for further analysis. Beginning with the entire genome and an α-value threshold of 0.05 ($\Pi_{ALL,0.05}$), which includes a large number of SNPs per phenotype, the WGCNA approach yielded 33 PRS clusters (or modules). As expected, biologically and clinically related traits clustered together. The number of modules identified, differed as a function of α-value threshold (12 modules from $\Pi_{ALL,5\times10^{-8}}$, 15 modules from $\Pi_{ALL,5\times10^{-6}}$, 15 modules from $\Pi_{ALL,1\times10^{-6}}$, 35 modules from $\Pi_{ALL,1}$, the rest are shown in **Table S3**), though module identities remained largely cohesive. Hub analysis was used to reveal the PRSs that were most interconnected within each module; hub PRSs at different thresholds included those for weight in the red module ($\Pi_{ALL,5\times10^{-8}}$, consisting of 22 PRSs), rheumatoid arthritis in the blue module ($\Pi_{ALL,5\times10^{-5}}$, consisting of 209 PRSs), and malignant melanoma in the black module ($\Pi_{ALL,1\times10^{-5}}$, consisting of 25 PRSs), and cancer of bronchus lung in the brown module ($\Pi_{ALL,0.01}$, consisting of 103 PRSs). Note that module color labels are arbitrarily assigned by WGCNA and not linked to particular sets or sizes of PRS (i.e. they are not preserved between runs of WGCNA). Some modules, such as the weight module, were largely preserved across nearly all α-value (except for 1×10^{-6} and 5×10^{-7}) thresholds (**Figures S12-S15**); this module was clustered with PRSs related to body mass, bone density, and coronary and heart diseases. Conversely, chronic ischemic heart disease in the yellow module ($\Pi_{ALL,5e-05}$, consisting of 21 PRSs) clustered along with other heart or coronary complications such as heart attack (myocardial infarction) and ischemic heart disease. Notably, this module was conserved across α-value thresholds of 0.01, 0.005, 5×10^{-5} and 5×10^{-7}, and was correlated with modules that included PRS$_{LOAD}$ and PRSs for family history of AD and all-cause dementia. Of note, PRS$_{LOAD}$ consistently clustered within the rheumatoid arthritis module in $\Pi_{ALL,0.001}$, $\Pi_{ALL,0.0005}$, $\Pi_{ALL,0.0001}$, $\Pi_{APOE-,0.001}$, $\Pi_{APOE-,0.0005}$, $\Pi_{APOE-,0.0001}$, $\Pi_{APOE-,5\times10^{-5}}$. **Figure S16** outlines the correlational structure of two selected modules (weight and high cholesterol).

Focusing specifically on the PRS$_{LOAD}$, we found it clustered within the pink module from $\Pi_{ALL,5e-08}$ (consisting of 19 PRSs), red module from $\Pi_{ALL,1e-07}$ (consisting of 23 PRSs) and

$\Pi_{ALL,5e-05}$ (consisting of 22 PRSs), brown module from $\Pi_{ALL,1e-06}$ (consisting of 68 PRSs)

and $\Pi_{ALL,5e-06}$ (consisting of 85 PRSs) and $\Pi_{ALL,1e-05}$ (consisting of 77 PRSs) along with

PRSs for all-cause dementia, family history of AD or AD-dementia, delirium, Apolipoprotein

levels, cholesterol levels, presence of cardiac and vascular implants and grafts – reinforcing

existing evidence for genetic correlation between these traits in an unbiased manner. History

of taking prescriptions to treat vascular and abnormal cholesterol levels, such as rosuvastatin,

ezerimibe, and HMG-CoA reductase inhibitor statins, also clustered with PRS_{LOAD} and other

dementias at lower α-value thresholds. Interestingly, the PRS_{LOAD} did not consistently cluster

with the UKB-derived family history of AD dementia (ICD code: G30.9), family history of

AD (ICD code: G30.9) or dementia (ICD code: F03), PRSs – demonstrating the divergence

of PRS derived from different cohorts. However, this divergence was only present at higher

α-value thresholds ($\Pi_{ALL,1}$, $\Pi_{ALL,0.01}$, $\Pi_{ALL,0.05}$, $\Pi_{ALL,0.01}$) reinforcing the importance of SNP

inclusion when using clumping and thresholding [40].

The removal of MHC and/or *APOE* loci resulted in several changes in module number and

membership across α-value thresholds. Most notably, when the *APOE* region was excluded

($\Pi_{APOE-,\alpha}$), the PRS_{LOAD} clustered either in the grey module (this module contains a set of

PRSs which have not been clustered in any module), or the turquoise module (this module

contains the highest number of PRSs), along with other dementia-related phenotypes (and

many others). This was true across all α-value thresholds except for the blue module

($\Pi_{APOE-,0.01}$ represented by weight, $\Pi_{APOE-,0.005}$ represented by BMI, $\Pi_{APOE-,10^{-7}}$

represented by family history of diabetes, $\Pi_{APOE-,10^{-5}}$ represented by prednisolone,

$\Pi_{APOE-,5\times10^{-5}}$ represented by Rheumatoid Arthritis (RA)), yellow module ($\Pi_{APOE-,0.001}$

represented by RA), and the brown module ($\Pi_{APOE-,0.0005}$ represented by RA) where the

PRS_{LOAD} was clustered with other PRSs of which none were PRS of dementia or family

history of AD. When the MHC region was also excluded, there were some modules where

PRS_{LOAD} was clustered with PRS of dementia or family history of AD or delirium. However,

this membership was not consistent and varied for different α-value thresholds. Furthermore,

PRS_{LOAD} appeared in the blue module ($\Pi_{MHC/APOE-,5\times10^{-6}}$ represented by presence of cardiac

and vascular implants and grafts, $\Pi_{MHC/APOE-,0.01}$ represented by weight).

Associations of ePRS with clinical and neuropathological traits

Following the network-based identification of correlated PRSs across the phenome, we evaluated the associations of derived ePRSs with late-life cognitive and neuropathological phenotypes. **Figure 2** summarizes these results across all α-value thresholds (and all PRS matrices), highlighting several modules strongly associated with our LOAD-related phenotypes at a false discovery rate (FDR) adjusted q-value ≤ 0.05. Notably, all significant ePRS associations were observed for the $\Pi_{ALL,\alpha}$ and $\Pi_{MHC-,\alpha}$ matrices, whereas none of the ePRSs calculated from $\Pi_{APOE-,\alpha}$ and $\Pi_{MHC-APOE-,\alpha}$ passed FDR correction. Full sets of ePRS association results are summarized in **Figures S18-S20**.

Among the top results were the pink ePRS from $\Pi_{All,5e-08}$, the red module from $\Pi_{All,1e-07}$, and the red module from $\Pi_{All,5e-05}$, $\Pi_{All,1e-06}$, which were each largely composed of PRSs for cholesterol levels, and strongly associated with all five LOAD phenotypes. EPRSs with hub PRSs of maximum carotid intima-medial thickness (IMT; based on ultrasound) were strongly associated with β-amyloid levels. With the removal of the MHC region, the top associated ePRSs with all five phenotypes were consistent in terms of overall PRSs in these modules. Though, more ePRSs represented by high cholesterol indicated association with all five phenotypes. Overall, ePRSs represented by hubs of high cholesterol levels, LDL, and maximum carotid IMT were found to be highly associated with the LOAD-related traits and neuropathologies.

Assessment of improvements in predictive performance of ePRS over established single-trait LOAD PRS

Having identified several ePRSs associated with LOAD-related traits and neuropathologies, we sought to test whether these ePRSs could significantly augment single-PRS models of only PRS$_{LOAD}$ and basic demographics. In some cases, the ePRSs were able to improve the variance explained by the base models by up to 1.7%. Modelling results for selected top ePRSs are shown in **Table 2**. The brown hub of $\Pi_{MHC-,1e-05}$, represented by the hub PRS for cholesterol levels, was able to improve predictive models over the single-trait PRS$_{LOAD}$ for both β-amyloid counts ($P_{likelihood-ratio\ test} = 5.29 \times 10^{-5}$) and global random slope of cognitive decline ($P_{likelihood-ratio\ test} = 3.16 \times 10^{-3}$). Similarly, the red module, represented by high cholesterol levels, from $\Pi_{All,1e-07}$, was able to improve the predictability power of the PRS$_{LOAD}$ in association with β-amyloid count ($P_{likelihood-ratio\ test} = 9.25 \times 10^{-6}$). **Figure 3** (**expanded in S21**) summarizes the overall performance comparison

548   between base models and the ePRS-inclusive full models (defined as model 2 and model 3).

549   Only ePRSs with FDR-adjusted q-value ≤ 0.05 for the likelihood ratio test between base and

550   full model are shown.

Figure 2. Eigen-PRS associations with LOAD-related phenotypes

A) Selected dendrogram (based on the WGCNA topological overlap matrix of $\Pi_{All,5e-06}$) produced by the WGCNA pipeline. **B)** Dotplots summarizingthe associations of ePRSs derived from Π_{All} with each LOAD phenotype. Hub PRSs for significantly associated ePRSs (FDR q-value ≤ 0.05) are labelled. The red dotted line represents uncorrected *p*=0.05. The module represented by high cholesterol (pink) at α-value threshold of 5×10^{-7} had the highest association with all five phenotypes. The modules, represented by high cholesterol levels (pink, purple) showed the strongest associations with all five aging phenotypes at α-value threshold of 1×10^{-6}, 1×10^{-7}, and 5×10^{-8}. A triangular dot shape indicates that the LOAD PRS calculated from Bellenguez et al. (2022) was a member of the PRS cluster from which a given ePRS was calculated. Refer to **Figure S18, S19, and S20** for summary results from PRS matrices from sensitivity analyses (Π_{MHC-}, Π_{APOE-}, $\Pi_{MHC-APOE-}$).

Figure 3. Summary of model improvements for ePRS-inclusive models over single-trait LOAD PRS models of LOAD-related phenotypes.

Barplots showing the results of the comparison of full models including ePRSs and single-trait PRS models of PRS_{LOAD}, calculated from $\Pi_{ALL,\alpha}$. For continuous response variables (Aβ burden, PHF tau, cognitive global random slope, cognitive global random slope at last visit) we report the .632 bootstrapped variation explained (R^2) by the covariates for the base (model 2) and full model (model 3). Results for ePRS derived from matrices excluding the MHC region are shown in **Figure S21**. Error bards shown are estimates of 95% confidence intervals for variance explained with bootstrapping. Significance of differences in model performance were determined using likelihood ratio tests between the base and full models prior to bootstrapping, followed by FDR correction (all shown results had corrected q-value < 0.01).

Table 2. Evaluation of significant ePRSs models against standard single-trait PRS_{LOAD}

models of Alzheimer's neuropathology and cognitive decline

PRS Matrix	Outcome	ePRS (module)	β_{ePRS}	Model Performance (95% CI) [##]	ΔModel performance P_{LRT} [###]	n
$\Pi_{All,5\times10^{-5}}$	Total Aβ	Red (High cholesterol)	5.105	0.084 (0.063,0.123)	0.008 4.69×10^{-4}	1238
$\Pi_{All,1\times10^{-6}}$	Total Aβ	Salmon (Mean Carotid)	3.634	0.085 (0.059,0.114)	0.002 8.63×10^{-3}	1238
$\Pi_{All,1\times10^{-7}}$	Global slope	Red (High Cholesterol)	-0.334	0.030 (0.017,0.048)	0.003 1.11×10^{-3}	1915
	Total Aβ		6.770	0.093 (0.067,0.125)	0.011 9.25×10^{-6}	1238
$\Pi_{All,5\times10^{-8}}$	Global slope	Pink (High Cholesterol)	-0.325	0.029 (0.013,0.047)	0.002 1.46×10^{-3}	1915
	Total Aβ		6.545	0.094 (0.071,0.126)	0.013 1.57×10^{-5}	1238
$\Pi_{MHC-,0.0001}$	Total Aβ	Green (High Cholesterol)	0.179	0.091 (0.073,0.129)	0.017 9.02×10^{-7}	1238
$\Pi_{MHC-,5\times10^{-5}}$	Global slope	Yellow (High Cholesterol)	-0.017	0.093 (0.071,0.121)	0.013 1.10×10^{-5}	1915
	Total Aβ		0.181	0.091 (0.059,0.121)	0.010 2.22×10^{-5}	1238
$\Pi_{MHC-,1\times10^{-5}}$	Global slope	Brown (High Cholesterol)	-0.015	0.085 (0.061,0.119)	0.005 3.16×10^{-3}	1915
	Total Aβ		0.163	0.085 (0.064,0.122)	0.012 5.29×10^{-5}	1238
$\Pi_{MHC-,1\times10^{-6}}$	Global slope	Brown (High Cholesterol)	-0.013	0.030 (0.015,0.048)	0.004 9.71×10^{-4}	1915
	Total Aβ		0.180	0.079 (0.054,0.118)	0.008 3.39×10^{-4}	1238
$\Pi_{MHC-,5\times10^{-7}}$	Global slope	Red (High Cholesterol)	-0.013	0.036 (0.024,0.053)	0.004 2.02×10^{-3}	1915
	Total Aβ		0.179	0.028 (0.013,0.048)	0.002 1.88×10^{-3}	1238
$\Pi_{MHC-,1\times10^{-7}}$	Total Aβ	Black (LDL)	0.205	0.084 (0.058,0.131)	0.003 4.16×10^{-3}	1238
$\Pi_{MHC-,5\times10^{-8}}$	Total Aβ	Green (LDL)	0.209	0.037 (0.023,0.052)	0.001 1.37×10^{-2}	1238

Model performance is provided as variance explained (R^2) after .632 bootstrapping, including 25% confidence interval. For Δmodel performance, the ΔR^2 value is shown as well as the *p*-value of the likelihood ratio test between the full and base models. *p = <0.05, **p = <0.01, ***p = 0.001. All *p*-values are reported at an uncorrected level. Global slope = slope of change in longitudinal global cognitive performance; n = sample size.

Discussion

We generated over 120,000 PRSs from 2,218 phenotypes in a sample of over 2,000 elderly individuals harnessing the latest GWAS summary statistics from the Pan-UK Biobank Consortium. We used a network-based approach to derive clusters of PRSs and identified meaningful patterns of biologically and clinically related phenotypes with relevance to cognitive decline and core LOAD-related neuropathologies. By including summary statistics for thousands of diagnoses, traits, biomarkers, behaviours, and health-related measures, rather than selecting major risk factors *a priori*, we build a richer understating of the relationship between the genetic structure of our elderly population and risk for LOAD.

In PCA-based exploration of the full set of PRSs at each alpha threshold, loadings on the top PCs uncovered both previously observed genetic correlations of certain phenotypes, as well as some new ones; at lower α-value thresholds, corresponding to genome-wide and near genome-wide significant SNPs only, the top PCs were largely dominated by PRSs for cholesterol levels and cardiovascular related health conditions, treatments, and procedures. At higher α-value thresholds, we found PRSs for pain medications, including opioids and general over-the-counter painkillers, such as paracetamol, dominating the loadings of top PCs. We hypothesize that this is due to the wide use of these medications for broad spectra of clinical and subclinical afflictions, and therefore genetic associations with their use would tag variation common across hundreds or thousands of included phenotypes (from minor injury risk, to cancers, to stress-related events and conditions). Unsurprisingly, top PCs explained smaller proportions of variance across PRS matrices that included more PRSs (i.e. those calculated using less stringent, higher alpha thresholds). However, the proportion of PCs (out of number of included PRSs) required to account for 80% of variance in all PRSs was largely consistent across larger α-value thresholds (5×10^{-6} to 1). Compared to similar work [21], the LOAD PRS consistently clustered PRS for cholesterol levels and statin prescription, with both PRSs contributing highly to their corresponding PC. However, we also noticed the influence of apolipoprotein biomarker and LDL in these PCs as well. While PCA provided useful insight into the primary axes of variability in our PRS matrices, the utility of the resulting PCs for genetic prediction are not clear; for modelling there are no reliable guidelines for which PCs to include at each α-value threshold, and the variation explained by the top PCs was less than 1% for the majority of the PCs.

Thus, we applied a clustering method which is traditionally used on gene co-expression data,

WGCNA. In most cases, ePRSs derived from the WGCNA pipeline that were significantly

associated with LOAD phenotypes were composed of a combination of PRSs for dementia-

related phenotypes (e.g. family history of LOAD, LOAD diagnosis, and all-cause dementia),

cholesterol levels, prescriptions for statins and other cardiovascular health medications,

diseases, and physiological measurements of carotid arteries. This reinforces the known links

between cardiovascular health and risk for LOAD [56], but also points toward the specific

aspects of cardiovascular health that may be more genetically important for determining this

risk than others. In sensitivity analyses removing the *APOE* and/or MHC regions from our

scores, we noted major impacts on the associations of derived ePRS on LOAD phenotypes. In

fact, we found no associations (q-value < 0.05) between ePRSs produced from Π_{APOE-} and

$\Pi_{MHC/APOE-}$ and any LOAD phenotypes. Even ePRSs that included PRS$_{LOAD}$ were not

associated with LOAD-related phenotypes, though interestingly PRS$_{LOAD}$ did cluster with

PRSs for rheumatoid arthritis, which has been shown to be genetically related to microglial

activation that is implicated in LOAD pathogenesis [32].

This study has several limitations. First, our conclusions related to the relationship between

heritable risk for our phenotypes of interest are based on summary statistics derived from the

UK Biobank (indeed, those of any PRS-based study). The UK Biobank has known healthy

volunteer bias [57], and our target ROS/MAP sample also has a known late-life resilience

bias as a result of survival to study entry without dementia and lifestyle factors associated

with part of its underlying religious population [25]. Therefore, the transfer of genetic risk

profiles from one population to the other may be subject to challenges with cross-sample

portability – most apparent when considering ancestral differences but extend to age, sex, and

measurement differences between groups - that may hide true genetic effects [58]. This issue

of portability is also made clear by the observation that our PRS$_{LOAD}$ did not always cluster

with UKB summary stats-derived LOAD and dementia PRSs; in part likely due to differences

in phenotypic definition between studies. Second, by clustering matrices across traits and

within SNP inclusion thresholds, we have assumed that meaningful correlations between

PRSs may not occur between different traits at different thresholds. Others have attempted to

resolve this, for example by calculating the first principal component of a set of PRSs derived

from the same set of summary statistics across multiple thresholds [42]. However, we sought

specifically to find relationships between the PRSs rather than finding the most optimized α-

value threshold of each PRS, since it would require availability of the same phenotypes in the
target population. Further, while we used a 5% threshold on estimated heritability in the UK
Biobank to ensure inclusion of only traits with a nonzero genetic basis, there is no guarantee
that PRS models based on these GWAS will have a significant predictive power. Therefore,
we acknowledge that our observed clusters of PRSs may have been biased by patterns of
unquantified within-trait predictive performance, which could only be estimated at scale in
fully-phenotyped (i.e. all 2000+ traits) out-of-sample data. Methods have been proposed for
addressing this limitation in PRS tuning [59] and they will be explored in future work.
Finally, we selected only sex non-specific phenotypes to generate their corresponding PRSs
using commonly used methods. Thus, our analysis omits information available for sex-
specific traits, such as those related to menopause or testicular cancers, which are known to
impact aging in important ways. Additionally, evidence suggests sex-specific effects of
LOAD genetic risk factors in the clinical progression of LOAD and the accumulation of
LOAD-related neuropathologies [60]; these effects were not explored here and are the focus
of ongoing work.
651
In sum, our study provides a whole-person view of the genetic risk landscape for LOAD in an
elderly population. We highlight the impacts of 1) varying SNP inclusion α-value thresholds
on this landscape using the most popular PRS calculation method, 2) the links between
several blood-based and ultrasound imaging-derived cardiovascular traits on LOAD risk, as
well as 3) the important contribution of the *APOE* region. Our results suggest that combining
genetic risk profiles for vascular and dementia-related phenotypes using latent methods may
offer improvements over existing single-trait PRS models of LOAD, a step toward achieving
the unrealized potential of PRS for risk stratification in populations and clinics.
660
661

References

1. 2021 Alzheimer's disease facts and figures. Alzheimers Dement. 2021;17: 327–406. doi:10.1002/alz.12328

2. Sims R, Hill M, Williams J. The multiplex model of the genetics of Alzheimer's disease. Nat Neurosci. 2020;23: 311–322. doi:10.1038/s41593-020-0599-5

3. Gatz M, Reynolds CA, Fratiglioni L, Johansson B, Mortimer JA, Berg S, et al. Role of genes and environments for explaining Alzheimer disease. Arch Gen Psychiatry. 2006;63: 168–174.

4. Andrews SJ, Fulton-Howard B, Goate A. Interpretation of risk loci from genome-wide association studies of Alzheimer's disease. Lancet Neurol. 2020;19: 326–335. doi:10.1016/S1474-4422(19)30435-1

5. Saez-Atienzar S, Masliah E. Cellular senescence and Alzheimer disease: the egg and the chicken scenario. Nat Rev Neurosci. 2020;21: 433–444. doi:10.1038/s41583-020-0325-z

6. Jones L, Harold D, Williams J. Genetic evidence for the involvement of lipid metabolism in Alzheimer's disease. Biochim Biophys Acta BBA - Mol Cell Biol Lipids. 2010;1801: 754–761. doi:10.1016/j.bbalip.2010.04.005

7. Jones L, Holmans PA, Hamshere ML, Harold D, Moskvina V, Ivanov D, et al. Genetic Evidence Implicates the Immune System and Cholesterol Metabolism in the Aetiology of Alzheimer's Disease. PLOS ONE. 2010;5: e13950. doi:10.1371/journal.pone.0013950

8. Voineskos AN, Lerch JP, Felsky D, Shaikh S, Rajji TK, Miranda D, et al. The Brain-Derived Neurotrophic Factor Val66Met Polymorphism and Prediction of Neural Risk for Alzheimer Disease. Arch Gen Psychiatry. 2011;68: 198–206. doi:10.1001/archgenpsychiatry.2010.194

9. Yokoyama JS, Wang Y, Schork AJ, Thompson WK, Karch CM, Cruchaga C, et al. Association Between Genetic Traits for Immune-Mediated Diseases and Alzheimer Disease. JAMA Neurol. 2016;73: 691–697. doi:10.1001/jamaneurol.2016.0150

10. Ferrucci L, Fabbri E. Inflammageing: chronic inflammation in ageing, cardiovascular disease, and frailty. Nat Rev Cardiol. 2018;15: 505–522. doi:10.1038/s41569-018-0064-2

11. Hersi M, Irvine B, Gupta P, Gomes J, Birkett N, Krewski D. Risk factors associated with the onset and progression of Alzheimer's disease: A systematic review of the evidence. Neurotoxicology. 2017;61: 143–187. doi:10.1016/j.neuro.2017.03.006

12. Loeffler DA. Modifiable, Non-Modifiable, and Clinical Factors Associated with Progression of Alzheimer's Disease. J Alzheimers Dis JAD. 2021;80: 1–27. doi:10.3233/JAD-201182

700   13.   Santiago JA, Potashkin JA. The Impact of Disease Comorbidities in Alzheimer's
701   Disease. Front Aging Neurosci. 2021;13. Available:
702   https://www.frontiersin.org/articles/10.3389/fnagi.2021.631770

703   14.   Lucey BP. It's complicated: The relationship between sleep and Alzheimer's disease in
704   humans. Neurobiol Dis. 2020;144: 105031. doi:10.1016/j.nbd.2020.105031

705   15.   Kivimäki M, Luukkonen R, Batty GD, Ferrie JE, Pentti J, Nyberg ST, et al. Body mass
706   index and risk of dementia: Analysis of individual-level data from 1.3 million
707   individuals. Alzheimers Dement J Alzheimers Assoc. 2018;14: 601–609.
708   doi:10.1016/j.jalz.2017.09.016

709   16.   Lennon MJ, Koncz R, Sachdev PS. Hypertension and Alzheimer's disease: is the picture
710   any clearer? Curr Opin Psychiatry. 2021;34: 142–148.
711   doi:10.1097/YCO.0000000000000684

712   17.   Feringa FM, van der Kant R. Cholesterol and Alzheimer's Disease; From Risk Genes to
713   Pathological Effects. Front Aging Neurosci. 2021;13. Available:
714   https://www.frontiersin.org/article/10.3389/fnagi.2021.690372

715   18.   Olmastroni E, Molari G, De Beni N, Colpani O, Galimberti F, Gazzotti M, et al. Statin
716   use and risk of dementia or Alzheimer's disease: a systematic review and meta-analysis
717   of observational studies. Eur J Prev Cardiol. 2022;29: 804–814.
718   doi:10.1093/eurjpc/zwab208

719   19.   Gong C-X, Liu F, Iqbal K. Multifactorial Hypothesis and Multi-Targets for Alzheimer's
720   Disease. J Alzheimers Dis JAD. 2018;64: S107–S117. doi:10.3233/JAD-179921

721   20.   UK Biobank. In: Neale lab [Internet]. [cited 13 Jan 2022]. Available:
722   http://www.nealelab.is/uk-biobank

723   21.   Tanigawa Y, Li J, Justesen JM, Horn H, Aguirre M, DeBoever C, et al. Components of
724   genetic associations across 2,138 phenotypes in the UK Biobank highlight adipocyte
725   biology. Nat Commun. 2019;10: 4064. doi:10.1038/s41467-019-11953-9

726   22.   Yan D, Hu B, Darst BF, Mukherjee S, Kunkle BW, Deming Y, et al. Biobank-wide
727   association scan identifies risk factors for late-onset Alzheimer's disease and
728   endophenotypes. 2018 Nov p. 468306. doi:10.1101/468306

729   23.   Andrews SJ, Fulton-Howard B, O'Reilly P, Marcora E, Goate AM, Consortium
730   collaborators of the ADG. Causal Associations Between Modifiable Risk Factors and
731   the Alzheimer's Phenome. Ann Neurol. 2021;89: 54–65. doi:10.1002/ana.25918

732   24.   Towards clinical utility of polygenic risk scores | Human Molecular Genetics | Oxford
733   Academic. [cited 13 Jan 2022]. Available:
734   https://academic.oup.com/hmg/article/28/R2/R133/5540980?login=true

735   25.   Bennett DA, Buchman AS, Boyle PA, Barnes LL, Wilson RS, Schneider JA. Religious
736   Orders Study and Rush Memory and Aging Project. J Alzheimers Dis. 2018;64: S161–
737   S189. doi:10.3233/JAD-179939

26. McKhann GM, Knopman DS, Chertkow H, Hyman BT, Jack Jr. CR, Kawas CH, et al.
The diagnosis of dementia due to Alzheimer's disease: Recommendations from the
National Institute on Aging-Alzheimer's Association workgroups on diagnostic
guidelines for Alzheimer's disease. Alzheimers Dement. 2011;7: 263–269.
doi:10.1016/j.jalz.2011.03.005

27. Schneider JA, Arvanitakis Z, Bang W, Bennett DA. Mixed brain pathologies account
for most dementia cases in community-dwelling older persons. Neurology. 2007;69:
2197–2204. doi:10.1212/01.wnl.0000271090.28148.24

28. Wilson RS, Boyle PA, Yu L, Barnes LL, Sytsma J, Buchman AS, et al. Temporal course
and pathologic basis of unawareness of memory loss in dementia. Neurology. 2015;85:
984–991. doi:10.1212/WNL.0000000000001935

29. Wilson RS, Yu L, Trojanowski JQ, Chen E-Y, Boyle PA, Bennett DA, et al. TDP-43
pathology, cognitive decline, and dementia in old age. JAMA Neurol. 2013;70: 1418–
1424. doi:10.1001/jamaneurol.2013.3961

30. Bennett: Decision rules guiding the clinical diagnosis... - Google Scholar. [cited 11 May
2022]. Available:
https://scholar.google.com/scholar_lookup?journal=Neuroepidemiology&title=Decision
+rules+guiding+the+clinical+diagnosis+of+Alzheimer%E2%80%99s+disease+in+two+
community-based+cohort+studies+compared+to+standard+practice+in+a+clinic-
based+cohort+study&author=DA+Bennett&author=JA+Schneider&author=NT+Aggar
wal&author=Z+Arvanitakis&author=RC+Shah&volume=27&issue=3&publication_yea
r=2006b&pages=169-76&pmid=17035694&

31. Wilson RS, Arnold SE, Schneider JA, Tang Y, Bennett DA. The relationship between
cerebral Alzheimer's disease pathology and odour identification in old age. J Neurol
Neurosurg Psychiatry. 2007;78: 30–35. doi:10.1136/jnnp.2006.099721

32. Felsky D, Roostaei T, Nho K, Risacher SL, Bradshaw EM, Petyuk V, et al.
Neuropathological correlates and genetic architecture of microglial activation in elderly
human brain. Nat Commun. 2019;10: 409. doi:10.1038/s41467-018-08279-3

33. De Jager PL, Ma Y, McCabe C, Xu J, Vardarajan BN, Felsky D, et al. A multi-omic
atlas of the human frontal cortex for aging and Alzheimer's disease research. Sci Data.
2018;5: 180142. doi:10.1038/sdata.2018.142

34. Taliun D, Harris DN, Kessler MD, Carlson J, Szpiech ZA, Torres R, et al. Sequencing
of 53,831 diverse genomes from the NHLBI TOPMed Program. Nature. 2021;590: 290–
299. doi:10.1038/s41586-021-03205-y

35. Marees AT, de Kluiver H, Stringer S, Vorspan F, Curis E, Marie-Claire C, et al. A
tutorial on conducting genome-wide association studies: Quality control and statistical
analysis. Int J Methods Psychiatr Res. 2018;27: e1608. doi:10.1002/mpr.1608

36. Allen NE, Sudlow C, Peakman T, Collins R, Biobank on behalf of U. UK Biobank
Data: Come and Get It. Sci Transl Med. 2014 [cited 14 Jan 2022].
doi:10.1126/scitranslmed.3008601

37. Zhou W, Nielsen JB, Fritsche LG, Dey R, Gabrielsen ME, Wolford BN, et al.
Efficiently controlling for case-control imbalance and sample relatedness in large-scale
genetic association studies. Nat Genet. 2018;50: 1335–1341. doi:10.1038/s41588-018-
0184-y

38. Bellenguez C, Küçükali F, Jansen IE, Kleineidam L, Moreno-Grau S, Amin N, et al.
New insights into the genetic etiology of Alzheimer's disease and related dementias. Nat
Genet. 2022;54: 412–436. doi:10.1038/s41588-022-01024-z

39. Choi SW, O'Reilly PF. PRSice-2: Polygenic Risk Score software for biobank-scale
data. GigaScience. 2019;8: giz082. doi:10.1093/gigascience/giz082

40. Privé F, Vilhjálmsson BJ, Aschard H, Blum MGB. Making the Most of Clumping and
Thresholding for Polygenic Scores. Am J Hum Genet. 2019;105: 1213–1221.
doi:10.1016/j.ajhg.2019.11.001

41. Association of accelerometer-derived sleep measures with lifetime psychiatric
diagnoses: A cross-sectional study of 89,205 participants from the UK Biobank | PLOS
Medicine. [cited 22 Jun 2022]. Available:
https://journals.plos.org/plosmedicine/article?id=10.1371/journal.pmed.1003782

42. Zhao H, Mitra N, Kanetsky PA, Nathanson KL, Rebbeck TR. A Practical Approach to
Adjusting for Population Stratification in Genome-wide Association Studies: Principal
Components And Propensity Scores (PCAPS). Stat Appl Genet Mol Biol. 2018;17:
/j/sagmb.2018.17.issue-6/sagmb-2017-0054/sagmb-2017-0054.xml. doi:10.1515/sagmb-
2017-0054

43. Belloy ME, Napolioni V, Greicius MD. A Quarter Century of APOE and Alzheimer's
Disease: Progress to Date and the Path Forward. Neuron. 2019;101: 820–838.
doi:10.1016/j.neuron.2019.01.056

44. Matzaraki V, Kumar V, Wijmenga C, Zhernakova A. The MHC locus and genetic
susceptibility to autoimmune and infectious diseases. Genome Biol. 2017;18: 76.
doi:10.1186/s13059-017-1207-1

45. Candore G, Balistreri CR, Colonna-Romano G, Lio D, Caruso C. Major
histocompatibility complex and sporadic Alzheimer's disease: a critical reappraisal. Exp
Gerontol. 2004;39: 645–652. doi:10.1016/j.exger.2003.10.027

46. Zhou X, Chen Y, Mok KY, Kwok TCY, Mok VCT, Guo Q, et al. Non-coding
variability at the APOE locus contributes to the Alzheimer's risk. Nat Commun.
2019;10: 3310. doi:10.1038/s41467-019-10945-z

47. Abdi H, Williams LJ. Principal component analysis. WIREs Comput Stat. 2010;2: 433–
459. doi:10.1002/wics.101

48. Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network
analysis. BMC Bioinformatics. 2008;9: 559. doi:10.1186/1471-2105-9-559

49. Bustamante AC, Armstrong DL, Uddin M. Epigenetic profiles associated with major
depression in the human brain. Psychiatry Res. 2018;260: 439–442.
doi:10.1016/j.psychres.2017.12.010

50. Jia R, Zhao H, Jia M. Identification of co-expression modules and potential biomarkers
of breast cancer by WGCNA. Gene. 2020;750: 144757.
doi:10.1016/j.gene.2020.144757

51. Levine ME, Langfelder P, Horvath S. A Weighted SNP Correlation Network Method
for Estimating Polygenic Risk Scores. Methods Mol Biol Clifton NJ. 2017;1613: 277–
290. doi:10.1007/978-1-4939-7027-8_10

52. Langfelder P, Zhang B, Horvath S. Defining clusters from a hierarchical cluster tree: the
Dynamic Tree Cut package for R. Bioinforma Oxf Engl. 2008;24: 719–720.
doi:10.1093/bioinformatics/btm563

53. Storey JD, Tibshirani R. Statistical significance for genomewide studies. Proc Natl Acad
Sci. 2003;100: 9440–9445. doi:10.1073/pnas.1530509100

54. Sun L, Craiu RV, Paterson AD, Bull SB. Stratified false discovery control for large-
scale hypothesis testing with application to genome-wide association studies. Genet
Epidemiol. 2006;30: 519–530. doi:10.1002/gepi.20164

55. Efron B, Tibshirani R. Improvements on Cross-Validation: The .632+ Bootstrap
Method. J Am Stat Assoc. 1997;92: 548–560. doi:10.2307/2965703

56. Samieri C, Perier M-C, Gaye B, Proust-Lima C, Helmer C, Dartigues J-F, et al.
Association of Cardiovascular Health Level in Older Age With Cognitive Decline and
Incident Dementia. JAMA. 2018;320: 657–664. doi:10.1001/jama.2018.11499

57. Fry A, Littlejohns TJ, Sudlow C, Doherty N, Adamska L, Sprosen T, et al. Comparison
of Sociodemographic and Health-Related Characteristics of UK Biobank Participants
With Those of the General Population. Am J Epidemiol. 2017;186: 1026–1034.
doi:10.1093/aje/kwx246

58. Privé F, Aschard H, Carmi S, Folkersen L, Hoggart C, O'Reilly PF, et al. Portability of
245 polygenic scores when derived from the UK Biobank and applied to 9 ancestry
groups from the same cohort. Am J Hum Genet. 2022;109: 12–23.
doi:10.1016/j.ajhg.2021.11.008

59. Zhao Z, Yi Y, Song J, Wu Y, Zhong X, Lin Y, et al. PUMAS: fine-tuning polygenic risk
scores with GWAS summary statistics. Genome Biol. 2021;22: 257.
doi:10.1186/s13059-021-02479-9

60. Fan CC, Banks SJ, Thompson WK, Chen C-H, McEvoy LK, Tan CH, et al. Sex-
dependent autosomal effects on clinical progression of Alzheimer's disease. Brain.
2020;143: 2272–2280. doi:10.1093/brain/awaa164

851
852

Supporting Information

The Supporting Document contains **Supplementary Methods**, **Tables S1-S3** (**Table S1** is stored as an external CSV file due to its size), and **Figures S1-S20**.

Declaration of Interests

The authors declare no competing interests.

Acknowledgements

DF is supported by the Michael and Sonja Koerner Family New Scientist Award, the Krembil Family Foundation, the Canadian Institutes of Health Research (CIHR), and the Centre for Addiction and Mental Health (CAMH) Discovery Fund. LS is supported by The Natural Sciences and Engineering Research Council of Canada (NSERC), RGPIN- 04934 and RGPAS-522594. ROS/MAP is supported by P30AG10161, P30AG72975, R01AG15819, R01AG17917. U01AG46152, U01AG61356.

Data and Code Availability

All phenotypes and omics data of the Religious Orders Study (ROS) and Memory Aging Project (MAP) Cohorts are available through the RADC Resource Sharing Hub at www.radc.rush.edu. ROS/MAP resources can be requested at https://www.radc.rush.edu. The UK Biobank GWAS summary statistics are available from Neale Lab, http://www.nealelab.is/ukbiobank. All scripts used for this manuscript are open-source and available at https://github.com/aminekhasteh/ePRS.