Ants dynamically adjust liquid foraging strategies in response to

biophysical constraints

Haruna FUJIOKA[1,2]*, Manon Marchand[3], Adria C. LeBoeuf[2]*

1. Faculty of Environmental and Life Science, Okayama University, Okayama, Japan.

2. Department of Biology, University of Fribourg, Fribourg, Switzerland.

3. Department of Physics, University of Fribourg, Fribourg, Switzerland.

Author correspondence* HF: fujioka.ha@gmail.com, AL: adria.leboeuf@gmail.com

Abstract

Ant foragers need to provide food to the rest of the colony, which often requires food transport over

long distances. Foraging for liquid is especially challenging because it is difficult to transport and

share. Many social insects store liquid food inside the crop to transport it to the nest, and then

regurgitate this fluid to distribute it to nestmates through a behaviour called trophallaxis. Some ants

instead transport fluids with a riskier behaviour –holding a drop of liquid between the mandibles

through surface tension– after which the ant shares this droplet with nestmates without ingestion or

regurgitation in a behaviour called pseudotrophallaxis. Here, we hypothesized that ants optimise their

liquid-collection approach depending on food quality and biophysical properties. Working with a

ponerine ant that uses both trophallaxis and pseudotrophallaxis, we investigated why each liquid-

collection behaviour might be favored under different conditions by measuring handling time and

liquid viscosity and reaction to food quality (i.e., sugar concentration and viscosity) using a viscosity

additive. We found that ants could collect more liquid food per unit time by mandibular grabbing than

by drinking. At high viscosities, which in nature correspond to high sugar concentrations, ants

switched their liquid collection method to mandibular grabbing in response to viscosity, and not to

sweetness. In addition, mandibular grabbing of liquid food allowed ants to carry more sugar per unit

time than drinking. Our results demonstrate that ants change not only their feeding preference but also

their transport and sharing methods according to viscosity–a proxy for sugar concentration in nature–

optimising the mass of sugar returned to the nest over time.

keywords: mandibular pseudotrophallaxis, social bucket, *Diacamma* cf. *indicum*, liquid

transportation, optimal foraging theory, biomechanics

Introduction

Efficient foraging is crucial for animals to survive, grow and reproduce. Organisms need to balance energy spent with energy gained [1,2]. Optimal foraging theory assumes that animals' foraging decision making has evolved to the point that the fitness of individuals has been maximised. Foraging provides energy to survive and reproduce; however, it has the cost of exposing the individual to being preyed upon by other animals and it costs energy, for example, due to the time spent exploiting, processing and transporting food.

Animals have a wide variety of foraging strategies. Ants are central-place foragers with diverse diets ranging from complete herbivory to complete predation [3–6]. Morphological and phylogenetic evidence suggests that the ancestral ant was a predator, and transitions to herbivory occurred several times in predatory lineages [6–8]. Plant-based food sources, such as plant nectar [9] or honeydew excreted by sapsucking Hemiptera and scale insects [10–12], are rich in carbohydrates. In many species and especially in ecologically dominant ant lineages, these sugary liquids are ants' main source of energy [13]. Additionally, liquid resources, such as honeydew or nectar, are less ephemeral relative to insect prey, and incur fewer risks for ant foragers relative to hunting. Thus, the use of plant-based food sources may lead to lower foraging time and less risk for foragers per calorie returned to the nest.

Transportation of liquid food is a foraging challenge. Many ants and bees transport liquid stored inside their crop, where it cannot easily be lost or stolen[14] during transport. Foragers regurgitate this fluid to distribute it to nestmates through a behaviour called trophallaxis. Many liquid-feeding ants have acquired morphological adaptations for trophallaxis. The ant crop is separated from their midgut by a variably developed proventriculus, which allows the crop to store a large amount of liquid in some species [15]. The structure of proventriculus varies considerably across taxa [16–18], and liquid-feeding ants often have a more elaborate proventriculus. The gaster and crop also need to be expandable to best store liquid food, either temporarily for transport or over the long term in the case of repletes. The extreme example of morphological specialisation are honeypot ants *Myrmecocystus* (Formicinae), where replete workers have a massive ball-like distended gaster full of food to the point where they can barely move [19]. Such species rely on trophallaxis to redistribute food from the repletes to the rest of the colony. Although trophallaxis is considered a safe and reliable liquid transportation method for ants, the crop load (i.e., liquid food intake) strongly depends on these morphological constraints.

Some ants do not have these morphological specialisations but nonetheless consume liquid food: Ectatomminae (*Ectatomma*), Ponerinae (*Diacamma, Neoponera Odontomachus, Paraponera, Pachycondyla, Rhytidoponera*). These ants typically use mandibular pseudotrophallaxis (hereafter called pseudotrophallaxis) as their method of liquid transport [15,20]. Instead of storing liquid inside the crop, foragers hold liquid food between heir mandibles where it forms a droplet because of surface tension. After foragers return to the nest, they pass the liquid food to nestmates without regurgitation.

Previous studies in ponerine ants have reported how this behaviour allows liquids to be distributed in
the nest [21,22]. This liquid transport method is sometimes referred to as the 'social bucket' method
and has been suggested to be an evolutionary precursor to 'true' trophallaxis [7].

Handling time is crucial for efficient foraging. For liquid food, the handling time includes
both the speed of food collection (i.e., drinking time or grabbing time) and the transport time to the
nest. Drinking time in ants has been shown to depend on food quality such as sugar concentration and
viscosity itself [23]. Previous studies found that drinking time increased linearly with increasing
sucrose concentration [23–27]. Individuals need to decide when to stop drinking, considering the
balance between energy gain and predation risk. The handling time of pseudotrophallaxis have not
been investigated. In the case of the transport time, once foragers store food in the crop, they can
transport the liquid food safely. When using pseudotrophallaxis, there is the possibility to lose the
liquid food along the return path. Also, keeping mandibles open adequately may reduce walking
speed and increase likelihood of predation.

In summary, to provide liquid resources for nestmates, ants must collect, transport, and share
liquid food through a series of behaviours. Generally, ants either drink, and internally store liquid and
share it through regurgitation and trophallaxis, or they grab liquid into a mandibular droplet and share
through pseudotrophallaxis (Figure 1). Some ants use both behaviours. The ponerine ant *Diacamma*
cf. *indicum* from Japan performs both trophallaxis and pseudotrophallaxis[28], and has a simple
proventricular morphology and has a rigid, non-extensible gaster. Thus, *Diacamma* cf. *indicum* is an
ideal model species to investigate efficient foraging strategies regarding liquid food because it allows
us to investigate foraging strategies without morphological specialisation.

The aim of the present study is to reveal what leads ants use the collection mode of
mandibular grabbing instead of drinking and whether ants' liquid collection modes mechanisms
maximise calorie intake rates for their colony. We hypothesize that viscosity triggers a switch in
collection behaviour between drinking and mandibular grabbing, where mandibular grabbing being
more efficient to collect high viscosity solutions. To test this hypothesis, we conducted behavioural
experiments to investigate 1) the volume and speed of liquid food collection depending on sugar
concentration, 2) whether ants change their transportation method depending on sugar concentration
or viscosity, and 3) the foraging efficiency for each approach by estimating the total amount of sugar
carried per trip.

Method

Colony collection and colony keeping

12 Colonies of *Diacamma* cf. *indicum* from Japan were collected from Kenmin-no-mori (Onna) and Sueyoshi park (Naha), Okinawa, Japan. The colonies were kept in plastic artificial nests filled with moistened plaster (9 cm diameter × 1.5 cm height). Each colony contained a mated gemma-possessing female (i.e., functional queen or gamergate), 50–150 workers, and brood. The artificial nests (90 mm in diameter) were placed in a plastic arena (diameter: cm, height: cm). Nests were maintained at 25 °C under a 12h/12h light-dark regime (light phase: 0800–2000 hours). Reared colonies were fed with chopped frozen crickets three times per week. Water and 10% sugar water were provided ad libitum.

Behaviour Definitions

The ethogram of the social bucket method (encompassing mandibular grabbing, transport and pseudotrophallaxis) of *Diacamma* ant is shown in Figure 1. Based on a previous study [29] three behaviours for liquid feeding and collection were defined 1) tasting: placing mandibles on a solution without drinking or antennating, 2) drinking: individuals drink (i.e., mouthpart, labrum, attached to a liquid solution), 3) grabbing: individuals use (open) mandibles to grab and pull at a liquid solution, occasionally succeeding in collecting a droplet. We confirmed that the focal ant species showed the same actions. However, tasting was difficult to see by video observation, and thus we only considered drinking and grabbing.

All behavioural experiments were conducted between 12:00-19:00 under the light condition at 25℃. Colonies were starved for 3-4 hours before experiments (starvation time based on preliminary behavioural observations). We placed an artificial nest on one side and a plate-shaped feeder (40 × 40 mm) on the other side of the foraging arena (460 × 260 × 100 mm) (Supp. figure 1). In the preliminary observation, all foraging trips and pseudotrophallaxis bouts were counted by watching the recorded video for 30 minutes (Supp. Figure 2). Ants clearly preferred sugar water over water.

Measurement of volume of water internally and externally carried

The volume of sugar water drunk was estimated as the difference in body mass before and after drinking. We chose ants outside of the nest and measured their body mass (before drinking). Then, each individual was separately placed into a plastic box (4.5 × 4.5 × 2 cm), containing a drop (approximately 100 μl) of sugar water. Ants were offered one of six different sucrose concentrations: 10, 20, 30, 40, 50, 60 % w/w. The concentrations we used were within the range reported for natural nectar sources for ants (extrafloral nectars: 4.7–76% w/w [30]). After 10 minutes, or when ants stopped drinking, we dried the ant mouths and measured the ant's body mass to determine the volume

of liquid food inside its crop. We estimated the volume (μL) of sugar water using the average weight

of sugar water per 1 μ l (Supp. table 1). Individuals were never tested more than once a day.

The volume of sugar water carried by mandibular grabbing was measured with a microcapillary tube.

We used the same setup of the experimental arena above (Supp. figure 1). Ants were offered one of

six different sucrose concentrations: 10, 20, 30, 40, 50, 60 % w/w. The ants freely accessed the

offered food. After an ant succeeded in grabbing a droplet of the solution, we collected the droplet

with a microcapillary tube during a return trip to the nest. The weight of the dispensed volume was

measured to calculate the volume of sugar water the ant carried. We estimated the volume (μL) of

sugar water using the average weight of sugar water per 1 μl (Supp. table 1).

Measurement of grabbing and drinking time

We placed an artificial nest on one side and a plate-shaped feeder (40 × 40 mm) on the other side of

the foraging arena (460 × 260 × 100 mm) (Supp. figure 1). Ants were offered one of the six sucrose

concentrations: 10, 20, 30, 40, 50, 60 % w/w. We video-recorded the area around the sucrose droplet

for 1 hour. We manually recorded one type of the interaction which foragers had with the droplet only

drinking, grabbing after drinking (both), or only grabbing and accumulated grabbing/drinking time for

each foraging trip by an observer analysing the videos.

Reaction to viscosity

To test the effects of sweetness and viscosity on foraging methods used, we modified the viscosity of

the solution using carboxymethylcellulose sodium salt (Medium viscosity) (Sigma-Aldrich). The

carboxymethylcellulose sodium salt (CMC) is a non-toxic inert viscosity modifier [31]. By adding

this product, we increase the viscosity of the solution without changing its sugar concentration. We

used: 10% w/w sugar solution with CMC 0.25% w/w (10CMC) as a viscosity-altered solution. We

confirmed that the sugar concentration of CMC additive solution was not changed using a

saccharimeter (Refractometer RBR32-ATC).

Dynamic viscosity measurement

We prepared six sugar solutions, 10, 20, 30, 40, 50, 60 % w/w, and in addition, two viscosity-altered

solutions, 10% sugar solution with CMC 0.25% (10CMC) and 30% sugar solution with CMC 0.25%

w/w (30T). We measured their dynamic viscosity at 25℃. The dynamic viscosity is determined in a

commercial stress-controlled rheometer (ANTON PAAR MCR 300) with a plate-plate geometry of 5

cm diameter and 0.5 mm gap. The bottom plate was roughened by sandblasting to prevent slip

artefacts and the temperature was fixed at 25°C by a Peltier hood. We applied constant shear rates,

and stress was computed when the steady-state regime had been reached for each shear rate. The

resulting stress versus shear rate experiments exhibited linear behaviour, as expected for Newtonian

liquids, and the dynamic viscosity was directly read from the slope evaluated by least-square

minimisation for each sample, for more detail see Rhee and Lee (1994).

Intake rate and estimation of sugar intake

Following a previous study [33], intake rate was calculated as the slope (μL/sec) of the linear

regression of crop load and corresponding feeding time. Thus, we measured these two for each sugar

concentration. For crop loads (μL), we used the same procedure as 'measurement of the volume of

liquid drunk'. For the corresponding feeding time, we filmed the behaviour of ants when drinking and

measured the time spent in contact with the droplet.

We calculated the total liquid load per trip based on the foraging action used. The total crop

load per trip was estimated by multiplying the intake rate by the accumulated time drinking. The

mandible load (collected by grabbing) was defined as the average volume of liquid carried for each

sugar concentration in the experiment 'Measurement of volume of water drunk and water carried'.

The average volume can change depending on sugar concentration. When ants performed both

drinking and grabbing, we summed the average volume of water carried and the estimated total crop

load. Using the average weight of sugar water per 1 μl (Supp. table 1), we converted the total liquid

load per trip (μL) to weight of the liquid load (mg). From the liquid load weight (mg), we calculated

the total sugar intake per trip (mg) for each sugar concentration.

Statistical analysis

Generalized linear regression models were used to investigate the relationship between the liquid

volume or loading speed with the food quality variables and foraging actions. Pairwise chi-square

tests with a Bonferroni correction were used for comparing foraging action on different sugar

concentrations. Generalized linear regression models were used to investigate the relationship of the

load with the food quality variables and foraging actions. A significance general level of 5% was used

in all comparisons. All analyses were run in R studio 2022.02.3 (package: ggplot2).

Results

To understand whether ants altered their liquid collection and transport behaviour according to the concentration of sugar they encounter, we measured multiple variables to find what the ants are optimising: volume curried, time spent per trip, frequencies of different foraging actions, and sugar load per trip.

First, we analysed the liquid food collection method and volume of liquid food collected by 637 ants from eight colonies feeding on six different sugar concentrations (Figure 2). We observed an interaction between sugar concentration and foraging action (Figure 2a, Table 1a, GLM, sugar $\times$ foraging action: $p < 0.001$), and therefore analysed the foraging methods separately. When ants drank liquid, the amount of sugar water imbibed decreased as sugar concentration increased (Figure 2a, Table 1b, LM: $p < 0.001$), suggesting that with increasing sugar concentration, drinking becomes more difficult. For the amount of liquid grabbed within the mandibles, there was no significant trend in the across the different sugar concentrations (Figure 2a. Table 1b, LM: $p = 0.41$). Thus, the most effective method to bring liquid food home to the colony, in terms of volume, depended on sugar concentration.

Regarding the collection time of drinking or mandible grabbing, we again found an interaction between sugar concentration and foraging action (Figure 2b, Table 1a, GLM, sugar $\times$ foraging action: $p < 0.001$). When ants drank, the drinking time also decreased with increasing sugar concentration (Figure 2b, Table 1b, LM: $p < 0.001$). Grabbing time slightly increased with sugar concentration (Figure 2b. Table 1b, LM: $p < 0.01$), though mandible grabbing generally took less time when compared to drinking.

Ants often collected sugar water in their mandibles after drinking sugar water and rarely performed only mandible grabbing without drinking (Figure 3). The proportion of these foraging actions was significantly different across different sugar concentrations (Figure 3a, chi-sq test with Bonferroni correction). The proportion of the mandible grabbing after drinking (both) and mandible grabbing alone (both of which results in pseudotrophallaxis) increased with increasing sugar concentration (Figure 3a). This indicates that ants switch to grabbing and pseudotrophallaxis when they feed on liquid food with higher concentrations of sugar. This could come about because this high-sugar food is more valuable or because high viscosity liquids are difficult for them to drink.

To test whether ants react to changes in sugar concentration or viscosity, we altered the viscosity of a low-sugar solution using a viscosity-modifying additive CMC (carboxymethylcellulose sodium salt). The viscosity level of 10% sugar water with CMC (CMC10) was comparable to the one between 40% and 50% sugar water (Supp. Table 2, Figure 4). When we offered ants CMC10, the proportion of drinking was significantly decreased and equivalent to the high viscosity 50% sugar solution (Figure 3b, chi-sq test, $p < 0.001$). This result suggests that ants switch collection method in response to viscosity, and not to sweetness.

To investigate whether this transition toward grabbing over drinking with increasing viscosity
was optimal for the colony, we estimated the total sugar load per trip by combining intake rate and
load per trip across the different sugar concentrations. The intake rate was the highest with 20% sugar
water (Supp. Table 3, Figure S3, 20%: 0.21 μL/min), and decreased dramatically with increasing
sugar concentration to just 8% of maximum at the highest tested viscosity (Supp. Table 3, Figure S3).
The total crop load per trip was estimated using the intake rate and time spent drinking. The total
liquid load had the largest volume when ants used both drinking and grabbing in the same trip (Figure
5a). There was a significant interaction between sugar concentration and foraging action on the liquid
load (Table 2a, GLM, sugar $\times$ foraging action: $p < 0.01$). The crop load decreased with increasing
sugar concentration (Table 2b, LM: $p < 0.001$). On most trips, the total liquid load of grabbing was
7.5 times larger than the crop load (Table 2b, GLM, foraging action: $p < 0.01$). To examine how much
energy ants can bring back to the nest through these methods, we transformed the liquid load to sugar
load. We found that the difference in efficiency between drinking and grabbing increased with sugar
concentration (Figure 5b). There was a significant interaction between sugar concentration and
foraging action (Table 2a, GLM, sugar $\times$ foraging action: $p < 0.01$). While there were no significant
linear associations between sugar concentration and drinking (Table 2b, LM: $p = 0.39$), the total sugar
load acquired by grabbing significantly increased with sugar concentration (Table 2b, LM: $p < 0.001$).
These results suggest that grabbing, and consequently pseudotrophallaxis, are more efficient methods
to collect high-viscosity liquid than drinking and regurgitation.

Discussion

Behaviour in a given species is highly adapted to that organism's context, and these behavioural adaptations often involve precise forms of behavioural plasticity. In this study, we analysed the flexibility of foraging behaviours in response to biophysical constraints in *Diacamma* cf. *indicum*. The ant used two liquid collection actions – drinking and grabbing – when collecting liquid food. In this study, we aimed to quantify dynamic switching between of two types of liquid food collection used by a single ant species. *Diacamma* cf. *indicum* is a part of a clade of ants that rarely specialise on liquid food. Given this species' phylogenetic context, these behaviours are likely to be relatively recent specialisations [15]. However, it remains unclear when ants use pseudotrophallaxis as opposed to true trophallaxis to collect, transport, and share liquids, whether the use of these behaviours varies according to food quality, and whether one is an evolutionary step toward another.

Viscosity dictates behaviour

Here, we clearly observed mandibular gabbing and pseudotrophallaxis in the lab in *Diacamma* cf. *indicum* and we see that their use of this collection behaviour changes with viscosity. Our results are consistent with previous study in other ponerine ants, where ants stopped drinking [34] and tended to use mandibular grabbing at higher sugar concentrations (> 40%) [29]. Our work revealed that ants made this switch in collection mode according to viscosity (Figure 3b). Viscosity has been seen to reduce the liquid intake rates in many insects, including ants [23,34–37], consistent with our results (Figure S3). The viscosity of the solution makes drinking more time-consuming and this causes the ant to switch toward grabbing behaviour. When *Diacamma* ants used mandibular grabbing, total sugar load clearly increased at higher sugar concentrations. We also found that ants used mandibular grabbing after drinking liquid (Figure 3a). This maximises liquid load per trip because ants can transport internally and externally. Multiple trips can be costly as they involve loss of energy and increased predation risk.

Why do so few ants perform mandibular grabbing at high sugar concentration?

Mandibular grabbing and pseudotrophallaxis are mostly performed by ponerine ants (Ponerinae), with only a few noted exceptions in other major ant subfamilies [38] One possibility is that mandibular grabbing is a risky, but high pay-off collection method. For example, if ants encounter predators, they might not react quickly enough, ending up lose their mandibular droplet and/or being preyed upon because they are less agile. Ants might not use mandibular grabbing in dangerous sites where they encounter predators. On the contrary, if there are competitors around the food site, ants need to compete against other ant species. *Pheidole megacephala* soldiers reacted to the presence of competitors. The soldiers performed more mandibular grabbing on the territory of other ant species in order to rapidly gathering and transporting large loads of liquid food [38] In our study, we did not measure transportation time from feeding site to the nest or impact of predation or competition.

Future studies with different distances and ecological contexts while analysing transport time could

help elucidate the cost of transportation of pseudotrophallaxis through surface tension.

Morphological adaptations and biophysical constraints

Another possible reason why some ants use pseudotrophallaxis and others use true trophallaxis is that

many ants have internal morphological adaptations for liquid intake, storage and regurgitation, such as

an expandable gaster, an elastic crop and a highly developed proventriculus [7,15]. These are likely to

speed intake rate, allow for larger internal capacity and possibly allowing for greater flexibility

regarding the intake of high viscosity solutions [16,39]. Thus, species with these morphological

adaptations may not need pseudotrophallaxis.

A third possibility is that there might be biophysical restrictions on whether an ant can collect

a liquid drop between her mandibles. It is likely that small body size makes interactions with liquid

droplets more dangerous due to the strong forces of surface tension [40]. We observed that ants make

a 'hasty' motion at the end of the extraction of the droplet. The strong force is likely needed to

break the droplet away. If the ants pull with a constant force or weak force, they struggle to grab a

droplet. For small ants, it may be difficult to exert force to extract the droplet. Whether any

relationship exists between the ability to perform pseudotrophallaxis and biophysical restrictions,

related to body size, has not yet been studied.

Not only body size, but also head and mandible shape could be relevant to the performance of

liquid collection. Like *Platythyrea conradti*, soldiers of *Pheidole megacephala* use corporal

pseudotrophallaxis wherein liquid is held under the head and thorax through surface tension [38],

while minor workers do not perform pseudotrophallaxis. Soldier of *Pheidole* are much larger and have

disproportionately large heads compared with minors. These morphological traits of soldiers might be

related to their performance of corporal pseudotrophallaxis. Some ant species have unique and/or

exaggerated mandible shapes, for example, some army ants [41], desert ants [42], and trap-jaw ants

[43]. Given that the trap-jaw ant *Odontomachus* can collect droplets using uniquely shaped mandibles

[22], this indicates that several types of mandible shapes allow ants to hold liquid between mandibles.

However, the interaction between ant head and mouth-part morphology, body size and liquid

collection modes is unclear, making this area well poised for study from a biomechanics perspective.

Share with nestmates: diffusion in the colony

After foragers go back to the nest, they share the liquid food through trophallaxis including

(regurgitation), or through pseudotrophallaxis. Trophallaxis should more rapidly distribute food in the

colony because a receiver can become a donor and continue to distribute liquid food by regurgitation,

also allowing the formation of a more complex social network[44–46]. In the carpenter ant

Camponotus, foragers give food to a receiver, proportional to the available capacity in the receiver's

crop. This trophallactic interaction helps the forager to sense colony satiation level and decide when

to leave the nest and bring in more food [47]. In case of pseudotrophallaxis, a donor can provide

liquid food to several receivers at same time. However, the distribution dynamics of liquid by

pseudotrophallaxis has not been studied. We also do not know whether *Diacamma* ants share both

internal and external liquid foods to nestmates. The observation of liquid distribution in the focal

species that use both trophallaxis and pseudotrophallaxis is needed to understand these dynamics of

liquid distribution and the regulation of foraging effort.

Share with nestmates: social circulatory system

Trophallaxis allows for medium- to long-term food storage before redistribution while

pseudotrophallaxis does not. Thus, ecological contexts and environmental harshness may also tilt an

ant to engage in one behaviour or another. Another valuable feature of trophallaxis is that donors can

alter the contents of what they pass to nestmates, either through partial digestion or through more

complex signaling [48,49], which may bias a species or even a single ant to use one or another

behaviour. Recent studies reveal that ants' regurgitated fluid contained more than food [48,49]. For

example, trophallactic fluid in carpenter ants contains hormones, nestmate recognition cues, small

RNAs, and various proteins. In *Diacamma*, it is unclear whether foragers regurgitate the contents of

their crop during pseudotrophallaxis, or if they only regurgitate when they do trophallaxis. Future

studies could examine whether they add any endogenous materials during these behaviours.

Table 1. Influence of sugar concentration and foraging action on volume and handling time.
Results of generalized and general linear regression (a) and linear regression model (b) between two
explanatory variables and volume of carried or drunk water and handling time. Sugar concentration
water is 10, 20, 30, 40, 50, and 60% (w/w) and foraging actions are drinking (DR) and mandibular
grabbing (GR).
(a) generalized linear regression

	Explanatory variables	Estimate	Std. Error	p-value
Volume				
	Sugar concentration	-0.028	0.003	< 0.001
	Foraging actions (DR or GR)	-1.153	0.231	< 0.001
	Sugar * Type of actions	0.030	0.005	< 0.001
Handling time				
	Sugar concentration	-0.999	0.172	< 0.001
	Foraging actions (DR or GR)	-85.573	13.375	< 0.001
	Sugar * Foraging actions	1.118	0.320	< 0.001

351 (b) linear regression

		Explanatory variables	Estimate	Std. Error	t-value	p-value
Volume	Drinking	(Intercept)	1.965	0.139	13..656	< 0.001
		Sugar concentration	-0.028	0.003	-7.647	< 0.001
	Grabbing	(Intercept)	0.812	0.118	6.832	< 0.001
		Sugar concentration	-0.999	0.172	0.839	0.41
Handling time	Drinking	(Intercept)	91.41	7.964	11.477	< 0.001
		Sugar concentration	-9.999	2.112	-4.733	< 0.001
	Grabbing	(Intercept)	5.836	1.884	3.097	< 0.01
		Sugar concentration	1.1822	0.4349	2.718	< 0.01

Table 2. Influence of sugar concentration and foraging action on liquid and sugar load.

Results of generalized and general linear regression (a) and linear regression model (b) between two
explanatory variables and liquid load or sugar load per trip. Sugar concentration water is 10, 20, 30,
40, 50, and 60% (w/w) and foraging actions are drinking (DR) and both drinking and mandibular
grabbing (Both).

(a) generalized linear regression

		Explanatory variables	Estimate	Std. Error	p-value
Liquid load		(Intercept)	1.035	0.032	< 0.001
		Sugar concentration	-0.0005	0.0007	0.451
		Foraging actions (DR or Both)	-0.784	0.038	< 0.001
		Sugar * Type of actions	-0.003	0.0009	< 0.001
Sugar load		(Intercept)	-0.035	0.01	< 0.001
		Sugar concentration	0.011	0.0002	< 0.001
		Foraging actions (DR or Both)	0.066	0.011	< 0.001
		Sugar * Foraging actions	-0.011	0.0003	< 0.001

360 (b) linear regression model

		Explanatory variables	Estimate	Std. Error	t-value	p-value
Liquid load	Drinking	(Intercept)	0.25	0.017	14.207	< 0.001
		Sugar concentration	-0.004	0.0005	-7.558	< 0.001
	Both	(Intercept)	1.035	0.038	27.062	< 0.001
		Sugar concentration	-0.0005	0.0008	-0.648	0.518
Sugar load	Drinking	(Intercept)	0.031	0.004	7.388	< 0.001
		Sugar concentration	-0.0001	0.0001	-0.860	0.391
	Both	(Intercept)	-0.035	0.0135	-2.599	0.01
		Sugar concentration	0.01	0.0003	37.494	< 0.01

(a)   (b)

(c)   (d)

Figure 1. Ethogram of mandibular pseudotrophallaxis (social bucket).

An ant collects (a, b), transports (c) and shares (d) liquid food through pseudotrophallaxis. The ant

touches a solution with her antennae and mouthparts (tasting). After tasting, the individual opens her

mandibles to grab and pull at the liquid solution (a). The ant occasionally succeeds in collecting a

droplet of liquid between her mandibles (b). The ant returns to her nest (c). Inside the nest, the doner

ant (right) shares the droplet with other nestmates (d). When the receiver (left) begs, their antennae

move rapidly. Several nestmates can drink from the donor's droplet at the same time. Illustrations by

Ken Naganawa.

Figure 2. Effect of sugar concentration on foraging action.

(a) The volume of liquid drunk or grabbed, and (b) the time spent drinking and grabbing, by an ant for

each concentration of sugar water (% w/w). Black circles and grey triangles indicate drinking and

grabbing, respectively. The shading is 95% confidence limits for the regression line.

Figure 3. The proportion of pseudotrophallaxis depends on sugar concentration (a) and

viscosity (b).

Grabbing (white), both (grey) and drinking (black) indicate the behaviour of only mandibular

grabbing, mandibular grabbing after drinking, and only drinking. The x-axis is sorted by viscosity (b).

Different letters on the top of the bar mean they were significantly different at $p < 0.05$ (chi-square

test with Bonferroni correction).

386 **Figure 4. Dynamic viscosities at 25°C for different sugar water solutions as a function of the**

387 **sugar concentration (w/w) and viscosity-altering materials.**

388 The different colors correspond to pure water and viscosity-altered solutions. The error bars denote

389 the quality of the linear fit applied to stress versus shear rate experiments for each solution (see

390 Methods). Noted that the error margin was very small in our measurement.

Figure 5. Foraging efficiency of different modes of liquid collection and transport.

Estimated total liquid(a) and sugar load per trip (b) are estimated based on drinking time and foraging

action. Dark grey, grey, and light grey indicate the action of drinking, both and grabbing. The shading

is 95% confidence limits for the regression line. The volume of liquid by grabbing is the mean of

carried volume by an ant, calculated in Figure 2a.

Acknowledgments

This study was funded by JSPS KAKENHI Grant number JP20J01766 to HF and Swiss National Science Foundation Grant PR00P3_179776 to ACL. We are grateful to Ken Naganawa for his amazing illustrations. We would like to thank Dr. Lavergne François and Satomi Koga for their contribution to data collection. We also thank to all member of social fluids lab at Fribourg university for discussions about the study and Dr. Isaac Planas-Sitjà and Marie-Pierre Meurville for providing feedback on early drafts of the manuscript.

References

1. Pyke GH, Pulliam HR, Charnov EL. 2015 Optimal Foraging: A Selective Review of Theory and Tests. *https://doi.org/10.1086/409852* **52**, 137–154. (doi:10.1086/409852)

2. Stephens DW, Krebs JR. 1987 *Foraging Theory*. Princeton University Press. (doi:10.1515/9780691206790)

3. Buckley RC. 1987 Interactions involving plants, Homoptera, and ants. *Annual review of ecology and systematics. Vol. 18* , 111–135. (doi:10.1146/ANNUREV.ES.18.110187.000551)

4. Davidson DW, Cook SC, Snelling RR, Chua TH. 2003 Explaining the abundance of ants in lowland tropical rainforest canopies. *Science (1979)* **300**, 969–972. (doi:10.1126/SCIENCE.1082074/SUPPL_FILE/DAVIDSON.D.SOM.PDF)

5. Heil M, Mckey D. 2003 Protective Ant-Plant Interactions as Model Systems in Ecological and Evolutionary Research. *Source: Annual Review of Ecology, Evolution, and Systematics* **34**, 425–453. (doi:10.1146/132410)

6. Nelsen MP, Ree RH, Moreau CS. 2018 Ant–plant interactions evolved through increasing interdependence. *Proceedings of the National Academy of Sciences* **115**, 12253–12258. (doi:10.1073/pnas.1719794115)

7. Hölldobler B, Wilson EO. 1990 *The Ants*. Cambridge: Harvard University Press.

8. Wilson EO. 1990 *The insect societies (Harvard paperbacks)*.

9. Calixto ES, Lange D, Del-Claro K. 2021 Net benefits of a mutualism: Influence of the quality of extrafloral nectar on the colony fitness of a mutualistic ant. *Biotropica* **53**, 846–856. (doi:10.1111/BTP.12925)

10. Pekas A, Tena A, Aguilar A, Garcia-Marí F. 2011 Spatio-temporal patterns and interactions with honeydew-producing Hemiptera of ants in a Mediterranean citrus orchard. *Agric For Entomol* **13**, 89–97. (doi:10.1111/j.1461-9563.2010.00501.x)

11. Pringle EG. 2021 Ant-Hemiptera Associations. *Encyclopedia of Social Insects* , 45–48. (doi:10.1007/978-3-030-28102-1_8)

12. Styrsky JD, Eubanks MD. 2006 Ecological consequences of interactions between ants and honeydew-producing insects. *Proceedings of the Royal Society B: Biological Sciences* **274**, 151–164. (doi:10.1098/RSPB.2006.3701)

13. Sempo G, Detrain C. 2004 Between-species differences of behavioural repertoire of castes in the ant genus Pheidole: A methodological artefact? *Insectes Soc* (doi:10.1007/s00040-003-0704-2)

14. Richard FJ, Dejean A, Lachaud JP. 2004 Sugary food robbing in ants: a case of temporal cleptobiosis. *C R Biol* **327**, 509–517. (doi:10.1016/J.CRVI.2004.03.002)

15. Meurville MP, LeBoeuf AC. 2021 Trophallaxis: the functions and evolution of social fluid exchange in ant colonies (Hymenoptera: Formicidae). *Myrmecol News* **31**, 1–30. (doi:10.25849/MYRMECOL.NEWS_031:001)

443   16.   Davidson DW, Cook SC, Snelling RR. 2004 Liquid-feeding performances of ants
444   (Formicidae): Ecological and evolutionary implications. *Oecologia* **139**, 255–266.
445   (doi:10.1007/S00442-004-1508-4/TABLES/4)

446   17.   Eisner T. 1957 A comparative morphological study of the proventriculus of ants
447   (Hymenoptera: Formicidae). *Bull Mus Comp Zool* **116**, 437–490.

448   18.   Eisner T, Brown WLJ. 1958 The evolution and social significance of the ant proventriculus.
449   *Proceedings Tenth International Congress of Entomology* **2**, 503–508.

450   19.   Robinson EJH, Feinerman O, Franks NR. 2009 Flexible task allocation and the organization of
451   work in ants. *Proceedings of the Royal Society B: Biological Sciences* **276**, 4373–4380.
452   (doi:10.1098/rspb.2009.1244)

453   20.   Hölldobler B. 1985 Liquid food transmission and antennation signals in ponerine ants. *Isr J*
454   *Entomol* , 89–99.

455   21.   A. Dejean, J.P. Suzzoni. 1997 Surface Tension Strengths in the Service of a Ponerine Ant: a
456   New Kind of Nectar Transport. *Naturwissenschaften* **84**, 76–79.

457   22.   Lachaud Jean-Paul, Alain Dejean. 1991 Food sharing in Odontomachus troglodytes (Santschi):
458   a behavioral intermediate stage in the evolution of social food exchange in ants. *An Biol* **17**,
459   53–61.

460   23.   Lois-Milevicich J, Schilman PE, Josens R. 2021 Viscosity as a key factor in decision making
461   of nectar feeding ants. *J Insect Physiol* **128**, 104164. (doi:10.1016/j.jinsphys.2020.104164)

462   24.   Bonser R, Wright PJ, Bament S, Chukwu UO. 1998 Optimal patch use by foraging workers of
463   Lasius fuliginosus, L. niger and Myrmica ruginodis. *Ecol Entomol* **23**, 15–21.
464   (doi:10.1046/J.1365-2311.1998.00103.X)

465   25.   Dejean A, Solano PJ, Ayroles J, Corbara B, Orivel J. 2005 Arboreal ants build traps to capture
466   prey. *Nature* **434**, 973–973. (doi:10.1038/434973a)

467   26.   Detrain C, Prieur J. 2014 Sensitivity and feeding efficiency of the black garden ant Lasius
468   niger to sugar resources. *J Insect Physiol* **64**, 74–80. (doi:10.1016/J.JINSPHYS.2014.03.010)

469   27.   Falibene A, de Figueiredo Gontijo A, Josens R. 2009 Sucking pump activity in feeding
470   behaviour regulation in carpenter ants. *J Insect Physiol* **55**, 518–524.
471   (doi:10.1016/J.JINSPHYS.2009.01.015)

472   28.   Fujioka H, Okada Y. 2019 Liquid exchange via stomodeal trophallaxis in the ponerine ant
473   Diacamma sp. from Japan. *J Ethol* **37**, 371–375. (doi:10.1007/s10164-019-00602-9)

474   29.   Jandt J, Larson HK, Tellez P, McGlynn TP. 2013 To drink or grasp? How bullet ants
475   (Paraponera clavata) differentiate between sugars and proteins in liquids. *Naturwissenschaften*
476   **100**, 1109–1114. (doi:10.1007/S00114-013-1109-3/FIGURES/2)

477   30.   Blüthgen N, Menzel F, Blüthgen N. 2006 Measuring specialization in species interaction
478   networks. *BMC Ecol* **6**, 9. (doi:https://doi.org/10.1186/1472-6785-6-9)

31.   VanGinkel CG, Gayton S. 1996 The biodegradability and nontoxicity of carboxymethyl
cellulose (DS 0.7) and intermediates. *Environ Toxicol Chem* **15**, 270–274.
(doi:10.1002/ETC.5620150307)

32.   Rhee BO, Lee S. 1999 Evaluation on Accuracy of the Rheological Data of PIM Feedstocks.
Journal of the Japan Society of Powder and Powder Metallurgy **46**, 830–836.
(doi:10.2497/jjspm.46.830)

33.   Josens RB, Farina WM, Roces F. 1998 Nectar feeding by the ant Camponotus mus: intake rate
and crop filling as a function of sucrose concentration. *J Insect Physiol* **44**, 579–585.
(doi:10.1016/S0022-1910(98)00053-5)

34.   Ávila Núñez JL, Naya M, Calcagno-Pissarelli MP, Otero LD. 2011 Behaviour of
Odontomachus chelifer (Latreille) (Formicidae: Ponerinae) Feeding on Sugary Liquids. *J*
Insect Behav **24**, 220–229. (doi:10.1007/S10905-010-9249-1/FIGURES/4)

35.   Borrell BJ. 2006 Mechanics of nectar feeding in the orchid bee Euglossa imperialis: pressure,
viscosity and flow. *Journal of Experimental Biology* **209**, 4901–4907.
(doi:10.1242/JEB.02593)

36.   Kim W, Gilet T, Bush JWM. 2011 Optimal concentrations in nectar feeding. *Proc Natl Acad*
Sci U S A **108**, 16618–16621. (doi:10.1073/PNAS.1108642108)

37.   Nicolson SW, de Veer L, Köhler A, Pirk CWW. 2013 Honeybees prefer warmer nectar and
less viscous nectar, regardless of sugar concentration. *Proceedings of the Royal Society B:*
Biological Sciences **280**. (doi:10.1098/RSPB.2013.1597)

38.   Dejean A, le Breton J, Suzzoni JP, Orivel J, Saux-Moreau C. 2005 Influence of interspecific
competition on the recruitment behavior and liquid food transport in the tramp ant species
Pheidole megacephala. *Naturwissenschaften* **92**, 324–327. (doi:10.1007/S00114-005-0632-
2/FIGURES/2)

39.   Josens RB, Farina WM, Roces F. 1998 Nectar feeding by the ant Camponotus mus: intake rate
and crop filling as a function of sucrose concentration. *J Insect Physiol* **44**, 579–585.
(doi:10.1016/S0022-1910(98)00053-5)

40.   Zhou A, Du Y, Chen J. 2020 Ants adjust their tool use strategy in response to foraging risk.
Funct Ecol **34**, 2524–2535. (doi:10.1111/1365-2435.13671)

41.   POWELL S, FRANKS NR. 2006 Ecology and the evolution of worker morphological
diversity: a comparative analysis with Eciton army ants. *Funct Ecol* **20**, 1105–1114.
(doi:10.1111/j.1365-2435.2006.01184.x)

42.   Molet M, Maicher V, Peeters C. 2014 Bigger Helpers in the Ant Cataglyphis bombycina:
Increased Worker Polymorphism or Novel Soldier Caste? *PLoS One* **9**, e84929.
(doi:10.1371/journal.pone.0084929)

43.   Larabee FJ, Suarez A v. 2014 The evolution and functional morphology of trap-jaw ants
(Hymenoptera: Formicidae). *Myrmecological News* **20**, 25–36.

44. Buffin A, Mailleux AC, Detrain C, Deneubourg JL. 2011 Trophallaxis in Lasius niger: A
variable frequency and constant duration for three food types. *Insectes Soc* **58**, 177–183.
(doi:10.1007/S00040-010-0133-Y/TABLES/2)

45. Greenwald E, Segre E, Feinerman O. 2015 Ant trophallactic networks: Simultaneous
measurement of interaction patterns and food dissemination. *Sci Rep* (doi:10.1038/srep12496)

46. Bles O, Deneubourg J-L, Nicolis SC. 2018 Food dissemination in ants: Robustness of the
trophallactic network against resource quality. *Journal of Experimental Biology* **221**.
(doi:10.1242/jeb.192492)

47. Greenwald EE, Baltiansky L, Feinerman O. 2018 Individual crop loads provide local control
for collective food intake in ant colonies. *Elife* **7**, e31730. (doi:10.7554/eLife.31730)

48. Leboeuf AC *et al.* 2016 Oral transfer of chemical cues, growth proteins and hormones in social
insects. *Elife* **5**. (doi:10.7554/eLife.20375)

49. Hakala SM, Meurville MP, Stumpe M, Leboeuf AC. 2021 Biomarkers in a socially exchanged
fluid reflect colony maturity, behavior, and distributed metabolism. *Elife* **10**.
(doi:10.7554/ELIFE.74005)

547 **Supp. Table 1. Weight of each concentration of sugar water per 1 μ L.**

548 Weight is the average weight of corresponding 1 μ L sugar water (n=10).

Sugar concentration % (w/w)	Weight (mg/μL)
10	0.955
20	0.968
30	1.020
40	1.107
50	1.143
60	1.094

550 **Supp. Table 2. Viscosity.**

551 CMC is a non-toxic inert viscosity modifier. 10 CMC and 30 CMC are 10% sugar solution with CMC

552 0.25% and 30% sugar solution with CMC 0.25% w/w.

Sugar concentration % (w/w)	Viscosity (mPa•sec)
10	1.158 ± 0.002
20	1.656 ± 0.002
30	2.641 ± 0.003
40	5.308 ± 0.005
50	11.919 ± 0.009
60	47.34 ± 0.004
10 CMC	8.97 ± 0.002
30 CMC	24.6 ± 0.2

554 **Supp. Table 3. Intake rate of each concentration of sugar water.**

555 Intake rate is the slope (μL/sec) of the linear regression of crop load and corresponding feeding time.

Sugar concentration % (w/w)	Intake rate (μL/sec)
10	0.00245
20	0.00355
30	0.00206
40	0.00136
50	0.00091
60	0.00027

558 **Supp. Figure 1. Experimental arena design.**

559 An artificial nest (diameter = 9 cm) is placed on a container (48* 26* 10 cm). The nest was covered

560 with a red film to darken inside the nest. The distance between the nest and food is about 30 cm.

561 Fluon was applied to the wall of the foraging arena to prevent ants from escaping.

Supp. Figure 2. Preference and foraging actions of different sugar concentrations on non-thirsty and thirsty colonies.

Colonies were starved for 3-4 hours before the experiment (thirsty condition). Ants were offered three different sugar concentrations: 0, 30, and 60 % w/w. (a) Total foraging bouts and (b) the number of pseudotrophallaxis that an ant grabbed a drop by mandibles and transported it were counted during 30-min observation. (c) The proportion of pseudotrophallaxis ants used was calculated for each colony. The color of white, grey, and dark grey represents 0, 30, and 60 % w/w sucrose concentrations. Asterisk indicates significant difference (Tukey–Kramer test, * $p < 0.05$, ** $p < 0.01$). Ants clearly preferred sugar water compared to water. The number of foraging trips was high in the thirsty condition.

Supp. Figure 3. Intake rate, plotting crop load (µL), and corresponding drinking time (sec).

(a-f) Each column indicates different sugar concentrations from 10% to 60%. Linear regression

models are shown inside the column.