Diacamma **ants adjust liquid foraging strategies in response to biophysical constraints**

Haruna FUJIOKA[1,2*], Manon Marchand[3], Adria C. LeBoeuf[2*]

1. Faculty of Environmental and Life Science, Okayama University, Okayama, Japan.

2. Department of Biology, University of Fribourg, Fribourg, Switzerland.

3. Department of Physics, University of Fribourg, Fribourg, Switzerland.

Author correspondence* HF: fujioka.ha@gmail.com, AL: adria.leboeuf@gmail.com

Abstract

Ant foragers need to provide food to the rest of the colony, which often requires transport

over long distances. Foraging for liquid is especially challenging because it is difficult to transport

and share. Many social insects store liquid food inside the crop to transport it to the nest, and then

regurgitate this fluid to distribute it to nestmates through a behaviour called trophallaxis. Some ants

instead transport fluids with a riskier behaviour called pseudotrophallaxis – holding a drop of liquid

between the mandibles through surface tension. Ants can then share this droplet with nestmates

without ingestion or regurgitation. Here, we hypothesised that ants optimise their liquid-collection

approach depending on viscosity. Working with a ponerine ant that uses both trophallaxis and

pseudotrophallaxis, we investigated why each liquid-collection behaviour might be favoured under

different conditions by measuring biophysical properties, collection time, and reaction to food quality

for typical and viscosity-altered sucrose solutions. We found that ants can collect more liquid food per

unit time by mandibular grabbing than by drinking. At high viscosities, which in nature correspond to

high sugar concentrations, ants switched their liquid collection method to mandibular grabbing in

response to viscosity, and not to sweetness. Our results demonstrate that ants change their transport

and sharing methods according to viscosity – a proxy for sugar concentration in nature – increasing

the mass of sugar returned to the nest per trip.

keywords: mandibular pseudotrophallaxis, social bucket, liquid transportation, liquid collection,

optimal foraging theory, biophysics

Introduction

Efficient foraging is crucial for animals to survive, grow and reproduce. Organisms need to balance energy spent with energy gained [1,2]. Optimal foraging theory assumes that animals' foraging decision making has evolved to the point that the fitness of individuals has been maximised. Foraging provides energy to survive and reproduce; however, it has the cost of exposing the individual to being preyed upon by other animals and it costs energy, for example, due to the time spent exploiting, processing and transporting food.

Animals have a wide variety of foraging strategies. Ants are central-place foragers with diverse diets ranging from complete herbivory to complete predation[3–6]. Morphological and phylogenetic evidence suggests that the ancestral ant was a predator, and transitions to herbivory occurred several times in predatory lineages [5,7,8]. Plant-based food sources, such as plant nectar [9] or honeydew excreted by sapsucking Hemiptera and scale insects [10–12], are rich in carbohydrates. In many species and especially in ecologically dominant ant lineages, these sugary liquids are ants' main source of energy [13]. Additionally, liquid resources, such as honeydew or nectar, are less ephemeral relative to insect prey and incur fewer risks for ant foragers relative to hunting. Thus, the use of plant-based food sources may lead to lower foraging time and less risk for foragers per calorie returned to the nest.

Transportation of liquid food is a foraging challenge. Many ants and bees transport liquid stored inside their crop, where it cannot easily be lost or stolen [14] during transport. Foragers regurgitate this fluid to distribute it to nestmates through a behaviour called trophallaxis. Many liquid-feeding ants have acquired morphological adaptations for trophallaxis. The ant crop is separated from their midgut by a variably developed proventriculus, which allows the crop to store a large amount of liquid in some species [15]. The structure of proventriculus varies considerably across taxa [16–18], and liquid-feeding ants often have a more elaborate proventriculus. The gaster and crop also need to be expandable to best store liquid food, either temporarily for transport or over the long term in the case of repletes. The extreme example of morphological specialisation are honeypot ants *Myrmecocystus* (Formicinae), where replete workers have a massive ball-like distended gaster full of food to the point where they can barely move [19]. Such species rely on trophallaxis to redistribute food from the repletes to the rest of the colony. Although trophallaxis is considered a safe and reliable liquid transportation method for ants, the crop load (i.e., liquid food intake) strongly depends on these morphological constraints.

Some ants do not have these morphological specialisations but nonetheless consume liquid food: Ectatomminae (*Ectatomma*), Ponerinae (*Diacamma, Neoponera, Odontomachus, Paraponera, Pachycondyla, Rhytidoponera*). These ants typically use mandibular pseudotrophallaxis (hereafter called pseudotrophallaxis) as their method of liquid transport [15,20]. Instead of storing liquid inside the crop, foragers hold liquid food between their mandibles where it forms a droplet maintaining its round shape through surface tension. After foragers return to the nest, they pass the liquid food to nestmates without regurgitation. Previous studies in ponerine ants have reported how this behaviour allows liquids to be distributed in the nest [21,22]. This liquid transport method is sometimes referred to as the 'social bucket' method and has been suggested to be an evolutionary precursor to 'true' trophallaxis [7].

Handling time is crucial for efficient foraging. For liquid food, the handling time includes both the speed of food collection (i.e., drinking time or grabbing time) and the transport time to the nest. Drinking time in ants has been shown to depend on food quality such as sugar concentration and viscosity itself [23]. Previous studies found that drinking time increased linearly with increasing sugar concentration [23–27]. Individuals need to decide when to stop drinking, considering the balance between energy gain and predation risk. The handling time of pseudotrophallaxis has not been investigated. Regarding transport success, once foragers store food in the crop, they can transport the liquid food safely back to the nest. When using pseudotrophallaxis, there is the possibility of losing the liquid food along the return path. Also, keeping mandibles open adequately may increase the likelihood of predation.

Some ants use both behaviours. The ponerine ant *Diacamma* cf. *indicum* from Japan performs both trophallaxis and pseudotrophallaxis [28], and has a simple proventricular morphology and a rigid, non-extensible gaster. Thus, *Diacamma* cf. *indicum* is an ideal model species to investigate efficient foraging strategies regarding liquid food because it allows us to investigate foraging strategies without morphological specialisation.

The aim of the present study is to reveal what leads ants to use the collection mode of mandibular grabbing instead of drinking and whether ants' liquid collection modes mechanisms maximise calorie intake rates per foraging trip. We hypothesise that viscosity triggers a switch in collection behaviour between drinking and mandibular grabbing, where mandibular grabbing is more efficient to collect high viscosity solutions. To test this hypothesis, we conducted experiments to investigate 1) physical properties of the liquids such as capillary length, viscosity and surface tension,

2) volume and speed of liquid food collection depending on sugar concentration, 3) whether ants

change their transportation method depending on sugar concentration or viscosity by dissociating

viscosity from sugar concentration, and 4) the foraging efficiency for each approach using walking

speed and the success rate of mandibular transport to estimate the total amount of sugar carried per

trip.

Methods

Colony collection and colony keeping

Colonies of *Diacamma* cf. *indicum* from Japan were collected from Kenmin-no-mori (Onna)

and Sueyoshi park (Naha), Okinawa, Japan. The colonies were kept in plastic artificial nests filled

with moistened plaster (9 cm diameter □×□1.5 cm height). Each colony contained a mated gemma-

possessing female (i.e., functional queen or gamergate), 50–150 workers, and brood. The artificial

nests (90 mm in diameter) were placed in a plastic arena (diameter: cm, height: cm). Nests were

maintained at 25 °C under a 12h/12h light-dark regime (light phase: 0800–2000 hours). Reared

colonies were fed with chopped frozen crickets three times per week. Water and 10% sugar water

were provided *ad libitum*. All behavioural experiments were conducted between 12:00-19:00 in well-

lit conditions at 25□ and 50-60% humidity.

Behaviour Definitions

The ethogram of the social bucket method (encompassing mandibular grabbing, transport and

pseudotrophallaxis) of *Diacamma* ant is shown in Figure 1. Based on a previous study [22] two

behaviours for liquid feeding and collection were defined 1) drinking: individuals drink (i.e.,

mouthpart, labrum, attached to a liquid solution), 2) grabbing: individuals open mandibles to grab and

pull at a liquid solution, occasionally succeeding in collecting a droplet.

Dynamic viscosity measurement

We prepared six sugar solutions, 10, 20, 30, 40, 50, 60 % w/w. The concentrations we used

were within the range reported for natural nectar sources (extrafloral nectars: 4.7–76% w/w [29]). In

addition, we used two viscosity-altered solutions using carboxymethylcellulose sodium salt (CMC)

(Medium viscosity, Sigma-Aldrich), 10% sugar solution with 0.25% CMC (10CMC), and 30% sugar

solution with 0.25% CMC (30CMC). The CMC is a non-toxic inert viscosity modifier [30]. By

adding this product, we increase the viscosity of the solution without changing its sugar concentration.

We confirmed that the sugar concentration of CMC additive solution was not changed using a

saccharimeter (Refractometer RBR32-ATC).

We measured these solutions' dynamic viscosity in a commercial stress-controlled rheometer

(ANTON PAAR MCR 300) with a plate-plate geometry of 5 cm diameter and 0.5 mm gap. The

bottom plate was roughened by sandblasting to prevent slip artefacts and the temperature was fixed at

25°C by a Peltier hood. We applied constant shear rates, and stress was computed when the steady-

state regime had been reached for each shear rate. The resulting stress versus shear rate experiments

exhibited linear behaviour characteristic of a Newtonian fluid. This was expected for sucrose/water

solutions since sucrose is a small molecule that dissolves well in water, though this was not obvious

for the CMC polymer solutions [31]. The dynamic viscosity was directly read from the slope

evaluated by least-square minimisation for each sample, for more detail see [32].

Measurement of surface tension and capillary length

Surface tension is the energy it costs to create a surface of a liquid. We measured the surface

tension of the different sugar/water concentrations with a commercial Du Noüy apparatus (Kibron

EZPIPlus). The laboratory room was regulated at 25°C +/- 1°C. Each measurement was repeated four

times and the reported error is the standard deviation.

Capillary length is the size above which a drop of liquid can no longer sustain its own shape

by surface tension but starts to be deformed and to flow under its own weight. In order to obtain the

capillary length, we used tabulated density measurements [33]. The capillary length is computed from

surface tension and density through the following formula

$\lambda_c = \sqrt{\dfrac{\gamma}{\rho g}}$

With γ the surface tension, ρ the density and g the gravitational acceleration [34].

Drinking speed

Drinking speed was calculated as the slope (μL/sec) of the linear regression of *crop load* and

corresponding *drinking time* as intake rate [35] . Thus, we measured these two values (crop load and

drinking time) for drinking events of 118 ants from six colonies, drinking seven different solutions: 10,

20, 30, 40, 50, 60% sugar w/w, and 10CMC. The crop load (volume of sugar water drunk) was

estimated as the difference in body mass before and after drinking (using a Fisherbrand analytical

balance, to 0.1 mg). We chose ants outside of the nest and measured their body mass (before drinking).

Then, each individual was separately placed into a plastic box (4.5 × 4.5 × 2 cm), containing a

drop (approximately 100 μl) of sugar water. To measure the corresponding drinking time, we video-

recorded the behaviour of ants when drinking. Ants were offered one of seven different solutions: 10,

20, 30, 40, 50, 60% sugar w/w, and 10CMC. When ants stopped drinking, we wiped the liquid off the

mouthpart with a tissue and measured the ant's body mass to determine the volume of liquid food

inside its crop. We estimated the volume (μL) of sugar water using the average weight (density) of the

relevant sugar solution per 1 μl (Supp. table 1). We weighed 100 μL droplets using a micropipette and

an electronic balance (Semi-Micro Analytical Balances, GH-202, A&D Company).

We measured the time spent in contact with the droplet by manual observation as the

corresponding drinking time. The drinking time was the total time the ant's mouthpart was attached to

a liquid solution. We plotted the crop load and corresponding drinking time and calculated the slope

(μL/sec) of the linear regression as the drinking speed (Supp. figure S2). Individuals were never tested

more than once a day.

Measurement of volume of sugar water internally and externally carried

We measured the volume of water internally (crop load) and externally carried (mandible

droplet), collected by 207 ants from 4 colonies. We measure the crop load, using the same procedure

as 'drinking speed'. Ants were offered one of six different sugar concentrations: 10, 20, 30, 40, 50,

60% w/w.

Colonies were starved for 3-4 hours before experiments (starvation time based on preliminary

behavioural observations, data not shown). The volume of sugar water carried by mandibular

grabbing was measured with a microcapillary tube. We placed an artificial nest on one side and a

plate-shaped feeder (40 × 40 mm) on the other side of the foraging arena (460 × 260 × 100 mm)

(Supp. figure 1). Ants were offered one of six different sugar concentrations: 10, 20, 30, 40, 50, 60 %

w/w. The ants freely accessed the offered food. After an ant succeeded in grabbing a droplet of the

solution, we collected the droplet with a microcapillary tube during a return trip to the nest. The

weight of the dispensed volume was measured to calculate the volume of sugar water the ant carried.

We estimated the volume (μL) of sugar water using the average weight of the relevant sugar water

(Supp. table 1).

Grabbing time, drinking time and foraging actions in the foraging arena

We measured accumulated grabbing/drinking time during 1047 foraging events of 572 ants

from nine colonies. To test the effects of sweetness and viscosity on the foraging methods used, we

used seven different sugar concentrations: 10, 20, 30, 40, 50, 60 % w/w, and 10CMC as a viscosity-

altered solution. When foragers found 10CMC, they started tasting and drinking as usual. Refusal or

avoidance-like behaviours were not observed toward 10CMC.

We placed an artificial nest on one side and a plate-shaped feeder (40 × 40 mm) on the other

side of the foraging arena (460 × 260 × 100 mm) (Supp. figure 1). Ants were offered one of the

seven sugar concentrations: 10, 20, 30, 40, 50, 60 % w/w, and 10CMC. We video-recorded the area

around the sugar water droplet for 1 hour. We manually recorded one type of foraging action that

foragers had with the droplet, 'only drinking,' 'grabbing after drinking (both),' or 'only grabbing.'

Accumulated grabbing/drinking time for each foraging trip was calculated by an observer analysing

the videos. The drinking time (sec) is the accumulated time an ant's mouthparts were attached to the

solution. We defined that the start of grabbing was when they opened the mandibles because their

mandibles are closed when this focal ant species drinks. The endpoint was when the ant succeeded in

grabbing the droplet (e.g., detached droplet from solution). When ants repeated the grabbing, we

included grabbing time until the last successful attempt.

Walking speed and success rate of mandibular transport

We measured ants' walking speed and success rate using each of the modes of liquid transport.

We placed an artificial nest on one side and a plate-shaped feeder (40 × 40 mm) on the other side of

the foraging arena (387 × 267 × 65 mm). Ants were offered one of the two sugar concentrations: 10%

w/w (low sugar/low viscosity) and 50 % w/w (high sugar/high viscosity). We continuously recorded

the foraging arena for 1 hour. The images were captured from above using a web camera (Logicool;

HD Pro Webcam C920t) at 10 frames per second. We automatically extracted the coordinates of each

individual using the video-tracking system, UMATracker [36]. We categorised ants' behaviours into

1) empty walking speed, 2) crop-full walking speed, and 3) crop- and mandible-full walking speed by

visual observation. We obtained walking speeds for 80 foraging trips from four colonies.

To investigate the success rate of mandibular liquid transport, we observed 30 foraging trips

from the feeder to the nest from three colonies and investigated whether ants lost the mandibular

droplet during the return trip. The size of the foraging arena ($387 \times 267 \times 65$ mm) and two types of

sugar water (10% and 50%) were used. If an ant entered the nest without dropping the mandibular

droplet, we recorded that as a foraging success. Returning to the nest after drinking was always

considered a foraging success as the liquid cannot be lost.

Estimation of sugar intake

To assess foraging efficiency, we estimated total sugar intake per foraging trip by combining

several forms of data: 1) observation of foraging action ants performed, 2) drinking or grabbing time,

3) drinking speed, and 4) average grabbing volume. First, we calculated the total liquid load per trip

based on the foraging action used (see the section '*Grabbing time, drinking time and foraging actions*

in the foraging arena' and Figure 4). The total crop load per trip was estimated by multiplying the

drinking speed (see the section '*Drinking speed*', Supp. figure 2) by the accumulated time drinking in

the foraging arena (see the section '*Grabbing time, drinking time and foraging actions in the foraging*

arena', Figure 3b). The mandible load (collected by grabbing) was defined as the average volume of

liquid carried for each sugar concentration in the experiment '*Measurement of volume of sugar water*

internally and externally carried' (see Figure 3a). When ants performed both drinking and grabbing,

we summed the average volume of liquid carried and the estimated total crop load. Using the average

weight of sugar water (Supp. table 1), we converted the total liquid load per trip (μL) to the weight of

the liquid load (mg). From the liquid load weight (mg), we calculated the total sugar intake per trip

(mg) for each sugar concentration.

Statistical analysis

Linear regression models were used to calculate the drinking speed. We compared regression

slopes using a Wilcoxon test with Bonferroni correction. Generalised linear regression models were

used to investigate the relationship between the liquid volume or loading speed with the food quality

variables and foraging actions. Pairwise chi-square tests with a Bonferroni correction were used for

comparing foraging action on different sugar concentrations. The relationship of the load with the

food quality variables and foraging actions was analysed by two-way analysis of variance (ANOVA),

followed by Tukey's honestly significant difference (HSD) test for multiple comparison of means. A

significance level of 5% was used in all comparisons. All analyses were run in R studio 2022.02.3

(package: ggplot2), except for viscosity measurements and drinking speed that were analysed in

python 3.9 and the linear regressions were done with the module scipy.optimize v1.7.1. [37].

Results

Diacamma ants forage for sugary liquids and use two forms of behaviour to collect liquid –

drinking and mandibular grabbing. To understand whether ants alter their liquid collection and

transport behaviour according to the contents and properties of the liquid they are transporting, we

measured multiple variables to find what ants are optimising: volume carried, time spent collecting,

frequencies of different foraging actions, walking speed, success rate of transport, biophysical

properties of the liquids and sugar load acquired per trip.

First, we analysed three liquid properties of a range of ecologically relevant sugary liquids

from 0-60% sucrose: dynamic viscosity, surface tension, and capillary length. Dynamic viscosity

increased with increasing sugar water concentration (Figure 2a). We altered the viscosity of a low-

sugar solution using the viscosity-modifying additive CMC (carboxymethylcellulose sodium salt).

The viscosity of 10% sugar water with CMC (10CMC) was comparable to the viscosity of 40-50%

sugar water (Supp. Table 2, Figure 2a). We measured an increase in surface tension (mN/m) with

increasing sugar concentration while the capillary length was not significantly affected (Table 1).

Combining video recording and weight measurements of ants before and after drinking, we

determined that drinking speed is inversely proportional to the viscosity of the sugar water solution

imbibed (Supp. Table 3, Figure 2b). If we approximate flow as if the alimentary canals of ants were

cylindrical pipes, we can use the Hagen-Poiseuille equation

$$1/Q = \frac{8\pi\eta L}{\Delta p\ A^2},$$

where Q is the drinking speed, L and A the length and cross-sectional area of the ant throats,

η the viscosity of the solutions and Δp the pressure drop that the ants generate by suction [38]. Thus,

high viscosity makes drinking more time-consuming and may cause the ants to switch toward

grabbing behaviour. Indeed, drinking speed for 10CMC was slower than that for 10% sugar water

(Wilcoxon test with Bonferroni correction, 10CMC vs. 10%: p = 0.013) and comparable to drinking

speed for 40% and 50% sugar water (Wilcoxon test with Bonferroni correction, 10CMC vs. 40: p =

0.58; 10CMC vs. 50: p = 0.37). Drinking speed for 10CMC falls onto the linear adjustment for

drinking speed for solutions without additives (Figure 2b).

We investigated the volume of liquid food collected by ants using two different collection

methods, feeding on different sugar concentrations (Figure 3). For the volume ants collected through

drinking or mandibular grabbing, we observed an interaction between sugar concentration and

foraging action (Figure 3a, Table 2a, GLM, sugar × collection method: $p < 0.001$), and therefore we

analysed the foraging actions separately. When ants drank liquid, the amount imbibed decreased as

viscosity increased (Figure 3b, Table 2b, LM: $p < 0.001$). For the amount of liquid grabbed within the

mandibles, there was no significant trend across the different viscosity (Figure 3c, Table 2b, LM: $p =$

0.14).

Since the volume of droplets collected by mandibular grabbing were more or less constant

(~1µL) at all sugar concentrations (Fig 3a), the energy required to extract the droplets should be solely

a function of the surface tension, where it should be easier to grab lower sugar concentration solutions

due to the lower surface tension (Table 1). In terms of capillary length, our measurements showed that

the increase in surface tension with the addition of sugar is balanced by the increase in density (Table

1). These values lead to an estimate of the maximum transportable volume as a drop of approximately

80 ± 10 µL, estimated from the capillary length translated into the volume of a sphere. This volume is

more than one order of magnitude above the maximum volume measured in this study (Table 1,

Figure 3a). Thus, at all sugar concentrations, the droplets transported by grabbing could sustain their

own shape, were not impacted by gravity and did not flow during transport. This suggests that the

limiting factor of the grabbing volume is not biophysical liquid properties, but more likely the ants'

morphological constraints.

Regarding the collection time for drinking or mandible grabbing, we found an interaction

between viscosity and foraging action (Table 2a, GLM, viscosity × foraging action: $p < 0.001$). When

ants drank, the drinking time decreased with increasing viscosity (Figure 3b, Table 2b, LM: $p <$

0.001). There was no linear relationship between grabbing time and viscosity (Figure 3c, Table 2b,

LM: $p = 0.14$). All in all, mandible grabbing took less time to collect liquid when compared to

drinking. Thus, mandible grabbing is a more effective method to collect liquid food, in terms of

volume per foraging time, across different sugar concentrations.

In spite of this, ants often collected sugar water in their mandibles after drinking and rarely

performed only mandible grabbing without drinking (Figure 4). The proportion of these foraging

actions was significantly different across sugar concentrations (Figure 4, $p < 0.05$, chi-sq test with

Bonferroni correction). The proportion of mandible grabbing after drinking (both) and mandible

grabbing alone (both of which result in pseudotrophallaxis) increased with increasing sugar

concentration. This indicates that ants switch to grabbing and pseudotrophallaxis when they feed on

liquid food with higher concentrations of sugar. This could come about because this high-sugar food

is more valuable or because high-viscosity liquids are difficult for them to drink, as indicated by the

drinking speed measurements (Figure 2). To test whether ants react to changes in sugar concentration

or viscosity, we offered ants our viscosity-altered solution 10CMC and observed which foraging

action was used. The proportion of ants drinking significantly decreased compared to those drinking

10% sugar water (Figure 4, $p < 0.05$, chi-sq test with Bonferroni correction). The proportion of ants

drinking 10CMC was equivalent to those drinking a high viscosity 50% sugar solution (Figure 4).

This result shows that ants switch collection methods in response to viscosity, and not to sweetness.

To investigate whether this transition toward grabbing over drinking with increasing viscosity

is related to foraging efficiency, we estimated the liquid load and sugar load per trip across the

different sugar concentrations. The liquid load per trip had the largest volume when ants used both

drinking and grabbing in the same trip (Figure 5a). There was a significant interaction between sugar

concentration and foraging action on the liquid load (Table 3, ANOVA, sugar × foraging action: $p <$

0.01). The total liquid load when grabbing was larger than the crop load (Table 3, ANOVA, foraging

action: $p < 0.05$). The crop load decreased with increasing sugar concentration (Tukey's HSD Test, p

< 0.001). To examine how much energy ants can bring back to the nest through these methods, we

transformed the liquid load to sugar load. We found that the difference in efficiency between drinking

and grabbing increased with sugar concentration (Figure 5b). There was a significant interaction

between sugar concentration and foraging action (Table 3, ANOVA, sugar × foraging action: $p <$

0.01). The sugar loads of drinking were higher in the 20-40% of sugar concentrations (Wilcoxon test

with Bonferroni correction, $p < 0.05$). The total sugar load acquired by grabbing significantly

increased with sugar concentration (Wilcoxon test with Bonferroni correction, $p < 0.05$). In addition,

for the high viscosity and low sugar water 10CMC, the total sugar load per trip was lower than 10%

sugar water (Figure 5b, Wilcox test with Bonferroni correction, $p < 0.05$). Compared to the solution of

similar viscosity (40% sugar water), the sugar intake of 10CMC was less per foraging trip (Wilcoxon

test with Bonferroni correction, $p < 0.05$). If ants were optimising sugar returned to the nest per unit

time while collecting (as opposed to per trip), grabbing only would be the optimal collection

behaviour across viscosities.

To understand why ants might drink as well as grab, we investigated how often failures occur

during mandibular transportation and whether carrying a mandibular droplet causes ants to walk more

slowly. In our experimental set-up, a flat and relative short foraging path (approximately 30 cm), all

ants succeeded in transportation by mandibular grabbing (Supp. Table 4). For both mean walking

speed and maximum walking speed, there were no significant differences across the three categories:

empty-crop, crop-full, and crop- and mandible-full (Supp. Fig. 4, Tukey-Kramer test). Even when

full-crop ants held a droplet in their mandibles, their walking speed did not decrease compared to ants

in empty-crop and full-crop conditions (Supp. Fig. 4).

Discussion

Behaviour in a given species is highly adapted to that organism's context, and these

behavioural adaptations often involve precise forms of behavioural plasticity. In this study, we

analysed the flexibility of foraging behaviours in response to viscosity in *Diacamma* cf. *indicum*. The

ant used two liquid collection actions – drinking and grabbing – when collecting liquid food. In this

study, we aimed to quantify dynamic switching between of two types of liquid food collection used by

a single ant species. *Diacamma* cf. *indicum* is a part of a clade of ants that rarely specialise on liquid

food. Given this species' phylogenetic context, these behaviours are likely to be relatively recent

specialisations [15]. However, it remains unclear when ants use pseudotrophallaxis as opposed to true

trophallaxis to collect, transport, and share liquids, whether the use of these behaviours varies

according to food quality, and whether one is an evolutionary step toward another.

Viscosity dictates behaviour

Here, we clearly observed mandibular gabbing and pseudotrophallaxis in the lab in

Diacamma cf. *indicum* and saw that their use of this collection behaviour changes with sugar

concentration and viscosity. Our results are consistent with previous studies in other ponerine ants,

where ants stopped drinking [39] and tended to use mandibular grabbing at higher sugar

concentrations (> 40%) [40]. By decorrelating viscosity from sugar content, our work revealed that

ants made this switch in collection mode according to viscosity only (Figure 4). Viscosity has been

seen to reduce the liquid intake rates in many insects, including ants [23,39,41–43], consistent with

our results (Figure S2). Here, we directly measured the drinking speed of the ants. We show a robust

linear relationship between the inverse of drinking speed and the viscosity of the solutions regardless

of the presence of the viscosity-altering agent, indicating that viscosity is the main factor regulating

drinking time. Our analyses of surface tension and capillary length indicate that the biophysical limits

for maximum droplet volume of these solutions that *Diacamma* ants could transport through surface

tension are never reached. Rather, the constraints on volume obtained by grabbing are more likely

related to the ants' size and mandible span, which could be explored in future studies for example in

wasps that are physically larger and that also transport fluids between their mandibles.

We found that ants used mandibular grabbing after drinking liquid (Figure 3a). This

maximises liquid load per trip because ants can transport both internally and externally. Multiple trips

may be costly as they involve loss of energy and increased predation risk. We observed no foraging

failures or reduced travel speeds when ants were internally and externally loaded. Thus, the key factor

in evaluating foraging efficiency would be calorie intake per trip. The data showed that the total sugar

load clearly increased at higher sugar concentrations when ants used mandibular grabbing, suggesting

that grabbing, and consequently pseudotrophallaxis, are more efficient methods to collect high-

viscosity liquid than drinking and regurgitation.

One possibility is that ants drink to fulfill individual energetic needs, but transport liquid

through grabbing for the community. The sugar load by drinking 10% sugar water was greater than

the high-viscosity 10CMC solution (Figure 5). That might be caused by the fact that ants can drink

less high-viscosity fluid per unit time than low-viscosity fluid. If the volume of liquid imbibed was

solely taken to cover individual energetic foraging costs, they would drink the same *sugar content*,

which was not the case here.

Why do they not always use both drinking and mandibular grabbing?

One possibility is that mandibular grabbing is a risky, but high pay-off collection method. For

example, if ants encounter predators, they might not react quickly enough, ending up to lose their

mandibular droplet and/or being preyed upon because they are less agile. Ants might not use

mandibular grabbing in dangerous sites where they encounter predators. On the contrary, if there are

competitors around the food site, ants need to compete against other ant species. *Pheidole*

megacephala soldiers reacted to the presence of competitors. The soldiers performed more

mandibular grabbing on the territory of other ant species in order to rapidly gather and transport large

loads of liquid food [44]. Future studies with different distances and ecological contexts while

analysing transport time could help elucidate the cost of transportation of pseudotrophallaxis through

surface tension.

In addition, the speed of sharing to nestmates may differ between trophallaxis and

pseudotrophallaxis. After foragers return to the nest, they share the liquid food through trophallaxis

(regurgitation), or through pseudotrophallaxis. Trophallaxis should more rapidly distribute food in the

colony because a receiver can become a donor and continue to distribute liquid food by regurgitation,

also allowing the formation of a more complex social network [45–47]. In the carpenter ant

Camponotus, foragers give food to a receiver, proportional to the available capacity in the receiver's

crop. This trophallactic interaction helps the forager to sense colony satiation level and decide when

to leave the nest and bring in more food [48]. However, the distribution dynamics of liquid by

pseudotrophallaxis have not been studied. The observation of liquid distribution in the focal species

that use both trophallaxis and pseudotrophallaxis is needed to understand these dynamics of liquid

distribution and the regulation of foraging effort.

Why do so few ant species perform mandibular grabbing?

Mandibular grabbing and pseudotrophallaxis are mostly performed by ponerine ants

(Ponerinae), with only a few noted exceptions in other major ant subfamilies [38] One possible reason

why some ants use pseudotrophallaxis and others use true trophallaxis is that many ants have internal

morphological adaptations for liquid intake, storage and regurgitation, such as an expandable gaster,

an elastic crop, a larger mouth and throat, and a highly developed proventriculus [7,15]. These

adaptations possibly allow for greater flexibility regarding the intake of high-viscosity solutions

[16,39]. Thus, species with these morphological adaptations may not need pseudotrophallaxis.

A second possibility is that there might be biophysical restrictions on whether an ant can

collect a liquid drop between her mandibles. It is likely that small body size makes interactions with

liquid droplets more dangerous due to the difficulty of getting out of a liquid with a high surface

tension [49]. We observed that ants make a 'hasty' motion at the end of the extraction of the droplet.

The force needed to break the droplet away is also directly dictated by the surface tension of the liquid.

For small ants, it may be more difficult to exert the force required to extract the droplet. Whether any

relationship exists between the ability to perform pseudotrophallaxis and biophysical restrictions,

related to body size, has not yet been studied, making this area well poised for study from a

biomechanics perspective.

Share with nestmates: socially transferred materials

Trophallaxis allows for medium- to long-term food storage before redistribution while pseudotrophallaxis does not. Thus, ecological contexts and environmental harshness may also tilt an ant to engage in one behaviour or another. Another valuable feature of trophallaxis is that donors can alter the contents of what they pass to nestmates, either through partial digestion or through more complex signalling [50–52], which may bias a species or even a single ant to use one or another behaviour. Recent studies reveal that ants' regurgitated fluid contained more than food [50,51]. For example, trophallactic fluid in carpenter ants contains hormones, nestmate recognition cues, small RNAs, and various proteins. In *Diacamma*, it is unclear whether foragers regurgitate the contents of their crop during pseudotrophallaxis, or if they only regurgitate when they do trophallaxis. Future studies could examine whether they add any endogenous materials during these behaviours.

Data accessibility

All data and codes are available at:

https://datadryad.org/stash/share/zYykgqIg6GgOgRjhPVqO4Lwq-Xhe8etdBlHl4qHQZhg

Fudging

This work was funded by JSPS KAKENHI [Grant number JP20J01766] and Myohara Maroko

research grant by the Zoological Society of Japan to HF, and Swiss National Science Foundation

Grant PR00P3_179776 to ACL.

Acknowledgments

We are grateful to Ken Naganawa for his amazing illustrations. We would like to thank Dr. François

Lavergne and Satomi Koga for their contribution to data collection and Dr. Andrew Brown for his

advice on statistical analysis. We also thank all members of the social fluids lab at University of

Fribourg for discussions about the study and Dr. Isaac Planas-Sitjà and Marie-Pierre Meurville for

providing feedback on early drafts of the manuscript.

459 **Tables**

461 Table 1. Surface tension and capillary length for water/sugar solutions.

Sugar concentration	Surface tension (mN/m)	Std. Error surface tension (mN/m)	Density (g/mL)	Capillary length (mm)
0	71.7	0.1	0.9982	2.71
10	70.7	0.5	1.022	2.66
20	74.7	0.4	1.0631	2.68
30	80.0	0.4	1.1132	2.70
40	77.8	0.3	1.1374	2.64
50	78.8	0.3	1.1975	2.59
60	84.5	0.3	1.2391	2.64

Table 2. Influence of sugar concentration and foraging action on volume and handling time. Results

of generalised and general linear regression (a) and linear regression model (b) between two

explanatory variables and volume of carried or drunk water and handling time. Sugar concentration

water is 10, 20, 30, 40, 50, and 60% (w/w) and foraging actions are drinking (DR) and mandibular

grabbing (GR). Viscosity is seven different sugar and viscosity solutions; 10, 20, 30, 40, 50, 60%

(w/w), and the viscosity-altered solution 10CMC.

(a) generalised linear regression

	Explanatory variables	Estimate	Std. Error	p-value
Volume	Sugar concentration	-0.026	0.003	< 0.001
	Foraging actions (DR or GR)	-1.05	0.21	< 0.001
	Sugar * foraging actions	0.028	0.0054	< 0.001
Handling time	Viscosity	-1325.942	193.878	< 0.001
	Foraging actions (DR or GR)	-62.635	6.594	< 0.001
	Viscosity * foraging actions	1386.844	310.709	< 0.001

472 (b) linear regression

		Explanatory variables	Estimate	Std. Error	t-value	p-value
Volume	Drinking	(Intercept)	1.81	0.13	3.6	0.001
		Sugar concentration	-0.026	0.0034	-7.57	0.001
	Grabbing	(Intercept)	0.76	0.11	7.19	< 0.001
		Sugar concentration	0.002	0.0027	0.75	0.41
Handling	Drinking	(Intercept)	75.4933	4.4055	17.136	< 0.001

	time					
		Viscosity	-1.3259	0.2322	-5.709	< 0.001
	Grabbing	(Intercept)	9.609	3.263	2.944	< 0.01
		Viscosity	2.069	1.403	1.475	0.14

476 Table 3. Influence of different concentrations and viscosity of sugar water and foraging action on

477 liquid and sugar load. Results of two-way ANOVA between two explanatory variables and liquid load

478 or sugar load per trip. Solutions are seven different sugar solutions 10, 20, 30, 40, 50, and 60% (w/w)

479 and 10CMC. Collection methods are drinking (DR) and both drinking and mandibular grabbing

480 (both).

	Explanatory variables	Df	Sum Sq	Mean Sq	F value	p-value
Liquid load	Solutions	6	8.66	1.44	60.12	< 0.001
	Collection methods (DR or both)	1	92.13	92.13	3838.72	< 0.001
	Solution * collection methods	6	1.55	0.26	10.74	< 0.001
	Residuals	515	12.36	0.02		
Sugar load	Solutions	6	13.758	2.293	1154.6	< 0.001
	Collection methods (DR or both)	1	14.902	14.902	7503.6	< 0.001
	Solution * collection methods	6	6.841	1.140	574.1	< 0.001
	Residuals	515	1.023	0.002		

Figures

Figure 1. Ethogram of the social bucket technique.

An ant touches a solution with the antennae and mouthparts (tasting). After tasting, an

individual opens mandibles to grab and pull at the liquid (a). The ant occasionally succeeds in

collecting a droplet of solution between the mandibles (b). The individual returns to the nest

(c). Inside the nest, the ant shares the drop with other nestmates (d). When the receiver begs

for the droplet, their antennae move quickly. Several nestmates can drink from the droplet at

the same time. Illustrations by Ken Naganawa.

Figure 2. Dynamic viscosity leads to drinking speed.

(a) Dynamic viscosities at 25°C for different sugar water solutions as a function of the sugar

concentration (w/w). The different colours correspond to pure water (circle) and viscosity-

altered solutions (triangle). The error bars denote the quality of the Newtonian fit applied to

each flow curve for each solution. The blue triangle indicates the viscosity-altered solutions

10CMC and 30CMC sugar solution with 0.25 CMC(w/w). (b) The relationship between

drinking speed and viscosity. The inverse of drinking speed increases linearly with dynamic

viscosity for sugar solutions with or without viscosity-altering additives.

Figure 3. Impact of sugar concentration and viscosity on liquid collection.

 (a) The amount of liquid collected by drinking (black) and grabbing (grey) at different concentrations of sugar water (% w/w). Time spent drinking (b) and grabbing (c) by ants at different concentrations and viscosities of sugar water (% w/w). Six sugar concentration solutions are shown in black circles, and the viscosity-altered solution 10CMC is shown with blue triangles. Statistical analysis can be found in Table 2. Linear regression shown in (b) is performed only on the unaltered solutions…

Figure 4. The proportion of pseudotrophallaxis depends on sugar concentration and viscosity.

Drinking (white), both (grey), and grabbing (dark grey) indicate the behaviour of only

pseudotrophallaxis, pseudotrophallaxis after drinking, and only drinking. The blue-ish bars indicate

the viscosity-altered solution 10CMC. Letters on the top of the bar mean they were significantly

different at $P < 0.05$ (chi-sq test with Bonferroni correction).

Figure 5. Foraging efficiency for the foraging action.

Estimated total liquid (a) and sugar load per trip (b) are estimated based on drinking time and foraging

action. White (circle) and dark grey (triangle) boxes indicate the action of drinking and both drinking

and grabbing. Blue indicates the 10CMC viscosity-altered solution. Statistical analysis can be found

in Table 3. Letters above points indicate significant differences of $p < 0.05$ (Wilcoxon test with

Bonferroni correction).

References

1.   Pyke GH, Pulliam HR, Charnov EL. 2015 Optimal Foraging: A Selective Review of Theory

and Tests. *https://doi.org/10.1086/409852* **52**, 137–154. (doi:10.1086/409852)

2.   Stephens DW, Krebs JR. 1987 *Foraging Theory*. Princeton University Press.

(doi:10.1515/9780691206790)

3.   Buckley RC. 1987 Interactions involving plants, Homoptera, and ants. *Annual review of*

ecology and systematics. Vol. 18 , 111–135. (doi:10.1146/annurev.es.18.110187.000551)

4.   Davidson DW, Cook SC, Snelling RR, Chua TH. 2003 Explaining the abundance of ants in

lowland tropical rainforest canopies. *Science (1979)* **300**, 969–972.

(doi:10.1126/SCIENCE.1082074/SUPPL_FILE/DAVIDSON.D.SOM.PDF)

5.   Nelsen MP, Ree RH, Moreau CS. 2018 Ant–plant interactions evolved through increasing

interdependence. *Proceedings of the National Academy of Sciences* **115**, 12253–12258.

(doi:10.1073/pnas.1719794115)

6.   Heil M, Mckey D. 2003 Protective Ant-Plant Interactions as Model Systems in Ecological and

Evolutionary Research. *Source: Annual Review of Ecology, Evolution, and Systematics* **34**,

425–453. (doi:10.1146/132410)

7.   Hölldobler B, Wilson EO. 1990 *The Ants*. Cambridge: Harvard University Press.

8.   Wilson EO. 1990 *The insect societies (Harvard paperbacks)*.

9.   Calixto ES, Lange D, Del-Claro K. 2021 Net benefits of a mutualism: Influence of the quality

of extrafloral nectar on the colony fitness of a mutualistic ant. *Biotropica* **53**, 846–856.

(doi:10.1111/BTP.12925)

10. Pekas A, Tena A, Aguilar A, Garcia-Marí F. 2011 Spatio-temporal patterns and interactions

with honeydew-producing Hemiptera of ants in a Mediterranean citrus orchard. *Agric For*

Entomol **13**, 89–97. (doi:10.1111/j.1461-9563.2010.00501.x)

11. Pringle EG. 2021 Ant-Hemiptera Associations. *Encyclopedia of Social Insects* , 45–48.

(doi:10.1007/978-3-030-28102-1_8)

12. Styrsky JD, Eubanks MD. 2006 Ecological consequences of interactions between ants and

honeydew-producing insects. *Proceedings of the Royal Society B: Biological Sciences* **274**,

151–164. (doi:10.1098/RSPB.2006.3701)

13.   Sempo G, Detrain C. 2004 Between-species differences of behavioural repertoire of castes in

554   the ant genus Pheidole: A methodological artefact? *Insectes Soc* (doi:10.1007/s00040-003-

555   0704-2)

556   14.   Richard FJ, Dejean A, Lachaud JP. 2004 Sugary food robbing in ants: a case of temporal

557   cleptobiosis. *C R Biol* **327**, 509–517. (doi:10.1016/J.CRVI.2004.03.002)

558   15.   Meurville MP, LeBoeuf AC. 2021 Trophallaxis: the functions and evolution of social fluid

559   exchange in ant colonies (Hymenoptera: Formicidae). *Myrmecol News* **31**, 1–30.

560   (doi:10.25849/MYRMECOL.NEWS_031:001)

561   16.   Davidson DW, Cook SC, Snelling RR. 2004 Liquid-feeding performances of ants

562   (Formicidae): Ecological and evolutionary implications. *Oecologia* **139**, 255–266.

563   (doi:10.1007/S00442-004-1508-4/TABLES/4)

564   17.   Eisner T. 1957 A comparative morphological study of the proventriculus of ants

565   (Hymenoptera: Formicidae). *Bull Mus Comp Zool* **116**, 437–490.

566   18.   Eisner T, Brown WLJ. 1958 The evolution and social significance of the ant proventriculus.

567   *Proceedings Tenth International Congress of Entomology* **2**, 503–508.

568   19.   Robinson EJH, Feinerman O, Franks NR. 2009 Flexible task allocation and the organization of

569   work in ants. *Proceedings of the Royal Society B: Biological Sciences* **276**, 4373–4380.

570   (doi:10.1098/rspb.2009.1244)

571   20.   Hölldobler B. 1985 Liquid food transmission and antennation signals in ponerine ants. *Isr J*

572   *Entomol* , 89–99.

573   21.   A. Dejean, J.P. Suzzoni. 1997 Surface Tension Strengths in the Service of a Ponerine Ant: a

574   New Kind of Nectar Transport. *Naturwissenschaften* **84**, 76–79.

575   22.   Lachaud Jean-Paul, Alain Dejean. 1991 Food sharing in Odontomachus troglodytes (Santschi):

576   a behavioral intermediate stage in the evolution of social food exchange in ants. *An Biol* **17**,

577   53–61.

578   23.   Lois-Milevicich J, Schilman PE, Josens R. 2021 Viscosity as a key factor in decision making

579   of nectar feeding ants. *J Insect Physiol* **128**, 104164. (doi:10.1016/j.jinsphys.2020.104164)

580   24.   Bonser R, Wright PJ, Bament S, Chukwu UO. 1998 Optimal patch use by foraging workers of

581   Lasius fuliginosus, L. niger and Myrmica ruginodis. *Ecol Entomol* **23**, 15–21.

582   (doi:10.1046/J.1365-2311.1998.00103.X)

25.   Dejean A, Solano PJ, Ayroles J, Corbara B, Orivel J. 2005 Arboreal ants build traps to capture
prey. *Nature* **434**, 973–973. (doi:10.1038/434973a)

26.   Detrain C, Prieur J. 2014 Sensitivity and feeding efficiency of the black garden ant Lasius
niger to sugar resources. *J Insect Physiol* **64**, 74–80. (doi:10.1016/J.JINSPHYS.2014.03.010)

27.   Falibene A, de Figueiredo Gontijo A, Josens R. 2009 Sucking pump activity in feeding
behaviour regulation in carpenter ants. *J Insect Physiol* **55**, 518–524.
(doi:10.1016/J.JINSPHYS.2009.01.015)

28.   Fujioka H, Okada Y. 2019 Liquid exchange via stomodeal trophallaxis in the ponerine ant
Diacamma sp. from Japan. *J Ethol* **37**, 371–375. (doi:10.1007/s10164-019-00602-9)

29.   Blüthgen N, Menzel F, Blüthgen N. 2006 Measuring specialization in species interaction
networks. *BMC Ecol* **6**, 9. (doi:https://doi.org/10.1186/1472-6785-6-9)

30.   VanGinkel CG, Gayton S. 1996 The biodegradability and nontoxicity of carboxymethyl
cellulose (DS 0.7) and intermediates. *Environ Toxicol Chem* **15**, 270–274.
(doi:10.1002/ETC.5620150307)

31.   Osswald T, Rudolph N. 2014 Polymer Rheology. In *Polymer Rheology*, pp. I–XI. München:
Carl Hanser Verlag GmbH & Co. KG. (doi:10.3139/9781569905234.fm)

32.   Rhee BO, Lee S. 1999 Evaluation on Accuracy of the Rheological Data of PIM Feedstocks.
Journal of the Japan Society of Powder and Powder Metallurgy **46**, 830–836.
(doi:10.2497/jjspm.46.830)

33.   Emmerich A. 1994 Density data for sucrose solutions. Contribution to the application of the
new values adopted by ICUMSA. *Zuckerindustrie (Germany)* **119**, 120–123.

34.   de Gennes P-G, Brochard-Wyart F, Quéré D. 2004 *Capillarity and Wetting Phenomena*. New
York, NY: Springer New York. (doi:10.1007/978-0-387-21656-0)

35.   Josens RB, Farina WM, Roces F. 1998 Nectar feeding by the ant Camponotus mus: intake rate
and crop filling as a function of sucrose concentration. *J Insect Physiol* **44**, 579–585.
(doi:10.1016/S0022-1910(98)00053-5)

36.   Yamanaka O, Takeuchi R. 2018 UMATracker: An intuitive image-based tracking platform.
Journal of Experimental Biology **221**. (doi:10.1242/JEB.182469/VIDEO-3)

37.   Virtanen P *et al.* 2020 SciPy 1.0: fundamental algorithms for scientific computing in Python.
Nat Methods **17**, 261–272. (doi:10.1038/s41592-019-0686-2)

38.   Jean Leonard Poiseuille. 1844 *Recherches experimentales sur le mouvement des liquides dans*

614   *les tubes de tres-petits diametres*. Imprimerie Royale. See

615   https://books.google.co.jp/books?id=uBN1Q-IRzTMC}.

616   39.   Ávila Núñez JL, Naya M, Calcagno-Pissarelli MP, Otero LD. 2011 Behaviour of

617   Odontomachus chelifer (Latreille) (Formicidae: Ponerinae) Feeding on Sugary Liquids. *J*

618   *Insect Behav* **24**, 220–229. (doi:10.1007/S10905-010-9249-1/FIGURES/4)

619   40.   Jandt J, Larson HK, Tellez P, McGlynn TP. 2013 To drink or grasp? How bullet ants

620   (Paraponera clavata) differentiate between sugars and proteins in liquids. *Naturwissenschaften*

621   **100**, 1109–1114. (doi:10.1007/S00114-013-1109-3/FIGURES/2)

622   41.   Borrell BJ. 2006 Mechanics of nectar feeding in the orchid bee Euglossa imperialis: pressure,

623   viscosity and flow. *Journal of Experimental Biology* **209**, 4901–4907.

624   (doi:10.1242/JEB.02593)

625   42.   Kim W, Gilet T, Bush JWM. 2011 Optimal concentrations in nectar feeding. *Proc Natl Acad*

626   *Sci U S A* **108**, 16618–16621. (doi:10.1073/PNAS.1108642108)

627   43.   Nicolson SW, de Veer L, Köhler A, Pirk CWW. 2013 Honeybees prefer warmer nectar and

628   less viscous nectar, regardless of sugar concentration. *Proceedings of the Royal Society B:*

629   *Biological Sciences* **280**. (doi:10.1098/RSPB.2013.1597)

630   44.   Dejean A, le Breton J, Suzzoni JP, Orivel J, Saux-Moreau C. 2005 Influence of interspecific

631   competition on the recruitment behavior and liquid food transport in the tramp ant species

632   Pheidole megacephala. *Naturwissenschaften* **92**, 324–327. (doi:10.1007/S00114-005-0632-

633   2/FIGURES/2)

634   45.   Buffin A, Mailleux AC, Detrain C, Deneubourg JL. 2011 Trophallaxis in Lasius niger: A

635   variable frequency and constant duration for three food types. *Insectes Soc* **58**, 177–183.

636   (doi:10.1007/S00040-010-0133-Y/TABLES/2)

637   46.   Greenwald E, Segre E, Feinerman O. 2015 Ant trophallactic networks: simultaneous

638   measurement of interaction patterns and food dissemination. *Sci Rep* **5**, 12496.

639   (doi:10.1038/srep12496)

640   47.   Bles O, Deneubourg J-L, Nicolis SC. 2018 Food dissemination in ants: Robustness of the

641   trophallactic network against resource quality. *Journal of Experimental Biology* **221**.

642   (doi:10.1242/jeb.192492)

643   48.   Greenwald EE, Baltiansky L, Feinerman O. 2018 Individual crop loads provide local control

644   for collective food intake in ant colonies. *Elife* **7**, e31730. (doi:10.7554/eLife.31730)

645   49.   Padday JF, Pitt AR, Pashley RM. 1975 Menisci at a free liquid surface: surface tension from

646   the maximum pull on a rod. *Journal of the Chemical Society, Faraday Transactions 1:*

647   *Physical Chemistry in Condensed Phases* **71**, 1919. (doi:10.1039/f19757101919)

648   50.   Leboeuf AC *et al.* 2016 Oral transfer of chemical cues, growth proteins and hormones in social

649   insects. *Elife* **5**. (doi:10.7554/eLife.20375)

650   51.   Hakala SM, Meurville MP, Stumpe M, Leboeuf AC. 2021 Biomarkers in a socially exchanged

651   fluid reflect colony maturity, behavior, and distributed metabolism. *Elife* **10**.

652   (doi:10.7554/ELIFE.74005)

653   52.   Hakala SM *et al.* 2022 Socially transferred materials: why and how to study them. *Trends Ecol*

654   *Evol* (doi:10.1016/j.tree.2022.11.010)

655