Molecular landscape and functional characterization of centrosome amplification in ovarian cancer

Carolin M. Sauer[1,2,$], James A. Hall[1,2], Dominique-Laurent Couturier[1,3], Thomas Bradley[1,2], Anna M. Piskorz[1,2], Jacob Griffiths[1,2], Ashley Sawle[1,2], Matthew D. Eldridge[1,2], Philip Smith[1,2], Karen Hosking[2], Marika A.V. Reinius[1,2,4], Lena Morrill Gavarró[1,2], Anne-Marie Mes-Masson[5], Darren Ennis[6], David Millan[7], Aoisha Hoyle[8], Iain A. McNeish[6], Mercedes Jimenez-Linan[4], Filipe Correia Martins[1,4], Julia Tischer[1,2], Maria Vias[1,2], James D. Brenton[1,2,4,$]

1. Cancer Research UK Cambridge Institute, University of Cambridge, Li Ka Shing Centre, Robinson Way, Cambridge CB2 0RE, UK.

2. Cancer Research UK Major Centre–Cambridge, University of Cambridge, Cambridge CB2 0RE, UK.

3. Medical Research Council Biostatistics Unit, University of Cambridge, Cambridge CB2 0SR, UK.

4. Cambridge University Hospital NHS Foundation Trust and National Institute for Health Research Cambridge Biomedical Research Centre, Addenbrooke's Hospital, Cambridge, UK.

5. Centre de Recherche du Centre Hospitalier de l'Université de Montréal (CRCHUM) and Institut du Cancer de Montréal (ICM), CRCHUM, 900 Saint-Denis, Montreal, QC, Canada.

6. Department of Surgery and Cancer, Ovarian Cancer Action Research Centre, Imperial College London, London, UK.

7. Department of Pathology, Queen Elizabeth University Hospital, Glasgow, UK.

8. Department of Pathology, University Hospital Monklands. NHS Lanarkshire, UK.

[$]Correspondence: carolin.sauer@cruk.cam.ac.uk; james.brenton@cruk.cam.ac.uk

Abstract

High grade serous ovarian carcinoma (HGSOC) is characterised by poor outcome and extreme chromosome instability (CIN). Therapies targeting centrosome amplification (CA), a key mediator of chromosome missegregation and CIN, may have significant clinical utility in HGSOC. However, the prevalence of CA in HGSOC, its relationship to genomic biomarkers of CIN and its potential impact on therapeutic response have not been defined. Using high-throughput multi-regional microscopy on 287 clinical HGSOC tumour tissues and 73 ovarian cancer cell lines, we show that CA through centriole overduplication is a highly recurrent and heterogeneous feature of HGSOC and is strongly associated with CIN and genome subclonality. Cell-based studies showed that high prevalence CA is phenocopied in ovarian cancer cell lines, and that high CA is associated with increased multi-treatment resistance; most notably to paclitaxel which is the most common treatment used in HGSOC. CA in HGSOC may therefore present a potential driver of tumour evolution and a powerful biomarker for response to standard-of-care treatment.

Keywords

Centrosomes; Chromosomal instability; high grade serous ovarian cancer; micronuclei; copy number aberrations; cell line characterisation

Introduction

High grade serous ovarian carcinoma (HGSOC) accounts for the majority of ovarian cancer cases and ovarian cancer related deaths with a five-year survival rate of less than 30%. Overall survival of HGSOC patients has not changed over the last two decades mainly due to the severe ongoing chromosomal instability (CIN) characterising and driving this disease. CIN is present in virtually all HGSOCs[1], drives resistance to anti-cancer therapies[2–6], and is the main cause of regenerating subclonal diversity in response to treatment-induced selective pressures[7]. The lack of a clear understanding of the drivers of this genomic complexity in HGSOC has significantly impeded the development of precision therapies, with the major exception being the successful targeting of BRCA1/2 dysfunction with PARP inhibitors. Patterns of copy number aberrations in HGSOC can identify specific mutational processes causing CIN and highlight potential therapeutic vulnerabilities[8,9], but the relationship between these signatures and other cellular causes of CIN, specifically centrosome amplification has not been investigated.

The single most prominent cause of CIN is chromosome missegregation, which can be caused by weakened mitotic checkpoint, improper chromosome attachment to the mitotic spindle and centrosome amplification (CA; typically defined as an abnormal number of centrosomes >1 in non-dividing cells)[10–12]. The centrosome, also known as the microtubule organising centre (MTOC), consists of two centrioles embedded within the pericentriolar material (PCM; a proteinaceous scaffold with microtubule-nucleating activities[13,14]), and guides spindle formation and accurate chromosome segregation during cell division[15]. Consequently, centrosome abnormalities, both structural and numerical, can lead to the missegregation of chromosomes resulting in aneuploidy and CIN [10,16,17]. Centrosome abnormalities, such as CA, may provide a novel therapeutic target in cancers, and indeed, several drugs targeting centrosome duplication[18] or associated survival mechanisms[19,20] are now in development or clinical trials[21]. However detailed studies of the prevalence, mechanisms and origins of CA in HGSOC and other cancers are now required for the clinical exploitation of this phenotype.

Most studies that have investigated mechanisms and consequences of CA have experimentally induced or inhibited CA in cell lines or other model systems predominantly in breast cancer models. However, the high degree of variability in centrosome numbers observed across cancer cell populations indicates the existence of a CA "set point" or equilibrium[22] and suggests that cell lines have different tolerance thresholds for CA and maintain centrosome numbers through an equilibrium of CA mechanisms and negative selection[23,24]. Consequently, the experimental induction or inhibition of CA may trigger a range of different cellular responses that would not otherwise be observed without these artificial perturbations. In addition, because of significant challenges associated with detecting centrosomes in clinical human tissue samples, very few studies have investigated CA in clinical tumour specimens[25,26], and the prevalence of CA in HGSOC (both in cell lines and tumour tissues) remains largely unclear.

To address these shortcomings, we here describe a systematic and detailed analysis of baseline CA along with its molecular and genomic associations in large cohorts of HGSOC tissues and ovarian cancer cell lines. We developed a high-throughput microscopy-based assay to reliably detect and quantify centrosomes in formalin-fixed paraffin-embedded (FFPE) tissues and analysed

centrosome profiles in >300 tissue samples containing 287 HGSOC tumours. These methods
uncovered a high prevalence of CA in HGSOC with marked intra-tumoural tissue heterogeneity. To
further probe the biological nature of CA in ovarian cancer, we extended our approaches to the in-depth
profiling of 73 ovarian cancer cell line models at single-cell resolution, and provide the first
comprehensive phenotypic, transcriptomic, and genomic characterisation of supernumerary
centrosomes. The results of this work confirm the high prevalence of CA arising from centriole
overduplication in ovarian cancer, directly link CA to CIN and genomic subclonality, and show that CA
dictates treatment response to anti-mitotic agents, most importantly paclitaxel, with clear clinical
implications. Finally, our work paves the way for future studies characterising centrosome abnormalities
in large cohorts of clinical tumour specimens and provides an important resource for future research
investigating CA and associated vulnerabilities for the treatment of HGSOCs.

Results

Supernumerary centrosomes are present in the majority of HGSOC tissues and show marked inter- and intra-tumour heterogeneity

To investigate the presence and extent of supernumerary centrosomes in HGSOC, we selected full-face FFPE tumour sections (n=93) from the prospective non-interventional cohort study (OV04) and tissue microarrays (TMAs; n=194) from the British Translational Research Ovarian Cancer Collaborative (BriTROC) study (total of n=287 tumour tissues; see Methods and **Supplementary Table 1**). We included matched and unmatched normal fallopian tube (FT) and other normal tissues as negative controls. Liver samples were included as a positive control for CA, since the liver is the only organ that has supernumerary centrosomes in normal cells[27–29]. To obtain accurate estimates of centrosome numbers in clinical HGSOC tissue samples, we developed an automated, high-throughput immunofluorescent microscopy-based imaging approach using the confocal Operetta CLS™ imaging system (**Fig. 1 and Supplementary Methods**). In total we profiled 3,632,389 nuclei across 11,811 imaging fields using up to 50 non-overlapping randomly placed imaging fields per tissue sample in 25 µm FFPE tissue sections. To allow comparisons across both cohorts and to account for potential batch effects, CA scores were normalised to the median CA score of normal control tissues within each cohort.

In both study cohorts, CA scores were significantly higher in tumour tissues than normal control tissues (FT, spleen and kidney; Kruskal-Wallis, $p = 1.9 \times 10^{-10}$). As expected, control liver tissues showed the highest CA score (**Fig. 1d**). Using a CA threshold based on the 95% confidence interval of the FT sample with the highest CA score (1.83; **Fig. 2**), 73% of OV04 and 59% of BriTROC tumour samples displayed significant CA (**Fig. 1d**). Only one tumour sample showed significantly lower CA scores than normal tissues (**Fig. 2**). Centrosomes in tumour tissues also had significantly larger PCM areas (measured as width × length determined from 2D maximum intensity projection images) than centrosomes detected in normal control tissues (Kruskal-Wallis, $p = 2.6 \times 10^{-8}$; **Fig. 1e**), indicating that structural centrosome abnormalities might also be a common feature in HGSOCs.

As expected, we observed marked inter-tumoural heterogeneity for CA, but also substantial intra-tumoural differences within cases. Notably, there was marked variability in the distribution of CA scores across imaging fields, even between cases with similar median CA scores (**Fig. 2**). To investigate this further, we developed a hierarchical statistical model able to account for intra-tissue dependence in mean and variance CA scores by means of random effects and allowing for heavy-tailed error distributions (see Methods for details). Model estimates confirmed the significant differences in average CA levels between FT and tumour tissues ($p \ll 0.001$), as well as the presence of marked intra-tumoural heterogeneity ($p \ll 0.001$). To quantify the observed heterogeneity in CA phenotypes, we estimated the standard deviation of log-transformed CA scores across all imaging fields for each tissue sample. Examples of CA high and low tissues with varying tissue heterogeneity scores are shown as spatial heatmaps in **Fig. 3a–b** and **Supplementary Fig. 1**. Heterogeneity scores were significantly higher in OV04 samples (full face sections) than BriTROC samples (TMA cores; Wilcoxon, $p = 6.6 \times 10^{-11}$) consistent with a much larger area sampled by random imaging fields across full face sections compared to adjacent imaging fields covering 1 mm TMA cores. We also observed significant differences in CA levels between tumour tissues collected from the same patient (n = 36; **Fig. 3c**). Using

the CA cutoff of 1.83, for the majority of patients (64%), analysed tissues fell into the same CA category (i.e. either significant CA or no CA), whereas for 36% of cases, tissues within either category were found. Importantly, no differences in CA scores were observed between different anatomical tissue sites (Kruskal-Wallis, p = 0.62; **Fig. 3d**), while tissue samples collected from the omentum showed higher CA tissue heterogeneity (Kruskal-Wallis, p = 0.0045; **Fig. 3e**). However, this variability could result from differences in how blocks were sampled from pelvic and omental tissues between the two study cohorts.

Using robust statistical modelling, these analyses confirm very high prevalence of CA in HGSOC cases and significant intra-tumoural and inter-tumoural heterogeneity of CA phenotypes.

Centrosome amplification is not a prognostic marker in HGSOC

CA has been frequently associated with increased disease aggressiveness and poor patient prognosis in various cancer types[25,30]. We therefore investigated whether CA was associated with clinical or molecular features and overall survival in ovarian cancer. For patients with multiple tissue samples, the median CA score across these tissues was used. The BriTROC cohort, contained four patients with endometrioid ovarian cancer, which has improved prognosis compared to HGSOC patients. These patients showed significantly lower CA compared to the remaining HGSOC patients from both cohorts (Wilcoxon, p = 0.02; **Fig. 4a**). No significant differences in CA scores were observed across other important clinical variables including tumour stage (Kruskal-Wallis, p = 0.19), *BRCA1* and *BRCA2* germline mutation status (Kruskal-Wallis, p = 0.36), or between treatment-naïve (immediate primary surgery [IPS]) and post-chemotherapy initiation specimens (delayed primary surgery [DPS]; Wilcoxon, P = 0.26; **Fig. 4b–d**). We did, however, find an increase in CA tissue heterogeneity scores in DPS compared to IPS samples (Wilcoxon, P = 0.015**; Fig. 4e**) as well as an increase in centrosome size (Wilcoxon, p = 0.00036; data not shown), suggesting that chemotherapy might influence tissue-wide CA characteristics.

The BriTROC study enrolled women who had relapsed following front-line therapy and who were well enough to undergo surgery or an image-guided biopsy[31]. As a result, the BriTROC cohort is enriched for better-outcome patients with a median survival of 4.9 years as compared median survival of 2.4 years in the OV04 cohort (**Supplementary Table 1**). We therefore performed survival analyses independently on both cohorts. Multivariable Cox proportional hazards showed that the presence of CA was not associated with overall survival in either of the two cohorts (OV04: p = 0.461, HR = 0.8, 95% CI 0.43–1.5; BriTROC: p = 0.09, HR = 1.59, 95% CI 0.93–2.7; **Fig. 4f–g**). This result was surprising given previous publications reporting CA as a poor prognostic variable in other epithelial cancers[25,30]. We further analysed the relationship between CA and survival using data from the Cancer Genome Atlas (TCGA; n = 9,721), using the CA20 gene expression signature as a proxy marker for CA[30,32]. This confirmed CA was highly prevalent in ovarian cancer which was amongst the highest CA20 expressing cancer type[30] (**Supplementary Fig. 2a**). CA20 scores were strongly correlated with the CIN gene expression signature, CIN25[33] (Spearman's R = 0.96, p = 2.2 × 10^{-16}; **Supplementary Fig. 2b**). This is consistent with previous studies reporting that experimentally induced CA can give rise to aneuploidy and CIN[10,34]. Importantly, we noticed that cancer types in which CA had previously been implicated as a poor prognostic variable[25,30] (highlighted by blue arrowheads in **Supplementary Fig. 2a**) generally

had lower than the pan-cancer average for CA20 and CIN25 signature expression (see **Supplementary Fig. 2a**). We hypothesised that the survival differences observed for these cancers is caused by CA identifying cases with higher CIN. We further reasoned that in cancer subtypes that are primarily driven by severe CIN, such as HGSOC[35], there is less CA effect on universally poor disease outcome. We therefore separated the TCGA cancer subtypes into CIN-high and CIN-low cancers (using a median split). Multivariable Cox proportional hazard analysis on the CIN-low group confirmed previous findings that high CA20 was associated with poor survival (P < 0.001, HR = 1.5, 95% CI 1.2–1.9; **Supplementary Fig. 2c**). However, this association was not observed in CIN-high cancers (P < 0.971, HR = 1.0, 95% CI 0.89–1.1; **Supplementary Fig. 2d**). In addition, high CA20 was also not associated with disease outcome in the TCGA HGSOC group (n = 304, patients divided by CA20 median; log-rank test, p = 0.63), confirming our observations in the BriTROC and OV04 cohorts.

In-depth characterisation of ovarian cancer cell lines shows frequent centrosome amplification through centriole overduplication

Our data confirms that CA is a highly prevalent and heterogenous feature in HGSOC tumours, but several unanswered questions remain. PCM proteins are highly dynamic and can form assemblies without centrioles[36]. Our tissue-based CA assay was limited to PCM proteins (PCNT and CDK5RAP2) owing to significant limitations of other antibodies in FFPE processed samples. It is therefore unknown whether supernumerary centrosomes observed in HGSOC specimens are functional and intact, or might alternatively be acentriolar PCM foci caused by PCM fragmentation. To investigate this possibility, we performed in-depth characterisation in a large panel of ovarian cancer cell lines by developing a semi-automated imaging approach, as summarised in **Fig. 5a**. Antibody panels were designed to allow quantification of centrosomes (PCNT, CETN3, and CEP164) and micronuclei (Hoechst) in single cells (cytokeratin stains were used to estimate cell boundaries), as well as the estimation of mitotic indices (pHH3), and the detection of DNA damage (γH2AX). Combined, we analysed a total of 388,748 cells from 73 screened ovarian cancer cell lines (**Fig. 5b**). These analyses confirmed that ovarian cancer cell lines recapitulate the high prevalence of CA observed in HGSOC tissue samples with a mean CA frequency of 26.2% (range 0.5–50.4%; non-mitotic cells with two or more centrosomes; **Fig. 6a, b**).

To examine whether supernumerary centrosomes were intact centrosomes or aggregates of PCM, we analysed co-staining of PCNT with the centriole markers Centrin 3 (CETN3) and CEP164. CEP164 is a distal appendage marker which can only be found on the mother centriole of mature centrosomes[37]. Accordingly, acentriolar PCM assemblies would be expected to be negative for both CEP164 and CETN3 markers. Extra centrosomes caused by the incorrect or premature segregation of centrioles would be expected to result in CETN3-positive PCM foci with approximately 50% showing positive CEP164 staining. Lastly, CA arising from true centriole overduplication would be expected to result in fully functional centrosomes containing two centrioles, a daughter and a CEP164-positive mother centriole, surrounded by PCM. We found that both markers, CETN3 and CEP164, were present in the large majority (>95%) of detected PCM foci (**Fig. 5b**). This confirms that CA in ovarian cancer cells is predominantly caused by centriole overduplication resulting in intact and therefore likely fully functional centrosomes.

Given the high degree of CIN observed in HGSOC in comparison to other ovarian cancer subtypes, we next asked whether HGSOC cell lines showed higher levels of CA, and whether CA correlated with other molecular features of interest, including DNA content and the presence of micronuclei. We detected micronuclei in the majority of analysed cell lines with micronuclei frequencies ranging from 2.7–48.9% (mean 15.6%; defined as single non-mitotic cells which contain at least one or more micronuclei). High level CA was associated with increased MN frequencies (Spearman's R = 0.42, p ≪ 0.001), providing further evidence that CA in non-perturbated ovarian cancer cell lines may induce MN formation through chromosome missegregation. In addition, we found moderate to strong correlations between CA frequencies and nuclei intensities (surrogate marker for DNA content; Spearman's R = 0.32, p = 0.006) as well as increased mitotic indices (Spearman's R = 0.38, p < 0.001). No correlation was observed between CA frequencies and DNA damage (measured as percentage of cells with γH2AX-positive nuclei Spearman's R = 0.14, p = 0.3; **Fig. 6c**). Importantly, we did also not observe significant differences in CA frequencies across different histological subtypes (Wilcoxon, p = 0.17; **Fig. 6d**) including detection of CA in the non-cancerous fallopian and ovarian surface epithelial cell lines, FT194, FT246 and IOSE4. These observations suggest that the transformation and/or immortalisation of normal cells may induce oncogenic features that include CA, and that these lines may not represent suitable normal controls in CA/CIN studies. In contrast, MN frequencies were significantly higher in HGSOC cell lines compared to all other histological cell line subtypes (Wilcoxon, p ≪ 0.0001; **Fig. 6e**). Nevertheless, our large-scale characterisation assay highlights CA through centriole overduplication as a highly prevalent feature and to be strongly correlated with the presence of MN in ovarian cancer cell lines.

Reduction of oxygen levels induces CA in HGSOC cell lines

Hypoxia can promote CA via HIF1α-dependent induction of PLK4[38]. Out of the 73 cell lines screened, 25 were derived and maintained in low oxygen growth conditions (5% O_2; termed here as low-O_2 cell lines). Low-O_2 lines had significantly higher CA and MN frequencies to those grown in normoxia (21% O_2; Wilcoxon, p = 0.0004 and p ≪ 0.0001, respectively; **Fig. 6f, g**) but also contained only HGSOC lines, which showed higher MN frequencies than lines from other ovarian cancer subtypes. To investigate whether increased CA and MN frequencies were caused by lower oxygen growth conditions, 8 HGSOC cell lines that are normally passaged in normoxia were transferred to 5% oxygen for 48 hours. Transient low oxygen growth conditions significantly increased CA (Wilcoxon, p = 2×10^{-6}) but had no effect on MN frequencies (Wilcoxon, p = 1; **Fig. 6h-i**). Longer exposure to 5% oxygen conditions might be required to observe MN formation but these findings suggest, that at least in this experimental setting CA precedes MN formation, or MN formation is independent of CA.

Cell lines with high centrosome amplification show upregulated survival signalling pathways

Having confirmed that HGSOC cell lines strongly phenocopied CA observed in patient samples, we next used RNA sequencing to investigate further potential drivers and consequences of CA. To understand possible confounding factors that might influence gene expression independent of CA and MN status, we first performed differential gene expression (DGE) analyses comparing different

histological subtypes and cell culture growth conditions (see **Supplementary Fig. 3**). We did not

observe significant upregulation of hypoxia hallmark/response genes in low-O_2 cell lines compared to

cell lines cultured at 21% oxygen, and no additive effect was detected between CA and oxygen growth

conditions when performing DGE model checks. Importantly, this suggests that it is the reduction of

oxygen levels (compared to baseline) that induces CA (see **Fig. 6h–i**), and that the higher frequencies

of CA and MN observed in low-O_2 cell lines are cell line intrinsic rather than caused by differing oxygen

growth conditions. Based on these results, we developed an additive model that accounted for

differences in histological subtypes to investigate differentially expressed genes in cell lines with high

vs. low CA, and high vs. low MN frequencies (**Fig. 7a, b**). Consistent with observation that CA and MN

are moderately correlated (Spearman's R = 0.38, p = 0.0015; **Fig. 7c**), cell lines with either high CA or

high MN frequencies showed similar gene expression signatures, including downregulation of E2F

target genes, and upregulation of genes involved in interleukin and interferon signalling (**Fig. 7c, d**).

The most upregulated pathway in MN-high cell lines was interferon alpha response signalling. In

contrast, the most significantly upregulated pathway in CA-high cell lines was TNFα via NFκB signalling

(**Fig. 7c–f**). NFκB/TNFα signalling pathways play pleiotropic roles in cell homeostasis and stress

response[39,40], and might therefore be required for cell survival in response to CA.

Centrosome amplification does not correlate with distinct patterns of copy number aberrations

but is associated with increased genome subclonality

280   Multiple studies have reported putative associations between centrosome abnormalities and

aneuploidy/CIN[10,30,34]. However, the underlying mechanisms for this association remain poorly

understood. We hypothesised that supernumerary centrosomes might present a previously

undescribed mutational process in HGSOC and induce distinct patterns of genomic aberrations. We

therefore performed low-coverage/shallow whole genome sequencing (sWGS) and absolute copy

number (ACN) fitting of HGSOC tissue samples to identify copy number signatures[8] that might be

associated with the CA phenotype. No significant correlations were observed between copy number

signatures and CA scores following p-value adjustment for multiple comparisons using samples with

high confidence ACN fits (n=54 OV04 and n=84 BriTROC samples; **Supplementary Fig. 4a, b**) Further,

no associations were found between CA and the number of breakpoints, copy number change points,

oscillating copy number or copy number segment size (**Supplementary Fig. 4c, d**). Importantly, we did

also not observe a correlation between CA scores and tumour ploidy (**Supplementary Fig. 4e**),

suggesting that the high prevalence of CA in HGSOC is not causally related to whole genome

duplication or cytokinesis failure.

294   ACN fitting from sWGS data from FFPE tumour tissue samples is challenging[41,42] and can be

confounded by poor sequencing quality as a result of FFPE processing, tumour heterogeneity and low

tumour purity. We therefore next used our cell line collection to further investigate potential CA-

associated patterns of genomic aberration. Importantly, our ovarian cancer cell lines were highly

representative of genomic features observed in the HGSOC cases (**Supplementary Fig. 5**)[8,43,44].

Consistent with our observations in HGSOC cases, we did not observe significant correlations between

ploidy and CA (Spearman's R = 0.13, p = 0.34, respectively; **Fig. 8a**). This further confirms that

cytokinesis failure and other processes impacting tumour ploidy may not be involved in driving CA in ovarian cancer.

To test the relationship between CA and CIN in our cell lines, we next quantified genome-wide copy number aberrations by estimating trimmed median absolute deviation from copy number neutrality (tMAD) scores[45] across all cell lines (**Fig. 8b–d**, **Supplementary Fig. 5**). Both CA and MN frequencies significantly correlated with tMAD scores (Spearman's R = 0.36, p = 0.0063; Spearman's R = 0.46, p = 0.00043, respectively); and cell lines which had high CA and high MN frequencies showed the most extreme genomic aberration (Spearman's R = 0.48, p = 0.00019; **Fig. 8b–d**). Consistent with our HGSOC cases, there were no significant associations between CA and copy number signatures (**Fig. 8e**). However, MN frequencies showed a significant negative correlation with signature 1 (Spearman's R = −0.52, p ≪ 0.0001, p.adj = 0.0003), and were positively correlated with signature 4 (Spearman's R = 0.42, p = 0.0011, p.adj = 0.007), and signature 5 (Spearman's R = 0.4, p = 0.0015, p.adj = 0.007; **Fig. 8f**). Given the compositional nature of copy number signatures, we also used a fixed effect model on the additive log ratio (ALR)-transformed copy number data to confirm these results and to test for global differential abundance of copy number signature exposures between both, CA high vs. low, and MN high vs. low cell lines (**Fig. 8g, h**). These analyses confirmed no statistically significant shift in copy number signature activities between CA high and low cell lines (Wald test, p = 0.913), but showed a significant change in copy number signature abundances observed in MN high vs. low cell lines (Wald test, p = 0.0083). This shift is most likely caused by the decrease in signature 1 (corresponding to the signature with the most extreme ALR coefficient) in MN high cell lines. Copy number signature 1 is characterised by a low number of breakpoints and large copy number segment size[8]. Consistent with these results, MN frequencies were strongly correlated with a higher number of breakpoints per chromosome arm (Spearman's R = 0.52, p ≪ 0.001) and negatively correlated with mean segment size (Spearman's R = −0.51, p = 0.000039). In contrast, we did not observe any correlations between CA frequencies and genomic features, including segment size, number of breakpoints, copy number change points, and oscillating copy number (**Supplementary Fig. 6**).

Finally, ongoing CIN is considered a principal mediator of intra-tumour heterogeneity[11]. Given the strong link between CA and CIN, we estimated fractions of subclonal segments (genome subclonality) from sWGS data as previously described[41,46] for all sequenced cell lines. Strikingly, CA frequencies were significantly correlated with genome subclonality (Spearman's R = 0.36, p = 0.0069, p = 0.054), while a similar trend was observed for MN (Spearman's R = 0.26, p = 0.054; **Fig. 7i–j**). This suggests that while CA is not associated with specific patterns of genome aberrations in our experimental setting (as read out by sWGS), it may play an important role in driving intra-tumour heterogeneity and tumour evolution.

Cell lines with high centrosome amplification show decreased chemosensitivity

Experimentally expanding the degree of aneuploidy in cells can increase their ability to adapt and develop resistance to chemotherapeutic agents[2,5,6] and in this study, CA showed strong correlations with both genome subclonality and CIN (**Fig. 7b–d, i**). We therefore hypothesised that CA might be an integrative and quantitative biomarker for predicting resistance to chemotherapy in ovarian cancer cell

lines. To test the relationship between CA and drug sensitivity, we selected seven CA-high and seven CA-low cell lines using a panel of eleven drugs focusing on standard-of-care chemotherapeutics as well as drugs targeting centrosome clustering, the spindle assembly checkpoint (SAC) and different components of the centrosome duplication and cell cycle (**Fig. 9a**). To control for variations in cell proliferation/division rates and assay duration, we used the growth rate inhibition (GR) metric[47,48] and estimated both potency (GR50, i.e., the concentration of a drug at which cell proliferation is reduced by 50%) and efficacy (GRmax, i.e., the maximum effect of a drug at the highest concentration tested) for each therapeutic agent (an ideal drug would have high potency and high efficacy).

We observed that drugs targeting SAC components or components of the centrosome duplication/cell cycle were highly effective in all fourteen cell lines inducing cytotoxic responses (**Fig. 9b**). By contrast, CW069, which specifically impedes centrosome clustering by inhibiting the microtubule motor protein HSET, was the least effective and only induced partial cytotoxicity in four out of the 14 lines. Paclitaxel, which is the standard of care treatment for both newly diagnosed and recurrent HGSOC, was the most potent drug. Strikingly, cell lines with high CA had decreased drug response compared to cell lines with low CA. This observation was most marked for paclitaxel treatment which induced cytotoxicity in low CA lines, but only partial cell growth inhibition in high CA lines (Spearman's R = 0.65, p = 0.014; **Fig. 9c**). Similarly, BI2536, Volasertib (both PLK1 inhibitors) and AZ3146 (Mps1 inhibitor) showed significantly lower potencies, while Oxaliplatin, and Barasertib (Aurora B inhibitor) showed significantly lower efficacies in CA-high compared to CA-low cell lines (**Fig. 9c**). These data confirm that CA predicts differential response to mitotic and centrosomal targeting agents and has clear relevance to current use of paclitaxel in the clinic.

Discussion

Centrosome amplification is highly prevalent in many cancer types. However, the in-depth characterisation of centrosome abnormalities and associated molecular features has been limited to small cohorts of tissue samples and cell lines, and previous approaches have not addressed the challenge of significant intra-tumoural heterogeneity. We developed a high-throughput microscopy-based approach to facilitate robust detection of centrosomes and to ensure adequate tissue representation. We found CA to be a highly prevalent and heterogeneous feature of HGSOC. Interestingly, a recent study in prostate cancer reported centrosome loss as another centrosomal abnormality that was associated with aneuploidy and disease progression[49], and studies on frozen ovarian cancer tissues suggest that regions of centrosome loss may be common in some tumour regions[26]. In our study, only one tissue sample showed significantly lower CA scores (tissue-wide) than those detected in normal control tissues. However, given the high-throughput data-driven approach deployed in this study, and the large amount of imaging data collected (>10,000 images across >300 FFPE tissues), we did not further inspect individual tissue regions for loss of centrosomes. Nevertheless, consistent with the previous study [26], we observed large variability in the spread of CA scores across all imaging fields within the same tissue, confirming that the CA phenotype displays marked inter- and intra-tissue heterogeneity in HGSOC tumours[22,26]. Both the high degree of CA heterogeneity and the confounding effect of stromal cells underscore the importance of acquiring multiple imaging fields per tumour sample and highlight a key strength of our imaging approach (see **Supplementary Methods**). In contrast to previous observations in other cancer types[25,30], we did not observe a correlation between the CA phenotype and disease outcome. This might be explained by the high degree of severe CIN observed in virtually all HGSOC cases[1] which is itself associated with poor outcome.

In addition, we conclusively confirmed that CA through centriole overduplication is a widespread phenomenon in ovarian cancer cells, recapitulating the high prevalence of CA observed in HGSOC tissues. Cell lines with high CA were associated with the presence of micronuclei, increased nuclei intensity (often used as a surrogate for DNA content) and mitotic indices, suggesting that cells with CA are delayed in mitosis. This is consistent with previous studies which have reported that cells with multiple centrosomes can take up to twice as long to complete mitosis, as additional time is required during the SAC to cluster centrosomes and align chromosomes in the metaphase plate[19,50–52].

HGSOC is characterised by frequent whole genome duplication and severe CIN. We recently reported the development of copy number signatures summarising distinct patterns of genomic features associated with CIN[8,9]. Importantly, we did not observe correlations between CA and ploidy in either our HGSOC tissue sections or our ovarian cancer cell lines. This strongly suggests that while whole genome duplication through cell fusion, mitotic slippage and/or cytokinesis failure may induce CA in some cells, it is not the main driver of CA in ovarian cancer cell lines and tissues. In this study, CA did also not correlate with known HGSOC copy number signatures, in either tumour tissues or ovarian cancer cell lines. Importantly, early and ubiquitous *TP53* mutations in HGSOC provide a permissive environment for a multiple mutational processes, including CA, to evolve and shape the HGSOC genome[8,12]. This makes it highly challenging to decipher out distinct mutational patterns linked to CA. Single-cell sequencing or third generation long-read sequencing technologies might therefore be

required to identify genomic patterns associated with CA. Nevertheless, our findings further support the notion that CA correlates with increased genomic aberration, can result in micronuclei formation, and thus likely presents a key driver of CIN[10,16,53] in HGSOC. Importantly, we also observed that high level CA is associated with increased genome subclonality. CA might therefore play an important role in driving intra-tumour heterogeneity and tumour evolution.

Ongoing CIN and tumour heterogeneity are associated with increased treatment resistance by providing tumours with the ability to more readily adapt to environmental stresses[2,4–6,11]. In line with these findings, we observed decreased sensitivities to several therapeutic agents in cell lines with higher levels of CA. This observation was most severe in cells treated with paclitaxel. While paclitaxel induced cytotoxic responses in CA low cell lines, CA high cell lines only showed partial growth inhibition. Importantly, supernumerary centrosomes have increased capacities for microtubule nucleation[54]. Consequently, higher concentrations of paclitaxel might be required in cells with high CA to inhibit spindle assembly and thus cell division. Centrosome-driven microtubule nucleation is regulated by the PCM, the size of which changes throughout the cell cycle and in preparation for spindle formation. We reported that HGSOC tumours frequently contain centrosomes with enlarged PCM foci in comparison to normal control tissues. However, due to FFPE staining limitations, we were not able to determine whether these size variations were true structural centrosome abnormalities or caused by staining artefacts and/or overlapping centrosomes that were not correctly segmented during spot counting. Further analyses on PCM size variations should be performed to investigate the role of enlarged PCM in CIN and paclitaxel sensitivities.

Lastly, RNA sequencing experiments revealed significant upregulation of TNFα/NFκB signalling in cell lines with high CA. Importantly, TNFα and NFκB signalling have been implicated in various aspects of oncogenesis[39,40,55,56]. Activation of NFκB results in the transcriptional upregulation of anti-apoptotic and antioxidant genes, promoting cell proliferation and survival. We therefore propose that NFκB signalling might be a key mediator of the CA "set point" allowing cells to tolerate higher levels of CA. Interestingly, a recent study illustrated CA-associated induction of the oxidative stress response via increased reactive oxygen species (ROS)[57]. High levels of ROS are toxic to cells and can induce cell senescence and apoptosis. Constitutive NFκB activity might therefore provide an important antioxidant mechanism resulting in cell survival and CA tolerance. NFκB-mediated upregulation of survival factors, together with increased levels of CIN and subclonality, might also contribute toward the CA-associated drug resistance observed in this study. In addition, NFκB signalling induces the transcription of many cytokines and chemokines, including IL6 and IL8, which regulate inflammatory responses, as well as cell invasion and migration[56]. Our RNA sequencing experiments showed that IL6 signalling was among the most upregulated signalling pathway in CA high cell lines. These findings further support the recent discovery that CA induces non-cell autonomous invasion through the secretion of pro-invasive factors including both, IL6 and IL8[57]. Together, these data highlight NFκB as a potential mediator of CA-associated oncogenic features, such as increased resistance and invasion. Future studies will be needed to further investigate the exact mechanisms and relationship between NFκB signalling and CA in HGSOC.

441   In summary, our large-scale characterisation of centrosome profiles in both ovarian cancer cell lines and tissue samples revealed that CA, through centriole overduplication, is a highly prevalent yet heterogeneous feature of HGSOC, and links supernumerary centrosomes to CIN, genome subclonality and thus tumour evolution. In addition, we highlight the centrosome and CA-associated survival mechanisms as promising targets for novel therapeutic approaches in HGSOC granting further investigation. In particular, our data shows that CA predicts differential response to standard-of-care paclitaxel with clear implications for the clinical disease management of HGSOCs. Finally, our work provides an important resource detailing the phenotypic, genomic, and transcriptomic characterisation of >70 ovarian cancer cell lines. This will facilitate the informed selection of suitable cell line models for future pre-clinical research investigating CIN, CA and associated vulnerabilities in ovarian cancer.

Methods

Clinical samples and primary tissue processing and selection

HGSOC cases were selected from the CTCR-OV04 study, which is a prospective non-interventional cohort study approved by the local research ethics committee at Addenbrooke's Hospital, Cambridge, UK, (REC reference numbers: 07/Q0106/63; and NRES Committee East of England - Cambridge Central 03/018). Tumour samples were processed following standardised operating protocols as outlined in the OV04 study design. For the generation of FFPE tissue blocks, tumour tissues were suspended in 10% neutral buffered formalin (NBF) for 24 hours and subsequently transferred into 70% ethanol for paraffin embedding and sectioning. Cases were chosen based on HGSOC histology and specimen availability from diagnosis, and represented real world clinical pathways with 29% from immediate primary surgery and 67% from delayed primary surgery. The sample set originally comprised a total of 120 FFPE tissues from 105 OV04 patients enrolled between January 2010 and April 2018. However, 27 samples were excluded prior to centrosome analyses owing to poor tissue quality and purity. The BriTROC-1 study enrolled patients with recurrent ovarian high grade serous or grade 3 endometrioid carcinoma who had relapsed following at least one line of platinum-based chemotherapy between January 2013 and September 2017 (REC reference number 12/EE/0349)[31,58]. BriTROC TMAs were generated using 247 specimens collected at diagnosis from 172 cases. Three 1 mm cores were collected from each donor block and placed into three sister TMAs with identical layout. Prior to centrosome analyses, H&E stains of resulting TMAs were inspected, and tissue cores/samples that mostly contained stromal or necrotic regions were excluded (n = 53). An overview of all remaining tumour samples (n = 287) and patients (n = 226) from both cohorts included in this study is shown in **Supplementary Table 1**. In addition to the 287 tumour samples, we also included 24 normal fallopian tube (FT) samples, one normal spleen, one normal kidney and five normal liver samples.

Cell Lines

Cell lines used in this study are listed in **Supplementary Table 2**, together with their associated growth conditions and culture media. In general, cell lines were grown at 37℃ and 5% CO_2 according to ATCC/ECACC recommendations. OSE medium is composed of 50:50 medium 199 (Sigma-5017) and medium 105 (Sigma-6395). Cells were tested for mycoplasma contaminations on a regular basis using the qPCR PhoenixDx Mycoplasma kit (Procomcure Biotech), and cell line identities were confirmed prior to DNA extractions using our in-house human short tandem repeat (STR) profiling cell authentication service.

Cell line growth curves

Cell line growth curves were generated to estimate optimal cell seeding densities for subsequent drug and centrosome characterisation assays. Cell lines were seeded into 96-well plates at different densities ranging from 5,000–20,000 cells/well. Plates were placed inside a real-time live-cell system (IncuCyte®, Sartorius) and monitored for four days. Phase contrast images were acquired at 3-hour intervals and confluence measures were collected. Cell seeding densities that resulted in 80–90% confluency at the end of the assay (i.e. four days) were chosen for each cell line.

Immunofluorescent centrosome staining

    1. FFPE tissue samples (also see **Supplementary Methods**)

Tissue samples were fixed in 10% NBF for 24 hours, embedded in paraffin and cut into 25 μm thick sections. Subsequently, tissue slides were deparaffinised and rehydrated by resuspending them into 1xPBS three times for 5 min each. Antigen retrieval was performed via heat-induced epitope retrieval (HIER) in universal antigen retrieval buffer (ab208572, Abcam). Deparaffinised tissue slides were immersed in antigen retrieval buffer which was brought to boil using a conventional microwave. Once boiling temperature was reached, tissue slides were boiled for 10 mins at medium-high power and subsequently incubated at room temperature for 25–30 mins to allow buffer and slides to cool down. Slides were washed three times in 1xPBS for 5 min each. Subsequently, tissue sections were permeabilised in permeabilisation buffer (5% BSA, 0.3% TritonX100) for 25–30 mins, and blocked in 5% BSA for 2 hours at room temperature.

    Primary antibodies (anti-Pericentrin, Abcam ab28144; anti-CDK5RAP2, Gergely Lab) were diluted in 5% BSA, added to the tissue slides and incubated at 4 °C overnight. The anti-CDK5RAP2 antibody was used at a concentration of 2 μg/ml and the ab28144 anti- Pericentrin antibody was used at a concentration of 5 μg/ml. Following incubation with primary antibodies, tissue slides were thoroughly washed three times in 0.1% Tween 20 in 1xPBS for 8 mins each. Secondary antibodies (Alexa Fluor 488 and Alexa Fluor 555, cat No.: A-11001 and A-21429, Invitrogen) were diluted in 5% BSA to a concentration of 4 μg/ml (1:500) and incubated for 1 hour at 37 °C. Subsequently, slides were washed again three times in 0.1% Tween 20 in 1xPBS for 8 mins each, rinsed with distilled water and counterstained with 1 μg/ml Hoechst (33342, B2261 Sigma-Aldrich) for 15 mins at room temperature. Tissue slides were mounted in glycerol / n-propyl gallate medium (4% n-propyl gallate (02370, Sigma-Aldrich), 8 ml 100% glycerol, and 2 ml 1× PBS) and incubated for 24–48 hours protected from light to allow tissue equilibration and diffusion of mounting media prior to imaging.

    2. Cell lines

Optical 96–well plates (CellCarrier-96 Ultra, Perkin Elmer) were coated with sterile Poly-L-Lysine (P4832, Sigma). Cells were seeded using previously estimated cell seeding densities and cultured for four days until confluent (see **Fig. 5a** for an overview of the experimental set-up). Cells were fixed in 100% ice-cold methanol for 5 mins at −20 °C, subsequently washed three times in 1×PBS and stored in 200 μl PBS at 4 °C until further use. Cells were rinsed with 0.1% Tween 20 in 1×PBS and permeabilised with 1% TritonX100 + 0.5% NP40 in 1xPBS for 5 mins at room temperature. Subsequently, cells were blocked for 1 hour at room temperature using 5% BSA in 1×PBS. Primary non-conjugated antibodies were diluted in blocking buffer, added to the cells and incubated overnight at 4 °C. An overview of antibodies, provider information, and concentrations for each of the four stains is shown in **Supplementary Table 3**. Following primary antibody incubation, cells were washed three times with 0.1% Tween 20 in 1×PBS. Secondary antibodies (Alexa Fluor 555 and Alexa Fluor 647, cat No.: A-21429 and A-21236, Invitrogen) were diluted to 1 μg/ml (1:2000) in blocking buffer, added to the cells and incubated for 1 hour at 37 °C. Subsequently, cells were washed three times with 0.1% Tween 20 in 1×PBS, and re-blocked with 5% BSA in 1×PBS for 30 mins at room temperature. Subsequently,

conjugated primary antibodies (Table 7.3) were diluted, added to the cells and incubated for 1 hour at room temperature, after which cells were washed again three times with 0.1% Tween 20 in 1×PBS. After the last washing step, cells were rinsed in Milli-Q water. Hoechst 33342 was diluted to 1 µg/ml in MilliQ water, added to the cells and incubated for 15-30 mins at room temperature. Cells were washed and stored in 1×PBS until further use.

Microscopy

1. SP8

For method optimisation purposes (see **Supplementary Methods**), sequential confocal images were acquired using a Leica TCS SP8 microscope. Z-stacks were collected at a step size of 0.35 µm. The pinhole diameter for channels containing centrosome staining signals was set to 0.5 AU. White light lasers with wavelengths of 488 and 555 nm were used to excite AlexaFluor 488 and AlexaFluor 555, respectively. Hoechst signal was excited using a laser diode 405.

2. Operetta

For large scale centrosome profiling experiments, the Operetta CLS™ high-content analysis system was used. For FFPE tissue sections, the Operetta PreciScan™ feature was used to allow automatisation of imaging processes – whole microscopy slides were initially scanned a 5× (PreScan), global images were filtered using Gaussian blur and tissue regions were identified based on Hoechst 33342 signal. The geometric centre of the tissue was then calculated and expanded to create a region of interest covering most of the tissue. Following morphological properties calculations, 50 independent non-overlapping imaging fields were placed into the tissue region of interest and subsequently imaged (ReScan) at high resolution using a confocal 63× 1.15NA water objective (see **Supplementary Methods**). For each imaging field, Z-stacks were generated at a step size of 0.5µm (100 planes in total) across all imaging channels (i.e. 405 nm, 488 nm and 555 nm wavelengths).

Cell lines were imaged in filtered 1 × PBS as mounting medium (200 µl/well) using a confocal 40× 1.1NA water objective. Images from five independent non-overlapping imaging fields were acquired from each well (see **Fig. 5a**). For each imaging field, Z-stack images were collected at a step size of 0.5 µm (28 planes in total) across all imaging channels (i.e. 405 nm, 488 nm, 555 nm and 647 nm wavelengths).

Image Analysis

Image analyses were performed using the Harmony software. Image analysis scripts were optimised for different sample types and immunofluorescent stains. In brief, images were reconstructed as maximum projections (collapsed Z-stacks) using basic brightfield correction. Image analyses included segmentation of nuclei, removal of border objects to only include whole cells, identification of cytoplasm and regions of interest, and subsequent centrosome detection and spot counting (see **Supplementary Methods** for a detailed description). Each image analysis step included a series of quality controls based on morphological and intensity properties of identified objects. The centrosome amplification score (CA score) was estimated as the total number of centrosomes divided by the total number of nuclei detected in each imaging field. To account for immunofluorescent staining background noise,

autofluorescence and the resulting rate of false positive centrosome detection, imaging fields with CA
scores < 0.1 were removed from downstream analysis (see **Supplementary Methods** and
Supplementary Methods Fig. 7 for more information and quality control cut-off estimation).

Statistical modelling of centrosome amplification score (CA score) variabilities
To model the distribution of CA scores across imaging fields within tissue samples and within tissue
types (displayed in **Fig. 2**), we developed a linear mixed model[59] allowing

1. to detect differences in average CA scores between tissue types by means of fixed effects,
2. to take the CA within-tissue dependence into account by means of normally distributed random
intercepts,
3. the spread (variance) of within-tissue CA scores to vary between tissues by means of gamma
distributed standard deviations and to vary between tissue types by means of fixed effects,
4. the mean and variance of within-tissue CA scores to be dependent by using a Gaussian copula
to model the association between the random intercepts of point 2 and random standard
deviations of point 3,
5. the model residuals to be heavy-tailed by considering the generalised T (taking the Gaussian
distribution as limit case) as conditional error distribution (i.e., distribution of the error terms
given the fixed and random effects described in points 1 to 4).

Parameter estimates and corresponding inference were respectfully obtained by means of the iterative
bootstrap[60] and the indirect inference method[61]. Monte Carlo simulations showed that the estimator was
consistent and that the confidence interval coverages were close to the 0.95 nominal values.

H&E purity estimation
H&E sections from FFPE tissues were sent to our pathologist for tumour marking and purity estimation.
In addition, H&E sections were scanned and subjected to HALO, an image analysis platform for
quantitative tissue analysis in digital pathology. HALO's random forest classifier was used to separate
the H&E image into tumour, stroma and microscope glass slide, allowing tumour purity estimation.

DNA and RNA extraction
1. FFPE tissue samples
For each FFPE sample, multiple sections at 10 µm thickness were cut depending on tissue size and
tumour cellularity assessed by a pathologist, who marked tumour areas on separate Haematoxylin and
Eosin (H&E) stained sections to guide microdissection for DNA extraction. Marked tumour areas from
unstained FFPE sections were scraped off manually using a scalpel, dewaxed in xylene and
subsequently washed with 100% ethanol. Following complete removal and evaporation of residual
ethanol (10 mins at 30℃) DNA was extracted using the AllPrep DNA/RNA FFPE Kit (Qiagen). DNA was
eluted in 40 µl Elution buffer.
2. Cell Lines
Cell pellets of approximately 1×10^6 cells were generated from cultured cells for each cell line outlined
above and stored at −80℃ until further use. DNA and RNA were extracted from cell pellets using the

Maxwell® RSC Cultured Cells DNA and RNA Kit (Promega, AS1620 and AS1390, respectively) with

the Maxwell® RSC 48 Instrument (Promega, AS8500).

DNA sequencing

1. Tagged-Amplicon Sequencing (TAmSeq)

Extracted DNA samples were diluted to a final concentration of 10 ng/ml using PCR certified water.

Tagged-Amplicon deep sequencing was performed as previously described[62]. In short, libraries were

prepared in 48.48 Juno Access Array Integrated Fluidic Circuits chips (Fluidigm, PN 101-1926) on the

IFC Controller AX instrument (Fluidigm) using primers designed to assess small indels and single

nucleotide variants across the coding region of the TP53 gene. Following target-specific amplification

and unique sample indexing, libraries were pooled and subsequently purified using AMPure XP beads.

Quality and quantity of the pooled library were assessed using a D1000 genomic DNA ScreenTape

(Agilent 5067-5582) on the Agilent 4200 TapeStation System (G2991AA), before submitting the library

for sequencing to the CRUK CI Genomics Core Facility using 150bp paired-end mode on either the

NovaSeq 6000 (SP flowcell) or HiSeq 4000 system. Sequencing reads were aligned to the 1000

Genomes Project GRCh37-derived reference genome (i.e. hs37d5) using the 'BWA-MEM' aligner.

OV04 tissue and cell line data analysis and variant calling was performed as previously described[62,63].

TAmSeq data and variant calls for the BriTROC study subset was obtained from Smith et al. (2022)[58].

2. Shallow Whole Genome Sequencing (sWGS)

DNA extractions were performed as described above, and quantified using Qubit quantification

(ThermoFisher, Q328851). DNA samples were diluted to 75 ng in 15 μl of PCR certified water and

sheared by sonication with a target of 200–250bp using the LE220-plus Focused-Ultrasonicator

(Covaris) with the following settings: 120 sec at room temperature; 30% duty factor; 180W peak incident

power; 50 cycles per burst. Sequencing libraries were prepared using the SMARTer Thruplex DNA-Seq

kit (Takara), with each FFPE sample undergoing 7 PCR cycles, and all other samples undergoing 5

PCR cycles for unique sample indexing and library amplification. AMPure XP beads were used following

manufacturer's recommendations to clean prepared libraries, which were subsequently eluted in 20 μl

TE buffer. sWGS libraries were quantified and quality-checked using D5000 genomic DNA

ScreenTapes (Agilent 5067-5588) on the Agilent 4200 TapeStation System (G2991AA) before pooling

the uniquely indexed samples in equimolar ratios. Pooled libraries were sequenced at low coverage

(~0.4 × coverage) on either NovaSeq 6000 S1 flowcells with paired-end 50 bp reads for clinical tissue

samples, or the HiSeq 4000 with single 50 bp reads, at the CRUK CI Genomic Core Facility. Resulting

sequencing reads were aligned to the 1000 Genomes Project GRCh37-derived reference genome (i.e.

hs37d5) using the 'BWA' aligner (v.0.07.17) with default parameters.

Copy number analyses

Relative copy number data for OV04 tissue samples and ovarian cancer cell lines was obtained using

the QDNAseq[64] R package (v1.24.0) to count reads within 30 kb bins, followed by read count correction

for sequence mappability and GC content, and copy number segmentation. QDNAseq data were then

subjected to downstream analyses using the Rascal R package[41] for ploidy and cellularity estimation

and absolute copy number fitting. Absolute copy number data for BriTROC tissue samples was obtained

from Smith et al. (2022) utilising comparable analysis approaches[58].

In all data sets, putative driver amplifications were detected and identified using the Catalogue

Of Somatic Mutations In Cancer (COSMIC; https://cancer.sanger.ac.uk/cosmic/help/cnv/overview)

definitions and thresholds for high level amplifications and homozygous deletions: *Amplification –*

average genome ploidy ≤2.7 and total copy number ≥5; or average genome ploidy >2.7 and total copy

number ≥9. *Deletion –* average genome ploidy ≤2.7 and total copy number = 0; or average genome

ploidy >2.7 and total copy number <(average genome ploidy 2.7). In contrast, copy number gains and

losses were defined as copy number > ploidy + 1, or copy number < ploidy − 1, respectively, and are

not defined as distinct copy number alterations in the same manner as amplifications and deletions.

Copy number signatures (CNS)

Copy number signatures were estimated as previously described[8] from absolute copy number data,

following computation of the distributions of six genomic features: I. the breakpoint count per 10 Mb, II.

the copy number of each segment, III. the copy number change points (i.e. the difference in copy

number between adjacent segments), IV. the breakpoint count per chromosome arm, V. the lengths of

chains of oscillating copy number segments, and VI. the size of segments.

Fixed effect model (for CNS analyses)

To account for the compositional nature of copy number signatures (the sum of copy number signature

exposures within each samples = 1), we performed additive log ratio (ALR) transformation[65] in a sample-

wise manner, choosing signature 7 as the denominator. Subsequently a fixed effect regression model

was applied to investigate differences in signature abundances between CA and MN high vs. low

samples. For determining global statistically significant shifts in copy number signature abundances we

used a Wald test.

tMAD scores

Trimmed median absolute deviation from CN neutrality (tMAD) scores were estimated as a genome-

wide summary measure of genomic abnormality as described before[45]ribed before[45]. The R code was

adjusted from https://github.com/sdchandra/tMAD. Samples were downsampled to $3{\times}10^6$ reads and

allocated into 500 kb non-overlapping bins as recommended by the authors. The diploid normal ovarian

surface epithelium cell line, IOSE4, which was the cell line with the least (no) copy number aberrations,

was chosen as normal/control sample for normalisation.

TCGA dataset analysis

Publicly available clinical data and RNAseq V2 data from The Cancer Genome Atlas (TCGA;

https://www.cancer.gov/about-nci/organization/ccg/research/structural-genomics/tcga) were

downloaded using Firebrowse (http://firebrowse.org/). Centrosome amplification gene expression

(CA20) signatures were calculated as previously described[30]. In brief, gene-level read counts were

quantile normalised using Voom[66] and the CA20 score was estimated for each sample as the sum of

the log2 median-centred gene expression levels of the CA20 signature genes: *AURKA*, *CCNA2*,
CCND1, *CCNE2*, *CDK1*, *CEP63*, *CEP152*, *E2F1*, *E2F2*, *LMO4*, *MDM2*, *MYCN*, *NDRG1*, *NEK2*, *PIN1*,
PLK1, *PLK4*, *SASS6*, *STIL*, and *TUBG1*. The CIN25 gene expression signature, as previously
defined[33], was estimated in the same manner using the following genes: *C20orf24*/*TGIF2*, *CCNB2*,
CCT5, *CDC45*, *CDC2*, *ESPL1*, *FEN1*, *FOXM1*, *H2AFZ*, *KIF20A*, *MAD2L1*, *MCM2*, *MCM7*, *MELK*,
NCAPD2, *PCNA*, *PRC1*, *RAD51AP1*, *RFC4*, *RNASEH2A*, *TOP2A*, *TPX2*, *TRIP13*, *TTK*, *UBE2C*.
697

RNA sequencing

699 Following RNA extractions, RNA integrity number (RIN) values were estimated using the Bioanalyzer
700 system (Agilent). RIN values ranged from 8.3–10 with a mean of 9.7. Library preparation and RNA
701 sequencing (RNAseq) were performed by the CRUK CI Genomics Core Facility: In brief libraries were
702 prepared using the Illumina TruSeq standard mRNA kit and sequenced using 50 bp paired-end mode
703 on the NovaSeq 6000 sequencer (S1 flowcell).
704

RNAseq analyses

Sequencing reads were mapped to the GRCh38 (hg38) reference genome using STAR aligner v2.7.6a
(https://github.com/alexdobin/STAR/releases). Following standard quality control (MultiQC[67]), raw
sequencing reads were quantified against the same reference genome (GRCh38) using Salmon
v1.4.0[68], https://combine-lab.github.io/salmon/). Quantified transcripts (gene-level) were imported into
R using the tximport[69] in preparation for downstream analyses. Low-abundance genes were filtered out,
keeping only genes with at least 25 reads in at least two samples. Gene count normalisation and
differential gene expression analyses were performed using DESeq2 v1.26.0[70]. Genes with p-value <
0.05, FDR < 0.1 and a fold-change of at least 1.5 were considered as differentially expressed. Likelihood
ratio tests were used to assess which model (simple, additive or interactive) was most appropriate for
each comparison. Data was visualised using the lfcShrink() function (DESeq2 package) and the
ComplexHeatmap R package v2.2.0[71]. Gene set enrichment analyses (GSEA) were performed using
the fgsea R package v1.12.0 (https://github.com/ctlab/fgsea). Genes were ranked by statistical
significance and fold change; and 50 hallmark sets from the human C5 MSigDB collection were used
for running GSEA. 1,000 permutations were used to calculate p-values. In addition, the clusterProfiler
R package v3.14.3[72] was used for gene ontology-based overrepresentation analyses.
721

Drug sensitivity assays – GR metrics

Cell lines were seeded into opaque-walled 384 well plates compatible with the CLARIOstar® Plus (BMG
Labtech) luminometer using a CombiDispenser (Thermo Scientific™). For each cell line, two plates
were generated – a control plate to correct for the number of divisions taking place over the course of
the assay, and a test plate. 24 hours after seeding, 10 μl of CellTiter-Glo® (Promega) were added to
the control plate using the CombiDispenser (Thermo Scientific™), and after an incubation time of 30
mins the plate was read using the CLARIOstar® Plus luminometer (BMG Labtech). At the same time
(24 hours post seeding), drugs were added to the test plate using a D300e Digital Dispenser (Tecan).
Plate layouts were randomised, normalised for DMSO volumes across wells, and drugs were dispensed

as 10-point half-log dilution series with a DMSO concentration of 0.01%. The following compounds were
used: oxaliplatin (Selleckchem S1224), paclitaxel (Selleckchem S1150), tozasertib (Selleckchem
S1048), alisertib (Selleckchem S1133), barasertib (Selleckchem S1147), cw069 (Selleckchem S7336),
bi2536 (Selleckchem S1109), volasertib (Selleckchem S2235), cfi400945 (Selleckchem S7552),
az3146 (Selleckchem S2731), and gsk923295 (Selleckchem S7090). The highest drug concentration
used for paclitaxel was 0.1 μM. For all other compounds, the highest tested concentration was 100 μM.
Drug screens were performed in triplicates. Following the addition of drugs, treatment plates were
incubated for 72 hours, after which cell viabilities were assessed as described above. To estimate drug
sensitivity, as well as drug potency and efficacy for each cell line and compound, the growth rate
inhibition metrics were estimated as previously described[47,48] using the GRmetrics R package v1.12.2
(https://bioconductor.org/packages/release/bioc/html/GRmetrics.html).

Acknowledgements

We thank all patients who participated in and donated tissue samples to this study. The Addenbrookes Human Research Tissue Bank is supported by the NIHR Cambridge Biomedical Research Centre. We also thank the OV04 study team for their help with clinical tissue samples. We would like to thank the Cancer Research UK Cambridge Institute Microscopy, Compliance & Biobanking, Genomics, Bioinformatics, and IT & Scientific Computing core facilities for their support with various aspects of this study. We would like to thank D. Provencher and A.-M Mes-Masson for kindly donating a subset of 25 ovarian cancer cell lines to us that were included in this study. These cell lines were derived with the support of the Banque de tissus et de données of the Réseau de recherche sur le cancer of the Fonds de recherche du Québec - Santé (FRQS) affiliated with the Canadian Tumor Repository Network (CTRNet). We also thank Fanni Gergely for her kind donation of the CDK5RAP2 antibody, and Gayathri Chandrasekaran for her help and advice on the centrosome staining protocol.

We acknowledge funding and support from Cancer Research UK, and the Cancer Research UK Cambridge Centre (A22905, A25177, A25117). L.M.G. was supported by the Wellcome Trust PhD programme in Mathematical Genomics and Medicine (grant number RG92770). This research was also supported by the Ovarian Cancer Action (grant number 006), and the NIHR Cambridge Biomedical Research Centre (BRC-1215-20014). Work in the Cancer Molecular Diagnostics Laboratory/Blood Processing Laboratory was supported by the NIHR Cambridge Biomedical Research Centre, Cancer Research UK Cambridge Centre and the Mark Foundation Institute for Integrated Cancer Medicine. The views expressed are those of the authors and not necessarily those of Cancer Research UK, the NIHR or the Department of Health and Social Care. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author Contributions

Conceptualization, C.M.S., J.T., M.V. and J.D.B.; Methodology, C.M.S. and D.L.C; Validation, C.M.S.; Formal Analysis, C.M.S. and D.L.C.; Investigation, C.M.S., J.A.H., A.M.P., J.G., J.T., M.V.; Resources, C.M.S., D.L.C., T.B., A.P., A.S., M.D.E., P.S., K.H., M.A.V.R., L.M.G, A-M.M-M., D.E., D.M., A.H., I.A.M., M.J.L., F.C.M., and J.T.; Data Curation, C.M.S., D.L.C., T.B., P.S., and K.H.; Writing - Original Draft, C.M.S.; Writing - Review & Editing, C.M.S., J.T., M.V., and J.D.B.; Visualisation, C.M.S.; Supervision, C.M.S., J.T., M.V., and J.D.B.; Project Administration, C.M.S. and M.V. Funding Acquisition, J.D.B.

Conflicts of Interest

J.D.B. is a co-founder and shareholder of Tailor Bio Ltd. There are no further conflicts of interests to declare.

References

1. Morden, C. R. *et al.* Chromosome instability is prevalent and dynamic in high-grade serous
ovarian cancer patient samples. *Gynecol Oncol* (2021) doi:10.1016/j.ygyno.2021.02.038.

2. Ippolito, M. R. *et al.* Gene copy-number changes and chromosomal instability induced by
aneuploidy confer resistance to chemotherapy. *Dev Cell* **56**, 2440-2454.e6 (2021).

3. Kim, C. *et al.* Chemoresistance Evolution in Triple-Negative Breast Cancer Delineated by Single-
Cell Sequencing. *Cell* **173**, 879-893.e13 (2018).

4. Lee, A. J. X. *et al.* Chromosomal instability confers intrinsic multidrug resistance. *Cancer Res*
71, 1858–1870 (2011).

5. Lukow, D. A. *et al.* Chromosomal instability accelerates the evolution of resistance to anti-cancer
therapies. *Dev Cell* **56**, 2427-2439.e4 (2021).

6. Shaikh, N. & McClelland, S. E. Diversity in chromosome numbers promotes resistance to
chemotherapeutics. *Dev Cell* **56**, 2399–2400 (2021).

7. Minussi, D. C. *et al.* Breast tumours maintain a reservoir of subclonal diversity during expansion.
Nature **592**, 302–308 (2021).

8. Macintyre, G. *et al.* Copy number signatures and mutational processes in ovarian carcinoma.
Nat Genet **50**, 1262–1270 (2018).

9. Drews, R. M. *et al.* A pan-cancer compendium of chromosomal instability. *Nature 2022 606:7916*
606, 976–983 (2022).

10. Ganem, N. J., Godinho, S. A. & Pellman, D. A mechanism linking extra centrosomes to
chromosomal instability. *Nature* **460**, 278–282 (2009).

11. McGranahan, N., Burrell, R. A., Endesfelder, D., Novelli, M. R. & Swanton, C. Cancer
chromosomal instability: Therapeutic and diagnostic challenges. *EMBO Rep* **13**, 528–538
(2012).

12. Tamura, N. *et al.* Specific Mechanisms of Chromosomal Instability Indicate Therapeutic
Sensitivities in High-Grade Serous Ovarian Carcinoma. *Cancer Res* **80**, 4946–4959 (2020).

13. Conduit, P. T. *et al.* The centrosome-specific phosphorylation of Cnn by Polo/Plk1 drives Cnn
scaffold assembly and centrosome maturation. *Dev Cell* **28**, 659–669 (2014).

14. Conduit, P. T., Wainman, A. & Raff, J. W. Centrosome function and assembly in animal cells.
Nat Rev Mol Cell Biol **16**, 611–624 (2015).

15. Nigg, E. A. & Holland, A. J. Once and only once: mechanisms of centriole duplication and their
deregulation in disease. *Nat Rev Mol Cell Biol* **19**, 297–312 (2018).

16. Godinho, S. A. & Pellman, D. Causes and consequences of centrosome abnormalities in cancer.
Philosophical Transactions of the Royal Society B: Biological Sciences **369**, 20130467–
20130467 (2014).

17. Silkworth, W. T., Nardi, I. K., Scholl, L. M. & Cimini, D. Multipolar spindle pole coalescence is a
major source of kinetochore mis-attachment and chromosome mis-segregation in cancer cells.
PLoS One **4**, (2009).

18. Mason, J. M. *et al.* Functional characterization of CFI-400945, a polo-like kinase 4 inhibitor, as
a potential anticancer agent. *Cancer Cell* **26**, 163–176 (2014).

817 19. Kwon, M. *et al.* Mechanisms to suppress multipolar divisions in cancer cells with extra
818   centrosomes. *Genes Dev* **22**, 2189–2203 (2008).

819 20. Marteil, G. *et al.* Over-elongation of centrioles in cancer promotes centriole amplification and
820   chromosome missegregation. *Nat Commun* **9**, (2018).

821 21. Tischer, J. & Gergely, F. Anti-mitotic therapies in cancer. *Journal of Cell Biology* **218**, 10–11
822   (2019).

823 22. Dias Louro, M. A., Bettencourt-Dias, M. & Bank, C. Patterns of selection against centrosome
824   amplification in human cell lines. *PLoS Comput Biol* **17**, 1–22 (2021).

825 23. Wong, Y. L. *et al.* Reversible centriole depletion with an inhibitor of Polo-like kinase 4. *Science*
826   *(1979)* **348**, 1155–1160 (2015).

827 24. Galofré, C. *et al.* Centrosome reduction in newly-generated tetraploid cancer cells obtained by
828   separase depletion. *Sci Rep* **10**, 1–12 (2020).

829 25. Chan, J. A clinical overview of centrosome amplification in human cancers. *Int J Biol Sci* **7**,
830   1122–1144 (2011).

831 26. Morretton, J.-P. *et al.* A catalog of numerical centrosome defects in epithelial ovarian cancers.
832   *EMBO Mol Med* **14**, e15670 (2022).

833 27. Donne, R., Sangouard, F., Celton-Morizur, S. & Desdouets, C. Hepatocyte polyploidy: Driver or
834   gatekeeper of chronic liver diseases. *Cancers (Basel)* **13**, 1–19 (2021).

835 28. Guidotti, J. E. *et al.* Liver cell polyploidization: A pivotal role for binuclear hepatocytes. *Journal*
836   *of Biological Chemistry* **278**, 19095–19101 (2003).

837 29. Margall-Ducos, G., Celton-Morizur, S., Couton, D., Brégerie, O. & Desdouets, C. Liver
838   tetraploidization is controlled by a new process of incomplete cytokinesis. *J Cell Sci* **120**, 3633–
839   3639 (2007).

840 30. de Almeida, B. P., Vieira, A. F., Paredes, J., Bettencourt-Dias, M. & Barbosa-Morais, N. L. Pan-
841   cancer association of a centrosome amplification gene expression signature with genomic
842   alterations and clinical outcome. *PLoS Comput Biol* **15**, 1–31 (2019).

843 31. Goranova, T. *et al.* Safety and utility of image-guided research biopsies in relapsed high-grade
844   serous ovarian carcinoma - Experience of the BriTROC consortium. *Br J Cancer* **116**, 1294–
845   1301 (2017).

846 32. Ogden, A., Rida, P. C. G. & Aneja, R. Prognostic value of CA20, a score based on centrosome
847   amplificationassociated genes, in breast tumors. *Sci Rep* **7**, 1–11 (2017).

848 33. Carter, S. L., Eklund, A. C., Kohane, I. S., Harris, L. N. & Szallasi, Z. A signature of chromosomal
849   instability inferred from gene expression profiles predicts clinical outcome in multiple human
850   cancers. *Nat Genet* **38**, 1043–1048 (2006).

851 34. Raff, J. W. & Basto, R. Centrosome Amplification and Cancer: A Question of Sufficiency. *Dev*
852   *Cell* **40**, 217–218 (2017).

853 35. Ciriello, G. *et al.* Emerging landscape of oncogenic signatures across human cancers. *Nat Genet*
854   **45**, 1127–1133 (2013).

855 36. Woodruff, J. B., Wueseke, O. & Hyman, A. A. Pericentriolar material structure and dynamics.
856   *Philos Trans R Soc Lond B Biol Sci* **369**, 20130459- (2014).

37.   Tischer, J., Carden, S. & Gergely, F. Accessorizing the centrosome: new insights into centriolar
appendages and satellites. *Curr Opin Struct Biol* **66**, 148–155 (2021).

38.   Mittal, K. *et al.* Hypoxia drives centrosome amplification in cancer cells via HIF-1α-dependent
induction of polo-like kinase 4. *Molecular Cancer Research* **4**, molcanres.MCR-20-0798-E.2020
(2021).

39.   Luo, J. L., Kamata, H. & Karin, M. IKK/NF-κB signaling: Balancing life and death - A new
approach to cancer therapy. *Journal of Clinical Investigation* **115**, 2625–2632 (2005).

40.   Tornatore, L. *et al.* Cancer-Selective Targeting of the NF-κB Survival Pathway With
Gadd45β/Mkk7 Inhibitors. *Cancer Cell* **26**, 495–508 (2014).

41.   Sauer, C. M. *et al.* Absolute copy number fitting from shallow whole genome sequencing data.
bioRxiv 2021.07.19.452658 (2021).

42.   Carter, S. L. *et al.* Absolute quantification of somatic DNA alterations in human cancer. *Nat
Biotechnol* **30**, 413–421 (2012).

43.   Bell, D. *et al.* Integrated genomic analyses of ovarian carcinoma. *Nature* **474**, 609–615 (2011).

44.   Bowtell, D. D. *et al.* Rethinking ovarian cancer II: reducing mortality from high-grade serous
ovarian cancer. *Nat Rev Cancer* **15**, 668–679 (2015).

45.   Mouliere, F. *et al.* Enhanced detection of circulating tumor DNA by fragment size analysis. *Sci
Transl Med* **10**, 1–14 (2018).

46.   van Dijk, E. *et al.* Chromosomal copy number heterogeneity predicts survival rates across
cancers. *Nat Commun* **12**, 1–12 (2021).

47.   Hafner, M., Niepel, M. & Sorger, P. K. Alternative drug sensitivity metrics improve preclinical
cancer pharmacogenomics. *Nat Biotechnol* **35**, 500–502 (2017).

48.   Hafner, M., Niepel, M., Chung, M. & Sorger, P. K. Growth rate inhibition metrics correct for
confounders in measuring sensitivity to cancer drugs. *Nat Methods* **13**, 521–527 (2016).

49.   Wang, M., Nagle, R. B., Knudsen, B. S., Cress, A. E. & Rogers, G. C. Centrosome loss results
in an unstable genome and malignant prostate tumors. *Oncogene* **39**, 399–413 (2020).

50.   Acilan, C. & Saunders, W. S. A Tale of Too Many Centrosomes. *Cell* **134**, 572–575 (2008).

51.   Basto, R. *et al.* Flies without Centrioles. *Cell* **125**, 1375–1386 (2006).

52.   Yang, Z., Lončarek, J., Khodjakov, A. & Rieder, C. L. Extra centrosomes and/or chromosomes
prolong mitosis in human cells. *Nat Cell Biol* **10**, 748–751 (2008).

53.   Vitre, B. D. & Cleveland, D. W. Centrosomes, chromosome instability (CIN) and aneuploidy.
Curr Opin Cell Biol **24**, 809–815 (2012).

54.   Godinho, S. a *et al.* Oncogene-like induction of cellular invasion from centrosome amplification.
Nature **510**, 167–171 (2014).

55.   Godwin, P. *et al.* Targeting nuclear factor-kappa B to overcome resistance to chemotherapy.
Front Oncol **3 MAY**, 1–10 (2013).

56.   Taniguchi, K. & Karin, M. NF-κB, inflammation, immunity and cancer: Coming of age. *Nat Rev
Immunol* **18**, 309–324 (2018).

57.   Arnandis, T. *et al.* Oxidative Stress in Cells with Extra Centrosomes Drives Non-Cell-
Autonomous Invasion. *Dev Cell* **47**, 409-424.e9 (2018).

58.   Smith, P.-S. *et al.* The genomic landscape of recurrent ovarian high grade serous carcinoma:
898   the   BriTROC-1   study.   *medRxiv*   2022.10.21.22280992   (2022)
899   doi:10.1101/2022.10.21.22280992.

900   59.   McCulloch, C. E., Searle, S. R. & Neuhaus, J. M. Generalized, linear, and mixed models, 2nd
901   edn. 424 (2008).

902   60.   Kuk, A. Y. C. Asymptotically Unbiased Estimation in Generalized Linear Models with Random
903   Effects. *Journal of the Royal Statistical Society: Series B (Methodological)* **57**, 395–407 (1995).

904   61.   Guerrier, S., Dupuis-Lozeron, E., Ma, Y. & Victoria-Feser, M. P. Simulation-Based Bias
905   Correction Methods for Complex Models. *https://doi.org/10.1080/01621459.2017.1380031* **114**,
906   146–157 (2018).

907   62.   Forshew, T. *et al.* Noninvasive identification and monitoring of cancer mutations by targeted
908   deep sequencing of plasma DNA. *Sci Transl Med* **4**, (2012).

909   63.   Piskorz, A. M. *et al.* Methanol-based fixation is superior to buffered formalin for next-generation
910   sequencing of DNA from clinical cancer samples. *Annals of Oncology* **27**, 532–539 (2016).

911   64.   Scheinin, I. *et al.* DNA copy number analysis of fresh and formalin-fixed specimens by shallow
912   whole-genome sequencing with identification and exclusion of problematic regions in the
913   genome assembly. *Genome Res* **24**, 2022–2032 (2014).

914   65.   Aitchison, J. The Statistical Analysis of Compositional Data. *Journal of the Royal Statistical
915   Society: Series B (Methodological)* **44**, 139–160 (1982).

916   66.   Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. Voom: Precision weights unlock linear model
917   analysis tools for RNA-seq read counts. *Genome Biol* **15**, 1–17 (2014).

918   67.   Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: Summarize analysis results for
919   multiple tools and samples in a single report. *Bioinformatics* **32**, 3047–3048 (2016).

920   68.   Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast and bias-
921   aware quantification of transcript expression. *Nat Methods* **14**, 417–419 (2017).

922   69.   Soneson, C., Love, M. I. & Robinson, M. D. Differential analyses for RNA-seq: Transcript-level
923   estimates improve gene-level inferences. *F1000Res* **4**, 1–19 (2016).

924   70.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for
925   RNA-seq data with DESeq2. *Genome Biol* **15**, 1–21 (2014).

926   71.   Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in
927   multidimensional genomic data. *Bioinformatics* **32**, 2847–2849 (2016).

928   72.   Yu, G., Wang, L. G., Han, Y. & He, Q. Y. ClusterProfiler: An R package for comparing biological
929   themes among gene clusters. *OMICS* **16**, 284–287 (2012).

Main Figures

Figure 1 – Centrosome characterisation in HGSOC tissue samples.

(**a**) Basic workflow of centrosome characterisation in clinical tissue samples (also see **Supplementary Methods**). Example confocal immunofluorescent images (max. intensity projections) of a fallopian tube and HGSOC tissue are shown in (**b**) and (**c**), respectively (left panel). Nuclei are shown in blue, centrosomes are shown in green. Individually detected centrosomes following automated image analysis are highlighted in rainbow colours (right panel; nuclei shown in grey). Scale bars = 50 μm. (**d**) and (**e**) show sample type comparison of centrosome amplification (CA) scores and centrosome size, respectively. Sample types are indicated by different colours. Dashed horizontal line indicates CA cutoff threshold of ~1.83.

Figure 2 – Centrosome amplification profiles across 318 tissue samples.

(**a**) Centrosome amplification scores across 287 HGSOC tissue samples, 24 fallopian tube, a spleen, a kidney, and
5 liver samples. Each boxplot represents CA scores across up to 50 imaging fields from individual tissue samples.
Different tissue types are indicated by different colours. Dashed vertical line indicates the CA cutoff threshold of
~1.83. Samples with median CA scores > CA cutoff were considered to have significant CA.

Figure 3 – Centrosome amplification shows inter- and intra-tissue heterogeneity.

Spatial heatmap of centrosome amplification in a normal fallopian tube (**a**) and HGSOC tumour tissue (**b**).
Additional examples are shown in **Supplementary Fig. 1**. CA scores are indicated by a colour gradient (purple =
low, yellow = high) for each imaging field. Points represent individual nuclei detected during image analyses and
were plotted in relation to their physical position (μm) on the microscope slide (x and y axis). Note that CMYK
printing may obscure differences on the CA score colour scale. (**c**) Comparison of CA scores across tissues
collected from individual patients in a total of 36 cases. Red stars indicate CA score medians for patient with multiple
tissue samples. (**d**) Comparison of CA scores across different tissue sites. (**e**) Comparison of CA tissue
heterogeneity across different tissue sites. Tissue sites are indicated by different shapes. Cohorts are indicated by
teal (OV04) and light blue (BriTROC) colours. Dashed horizontal lines indicate the CA score cutoff of ~1.83.

Figure 4 – Centrosome amplification is not associated with clinical features and disease
outcome in HGSOC.

Comparison of CA scores across different (**a**) histotypes, (**b**) disease stages at diagnosis, (**c**) germline
BRCA1/BRCA2 mutation status, and (**d**) surgery types during sample collection (IPS = immediate primary surgery;
DPS = delayed primary surgery). (**e**) Comparison of CA tissue heterogeneity in IPS vs. DPS cases. Forest plots of
multivariable Cox proportional hazard modelling on overall survival for (**f**) OV04 and (**g**) BriTROC patients with and
without CA. Adjusted covariates included surgery type, stage and age. OV04 samples are shown in teal, BriTROC
samples are shown in light blue. Note that for patients with multiple tissue samples, the median CA score across
these tissues was used.

Figure 5 – In-depth imaging-based characterisation of ovarian cancer cell lines.

(a) Cell line characterisation workflow. Growth curves were estimated for each cell line to determine optimal seeding
densities. Cells were then seeded according to estimated seeding densities into an optical, poly-L-lysine coated 96
well plate and incubated for 4 days until they reached confluence (late Log/early Stationary phase). Cells were
fixed with 100% methanol and stained against indicated proteins. For each well, five independent fields were

imaged using the indicated Operetta CLS™ system (40× confocal water objective). (**b**) Immunofluorescent imaging
results for 68 screened cell lines ordered by the median percentage of non-mitotic cells with 2 or more centrosomes
(boxplots). Cell lines highlighted by grey arrowheads are depicted in **Fig. 6a-b**. Panel 1 shows the average number
of nuclei included in each of the four staining screens. Bar plots are coloured by cell line subtype (from publicly
available information (Cellosaurus) and Domcke et al. (2013)). Panel 2 indicates the mitotic index for each cell line.
The fraction of non-mitotic cells with two or more centrosomes is shown in panel 3 (boxplots). Each point represents
an individual imaging field. The mean (population wide) CA score (= $\frac{\sum \text{Centrosomes}}{\sum \text{Nuclei}}$) is shown by light blue bars. Panel
4 shows the centrosome size (estimated from max. intensity projection images), and panel 4 indicates the
percentage of CEP164-positive centrosomes. Micronuclei (MN) frequencies, estimated as percentage of non-
mitotic cells with at least one or more MN, are shown in panel 6.

Figure 6 – Centrosome amplification is associated with increased micronuclei frequencies and

can be induced by low oxygen growth conditions.

Confocal immunofluorescent staining images (zoomed in from 40× max. intensity projection images) for (**a**) CA-low and (**b**) CA-high cell lines as highlighted in **Fig. 5b**. Cells were stained for Pericentrin (yellow), pHH3 (mitotic cells; red), DNA (Hoechst; blue) and cytokeratin (not shown for easier centrosome visualisation). Scale bar = 50 μm. (**c**) Cell line characterisation correlation plot whosing Spearman's rank correlations of analysed immunofluorescent staining results. P-values: *** P < 0.001; ** P < 0.01; * P < 0.05; . P < 0.1. (**d-e**) Cell line subtype comparison of centrosome amplification and micronuclei frequencies, respectively. Histological subtypes are

994   indicated by different colours. Note that the three "normal" cell lines were grouped together with other subtypes, as
995   these cell lines are transformed and show some cancer characteristics. (**f-g**) Centrosome amplification and
996   micronuclei frequencies, respectively, in cell lines grown at 21% vs. 5% oxygen. (**h-i**) Cell lines, which are normally
997   grown in normoxic conditions (21% oxygen), were transferred into 5% oxygen and centrosome amplification (**h**)
998   and micronuclei frequencies (**i**) were estimated.
999

Figure 7 – Differential gene expression in cells with high centrosome amplification and
micronuclei frequencies.

1002

Differential gene expression volcano plots of all genes significantly regulated by (**a**) CA and (**b**) MN frequencies
(high vs. low; median split). FDR < 0.1 highlighted in lightblue/blue. The top 25 most significantly up- or
downregulated genes are labelled. (**c-d**) Gene set enrichment analysis (GSEA) of centrosome and micronuclei
high vs. low cell lines, showing hallmark pathways for which p < 0.05 and FDR < 0.1. Examples of enrichment
score plots for centrosome and micronuclei GSEA pathway results are shown in (**e**) and (**f**), respectively. Light

1008   blue/blue lines illustrate the running sum for each gene set shown. Maximum and minimum enrichment score (ES)
1009   are indicated by orange dotted lines. Leading-edge gene subset is indicated by vertical black lines.

1010 **Figure 8 – Genomic characterisation of centrosome amplification and micronuclei.**

1011

(**a**) Correlation of centrosome amplification (CA) with cell line ploidy. Coefficients were estimated using Spearman's
rank correlations. (**b-d**) Correlations of cell line tMAD scores with CA frequencies, micronuclei (MN) frequencies
and CA × MN (combined) frequencies, respectively. tMAD = trimmed median absolute deviation from copy number
neutrality. Note that normal cell lines were excluded from this analysis. (**e-f**) Correlations of copy number signature
exposures with CA and MN frequencies, respectively. Different signatures are indicated by colours. Intercepts and
coefficients (slopes) and their standard errors of the fixed effect model of ALR-transformed copy number signature
data for CA and MN frequencies are shown in (**g**) and (**h**). (**i-j**) Correlations of genome subclonality estimated from
ACN fits with CA and MN frequencies. All correlation coefficients were estimated using Spearman's rank
correlations.

1021

Figure 9 – Cell lines with high centrosome amplification show decreased drug sensitivities.

1023

(**a**) Ovarian cancer cell line drug-screen overview showing drugs used in our drug sensitivity assay and their
respective targets within the cell cycle. (**b**) Variation in potency and efficacy of selected drugs a cross all tested
ovarian cancer cell lines (n=14). Colours of box plots indicate drug targets as shown in panel a. (**c**) Correlation of
centrosome amplification frequencies and drug potency (GR50; top panel) and drug efficacy (GRmax; bottom
panel). Spearman's rank correlations are shown for each drug. Drug agents are colour-coded according to their
respective drug target groups. Note that higher GR values correspond to lower potency or efficacy. The signs of
GRmax values indicated drug response: GRmax > 0 indicates partial growth inhibition; GRmax = 0 indicates
complete cytostasis; GRmax < 0 indicates cell death (cytotoxicity).