Real-time TIRF imaging of single adiponectin vesicle exocytosis in 3T3-L1 adipocytes

stably expressing mCherry fused to human adiponectin

Man Mohan Shrestha[1], Sebastian Barg[2] and Charlotta S. Olofsson[1]

[1]Department of Physiology/Metabolic Physiology, Institute of Neuroscience and Physiology, Institute

of Biochemistry, The Sahlgrenska Academy, University of Gothenburg, Göteborg, Sweden.

[2]Department of Medical Cell Biology, Uppsala University, Uppsala, Sweden.

Running title: *Imaging hormone exocytosis in live adipocytes*

12
13

To whom correspondence should be addressed: Dr Man Mohan Shrestha and Associate Professor

Charlotta Olofsson, University of Gothenburg, The Sahlgrenska Academy, Dept. Neuroscience and

Physiology, Section of Metabolic Physiology, Postal address: Box 432, SE-405 30 Göteborg, Sweden,

E-mail: Man Mohan Shrestha: man.shrestha@gu.se. Charlotta Olofsson: charlotta.olofsson@gu.se

Keywords: adiponectin, exocytosis, TIRF imaging, white adipocyte, chimeric protein expression

Abstract

Adiponectin is a peptide hormone abundantly released from adipocytes, and reduced

circulating levels are associated with obesity-related diseases, such as type 2 diabetes, insulin

resistance and cardiovascular disease. Adiponectin is released by regulated exocytosis of

secretory vesicles, but traditional molecular biology and imaging techniques lack the

specificity and time resolution to adequately quantify exocytosis and trafficking of adiponectin-

containing vesicles. Here we generated 3T3-L1 adipocytes that stably express mCherry-tagged

human adiponectin, resulting in robust labelling of small adiponectin vesicles with a diameter

of 200-300 nm, in live cells. Total internal reflection fluorescence (TIRF) microscopy was used

to visualise and quantify exocytosis and adiponectin release in real-time, observed as rapid

disappearance of the fluorescence of individual vesicles. Bulk adiponectin secretion

measurements confirmed that the labelled adiponectin was secreted to the surrounding solution

under these conditions, and expressed in the same vesicle population as endogenous

adiponectin. In contrast to previous electrophysiological results, elevation of cytosolic Ca^{2+}

alone was sufficient to induce exocytosis, although at a lower rate compared to elevated

cytosolic cAMP. We conclude that the adiponectin-mCherry-labelled cells are useful for

studying adiponectin exocytosis at the single vesicle level, and that an intracellular elevation

of either cAMP or Ca^{2+} can trigger adiponectin vesicle release.

Introduction

The adipocyte-derived hormone adiponectin has anti-diabetic, anti-atherogenic and anti-inflammatory effects and acts as an insulin sensitizer. Thus, adiponectin plays crucial roles in metabolic disorders, including insulin resistance, type 2 diabetes and cardiovascular disease (1, 2). Overexpression of adiponectin is associated with protection against both acute and chronic effects of high fat diet-induced lipotoxic effects of lipid accumulation, and enhances the metabolic flexibility of adipose tissue (3).

A number of studies support that adiponectin is secreted via regulated exocytosis of pre-formed adiponectin-containing vesicles. Lim et al. demonstrated the vesicular localisation of adiponectin and that insulin stimulates release of measurable amounts of adiponectin at time-points between 60 and 120 minutes (4). The insulin-induced secretion was unaffected by pre-incubation with the protein synthesis inhibitor cyclohexamide, indicating that the secretagogue stimulates release of vesicles belonging to a pool of already synthesised vesicle. Our own work has defined an alternate pathway for adiponectin exocytosis: By combining electrophysiological and biochemical techniques with mathematical modelling, we have demonstrated that adiponectin exocytosis is triggered by an increase of intracellular cAMP and activation of exchange factor directly activated by cAMP, isoform 1 (Epac1). The cAMP/Epac1-dependent adiponectin release occurs in the absolute absence of Ca^{2+} (shown by buffering with the Ca^{2+} chelator BAPTA), but is none the less potentiated by an elevation of cytosolic Ca^{2+}. This regulation of adiponectin exocytosis exists in cultured 3T3-L1 adipocytes as well as in primary mouse and human adipocytes. Membrane capacitance recordings (that measure exocytosis as the increase in plasma membrane area that occurs upon vesicle fusion) showed that the cAMP/Ca^{2+}-triggered adiponectin exocytosis can be detected within 1-2 min after intracellular addition of stimuli. The stimuli consisted of inclusion of cAMP with or without Ca^{2+} in the patch pipette solution washing into the recorded cell, or extracellular addition of the cAMP-elevating agent forskolin and/or the Ca^{2+} ionophore ionomycin (5-7). Later work demonstrated that pre-treatment of adipocytes with cyclohexamide was without effect also on the cAMP-triggered adiponectin secretion (8). Collectively, our work proposes that pre-stored adiponectin vesicles undergo rapid exocytosis upon stimulation with cAMP in both the presence and absence of Ca^{2+}, in agreement with the cell model in Fig. 1.

Total internal reflection fluorescence (TIRF) microscopy is an optical technique that enables the excitation of fluorophores in a very narrow optical section. TIRF microscopy provides a means to selectively excite fluorophores near the adherent cell surface, while counter-

minimizing fluorescent signals from other intracellular regions (10). The narrow TIRF

excitation field allows high-resolution visualisation of processes occurring near or in the

plasma membrane, in live cells (11). Thus, by applying TIRF microscopy to adipocytes

expressing a fluorophore localised to adiponectin vesicles, their exocytosis can be visually

observed at single cell and single vesicle level, at high spatial and temporal resolution.

Here, we describe the generation of 3T3-L1 cells that stably express mCherry fused to the C-

terminal end of human adiponectin, to produce a genetically homogenous clonal cell population

that over-express the recombinant protein of interest. We demonstrate that the adiponectin-

mCherry expressing cells can be differentiated into mature adipocytes, as judged by lipid-

filling. Using TIRF microscopy, we can visualise and quantify Ca^{2+} and cAMP-triggered

adiponectin exocytosis in real-time, at the single-vesicle resolution.

Material and methods

Safety and Handling of Recombinant Lentiviruses

87 All the lentiviruses were generated using a third generation packaging system as per Biosafety

88 Level 2 and institutional guidelines of the University of Gothenburg. Disposable plastic

89 pipettes were used and liquid waste was decontaminated with 10 % bleach. Laboratory

90 materials in contact with viral particles were treated as biohazardous waste and autoclaved.

Cell culture

HEK293T (ATCC) cells were cultured in high glucose DMEM (Invitrogen, GIBCO)

supplemented with 10 % fetal bovine serum (FBS) and 1 % penicillin-streptomycin at 37 ^{0}C

and 5 % CO2. Cultured clonal 3T3-L1 fibroblasts (Zenbio) were seeded and differentiated into

mature adipocytes and used for experiments at day 8 or 9 according to previous establish

protocol (7). In brief, 3T3-L1 fibroblasts were cultured in high glucose DMEM (Invitrogen,

GIBCO) supplemented with 10 % new born calf serum at 37 ^{0}C and 5 % CO2. Two days after

confluence, cells were washed twice with warm PBS and a differentiation medium containing

10 % FBS, 1 % penicillin-streptomycin, 0.85 μM insulin, 0.5 mM 3-isobutyl-1-methylxanhine

and 1 μM dexamethasone was added for 2 days. The differentiation medium was there after

replaced with medium containing 10 % FBS and insulin for another 2 days. Cells were then

maintained in DMEM with 10 % FBS. Adipocytes were used for experiments on day 8 or 9

post differentiation. Only 3T3-L1 adipocytes that appeared mature, as judged visually by
accumulated lipid, were studied.

Generation of lentiviral constructs

Three candidate hairpin sequences for mouse adiponectin (NM_009605.5) each containing 21
sense bases that are identical to the target gene, a loop containing an XhoI restriction site and
21 antisense bases that are complementary to the sense bases followed by a poly T termination
sequence, were designed using an small interfering RNA (siRNA) selection tool hosted by the
Whitehead Institute for Biomedical Research (http://sirna.wi.mit.edu/) (12). Starting at 25nt
downstream of the start codon (ATG), we searched for 21nt sequences that matched the pattern
AA(N19). It was ascertained that the G-C content was >35 %. To minimize degradation of off-
target mRNAs, the NCBI's BLAST program was used. Since some sequences are likely more
effective than others, multiple target sequences were selected for the *Acrp30* gene (Table 1).

The annealed oligos were inserted into an empty backbone third generation transfer plasmid
(pLKO.1-TRC cloning vector; Plasmid #10878; addgene) using AgeI (5' cloning site) and
EcoRI (3' cloning site) restriction digestion sites. Similarly, a candidate hairpin sequence for
green fluorescent protein (GFP) was used for a control/scramble (pLKO.1 GFP; Plasmid
#30323; addgene). mCherry fused to human adiponectin (hAcrp30-mCherry) was cloned into
a pLenti vector using NheI and HpaI restriction digestion sites to generate lentiviral constructs
(Supplementary Fig. 1A). The sequences of the knockdown and lentiviral constructs were
verified using restriction digestion and Sanger's Sequencing from Eurofins Genomics.

Lentivirus Packaging and Infection

Lentiviruses were produced by co-transfection of HEK293T cells with lentiviral expression
and packaging plasmids using the calcium phosphate transfection method (13, 14). Viral
supernatant was collected after 48 hours of transfection, centrifuged at 1,000 x g for 5 minutes
and filtered through 0.45 µm filter. The viruses were further concentrated by ultracentrifugation
using Ti45 rotor in Beckman Coulter Optima-L-100 XP at 116,000 x g for 2 hours at 4 °C. For
gene silencing, 3T3-L1 fibroblasts were transduced with titrated viruses in presence of 8 µg/mL
of polybrene (Hexadimethrine bromide, Cat. No: 107689-10G, Sigma-ALDRICH) and
subjected to antibiotic selection using 4 µg/mL puromycin after 48 hours. The successful stable
expression was verified by fluorescence microscopy for fusion proteins whereas knockdown
efficiency was assessed by Western blotting and RT-qPCR at protein and mRNA level

respectively. For generation of a stably expressing mCherry fused adiponectin cell line, 3T3-L1 fibroblasts were infected with optimum amount of viral particles in the presence of polybrene. The growth medium was changed 24 hours after infection.

Adiponectin secretion

3T3-L1 cells stably expressing adiponectin-mCherry were seeded and differentiated into mature adipocytes on 35 mm glass bottom dish (D35-20-1.5-N, Cellvis). The mature adipocytes were incubated with an extracellular solution containing 140 mM NaCL, 3.6 mM KCL, 0.5 mM $MgSO_4$, 0.5 mM NaH_2PO_4, 2.0 mM $NaHCO_3$ and 5.0 mM Hepes (pH 7.4) supplemented with 2.6 mM $CaCl_2$ and 5.0 mM glucose for 30 minutes at 32 °C. Test substances were added for 30 min at 32 °C under gentle shaking. At termination of the incubation, the medium containing secreted product was collected, and the adipocytes were lysed using PBS containing 2 % SDS and protease inhibitor (1 tablet per 10 mL; cOmplete Mini, Roche Diagnotics). Secretion samples were centrifuged for 5 minutes at 2000 rpm and 4 ^{0}C, and the supernatant was stored at -80 °C. Secreted HMW and total adiponectin (measured with HMW and Total Adiponectin ELISA (human) kit, 47-ADPMS-E01, Alpco) were expressed in relation to total protein content. Cell lysates were centrifuged at 14000 rpm for 10 minutes at 4 ^{0}C and total protein contents were measured by Bradford protein assay.

Lipolysis Assays

Matured adipocytes were incubated with an extracellular (EC) solution for 30 minutes at 32 °C. Lipolysis was stimulated by adding the adenylate cyclase activator Forskolin (10 μM) and the phosphodiesterase inhibitor isobutylmethylxanthine (IBMX) (200 μM) or Ionomycin (1 μM) in serum-free extracellular (EC) solution for 30 min at 32 °C under gentle shaking. At termination of the incubation, the medium containing secreted product was collected, and the adipocytes were lysed using PBS containing 2% SDS and protease inhibitor (1 tablet per 10 mL; cOmplete Mini, Roche Diagnotics). Cell lysates were centrifuged at 14000 rpm for 10 minutes at 4 ^{0}C and total protein contents were measured by Bradford protein assay. Secretion samples were centrifuged for 5 minutes at 2000 rpm and 4 ^{0}C, and the supernatant was collected. Lipolysis was assessed from the release of FFA and glycerol in the secretion samples using non-esterified fatty acids measurement kit (FUJIFILM Wako Chemicals Europe GmbH) and free glycerol reagents (G7793 and F6428, Sigma) respectively and normalised to total protein contents.

Real-Time Quantitative PCR

RNA was extracted and purified using QIAzol (Qiagen) and ReliaPrep RNA Cell MiniPrep System (Promega). 1 µg of RNA was used for reverse transcription. cDNA synthesis was performed according to the protocol as described in Invitrogen™ SuperScript™ II Reverse Transcriptase kit (Thermo Fisher Scientific #18064022) in presence of 40 units of RNaseOUT™ Recombinant Ribonuclease Inhibitor (Thermo Fisher Scientific #10777-019) and 200 units of Reverse Transcriptase per 20 µL of reaction mixture. The real-time quantitative PCR (qPCR) assays were performed using Fast SYBR Master Mix (4385612, Applied Biosystems). The relative expression of mRNA abundance was calculated using $2^{-\Delta\Delta Ct}$ and was normalized to β-actin. Data are represented as mean $\pm$ SEM, ****$p<0.0001$ (n=3).

Western blotting

3T3-L1 adipocytes were lysed with TNET buffer (50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 10 mM NaF, 1 mM EDTA and 1% Triton X-100) containing 2 mM Na_3VO_4 (Sodium Orthovanadate) and 1 mM pmsf (phenylmethylsulfonyl fluoride) and 1x protease inhibitor cocktail tablet (Roche). Whole cell lysates were centrifuged at 12,000 rpm at 4 ^{0}C for 10 minutes. The protein concentration was determined using Bradford protein assay. Proteins from whole cell lysate (20 µg) were separated by 8 % sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS/PAGE) and then transferred onto polyvinylidene difluoride (PVDF) membranes (Millipore) at 120 V for 2 hours using transfer buffer (25 mM Tris-base, pH 8.3, 190 mM glycine, 20 % methanol and 0.1 % SDS). After blocking in blocking buffer (2 % BSA in PBS containing 0.2 % Tween-20 and 0.01 % sodium azide) for 1 hour at room temperature, the membrane was incubated overnight at 4 ^{0}C with primary antibody against adiponectin (ab3455, Abcam) and tubulin (T9026, Sigma) as loading control at 1:1000 dilution. The primary antibodies were probed using HRP-conjugated secondary antibodies at 1:5000 dilution and detection was performed with enhanced chemiluminescence.

Live Cell imaging using TIRF microscopy and analysis of vesicle exocytosis

3T3-L1 adipocytes expressing hAcrp30-mCherry were pre-treated with extracellular solution for 30 minutes at 32 °C and kept in the same solution supplemented with 2.6 mM $CaCl_2$ and 5.0 mM glucose during imaging. Image acquisition was performed by TIRF Observer Z1 microscope with alpha Plan-Apochromat 100×/1.46 Oil objective (both Carl Zeiss) and an EMCCD camera (Evolve 512 delta, Photometrics). Images were acquired using Zeiss Zen 2.6

(blue edition). Excitation laser of wavelength 561 nm with exposure time 25–100 ms was used for mCherry. Forskolin together with the phosphodiesterase inhibitor 3-isobutyl-1-methylxanthine (FSK/IBMX) or ionomycin alone were added 1–2.5 minutes after acquisition start and the time point was noted. TIRF images were acquired over 15 minutes at 2 Hz through 1801 total readings. Circles of region of interest (ROI) around the vesicles were used to measure the fluorescence intensities of images over time. All the readings were normalized to the intensity measured at time point zero, after removal of the background fluorescence. Values were plotted against time, so as to indicate the time course of adiponectin containing vesicle exocytosis. Vesicle size was estimated using fluorescent polystyrene beads of known diameter. Vesicle locations were found by eye, a 21 pxl long line was centered on this location, and the brightness along this line was read out ("linescan"). The following piecewise linear function describing a trapezoid shape was then fitted to the linescan to estimate spot width: if (x1< x<x1+bl) y=a+s*(x-x1); if (x2-bl< x<x2) y=a+s*(x2-x); if (x1+bl<x<(x2-bl) y=a+s*bl; else y=a, where x1 and x2 are boundaries of the peak, a the peak amplitude, s the slope, bl the width of the two sloped sections, and y the brightness. The apparent width is w=x2-x1-bl, which is the full width at half maximum (FWHM) of the trapezoid. bl was fixed to 1.5pxl (240nm), the theoretical resolution of the system. Beads with known diameters were imaged and measured in the same way, to create a reference scale.

Analysis

Images were analysed using Zeiss Zen 2.6 (blue edition) and OriginPro (OriginLab Corporation). All data are presented as mean values ± SEM for designated number of experiments. Significance of difference was tested by Student's t-test and ANOVA. Statistical analysis were performed using Prism 9.0 (GraphPad, San Diego, CA).

Results

Knock down and rescue of endogenous adiponectin in 3T3-L1 adipocytes

3T3-L1 cells with low passage number were used and it was ascertained that the cells were able to express high levels of endogenous adiponectin when differentiated into adipocytes (Fig. 2A). To avoid that native adiponectin interferes with the expression of human adiponectin-mCherry, the endogenous adiponectin was knocked down by short hairpin RNA (shRNA). Three different target sequences for shRNA mouse adiponectin knockdown were

designed: shAcrp30-A, shAcrp30-B and shAcrp30-C (Table 1). The targets were inserted into

a lentiviral plasmid and 3T3-L1 fibroblasts were transduced with the viral particles. The

transduced cells were thereafter differentiated into mature adipocytes. For two out of three

knockdowns (shAcrp30-A and shAcrp30-C) transduced cells could be cultured into mature

adipocytes, whereas cells transduced with shAcrp30-B did not survive. The successful

knockdown was confirmed by protein and gene expression, where adiponectin gene

knockdown was compared with scrambled shRNA targeting GFP (Fig. 2B-D).

Cells that displayed the most efficient knockdown (shAcrp30-A), were rescued with hAcrp30-

mCherry alone (plasmid map in Suppl. Fig. 1) or with a FLAG tag either at the N- or C-terminal

in the fibroblast stage. (The FLAG was added to allow future protein purification by affinity

chromatography as well as localisation of the fusion protein in live cells.) The rescued cells

were seeded and differentiated into mature adipocytes, and the over-expression of hAcrp30

was verified by Western blotting using antibodies against adiponectin or FLAG. The hAcrp30-

mCherry tagged with FLAG at the C-terminal was found to be highly over-expressed compared

to the other two variants under both reduced (Fig. 3A) and non-reduced (Fig. 3B) conditions.

The over-expression of adiponectin-mCherry was visualized by inverted fluorescence

microscope (Fig. 3C) and TIRF microscopy (Fig. 3D); TIRF imaging demonstrated a punctate

pattern of fluorescence, as typical for vesicular compartments. Most of the labelled vesicles

appeared diffraction limited or slightly larger in size. Comparing the apparent width (FWHM)

of the vesicles with that of fluorescently labelled beads of known diameter allowed us to

estimate their median diameter to ~200-300 nm (Fig. 3E, F).

An elevation of intracellular cAMP or Ca^{2+} triggers exocytosis of mCherry-labelled

adiponectin vesicles

We have shown that a combination of the adenylyl cyclase activator forskolin and the

phosphodiesterase inhibitor 3-isobutyl-1-methylxanthine (FSK/IBMX) increases 3T3-L1

adipocyte cAMP levels >6-fold (7), and that the Ca^{2+} ionophore ionomycin elevates the

intracellular Ca^{2+} level ($[Ca^{2+}]_i$; (6). Thus, we investigated adiponectin vesicle exocytosis in

differentiated 3T3-L1 adipocytes stably expressing adiponectin-mCherry exposed to

FSK/IBMX or ionomycin. As shown in Fig. 4A, addition of FSK/IBMX led to disappearance

of mCherry-labelled vesicles (bottom images in Fig. 4A and Suppl. video 1). A rapid loss of

fluorescence puncta was observed upon stimulation, which we interpreted as exocytosis events

(top row Fig. 4C and Fig. 4D). Approx. 60% of the vesicles observed by TIRF microscopy

fused with the plasma membrane after addition of the stimulatory agents, while the remaining

40% were stable throughout the recording (middle row Fig. 4C and Fig. 4E). This is expected

since it is unlikely that all adiponectin vesicles that reside close to the plasma membrane, within

the TIRF zone, are primed for exocytosis (15). In experiments with adipocytes exposed to the

control solution during the entire experiment (Fig. 4B and Suppl. video 2), the labelled

adiponectin vesicles remained stable throughout the TIRF recording (bottom row Fig. 4C).

Calculation of the fraction of released vesicles confirmed that a larger number of vesicles were

exocytosed upon exposure to the cAMP-elevating agents compared to control (Fig. 4F). At the

end of 15 min recordings, significantly fewer adiponectin-mCherry labelled vesicles remained

in adipocytes exposed to FSK/IBMX, compared to control solution (Fig. 4G).

Ionomycin likewise triggered the exocytosis of adiponectin-containing vesicles (Fig. 5A, and

Suppl. video 3). Similar to the experiments with FSK/IBMX, a rapid loss of fluorescence

puncta was observed upon stimulation, indicating vesicle exocytosis (top row Fig. 5C and

Fig. 5D) and a fraction of adiponectin-containing vesicles remained stable throughout the

recording (middle row Fig. 5C and Fig. 5E). It was again confirmed that adiponectin-mCherry

vesicles remained stable in adipocytes maintained in control solution (Fig. 5B, bottom row

Fig. 5C and Suppl. video 4). The fraction of released vesicles was significantly larger in the

ionomycin-exposed adipocytes compared to control experiments, at all investigated time-

points (Fig. 5F). Significantly fewer adiponectin-mCherry labelled vesicles remained in

ionomycin-exposed adipocytes compared to control cells, at the end of 15 minute recordings

(Fig. 5G). As can be seen upon comparison of Figs. 4F and 5F, whereas 60% of the vesicles

residing in the TIRF zone were released after 10 min exposure to FSK/IBMX, the

corresponding fraction of secreted vesicles amounted to only 15% in ionomycin-treated

adipocytes.

Measurements of released adiponectin, FFA and glycerol in 3T3-L1 adipocytes stably

expressing adiponectin-mCherry

To verify the adiponectin secretion capacity of the stably expressing 3T3-L1 cells, the matured

adipocytes were incubated in the presence of FSK/IBMX or ionomycin and product secreted

to the medium was measured by ELISA. In agreement with previous results (6, 7) the

secretagogues stimulated the release of mouse adiponectin in native 3T3-L1 adipocytes

(Fig. 6A). The 3T3-L1 adipocytes stably expressing adiponectin-mCherry likewise secreted

adiponectin in response to both FSK/IBMX and ionomycin and both total (Fig. 6B) and the

high-molecular weight (HMW) form (Fig. 6C) of human adiponectin were released over basal

(unstimulated control cells). The data confirm that the stably expressing adipocytes secrete

smaller and larger forms of the adiponectin-mCherry fusion protein in response to cAMP or

Ca^{2+} stimulation, at levels similar to that observed in mouse native adipocytes. To further

validate the functionality of in the adiponectin-mCherry adipocytes, we measured release of

free fatty acids (FFA) and glycerol. FSK/IBMX elevated levels of both FFA and glycerol in

the incubation medium whereas ionomycin was without effect of release of either of the lipid

metabolites (Figs. 6D, E).

Discussion

Pre-stored neurotransmitter- or hormone-containing vesicles typically release their content to

the extracellular space upon receiving the correct triggering signal. Classically, an elevation of

cytosolic Ca^{2+}, released from intracellular stores or entering via Ca^{2+}-permeable ion channels

in the plasma membrane, stimulates vesicle exocytosis. Other intracellular mediators such as

cAMP and ATP also have important roles for vesicle maturation steps (15). Peptide hormone

exocytosis has been detailed in a number of endocrine cell types, including insulin secreting

beta-cells (16) and catecholamine-releasing chromaffin cells (17). However, the control of

hormone release in white adipocytes is a research field that has received little attention.

Adiponectin, one of the chief hormones secreted from white adipocytes and the most abundant

protein present in serum (18), has in several studies been reported to have favourable metabolic

effects and to protect from development of obesity-associated metabolic diseases (1, 2).

Although the general control of adiponectin secretion over shorter and longer time-periods has

been addressed in a small number of studies (4, 19-21), adiponectin vesicle exocytosis – the

molecular and cellular process in which adiponectin-containing vesicles are recruited to and

fuse with the adipocyte plasma membrane to release their cargo - is a largely unexplored field.

Here we have generated 3T3-L1 cells stably expressing human adiponectin fused with the

fluorescent protein mCherry, and we use the overexpressing matured adipocytes to visualise

and quantify adiponectin exocytosis. We demonstrate that adiponectin vesicle trafficking and

exocytosis can be studied at the single vesicle level, and we determine that an elevation of

intracellular Ca^{2+} stimulates adiponectin vesicles exocytosis in the absence of a concomitant

increase of cAMP. Below we discuss the usefulness of our cells for future studies of

adiponectin vesicle behaviour, and we reconsider the Ca^{2+}-dependence of adiponectin
exocytosis.

The adiponectin-mCherry protein is targeted to endogenous adiponectin vesicles that
resemble hormone-containing vesicles in other endocrine cell types
The addition of a fluorescent tag to a protein may disturb the correct targeting of the product.
Our TIRF data propose that the chimeric protein is present in vesicular compartments, indicated
by the punctate appearance of the fluorescence signal. Moreover, the real-time recordings
suggest that the overexpressing adipocytes are functional, and that adiponectin-mCherry is
targeted to the correct vesicle population; this is shown by that the labelled vesicles fuse with
the plasma membrane in response to signals known to stimulate adiponectin exocytosis. In
further support of this, measurements of released human adiponectin show that the fusion
protein is secreted in response to an elevation of cAMP or Ca^{2+}. Our estimated diameter of
200-300 nm for the adiponectin-labelled vesicles is similar to hormone-containing vesicles in
other endocrine cell types, such as insulin granules with a diameter of ~300 nm (26).

We have previously demonstrated that secretion of the hormones leptin, apelin and adipsin is
unaffected by cAMP in 3T3-L1 adipocytes (5) and that resistin is co-secreted with adiponectin
(contained within the same vesicles (37)). Collectively, our data propose that the rapid loss of
fluorescence puncta in response to cAMP or Ca^{2+} reflects exocytosis of single vesicles
belonging to the endogenous adiponectin vesicle population.

Both cAMP and Ca^{2+} trigger exocytosis of adiponectin-mCherry labelled vesicles
The real-time imaging data in Figs. 4 and 5, as well as Suppl. videos 1 and 3, confirm the
previously shown importance of cAMP and Ca^{2+} for stimulation of adiponectin vesicle release
(5-7, 27). However, the results in Fig. 5 and Suppl. video 3, where ionomycin triggers
exocytosis in the absence of a cAMP increase, can appear to contradict previous experiments
showing that Ca^{2+} alone is unable to induce exocytosis in 3T3-L1 adipocytes (6, 7). Notably,
there are some important difference between our previous patch-clamp data and the current
TIRF recordings: During the electrophysiological experiments, the interior of the adipocyte is
controlled by the pipette solution – thus by exclusion of cAMP in the pipette solution washing
into the cell, the experimental approach applied in (6, 7), the 3T3-L1 adipocytes can essentially
be depleted of cAMP. In contrast, the adipocytes used for TIRF imaging are metabolically
intact and have a continuous endogenous production of cAMP, maintaining basal levels of the

cyclic nucleotide. We suggest that the elevated Ca^{2+} acts together with preserved endogenous
levels of cAMP, to stimulate adiponectin exocytosis in micro-domains where cAMP is
sufficiently high. The finding that ionomycin on its own stimulates adiponectin secretion in
metabolically intact adipocytes is in agreement with data on secretion of the adipocyte hormone
resistin (28); we recently showed that resistin is co-secreted with adiponectin in mouse
adipocytes (thus contained within the same vesicles) and that ionomycin induced the release of
resistin in the absence of a cAMP-elevating agent (29). An alternative explanation to the effect
of ionomycin alone is that the increase in intracellular Ca^{2+} activates Ca^{2+}-dependent adenylyl
cyclases leading to production of cAMP. In Musovic et al. 2021, we were unable to demonstrate
a significant increase of cAMP levels in 3T3-L1 adipocytes exposed to ionomycin (Fig. 1C of
(29)). However, cAMP was measured in cell lysates, with no information of changes of cAMP
confined to specific near-membrane regions. Thus, ionomycin might stimulate local elevations
of cAMP, at sites of adiponectin vesicle exocytosis. It is well known that cAMP is
compartmentalised in adipocytes and that this regulates several intracellular processes (30). A
role of caveolae for the spatial control of cAMP has been demonstrated in cardiac myocytes,
and proposed to be relevant for βAR signalling (31). We have previously shown that caveolae
are necessary for the organisation of signalling pathways involved in the regulation of HMW
adiponectin exocytosis (32).

Vesicles that contain peptide/protein hormones typically undergo a series of maturation steps
including physical movement, docking (stable attachment to the plasma membrane) and
priming (chemical modifications that make docked vesicles release competent) before they can
discharge their cargo upon fusion with the plasma membrane (15). The data in Figs. 4 and 5 as
well as supplementary videos 1 and 3, show that both cAMP and Ca^{2+} trigger exocytosis of
adiponectin vesicles that are already docked at the adipocyte plasma membrane; no new
vesicles (newcomers) are observed to enter the TIRF zone. This observation seems to stand in
conflict with our previous electrophysiological data demonstrating that an elevation of Ca^{2+}
maintains adiponectin secretion over time, and with our conclusion that Ca^{2+} induces vesicle
replenishment. In this context, it is essential to keep in mind that we image only a small part
of/near the adipocyte plasma membrane, the ventral side of the cell attached to the culture dish,
whereas the capacitance recording reflect exocytosis that occur over the entire cell membrane.
Consequently, vesicles that move to and undergo exocytosis at the apical side of the adipocyte,
outside of the TIRF imaging zone, are not visualised.

Choice of cells and fluorescent protein

3T3-L1 cells were used to generate the adiponectin-mCherry expressing adipocytes. A number of studies have demonstrated that *in vitro* differentiated 3T3-L1 adipocytes are a relevant cell model for investigations of the adiponectin secretion process (5-7, 9, 22). In fact, all mechanisms and mediators determined to regulate adiponectin exocytosis in 3T3-L1 adipocytes have been confirmed in primary mouse and human adipocytes ((7-9) and Musovic & Olofsson, unpublished data). The 3T3-L1 cells are easy to culture and differentiate, and their functionality with regard to adipocyte metabolic processes, such as lipolysis (23) and glucose uptake (24), is well determined. Nonetheless, they are clonal cells and we acknowledge that results must be continuously validated in experiments with primary, preferably human, adipocytes. A disadvantage with the 3T3-L1 adipocytes is that their ability to differentiate into mature adipocytes ceases with increased cell passage. Although multiple vials of low passage cells can be stored, it will eventually be necessary to produce new stably expressing 3T3-L1 cells. However, we are not aware of any adipocyte cell line that does not have this limitation. Collectively, we regard the 3T3-L1 adipocytes to be a good choice for our studies.

A large number of fluorescent proteins are available for protein tagging and real-time studies of their specific function. The continuous development of fluorescent proteins with enhanced or altered qualities have yielded a supply that covers the full visible spectrum, with diverse advantages and disadvantages. Introduced nearly a decade ago, mCherry remains a frequently used fluorescent marker for cellular studies. The mCherry has several advantages: It is bright and it matures rapidly, enabling experiments to be performed soon after initiation of expression. It is also highly photostable and resistant to photobleaching (25). Since experiments must be carried out at high time-resolution and over an extended time-period (up to 30 min) the listed qualities are important. We therefore labelled adiponectin with mCherry, expressed under the CMV promoter that has high housekeeping expression. Human adiponectin was chosen to enable us to biochemically distinguish between endogenous mouse adiponectin and the overexpressed fusion protein. Nonetheless, there are some disadvantages related to using mCherry for studies of adiponectin exocytosis: Red fluorescent proteins display limited pH sensitivity compared to e.g. enhanced green fluorescent protein (EGFP) or pHluorin. Upon vesicle fusion with the plasma membrane the pH of the vesicle lumen (which is acidic) increases; pH-sensitive fluorescent proteins will consequently increase in brightness already upon formation of the vesicle-plasma membrane fusion pore, thus confirming vesicle fusion.

For this reason, we initially evaluated the use of EGFP for studies of adiponectin exocytosis,

but both the more rapid photobleaching of EGFP, and lipid autofluorescence interfering with

the green signal complicated the studies. With regard to pHluorin, this optical indicator is non-

fluorescent at low pH and thus does now allow investigations of events that occur prior to

vesicle fusion with the plasma membrane. We nonetheless acknowledge the advantage of using

a pH-sensitive fluorescent tag for studies of vesicle fusion and cargo release and we will

evaluate their use for future investigations of the adiponectin exocytosis process. We also

recognise that tagging adiponectin itself might interfere with both protein mulitmerisation and

the secretion process. However, the secretion data confirm that the stably expressing adipocytes

release abundant amounts of human HMW adiponectin, verifying multimerisation of the

chimeric protein. Regardless of this, it would be valuable to assess if labelling of another

exocytosis vesicle marker such as Neuropeptide Y (NPY), a proven reliable marker for insulin

granule exocytosis (25), is useful for real-time imaging of the adiponectin vesicle release

process.

What can we learn about the control of adiponectin secretion by real-time TIRF imaging

of adiponectin exocytosis?

Our previous work has begun to detail the white adipocyte adiponectin exocytosis process (5-

9, 22, 27, 33). However, the combination of methods used have temporal and spatial limitations

and detailed knowledge of adiponectin vesicle dynamics and release can only be gained by

high-resolution imaging in live cells. The finding that adiponectin exocytosis is similarly

controlled in 3T3-L1 and primary mouse and human adipocytes (5, 7-9, 32, 33) open

opportunities to use the stably expressing cells to achieve a comprehensive understanding of

mechanisms and mediators that control adiponectin vesicle exocytosis, at the cellular and

molecular level. TIRF movies can be acquired at high time resolution (10-20 FPS), thus

allowing real time direct visualisation of the adiponectin vesicle exocytosis process.

Consequently, the control and rate of docking and exocytosis can be determined.

In summary, TIRF imaging of cultured or primary live adipocytes with fluorescently labelled

adiponectin vesicles will enable us, and others, to define currently unknown regulatory steps

of adiponectin vesicle dynamic, maturation and exocytosis, at a level of detail that has not been

studied previously. How are adiponectin vesicles recruited to the exocytosis site? Are the

vesicles present before, during and after exocytosis? Do vesicles interact with each other during

the exocytosis process? Those are questions that can be answered with the type of research

presented here.

References

1.   Straub LG, Scherer PE. Metabolic Messengers: Adiponectin. Nat Metab. 2019;1(3):334-
9.
2.   Iwabu M, Okada-Iwabu M, Yamauchi T, Kadowaki T. Adiponectin/AdipoR Research and
Its Implications for Lifestyle-Related Diseases. Front Cardiovasc Med. 2019;6:116.
3.   Combs TP, Pajvani UB, Berg AH, Lin Y, Jelicks LA, Laplante M, et al. A transgenic mouse
with a deletion in the collagenous domain of adiponectin displays elevated circulating adiponectin and
improved insulin sensitivity. Endocrinology. 2004;145(1):367-83.
4.   Lim CY, Hong W, Han W. Adiponectin is released via a unique regulated exocytosis
pathway from a pre-formed vesicle pool on insulin stimulation. Biochem J. 2015;471(3):381-9.
5.   Brannmark C, Lovfors W, Komai AM, Axelsson T, El Hachmane MF, Musovic S, et al.
Mathematical modeling of white adipocyte exocytosis predicts adiponectin secretion and quantifies
the rates of vesicle exo- and endocytosis. J Biol Chem. 2017;292(49):20032-43.
6.   El Hachmane MF, Komai AM, Olofsson CS. Cooling Reduces cAMP-Stimulated
Exocytosis and Adiponectin Secretion at a Ca2+-Dependent Step in 3T3-L1 Adipocytes. PLoS One.
2015;10(3):e0119530.
7.   Komai AM, Brannmark C, Musovic S, Olofsson CS. PKA-independent cAMP stimulation
of white adipocyte exocytosis and adipokine secretion: modulations by Ca2+ and ATP. J Physiol.
2014;592(Pt 23):5169-86.
8.   Komai AM, Musovic S, Peris E, Alrifaiy A, El Hachmane MF, Johansson M, et al. White
Adipocyte Adiponectin Exocytosis Is Stimulated via beta3-Adrenergic Signaling and Activation of
Epac1: Catecholamine Resistance in Obesity and Type 2 Diabetes. Diabetes. 2016;65(11):3301-13.
9.   Musovic S, Komai AM, Said MK, Shrestha MM, Wu Y, Wernstedt Asterholm I, et al.
Noradrenaline and ATP regulate adiponectin exocytosis in white adipocytes: Disturbed adrenergic and
purinergic signalling in obese and insulin-resistant mice. Mol Cell Endocrinol. 2022;549:111619.
10.   Mattheyses AL, Simon SM, Rappoport JZ. Imaging with total internal reflection
fluorescence microscopy for the cell biologist. J Cell Sci. 2010;123(Pt 21):3621-8.
11.   Burchfield JG, Lopez JA, Mele K, Vallotton P, Hughes WE. Exocytotic vesicle behaviour
assessed by total internal reflection fluorescence microscopy. Traffic. 2010;11(4):429-39.
12.   Moffat J, Sabatini DM. Building mammalian signalling pathways with RNAi screens.
Nature reviews Molecular cell biology. 2006;7(3):177-87.
13.   Li M, Rossi JJ. Lentiviral vector delivery of siRNA and shRNA encoding genes into
cultured and primary hematopoietic cells. Methods in molecular biology. 2008;433:287-99.
14.   Carlotti F, Bazuine M, Kekarainen T, Seppen J, Pognonec P, Maassen JA, et al. Lentiviral
vectors efficiently transduce quiescent mature 3T3-L1 adipocytes. Mol Ther. 2004;9(2):209-17.
15.   Burgoyne RD, Morgan A. Secretory granule exocytosis. Physiol Rev. 2003;83(2):581-
632.
16.   MacDonald PE, Rorsman P. The ins and outs of secretion from pancreatic beta-cells:
control of single-vesicle exo- and endocytosis. Physiology (Bethesda). 2007;22:113-21.
17.   Garcia AG, Garcia-De-Diego AM, Gandia L, Borges R, Garcia-Sancho J. Calcium signaling
and exocytosis in adrenal chromaffin cells. Physiol Rev. 2006;86(4):1093-131.
18.   Wang ZV, Scherer PE. Adiponectin, the past two decades. J Mol Cell Biol. 2016;8(2):93-
100.
19.   Cong L, Chen K, Li J, Gao P, Li Q, Mi S, et al. Regulation of adiponectin and leptin
secretion and expression by insulin through a PI3K-PDE3B dependent mechanism in rat primary
adipocytes. Biochem J. 2007;403(3):519-25.

20.   Bogan JS, Lodish HF. Two compartments for insulin-stimulated exocytosis in 3T3-L1
adipocytes defined by endogenous ACRP30 and GLUT4. J Cell Biol. 1999;146(3):609-20.
21.   Xie L, O'Reilly CP, Chapes SK, Mora S. Adiponectin and leptin are secreted through
distinct trafficking pathways in adipocytes. Biochim Biophys Acta. 2008;1782(2):99-108.
22.   Lovfors W, Simonsson C, Komai AM, Nyman E, Olofsson CS, Cedersund G. A systems
biology analysis of adrenergically stimulated adiponectin exocytosis in white adipocytes. J Biol Chem.
2021;297(5):101221.
23.   Zhang X, Heckmann BL, Liu J. Studying lipolysis in adipocytes by combining siRNA
knockdown and adenovirus-mediated overexpression approaches. Methods Cell Biol. 2013;116:83-
105.
24.   Nugent C, Prins JB, Whitehead JP, Savage D, Wentworth JM, Chatterjee VK, et al.
Potentiation of glucose uptake in 3T3-L1 adipocytes by PPAR gamma agonists is maintained in cells
expressing a PPAR gamma dominant-negative mutant: evidence for selectivity in the downstream
responses to PPAR gamma activation. Mol Endocrinol. 2001;15(10):1729-38.
25.   Gandasi NR, Vesto K, Helou M, Yin P, Saras J, Barg S. Survey of Red Fluorescence
Proteins as Markers for Secretory Granule Exocytosis. PLoS One. 2015;10(6):e0127801.
26.   Olofsson CS, Gopel SO, Barg S, Galvanovskis J, Ma X, Salehi A, et al. Fast insulin secretion
reflects exocytosis of docked granules in mouse pancreatic B-cells. Pflugers Arch. 2002;444(1-2):43-
51.
27.   Komai AM, Shrestha MM, Musovic S, Olofsson CS. An Approach to Monitor Exocytosis
in White Adipocytes. Methods in molecular biology. 2021;2233:203-22.
28.   Park HK, Ahima RS. Resistin in rodents and humans. Diabetes Metab J. 2013;37(6):404-
14.
29.   Musovic S, Shrestha MM, Komai AM, Olofsson CS. Resistin is co-secreted with
adiponectin in white mouse adipocytes. Biochem Biophys Res Commun. 2021;534:707-13.
30.   Rogne M, Tasken K. Compartmentalization of cAMP signaling in adipogenesis,
lipogenesis, and lipolysis. Hormone and metabolic research = Hormon- und Stoffwechselforschung =
Hormones et metabolisme. 2014;46(12):833-40.
31.   Calaghan S, Kozera L, White E. Compartmentalisation of cAMP-dependent signalling by
caveolae in the adult cardiac myocyte. J Mol Cell Cardiol. 2008;45(1):88-92.
32.   Brannmark C, Kay EI, Ortegren Kugelberg U, Chanclon B, Shrestha MM, Wernstedt
Asterholm I, et al. Adiponectin is secreted via caveolin 1-dependent mechanisms in white adipocytes.
J Endocrinol. 2020;247(1):25-38.
33.   Musovic S, Olofsson CS. Adrenergic stimulation of adiponectin secretion in visceral
mouse adipocytes is blunted in high-fat diet induced obesity. Sci Rep. 2019;9(1):10680.

Author contributions

MMS and CSO outlined the study and designed the experiments. MMS carried out all

experimental work and analysis. CSO and MMS wrote the manuscript. SB provided method

and analysis expertise and gave valuable input on the manuscript. All authors approved the

final version of the manuscript.

Funding

This study was funded by the Swedish Research Council (2013-07107 and 2019-01239).

Legends

Fig. 1 Effects of an elevation of cAMP and Ca^{2+} on adiponectin exocytosis in white adipocytes.

Forskolin and 3-isobutyl-1-methylxanthine (IBMX) elevate the intracellular concentration of cAMP through the activation of adenylyl cyclase (AC) and inhibition of phosphodiesterase, respectively. The elevation of cAMP results in activation of the downstream target Epac1 and fusion of adiponectin containing vesicles with the plasma membrane with subsequent adiponectin release. Ionomycin, a calcium ionophore, facilitates the transport of Ca^{2+} across the plasma membrane and the elevation in intracellular Ca^{2+} augments the fusion of adiponectin-containing vesicles with the plasma membrane. Ca^{2+} also induces replenishment of release-competent adiponectin vesicles, to maintain secretion of adiponectin over longer time periods.

Fig. 2 Expression of adiponectin and shRNA-mediated adiponectin knockdown in 3T3-L1 adipocytes.

A) Gene expression of endogenous adiponectin in 3T3-L1 cells/adipocytes measured at every 48 hours after initiation of differentiation until day 10. Western blot **(B)** and densitometric analysis **(C)** of adiponectin protein expression after shRNA-mediated knockdown of endogenous adiponectin in 3T3-L1 cells. The cells were infected with lenti-virus-expressing control shRNA targeting GFP (scrambled) and two variants of shRNA targeting murine adiponectin (shAcrp30-A and shAcrp30-C), and differentiated into mature adipocytes after puromycin selection. **D)** mRNA levels of adiponectin in shRNA-mediated knockdown compared with control cells. Statistical analysis: Brown-Forsythe and Welch ANOVA tests (Dunnett's T3 multiple comparisons test), assuming unequal SDs at α – level of 0.05 for A and Ordinary One-way ANOVA tests (Dunnett's multiple comparisons test), assuming equal SDs at α – level of 0.05 for C & D. Data are presented as mean ± SEM of three independent experiments. *P<0.05, ***P<0.001 and ****P<0.0001.

Fig. 3 Rescue of shRNA-mediated adiponectin knockdown with mCherry fused with human adiponectin and estimation of vesicle size

Western blot for **A)** reduced and **B)** non-reduced adiponectin. The expression of mCherry-fused adiponectin in the 3T3-L1 adipocytes was observed under **C)** Fluorescence (Scale bar=100 μm) and **D)** TIRF microscope (Scale bar=10 μm). The 3T3-L1 cells, with shRNA-mediated silencing of endogenous adiponectin expression, were infected with lenti-virus for stably expressing human adiponectin fused with mCherry at C-terminal alone or with FLAG-tag at either C- or N- terminal. The lenti-virus infected cells were differentiated and the over-expression of recombinant adiponectin was analysed. **E-F)** Estimation of adiponectin-mCherry vesicle size (n=267), as described in Methods. Fluorescent polystyrene beads with a diameters of 2, 1, 0.5, 0.2 and 0.1 μm were imaged. Scale bars=10 μm.

Fig. 4 TIRF imaging of cAMP-stimulated 3T3-L1 adipocyte adiponectin exocytosis.

3T3-L1 adipocytes stably expressing recombinant adiponectin were serum starved for 30 min at 32^{0}C and treated with **A)** 10 μM Forskolin and 200 μM IBMX or **B)** DMSO (Control) for 15 minutes during which images were taken at 2 FPS using TIRF microscopy. A laser wavelength of 561 nm with exposure time between 30 - 100 ms was used to excite mCherry.

Note in inset region at lower panel that some vesicles are released (dotted circles) while others remain stable (whole circles). Scale bar = 10 μm. **C)** Examples of FSK/IBMX-induced single vesicle dynamics over time, in the stimulated adipocyte in (A) and in the control adipocyte in (B). Scale bar = 2 μm. **D-E)** Relative average fluorescence intensity change vs. time in a single vesicle stimulated with FSK/IBMX. **F)** Cumulative frequency of percentage of vesicles released at the end of a 10 min FSK/IBMX stimulation. **G)** Vesicle count before and 15 min after stimulation with FSK/IBMX. Statistical analysis: 2way ANOVA (Sidak's multiple comparisons test). Data are presented as mean ± SEM of eight independent experiments. *P<0.05; **P<0.01.

Fig. 5 TIRF imaging of Ca^{2+}-stimulated 3T3-L1 adipocyte adiponectin exocytosis.
3T3-L1 adipocytes stably expressing recombinant adiponectin were serum starved for 30 min at 32^0C and treated with **A)** 1 μM ionomycin or **B)** DMSO (Control) for 15 minutes during which images were taken at 2 FPS using TIRF microscopy. A laser wavelength of 561 nm with exposure time between 30 - 100 ms was used to excite mCherry. Note in inset region at lower panel that some vesicles are released (dotted circles) while others remain stable (whole circles). Scale bar = 10 μm. **C)** Examples of ionomycin-induced single vesicle dynamics over time, in the stimulated adipocyte in (A) and in the control cell in (B). Scale bar = 2 μm. **D-E)** Relative average fluorescence intensity change vs. time in a single vesicle upon ionomycin stimulation. **F)** Cumulative frequency of percentage of vesicles released by ionomycin at the end of 10 min of stimulation. **G)** Vesicles count before and 15 min after stimulation with ionomycin. Statistical analysis: 2way ANOVA (Sidak's multiple comparisons test). Data are presented as mean ± SEM of eight independent experiments. *P<0.05; **P<0.01.

Fig.6 cAMP- and Ca^{2+}-stimulated adiponectin secretion in native and adiponectin-mCherry-expressing 3T3-L1 adipocytes.
A) Native mature 3T3-L1 adipocytes were serum starved for 30 min at 32^0C and treated with DMSO, FSK/IBMX, or ionomycin for 30 min at 32^0C. The extracellular solutions were collected and analysed for endogenous total adiponectin release using ELISA. Statistical analysis: Brown-Forsythe and Welch ANOVA (Dunnett's T3 multiple comparisons test), assuming unequal SDs at α – level of 0.05. N=6 for Control and ionomycin; N= 5 for FSK/IBMX. **B)** Human total adiponectin and **C)** human HMW adiponectin secretion in stably expressing adipocytes, measurement using ELISA. Statistical analysis: Ordinary One-way ANOVA (Holm-Sidak's multiple comparisons test), assuming equal SDs at α – level of 0.05. N= 4. Data are presented as mean ± SEM. *P<0.05, **P<0.01 and ***P<0.001. Lipolysis measured as FFA **D)** and glycerol **E)** release in 3T3-L1 adipocytes stably expressing recombinant adiponectin under DMSO (Control), FSK/IBMX or ionomycin stimulation. Note that only co-stimulation with FSK/IBMX significantly increased the secretion of FFA and glycerol compared to the control. Statistical analysis: Ordinary One-way ANOVA (Dunnett's multiple comparisons test), assuming equal SDs at α – level of 0.05. N=5. Data are presented as mean ± SEM. ****P<0.0001 and ns = non-significant.

Table 1

Primer	Forward	Reverse
shAcrp30-A	CCGGCTACAACCAACAGAATCAT TACTCGAGTAATGATTCTGTTGGT TGTAGTTTTTG	AATTCAAAAACTACAACCAA CAGAATCATTACTCGAGTAAT GATTCTGTTGGTTGTAG
shAcrp30-B	CCGGGACAAGGCCGTTCTCTTCA CCCTCGAGGGTGAAGAGAACGGC CTTGTCTTTTTG	AATTCAAAAAGACAAGGCCG TTCTCTTCACCCTCGAGGGTG AAGAGAACGGCCTTGTC
shAcrp30-C	CCGGCCACAATGGCACACCAGGC CGCTCGAGCGGCCTGGTGTGCCA TTGTGGTTTTTG	AATTCAAAAACCACAATGGC ACACCAGGCCGCTCGAGCGG CCTGGTGTGCCATTGTGG
Acrp30	CCCAAGGGAACTTGTGCAGGTTG GATG	GTTGGTATCATGGTAGAGAAG AAAGCC
β-Actin	TTCTTTGCAGCTCCTTCG	TTCTGACCCATTCCCACC

Supplementary Videos

Supplementary video 1

3T3-L1 adipocytes stably expressing adiponectin-mCherry were treated with 10 μM Forskolin and 200 μM IBMX for 15 minutes during which images were taken at 2 FPS using TIRF microscopy. The video is condensed to show every 10th frame. Time point of FSK/IBMX addition = 30 s. Scale bar = 10 μm.

Supplementary video 2

3T3-L1 adipocytes stably expressing recombinant adiponectin were treated with DMSO (control for Forskolin/IBMX recording) for 15 minutes during which images were taken at 2 FPS using TIRF microscope. The video is condensed to show every 10th frame. Scale bar = 10 μm.

Supplementary video 3

3T3-L1 adipocytes stably expressing recombinant adiponectin were treated with 1 μM ionomycin for 15 minutes during which images were taken at 2 FPS using TIRF microscope. The video is condensed to show every 10th frame. Time point of ionomycin addition = 30 s. Scale bar = 10 μm.

Supplementary video 4

3T3-L1 adipocytes stably expressing recombinant adiponectin were treated with DMSO (control for ionomycin recording) for 15 minutes during which images were taken at 2 FPS using TIRF microscope. The video is condensed to show every 10th frame. Scale bar = 10 μm.

Abbreviation

AAV = Adeno-Associated Virus

ANOVA = Analysis of variance

ATCC = American Type Culture Collection

cAMP = Cyclic Adenosine Monophosphate

CMV = Cytomegalovirus

DMEM = Dulbecco's Modified Eagle Medium

DMSO = Dimethyl sulfoxide

EDTA = Ethylenediaminetetraacetic acid

EGFP = Enhanced Green Fluorescent Protein

ELISA = Enzyme-linked immunosorbent assay

Epac1 = Exchange factor directly activated by cAMP 1

Epac2 = Exchange factor directly activated by cAMP 2

EC = Extracellular

FBS = Fetal bovine serum

FPS = Frames per second

FSK = Forskolin

GFP = Green Fluorescent Protein

HMW = High Molecular Weight

IBMX = 3-isobutyl-1-methylxanthine

NPY = Neuropeptide Y

NS = Not significant

PAGE = Polyacrylamide gel electrophoresis

PVDF = Polyvinylidene fluoride

qPCR = quantitative polymerase chain reaction

ROI = Region of interest

SD = Standard deviation

SDS = Sodium dodecyl-sulfate

SEM = Standard error of mean

shRNA = short hairpin ribonucleic acid

TIRF = Total internal reflection fluorescence

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Plasmid map of lenti-viral construct for human adiponectin fused with mCherry at C-terminal created with SnapGene. 3T3-L1 adipocytes expressing recombinant adiponectin were serum starved for 30 minutes at 32 ^{0}C and treated with 10 µM Forskolin and 200 µM IBMX, 1 µM ionomycin and DMSO as Control for 15 minutes at 32 ^{0}C during which images were taken at 2 FPS using TIRF microscope.