Title: Involvement of an IgE/Mast cell/B cell amplification loop in abdominal aortic aneurysm progression

Alexia Loste[1,2], Marc Clément, PhD[1,2], Sandrine Delbosc[1,2], Kevin Guedj[1,2], Jean Sénémaud[1,2,3], Anh-Thu Gaston[1,2], Marion Morvan[1,2], Guillaume Even[1,2], Grégory Gautier[2,4], Alexander Eggel[5], Michel Arock[6], Emanuele Procopio[1,2], Catherine Deschildre[1,2], Liliane Louedec[1,2], Jean-Baptiste Michel[1,2], Lydia Deschamps[7], Yves Castier[3], Raphaël Coscas[1,8], Jean-Marc Alsac[9], Pierre Launay[2,4], Giuseppina Caligiuri[1,2,10], Antonino Nicoletti[1,2], Marie Le Borgne[1,2,*]

[1] Université Paris Cité and Université Sorbonne Paris Nord, INSERM, LVTS, F-75018 Paris, France.

[2] DHU FIRE, Paris, France.

[3] Department of Vascular and Thoracic Surgery, AP-HP, Bichat Hospital, Université Paris Cité, Paris, France.

[4] INSERM UMRS 1149, Centre de Recherche sur l'Inflammation (CRI), Université Paris Cité, Paris, France.

[5] Department for BioMedical Research, University of Bern, Bern, Switzerland.

[6] Department of Biology and CNRS UMR8113, Ecole Normale Supérieure de Paris-Saclay, Saclay, France.

[7] Department of Pathology, AP-HP, Bichat Hospital, Université Paris Cité, Paris, France.

[8] Department of Vascular Surgery, AP-HP, Ambroise Paré University Hospital, Université Paris Cité, Boulogne-Billancourt, France.

[9] Department of Vascular Surgery, AP-HP, Hôpital Européen Georges Pompidou, Université

Paris Cité, Paris, France.

[10] Department of Cardiology, AP-HP, Bichat Hospital, Université Paris Cité, Paris, France.

Address for correspondence:

[*] Corresponding author

e-mail: marie.le-borgne-moynier@inserm.fr

Abstract

Aims: IgE type immunoglobulins and their specific effector cells, mast cells (MCs), are associated with abdominal aortic aneurysm (AAA) progression. In parallel, immunoglobulin-producing B cells, organised in tertiary lymphoid organs (TLOs) within the aortic wall, have also been linked to aneurysmal progression. We aimed at investigating the potential role and mechanism linking local MCs, TLO B cells, and IgE production in aneurysmal progression.

Methods and Results: Through histological assays conducted on human surgical samples from AAA patients, we uncovered that activated MCs were enriched at sites of unhealed haematomas, due to subclinical aortic wall fissuring, in close proximity to adventitial IgE+ TLO B cells. Remarkably, *in vitro* the IgEs deriving from these samples enhanced MC production of IL-4, a cytokine which favors IgE class-switching and production by B cells. Finally, the role of MCs in aneurysmal progression was further analysed *in vivo* in ApoE$^{-/-}$ mice subjected to angiotensin II infusion aneurysm model, through MC-specific depletion after the establishment of dissecting aneurysms. MC-specific depletion improved intramural haematoma healing and reduced aneurysmal progression.

Conclusions: Our data suggest that MC located close to aortic wall fissures are activated by adventitial TLO B cell-produced IgEs and participate to their own activation by providing support for further IgE synthesis through IL-4 production. By preventing prompt repair of aortic subclinical fissures, such a runaway MC activation loop could precipitate aneurysmal progression, suggesting that MC-targeting treatments may represent an interesting adjunctive therapy for reducing AAA progression.

Keywords: Abdominal aortic aneurysm, mast cells, IgE antibodies, tertiary lymphoid organs, B lymphocytes, aortic dissection.

1 Abbreviations

2 AAA = abdominal aortic aneurysm; AngII = Angiotensin II; ApoE = apolipoprotein E; DT =

3 diphtheria toxin; MC = mast cell; NAA = non-aneurysmal aortas; RMB = red mast cells and

4 basophil; TLO = tertiary lymphoid organ.

1   Introduction

Arteries are subjected to recurrent mechanical insults which rise from the luminal side of

vessels (1). Vessels have intrinsic capacities to ensure prompt healing of local injuries and are

assisted in this task by resident and recruited inflammatory cells (2). However, the persistence

of vascular inflammation can eventually amplify the arterial damage and lead to severe and life-

threatening conditions including coronary artery disease, strokes or abdominal aortic

aneurysms (AAAs) (3). Promoting arterial healing by targeting inflammation thus constitutes a

major challenge in modern medicine.

In this study, we focused on AAA progression. Monitoring AAA enlargement is problematic

due to the discontinuous, so-called 'staccato' growth where month-lasting no-growth/healing

periods can be succeeded by a sudden enlargement and again a no-growth/healing period (4).

So far, no pharmacological treatments have been identified to reduce or stop AAA expansion

and subsequent arterial rupture, which causes up to 200,000 deaths worldwide each year (5).

The only therapeutic option when the aneurysm diameter exceeds a certain value (55mm in

men, 50mm in women) is aorta surgery. Therefore, deciphering the molecular pathways

involved in AAA healing and/or progression is essential to set up pharmacological alternatives.

Chronic immune responses involve adaptive and innate immunity and their relationship. In

particular, we and others have observed that chronic immune stimulation in AAAs leads to local

tertiary lymphoid organ (TLO) development within the adventitia (6-9). TLOs contain germinal

centre B cells corresponding to B cells undergoing differentiation into plasma cells (6). We

have reported that TLO development is associated with increased levels of antibodies in the

adventitia of human AAAs, in particular IgEs (6), which are also involved in chronic

inflammation. Notably, increase in IgE blood concentration correlates with the progression of

AAAs and other arterial diseases in patients (10, 11) and *in vivo* studies suggest a pathogenic

role for IgEs in experimental AAA (12).

In parallel, triggered by the binding of IgE at their surface, activated mast cells (MCs) release

potent proteases, cytokines and vasoactive molecules, such as leukotriene and histamine,

directly or indirectly favouring the detersion of the extracellular matrix and a massive loss of

the contingent of medial smooth muscle cells, that could eventually lead to aneurysmal

expansion (13-15). Interestingly, MC number increases in diseased arteries (16, 17).

Furthermore, MCs can promote B cell effector functions (18). As a consequence, MCs have

been surmised to be detrimental in AAAs (13, 14, 19). Studies aiming at inhibiting MC

activation with pemirolast, a histamine H1 antagonist, were inconclusive regarding their effect

on AAA reduction (20). However, pemirolast cannot uncouple IgE-mediated effector

processes, which are by far the most potent and specific trigger of MC activation. Hence, to

date, the role of MCs in aneurysmal progression remains elusive.

Here, we aimed at investigating the potential mechanism linking MC activity to IgEs and TLO

B cell in aneurysmal progression. To do so, we performed histological and *in vitro* assays using

human surgical samples from AAA patients. Furthermore, the putative pathogenic role of MCs

in aneurysmal growth was evaluated *in vivo*, by inducing MC depletion in Red Mast cell and

Basophil (RMB) - ApoE[-/-] mice subjected to angiotensin II infusion during the phase of

aneurysm progression.

1   Materials and Methods

2   Mice

Apolipoprotein E-deficient (ApoE$^{-/-}$) mice (RRID:IMSR_JAX:002052) were crossed to RMB

(B6.Ms4a2^{tm1Mal}) mice (21). RMB mice carry an additive transgene containing the promoter of

the high affinity IgE receptor β subunit controlling the human diphtheria toxin (DT) receptor

and the tomato red fluorescent protein (21). The heterozygous offspring were then intercrossed

to generate homozygous ApoE$^{-/-}$ RMB mice. Mice were maintained on a C57Bl/6J background

and fed a regular chow diet. Induction of AAAs, histology and flow cytometry were performed

as described below. All investigations on mice conformed to the Directive 2010/63/EU of the

European Parliament, and review and approval of the study was obtained from the Comité

d'Ethique Paris Nord #121 (APAFIS #12027).

12   Human samples

AAA tissues and blood were obtained from patients undergoing surgery and enrolled in the

RESAA protocol (REflet Sanguin de l'évolutivité des Anévrysmes de l'aorte abdominale) (22).

As expected (5), AAA patients were predominantly men, and presented risk factors such as age,

hypertension, hyperlipidaemia and smoking (Table S1). All patients gave written informed

consent, and the protocol was approved by the Comité Consultatif de Protection des Personnes

dans la Recherche Biomédicale (CCPPRB, Paris-Cochin, approval no. 2095). Our study

complies with the Declaration of Helsinki. Control aortas were sampled from dead organ donors

with the authorisation of the French Biomedicine Agency (PFS 09-007, Table S2). Depending

on their sizes, samples were cut in several pieces that were used to prepare conditioned medium,

and/or digested for flow cytometry analysis, and/or fixed in 3.7% paraformaldehyde (PFA) for

histological studies, and stored in the Inserm human CV biobank (BB-0033-00029), included

in the European network BBMRI-ERIC.

Histology and immunofluorescence on human samples

PFA-fixed aortic tissues were paraffin-embedded. Four µm-thick sections were deparaffinised

in toluene and rehydrated in ethanol. Sections were subjected to Carstair's stain, orcein stain or

Perl's + diaminobenzidine (DAB) stain. Alternatively, sections were incubated with retrieval

reagent (R&D Systems), then immunostained using antibodies against IgE (goat polyclonal,

Vector Laboratories), MC tryptase (rabbit monoclonal, clone EPR8476, abcam; or mouse

monoclonal, clone AA1, abcam), CD20 (mouse monoclonal, clone L26, Dako; or goat

polyclonal, Thermofisher), CD117 (rabbit polyclonal, Dako), or glycophorin A (rabbit

monoclonal, clone EPR8200, abcam). Immunostaining with isotypes were conducted to verify

the specificity of the primary antibodies. Overnight incubation at 4°C with primary antibodies

were followed by incubation with fluorophore-coupled anti-species antibodies (Jackson

Immunoresearch) for one hour at room temperature. References for antibodies are shown in

Supplementary File 1. Nuclei were then stained with Hoechst 53542, and slides were mounted

with fluorescent mounting medium (ProLong Gold, ThermoFisher). Slides were kept in the

dark at 4°C. Blind analysis was performed for the presence of IgE+ TLOs, MCs and

haematomas.

Preparation of conditioned medium and immunodetection of soluble molecules

For the preparation of conditioned medium, the adventitia of AAA and control aortas were

separated from the media and cut into small pieces (5 mm^3). The samples were then incubated

for 24 hours at 37°C in a standardised volume (6 mL/g of tissue) of RPMI 1640 medium (Gibco)

supplemented with antibiotics and antimycotics. The conditioned medium was then centrifuged

and the supernatant aliquoted and frozen at -80°C until use.

The IL-4 concentration in conditioned medium from adventitial tissues and serum was analysed

by CBA (BD biosciences). Bead fluorescence was recorded on a LSRII flow cytometer (BD).

Tryptase was quantified by ELISA (USCN). Immunoglobulins were analysed using a bio-plex

pro human isotyping assay (Bio-Rad) according to manufacturer instructions. Beads were

analysed on a Bioplex-200 analyser (Bio-Rad). IL-4 and Tryptase were analysed on a different

set of patients.

ROSA mast cell line

The ROSA$^{KIT\ WT}$ human cell line (RRID:CVCL_5G49) was cultured in the presence of

recombinant human SCF (80 ng/mL, R&D Systems), as described previously (23). Cells were

stimulated in the presence of conditioned medium from AAA and control adventitia, or PMA

(10ng/mL, Sigma-Aldrich) and ionomycin (1μM, Sigma-Aldrich) (hereafter P+I condition) or

IgE (2μg/mL, BeckmanCoulter) for 15 minutes followed by anti-IgE antibody (5μg/mL). After

1 hour of stimulation, the expression of CD63 in MCs was analysed by flow cytometry.

Alternatively, after 4 hours of stimulation, cells were collected and dry pellets were frozen for

gene expression analysis. In some experiments, 5μM DARPin® protein bi53_79 (24) was added

to the conditioned medium. Experiment was repeated twice with similar results.

Gene expression analysis

Total RNAs were extracted from ROSA MCs using the PureLink RNA mini kit (Invitrogen),

and mRNA reverse transcription was performed using iScript reverse transcriptase (Bio-Rad).

Real time quantitative PCR (RT-qPCR) was performed on a CFX 100 thermocycler (Bio-Rad)

using the following primers: forward IL-4: GCAGTTCTACAGCCACCAT; reverse IL-4:

ACTCTGGTTGGCTTCCTTCA . One nanogram of cDNA from each sample was mixed with

forward and reverse primers (300 nM) and SYBR Green master mix (Bio-Rad). The

amplification program was as follows: A first step of initial denaturation at 95°C for 3 minutes,

then 40 cycles of 3 steps: denaturation at 95°C 15 seconds, annealing at 57°C for 15 seconds,

and a final extension at 72°C for 30 seconds. The data were analysed using the $2^{-\Delta\Delta Ct}$ formulas:

the Ct values of IL-4 were normalised to the average Ct values of RPS18 (forward RPS18:

GCGGCGGAAAATAGCCTTTG; reverse RPS18: GATCACACGTTCCACCTCATC) and

ACTB   (forward   ACTB: TCCCTGGAGAAGAGCTACG;   reverse   ACTB:

TTTCGTGGATGCCACAGGAC) and non-treated MCs were used as reference.

Ang-II abdominal aortic pseudoaneurysms in mice

28-week-old ApoE-RMB males were used for the experiments, as males are more susceptible

to develop AAAs after Angiotensin II (AngII)   infusion   (25).   AngII   (#A9525,   1

mg/kg/day, Sigma-Aldrich, St Louis, Missouri) was continuously infused into the experimental

mice via osmotic mini-pumps (Model 2004, Alzet, Charles River Laboratories) that were

surgically implanted subcutaneously in the interscapular region under anaesthesia induced by

intraperitoneal injection of ketamine (100 mg/kg) and xylazine (20 mg/kg). Surgery was

followed by a buprenorphine intraperitoneal injection (0.1 mg/kg) for analgesia. Three mice

(8%) died within the first 10 days. Fourteen days after inducing AAAs through AngII infusion

(26), mice received two intraperitoneal injection of DT (1µg/ injection, n=18) at a one-day

interval (or PBS in the control group, n=17) in order to deplete MCs. Randomization was

achieved by having mice from each experimental group in each cage. Mice were sacrificed at

day 28 by intracardiac exsanguination under overdose of anaesthesia (intraperitoneal injection

of 150 mg/kg ketamine and 30 mg/kg xylazine). Before the exsanguination, a cell suspension

from the peritoneum was obtained by peritoneal lavage with 5 mL of ice-cold PBS. Blood was

withdrawn from the right ventricle of the heart and collected in EDTA tubes for blood cell

analysis. The heart and aorta were dissected, photographed and mounted on cryomolds for

further histomorphological analysis based on cryosections. The experiment was repeated twice

with similar results. Pooled number of mice from both experiments who survived and

developed aneurysms in each group are summarized in Table S3. DT injection had no effect on

blood basophils, peritoneal MCs, and aortic tissue MCs in control ApoE$^{-/-}$ mice (data not

shown).

Histology on mouse tissues

MC depletion was analysed on cryosections of mouse aortic roots stained with Toluidine Blue.

Images were captured on a Zeiss Axio Observer Z7 inverted microscope, and MCs were

counted manually. A detailed blind analysis of pseudoaneurysms was performed for mouse

abdominal aortas displaying macroscopic evidence of the occurrence of a pseudeoaneurysm.

Cross-cryosections (10 μm) of the aortic segments taken at different levels of the

pseudoaneurysms (every 300 μm) were stained for collagen with picrosirius red. Images were

acquired under polarised light. The size of the different layers (media, haematoma, adventitial

fibrous cap) and the extent of collagen deposition were quantitatively assessed using Image J.

Image acquisition

14   Images were digitally captured using an AxioObserver epi-fluorescent microscope equipped

15   with a Colibri 7 LED generator (Zeiss) and an ApoTome system and running Zen Software

16   (Zeiss). Macroscopic images of the human samples were acquired using the NanoZoomer

17   Digital slide scanner (Hamamatsu Photonics).

Flow cytometry analysis

Mouse blood samples collected in EDTA were incubated in Ammonium-Chloride-Potassium

lysis buffer for 5 minutes at room temperature to lyse red blood cells. Cells from blood samples

or 1 mL of peritoneal lavage were stained for dead cells with Live/Dead fixable far red Dead

cell stain kit (Invitrogen) for 30 minutes at 4°C, then incubated with a purified rat anti-mouse

CD16/32 (FcBlock, BD Biosciences) for 15 minutes at 4°C. Cells were then incubated for 20

minutes at 4°C with the following antibodies (from BD Biosciences unless stated otherwise):

BV605 rat anti-mouse CD19 (clone 1D3), FITC rat anti-mouse CD117 (clone 2B8), PerCP rat

anti-mouse CD45 (clone 30-F11), APC hamster anti-mouse FcεRIα (clone MAR-1,

Biolegend), AF700 rat anti-mouse CD3 (clone 172A). References for antibodies are shown in

Supplementary File 1.

For human tissue analysis, fresh adventitial layer samples were weighed, cut into small pieces

(<1 mm) and digested using a previously described protocol (27). After a wash step, the cells

were stained for dead cells with Live/Dead fixable far red Dead cell stain kit (Invitrogen) for

30 minutes at 4°C, then incubated for 20 minutes at 4°C with a combination of the following

mouse anti-human antibodies: AF488 anti-CD3 (clone UCHT1, BD Biosciences), PE anti-

CD203c (clone NP4D6, Biolegend), PerCP anti-FcεRIα (clone AER-37, Biolegend), BV421

anti-CD117 (clone 104D2, BD Biosciences), BV500 anti-CD45 (clone HI30, BD Biosciences),

PE-Cy7 anti-CD63 (clone H5C6, BD Biosciences), AF700 anti-CD19 (clone HIB19, BD

Biosciences), AF647 anti-CD107a (clone H4A3, BD Biosciences), BV421 anti-CD27 (clone

O323, Biolegend), BV570 anti-HLA-DR (clone L243, Biolegend), BV785 anti-CD19 (clone

HIB19, Biolegend), APC anti-IgD (clone IA6-2, BD Biosciences), FITC anti-CD38 (clone

HIT2, BD Biosciences), PE anti-CD45 (clone HI30, BD Biosciences), BV605 anti-CD45

(clone HI30, BD Biosciences).

For mouse peritoneal cells and human adventitial cells, flow-count fluorospheres (Beckman

Coulter) were added to the samples before the acquisition on the cytometer in order to calculate

absolute count of cells. Data were acquired on a LSRII flow cytometer or an ARIA III cell

sorter (BD Biosciences) and analysed using FACSDiva (BD Biosciences) and FlowJo

(TreeStar) softwares.

Statistical analysis

Statistical analysis was performed using the JMP9 and GraphPad Prism 9.0 software. To

compare two populations, we used Student t-tests or non-parametric Mann-Whitney tests when

sample size was too small (n<10) or when values were not normally distributed. Paired values

were compared using Wilcoxon matched-pairs signed rank test. Pearson correlation coefficients

or p-values of chi-square contingency tests were calculated to assess correlation between

continuous or categorical variables, respectively. A mixed model that uses a compound

symmetry covariance matrix was fitted in GraphPad Prism using Restricted Maximum

Likelihood (REML) to analyse effects of two categorical variables with repeated measures on

quantitative variables. Quantitative data are expressed as mean +/- standard error.

Results

IgE producing-TLOs are associated with aneurysmal wall unhealed fissures in patients

Tissue infiltrated IgE were enriched in aneurysmal samples and soluble IgE plasma levels were elevated in AAA patients compared to control non-aneurysmal aortas (NAAs, Fig S1). In parallel, TLOs were found in a majority of AAA samples (20 out of 25, Fig 1A) whereas they were absent in all NAA, in agreement with previous findings (6-9). In particular, adventitial TLOs were consistently found in samples from patients with larger aneurysms (Fig 1B), suggesting their higher frequency in these patients. In these adventitial TLOs, the proportion of germinal centre B cells was dramatically increased (Fig S2), suggesting that adventitial TLO B cells in large aneurysms are engaged into an Ab-producing program from which diverse Ig isotypes could emerge. Interestingly, we observed a strong staining for IgEs within the TLO "light zone" (where the centrocytes, i.e. B cells differentiating in immunoglobulins-producing cells, are located (28); Fig 1A) in 45% (n=9) of the samples. Altogether these results suggest that germinal centre B cells within adventitial TLOs situated within large AAAs may actively produce IgE immunoglobulins.

Intriguingly, combined histological analysis of Carstair's, Perl's + DAB and Orcein staining revealed the presence of recent or past wall fissuring with haematoma formation in the media in 92% AAA samples (Fig 1C, Fig S3, and Table 1). Indeed, 84% of total AAA tissues (n=21) presented an accumulation of red blood cells (Fig 1C), reflecting the recent occurrence of intramural haematomas. Of note, the diameter of the aneurysmal samples was significantly larger than 3 cm, implying a greater tension and mechanical stress, than the one of the non-aneurysmal aortas (diameter < 3 cm) (5) and an attentive evaluation of pre-surgery tomography angiograms consistently revealed the presence of at least one detectable aneurysmal wall

macroscopic fissure, with radiologic contrast penetrating *via* blood disruptions from the aortic

lumen to the aortic wall through the intraluminal thrombus, in 78% of patients with intramural

haematomas (Fig 1D and Fig S3D). In some samples, we were able to observe the entry site of

blood from the lumen on AAA histological sections, suggesting that the fissures leading to

intramural haematomas were provoked by tears (micro-fissures) initiated from the aorta lumen

(Fig S4A-B and S4D-E). Unhealed intramural haematomas could also be evidenced as large

areas of ferrous iron deposits next to modest red blood cell accumulation in the media (Fig S3B)

in 8% (n=2) of samples. Importantly, IgE+ TLOs were frequently localized near these

intramural haematomas (Fig 2A-C), suggesting that the maturation of B cells towards IgE-

producing cells within adventitial TLOs and the occurrence of micro-fissures are linked.

MCs are enriched at sites of aneurysmal wall fissures and IgE-producing TLOs in

human AAAs

In parallel, we observed that the adventitia of AAA patients was significantly enriched in MCs

compared to the adventitia of NAA organ donors (Fig 2E-F and Fig S5), in agreement with

previous observations (17). Moreover, MCs were often located next to IgE+ TLOs (Fig 2B-C),

and extracellular tryptase+ granules reflecting recent MC activation could be observed around

these MCs by immunostaining on AAA sections (Fig 2G-H). Indeed, as compared to NAA

samples, adventitial MCs in AAA samples were degranulating, as documented by the

expression of the degranulation/MC activation marker CD107a analysed by flow cytometry

(Fig 2E-F). Strikingly, MCs enriched in AAA tissues were consistently found at sites of

haematomas on AAA sections (Fig 2D and Fig S4C and S4F) where their granule enzymes

could favour arterial wall rupture. Altogether, these results suggest that MCs and IgE+ TLOs

could concur and be linked together with the occurrence of arterial wall injuries and aneurysmal

progression.

Soluble molecules from human AAA adventitia induce MC degranulation and IgE-

dependent production of IL-4

Given that MCs displayed an activated phenotype only in the adventitia of AAA patients (Fig

2), we assessed the putative presence of MC activation triggers in the conditioned medium

prepared from the adventitia of AAA as compared to control NAA tissues. We observed that

conditioned medium from AAA adventitias induced more surface expression of the

degranulating marker CD63 and more IL-4 mRNA transcription than conditioned medium from

NAA adventitia in cultured ROSA human MCs (Fig 3A-B). This is important because the

cytokine IL-4 is required to carry out an IgE antibody class switch recombination and hence

allow the production of IgE by B cells (29). Therefore, these results point at an unforeseen

function of MCs in AAAs, whereby MCs could orientate the local adaptive immune response

generated in adventitial TLOs toward IgE production. In turn, the binding of adventitial IgE on

their high affinity receptor on MCs could be responsible for triggering MC activation. To

directly assess this hypothesis, we pre-incubated the conditioned medium with anti-IgE

DARPin® protein bi53_79 (5 μM) before adding it to MCs (Fig 3C-D). DARPin® protein

bi53_79 is a soluble molecule that specifically binds to the IgE-Fc portion thereby preventing

IgE binding to FcεRI (24). Whereas treatment of conditioned medium with bi53_79 did not

prevent MC degranulation (Fig 3C), it decreased IL-4 mRNA production compared to untreated

conditioned medium from AAA tissues (Fig 3D). IgEs contained in the conditioned medium

from AAA adventitias therefore emerge as instrumental in promoting MC IL-4 production.

To assess if IL-4 production, MCs, and IgEs are also linked *in vivo*, we measured IgE, tryptase,

and IL-4 quantities in the conditioned medium from AAA adventitias. We found a statistically

significant positive correlation between the concentration of IgEs and IL-4 (p<0.001), as well

as between the concentration of IL-4 and tryptase (p<0.05, Fig 3E-F). No such correlations

were observed in the plasma of paired patients (Fig S6). These observations support the

existence of a local amplification loop in the adventitia of AAAs involving IL-4 producing MCs

and IgE-producing B cells.

9   MCs aggravates aneurysmal progression in mice

Previous studies have identified MCs as directly involved in provoking arterial damage (13, 14,

19). Consistently, the presence of medial/adventitial MCs with an activated phenotype was

associated to micro-fissures in our human AAA samples, suggesting that MCs could play a role

in the aneurysmal remodelling of the aortic wall, upon the occurrence of tissue fissuring.

Therefore, we asked whether specific depletion of MCs could prevent the aneurysmal

remodelling in ApoE$^{-/-}$ mice subjected to chronic infusion of Angiotensin II (AngII), a well-

known mouse model of aortic dissection eventually followed by aneurysmal progression (26).

In this setting, pseudoaneurysm of the abdominal aorta starts within 10 days upon the formation

of large intramural haematomas due to the occurrence of multiple medial tears originating from

the lumen, at the origin of the main side branches (30). In order to conditionally induce specific

MC depletion by DT injection after the occurrence of the aortic fissuring, we crossed ApoE$^{-/-}$

mice with RMB mice (ApoE-RMB mice) (21). ApoE-RMB mice were subjected to infusion

with AngII for 28 days. We injected DT (or PBS as a control) 14 days after the beginning of

AngII infusion. Thus, MCs were depleted after the occurrence of the dissection in this model

(26), allowing to evaluate the effect of their depletion on the remodelling of the dissected aortas

and their potential effect on aneurysmal progression 14 days later (Fig 4A).

Survival was similar in the two groups (Table S3). As anticipated, the proportion of mice

presenting an intramural haematoma at day 28 was equivalent in DT-treated mice (44%, n=

8/18) and in the control group (47%, n=7/17). DT injection almost completely depleted MCs

from the aortic tissue (Fig 4B-C) and the peritoneal cavity (Fig S7). It should be noted that in

RMB mice, DT induces transient depletion of both MCs and basophils. While basophils fully

repopulate the bloodstream within 12 days, the repopulation of peripheral tissues such as skin

or peritoneum by MCs is much slower (21, 31). As a result, two weeks after DT administration

(Day 28 of AngII infusion), MCs were still completely absent from the peritoneal cavity,

whereas the basophil population had fully replenished the blood of ApoE-RMB mice (Fig S7).

DT injection reduced the expansion of the pseudoaneurysm (Fig 4D-F), comprising the size of

the intramural haematoma and of the perivascular adventitial cuff (Fig S8A-C). Furthermore,

on the cross-sections of the dissected aorta segments stained with picrosirius red we observed

that the aortic wall of the pseudoaneurysm of DT-treated mice presented an increased density

of collagen compared to control mice (Fig 4E and 4G), especially in the adventitial fibrous cuff

(Fig S8D). Our data therefore suggest that MC depletion improves the adventitia fibrotic

remodelling which is critical for providing an adequate stiffness and strength subsequently

constraining the expansion of the haematoma and the reduction of the aneurysmal progression

(32).

Discussion

This study reveals the existence of a potential amplification loop involving IgE$^+$ TLO B cells and MCs at site of tissue fissuring, which could constitute an inflection point in AAA evolution, precipitating aneurysmal progression (Graphical Abstract).

To our knowledge, this is the first time IgE$^+$ germinal centre B cells have been reported in TLOs. Indeed, while they appear early in primary immune responses, coinciding with the peak of IL-4 production, IgE$^+$ cells are unable to populate the long-lived B cell compartment, which explains why IgE$^+$ germinal centres are rarely seen in lymphoid organs (33). Our data show that the observed IgE$^+$ TLOs develop close to aortic wall fissures. Strikingly, the proximity of MCs and IgE$^+$ TLOs at site of tissue injury in the wall of progressive aneurysms supports a possible role for these highly reactive inflammatory cells in the progression of the disease and in the maintenance of the pathologic loop, through their production of IL-4 in response to locally produced IgEs, which in turn sustains the IgE$^+$ germinal centre cells in local TLOs. The correlation observed between IgE and IL-4 concentrations, as well as between IL-4 and tryptase concentrations, provides compelling evidence for the interrelated activities of MCs and B cells. It is worth noting that other cell types besides MCs may also produce IL-4 and respond to IgEs (12, 34). Interestingly, IL-4 may play opposing roles at different stages of the pathology. Experimental models suggest that IL-4 produced by eosinophils may protect against the development of pseudoaneurysms (34). However, our data suggest that IL-4 may accelerate the rupture of established aneurysms.

Furthermore, it should be emphasized that the ROSA human MC line requires cross-linking of purified IgEs by an anti-IgE antibody for activation (23). However, we did not induce such cross-linking with the conditioned medium. The activation of MCs under these experimental

conditions suggests that the IgEs present in the conditioned medium were likely cross-linked

by their cognate antigens. In contrast, our data indicate that blood IgEs (not shown) failed to

activate MCs, suggesting that they were not circulating as immune complexes. This strongly

suggests that the antigen recognized by IgEs may be specifically present in the aneurysmal

aorta. Additionally, locally produced IgEs may contribute to AAA progression by promoting

the senescence of smooth muscle cells (35).

Whereas MC IL-4 production was dependent on IgEs present in the conditioned medium from

aneurysmal aorta, degranulation was not. This indicates that besides IgEs, other molecules from

the adventitia stimulate MCs. Indeed, other classes of immunoglobulins, which are also

enriched in aneurysmal aortas (6), cytokines, DAMP-containing molecules and inflammatory

mediators released by the injured vascular stroma could also trigger MC degranulation. Further

studies focusing on the relationship between IgEs and MC activation could lead to novel

therapeutic strategies to reduce chronic inflammatory disorders.

Our pre-clinical *in vivo* data directly support a role for MCs in the fate of aneurysmal

development of arterial wall subjected to fissuring and formation of an intramural haematoma.

Importantly, for reaching this conclusion, we used the ApoE-RMB mouse model in which the

depletion of MCs, conditioned by the use of DT, started only after the initial trigger (aortic

dissection). Instead, the rodent models previously used to address this question were

constitutively defective in MCs (13, 15, 17), implying that the deficiency of MCs preceded, and

possibly biased, the experimental intervention itself. Furthermore, constitutive MC-deficient

models such as $Kit^{W/Wv}$ and $Kit^{W-sh/W-sh}$ mice display other hematopoietic abnormalities, notably

neutrophilia, thrombocytosis, and macrophage defects (21, 36) which could be responsible for

some of the effects attributed to MCs. In addition, as resident MCs are present in healthy aortas,

constitutive MC depletion could impact aorta homeostasis as well as the

formation/progression/rupture of AAAs in these models. By triggering MC depletion after the

occurrence of dissections in our ApoE$^{-/-}$ RMB mouse model, we were able to specifically tackle

the role played by MCs during the healing processes associated with the progression of AAAs

in this model. We found that AAAs in MC-depleted animals displayed improved arterial

remodelling with an increased collagen content of the aneurysmal wall, supporting an active

role for MCs in collagen degradation and expansion of AAA (37, 38). Mechanistically, MC

chymase and tryptase could increase collagen degradation by converting collagenases from

their inactive to their active form (15). MCs could also inhibit the production of collagen by

smooth muscle cells (39). Hence, MCs could actively contribute to weaken the arterial wall at

sites of micro-fissures and favour the iterative expansion of aneurysms, under the biomechanic

stress of intraluminal thrombus (40, 41). Furthermore, the unhealed micro-fissures could allow

the accumulation of plasma proteins into the arterial wall, which is associated to AAA growth

(42).

Importantly, the experimental setting we used here is similar to the clinical settings of a recently

undertaken clinical trial investigating whether pemirolast, a histamine H1 antagonist acting as

a MC stabilizer in allergic conditions, could retard the growth of medium-sized AAAs (20).

This trial in which the major end-point was the change in aortic diameter as assessed by ultra-

sound imaging, failed to report a significant improvement in the patients receiving the MC

inhibitor. Although this study was elegantly designed and performed, it did not however bring

a definitive conclusion regarding MC involvement in AAA pathophysiology because i/ the

pemirolast treatment failed to decrease the plasma tryptase levels, indicating a critical dosing

issue, ii/ the study interval might have been too short, and iii/ contrary to our study, the effect

that MC inhibition might have exerted on tissue composition could not be studied in this clinical

study. Hence, in light of our results, we believe it would be interesting to re-evaluate the effect

of perimolast (and/or other MC-inhibiting drugs) on AAA growth and rupture rate by targeting

patients presenting elevated levels of IgEs and focusing on adventitial remodelling. Inhibiting

other key steps of the amplification loop, for instance by using anti-IgE antibodies, could be

another promising therapeutic strategy for such patients.

The present study has several limitations that should be acknowledged. First, our conclusion on

the pathological role of a MC and B cell amplification loop in AAA is based on observations

of human samples and experimental models. Intervention studies are needed to formally

demonstrate the involvement of such a loop in patients. Second, in our experimental mouse

model, basophils are also depleted, even if transiently, which means that an involvement of

basophils in the fibrotic remodelling in the adventitia and the progression of aneurysms cannot

be ruled out. Finally, the pseudoaneurysm model of AngII infusion in ApoE-/- mice lacks TLO

in the adventitia of the aorta, which limits its ability to test the impact of a local amplification

loop involving TLO B cells, even though it may still be useful in investigating the involvement

of MCs in preventing aneurysm worsening.

In spite of the aforementioned limitations, our findings strongly suggest that a local

amplification loop involving IgE, MCs, and TLO B cells plays a critical role in the development

and progression of AAA. This self-sustaining loop, driven by repetitive MC degranulation,

could constitute a turning point in the course of the disease, leading to impaired arterial healing

and ultimately AAA rupture. Our study provides a valuable framework for further

investigations into the pathological mechanisms underlying AAA and may inform the

development of novel therapeutic strategies targeting this debilitating condition (resumed in

Graphical Abstract).

1 Acknowledgments

2 The authors thank Melanie Gettings for her help with manuscript editing, and Emilie Monnet

3 for her help with the animals.

5 Materials availability statement

6 Use of DARPin® protein bi53_79 and RMB mice requires a MTA. The data underlying this

7 article will be shared on reasonable request to the corresponding author.

1 References

1.   Caligiuri G, Levy BP, Nicoletti A, Michel J-B. Role of Biomechanical Stress in the

Pathology of the Aorta. In: Stanger OH, Pepper JR, Svensson LG, editors. Surgical

Management of Aortic Pathology: Current Fundamentals for the Clinical Management of

Aortic Disease. Vienna: Springer Vienna; 2019. p. 163-80.

2.   Dutertre CA, Clement M, Morvan M, Schakel K, Castier Y, Alsac JM, et al. Deciphering

the stromal and hematopoietic cell network of the adventitia from non-aneurysmal and

aneurysmal human aorta. PLoS One. 2014;9(2):e89983.

3.   Libby P, Loscalzo J, Ridker PM, Farkouh ME, Hsue PY, Fuster V, et al. Inflammation,

Immunity, and Infection in Atherothrombosis: JACC Review Topic of the Week. Journal of the

American College of Cardiology. 2018;72(17):2071-81.

4.   Kurvers H, Veith FJ, Lipsitz EC, Ohki T, Gargiulo NJ, Cayne NS, et al. Discontinuous,

staccato growth of abdominal aortic aneurysms. J Am Coll Surg. 2004;199(5):709-15.

5.   Sakalihasan N, Michel JB, Katsargyris A, Kuivaniemi H, Defraigne JO, Nchimi A, et

al. Abdominal aortic aneurysms. Nat Rev Dis Primers. 2018;4(1):34.

6.   Clement M, Guedj K, Andreata F, Morvan M, Bey L, Khallou-Laschet J, et al. Control

of the T follicular helper-germinal center B-cell axis by CD8(+) regulatory T cells limits

atherosclerosis and tertiary lymphoid organ development. Circulation. 2015;131(6):560-70.

7.   Guedj K, Khallou-Laschet J, Clement M, Morvan M, Delbosc S, Gaston AT, et al.

Inflammatory micro-environmental cues of human atherothrombotic arteries confer to vascular

smooth muscle cells the capacity to trigger lymphoid neogenesis. PLoS One.

2014;9(12):e116295.

8.   Mohanta SK, Yin C, Peng L, Srikakulapu P, Bontha V, Hu D, et al. Artery tertiary

lymphoid organs contribute to innate and adaptive immune responses in advanced mouse

atherosclerosis. Circulation research. 2014;114(11):1772-87.

9.   Nicoletti A, Khallou-Laschet J, Guedj K, Clement M, Gaston AT, Morvan M, et al. L19.

Lymphoid neogenesis in vascular chronic inflammation. Presse Med. 2013;42(4 Pt 2):558-60.

10.   Guo X, Yuan S, Liu Y, Zeng Y, Xie H, Liu Z, et al. Serum IgE levels are associated

with coronary artery disease severity. Atherosclerosis. 2016;251:355-60.

11.   Tuleta I, Skowasch D, Aurich F, Eckstein N, Schueler R, Pizarro C, et al. Asthma is

associated with atherosclerotic artery changes. PLoS One. 2017;12(10):e0186820.

12.   Wang J, Lindholt JS, Sukhova GK, Shi MA, Xia M, Chen H, et al. IgE actions on CD4+

T cells, mast cells, and macrophages participate in the pathogenesis of experimental abdominal

aortic aneurysms. EMBO Mol Med. 2014;6(7):952-69.

13.   Sun J, Sukhova GK, Yang M, Wolters PJ, MacFarlane LA, Libby P, et al. Mast cells

modulate the pathogenesis of elastase-induced abdominal aortic aneurysms in mice. J Clin

Invest. 2007;117(11):3359-68.

14.   Sun J, Zhang J, Lindholt JS, Sukhova GK, Liu J, He A, et al. Critical role of mast cell

chymase in mouse abdominal aortic aneurysm formation. Circulation. 2009;120(11):973-82.

15.   Swedenborg J, Mayranpaa MI, Kovanen PT. Mast cells: important players in the

orchestrated pathogenesis of abdominal aortic aneurysms. Arteriosclerosis, thrombosis, and

vascular biology. 2011;31(4):734-40.

16.   Kaartinen M, Penttila A, Kovanen PT. Accumulation of activated mast cells in the

shoulder region of human coronary atheroma, the predilection site of atheromatous rupture.

Circulation. 1994;90(4):1669-78.

17.   Tsuruda T, Kato J, Hatakeyama K, Kojima K, Yano M, Yano Y, et al. Adventitial mast

cells contribute to pathogenesis in the progression of abdominal aortic aneurysm. Circulation

research. 2008;102(11):1368-77.

18. Merluzzi S, Frossi B, Gri G, Parusso S, Tripodo C, Pucillo C. Mast cells enhance proliferation of B lymphocytes and drive their differentiation toward IgA-secreting plasma cells. Blood. 2010;115(14):2810-7.

19. Furubayashi K, Takai S, Jin D, Muramatsu M, Ibaraki T, Nishimoto M, et al. The significance of chymase in the progression of abdominal aortic aneurysms in dogs. Hypertens Res. 2007;30(4):349-57.

20. Sillesen H, Eldrup N, Hultgren R, Lindeman J, Bredahl K, Thompson M, et al. Randomized clinical trial of mast cell inhibition in patients with a medium-sized abdominal aortic aneurysm. Br J Surg. 2015;102(10):1295.

21. Dahdah A, Gautier G, Attout T, Fiore F, Lebourdais E, Msallam R, et al. Mast cells aggravate sepsis by inhibiting peritoneal macrophage phagocytosis. J Clin Invest. 2014;124(10):4577-89.

22. Caligiuri G, Rossignol P, Julia P, Groyer E, Mouradian D, Urbain D, et al. Reduced immunoregulatory CD31+ T cells in patients with atherosclerotic abdominal aortic aneurysm. Arteriosclerosis, thrombosis, and vascular biology. 2006;26(3):618-23.

23. Saleh R, Wedeh G, Herrmann H, Bibi S, Cerny-Reiterer S, Sadovnik I, et al. A new human mast cell line expressing a functional IgE receptor converts to tumorigenic growth by KIT D816V transfection. Blood. 2014;124(1):111-20.

24. Eggel A, Baravalle G, Hobi G, Kim B, Buschor P, Forrer P, et al. Accelerated dissociation of IgE-FcepsilonRI complexes by disruptive inhibitors actively desensitizes allergic effector cells. J Allergy Clin Immunol. 2014;133(6):1709-19 e8.

25. Manning MW, Cassi LA, Huang J, Szilvassy SJ, Daugherty A. Abdominal aortic aneurysms: fresh insights from a novel animal model of the disease. Vasc Med. 2002;7(1):45-54.

26.   Saraff K, Babamusta F, Cassis LA, Daugherty A. Aortic dissection precedes formation

of aneurysms and atherosclerosis in angiotensin II-infused, apolipoprotein E-deficient mice.

Arteriosclerosis, thrombosis, and vascular biology. 2003;23(9):1621-6.

27.   Fletcher AL, Malhotra D, Acton SE, Lukacs-Kornek V, Bellemare-Pelletier A, Curry

M, et al. Reproducible isolation of lymph node stromal cells reveals site-dependent differences

in fibroblastic reticular cells. Frontiers in immunology. 2011;2:35.

28.   Pipi E, Nayar S, Gardner DH, Colafrancesco S, Smith C, Barone F. Tertiary Lymphoid

Structures: Autoimmunity Goes Local. Frontiers in immunology. 2018;9:1952.

29.   Saunders SP, Ma EGM, Aranda CJ, Curotto de Lafaille MA. Non-classical B Cell

Memory of Allergic IgE Responses. Frontiers in immunology. 2019;10:715.

30.   Trachet B, Aslanidou L, Piersigilli A, Fraga-Silva RA, Sordet-Dessimoz J, Villanueva-

Perez P, et al. Angiotensin II infusion into ApoE-/- mice: a model for aortic dissection rather

than abdominal aortic aneurysm? Cardiovascular research. 2017;113(10):1230-42.

31.   Ngo Nyekel F, Pacreau E, Benadda S, Msallam R, Abrink M, Pejler G, et al. Mast Cell

Degranulation Exacerbates Skin Rejection by Enhancing Neutrophil Recruitment. Frontiers in

immunology. 2018;9:2690.

32.   McGloughlin TM, editor Biomechanics and mechanobiology of aneurysms2011.

33.   He JS, Meyer-Hermann M, Xiangying D, Zuan LY, Jones LA, Ramakrishna L, et al.

The distinctive germinal center phase of IgE+ B lymphocytes limits their contribution to the

classical memory response. J Exp Med. 2013;210(12):2755-71.

34.   Liu CL, Liu X, Zhang Y, Liu J, Yang C, Luo S, et al. Eosinophils Protect Mice From

Angiotensin-II Perfusion-Induced Abdominal Aortic Aneurysm. Circulation research.

2021;128(2):188-202.

35.   Guo W, Gao R, Zhang W, Ge W, Ren M, Li B, et al. IgE Aggravates the Senescence of

Smooth Muscle Cells in Abdominal Aortic Aneurysm by Upregulating LincRNA-p21. Aging

Dis. 2019;10(4):699-710.

36.   Nigrovic PA, Gray DH, Jones T, Hallgren J, Kuo FC, Chaletzky B, et al. Genetic

inversion in mast cell-deficient (Wsh) mice interrupts corin and manifests as hematopoietic and

cardiac aberrancy. Am J Pathol. 2008;173(6):1693-701.

37.   Abdul-Hussien H, Soekhoe RG, Weber E, von der Thusen JH, Kleemann R, Mulder A,

et al. Collagen degradation in the abdominal aneurysm: a conspiracy of matrix

metalloproteinase and cysteine collagenases. Am J Pathol. 2007;170(3):809-17.

38.   Dobrin PB, Baker WH, Gley WC. Elastolytic and collagenolytic studies of arteries.

Implications for the mechanical properties of aneurysms. Archives Of Surgery.

1984;119(4):405-9.

39.   Wang Y, Shiota N, Leskinen MJ, Lindstedt KA, Kovanen PT. Mast cell chymase

inhibits smooth muscle cell growth and collagen expression in vitro: transforming growth

factor-beta1-dependent and -independent effects. Arteriosclerosis, thrombosis, and vascular

biology. 2001;21(12):1928-33.

40.   Davis FM, Daugherty A, Lu HS. Updates of Recent Aortic Aneurysm Research.

Arteriosclerosis, thrombosis, and vascular biology. 2019;39(3):e83-e90.

41.   Wilson JS, Virag L, Di Achille P, Karsaj I, Humphrey JD. Biochemomechanics of

intraluminal thrombus in abdominal aortic aneurysms. J Biomech Eng. 2013;135(2):021011.

42.   Behr Andersen C, Lindholt JS, Urbonavicius S, Halekoh U, Jensen PS, Stubbe J, et al.

Abdominal Aortic Aneurysms Growth Is Associated With High Concentrations of Plasma

Proteins in the Intraluminal Thrombus and Diseased Arterial Tissue. Arteriosclerosis,

thrombosis, and vascular biology. 2018;38(9):2254-67.

1   **Figures**

Fig 1

2   Fig 1. IgE-producing TLOs are present in the adventitia of human AAAs.

(A) Cross-sections of AAAs were stained for CD20 (magenta) and IgE (green). Dense DAPI[+]

lymphoid aggregates highly enriched in CD20+ B cells correspond to TLOs. Several TLOs

displayed a strong IgE staining projecting in the CD20+ B cells comprised in the DAPI "light

zone" of TLOs (white arrowheads). The magnified inset shows 5 adventitial TLOs (cluster of

DAPI+ CD20+ B cells), 3 of which are IgE+. adv: adventitia, m: media, thr; thrombus.

Representative image of 9 different samples. Scale bar: 2.5 mm. (B) Maximum diameter size

of AAAs depending on the absence (no) or the presence of TLOs. p value (Mann-Whitney test)

is indicated on the plot. (C) Serial cross-sections of AAAs were coloured with Carstairs'

staining, where fibrin appears in bright pink, collagen in blue, red blood cells in yellow and

nuclei in dark violet. Perl's+DAB staining highlight redox-active iron in dark brown, nuclei

have been counterstained in nuclear fast red. Orcein staining permits to detect elastin fibres in

dark violet. The lumen is at the bottom of each picture and the media (m) is indicated by the

blue bar on the side of the pictures. Black arrowheads point at red blood cells outside blood

1   vessels, reminiscent of recent intramural haematomas. Scale bar: 1 mm. Right panels:

2   magnified insets. (D) Contrast-enhanced tomography angiograms of an AAA displaying blood

3   disruption from the aortic lumen to the aortic wall through the intraluminal thrombus (white

4   arrow).

Fig 2

Fig 2. IgE-producing TLOs, activated MCs and intramural haematomas are in

close proximity in human AAAs.

(A) Cross-sections of AAAs were coloured with Perl's+DAB staining. TLOs (circled with

dashed line) are found at proximity of haematomas (arrowheads). Scale bar: 500 μm. (B)

Consecutive sections were stained for CD20 (magenta), IgE (green), CD117 (red) and DAPI

(grey). B cells (CD20$^+$ and DAPI$^+$) appear in light pink, autofluorescent red blood cells (low

intensity in the CD20 channel and DAPI$^-$) appear in magenta, and MCs (CD117$^+$ IgE$^+$) appear

in yellow. Scale bar: 250 μm. (C) Magnified inset highlighting the close proximity of IgE$^+$

TLOs (dashed circle), red blood cells (white arrowheads) and MCs (blue arrows). (D)

Magnified inset highlighting MCs (blue arrows) at proximity to intramural haematoma. (E-F)

Adventitia from NAA organ donors and AAA patients were digested and analysed by flow

cytometry after the addition of fluorescent count beads. Singlet, autofluorescent, live CD45$^+$

cells were selected, and MCs were identified as CD3$^-$ CD19$^-$ FcεRIa$^+$ CD117$^+$ cells (E). The

number of MCs (F, left) and activated (CD107a^{hi}) MCs (F, right) was calculated in each sample.

Error bars represent mean +/- standard error; Mann-Whitney test (***, p < 0.001). (G-H) Cross-

sections of AAAs were stained for CD20 (magenta), IgE (green), and tryptase (red). MCs

(tryptase$^+$ IgE$^+$) were detected around TLOs. Scale bar: 100 μm. The magnified inset (H) shows

a degranulating tryptase$^+$ MC at proximity of a TLO.

19
20

Fig 3

Fig 3. MC degranulation and IgE-dependent IL-4 production in response to conditioned medium from AAA adventitia.

(A-D) ROSA MCs were cultured in the presence of conditioned medium from adventitia of control organ donors (NAA) and AAA patients. DARPin® protein bi53_79 (5 μM) was added (+D) or not (-D) to inhibit IgE binding to MCs' FcεRI (C-D). (A, C) After 1 hr, degranulation (MFI CD63) was assessed by flow cytometry. The dotted line indicates CD63 MFI in non-stimulated cells. (B, D) After 4 hrs, mRNA IL-4 level was analysed by RT-PCR ($2^{-\Delta\Delta Ct}$, normalised to RP18S and non-stimulated cells). A,B: Mann-Whitney test; C,D: Wilcoxon matched-pairs signed rank test. (E, F) Concentrations of IL-4, tryptase, and IgE were measured in conditioned medium from AAA adventitia. Pearson correlation analysis (***, $p < 0.001$).

Fig 4

Fig 4. MC depletion decreases aneurysmal growth after dissections in ApoE-RMB

mice.

(A) Experimental protocol, showing the normal course of pseudoaneurysm development in the

Ang II infusion model, and the presence or absence of MCs depending on the treatment (PBS

in blue or DT in red). (B) Toluidine blue stain on aortic root cross-sections of DT- and PBS-

treated mice. The arrow points to one of the MCs detected in the aortic root adventitia of a PBS

control mouse. Scale bar: 250 μm. (C). Quantification of MCs in aortic roots. ***, p < 0.001,

Mann-Whitney test. (D) Macroscopic images of the hearts and aortas of DT- and PBS-treated

mice displaying aneurysms (indicated by arrows). (E) Consecutive cross-sections stained with

Masson Trichrome (top, scale bar: 500 μm) or with Sirius red (bottom; polarised light). Aorta

outer diameter (F) and collagen density of the aortic wall (G) for mice presenting

pseudoaneurysms (PBS: n=7; DT; n=6) were calculated by computer-assisted morphometry on

cross-sections stained with Sirius red. Data from sections taken at different levels (every 300

μm) from each pseudoaneurysm were used for calculations (Fig S8A), and aligned to the layer

with the largest haematoma. Mean +/- standard error; p-values for temperature effect in mixed-

model (REML) analysis.

Graphical Abstract

Graphical Abstract:

Locally produced IgEs, complexed with lesion-specific antigens, activate MCs and trigger a

self-sustained loop involving TLO B cells that may drive the progression of dissecting AAAs.

MC-derived IL-4 promotes the synthesis of more IgEs, which exacerbates the pathology,

potentially leading to AAA rupture. These findings highlight the importance of investigating

the involvement of the IgE/MC/B cell axis in the development and progression of AAA, and

may inform the development of novel therapeutic strategies targeting this pathway.

1 **Table 1: Patients' characteristics according to the presence of intramural**

2 **haematomas.**

	None (n=2, 8%)	Ancient (n=2, 8%)	Recent (n=21, 82%)
Age (years)	70 +/- 7	67+/- 1	71 +/- 2
Male (%)	100%	100%	90%
Max. aortic diameter (cm)	63 +/- 3	66 +/- 14	61 +/- 4
Diabetes (%)	0%	0%	7%
Hypertension (%)	100%	100%	87%
Hyperlipidaemia (%)	0%	50%	53%
Smoking (%)	50%	50%	93%

3 Values are mean +/- SEM or %. Ancient intramural haematomas are characterised by ferrous

4 iron deposits with few red blood cells in the media (Supplemental Figure 3B) and recent

5 intramural haematomas are characterized by abundant red blood cells in the media (Figure 1C).

1 Supporting information

Table S1: Patients'characteristics.

Table S2: NAA healthy donors'characteristics.

Table S3: ApoE-RMB survival and pseudoaneurysm occurrence.

Fig S1. IgEs are elevated in the plasma and adventitia of AAA patients.

Fig S2. GC B cells and plasma cells are elevated in AAA adventitia.

Fig S3. AAAs with ancient or no intramural haematomas.

Fig S4. Proximity of micro-fissures and MC degranulation in human AAAs.

Fig S5. MCs accumulate in the adventitia of human AAAs.

Fig S6. No correlation between tryptase, IgE and IL-4 in the plasma of AAA patients.

Fig S7. Repopulation of basophils in the blood and MCs in the peritoneum after DT depletion

in ApoE RMB mice.

Fig S8. MCs promote pseudoaneurysm expansion after dissection in ApoE-RMB mice.

Supplementary File 1. Antibodies.