Structural evolution of an immune evasion determinant shapes Lyme borreliae host tropism

Ashley L. Marcinkiewicz[1#], Kalvis Brangulis[2,3#*], Alan P. Dupuis, II[1], Thomas M. Hart[1,4†],

Maxime Zamba-Campero[1], Tristan A. Nowak[1,5], Jessica L. Stout[1], Inara Akopjana[2], Andris

Kazaks[2], Janis Bogans[2], Alexander T. Ciota[1,5], Peter Kraiczy[6], Sergios-Orestis

Kolokotronis[7,8,9,10*], Yi-Pin Lin[1,5*]

[1]Division of Infectious Diseases, Wadsworth Center, NYSDOH, Albany, NY, USA; [2]Latvian

Biomedical Research and Study Centre, Riga, Latvia; [3]Department of Human Physiology and

Biochemistry, Riga Stradins University, Riga, Latvia; [4]Department of Biological Sciences,

SUNY Albany, Albany, NY, USA; [5]Department of Biomedical Sciences, SUNY Albany,

Albany, NY, USA; [6]Institute of Medical Microbiology and Infection Control, University

Hospital of Frankfurt, Goethe University Frankfurt, Frankfurt, Germany; [7]Department of

Epidemiology and Biostatistics, School of Public Health; [8]Institute for Genomics in Health;

[9]Division of Infectious Diseases, Department of Medicine, College of Medicine; [10]Department of

Cell Biology, College of Medicine, SUNY Downstate Health Sciences University, Brooklyn,

NY, USA.

Short title: Structural and evolutionary insights into pathogen host tropism

Key words: Lyme disease, CspZ, *Borrelia*, Factor H, Complement

#These authors have contributed equally to this work.

†current address: Yale University School of Medicine, New Haven, CT

24   *correspondence: Kalvis Brangulis, Ph.D.
25   Latvian Biomedical Research and Study Centre, Riga, Latvia LV-1067
26   Telephone: +371 6780 8200
27   Email: kalvis@biomed.lu.lv
28
29   *correspondence: Sergios-Orestis Kolokotronis, Ph.D.
30   Department of Epidemiology and Biostatistics. School of Public Health
31   SUNY Downstate Health Sciences University
32   450 Clarkson Avenue, MSC43A, Brooklyn, NY 11203, USA
33   Telephone: +1 718-270-6741
34   Email: sok@downstate.edu
35
36   *correspondence: Yi-Pin Lin, Ph.D.
37   Wadsworth Center, NYSDOH, Albany, NY 12208
38   Telephone: +1 518-402-2233
39   Email: Yi-Pin.Lin@health.ny.gov
40

ABSTRACT (186 words)

Lyme disease presents a suitable model for the investigation of host tropism – a pathogen's ability to colonize and survive in different host species – since its causative agent, the spirochete *Borrelia burgdorferi* (*Bb*) is transmitted by ticks and maintained in rodent and bird reservoir hosts.

In order to survive in vertebrates and escape from killing by complement, a first-line host immune defense, *Bb* produces the outer surface protein CspZ that binds the complement inhibitor factor H (FH) to promote infection. Protein sequence conservation seems to be linked to FH-binding activity divergence, raising the hypothesis that even minor variation can confer host-specific, FH-binding-mediated infectivity. Our work shows that that this minor variation is located in a loop in the CspZ protein localized in the CspZ-FH binding interface. Our functional experiments prove that this loop promotes bird- or rodent-specific FH-binding activity and infectivity. Swapping loops between rodent- and bird-associated CspZ variants alters their capability to confer host-specific dissemination. We further investigated the evolutionary mechanisms driving the emergence of the CspZ loop-mediated, host-dependent complement evasion. This multifaceted work demonstrates how a single, short protein motif can significantly impact host tropism.

INTRODUCTION

The emergence of most infectious disease outbreaks often involves changes in host tropism, the preferential adaptation of pathogens to selectively invade and persist in hosts (1, 2). Such host tropism is often the result of ongoing host-pathogen interactions (3). Evolution theoretically favors the emergence of host-specializing pathogens, but host-generalist strategies can be advantageous in environments when pathogens have the potential to regularly interact with multiple hosts (4). Specialism vs. generalism can be attributed to polymorphisms within pathogen proteins that differentially interact with host ligands (3). Such polymorphism-mediated host ranges are often thought to require complex host-specific adaptive mechanisms (e.g., (5, 6)) that can be conferred by merely a few amino acids (7-9). Understanding how such minor differences impact diverse host-adapted phenotypes can elucidate the mechanistic insights for the emergence of modern infectious diseases, allowing the development of earlier and more efficient public health interventions.

Species within the *Borrelia* (also known as *Borreliella*) *burgdorferi* sensu lato genospecies complex cause Lyme disease. These spirochetal bacteria are transmitted by ticks and maintained by several reservoir hosts, primarily rodents and birds (10, 11). Lyme borreliae have been genotyped using different polymorphic loci, such as *ospA*, *ospC*, the 16S-23S rRNA intergenic spacer (RST[yping]), and the loci included in multilocus sequence typing (MLST) (12, 13). Laboratory and field studies have shown that not only *Borrelia* species differ in the host species they infect, but that individual genotypes within single spirochete species display distinct preferential host associations, particularly within *B. burgdorferi* sensu stricto (hereafter *B. burgdorferi*), the primary Lyme disease agent in North America (14, 15). This species- and

genotype -specific host selectivity distinguishes Lyme borreliae as a model system for the study

of the molecular basis and evolutionary history of host tropism (14, 15).

Lyme borreliae host associations require the spirochetes to transmit from infected ticks to

hosts, establish an infection at tick biting sites, and disseminate hematogenously to persist at distal

tissues (14, 15). The survival of Lyme borreliae during these discrete events necessitate evasion of

multiple host immune responses, including complement, a first-line innate defense mechanism of

vertebrate extracellular fluid and blood (16-18). This powerful immune mechanism can be

activated by the classical, lectin, or alternative pathways (19). The activation of these pathways

results in the cascading cleavage and recombination of multiple complement components,

ultimately resulting in phagocytic clearance, inflammation, and pore formation on the pathogen

surface by the C5b-9 protein complex to lyse the cells (20). Complement is downregulated by

diverse regulatory proteins to inhibit activation, preventing native cell damage in the absence of

pathogens (21). For example, the alternative pathway is inhibited by factor H (FH), which is

comprised of 20 individual short consensus repeat (SCR) domains (22, 23).

Similar to many bloodborne pathogens that evolved to circumvent complement-mediated

clearance (20), Lyme borreliae produce several complement-inhibitory outer surface proteins (16-

18). These proteins bind and recruit complement components and/or regulatory proteins on

spirochete surface to inactivate complement (16-18). One of these proteins is a FH-binding protein,

CspA (also known as Complement Regulator Acquiring Surface Protein 1, CRASP-1) that we

demonstrated to confer survival of spirochetes via complement evasion in the tick bloodmeal (24).

However, this protein is downregulated after spirochetes infect hosts suggesting a possibilities of

other functionally redundant proteins to facilitate spirochete survival at this infection stage. In fact,

CspZ (also known as CRASP-2), which binds to the 6[th] and 7[th] domains of SCR from FH (SCR6-

7), is upregulated when spirochetes reside in hosts (25, 26). This protein is highly conserved both

between and within different Lyme borreliae species (>85% amino acid (aa) identity between

species and 98% between *B. burgdorferi* strains) (27-30). We and others have previously reported

that FH-binding activity of one CspZ variant promotes spirochete survival in vertebrate sera, and

consequently promotes host infectivity when non-physiologically relevant infection routes are

used (i.e., not tick-transmitted) (31, 32). However, CspZ variants differ in their human FH-binding

activity (27, 28), and vertebrate complement components or regulatory proteins vary between host

taxa (e.g., ~40% aa identity between mammalian and avian FH) (33). These findings raise several

intriguing questions: Could such a minor divergence among CspZ variants confer host-specific

differences in FH-binding activity, resulting in varying host infectivity phenotypes? Further, if

those minor divergences are indeed a determinant of host tropism, how did such divergences

evolve to impact Lyme borreliae-host associations?

In this study, we solved the high-resolution structure of the CspZ-human FH complex,

identified a polymorphic CspZ motif within the FH-binding interface, and defined its contribution

to host-specific FH-binding activity. We then examined the role of this motif in dictating strain-

specific, host-dependent dissemination using mice and quail as rodent and avian models,

respectively. Paired with evolutionary analyses of *cspZ*, we further elucidated how the

evolutionary history behind a minor divergence in an immune evasion determinant can impact host

tropism.

RESULTS

A polymorphic CspZ loop structure defines human FH-binding activity. We set to identify

the regions driving the differences in human FH-binding activity among CspZ variants. We first

pinpointed the amino acids that are involved in FH-binding by co-crystalizing the SCR6-7 of

human FH and a CspZ variant with human FH-binding activity from *B. burgdorferi* strain B408

(CspZ$_{B408}$) (28). The resulting crystal structure at 2.59Å showed an extensive binding interface

between CspZ$_{B408}$ and both SCR6 and SCR7 (**Fig. 1A**). Specifically, Asp47, Tyr50, Asn51, Thr54,

Asn58, and Thr62 in helix-B, Arg142 in helix-F, Asn183 from helix-G, and Tyr214 in helix-I of

CspZ$_{B408}$ interacted with SCR7 (**Fig. 1B, Table S1**). Additionally, Asp71, from the loop between

helix-B and -C, and Asp73 and Ser75 from the same region, bound to SCR7 and SCR6,

respectively (**Fig. 1B, Table S1**).

To locate the residues that impact CspZ FH-binding activity, we aligned the sequences of

CspZ$_{B408}$ with another CspZ variants that binds to human FH (i.e., CspZ from *B. burgdorferi* B31,

CspZ$_{B31}$), as well as CspZ that lacks human FH-binding ability from strain B379 (CspZ$_{B379}$)(28).

All the aforementioned residues involved in human FH-binding are conserved among all three

variants, except Asn51 and Asp71 (**Fig. S1A**). Such conservation also reflects to the three-

dimensional overlayed structure of the previously resolved CspZ$_{B31}$ (30) with our newly-resolved

crystal structures of CspZ$_{B379}$ and CspZ$_{B408}$ (2.10 and 2.45Å, respectively) (**Fig. 1A**). For those

two non-conserved amino acids, Asn51 of CspZ$_{B408}$ is substituted to serine in CspZ$_{B379}$, but the

superimposed structure suggests this would not inhibit human FH-binding in CspZ$_{B379}$ (**Fig. 1B**).

Asp71 of CspZ$_{B408}$ interacts with Lys388 and Tyr390 of SCR6, but this residue is part of an

insertion that is not present in CspZ$_{B31}$ and CspZ$_{B379}$ (**Fig. 1B**). Therefore, these non-conserved

residues alone cannot explain the differential human FH binding ability of these CspZ variants.

However, the C-terminal helix-B and the following loop region between helix-B and -C do harbor

polymorphisms (**Fig. 1A**), with the unique duplication of the last residues in helix-B (Ile60, Met61,

Thr62, and Tyr63) of CspZ$_{B379}$ (**Fig. S1A**), resulting in an extended helix-B in this variant (**Fig.**

1B). Such a structural extension leads to steric hinderance between $CspZ_{B379}$ and human FH to

selectively prevent human FH-binding activity by $CspZ_{B379}$. Overall, our results reveal variant-

specific CspZ structural differences in a loop and the adjacent helix-B (hereafter, "loop structures"),

providing mechanistic insights into the CspZ-mediated, polymorphic human FH-binding activity.

CspZ loop structures determine host-specific FH-binding activity. Similar to CspZ, SCR6-7 is

divergent among vertebrate species (**Fig. S1B**), suggesting a role of the CspZ loop structures in

dictating host-specific FH-binding activity. We found that $CspZ_{B31}$ and $CspZ_{B408}$ –but not

$CspZ_{B379}$– bind to mouse FH, whereas $CspZ_{B31}$ and $CspZ_{B379}$ –but not $CspZ_{B408}$– bind to quail FH

(**Fig. 2A and S2, Table S2**). We then searched for structural evidence of this FH-binding activity

by superimposing the complex structure of $CspZ_{B408}$-human FH with $CspZ_{B31}$, $CspZ_{B379}$, as well

as the previously resolved structures of mouse SCR6-7, and the AlphaFold-predicted structure of

quail SCR6-7 (pLDDT > 70) (34, 35) (**Fig. 2C**). We noted the similar tertiary structures between

mouse and human SCR6-7, consistent with the ability of FH from both origins selectively binds

to $CspZ_{B31}$ and $CspZ_{B408}$ (28) (**Table S2**). Similar to human FH, the $CspZ_{B379}$-specific duplication

by the extended C-terminal helix-B showed potential structural hinderance in the mouse FH-

binding interface (**Fig. 2D**). However, compared to a loop region within human and mouse SCR7,

the equivalent region in quail SCR7 (median pLDDT: 92) is positioned away from the CspZ-FH-

binding interface, allowing sufficient space for extended helix-B of $CspZ_{B379}$ to interact with quail

FH (**Fig. 2D**). Further, the crystal structure of $CspZ_{B408}$-SCR6-7 superimposed with the predicted

structure of quail SCR6-7 showed that Asp71 within the loop of $CspZ_{B408}$ may collide with the

opposite loop region of quail FH, while the equivalent region in mouse SCR7 would not interfere

with the binding of $CspZ_{B408}$ (**Fig. 2D**).

These structural differences among FH from mouse/human vs. quail, pairing with the respective

impacting AAs from different CspZ variants, suggest a possibility of the loop structures of CspZ

to determine CspZ allelically different, host-specific FH-binding activity. To test this possibility,

we swapped the loop regions of the two CspZ variants with distinct host-specific FH-binding

ability to generate two chimeric proteins: $CspZ_{B379}L_{B408}$ has the backbone of a quail FH-binder

($CspZ_{B379}$) and the loop structures of a mouse FH-binder ($CspZ_{B408}$), whereas $CspZ_{B408}L_{B379}$ has

the backbone of $CspZ_{B408}$ and the loop structures of $CspZ_{B379}$ (**Fig. S1A**). We found that

$CspZ_{B379}L_{B408}$ and $CspZ_{B408}L_{B379}$ selectively bound to mouse and quail FH, respectively,

demonstrating the CspZ loop structures are a determinant of host-specific FH-binding activity (**Fig.**

2B and S2).

The CspZ loop structures dictate spirochete strain- and host-specific complement

inactivation. We next examined the host-dependent complement inactivation of these CspZ

variants produced on the spirochete surface. We thus obtained a wild-type *B. burgdorferi* strain

B31-A3 (WT B31-A3) and a *cspZ*-deficient mutant strain in this background harboring an empty

vector (Δ*cspZ*). We complemented this mutant with a plasmid encoding *cspZ_{B31}* (pCspZ_{B31}) (32),

cspZ_{B379} (pCspZ_{B379}), or *cspZ_{B408}* (pCspZ_{B408}), or one of the two loop-swapped mutants

(pCspZ_{B379}L_{B408} and pCspZ_{B408}L_{B379}). We verified there was a similar generation time of these

strains (**Table S3A**), and that there were no differences in the CspZ surface production level (**Fig.**

S3B). Additionally, surface-produced CspZ bound FH in the same host-specific manner as the

recombinant proteins (**Fig. S4**). We then determined the ability of each of these CspZ variants and

mutants to inactivate mouse and quail complement by measuring the deposition levels of mouse

C5b-9 and quail C8, respectively, in the presence of sera from each animal using flow cytometry

(**Fig. 3A-B**). All strains had levels of mouse C5b-9 or quail C8 deposition significantly lower than

the high passage, non-infectious, and mouse and quail complement-susceptible control strain *B.*

burgdorferi B313 (**Fig. 3C-D**)(26). The Δ*cspZ* strain had significantly greater levels of mouse

C5b-9 and quail C8 deposition than WT B31-A3 or pCspZ$_{B31}$ (**Fig. 3C-D**)(32, 36). Compared to

Δ*cspZ*, pCspZ$_{B379}$ had indistinguishable levels of mouse C5b-9 deposition but significantly lower

levels of quail C8, whereas pCspZ$_{B408}$ harbored significantly lower levels of mouse C5b-9 but

indistinguishable levels of quail C8 (**Fig. 3C-D**). Further, pCspZ$_{B379}$L$_{B408}$ recruited significantly

lower levels of mouse C5b-9 but indistinguishable levels of quail C8 than Δ*cspZ*, while

pCspZ$_{B408}$L$_{B379}$ bound indistinguishable levels of mouse C5b-9 but significantly lower levels of

quail C8 than Δ*cspZ* (**Fig. 3C-D**). These results correspond with these variants' host-specific FH-

binding activity, suggesting that the CspZ loop structures-driven host-specific FH-binding activity

confers host-specific complement inactivation.

We also measured the survival of these strains in rodent and quail sera. Note that sera from

white-footed mice (*Peromyscus leucopus*)*,* rather than house mice (*Mus musculus)*, were used to

represent rodent sera, as complement of house mouse is labile *in vitro*, leading to inconsistent

results (37, 38). Both WT B31-A3 and pCspZ$_{B31}$ survived in white-footed mouse and quail sera

more efficiently than Δ*cspZ* (**Fig. 3E-F**) (32). Compared to Δ*cspZ*, pCspZ$_{B379}$ survived at

significantly greater levels in quail but not white-footed mouse sera, whereas pCspZ$_{B408}$ survived

at significantly higher levels in white-footed mouse but not quail sera (**Fig. 3E-F**). Relative to

Δ*cspZ*, pCspZ$_{B379}$L$_{B408}$ survived significantly higher in white-footed mouse but not in quail sera,

whereas pCspZ$_{B408}$L$_{B379}$ survived significantly higher in quail but not in white-footed mouse sera

(**Fig. 3E-F**). However, these differences were not observed in the presence of sera treated with

Cobra Venom factor (CVF) or *Ornithodorus moubata* complement inhibitor (OmCI) that deplete

functional rodent and quail complement, respectively (**Fig. S5**) (39, 40), suggesting that CspZ loop

structures determine spirochete host-specific serum survival.

The CspZ loop structures define host-specific, complement-dependent, early onset

dissemination. We further evaluated if the CspZ loop structure variants determine host infectivity

by allowing *Ixodes scapularis* nymphs carrying similar loads of each of the aforementioned *B.*

burgdorferi strains to feed on mice and quail (**Fig. S6A, S6D**). The burdens of $\Delta cspZ$ were

significantly lower than WT B31-A3 at 7- and 10-days post feeding (dpf) in mouse blood, and at

10 dpf in distal tissues (joints, heart, and bladder), and at 9 dpf in quail blood and distal tissues

(brain and heart) (**Fig. S7**). There were no differences in burdens by a later time point (i.e., 14 dpf,

Fig. S7). We thus performed subsequent experiments on 10 and 9 dpf in mouse and quail,

respectively, for the other of *cspZ*-complemented strains. In mice, we found indistinguishable

spirochete burdens at initial infection sites (tick bite sites) between all seven strains tested at 10dpf

(**Fig. 4A**). The strains $pCspZ_{B31}$, $pCspZ_{B408}$, and $pCspZ_{B379}L_{B408}$, but neither $pCspZ_{B379}$ nor

$pCspZ_{B408}L_{B379}$, colonized mouse blood and distal tissues at significantly higher levels than $\Delta cspZ$

(**Fig. 4B-E**). At 9dpf in quail, we found no significantly different spirochete burdens at tick bite

sites among strains. However, $pCspZ_{B31}$, $pCspZ_{B379}$, and $pCspZ_{B408}L_{B379}$, but neither $pCspZ_{B408}$

nor $pCspZ_{B379}L_{B408}$, colonized quail blood and distal tissues at significantly greater levels than

$\Delta cspZ$ (**Fig. 4F-I**). There were no significantly different burdens of any strain in any tissues from

mice and quail with complement deficiency ($C3^{-/-}$ mice or OmCI-treated quail) (**Fig. S8**). These

results demonstrate that the CspZ loop structures define complement-dependent, spirochete strain-

and host-specific early onset dissemination.

The population-wide CspZ loop structures evolved from a variant with versatile FH-binding ability. To investigate the evolutionary history of the CspZ loop structures, we mined *B. burgdorferi cspZ* from NCBI GenBank and sequence read archive (totaling 174 high-quality *cspZ* isolates originated from ticks, reservoir hosts, and patients, across North America, Europe, and Asia). Phylogenetic analysis revealed three lineages with uncertain grouping among them, suggesting insufficient variation of *cspZ* to allow for well-established phylogenetic groups to emerge (data not shown). However, a haplotype-based phylogenetic network distinctly separated these three individual lineages, each of which contains one of the above-tested *cspZ* alleles: $cspZ_{B31}$, $cspZ_{B379}$, or $cspZ_{B408}$ (**Fig. S9**). Isolates within the same lineage had over 99% sequence identity and the same loop structure compared to $cspZ_{B408}$, $cspZ_{B379}$, or $cspZ_{B31}$ (containing the duplication, insertion, or neither at the loci encoding loop structure, respectively) (**Fig. 5A**). We pinpointed single amino acid polymorphisms (SAPs) in the isolates and found none of the SAPs within each lineage were located in the FH-binding interface (**Table S1**), with the exception of a SAP in one isolate in the $CspZ_{B379}$ lineage (**Table S4**)). Rather, the majority of the SAPs were between lineages, and were located in the loop structures that are driving the FH-binding abilities. While this present study has been the only one to investigate the abilities of CspZ variants to bind house mouse and quail FH, the human FH-binding abilities of 13% of these 174 isolates have been evaluated (27, 28, 32, 41). Incorporating these findings with the phylogeny, we found a 100% correlation of known host-specific FH-binding ability with lineage (**Fig 5A**). These results suggest each CspZ lineage with distinct loop structures can be linked to CspZ-specific, host-dependent FH-binding ability.

Additionally, we explored the evolutionary mechanisms that could have led to this minor loop structure-dependent FH-binding specificity. No evidence was found for recombination or past

changes to the effective population size (data not shown). Although there was no evidence of

selection for *cspZ* in its entirety (data not shown), several codons were undergoing either positive

or negative selection (**Fig. S1A**). We reconstructed the possible last common ancestor states

("LCAS") of the entire *B. burgdorferi* CspZ population using the chromosome mutation rate

estimated in a previous study (42). We found that the diversification of the three lineages was

estimated to be 261 to 784 years ago (y.a.) whereas the diversification of the LCAS likely occurred

approximately 2166y.a. (896-5718 $HPD_{5-95\%}$) (**Table S5**). The LCAS variants have a loop region

resembling that of $CspZ_{B31}$ both in AAs and structure (**Fig. S1A, S10**), and versatilely bound to

human, mouse, and quail FH, similar to $CspZ_{B31}$ (**Fig. 5B, Table S6**). These results suggest the

diversification of CspZ is arisen relatively recently from a generalist variant with versatile FH-

binding features.

DISCUSSION

Host tropism is an outcome of ongoing host-pathogen interactions (15). In vector-borne

zoonotic pathogens, such interactions include vector-to-host transmission, pathogen dissemination

and persistence, and host-to-vector acquisition (43). For Lyme borreliae to survive throughout

these infection steps, spirochetes need to overcome numerous host immunological mechanisms,

including complement-mediated killing (17, 18, 44). In fact, Lyme borreliae inactivate

complement in the tick blood meal for tick-to-host transmission and in the host bloodstream for

dissemination and persistence (32, 45-47). In this study, we examined this inactivation conferred

by a Lyme borreliae FH-binding protein, CspZ. We showed early colonization defects of a tick-

introduced, *cspZ*-deficient mutant at distal tissues, defining this polymorphic protein as a

contributor to the early stages of dissemination. Superimposing the crystal structures of three

polymorphic CspZ variants from genotypically-diverse *B. burgdorferi* strains revealed a variable

motif of a short AA stretch (30). Our newly-resolved and software-predicted complex structures

of CspZ with human, mouse, and quail SCR6-7 further linked those loop-encoded variable residues

to distinct host-specific FH binding activity. These results are congruent with the ability of CspZ

variants and their associated mutants with swapped loop structures to promote host-specific FH-

binding activity, complement evasion, and dissemination, defining this CspZ loop structure as a

determinant of host tropism. Our findings thus demonstrate the concept that minor variation could

functionally impact host-adapted phenotypes, potentially modulating host tropism (7-9). Our

findings can also be attributed to a structurally unique FH-binding mechanism, as the FH-binding

interface of CspZ is significantly different from that of the only other structurally-characterized

SCR6-7-binding pathogen protein, *Neisseria meningitidis* Fhbp (48)(**Fig. S11**). Taken together,

these results provide a platform of using structure-guided approaches in identifying pathogens'

host tropism determinants and their biological functions.

We also integrated an evolutionary approach to investigate the emergence of such host

tropism determinants, an approach whose importance has been recently demonstrated to

understand and track modern disease emergence (e.g., (49, 50)). Our findings demonstrate that the

B. burgdorferi CspZ variants diversified from a last common ancestor that could bind human,

mouse, and quail FH. This diversification occurred relatively recently (**Table S10**), compared to

the previous finding showing that *B. burgdorferi* as a species evolved about 60,000 years BP, and

the other FH-binding protein, CspA, diversified about 25,000 years y.a. (42, 45). The fact that

CspZ has no orthologs in non-*Borrelia* organisms also supports the young age of this protein,

suggesting CspZ may be undergoing weaker purifying and more variable selective pressure (30,

51). We did not detect selective pressure on the entirety of the *cspZ* locus, but rather only in a few

individual codons, none of which were in the loop structures driving FH-binding phenotypes. The

loop structures, which are the most variable part of these CspZ variants, contain indels, so the

detection of gene-wide selective pressures was likely masked by the exclusion of gaps by the

selection tests. In fact, other polymorphic anti-complement loci, particularly *ospC*, are undergoing

balancing selection (52), (53). Both Multiple Niche Polymorphism (MNP) and Negative

Frequency Dependent Selection (NFDS) have been proposed as mechanisms driving this selection.

On the one hand, NFDS proposes that a high diversity of alleles emerge through the negative

correlation between spirochete fitness and the frequency of any allele in the overall population

(54). In Lyme borreliae, NFDS may explain the greater antibody-mediated clearance of spirochete

strains harboring *ospC* alleles of higher frequency loci (55, 56). On the other hand, MNP proposes

that the diversity of alleles/genotypes is maintained by fitness variation of these spirochetes across

reservoir hosts (53, 57). Though it is possible that lack of selective pressure detection reflects the

neutral selection experienced in *cspZ*, as a systemically redundant anti-complement protein (58),

the possibility that *cspZ* is undergoing the same selective pressures as other functionally redundant

proteins (e.g., *ospC*) cannot be excluded. In fact, balancing selection on one locus can increase

diversity in another genetically-linked locus (59), and there is genetic linkage between CspZ types

and *ospC* types (27, 28, 36). This may raise an intriguing possibility that the balancing selection

on *ospC* may allow diversification through drift in other functionally redundant genes like *cspZ*

without deleterious effect to the spirochetes, which warrants further investigations.

Some Lyme borreliae species or strains carry determinants that promoting host-specific

phenotypes that differ from the known host range of these species or strains (36, 60-62). For

example, the *B. burgdorferi* strain 297 is highly infectious in mice, but its CspZ loop-structures

are identical to those in $CspZ_{B379}$ from the strain B379 that is not mouse adapted (36, 62). This

reflects the fact that most conclusions from this study were drawn by using spirochete strains with

the same genetic background but producing the CspZ variants of interest, which may not account

for polygenic contributions of other proteins in different strains (36). One possibility that addresses

this discrepancy is that host-adapted phenotypes are conferred by not only CspZ but other anti-

complement proteins, and the functional contribution from each of these proteins differ (and/or are

host-dependent). In fact, the colonization defects of the *cspZ*-deficient *B. burgdorferi* were not

found in the initial infection site at early infection stages, or in the distal tissues at later timepoints,

consistent with the concurrent production of other FH-binding proteins (25, 45, 63-65).

Additionally, non-FH-binding anti-complement proteins (66-68) and/or tick salivary proteins with

anti-complement functions (69-71) could also drive complement-based phenotypes. Further,

pathogen evasion to complement-independent mechanisms (e.g., antibodies), or even the host

immune evasion-independent adaptive phenotypes (e.g., adhesion), can occur simultaneously with

the anti-complement mechanisms (72), all contributing to the overall host-adapted phenotypes (6).

All these confounding factors may complicate the delineation of the roles for CspZ at later stages

of infection. Thus, our results do not rule out the possibility of CspZ-mediated complement evasion

at distal tissues during post-dissemination timepoints, but rather highlight the importance of this

protein immediately after spirochetes begin to disseminate. Despite such caveats, the strength of

these isogenic strains needs to be emphasized in delineating the roles of a particular pathogen

determinant from an enormous complexity of functionally redundant proteins (24, 45, 73, 74).

Such potential polygenic or multi-factorial, and host-specific phenotypes should consider the

complexity of numerous determinants across different mechanisms in defining the effect on host

tropism.

In this study, we used Lyme borreliae as a model to apply structural, microbiological, and

evolutionary approaches to identify a determinant of host-adapted phenotypes and the specific

mechanism impacting host tropism. It should be noted that distinct *B. burgdorferi* infectivity have

been observed for host species within the same taxa (e.g., house mice vs. the reservoir white-footed

mice) (6, 75, 76). Thus, we do not intend to extend the host-specific phenotypes from these

laboratory models to every individual reservoir animal in the same host taxa. Rather, our results

establish a platform with more accessibility of tools to facilitate the identification of host tropism

determinants and their underlying mechanisms. Such results build the foundation to further

examine similar concepts in reservoir animals to recapitulate field findings in understanding the

mechanisms of host tropism in a controllable laboratory setting. The information and the platform

established in this multi-disciplinary study establishes greater insights of pathogen-host

interactions, facilitating the understanding of host tropism as the cause of newly emergent

infectious diseases.

MATERIALS AND METHODS

Ethics statement. All mouse and quail experiments were performed in strict accordance with all

provisions of the Animal Welfare Act, the Guide for the Care and Use of Laboratory Animals, and

the PHS Policy on Humane Care and Use of Laboratory Animals. The protocol was approved by

the Institutional Animal Care and Use Committee (IACUC) of Wadsworth Center, New York State

Department of Health (Protocol docket number 19-451). All efforts were made to minimize animal

suffering.

Mice, quail, ticks, bacterial strains, animal sera, OmCI, and FH. BALB/c and Swiss Webster

mice were purchased from Taconic (Hudson, NY). C3$^{-/-}$ mice in the BALB/c background were

generated from the C3$^{-/-}$(C57BL/6) from Jackson Laboratory (Bar Harbor, ME) as described (24).

Coturnix quail were purchased from Cavendish Game Birds Farm (Springfield, VT). *Ixodes*

scapularis tick larvae were purchased from National Tick Research and Education Center,

Oklahoma State University (Stillwater, OK) or obtained from the CDC through BEI Resources

(Manassas, VA).

The *Escherichia coli, Pichia pastoris* and *Borrelia* strains used in this study are described

in **Table S7**. *E. coli* strains DH5α, BL21(DE3), and derivatives were grown in LB broth or agar,

supplemented with kanamycin (50µg/ml), ampicillin (100µg/ml), or no antibiotics as appropriate.

P. pastoris strain X-33 was grown on YPD plates supplemented with zeocin (800µg/ml) or BMGY

medium supplemented with 1% methanol. All *B. burgdorferi* strains were grown in BSK-II

completed medium supplemented with kanamycin (200µg/mL), streptomycin (50µg/mL), or no

antibiotics (**Table S7**).

Mouse FH was purchased from MyBiosource (San Diego, CA). Quail FH and recombinant

OmCI proteins were generated as described previously (24, 32, 36, 39). The mouse and quail sera

were obtained from Southern Biotech, Inc (Birmingham, AL) and Canola Live Poultry Market

(Brooklyn, NY), respectively. The sera from white-footed mice were obtained previously (6). Prior

to being used, all these sera were screened for antibodies against the C6 peptide of the *B.*

burgdorferi protein VlsE (77) with the C6 Lyme ELISA kit (Diamedix, Miami Lakes, FL) to

ensure the mice did not have prior exposure to *B. burgdorferi*.

Generation of recombinant CspZ proteins and recombinant human FH SCR6-7. To generate recombinant CspZ proteins for crystallization, $cspZ_{B379}$ (GenBank: FJ911671.1) and $cspZ_{B408}$ (GenBank: FJ911677.1) were amplified by PCR from the genomic DNA of *B. burgdorferi* strain B379 and B408 using the primers listed in **Table S8**. Note that B408 has two copies of *cspZ*, but they are functionally identical with only a single synonymous SNP at nucleotide 699 (36, 78). Based on the prediction by SignalP 4.1 (79) and according to our previous structural data from $CspZ_{B31}$ (30), the lipoprotein signal peptide (residues 1-22) was excluded from the amplified gene. The introduced *NcoI* and *NotI* restriction sites were used for ligation of the amplified fragments into the pETm-11 expression vector which contains the coding region for an N-terminal 6xHis tag and a tobacco etch virus (TEV) protease cleavage site. Expression in *E. coli*, purification by affinity chromatography, and 6xHis tag cleavage by TEV protease of both proteins $CspZ_{B379}$ and $CspZ_{B408}$ were performed similarly as described previously for $CspZ_{B31}$ (30). The purified and cleaved proteins were buffer exchanged into 10 mM Tris-HCl (pH 8.0) and concentrated to 11 mg/ml using an Amicon centrifugal filter unit (Millipore, Burlington, MA, USA).

To produce recombinant human FH for crystallization, the gene encoding the SCR6-7 of human FH was synthesized by BioCat GmbH (Heidelberg, Germany) and cloned into pPICZαA vector behind the α-factor secretion signal using *XhoI* and *NotI* restriction sites in a way to restore the Kex2 signal cleavage site. The plasmid was linearized with *PmeI* and transformed by electroporation into *Pichia pastoris* (reassigned as *Komagataella phaffii*) strain X-33. Transformants were obtained on YPD agar plates containing 800µg/ml of the antibiotic zeocin. The selected clone was cultivated 24-h in BMGY medium at 30°C with aeration (250rpm) following addition of 1% methanol daily, and cultivation was continued for three more days. The cell pellet was removed by low-speed centrifugation. Supernatant was buffer-exchanged into

50mM sodium phosphate (pH 6.0) by Sephadex G-25 Fine column (bed volume 360ml) (Cytiva,

Marlborough, MA, USA) in 100ml portions at a flow rate of 20ml/min. Two liters of supernatant

was passed through the CaptoS Improved Resolution column (bed volume 20ml) (Cytiva,

Marlborough, MA, USA) and bound material was eluted with a linear salt gradient at a flow rate

of 6ml/min. Target protein fractions were selected based on SDS-PAGE. The relevant fractions

were pooled and buffer-exchanged into 20mM Tris-HCl (8.0), 50mM NaCl and 10mM NaH_2PO_4

using an Amicon filter device (Millipore, Burlington, MA, USA).

To generate recombinant $CspZ_{B379}$ and $CspZ_{B408}$ for the studies other than crystallization,

the region encoding *cspZ_{B379}* or *cspZ_{B408}* without the signal peptide was amplified as described

above and engineered to encode *BamHI* and *SalI* sites at the 5' and 3' ends, respectively, allowing

subsequent cloning into the pJET cloning vector (Thermo Fisher Scientific, Waltham, MA). These

pJET-derived plasmids encoding *cspZ_{B379}* or *cspZ_{B408}* were used as template for site-directed,

ligase-independent mutagenesis (SLIM) (**Table S7**) to generate plasmids producing $CspZ_{B379}$-

L_{B408} and $CspZ_{B408}$-L_{B379} (80). After verifying the sequences of all the plasmids (Wadsworth

ATGC facility), the DNA fragments were subsequently excised using *BamHI* and *SalI* and then

inserted into the same sites in pGEX4T2 (GE Healthcare, Piscataway, NJ) (32). The pGEX4T2-

derived plasmids were then transformed into the *E. coli* strain BL21(DE3). The GST-tagged CspZ

proteins were produced and purified by affinity chromatography. These proteins were verified for

their secondary structures not impacted by the mutagenesis using CD (**Fig. S12**), as described in

the section "Circular dichroism (CD) spectroscopy."

To generate recombinant CspZ from the last common ancestor states, pET-28a+ encoding

these states flanked by *BamHI* and *SalI* sites at the 5' and 3' ends, respectively, were cloned

(Synbio Technologies, Monmouth Junction, NJ). The plasmids were transformed into the *E. coli*

strain BL21(DE3), and the His-tagged CspZ proteins were produced and purified by affinity

chromatography.

Crystallization and structure determination. For crystallization of CspZ$_{B379}$ and CspZ$_{B408}$, 96-

well sitting drop plates were set using a Tecan Freedom EVO100 workstation (Tecan Group,

Männedorf, Switzerland) by mixing 0.4μl of protein with 0.4μl of precipitant using the 96-reagent

sparse-matrix screens JCSG+ and Structure Screen 1&2 (Molecular Dimensions, Newmarket, UK).

The crystals for CspZ$_{B379}$ were obtained in 0.2M Ammonium citrate and 24% PEG 3350. For

CspZ$_{B408}$, the crystals were formed in 0.2M potassium acetate, 0.1M Tris-HCl (pH 8.0) and 28%

PEG 3350. Prior to the data collection, the crystals were frozen in liquid nitrogen. An additional

20% glycerol was used as a cryoprotectant for CspZ$_{B379}$ crystals, whereas the respective precipitant

with an additional 14% glycerol was used as cryoprotectant for CspZ$_{B408}$ crystals.

CspZ$_{B408}$ (4mg/ml) and human SCR6-7 (3mg/ml) were mixed together at a molar ratio of

1:2 and loaded on a HiLoad 16/600 Superdex 200 prep grade column (GE Healthcare, Chicago,

IL, USA) pre-equilibrated with 20mM Tris-HCl (8.0), 50mM NaCl and 10mM NaH$_2$PO$_4$. The

flow rate was set to 2ml/min. Size exclusion chromatography resulted in one major peak containing

the complex, confirmed by SDS-PAGE. Crystallization was set as described earlier for CspZ$_{B379}$

and CspZ$_{B408}$ by mixing 0.4μl of protein with 0.4μl of precipitant and using the 96-reagent sparse-

matrix screens. The crystals for CspZ$_{B408}$-SCR6-7 complex were obtained in 0.2M Zinc acetate,

0.1M imidazole (pH 7.4) and 10% PEG 3000. Crystals were frozen in liquid nitrogen by using

20% glycerol as a cryoprotectant.

Diffraction data for CspZ$_{B379}$, CspZ$_{B408}$ and CspZ$_{B408}$-SCR6-7 complex were collected at

the MX beamline instrument BL 14.1 at Helmholtz-Zentrum, Berlin (81). Reflections were

indexed by XDS and scaled by AIMLESS from the CCP4 suite (82-84). Initial phases for $CspZ_{B379}$

and $CspZ_{B408}$ were obtained by molecular replacement using Phaser (85), with the crystal structure

of the orthologous protein $CspZ_{B31}$ was used as a searching model (97% sequence identity, PDB:

4CBE). For $CspZ_{B408}$-SCR6-7 complex, the phases were determined using $CspZ_{B408}$ (PDB: 7ZJK,

RMSD 0.98 Å) and human FH SCR6-7 (PDB: 4AYD-A, RMSD 0.98 Å) as the searching models.

After molecular replacement, the protein models were built automatically in BUCCANEER (86).

The crystal structures were improved by manual rebuilding in COOT (87). Crystallographic

refinement was performed using REFMAC5 (88). A summary of the data collection, refinement

and validation statistics for $CspZ_{B379}$, $CspZ_{B408}$ and $CspZ_{B408}$-SCR6-7 complex are given in **Table**

S9.

Protein 3D structure prediction using AlphaFold. AlphaFold v2.0 (34) was used to predict the

3D structure for quail FH SCR6-7 extrapolated from the sequences of *Coturnix japonica*

complement FH (GenBank: XM_015869474.2). Structure prediction with AlphaFold v2.0 was

performed according to the default parameters as indicated at the website

https://github.com/deepmind/alphafold/ running on AMD Ryzen Threadripper 2990WX 32-Core;

128 GB RAM; 4 x NVIDIA GeForce RTX 2080, and using the full databases downloaded on

2021-09-25. For further structural analysis, only the predicted structure with the highest

confidence was used (as ranked by using LDDT (pLDDT) scores).

Circular dichroism (CD) spectroscopy. CD analysis was performed on a Jasco 810

spectropolarimeter (Jasco Analytical Instrument, Easton, MD) under nitrogen. CD spectra were

measured at room temperature (RT, 25°C) in a 1mm path length quartz cell. Spectra of each of the

CspZ proteins (10μM) were recorded in phosphate based saline buffer (PBS) at RT, and three far-

UV CD spectra were recorded from 190 to 250nm for far-UV CD in 1nm increments. The

background spectrum of PBS without proteins was subtracted from the protein spectra. CD spectra

were initially analyzed by the software Spectra Manager Program (Jasco). Analysis of spectra to

extrapolate secondary structures were performed using the K2D3 analysis programs (89).

ELISAs. Quantitative ELISA was used to determine FH-binding by CspZ proteins, or ancestral

proteins, as described previously (24, 90), with the following modifications: Mouse anti-GST tag

or mouse anti-His tag 1:200× (Sigma-Aldrich) and HRP-conjugated goat anti-mouse IgG 1:2,000×

(Seracare Life Sciences) were used as primary and secondary antibodies, respectively, to detect

the binding of GST- or histidine-tagged proteins.

Surface Plasmon Resonance (SPR). Interactions of CspZ proteins with FH were analyzed by

SPR using a Biacore T200 (Cytiva, Marlborough, MA). Ten micrograms of mouse or quail FH

were conjugated to a CM5 chip (Cytiva) as described previously (90). For quantitative SPR

experiments, 10μL of increasing concentrations (0.08, 0.03125, 0.0125, 0.5, 2μM) of each of the

CspZ proteins were injected into the control cell and the flow cell immobilized with FH at 10μl/min,

25°C. To obtain the kinetic parameters of the interaction, sensogram data were fitted by means of

BIAevaluation software version 3.0 (GE Healthcare), using the one step biomolecular association

reaction model (1:1 Langmuir model), resulting in optimum mathematical fit with the lowest Chi-

square values.

Shuttle vector construction and plasmid transformation into *B. burgdorferi*. "Loop swapped"

CspZ variants (i.e., $CspZ_{B379}L_{B408}$ and $CspZ_{B408}L_{B379}$) were designed based on the full-length

sequences (B379 accession: OM643341; B408: accession: OM643340) and purchased as double-

stranded DNA fragments flanked by *BamHI* and *SalI* on the 5' and 3', respectively (Integrated

DNA Technologies, Inc., Coralville, IA). B31-A3Δ*cspZ* was complemented with these variants,

or with native CspZ from B379 and B408 flanked by the same restriction enzyme sites (**Table S7**),

in the same manner as the previously published strains of B31-A3Δ*cspZ*/pKFSS and B31-

A3Δ*cspZ*/pCspZ$_{B31}$ (32). The plasmid profiles of these spirochetes were examined to ascertain

identical profiles between these strains and their parental strain B31-A3 (91). The generation time

of these transformants was calculated as previously described (32).

Flow cytometry. CspZ production and FH-binding on spirochete surface were determined as

described (24), including blood-treatment to induce the production of CspZ (32). To determine the

levels of mouse C5b-9 or quail C8 deposition on the surface of spirochetes, mouse or quail sera

were incubated with 1×10^7 spirochetes in PBS at a final concentration of 20% at 25°C for one hour

and the detection procedure has been described previously (36). Basically, after incubation,

spirochetes were washed then resuspended in HBSC-DB (25mM Hepes acid, 150mM sodium

chloride, 1mM $MnCl_2$, 1mM $MgCl_2$, 0.25mM $CaCl_2$, 0.1% glucose, and 0.2% BSA). Rabbit anti-

mouse C5b-9 polyclonal IgG (1:250x) (Complement Technology, Tyler, TX) or mouse anti-quail

C8 polyclonal sera (1:250x) (36) were used as the primary antibodies. An Alexa 647-conjugated

goat anti-rabbit (ThermoFisher) or a goat anti-mouse IgG (ThermoFisher) (1:250x) was used as

the secondary antibody. After staining, the spirochetes were fixed with 0.1% formalin. The

resulting fluorescence intensity of spirochetes was measured and analyzed by flow cytometry

using a FACSCalibur (BD Bioscience) as described (32, 36).

Serum resistance assays. The serum resistance of *B. burgdorferi* was measured as described with

modifications (6, 24, 32). Cultures in mid-log phase of each strain treated with human blood (32),

as well as the high passaged, non-infectious, and serum-sensitive human blood-treated *B.*

burgdorferi strain B313 (control), were cultivated in triplicate and diluted to a final concentration

of 5×10^6 cells/mL in BSK-II medium without rabbit sera. The cell suspensions were mixed with

sera collected from naïve white-footed mice or quail (60% spirochetes and 40% sera) in the

presence or absence of 2μM of CVF (Complement Technology) or recombinant OmCI, to deplete

complement from mouse and quail sera, respectively. Heat-inactivated sera (65°C for 2-h) were

also included in each of the aforementioned combinations as a control (all strains survived equally;

data not shown). To determine the bacteria survival, the number of motile spirochetes was counted

under dark field microscopy at 0- and 4-h following incubation with sera as described (32), as we

have shown that motile spirochetes determined using this methodology accurately reflect results

using live-dead staining assays (39). The percent survival of *B. burgdorferi* was calculated by the

normalization of motile spirochetes at 4-h post incubation to that immediately after incubation

with sera.

Mouse and quail infection by ticks. Generating flat, infected *I. scapularis* nymphs has been

described previously (24, 92). The infected nymphs were placed in a chamber to feed on 4- to 6-

week old male and female BALB/c or C3$^{-/-}$ mice in BALB/c background, or on four- to six-week

old male and female untreated or OmCI-treated quail, as described previously (45). For OmCI-

treatment, the quail were subcutaneously injected with OmCI (1mg/kg of quail) one day prior to

the nymph feeding. The engorged nymphs were obtained from the chambers at four days post tick

feeding. Animals were sacrificed and tissues collected from the mice at 7-, 10-, or 14- (blood, tick

bite site of the skin, heart, bladder, tibiotarsus joint), and quail (blood, tick bite site of the skin,

heart, brain) at 9-, or 14-, days post nymph feeding. To ensure OmCI was still functional at these

timepoints, quail were subcutaneously injected with OmCI (1mg/kg of quail) or PBS buffer

(control), and the sera were collected at 10 days post injection (equivalent to 11 days post nymph

feeding). The lack of the ability of this serum to kill the sera-sensitive *B. burgdorferi* strain B313

(i.e., to ensure complement was still depleted) was assessed as described in the section "Serum

resistance assays." (**Fig. S13**)

Quantification of spirochete burden The DNA from tissues, blood, and ticks was extracted as

described previously (45). qPCR was then performed to quantitate bacterial loads. Spirochete

genomic equivalents were calculated using an ABI 7500 Real-Time PCR System (ThermoFisher

Scientific) in conjunction with PowerUp SYBR Green Master Mix (ThermoFisher Scientific)

based on amplification of the Lyme borreliae *recA* gene using primers BBRecAfp and BBRecArp

(**Table S8**) with the amplification cycle as described previously (32). The number of *recA* copies

was calculated by establishing a threshold cycle (Cq) standard curve of a known number of *recA*

gene extracted from *B. burgdorferi* strain B31-A3, then comparing the Cq values of the

experimental samples.

Genomic analyses. To generate the *cspZ* phylogenetic trees, we mined all publicly available *cspZ*

sequences on NCBI as of September 2021, including assembled genomes, nucleotides, and

unassembled genomes on the SRA. To pull *cspZ* from unassembled genomes, the short reads were

aligned to *cspZ* from B31, B379, or B408 with UGENE v39.0 using BWA-MEM at defaults (93,

94). Strains were removed from the analyses if the coverage was too low, there was evidence of

PCR/sequencing errors (e.g., non-conserved homopolymer length) or evidence of multiple CspZ

variants within one strain. All resulting 174 *cspZ* sequences, plus the outgroup strains (*B.*

spielmanii A14s accession: EU272854.1; *B. afzelii* FEM4 accession: OM243915; *B. afzelii* VS461

accession: MN809989.1; *B. garinii* PBr accession: CP001307.1; *B. bissettii* DN127 accession:

NC_015916.1; *B. bissettii* CO275 accession: JNBW01000464.1), were aligned by codons using

TranslatorX (95). All isolates were collapsed into haplotypes in FaBox v1.61, and these haplotypes

were used with the *B. bissettii* outgroup to generate a NeighborNet network in SplitsTree v4 (96,

97). Phylogenetic trees were estimated using likelihood as optimality criterion in IQ-tree v1.6.12

(98) and a full substitution model search procedure in ModelFinder (99). Internode branch support

was estimated with 10,000 replicatess of both ultrafast bootstrapping and the SH-aLRT branch test

(98-100). All resulting phylogenetic trees were visualized in iTOL v6.4.3 (101). The pairwise

sequence similarity for each of the 174 *B. burgdorferi cspZ* isolates relative to $CspZ_{B31}$, $CspZ_{B408}$,

or $CspZ_{B379}$ was determined in MEGA-X with default settings (102). Putative recombination

breakpoints were analyzed with GARD (103), and evidence of selection for individual codons or

branches on the *cspZ* gene tree were inferred with FUBAR (104), FEL (105) and MEME (106),

all on the Datamonkey server (107). Selection tests for the entire *cspZ* gene included Tajima's

neutrality test in MEGA-X (102) and all tests available in DnaSP v5 with 10,000 coalescent

simulations (108), all at default settings. The ancestor state for the entire *B. burgdorferi* ingroup

was reconstructed using the LG model in GRASP 2020.05.05 (109, 110), as well as FireProt-ASR

with default settings (111) using both full and haplotype phylogenies, a multitaxon outgroup and

solely *B. bissettii* as outgroup. Divergence dating was carried out in BEAST v1.10.4 (112) using

the HKY+Γ_4 substitution model(113, 114), a coalescent Bayesian skyline coalescent model, and a

strict clock with a uniform prior on the substitution rate using a previously determined rate of

4.75e-06 substitutions/site/year(42). A Markov chain Monte Carlo chain length of 100 million

steps was used with a 10,000-step thinning, resulting in effective sample sizes greater than 200, an

indication of an adequate chain mixing. The analyses were ran in triplicate and combined after

removing a 10% burn-in.

The amino acids encoding SCRs 6-7 from human (GenBank accession U56979.1), mouse

(NM_009888.3), or quail (XM_015869474.2) FH or CspZ$_{B31}$, CspZ$_{B379}$, and CspZ$_{B408}$, the loop-

swapped variants, and the reconstructed ancestral CspZ sequences were aligned in MEGA-X using

ClustalW with default settings, analyzed with ESPript v3.0, and visualized with Jalview v2.11.0

(102, 115, 116).

Statistical analysis. Samples were compared using the Mann-Whitney *U* test or the Kruskal-

Wallis test with the two-stage step-up method of Benjamini, Krieger, and Yekutieli (117).

Accession numbers. The coordinates and the structure factors for CspZ$_{B379}$, CspZ$_{B408}$, and human

SCR-CspZ$_{B408}$ have been deposited in the Protein Data Bank with accession codes 7ZJJ, 7ZJK,

and 7ZJM, respectively.

ACKNOWLEDGEMENTS

We thank Simon Starkey, Quinton Smith, Deirdre Torrisi, and Joey Anderson from the Wadsworth

Veterinary Sciences facility for animal husbandry; Richard Marconi, Utpal Pal and Patricia Rosa

for providing the *B. burgdorferi* strains; David Vance and Nicholas Mantis for sharing the *E. coli*

strain; Timothy Czajka for the assistance of SPR analysis; Nikhat Parveen, Klemen Strle, and

Grace Chen for insightful comments; and Laurel Lown for assistance with running PCRs. We also

appreciate Leslie Eisele and Renjie Song of Wadsworth Biochemistry and Immunology Core for

CD spectroscopy, SPR, and flow cytometry; and Karen Chave of the Wadsworth Protein

Expression Core for the help of the materials to purify factor H. We thank Wadsworth ATGC core

for plasmid sequencing. Diffraction data for *B. burgdorferi* CspZ proteins and CspZ-human FH

were collected on BL14.1 at the BESSY II electron storage ring operated by the Helmholtz-

Zentrum, Berlin. We would particularly like to acknowledge the help and support of

Manfred S. Weiss and Jan Wollenhaupt during the diffraction data collection. This work was

supported by NSF IOS1754995 (SOK), NSF IOS1755286 (YPL, ALM, TAN, ADII, JS, and AC),

DoD TB170111, NIHR21AI144891, NIH R21AI146381, New York State Department of Health

Wadsworth Center Start-Up Grant (ALM, TAN, YPL), and LOEWE Center DRUID Novel Drug

Targets against Poverty-Related and Neglected Tropical Infectious Diseases, project C3 (PK). The

funders had no role in study design, data collection and analysis, decision to publish, or preparation

of the manuscript. The authors have declared that no competing interests exist.

REFERENCES

1.   Trovato M, Sartorius R, D'Apice L, Manco R, De Berardinis P. Viral Emerging Diseases:

Challenges in Developing Vaccination Strategies. Frontiers in immunology. 2020;11:2130.

2.   Cui J, Li F, Shi ZL. Origin and evolution of pathogenic coronaviruses. Nature reviews

Microbiology. 2019;17(3):181-92.

3.   Douam F, Gaska JM, Winer BY, Ding Q, von Schaewen M, Ploss A. Genetic Dissection

of the Host Tropism of Human-Tropic Pathogens. Annual review of genetics. 2015;49:21-45.

4.   Kurtenbach K, Hanincova K, Tsao JI, Margos G, Fish D, Ogden NH. Fundamental

processes in the evolutionary ecology of Lyme borreliosis. Nature reviews Microbiology.

2006;4(9):660-9.

5.   Longdon B, Brockhurst MA, Russell CA, Welch JJ, Jiggins FM. The evolution and

genetics of virus host shifts. PLoS pathogens. 2014;10(11):e1004395.

6.   Lin YP, Tufts, D.M., Combs, M., Dupuis II, A.P., Marcinkiewicz, A.L., Hirsbrunner, A.D.,

Diaz, A.J., Stout, J.L., Blom, A.M., Strle, K., Davis, A.D., Kramer, L.D., Kolokotronis, S.O., Diuk-

Wasser, M.A. Cellular and immunological mechanisms influence host-adapted phenotypes in a

vector-borne microparasite. Proc Biol Sci. 2022. 289(1969):20212087.

7.   Wollert T, Pasche B, Rochon M, Deppenmeier S, van den Heuvel J, Gruber AD, et al.

Extending the host range of *Listeria monocytogenes* by rational protein design. Cell.

2007;129(5):891-902.

8.   Viana D, Comos M, McAdam PR, Ward MJ, Selva L, Guinane CM, et al. A single natural

nucleotide mutation alters bacterial pathogen host tropism. Nature genetics. 2015;47(4):361-6.

9.   Sabrina Pankey M, Foxall RL, Ster IM, Perry LA, Schuster BM, Donner RA, et al. Host-

selected mutations converging on a global regulator drive an adaptive leap towards symbiosis in

bacteria. Elife. 2017;6.

10.   Radolf JD, Strle K, Lemieux JE, Strle F. Lyme Disease in Humans. Curr Issues Mol Biol.

2021;42:333-84.

11.   Steere AC, Strle F, Wormser GP, Hu LT, Branda JA, Hovius JW, et al. Lyme borreliosis.

Nature reviews Disease primers. 2016;2:16090.

12.   Schwartz I, Margos G, Casjens SR, Qiu WG, Eggers CH. Multipartite Genome of Lyme

Disease *Borrelia*: Structure, Variation and Prophages. Curr Issues Mol Biol. 2021;42:409-54.

13.   Margos G, Gatewood AG, Aanensen DM, Hanincova K, Terekhova D, Vollmer SA, et al.

MLST of housekeeping genes captures geographic population structure and suggests a European

origin of *Borrelia burgdorferi*. Proceedings of the National Academy of Sciences of the United

States of America. 2008;105(25):8730-5.

14.   Wolcott KA, Margos G, Fingerle V, Becker NS. Host association of *Borrelia burgdorferi*

sensu lato: A review. Ticks and tick-borne diseases. 2021;12(5):101766.

15.   Tufts DM, Hart TM, Chen GF, Kolokotronis SO, Diuk-Wasser MA, Lin YP. Outer surface

protein polymorphisms linked to host-spirochete association in Lyme borreliae. Molecular

microbiology. 2019;111(4):868-82.

16.   Skare JT, Garcia BL. Complement Evasion by Lyme Disease Spirochetes. Trends in

microbiology. 2020. 28(11):889-899.

17.   Lin YP, Diuk-Wasser MA, Stevenson B, Kraiczy P. Complement Evasion Contributes to

Lyme Borreliae-Host Associations. Trends in parasitology. 2020;36(7):634-45.

18.   Dulipati V, Meri S, Panelius J. Complement evasion strategies of *Borrelia burgdorferi*

sensu lato. FEBS letters. 2020;594(16):2645-56.

19.   Reis ES, Mastellos DC, Hajishengallis G, Lambris JD. New insights into the immune

functions of complement. Nature reviews Immunology. 2019;19(8):503-16.

20.   Merle NS, Church SE, Fremeaux-Bacchi V, Roumenina LT. Complement System Part I -

Molecular Mechanisms of Activation and Regulation. Frontiers in immunology. 2015;6:262.

21.   Blom AM. The role of complement inhibitors beyond controlling inflammation. Journal of

internal medicine. 2017;282(2):116-28.

22.   Jozsi M. Factor H Family Proteins in Complement Evasion of Microorganisms. Frontiers

in immunology. 2017;8:571.

23.   Najafpour B, Cardoso JCR, Canario AVM, Power DM. Specific Evolution and Gene

Family Expansion of Complement 3 and Regulatory Factor H in Fish. Frontiers in immunology.

2020;11:568631.

24.   Hart T, Nguyen NTT, Nowak NA, Zhang F, Linhardt RJ, Diuk-Wasser M, et al.

Polymorphic factor H-binding activity of CspA protects Lyme borreliae from the host complement

in feeding ticks to facilitate tick-to-host transmission. PLoS pathogens. 2018;14(5):e1007106.

25.   Bykowski T, Woodman ME, Cooley AE, Brissette CA, Brade V, Wallich R, et al.

Coordinated expression of *Borrelia burgdorferi* complement regulator-acquiring surface proteins

during the Lyme disease spirochete's mammal-tick infection cycle. Infection and immunity.

2007;75(9):4227-36.

26.   Hartmann K, Corvey C, Skerka C, Kirschfink M, Karas M, Brade V, et al. Functional

characterization of BbCRASP-2, a distinct outer membrane protein of *Borrelia burgdorferi* that

binds host complement regulators factor H and FHL-1. Molecular microbiology. 2006;61(5):1220-

36.

27.   Rogers EA, Marconi RT. Delineation of species-specific binding properties of the CspZ

protein (BBH06) of Lyme disease spirochetes: evidence for new contributions to the pathogenesis

of *Borrelia* spp. Infection and immunity. 2007;75(11):5272-81.

28.   Rogers EA, Abdunnur SV, McDowell JV, Marconi RT. Comparative analysis of the

properties and ligand binding characteristics of CspZ, a factor H binding protein, derived from

Borrelia burgdorferi isolates of human origin. Infection and immunity. 2009;77(10):4396-405.

29.   Marcinkiewicz AL, Lieknina I, Yang X, Lederman PL, Hart TM, Yates J, et al. The Factor

H-Binding Site of CspZ as a Protective Target against Multistrain, Tick-Transmitted Lyme Disease.

Infection and immunity. 2020;88(5).

30.   Brangulis K, Petrovskis I, Kazaks A, Bogans J, Otikovs M, Jaudzems K, et al. Structural

characterization of CspZ, a complement regulator factor H and FHL-1 binding protein from

Borrelia burgdorferi. The FEBS journal. 2014;281(11):2613-22.

31.   Lin T, Gao L, Zhang C, Odeh E, Jacobs MB, Coutte L, et al. Analysis of an ordered,

comprehensive STM mutant library in infectious *Borrelia burgdorferi*: insights into the genes

required for mouse infectivity. PloS one. 2012;7(10):e47532.

32.   Marcinkiewicz AL, Dupuis AP, 2nd, Zamba-Campero M, Nowak N, Kraiczy P, Ram S, et

al. Blood treatment of Lyme borreliae demonstrates the mechanism of CspZ-mediated complement

evasion to promote systemic infection in vertebrate hosts. Cellular microbiology.

2019;21(2):e12998.

33.   Nakao M, Somamoto, T. The Evolution of Complement Sysstem Functions and Pathways

in Vetebrates. Malagoli D, editor: Academic Press; 2016.

34.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly accurate

protein structure prediction with AlphaFold. Nature. 2021;596(7873):583-9.

35.   Tunyasuvunakool K, Adler J, Wu Z, Green T, Zielinski M, Zidek A, et al. Highly accurate

protein structure prediction for the human proteome. Nature. 2021;596(7873):590-6.

36.   Combs M, Marcinkiewicz AL, Dupuis AP, 2nd, Davis AD, Lederman P, Nowak TA, et al.

Phylogenomic Diversity Elucidates Mechanistic Insights into Lyme Borreliae-Host Association.

mSystems. 2022:e0048822.

37.   Caine JA, Coburn J. A short-term Borrelia burgdorferi infection model identifies tissue

tropisms and bloodstream survival conferred by adhesion proteins. Infection and immunity.

2015;83(8):3184-94.

38.   Ristow LC, Miller HE, Padmore LJ, Chettri R, Salzman N, Caimano MJ, et al. The beta(3)-

integrin ligand of *Borrelia burgdorferi* is critical for infection of mice but not ticks. Molecular

microbiology. 2012;85(6):1105-18.

39.   Frye AM, Hart TM, Tufts DM, Ram S, Diuk-Wasser MA, Kraiczy P, et al. A soft tick

Ornithodoros moubata salivary protein OmCI is a potent inhibitor to prevent avian complement

activation. Ticks and tick-borne diseases. 2020;11(2):101354.

40.   Finnie JA, Stewart RB, Aston WP. A comparison of cobra venom factor-induced depletion

of serum C3 in eight different strains of mice. Dev Comp Immunol. 1981;5(4):697-701.

41.   Marcinkiewicz AL, Lieknina I, Kotelovica S, Yang X, Kraiczy P, Pal U, et al. Eliminating

Factor H-Binding Activity of *Borrelia burgdorferi* CspZ Combined with Virus-Like Particle

Conjugation Enhances Its Efficacy as a Lyme Disease Vaccine. Frontiers in immunology.

2018;9:181.

42.   Walter KS, Carpi G, Caccone A, Diuk-Wasser MA. Genomic insights into the ancient

spread of Lyme disease across North America. Nat Ecol Evol. 2017;1(10):1569-76.

43.   Coburn J, Garcia B, Hu LT, Jewett MW, Kraiczy P, Norris SJ, et al. Lyme Disease

Pathogenesis. Curr Issues Mol Biol. 2021;42:473-518.

44.   Garcia BL, Summers BJ, Lin Z, Ramyar KX, Ricklin D, Kamath DV, et al. Diversity in

the C3b [corrected] contact residues and tertiary structures of the staphylococcal complement

inhibitor (SCIN) protein family. The Journal of biological chemistry. 2012;287(1):628-40.

45.   Hart TM, Dupuis AP, 2nd, Tufts DM, Blom AM, Starkey SR, Rego ROM, et al. Host

tropism determination by convergent evolution of immunological evasion in the Lyme disease

system. PLoS pathogens. 2021;17(7):e1009801.

46.   Lawrenz MB, Wooten RM, Zachary JF, Drouin SM, Weis JJ, Wetsel RA, et al. Effect of

complement component C3 deficiency on experimental Lyme borreliosis in mice. Infection and

immunity. 2003;71(8):4432-40.

47.   Woodman ME, Cooley AE, Miller JC, Lazarus JJ, Tucker K, Bykowski T, et al. *Borrelia*

burgdorferi binding of host complement regulator factor H is not required for efficient mammalian

infection. Infection and immunity. 2007;75(6):3131-9.

48.   Schneider MC, Prosser BE, Caesar JJ, Kugelberg E, Li S, Zhang Q, et al. *Neisseria*

meningitidis recruits factor H using protein mimicry of host carbohydrates. Nature.

2009;458(7240):890-3.

49.   Singh D, Yi SV. On the origin and evolution of SARS-CoV-2. Exp Mol Med.

2021;53(4):537-47.

50.   Xiang Y, White A. Monkeypox virus emerges from the shadow of its more infamous

cousin: family biology matters. Emerging microbes & infections. 2022;11(1):1768-77.

51.   Vishnoi A, Kryazhimskiy S, Bazykin GA, Hannenhalli S, Plotkin JB. Young proteins

experience more variable selection pressures than old proteins. Genome Res. 2010;20(11):1574-

81.

52.   Wang IN, Dykhuizen DE, Qiu W, Dunn JJ, Bosler EM, Luft BJ. Genetic diversity of *ospC*

in a local population of *Borrelia burgdorferi* sensu stricto. Genetics. 1999;151(1):15-30.

53.   Brisson D, Dykhuizen DE. ospC diversity in *Borrelia burgdorferi*: different hosts are

different niches. Genetics. 2004;168(2):713-22.

54.   Haven J, Vargas LC, Mongodin EF, Xue V, Hernandez Y, Pagan P, et al. Pervasive

recombination and sympatric genome diversification driven by frequency-dependent selection in

Borrelia burgdorferi, the Lyme disease bacterium. Genetics. 2011;189(3):951-66.

55.   Haven J, Magori K, Park AW. Ecological and inhost factors promoting distinct parasite

life-history strategies in Lyme borreliosis. Epidemics. 2012;4(3):152-7.

56.   Qiu WG, Martin CL. Evolutionary genomics of *Borrelia burgdorferi* sensu lato: findings,

hypotheses, and the rise of hybrids. Infection, genetics and evolution : journal of molecular

epidemiology and evolutionary genetics in infectious diseases. 2014;27:576-93.

57.   Bown KJ, Lambin X, Ogden NH, Begon M, Telford G, Woldehiwet Z, et al. Delineating

Anaplasma phagocytophilum ecotypes in coexisting, discrete enzootic cycles. Emerging infectious

diseases. 2009;15(12):1948-54.

58.   Ghosh S, O'Connor TJ. Beyond Paralogs: The Multiple Layers of Redundancy in Bacterial

Pathogenesis. Frontiers in cellular and infection microbiology. 2017;7:467.

59.   Charlesworth D. Balancing selection and its effects on sequences in nearby genome regions.

PLoS genetics. 2006;2(4):e64.

60.   Wang G, Ojaimi C, Iyer R, Saksenberg V, McClain SA, Wormser GP, et al. Impact of

genotypic variation of *Borrelia burgdorferi* sensu stricto on kinetics of dissemination and severity

of disease in C3H/HeJ mice. Infection and immunity. 2001;69(7):4303-12.

61.   Chan K, Awan M, Barthold SW, Parveen N. Comparative molecular analyses of *Borrelia*

burgdorferi sensu stricto strains B31 and N40D10/E9 and determination of their pathogenicity.

BMC microbiology. 2012;12:157.

62.   Masuzawa T, Beppu Y, Kawabata H, Yanagihara Y, Iwamoto Y, Shimizu T, et al.

Experimental *Borrelia burgdorferi* infection of outbred mice. Journal of clinical microbiology.

1992;30(11):3016-8.

63.   von Lackum K, Miller JC, Bykowski T, Riley SP, Woodman ME, Brade V, et al. *Borrelia*

burgdorferi regulates expression of complement regulator-acquiring surface protein 1 during the

mammal-tick infection cycle. Infection and immunity. 2005;73(11):7398-405.

64.   Hellwage J, Meri T, Heikkila T, Alitalo A, Panelius J, Lahdenne P, et al. The complement

regulator factor H binds to the surface protein OspE of *Borrelia burgdorferi*. The Journal of

biological chemistry. 2001;276(11):8427-35.

65.   Lin YP, Frye AM, Nowak TA, Kraiczy P. New Insights Into CRASP-Mediated

Complement Evasion in the Lyme Disease Enzootic Cycle. Frontiers in cellular and infection

microbiology. 2020;10:1.

66.   Garcia BL, Zhi H, Wager B, Hook M, Skare JT. *Borrelia burgdorferi* BBK32 Inhibits the

Classical Pathway by Blocking Activation of the C1 Complement Complex. PLoS pathogens.

2016;12(1):e1005404.

67.   Caine JA, Lin YP, Kessler JR, Sato H, Leong JM, Coburn J. *Borrelia burgdorferi* outer

surface protein C (OspC) binds complement component C4b and confers bloodstream survival.

Cellular microbiology. 2017.

68.   Pereira MJ, Wager B, Garrigues RJ, Gerlach E, Quinn JD, Dowdell AS, et al.

Lipoproteome screening of the Lyme disease agent identifies inhibitors of antibody-mediated

complement killing. Proceedings of the National Academy of Sciences of the United States of

America. 2022;119(13):e2117770119.

69.   Tyson KR, Elkins C, de Silva AM. A novel mechanism of complement inhibition

unmasked by a tick salivary protein that binds to properdin. Journal of immunology.

2008;180(6):3964-8.

70.   Schuijt TJ, Coumou J, Narasimhan S, Dai J, Deponte K, Wouters D, et al. A tick mannose-

binding lectin inhibitor interferes with the vertebrate complement cascade to enhance transmission

of the lyme disease agent. Cell host & microbe. 2011;10(2):136-46.

71.   Schuijt TJ, Hovius JW, van Burgel ND, Ramamoorthi N, Fikrig E, van Dam AP. The tick

salivary protein Salp15 inhibits the killing of serum-sensitive *Borrelia burgdorferi* sensu lato

isolates. Infection and immunity. 2008;76(7):2888-94.

72.   Bockenstedt LK, Wooten RM, Baumgarth N. Immune Response to *Borrelia*: Lessons from

Lyme Disease Spirochetes. Curr Issues Mol Biol. 2021;42:145-90.

73.   Lin YP, Benoit V, Yang X, Martinez-Herranz R, Pal U, Leong JM. Strain-specific variation

of the decorin-binding adhesin DbpA influences the tissue tropism of the lyme disease spirochete.

PLoS pathogens. 2014;10(7):e1004238.

74.   Lin YP, Tan X, Caine JA, Castellanos M, Chaconas G, Coburn J, et al. Strain-specific joint

invasion and colonization by Lyme disease spirochetes is promoted by outer surface protein C.

PLoS pathogens. 2020;16(5):e1008516.

75.   Barthold SW, Persing DH, Armstrong AL, Peeples RA. Kinetics of *Borrelia burgdorferi*

dissemination and evolution of disease after intradermal inoculation of mice. The American

journal of pathology. 1991;139(2):263-73.

76.   Hanincova K, Ogden NH, Diuk-Wasser M, Pappas CJ, Iyer R, Fish D, et al. Fitness

variation of *Borrelia burgdorferi* sensu stricto strains in mice. Applied and environmental

microbiology. 2008;74(1):153-7.

77.   Lawrenz MB, Hardham JM, Owens RT, Nowakowski J, Steere AC, Wormser GP, et al.

Human antibody responses to VlsE antigenic variation protein of *Borrelia burgdorferi*. Journal of

clinical microbiology. 1999;37(12):3997-4004.

78.   Casjens SR, Gilcrease EB, Vujadinovic M, Mongodin EF, Luft BJ, Schutzer SE, et al.

Plasmid diversity and phylogenetic consistency in the Lyme disease agent *Borrelia burgdorferi*.

BMC Genomics. 2017;18(1):165.

79.   Petersen TN, Brunak S, von Heijne G, Nielsen H. SignalP 4.0: discriminating signal

peptides from transmembrane regions. Nature methods. 2011;8(10):785-6.

80.   Chiu J, March PE, Lee R, Tillett D. Site-directed, Ligase-Independent Mutagenesis (SLIM):

a single-tube methodology approaching 100% efficiency in 4 h. Nucleic acids research.

2004;32(21):e174.

81.   Mueller U, Darowski N, Fuchs MR, Forster R, Hellmig M, Paithankar KS, et al. Facilities

for macromolecular crystallography at the Helmholtz-Zentrum Berlin. J Synchrotron Radiat.

2012;19(Pt 3):442-9.

82.   Winn MD, Ballard CC, Cowtan KD, Dodson EJ, Emsley P, Evans PR, et al. Overview of

the CCP4 suite and current developments. Acta crystallographica Section D, Biological

crystallography. 2011;67(Pt 4):235-42.

83.   Evans PR. An introduction to data reduction: space-group determination, scaling and

intensity statistics. Acta crystallographica Section D, Biological crystallography. 2011;67(Pt

4):282-92.

84.   Kabsch W. Xds. Acta crystallographica Section D, Biological crystallography. 2010;66(Pt

2):125-32.

85. McCoy AJ, Grosse-Kunstleve RW, Adams PD, Winn MD, Storoni LC, Read RJ. Phaser

crystallographic software. J Appl Crystallogr. 2007;40(Pt 4):658-74.

86. Cowtan K. The Buccaneer software for automated model building. 1. Tracing protein

chains. Acta crystallographica Section D, Biological crystallography. 2006;62(Pt 9):1002-11.

87. Emsley P, Cowtan K. Coot: model-building tools for molecular graphics. Acta

crystallographica Section D, Biological crystallography. 2004;60(Pt 12 Pt 1):2126-32.

88. Murshudov GN, Vagin AA, Dodson EJ. Refinement of macromolecular structures by the

maximum-likelihood method. Acta crystallographica Section D, Biological crystallography.

1997;53(Pt 3):240-55.

89. Kelly SM, Jess TJ, Price NC. How to study proteins by circular dichroism. Biochimica et

biophysica acta. 2005;1751(2):119-39.

90. Lin YP, Greenwood A, Nicholson LK, Sharma Y, McDonough SP, Chang YF. Fibronectin

binds to and induces conformational change in a disordered region of leptospiral immunoglobulin-

like protein B. The Journal of biological chemistry. 2009;284(35):23547-57.

91. Purser JE, Norris SJ. Correlation between plasmid content and infectivity in *Borrelia*

burgdorferi. Proceedings of the National Academy of Sciences of the United States of America.

2000;97(25):13865-70.

92. Hart T, Yang X, Pal U, Lin YP. Identification of Lyme borreliae proteins promoting

vertebrate host blood-specific spirochete survival in *Ixodes scapularis* nymphs using artificial

feeding chambers. Ticks and tick-borne diseases. 2018.

93. Rose R, Golosova O, Sukhomlinov D, Tiunov A, Prosperi M. Flexible design of multiple

metagenomics classification pipelines with UGENE. Bioinformatics. 2019;35(11):1963-5.

94.   Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

arXiv. 2013;1303:3997.

95.   Abascal F, Zardoya R, Telford MJ. TranslatorX: multiple alignment of nucleotide

sequences guided by amino acid translations. Nucleic acids research. 2010;38(Web Server

issue):W7-13.

96.   Huson DH, Bryant D. Application of phylogenetic networks in evolutionary studies. Mol

Biol Evol. 2006;23(2):254-67.

97.   Villesen P. FaBox: an online toolbox for fasta sequences. Mol Ecol Notes. 2007;7(6):965-

8.

98.   Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective

stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol.

2015;32(1):268-74.

99.   Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. ModelFinder:

fast model selection for accurate phylogenetic estimates. Nature methods. 2017;14(6):587-9.

100.   Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. UFBoot2: Improving the

Ultrafast Bootstrap Approximation. Mol Biol Evol. 2018;35(2):518-22.

101.   Letunic I, Bork P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree

display and annotation. Nucleic acids research. 2021;49(W1):W293-W6.

102.   Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular Evolutionary

Genetics Analysis across Computing Platforms. Mol Biol Evol. 2018;35(6):1547-9.

103.   Kosakovsky Pond SL, Posada D, Gravenor MB, Woelk CH, Frost SD. Automated

phylogenetic detection of recombination using a genetic algorithm. Mol Biol Evol.

2006;23(10):1891-901.

104.   Murrell B, Moola S, Mabona A, Weighill T, Sheward D, Kosakovsky Pond SL, et al.

FUBAR: a fast, unconstrained bayesian approximation for inferring selection. Mol Biol Evol.

2013;30(5):1196-205.

105.   Kosakovsky Pond SL, Frost SD. Not so different after all: a comparison of methods for

detecting amino acid sites under selection. Mol Biol Evol. 2005;22(5):1208-22.

106.   Murrell B, Wertheim JO, Moola S, Weighill T, Scheffler K, Kosakovsky Pond SL.

Detecting individual sites subject to episodic diversifying selection. PLoS genetics.

2012;8(7):e1002764.

107.   Weaver S, Shank SD, Spielman SJ, Li M, Muse SV, Kosakovsky Pond SL. Datamonkey

2.0: A Modern Web Application for Characterizing Selective and Other Evolutionary Processes.

Mol Biol Evol. 2018;35(3):773-7.

108.   Librado P, Rozas J. DnaSP v5: a software for comprehensive analysis of DNA

polymorphism data. Bioinformatics. 2009;25(11):1451-2.

109.   Le SQ, Gascuel O. An improved general amino acid replacement matrix. Mol Biol Evol.

2008;25(7):1307-20.

110.   Ross CM, Foley G, Boden M, Gillam EMJ. Using the Evolutionary History of Proteins to

Engineer Insertion-Deletion Mutants from Robust, Ancestral Templates Using Graphical

Representation of Ancestral Sequence Predictions (GRASP). Methods in molecular biology.

2022;2397:85-110.

111.   Musil M, Khan RT, Beier A, Stourac J, Konegger H, Damborsky J, et al. FireProtASR: A

Web Server for Fully Automated Ancestral Sequence Reconstruction. Brief Bioinform. 2021;22(4).

112.   Suchard MA, Lemey P, Baele G, Ayres DL, Drummond AJ, Rambaut A. Bayesian

phylogenetic and phylodynamic data integration using BEAST 1.10. Virus Evol.

2018;4(1):vey016.

113.   Hasegawa M, Kishino H, Yano T. Dating of the human-ape splitting by a molecular clock

of mitochondrial DNA. J Mol Evol. 1985;22(2):160-74.

114.   Yang Z. Maximum likelihood phylogenetic estimation from DNA sequences with variable

rates over sites: approximate methods. J Mol Evol. 1994;39(3):306-14.

115.   Robert X, Gouet P. Deciphering key features in protein structures with the new ENDscript

server. Nucleic acids research. 2014;42(Web Server issue):W320-4.

116.   Waterhouse AM, Procter JB, Martin DM, Clamp M, Barton GJ. Jalview Version 2--a

multiple sequence alignment editor and analysis workbench. Bioinformatics. 2009;25(9):1189-91.

117.   Benjamini YK, A. M. Yekutieli, D. . Adaptive linear step-up procedures that control the

false discovery rate. Biometrika. 2006;93:491-507.

118.   Kraiczy P, Skerka C, Brade V, Zipfel PF. Further characterization of complement

regulator-acquiring surface proteins of *Borrelia burgdorferi*. Infection and immunity.

2001;69(12):7800-9.

119.   Kraiczy P, Skerka C, Kirschfink M, Brade V, Zipfel PF. Immune evasion of *Borrelia*

burgdorferi by acquisition of human complement regulators FHL-1/reconectin and Factor H.

European journal of immunology. 2001;31(6):1674-84.

120.   Hallstrom T, Siegel C, Morgelin M, Kraiczy P, Skerka C, Zipfel PF. CspA from *Borrelia*

burgdorferi inhibits the terminal complement pathway. mBio. 2013; 4(4):e00481-13.

121.   Elias AF, Stewart PE, Grimm D, Caimano MJ, Eggers CH, Tilly K, et al. Clonal

polymorphism of *Borrelia burgdorferi* strain B31 MI: implications for mutagenesis in an

infectious strain background. Infection and immunity. 2002;70(4):2139-50.

122.   Coleman AS, Yang X, Kumar M, Zhang X, Promnares K, Shroder D, et al. *Borrelia*

burgdorferi complement regulator-acquiring surface protein 2 does not contribute to complement

resistance or host infectivity. PloS one. 2008;3(8):3010e.

123.   Benoit VM, Fischer JR, Lin YP, Parveen N, Leong JM. Allelic variation of the Lyme

disease spirochete adhesin DbpA influences spirochetal binding to decorin, dermatan sulfate, and

mammalian cells. Infection and immunity. 2011;79(9):3501-9.

124.   Frank KL, Bundle SF, Kresge ME, Eggers CH, Samuels DS. *aadA* confers streptomycin

resistance in Borrelia burgdorferi. Journal of bacteriology. 2003;185(22):6723-7.

SUPPLEMENTAL TABLES

Table S1: The list of CspZ$_{B408}$ residues that binds to human FH based on CspZB408-human

SCR6-7 complex structure and their equivalent residues of CspZ$_{B31}$ and CspZ$_{B379}$.

CspZ$_{B408}$	CspZ$_{B31}$	CspZ$_{B379}$
Asp47	Asp47	Asp47
Tyr50	Tyr50	Tyr50
Asn51	Asn51	Ser51
Thr54	Thr54	Thr54
Asn58	Asn58	Asn58
Thr62	Thr62	Thr62
Asp71	-	-
Asp73[a]	Asp70[a]	Asp74[a]
Ser75[a]	Ser72[a]	Ser76[a]
Arg142	Arg139	Arg143
Asn183	Asn180	Asn184
Tyr214	Tyr211	Tyr215

973 [a]Interacts with SCR6.

975 **Table S2**. CspZ variants differ in binding to Factor H from different animals

CspZ variant	Factor H source	ELISA[a] K_D (μM)	----- Surface Plasmon Resonance[b] ----- K_D (μM)	k_{on} ($10^3 s^{-1} M^{-1}$)	k_{off} (s^{-1})
CspZ$_{B31}$	Mouse	0.43±0.74[c]	0.20±0.02	45.78±15.36	0.0084±0.0022
	Quail	0.91±0.08[d]	0.81±0.01	117.43±3.25	0.095±0.0032
CspZ$_{B379}$	Mouse	n.b.[d]	n.b.	n.b.	n.b.
	Quail	0.59±0.037	0.75±0.15	119.83±7.08	0.088±0.012
CspZ$_{B379}$L$_{B408}$	Mouse	1.39±0.12	0.90±0.12	21.16±10.58	0.018±0.0094
	Quail	n.b.	n.b.	n.b.	n.b.
CspZ$_{B408}$	Mouse	0.68±0.05	0.20±0.02	22.13±1.31	0.008±0.002
	Quail	n.b.	n.b.	n.b.	n.b.
CspZ$_{B408}$L$_{B379}$	Mouse	n.b.	n.b.	n.b.	n.b.
	Quail	1.38±0.47	0.99±0.06	72.43±2.14	0.072±0.006
GST[e]	Mouse	n.b.	n.d.	n.d.	n.d.
	Quail	n.b.	n.d.	n.d.	n.d.

All values represent the mean ± SEM of three experiments.
[a]Determined using GST tagged CspZ variants or mutant proteins.
[b]Determined using untagged CspZ variants or mutant proteins.
[c]Reported previously in (41).
[d]No binding activity was detected.
[e]GST was included as a negative control.

Table S3. The generation time for *B. burgdorferi* strains used in this study.

Strain	Generation time[a, b]
B31-A3	16.45 ± 1.38
B31-A3$\Delta cspZ$/vector	16.42 ± 1.79
B31-A3$\Delta cspZ$/pCspZ$_{B31}$	17.12 ± 1.05
B31-A3$\Delta cspZ$/pCspZ$_{B379}$	16.62 ± 2.07
B31-A3$\Delta cspZ$/pCspZ$_{B379}$L$_{B408}$	16.40 ± 0.83
B31-A3$\Delta cspZ$/pCspZ$_{B408}$	17.31 ± 0.62
B31-A3$\Delta cspZ$/pCspZ$_{B408}$L$_{B379}$	18.66 ± 2.06

983 [a]The generation time was calculated as described previously (18)
984 [b]There were no significant differences between generation time of any strain (Kruskal-Wallis test with the two-stage step-up
985 method of Benjamini, Krieger, and Yekutieli)
986

987   **Table S4: The percentage of SAPs in CspZ variants**

Loop type	Position	AA	%	AA	%
CspZ$_{B408}$	20	Asp	64.00	Asn	36.00
	30	Asp	88.00	Asn	36.00
	41	Val	56.00	Phe	44.00
	84	Phe	52.00	Leu	48.00
	95	Lys	92.00	Asn	8.00
	107	Met	56.00	Ile	44.00
	183	Asp	52.00	Asn	48.00
CspZ$_{B31}$	30	Asn	88.30	Asp	11.70
	66[a]	Gly	80.85	Val	19.15
	68[a]	Phe	80.85	Tyr	19.15
	81	Phe	88.30	Leu	11.70
	88	Val	98.94	Ala	1.06
	131	Val	94.68	Ala	5.32
	154	Ser	98.64	Pro	1.06
	161	Lys	96.81	Glu	3.19
	203	Ser	98.94	Lys	1.06
	204	Arg	98.94	Leu	1.06
	208	Asn	67.02	Asp	32.98
CspZ$_{B379}$	30	Asp	87.50	Asn	12.50
	50[b]	Tyr	97.92	His	2.08
	235	Ile	97.92	Ser	2.08

988   [a]Part of the loop structures

989   [b]Predicted to directly interact with FH

991   **Table S5. Estimated diversification times for each lineage**.

Lineage	Median[a]	HPD$_{5\%}$[a]	HPD$_{95\%}$[a]
CspZ-B31	784	263	1860
CspZ-B379	261	59	741
CspZ-B408	671	179	1679

992   [a]Years before present

994 **Table S6**. **CspZ variants of last common ancestral states display versatile binding ability to**
995 **human, mouse, and quail FH**

CspZ variant	Factor H source	K_D (μM)[a]
CspZ$_{B31}$	Human	0.31±0.03
	Mouse	0.38±0.07
	Quail	0.74±0.15
CspZ-LCAS1	Human	0.34±0.10
	Mouse	0.41±0.07
	Quail	0.77±0.09
CspZ-LCAS2	Human	0.23±0.02
	Mouse	0.36±0.05
	Quail	0.94±0.09
CspZ-LCAS3	Human	0.25±0.02
	Mouse	0.53±0.12
	Quail	0.73±0.06
CspZ-LCAS4	Human	0.30±0.01
	Mouse	0.40±0.01
	Quail	0.93±0.06
DbpA[b]	Human	n.b.[c]
	Mouse	n.b.
	Quail	n.b.

996 All values represent the mean ± SEM of three experiments.
997 [a]Determined using histidine tagged CspZ variants or mutant proteins.
998 [b]DbpA was included as a negative control
999 [c]No binding activity was detected.

 Table S7. The strains and plasmids used in this study.

Strain or plasmid	Genotype or characteristic	Source
B. burgdorferi		
B313	High-passage *B. burgdorferi* B31 missing lp5, lp17, lp21, lp25, lp28-1, lp28-2, lp28-3, lp28-4, lp36, lp38, lp54, lp56, cp9, cp32-4, cp32-6, cp32-8, cp32-9	[120]
B31-A3	Clone of *B. burgdorferi* B31 missing cp9 RST type 1, *ospC* type A	[121]
B379	Clone of *B. burgdorferi* B379 isolated from humans with erythema migrans. RST Type 2, *ospC* type K	[21]
B408	Clone of *B. burgdorferi* B379 isolated from humans with erythema migrans. RST type 3, *ospC* type K	[21]
B31-A3Δ*cspZ*	B31-A3Δ*cspZ*::KanR[a]	[122]
B31-A3Δ*cspZ*/Vector	B31-A3Δ*cspZ*::KanR carrying plasmid pKFSS	[32]
B31-A3Δ*cspZ*/pCspZ$_{B31}$	B31-A3Δ*cspZ*::KanR complemented with intact *cspZ* driven by the promoter of *cspZ$_{B31}$*	[32]
B31-A3Δ*cspZ*/pCspZ$_{B379}$	B31-A3Δ*cspZ*::KanR complemented with intact *cspZ$_{B379}$* driven by the promoter of *cspZ$_{B31}$*	This study
B31-A3Δ*cspZ*/pCspZ$_{B408}$	B31-A3Δ*cspZ*::KanR complemented with intact *cspZ$_{B408}$* driven by the promoter of *cspZ$_{B31}$*	This study
B31-A3Δ*cspZ*/pCspZ$_{B379}$L$_{B408}$	B31-A3Δ*cspZ*::KanR complemented with intact *cspZ $_{B379}$* except the residues 190 to 216 replaced by residues 190 to 213 from *cspZ$_{B408}$* ,driven by the promoter of *cspZ$_{B31}$*	This study

B31-A3Δ*cspZ*/pCspZ$_{B408}$L$_{B379}$	B31-A3Δ*cspZ*::KanR complemented with intact *cspZ $_{B408}$* except the residues 190 to 213 replaced by residues 190 to 216 from *cspZ$_{B379}$*, driven by the promoter of *cspZ$_{B31}$*	This study
E. coli		
DH5α	F- Φ80lacZΔM15 Δ(lacZYA-argF) U169 recA1 endA1 hsdR17(rk-, mk+) phoA supE44 thi-1 gyrA96 relA1 λ-	ThermoFisher
BL21(DE3)	F−, ompT hsdSB (rB− mB−) gal dcm (DE3)	Novagen
BL21(DE3)/pGEX4T2-CspZ$_{B31}$	BL21(DE3) producing GST-tagged residues 58 to 711 of CspZ$_{B31}$	(32)
BL21(DE3)/pGEX4T2-CspZ$_{B379}$	BL21(DE3) producing GST-tagged residues 58 to 723 of CspZ$_{B379}$	This study
BL21(DE3)/pGEX4T2-CspZ$_{B408}$	BL21(DE3) producing GST-tagged residues 58 to 720 of CspZ$_{B408}$	This study
BL21(DE3)/pGEX4T2-CspZ$_{B379}$L$_{B408}$	BL21(DE3) producing GST-tagged residues 58 to 723 of CspZ$_{B379}$ except the residues 190 to 216 replaced by residues 190 to 213 from CspZ$_{B408}$	This study
BL21(DE3)/pGEX4T2-CspZ$_{B408}$L$_{B379}$	BL21(DE3) producing GST-tagged residues 58 to 720 of CspZ$_{B408}$ except the residues 190 to 213 replaced by residues 190 to 216 from CspZ$_{B379}$	This study
BL21(DE3)/pET15b-DbpA$_{VS461}$	BL21(DE3) producing histidine-tagged residues 22 to 170 of DbpA$_{VS461}$	(123)
BL21(DE3)/pET28a-CspZ-LCAS1	BL21(DE3 producing histidine tagged residue 58 to 711 of CspZ-LCAS1	This study
BL21(DE3)/pET28a-CspZ-LCAS2	BL21(DE3 producing histidine tagged residue 58 to 711 of CspZ-LCAS2	This study

BL21(DE3)/pET28a-CspZ-LCAS3	BL21(DE3 producing histidine tagged residue 58 to 711 of CspZ-LCAS3	This study
BL21(DE3)/pET28a-CspZ-LCAS4	BL21(DE3 producing histidine tagged residue 58 to 711 of CspZ-LCAS4	This study
Rosetta-gami(DE3)	F– ompT hsdSB (rB– mB–) gal dcm lacY1 ahpC (DE3) gor522::Tn10 trxB pRARE (CamR, KanR, TetR)	MilliporeSigma

P. pastoris

X-33/FH SCR6-7	Wild-type Mut+ Pichia strain for expression of FH SCR6-7 (residues 321 to 444 of human FH)	Invitrogen

Plasmids

pJET1.2/Blunt	AmpR[a]; PCR cloning vector	ThermoFisher
pGEX4T2	AmpR; GST-tagged protein expression vector	Qiagen
pGEX4T2-CspZ	AmpR; pGEX4T2 encoding GST fusion protein residue 58 to 711 of CspZ	(32)
pGEX4T2-CspZ- $CspZ_{B379}$	AmpR; pGEX4T2 encoding GST fusion protein residue 58 to 723 of $CspZ_{B379}$	This study
pGEX4T2-CspZ- $CspZ_{B408}$	AmpR; pGEX4T2 encoding GST fusion protein residue 58 to 720 of $CspZ_{B408}$	This study
pGEX4T2-CspZ- $CspZ_{B379}L_{B408}$	AmpR; pGEX4T2 encoding GST fusion protein residue 58 to 723 of $CspZ_{B379}$ except the residues 190 to 216 replaced by residues 190 to 213 from $CspZ_{B408}$	This study
pGEX4T2-CspZ- $CspZ_{B408}L_{B379}$	AmpR; pGEX4T2 encoding GST fusion protein residue 58 to 720 of $CspZ_{B408}$ except the residues 190 to 213 replaced by residues 190 to 216 from $CspZ_{B379}$	This study

pET28a	KanR[b]; Histidine-tagged protein expression vector	EMD Millipore
pET28a-CspZ-LCAS1	KanR; pET28a encoding histidine protein residues 58 to 711 of the CspZ-LCAS1	This study
pET28a-CspZ-LCAS2	KanR; pET28a encoding histidine protein residues 58 to 711 of the CspZ-LCAS2	This study
pET28a-CspZ-LCAS3	KanR; pET28a encoding histidine protein residues 58 to 711 of the CspZ-LCAS3	This study
pET28a-CspZ-LCAS4	KanR; pET28a encoding histidine protein residues 58 to 711 of the CspZ-LCAS4	This study
pKFSS-1	StrR[c]; *Borrelia* shuttle vector	(124)
pKFSS/pCspZ$_{B31}$	StrR; pKFSS-1 encoding intact *cspZ$_{B31}$*, driven by the promoter of *cspZ$_{B31}$*	(32)
pKFSS/pCspZ$_{B379}$	StrR; pKFSS-1 encoding intact *cspZ$_{B379}$*, driven by the promoter of *cspZ$_{B31}$*	This study
pKFSS/pCspZ$_{B408}$	StrR; pKFSS-1 encoding intact *cspZ$_{B408}$*, driven by the promoter of *cspZ$_{B31}$*	This study
pKFSS/pCspZ$_{B379}$L$_{B408}$	StrR; pKFSS-1 encoding intact *cspZ$_{B379}$* except the residues 190 to 216 replaced by residues 190 to 213 from *cspZ$_{B408}$*, driven by the promoter of *cspZ$_{B31}$*	This study
pKFSS/pCspZ$_{B408}$L$_{B379}$	StrR; pKFSS-1 encoding intact *cspZ$_{B408}$* except the residues 190 to 213 replaced by residues 190 to 216 from *cspZ$_{B379}$*, driven by the promoter of *cspZ$_{B31}$*	This study
pETm-11/pCspZ$_{B379}$	KanR; 6xHis tag expression vector encoding CspZ$_{B379}$ residues 23-236	This study
pETm-11/pCspZ$_{B408}$	KanR; 6xHis tag expression vector encoding CspZ$_{B408}$ residues 23-236	This study
pPICZα-FH		This study

Zeocin resistant; yeast expression vector
encoding SCR6-7 of human FH under
αMF secretion signal

1001   [a]Ampicillin resistant

1002   [b]Kanamycin resistant

1003   [c]Streptomycin resistant

1004

1005 **Table S8. Primers used in this study.**

Purpose	Primer	Sequence
qPCR spirochete burden: *recA*	BBRecAfp	GTGGATCTATTGTATTAGATGAGGCTCTCG
	BBRecArp	CAGCAACATGTCTGGCATTAGACAC
Generate B379LB408: SLIM step 1	B379LB408_A-1_mtsengtd	CAGAAGATGTGTTACCTTCCGAAGTCATATAAGTCATAATATCATTATATGCTCCTGTA
	B379LB408_B-1_mtsengtd	TATGACTTATATGACTTCGGAAGGTAACACATCTTCTGATAAAAGTAAGGTTAATCAAG
	B379LB408_C-1_mtsengtd	ATATCATTATATGCTCCTGTA
	B379LB408_D-1_mtsengtd	ATAAAAGTAAGGTTAATCAAG
Generate B379LB408: SLIM step 2	B379LB408_A-2_tsenngtd	TTTTATCAGAATCTGTGTTATTACCTTCCGAAGTATAAGTCATAATATCATTATATGCTCCTGTA
	B379LB408_B-2_tsenngtd	TATGACTTATACTTCGGAAGGTAATAACACAGATTCTGATAAAAGTAAGGTTAATCAAGCTATAT
	B379LB408_C-2_tsenngtd	ATATCATTATATGCTCCTGTA
	B379LB408_D-2_tsenngtd	GTAAGGTTAATCAAGCTATAT
Generate B379LB408: SLIM step 3	B379LB408_A-3_sevnnvtd	CAGAATCTGTAACGTTATTAACTTCCGAATAAGTCATAATATCATTATATGCTCCTGTA
	B379LB408_B-3_sevnnvtd	TATGACTTATTCGGAAGTTAATAACGTTACAGATTCTGATAAAAGTAAGGTTAATCAAG
	B379LB408_C-3_sevnnvtd	ATATCATTATATGCTCCTGTA
	B379LB408_D-3_sevnnvtd	ATAAAAGTAAGGTTAATCAAG
Generate B408LB379: SLIM step 1	B408LB379_A-1_tsenngtd	CAGAATCTGTACCGTTATTTTCCGAAGTATAAGTCATAATATCATTATATGCTTCTGTA
	B408LB379_B-1_tsenngtd	TATGACTTATACTTCGGAAAATAACGGTACAGATTCTGATAAAAGTAAGGTTAATCAAG
	B408LB379_C-1_tsenngtd	ATATCATTATATGCTTCTGTA
	B408LB379_D-1_tsenngtd	TAAAAGTAAGGTTAATCAAG
Generate B408LB379: SLIM step 2	B408LB379_A-2_mtsengts	TTTTATCAGAAGATGTACCGTTTTCCGAAGTCATATAAGTCATAATATCATTATATGCTTCTGTA

	B408LB379_B-2_mtsengts	TATGACTTATATGACTTCGGAAAACGGTACATCTTCTGATAAA AGTAAGGTTAATCAAGCTATAT
	B408LB379_C-2_mtsengts	ATATCATTATATGCTTCTGTA
	B408LB379_D-2_mtsengts	GTAAGGTTAATCAAGCTATAT
Generate B408LB379: SLIM step 3	B408LB379_A-3_imtysegts	AAGATGTACCTTCCGAATAAGTCATAATATAAGTCATAATATC ATTATATGCTTCTGTA
	B408LB379_B-3_imtysegts	TATGACTTATATTATGACTTATTCGGAAGGTACATCTTCTGATA AAAGTAAGGTTAATC
	B408LB379_C-3_imtysegts	ATATCATTATATGCTTCTGTA
	B408LB379_D-3_imtysegts	CTGATAAAAGTAAGGTTAATC
Generate CspZB379 residues 23-236: structural studies	*cspZB379* Forw.	CATGCCATGGGCAGATTAAATCAGAGAAAT
	cspZB379 Rev.	GCTTGCGGCCGCTTATAATAAAGTTTGCTTAAT
Generate CspZB408 residues 23-236: structural studies	*cspZB408* Forw.	CATGCCATGGGCAGATTAAATCAGAGAAAT
	cspZB408 Rev.	GCTTGCGGCCGCTTATAATAAAGTTTGCTTAAT

1006

 Table S9. Data processing, refinement, and validation statistics of crystal structures.

Dataset	CspZ$_{B408}$	CspZ$_{B379}$	CspZ$_{B408}$-SCR6-7
X-ray diffraction data			
PDB entry	7ZJK	7ZJJ	7ZJM
Beamline	BESSY II beamline 14.1	BESSY II beamline 14.1	BESSY II beamline 14.1
Space group	P2$_1$	P22$_1$2$_1$	P2$_1$2$_1$2$_1$
a, b, c (Å)	47.63, 87.49, 48.86	53.60, 59.95, 61.05	42.10, 71.07, 147.75
α, β, γ (°)	90.0, 97.1, 90.0	90.0, 90.0, 90.0	90.0, 90.0, 90.0
Wavelength (Å)	0.9798	0.9762	0.9798
Resolution (Å)	48.49-2.45	61.05-2.10	73.87-2.59
Highest resolution bin (Å)	2.55-2.45	2.16-2.10	2.65-2.59
No. of reflections	97062	127094	101717
No. of unique reflections	14406	11805	14404
Completeness (%)	98.2 (88.4)[a]	98.9 (99.5)	99.8 (99.8)
R$_{merge}$	0.09 (0.38)	0.10 (0.35)	0.11 (0.38)
CC$_{1/2}$	0.997 (0.940)	0.998 (0.985)	0.988 (0.926)
I/σ (I)	12.1 (4.2)	14.8 (6.3)	12.8 (4.9)
Multiplicity	6.7 (6.2)	10.8 (11.1)	7.1 (7.5)
Refinement			
R$_{work}$	0.193 (0.248)	0.208 (0.371)	0.217 (0.270)
R$_{free}$	0.262 (0.399)	0.262 (0.431)	0.275 (0.354)
Average B-factor (Å^2)			
Overall	45.1	33.0	31.9
From Wilson plot	36.9	15.9	24.1
No. of atoms			
Protein	3592	1769	1914
RMS deviations from ideal			
Bond lengths (Å)	0.007	0.008	0.010
Bond angles (°)	1.435	1.541	1.548
Ramachandran outliers (%)			
Residues in most favored regions (%)	93.74	95.31	94.93
Residues in allowed regions (%)	5.10	4.69	4.17

Outliers (%)	1.16	0.00	0.90

1030 **FIGURES**

1031

Figure 1. Superimposed crystal structures of CspZ_{B408}-SCR6-7 complex, CspZ_{B31}, and

CspZ_{B379} reveals the FH-binding mechanisms of CspZ variants. (**A**) The crystal structure of

human FH domains SCR6 (light grey) and SCR7 (dark grey) in complex with CspZ_{B408} (orange)

was superimposed with that of CspZ_{B31} (purple) and CspZ_{B379} (green). (**B**) The FH-binding

interface with the location of the CspZ_{B408} residues involved in human SCR6-7 binding, and the

equvalent residues in CspZ_{B31} and CspZ_{B379}. The numbering of the residues is given for CspZ_{B408}.

α-helices in CspZ are labelled from A to I starting from the N-terminus. The polymorphic loop

region between α-helices B and C in CspZ$_{B408}$ (residues 60-IMTYSEVNNVTD-71), CspZ$_{B31}$

(residues 60-IMTYSEGT-67) and CspZ$_{B379}$ (residues 60-IMTYIMTYSEGT-71) are indicated.

1041

1042

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061

Figure 2. The polymorphic loop region in CspZ promotes host-specific FH-binding ability.

(A, B) Ten micrograms of mouse or quail FH were conjugated on a SPR chip. 0.008 to 2μM of

untagged **(A)** CspZ$_{B31}$, CspZ$_{B379}$, CspZ$_{B408}$, **(B)** CspZ$_{B379}$L$_{B408}$, or CspZ$_{B408}$L$_{B379}$ was flowed over

the chip surface. Binding was measured in response units (RU) by SPR. The experiments were performed with a single preparation of recombinant proteins tested in three independent replicates with samples ran in duplicate. The k_{on}, k_{off}, and K_D values were determined from the average of these three experiments (**Table 1**). Shown is one representative experiment. (**C**) The crystal structure of CspZ$_{B408}$-SCR6-7 where human SCR6-7 (grey) is superimposed with mouse SCR6-7 (gold, PDB: 2YBY), and the predicted structure of quail SCR6-7 (brown), while CspZ$_{B408}$ (orange) is superimposed with CspZ$_{B31}$ (purple) and CspZ$_{B379}$ (green). (**D**) The loop region in CspZ proteins located between α-helices B and C. The Asp71 in CspZ$_{B408}$, and residues 60-IMTYIMTYSEGT-71 in CspZ$_{B379}$ are highlighted. The polymorphic loop region between α-helices B and C in CspZ variants, as well as the loop region in human, mouse and quail SCR6-7 located at the interface with helix-B in CspZ, is showed with a striped pattern.

1086

Figure 3. The CspZ loop-driven, variable FH-binding activity confers host-specific

complement inactivation on spirochete surface and serum resistance. (A to D) *B. burgdorferi*

strains B313 (negative control), B31-A3, B31-A3Δ*cspZ* harboring the empty vector pKFSS

("Δ*cspZ*/Vector"), or this *cspZ* mutant producing CspZ$_{B31}$, CspZ$_{B379}$, CspZ$_{B379}$L$_{B408}$, CspA$_{B408}$, or

CspZ$_{B408}$L$_{B379}$, were incubated with mouse or quail sera at a final concentration of 20%. The

bacteria were stained with antibodies that recognize mouse C5b-9 or quail C8 prior to analysis by

flow cytometry. Shown are the representative histograms of the analysis presenting the deposition

levels of **(A)** mouse C5b-9 or **(B)** quail C8 on the surface of the indicated *B. burgdorferi* strains.

The deposition levels of **(C)** mouse C5b-9 or **(D)** quail C8 on the surface of the indicated strains

were measured by flow cytometry and presented as mean fluorescence index (MFI). Each bar

represents the mean of three independent experiments ± SEM. Significant differences ($p < 0.05$,

Kruskal-Wallis test with the two-stage step-up method of Benjamini, Krieger, and Yekutieli) in

the deposition levels of mouse C5b-9 or quail C8 relative to B313 ("Φ"), Δ*cspZ*/Vector ("*"), or

between two strains relative to each other ("#"), are indicated. **(E, F)** The *B. burgdorferi* strains

were incubated for 4-h with **(E)** white-footed mouse sera or **(F)** quail sera, to a final concentration

of 40%. The number of motile spirochetes was assessed microscopically. The precent survival of

the strains was calculated using the number of motile spirochetes at 4-h post incubation normalized

to that prior to incubation with sera. Each bar represents the mean of three independent experiments

± SEM. Significant differences ($p < 0.05$, Kruskal-Wallis test with the two-stage step-up method

of Benjamini, Krieger, and Yekutieli) in the percent survival of spirochetes relative to the

Δ*cspZ*/Vector ("*") or between two strains relative to each other ("#") are indicated.

1108

1109

1110

1111

1112

1113

1114

Figure 4. The CspZ loop-mediated FH-binding activity defines host-specific early hematogenous dissemination in a complement-dependent manner. *I. scapularis* nymphs carrying B31-A3, B31-A3Δ*cspZ* harboring the empty vector pKFSS ("Δ*cspZ*/Vector"), or this *cspZ* mutant strain producing CspZ$_{B31}$, CspZ$_{B379}$, CspZ$_{B379}$L$_{B408}$, CspZ$_{B408}$, or CspZ$_{B408}$L$_{B379}$ were allowed to feed until repletion on **(A to E)** BALB/c mice or **(F to I)** quail. The bacterial loads in the indicated distal tissues were determined by qPCR at 10 days post nymphs feeding (dpf) in mice and 9dpf in quail. The bacterial loads were normalized to 100ng total DNA. Shown are the

1122 geometric mean of bacterial loads ± SEM of five mice or quail per group, except for the blood

1123 from BALB/c mice, which has nine mice per group. Significant differences ($p < 0.05$, Kruskal-

1124 Wallis test with the two-stage step-up method of Benjamini, Krieger, and Yekutieli) in the

1125 spirochete burdens relative to the $\Delta cspZ$/Vector ("*") or between two strains relative to each other

1126 ("#") are indicated.

1127

1128

1129

1130

1131

1132

1133

1134

1135

1136

1137

1138

1139

1140

1141

1142

1143

Figure 5. Phylogeny and sequence comparisons support polymorphic CspZ loop arising from host-specific adaptation of Lyme borreliae. (**A**) Unrooted likelihood tree of 174 *B. burgdorferi cspZ* isolates generated with IQ-tree and visualized in iTOL. The names of the isolates are highlighted based on their CspZ loop type (CspZ$_{B379}$, CspZ$_{B408}$, and CspZ$_{B31}$ as green, orange, and purple, respectively). The percent identity of each isolate relative to CspZ$_{B31}$ (purple), CspZ$_{B408}$ (orange), or CspZ$_{B379}$ (green) is indicated on the outer ring. The known FH-binding activity is marked: black or gray stick figures, mice, and quail indicate FH binding or lack of thereof, respectively, as determined herein and from previously published studies (27-29, 118, 119). (**B**)

The indicated concentrations of recombinant histidine-tagged, predicted last common ancestor states of CspZ ("CspZ-LCAS1, CspZ-LCAS2, CspZ-LCAS3, and CspZ-LCAS4), $CspZ_{B31}$, or DbpA (negative control) were added to triplicate wells coated with FH from human, mouse or quail, and protein binding was quantitated by ELISA. The experiments were performed with a single preparation of recombinant proteins tested in three independent iterations, in which samples were ran in duplicate. The K_D values **(Table S6)** representing the FH-binding affinity of each protein were determined from the average of three experiments. Shown is a representative iteration averaging the duplicates.

SUPPLEMENTAL FIGURES

Figure S1. Amino acid alignments of the CspZ and FH variants. Amino acid alignments of **(A)** the indicated CspZ variants, mutants, and reconstructed ancestral states of variants or **(B)** human, mammalian, and avian FH SCR 6-7. (**A**) The CspZ amino acids accounting for the loop structures are indicated with the box, and the alpha-helices labeled above the sequences are extrapolated from the high-resolution structure of CspZ$_{B408}$. The yellow and blue shading are indicative of loci showing evidence of positive and negative selection, respectively.

1183

Figure S2. The polymorphic loop structures in recombinant CspZ proteins promote host-

specific FH-binding ability determined by ELISA. The indicated concentrations of recombinant

GST-tagged $CspZ_{B379}$, $CspZ_{B408}$, $CspZ_{B379}L_{B408}$, or $CspZ_{B408}L_{B379}$, or GST (negative control) were

added to triplicate wells coated with FH from mouse or quail, and protein binding was quantitated

by ELISA. The experiments were performed with a single preparation of recombinant proteins

tested in three independent iterations, in which samples were ran in duplicate. The K_D values

(Table S2) representing the FH-binding affinity of each protein were determined from the average

of three experiments. Shown is a representative iteration averaging the duplicates.

1192

1193

1194

1195

1196

1197

1198

1199

1200

1201

Figure S3. Indistinguishable surface production of CspZ among *B. burgdorferi* strains was observed using flow cytometry. Flow cytometry analysis of CspZ localized on the surface of *B. burgdorferi* strains B31-A3, B31-A3Δ*cspZ* harboring the vector pKFSS ("Δ*cspZ*/Vector"), or this *cspZ* mutant strain producing CspZ$_{B31}$, CspZ$_{B379}$, CspZ$_{B379}$L$_{B408}$, CspA$_{B408}$, or CspZ$_{B408}$L$_{B379}$. **(A)** Representative histograms of flow cytometry analysis showing the levels of CspZ surface production on the indicated strains. **(B)** The production of FlaB (negative control) and CspZ on the surface of indicated *B. burgdorferi* strains was detected by flow cytometry. Values are shown normalized to the production levels of FlaB or CspZ on the surface of permeabilized B31-A3. Each bar represents the mean of four independent experiments ± the standard deviation. An asterisk (*) indicates that relative surface production of the indicated proteins was significantly lower (p <

1212   0.05, Kruskal-Wallis test with the two-stage step-up method of Benjamini, Krieger, and Yekutieli)

1213   than that of permeabilized FlaB or CspZ by B31-A3.

1214

1215

1216

1217

1218

1219

1220

1221

1222

1223

1224

1225

1226

1227

1228

1229

1230

1231

1232

1233

Figure S4. The polymorphic CspZ loop determines the host-specific, allelically variable FH-binding activity on spirochete surface. *B. burgdorferi* strains B313 (negative control), B31-A3, B31-A3Δ*cspZ* harboring the empty vector pKFSS ("Δ*cspZ*/Vector"), or this mutant strain producing CspZ$_{B31}$, CspZ$_{B379}$, CspZ$_{B379}$L$_{B408}$, CspZ$_{B408}$, or CspZ$_{B408}$L$_{B379}$, was incubated with mouse or quail FH. The bacteria were stained with antibodies that recognize these FH variants prior to flow cytometry. Shown are the representative histograms of flow cytometry analysis presenting the levels of FH from **(A)** mouse or **(B)** quail binding to each *B. burgdorferi* strain. The levels of **(C)** mouse or **(D)** quail FH-binding were measured by flow cytometry and presented as mean fluorescence index (MFI). Each bar represents the mean of three independent experiments ± SEM. Significant differences (p < 0.05, Kruskal-Wallis test with the two-stage step-up method of Benjamini, Krieger, and Yekutieli) in the levels of FH-binding relative to the B313/Vector ("Φ"), Δ*cspZ*/Vector ("*"), or between two strains relative to each other ("#") are indicated.

1246

1247

1248

1249

Figure S5. The CspZ loop-driven, host-specific serum resistance is recovered in complement-

depleted sera. *B. burgdorferi* strains B313 (negative control), B31-A3, B31-A3Δ*cspZ* harboring

the empty vector pKFSS ("Δ*cspZ*/Vector"), or this mutant strain producing CspZ$_{B31}$, CspZ$_{B379}$,

CspZ$_{B379}$L$_{B408}$, CspZ$_{B408}$, or CspZ$_{B408}$L$_{B379}$were incubated for 4-h with **(A)** CVF-treated white-

footed mouse or **(B)** OMCI-treated quail sera, to a final concentration of 40%. The number of

motile spirochetes was assessed microscopically. The precent survival of the *B. burgdorferi* strains

was calculated using the number of motile spirochetes at 4-h post incubation normalized to that

prior to incubation with sera. Each bar represents the mean of three independent experiments ±

SEM. There were no significant differences (p < 0.05, Kruskal-Wallis test with the two-stage step-

up method of Benjamini, Krieger, and Yekutieli) between the percent survival of any strain.

1260

1261

1262

1263

Figure S6. *B. burgdorferi* **strains exhibit similar burdens in flat and fed nymphs.** *B. burgdorferi*-infected flat nymphs were allowed to feed to repletion on **(B, E)** BALB/c or **(G)** C3-/- BALB/c mice, or **(C, F)** quail or **(H)** OmCI-treated quail. The spirochete loads in **(A, D)** flat or **(B, C, E, F, G, H)** replate nymphs were determined by qPCR. Shown are the geometric mean ± geometric standard deviation of at least five nymphs per group. There was no statistical difference (p > 0.05) of the spirochete burdens between different groups of the replete ticks using a **(A to C)** Mann-Whitney test or **(D to H)** Kruskal-Wallis test with the two-stage step-up method of Benjamini, Krieger, and Yekutieli.

1272

1273

1274

Figure S7. CspZ facilitates early bacteremia and distal tissue colonization during tick infection. The *I. scapularis* nymphs carrying *B. burgdorferi* strain B31-A3 or B31-A3Δ*cspZ* harboring the vector pKFSS ("Δ*cspZ*/Vector") were allowed to feed until repletion on **(A to O)** BALB/c mice or **(P to W)** quail. The mice were euthanized at **(A to E)** 7, **(F to J)** 10, or **(K to O)** 14 days post nymphs feeding (dpf), whereas the quail were euthanized at **(P to S)** 9 or **(T to W)** 14dpf. **(A, F, K)** The site of the skin where nymphs fed ("Bite site"), **(B, G, L)** blood, **(C, H, M)** tibiotarsus joints, **(D, I, N)** heart, and **(E, J, O)** bladder of mice; and **(P, T)** the bite site, **(Q, U)** blood, **(R, V)** brain, and **(S, W)** heart of quail, were collected immediately after euthanasia and spirochete loads were determined by qPCR. The burdens were normalized to 100ng total DNA.

1285   Shown are the geometric mean of bacterial loads ± SEM of five mice or quail per group, except

1286   for samples from the mouse blood at 7 and 10dpf where the results are from six and nine mice,

1287   respectively. Significant differences ($p < 0.05$, Mann-Whitney test) in the spirochete burdens

1288   between two strains relative to each other ("#") are indicated.

1289

1290

1291

1292

1293

1294

1295

1296

1297

1298

1299

1300

1301

1302

1303

1304

1305

1306

Figure S8. The CspZ loop-mediated early hematogenous dissemination is recovered when complement is depleted from hosts. *I. scapularis* nymphs carrying B31-A3, B31-A3Δ*cspZ* harboring the empty vector pKFSS ("Δ*cspZ*/Vector"), or this *cspZ* mutant strain producing CspZ$_{B31}$, CspZ$_{B379}$, CspZ$_{B379}$L$_{B408}$, CspZ$_{B408}$, or CspZ$_{B408}$L$_{B379}$ were allowed to feed until repletion on **(A to E)** C3$^{-/-}$ mice in a BALB/c background or **(F to I)** OMCI-treated quail, both of which deplete complement in the respective hosts. The bacterial loads in the indicated distal tissues were determined by qPCR and were normalized to 100ng total DNA, at 10 days post nymphs feeding (dpf) in mice and 9dpf in quail. Shown are the geometric mean of bacterial loads ± SEM of five

1315   mice or quail per group, except for the blood from BALB/c mice, which has nine mice per group.

1316   There were no significant differences ($p < 0.05$, Kruskal-Wallis test with the two-stage step-up

1317   method of Benjamini, Krieger, and Yekutieli) in the spirochete burdens for any strain.

1318

1319

1320

1321

1322

1323

1324

1325

1326

1327

1328

1329

1330

1331

1332

1333

1334

1335

1336

1337 **Figure S9. Phylogenetic network of *cspZ* haplotypes.** Edges are colored based on their loop

1338 structure (CspZ$_{B379}$, CspZ$_{B408}$, and CspZ$_{B31}$ in green, orange, and purple, respectively).

1339

1340

1341

1342

1343

1344

1345

1346

1347

1348

1349

1350

Figure S10. The CspZ last common ancestor states are predicted to versatiley bind FH. The

crystal structure of CspZ$_{B408}$-SCR6-7 where human SCR6-7 (grey) is superimposed with mouse

FH SCR6-7 (gold, PDB: 2YBY) and the predicted structure of quail SCR6-7 (brown). CspZ$_{B408}$

(orange) is superimposed with CspZ$_{B31}$ (purple) and the last common ancestor states: LCAS1

(yellow), LCAS2 (blue), LCAS3 (green), and LCAS4 (pink). Residues that differ between CspZ$_{B31}$

and the LCAS variants are labelled.

1357

Figure S11. Structural comparison between CspZ$_{B408}$-human SCR6-7 and the *N. meningitidis*

Fhbp-human SCR6-7 complexes. Human FH SCR6-7 (light grey) from the complex structure

with CspZ$_{B408}$ (orange) was superimposed with human FH SCR6-7 (dark grey) from the complex

structure with Fhbp (dark blue, PDB: 2W81), the FH-binding protein from *N. meningitidis*. α-

helices in CspZ$_{B408}$ are labelled from A to I starting from the N-terminus.

 Figure S12. CD spectra demonstrate no impacts of secondary structures by swapping the

 loops. Far-UV CD analysis of $CspZ_{B379}$, $CspZ_{B408}$, $CspZ_{B379}L_{B408}$, and $CspZ_{B408}L_{B379}$. The molar

 ellipticity, Φ, was measured from 190 to 250nm for 10μM of each protein in PBS.

1379

Figure S13. OmCI prevents quail serum-mediated killing of a complement-sensitive spirochete strain at 11 days post injection. *Coturnix* quail were subcutaneously inoculated with OmCI (1mg/kg of quail) or PBS buffer. Untreated (filled bars) or heat-treated (hatched bars) sera collected from these quail at 11 days post inoculation (dpi) were incubated with a serum-sensitive, highly passaged *B. burgdorferi* strain B313. The number of motile spirochetes were quantified microscopically and the survival percentage of the spirochetes was calculated using the number of mobile spirochetes at 4 h post incubation normalized to that at 0 h. Each bar represents the mean ± SEM of three independent experiments from sera from four quail per group. Significant differences ($p < 0.05$, Mann-Whitney test) in the percentage survival of spirochetes are indicated ("#").

1390

1391