Siderophore-mediated iron partition promotes dynamical coexistence between

cooperators and cheaters

Jiqi Shao[1], Nan Rong[1], Zhenchao Wu[2], Beibei Liu[2], Ning Shen[2], and Zhiyuan Li[1,3*]

[1] Center for Quantitative Biology, Academy for Advanced Interdisciplinary Studies, Peking

University, Beijing, 100871, China

[2] Department of Pulmonary and Critical Care Medicine, Peking University Third Hospital,

Beijing 100191, China

[3] Peking-Tsinghua Center for Life Sciences, Academy for Advanced Interdisciplinary

Studies, Peking University, Beijing, 100871, China

* To whom correspondence should be addressed. Tel: +86 10 62750190; Email:

zhiyuanli@pku.edu.cn

Abstract

Microbes shape their habitats through consuming resources, as well as actively producing
and secreting diverse chemicals. These chemicals serve various niche-construction
functions and can be considered "public good" for the community. Most microorganisms,
for instance, release small molecules known as siderophores to scavenge irons from the
extracellular environment. Despite being exploitable by cheaters, biosynthetic genes
producing such molecules widely exist in nature, invoking active investigation on the
possible mechanisms for producers to survive cheater invasion. In this work, we utilized
the chemostat-typed model to demonstrate that the division of the iron by private and
public siderophores can promote stable or dynamical coexistence between the cheater
and "partial cooperators", an adaptive strategy with the production of both public and
private siderophores. Further, our analysis revealed that when microbes not only consume
but also produce resources, this type of "resource partition model" exhibit different stability
criteria than that of the classical consumer resource model, allowing more complex
systems dynamics.
29
30   Keywords: siderophore, oscillation, cheaters and cooperators, resource competition and
31   partition

Introduction

Microbes interact by shaping their microhabitats (Bajić, Rebolleda-Gómez et al. 2021). Essential to this intricate ecological loop is the "chemical environment": the substances in the local inhabitants that directly influence and are influenced by microbes(Ley, Peterson et al. 2006, Delmont, Robe et al. 2011). Common examples of such substances include "resources" such as carbon, nitrogen, and oxygen, which are supplied into the extracellular environment and are consumed by microorganisms for their own growth(Boyd and Ellwood 2010, Miransari 2013, Dutkiewicz, Cermeno et al. 2020). In theoretical ecology, the consumer resource model has been utilized for decades to explore the feedback between species and resources(Lafferty, DeLeo et al. 2015), whereby the dimension of the chemical space has been demonstrated to be of key relevance: MacArthur et al. validated the so-called competitive exclusion principle (CEP) that the number of stably coexisting species cannot exceed the number of resources for which they are competing(Hardin and G. 1960, Macarthur and Levins 1964); In the chemical space of dimension two, Tilman et al. demonstrated that the zero-net growth isoclines and the vector of consumption determine the outcome of competition(Tilman 1982); And in a multi-species resource competition system with a minimal chemical dimension of three, Huisman et al. proved that oscillatory and even chaotic dynamics could emerge to permit dynamical coexistence that exceeds the upper bounds of CEP(Huisman and Weissing 1999, Huisman and Weissing 2001). Actually, numerous research efforts have been devoted to conciliating the conflicts between CEP and the apparent biodiversity in nature, most of which incorporate additional factors such as spatial or temporal heterogeneity(Erez, Lopez et al. 2020, Ho, Good et al. 2022). No definitive answer has yet been achieved (Roy and Chattopadhyay 2007, Gupta, Garlaschi et al. 2021).

Microorganisms have their means of lifting the upper limit of competitive exclusion. By leaking metabolic byproducts or actively producing and secreting secondary metabolites, cells can expand the chemical diversity of their habitats(Schmidt, Ulanova et al. 2019). Such activities, in which microorganisms generate new chemical dimensions, have profound implications for microbial ecology (Fischbach and Segre 2016, Estrela, Diaz-Colunga et al. 2022). For example, pervasive cross-feeding through metabolic byproducts has been shown suggested to promote stable coexistence even under a single carbon source(Goldford, Lu et al. 2018). Meanwhile, the vast variety of bacteriocins and antibiotics escalates microbial chemical warfare (Czaran, Hoekstra et al. 2002, Granato, Meiller-Legrand et al. 2019, Niehus, Oliveira et al. 2021). For microbial cooperation, various quorum sensing molecules coordinate the collective decision-making (Ross-Gillespie and Kümmerli 2014), and niche-construction molecules such as siderophores or secretory proteases serve as "public goods" that improve the microenvironment for the whole population(Leventhal, Ackermann et al. 2019, Smith and Schuster 2019). In ecological theories, however, the production of secondary metabolites confronts the same dilemma as public goods in "the tragedy of the commons" (Hardin 1968): if a cheater strain can always take advantage of public goods without paying for

76 them, how can the cooperator strains accountable for these extra chemical dimensions
77 remain competitive?
78
Big questions in biology may find diverse solutions in particular systems. Siderophore, a
diverse family of microbial secondary metabolites for iron-scavenging, serves as a nice
model system to study the self-generated chemical dimensions and the game between
cooperators and cheaters (Cremer, Melbinger et al. 2019). Iron is among one of the most
limiting resources for microbes(Andrews, Robinson et al. 2003). It is necessary for
processes like energy metabolism and DNA synthesis(Andrews, Robinson et al. 2003),
yet the concentration of bioavailable iron is orders of magnitude lower than what is
required for normal microbial growth in most environments (Boyd and Ellwood 2010,
Emerson, Roden et al. 2012). Most microorganisms acquire iron via siderophores, a type
of small molecule with a strong iron-binding affinity (Buckling, Harrison et al. 2010).
Previous research suggested that siderophores are secreted into the extracellular
environment to chelate trivalent iron, and that the iron-siderophore complex is then
acquired by membrane receptors so that cells can absorb iron (Kramer, Özkaya et al.
2020). The production of siderophores, on the other hand, comes at a considerable
metabolic burden and slows down microbial growth (Lv, Hung et al. 2014, Sexton and
Schuster 2017). Therefore, siderophore acts as a costly public good that invokes intricate
ecological games. Many theories have been proposed to explain the prevalence of
siderophore synthesis pathways (Butaitė, Baumgartner et al. 2017, Gu, Wei et al. 2020),
most of which entail spatial factors that facilitate kin selection or group selection (Allison
2005, Ross-Gillespie, Gardner et al. 2007, Julou, Mora et al. 2013). Yet, in well-mixing
environments such as the ocean, microorganisms continue to actively produce
siderophores of various types (Cordero, Ventouras et al. 2012, Hagstrom and Levin
2017).
102
Recent experiments indicated that private siderophores, which are solely accessible to
their cooperators, may be crucial to the survival of siderophore-producing microorganisms
(Scholz and Greenberg 2015). In some organisms, during the multi-step process of
producing siderophores, certain modifications can transform the secretory molecules into
a membrane-attached form (Martinez, Carter-Franklin et al. 2003, Scholz and Greenberg
2015, Niehus, Picot et al. 2017). Such privatization avoids diffusion losses and cheater

exploration, resembling the "snowdrift" scenario in game theory (Souza, Pacheco et al.

2009): Despite that cheaters continue to take advantage of public goods without paying,
cooperators have prioritized access to the goods to preserve coexistence (Gore, Youk et
al. 2009). Nevertheless, membrane-attached siderophores suffer a considerably lower
diffusion radius in scavenging iron (Leventhal, Ackermann et al. 2019, Kramer, Özkaya et
al. 2020). Whether the marginal benefits conferred by membrane-attached siderophores
are adequate to provide cooperators with a sufficient advantage over cheaters has not
been quantitatively assessed by ecological models. Many questions remain to be
systematically explored by mathematical formulation, such as whether and how
cooperators and cheaters coexist, how different allocation strategies between

membrane-attached and public-shareable secretory siderophores change the system
dynamics, and which strategies might be "optimal" considering within species and
between species. Further, may this siderophore-mediated iron competition helps address
a deeper question in ecology, namely whether such self-generated resource dimensions
change the properties of the system fundamentally?
124
In this work, we utilized the chemostat-type "resource partition model" to examine the
siderophore-mediated iron competition, taking into account the allocation of limited
cellular resources between the biomass accumulation and the production of
membrane-attached (private) and public-sharable (public) siderophores. With the
inclusion of private siderophores into the resource allocation strategies, new classes of
strategies emerge, such as "partial cooperators" that produce both types of siderophores,
and "self-seekers" that produce only membrane-attached siderophores. We confirmed
mathematically that private siderophores play a vital role in coexistence: private
siderophores provide partial cooperators with an advantageous growth zone over
cheaters to enable coexistence, in contrast to pure cooperators who always lose to
cheater invasion. Interestingly, such two-species coexistence can occur via dynamical
oscillation. Further stability analysis revealed that the action of species creating new
resource dimensions, i.e. the production of siderophores in our model, modifies the
stability criteria of the classical consumer resource model, allowing for rich dynamics in
the parameter regions of stable equilibrium for the classical model. In summary, our model
in iron competition suggested that the division of the iron resource by siderophores raises
the upper-bounds of coexistence, and the privatization of siderophores in cellular
strategies provides advantages to partial cooperators to realize coexistence. Our analysis
of the stability criteria revealed that the microbial niche-construction by creating new
resource dimensions adds more dynamics than the classical model, which may contribute
to the diversity and complexity of the microbial world.
146

Result

A resource partition model with trade-offs between growth and

siderophore-production

In the field of theoretical ecology, the consumer resource model mimics ecosystems with
constant nutrient supplies and extensive mixings, such as lakes and oceans, where
species compete by consuming the supplied resources. Nonetheless, in the microbial
world, species not only consume but also produce resources that can be shared by the
community, such as siderophores for iron scavenging. To examine the ecological
consequences of siderophore production and privatization, we developed a mathematical
model resembling a chemostat with trade-offs between growth and siderophore

production, with the following three assumptions:

1. Chemostat-typed resource partition: we used a chemostat-type model, in which
the volume is maintained constant by maintaining the same rate of in-and-out
fluxes. Parameters about the concentration of iron in the influx $R_{\text{iron,supply}}$,
together with the dilution rate d, are referred to as "chemostat conditions" (Fig.
1A). Within the chemostat, the chemical environment cells directly facing can be
quantified by two variables: concentration of the iron (R_{iron}) and of the public
siderophores (R_{sid}). They serve as both "resources" for microbial growth;
meanwhile, the public siderophore is also a "product" released by bacteria. We
coined the term "resource partition model" to refer to models in which organisms
generate more chemical dimensions than are supplied from the external influx,
and partition the externally supplied resources by these secondary metabolites.
2. The growth rate linearly scales with the iron uptake rate: Assuming iron was the
limiting resource, we set the growth rate to a value that linearly scales the total
iron fluxes obtained via two types of siderophores: the flux from secretive public
siderophores (J_{public}, depicted on the left side of Fig. 1B) and the flux from
membrane-bounded private siderophore (J_{private}, depicted on the right side of Fig.
1B).
3. Trade-offs in resource allocation between growth and siderophore-production:
Given the limited amount of proteins and energies in a microorganism, we
assumed that there are trade-offs between different biological processes. In an
iron-limited environment, we focused on resource partitioning into biomass
accumulation and production of private and public siderophores. Each partition
corresponds to an allocation strategy $\vec{\alpha} = \left(\alpha_{\text{growth}}, \alpha_{\text{private}}, \alpha_{\text{public}}\right)$, where
α_{private} and α_{public} denote the percentage of resources used to produce private
and public siderophores, respectively, and α_{growth} donates the percentage of
resources devoted to biomass accumulation. The summation of all allocations
was fixed as $\alpha_{\sigma,\text{growth}} + \alpha_{\sigma,\text{private}} + \alpha_{\sigma,\text{public}} = 1$. Various species may differ in
their abilities to produce and scavenge siderophores, but different strains σ only

differ in their resource allocation strategies $\vec{\alpha}$.

Under this trade-off, all possible strategies of a species can be located in a ternary
graph (Fig. 1C). There are four types of typical strategies (Fig. 1C-D): pure
cooperators, the pure cheater, partial cooperators, and self-seekers. Pure
cooperators produce public siderophores but not private siderophores ($\alpha_{\text{private}} =$
$0, \alpha_{\text{public}} > 0$), while the pure cheater allocates all resources to growth and
exploits only the public siderophores ($\alpha_{\text{growth}} = 1$). On the other hand, partial
cooperators ($\alpha_{\text{public}} > 0, \alpha_{\text{private}} > 0$) and self-seekers($\alpha_{\text{public}} = 0, \alpha_{\text{private}} > 0$)
have access to both public and private siderophores due to non-zero α_{private}. The
difference between the two strategies is that partial cooperators still produce
public siderophores while self-seekers do not. In the ternary graph (Fig. 1C), pure
cooperators' strategies span the triangle's base, while the pure cheater's strategy
locates on the right-vertex. The right side is occupied by self-seekers, and each
strategy contained within the triangle can be considered a partial cooperator.

With the three assumptions above, the microbe σ with biomass concentration m_σ and
strategy $\vec{\alpha}_\sigma$ shapes its chemical environment in two ways:

1. Producing public siderophore with the out flux $\epsilon \frac{m_\sigma}{r} \alpha_{\sigma,\text{public}}$, where ϵ is the

production coefficient of public siderophores, and the constant r represents the
biomass per cell volume (Details in SI). Assuming the public siderophores can be
fully recycled for simplicity, the changing rate for R_{sid} can be written as:

$$\frac{dR_{\text{sid}}}{dt} = -dR_{\text{sid}} + \sum_\sigma \epsilon \frac{m_\sigma}{r} \alpha_{\sigma,\text{public}}. \tag{1}$$

208   2. Consuming iron by public siderophores and private siderophores.
209   For iron-uptake, the fluxes of $J_{\sigma,\text{private}}$ and $J_{\sigma,\text{public}}$ take the Monod form with the
210   environmental iron concentration R_{iron}. We also assumed that both fluxes linearly
211   scale with the concentration of corresponding siderophores. Taken together, there
212   are:

$$J_{\sigma,\text{private}} = \frac{v_{\text{m}} \beta \alpha_{\sigma,\text{private}} R_{\text{iron}}}{K_{\text{m}} + R_{\text{iron}}}, \tag{2}$$

$$J_{\sigma,\text{public}} = \frac{v_{\text{l}} R_{\text{sid}} R_{\text{iron}}}{K_{\text{l}} + R_{\text{iron}}}. \tag{3}$$

213   In the equations above, v_{m} and v_{l} are the rate coefficients for the two fluxes; K_{m}
214   and K_{l} are the affinity coefficients of the two kinds of siderophores for intaking
215   iron; β is the efficiency coefficient of the private siderophores. In total, the
216   changing rate for iron is written as:

$$\frac{dR_{\text{iron}}}{dt} = d\left(R_{\text{iron,supply}} - R_{\text{iron}}\right) - \sum_\sigma \frac{m_\sigma}{r} \left(J_{\sigma,\text{private}} + J_{\sigma,\text{public}}\right), \tag{4}$$

217   It can be proved that other forms of iron uptake-fluxes do not qualitatively affect
218   the result of this work (See Supplementary Material, Section 4.2).

220 Meanwhile, the microbe σ has its growth rate affected by its chemical environment

221     $[R_{\mathrm{iron}}, R_{sid}]$ and allocation strategy $\vec{\alpha}_\sigma$ as:

$$g(R_{\mathrm{iron}}, R_{sid}, \vec{\alpha}_\sigma) = \gamma \alpha_{\sigma,\mathrm{growth}}(J_{\sigma,\mathrm{private}} + J_{\sigma,\mathrm{public}}), \tag{5}$$

222   where γ here represents the growth coefficient.

223   In this iron-partition model, the changing rate for the biomass for microbe σ can be
224   written as:

$$\frac{dm_\sigma}{dt} = m_\sigma \cdot (g(R_{sid}, R_{\mathrm{iron}}, \vec{\alpha}_\sigma) - d). \tag{6}$$

Privatization of siderophore enables the coexistence between partial cooperators

and pure cheaters

The graphical approaches developed for consumer resource models enabled an intuitive
assessment of ecological consequences in the chemical space (Tilman 1982) (Fig. S1).
This approach consists of two elements: the growth contour, i.e. the zero net growth
isoclines (ZNGI), and the consumption vector. After setting Eq. (6) to zero for a single
subpopulation, we obtained the growth contours. The growth contour shows all possible
chemical environments $[R_{\mathrm{iron}}, R_{\mathrm{sid}}]$ that the microbe could reach the steady-state growth
as the dilution rate. Any environment above the growth contour belongs to the "invasible
zone" where the microbe can invade (Li, Liu et al. 2020).

This graphical approach demonstrates intuitively why a pure cooperator cannot coexist
with a pure cheater. For two strains, the possible chemical environment allowing for stable
coexistence must locate at the intersection of their growth contours, indicated by the "*".
As illustrated in Figure 1E, the growth contour of the pure cheater entirely encloses the
growth contour of the pure cooperators without an intersection. Intuitively, it is due to that
the pure cheater grows faster than the pure cooperator in any chemical environment: they
have the same iron-influx by public siderophores J_{public}, yet the cooperator invests less in
α_{growth} (See rigorous proof in the Supplementary Material, Section 2.2). However,
because the pure cheater can only utilize public siderophores and cannot survive
independently, it can be observed in the simulation that the pure cheater eventually takes
the entire population and becomes extinct due to a lack of public siderophores. This
approach resembles "the tragedy of the commons".

Investing in private siderophores can change the game. The transition from pure
cooperators to partial cooperators enables the growth contours to intersect with those of
pure cheaters, hence enabling coexistence (Fig. 1F). In regions where the public
siderophore is abundant but iron is scarce (upper-left region of Figure 1F filled in white),
the cheater still retains a growth advantage over the partial cooperator; however, in
regions where the public siderophore is scarce but free iron is abundant (lower-right
region of Figure 1F filled in blue-green color), the partial cooperator gains a growth
advantage over the pure cheater because their membrane siderophores sustain
iron-dependent growth. Under this model setting, the game enters the regime of "snowdrift"
(Souza, Pacheco et al. 2009): despite the fact that producing siderophores incurs costs,

cooperators gain by holding portions of them as private goods, particularly in
environments where public siderophores are scarce. The fact that the superior strategy
varies according to the chemical environment enables the intersection of their growth
contours, which is necessary for coexistence.
Mathematically, it can be analytically proven that the presence of intersection between
growth contours of strain 1 and strain 2, under the setting that strain 1 invests less in its
own growth $\alpha_{1,\text{growth}} < \alpha_{2,\text{growth}}$, requires:

$$\alpha_{1,\text{private}} > \alpha_{2,\text{private}}. \tag{7}$$

266   Equation (7) shows that for any two strains to coexist, the strain that invests fewer
267   resources in growth must invest more in private siderophores (Details can be found in
268   Supplementary Material, Section 2.3).
269   Meanwhile, the non-zero biomass of two strains imposes constraints on the iron supply
270   concentration $R_{\text{iron,supply}}$ in that

$$R_{\text{iron}}^* + \frac{1}{k_2} R_{\text{sid}}^* > R_{\text{iron,supply}} > R_{\text{iron}}^* + \frac{1}{k_1} R_{\text{sid}}^*. \tag{8}$$

with $k_1 = \frac{\epsilon\gamma}{d}\alpha_{1,\text{public}}\alpha_{1,\text{growth}}$, $k_2 = \frac{\epsilon\gamma}{d}\alpha_{2,\text{public}}\alpha_{2,\text{growth}}$.

Graphically, inequality Eq. (8) requires the supply point $[R_{\text{iron,supply}}, 0]$, locating within the
region bounded by the reverse extensions of the consumption vectors in the chemical
space (See Supplementary Material, Section 2 for detailed proofs)
However, the presence of the intersection of growth contours is only one of the necessary
conditions for coexistence; the feasibility of coexistence is further determined by how each
strategy shapes the chemical environment for the whole community (Li, Liu et al. 2020). In
the system illustrated in Figure 1F, where the partial cooperator coexists stably with the
pure cheater (competition dynamics shown in insert), the partial cooperator creates a
public-siderophore-rich environment in the upper-left area of the chemical space where
the cheater can invade. Meanwhile, the pure cheater cannot survive on its own and has
no corresponding stable point, but its competition for iron reduces the abundance of the
partial cooperator, causing an environment deficient in public siderophores to favor the

partial cooperator. This interaction between the partial cooperator and the pure cheater

resembles mutual invasion and allows for stable coexistence.
Such stable coexistence does, however, occur under a specific chemostat condition
(quantified as the iron influx $R_{\text{iron,supply}}$ and the dilution rate d). The presence of an area
in the chemical space in which the partial cooperator thrives prompted us to wonder
whether other chemostat conditions would allow for even more interesting ecological
dynamics.

The partial cooperators and cheaters can generate rich ecological dynamics

In classical consumer resource models, oscillatory dynamics can be generated among a
minimum of three species that preferentially consume the resources for which they have
intermediate requirements (Huisman and Weissing 1999, Huisman and Weissing 2001).
To our surprise, oscillations between the partial cooperator and the pure cheater can be

detected in this resource partition model at intermediate levels of iron supply (Fig. 2A).
In comparison to the steady coexistence depicted in Figure 1F, the partial cooperator
generates an environment susceptible to cheater invasion. The intersection of the two
growth contours, on the other hand, becomes unstable.
300
Figure 2C explains the oscillation visually with a phase diagram. In the diagram's
upper-left quadrant, where the public siderophore concentration is high and the iron
concentration is low, the rapid proliferation of cheaters effectively eliminates partial
cooperators (cooperator -, cheater+). The decline in partial cooperators results in a
decrease of public siderophores, which forces the system into the lower-left quadrant,
where the abundance of both partial cooperators and cheaters falls (cooperator-,
cheater-). Reduced total population promotes iron concentration recovery, allowing the
system to enter the advantageous growth zone of partial cooperators, where public
siderophores remain low but iron is abundant. Iron influx through private siderophores
causes partial cooperators to resume positive growth in this lower-right quadrant, whereas
pure cheaters continue to fall (cooperator+, cheater-). The rise of partial cooperators
increases the concentration of public siderophores, hence inhibiting the decrease of pure
cheaters and restoring their growth, as shown in the upper-right quadrant (cooperator+,
cheater+). However, the continued growth of pure cheaters depletes iron, forcing partial
cooperators into negative net growth again (cooperator-, cheater+). The preceding
process generates oscillations, mostly owing to the private siderophores providing an
advantageous growth zone for partial cooperators in the absence of public siderophores.
318
319   When the partial cooperator is more capable of establishing a high-iron,
320   low-public-siderophores environment, the dynamics become even more skewed in favor
321   of partial cooperators. For instance, if the steady-state environment formed by partial
322   cooperators goes outside the development contour of pure cheaters, the partial
323   cooperators can effectively exclude the pure cheater (Fig. 2B).
324
Scanning across the chemostat conditions revealed the bifurcation into and out of
oscillation. In the system with one pure cheater and one partial cooperator, as the dilution
rate increases, a Hopf bifurcation drives the transition from stable coexistence into
oscillation (Fig. 2D-E). Similarly, by increasing the iron supply concentration, the
oscillation shifted into stable coexistence. The phase diagram of chemostat conditions
(Fig. S2) demonstrates that a "better" environment (low dilution rate or high iron supply)
reduces the relative difference in growth between partial cooperators and cheaters,
thereby increasing the tendency of coexistence; on the other hand, a harsher environment
(high dilution rate or low iron supply) increases the relative difference in growth between
the two, thereby increasing the likelihood of oscillations.
335
Oscillatory dynamics demand a minimum of three species in a classical consumer
resource model (Huisman and Weissing 1999, Huisman and Weissing 2001). The
observed oscillation between the pure cheater and the partial cooperator not only sheds
light on the stability of cooperation in an iron-related snowdrift game, but also motivates us

to dig deeper into the effect of the self-generated chemical dimension on the resource
partition models.

The self-generated chemical dimension changes the stability criteria of classical

consumer resource models.

Similar to a classical consumer resource model (represented by Tilman's model (Tilman
1982)), the generalized form of microbes interacting with two resources R_1 and R_2 can
be represented as:

$$\frac{dm_i}{dt} = m_i(g_i(R_1, R_2) - d), \text{ for } i = 1,2 \tag{9}$$

$$\frac{dR_j}{dt} = d(R_{j,\text{supply}} - R_j) - \sum_{i=1}^{2} m_i h_{ij}(R_1, R_2)g_i(R_1, R_2), \text{ for } j = 1,2, \tag{10}$$

where $R_{j,\text{supply}}$ is the supply concentration of resource j, and h_{ij} is the function
describing the amount of resource j impacted by microbe i per-biomass.
When h_{ij} describes resource uptake and has a positive sign. Eq. (9)-(10) above
represent the classical consumer resource model. When at least one of the h_{ij} describes
resource production and exhibits a negative sign, these equations represent the broader
"resource partition model" and exhibit different criteria for the stability of coexistence.
353
354 The stability of the fixed point (*) can be deduced from the Routh-Hurwitz (RH) criterion
355 (May and Allen 1977). From the Jacobian matrix of the model:

$$J = \begin{bmatrix} 0 & 0 & v_{11} & v_{12} \\ 0 & 0 & v_{21} & v_{22} \\ -w_{11} & -w_{12} & -x_{11} & -x_{12} \\ -w_{21} & -w_{22} & -x_{21} & -x_{22} \end{bmatrix}, \tag{11}$$

356 with

$$v_{ij} = \left(\frac{\partial \dot{m}_i}{\partial R_j}\right)^*, \ w_{ij} = -\left(\frac{\partial \dot{R}_i}{\partial m_j}\right)^*, \ x_{ij} = -\left(\frac{\partial \dot{R}_i}{\partial R_j}\right)^*,$$

357 For simplicity, we set the abbreviation as:

$$P_1 = \left(\partial_{R_1} g_1\right)^*, P_2 = \left(\partial_{R_2} g_1\right)^*,$$

$$P_3 = \left(\partial_{R_1} g_2\right)^*, P_4 = \left(\partial_{R_2} g_2\right)^*, \tag{12}$$

$$c_{11} = (h_{11})^*, c_{12} = (h_{21})^*,$$

$$c_{21} = (h_{12})^*, c_{22} = (h_{22})^*.$$

358 Here, P can be interpreted as the growth rate dependency on "resources" at the fixed
359 point, and c_{ij} represents how microbe i impacts resource j at the fixed point.
360 So we have elements in the Jacobian matrix expressed as:

$$\begin{aligned} v_{11} &= m_1^* P_1, & v_{12} &= m_1^* P_2, \\ v_{21} &= m_2^* P_3, & v_{22} &= m_2^* P_4 \end{aligned} \tag{13}$$

$$w_{11} = c_{11}d, \qquad w_{12} = c_{12}d,$$
$$w_{21} = c_{21}d, \qquad w_{22} = c_{22}d.$$
$$x_{11} = d, x_{12} = 0,$$
$$x_{21} = m_1^* c_{21} P_1 + m_2^* c_{22} P_3, x_{22} = d + m_1^* c_{21} P_2 + m_2^* c_{22} P_4$$

361 At the steady state, $g_1^* = g_2^* = d$. With definitions in Eq.(12)-(13), the characteristic
362 equation of eigenvalue λ for the Jacobian matrix can be expressed as:

$$\lambda^4 + (x_{11} + x_{22})\lambda^3 + (q_1 + q_4 + x_{11}x_{22} - x_{12}x_{21})\lambda^2$$
$$+ (x_{11}q_4 + x_{22}q_1 - x_{12}q_3 - x_{21}q_2)\lambda \qquad (14)$$
$$+ (q_1 q_4 - q_2 q_3) = 0,$$

363 the coefficients of this quartic equation are: $a_0 = 1$, $a_1 = x_{11} + x_{22}$, $a_2 = q_1 + q_4 -$
364 $x_{12}x_{21} + x_{11}x_{22}$, $a_3 = q_4 x_{11} - q_3 x_{12} - q_2 x_{21} + q_1 x_{22}$, $a_4 = -q_2 q_3 + q_1 q_4$, with q_i defined
365 as:

$$q_1 = w_{11}v_{11} + w_{12}v_{21},$$
$$q_2 = w_{11}v_{12} + w_{12}v_{22},$$
$$q_3 = w_{21}v_{11} + w_{22}v_{21}, \qquad (15)$$
$$q_4 = w_{21}v_{12} + w_{22}v_{22}.$$

The Routh-Hurwitz (RH) criterion states the necessary and sufficient condition for the
stability of a dynamical system (May and Allen 1977): for a quartic equation for the
eigenvalue λ, $a_0\lambda^4 + a_1\lambda^3 + a_2\lambda^2 + a_3\lambda + a_4 = 0$, if the fixed point for coexistence is
stable, then the coefficients must satisfy: (1) $a_1 > 0$, (2) $a_4 > 0$, (3) $a_3 > 0$, (4)
$a_1 a_2 a_3 - a_1^2 a_4 - a_3^2 > 0$.
Based on the four RH criteria, we compared the difference between the classical
consumer resource model and the resource partition model on its stability. Details on the
comparison can be found in Supplementary Material Section 3.1.
374
375 First, the criteria (1) $a_1 > 0$ holds for both the classical consumer resource model and the
376 iron-partition model.
377 Second, The criteria (2) $a_4 > 0$ is central to the classical model. By definitions in Eq. (12),
378 the original form $a_4 = -q_2 q_3 + q_1 q_4$ expands into:

$$a_4 = dm_1^* m_2^* (c_{12}c_{21} - c_{11}c_{22})(P_2 P_3 - P_1 P_4), \qquad (16)$$

379 There are two possible situations for $(c_{12}c_{21} - c_{11}c_{22})(P_2 P_3 - P_1 P_4) > 0$:

situation 1: $$\frac{P_2}{P_1} > \frac{P_4}{P_3} , \frac{c_{11}}{c_{21}} < \frac{c_{12}}{c_{22}}, \qquad (17)$$

or situation 2: $$\frac{P_2}{P_1} < \frac{P_4}{P_3} , \frac{c_{11}}{c_{21}} > \frac{c_{12}}{c_{22}}. \qquad (18)$$

380 In the classical consumer resource model, these two inequalities can be interpreted as
381 "each species must consume more of the one resource which more limits its own growth
382 rate."
383 In the resource partition model, using the iron-partition model at Eq. (1)-(6) as the
384 example, parameters in Eq.(12) can be specified into:

$$P_1 = \left(\partial_{R_{\text{sid}}} g_1\right)^*, \qquad P_2 = \left(\partial_{R_{\text{iron}}} g_1\right)^*,$$
$$P_3 = \left(\partial_{R_{\text{sid}}} g_2\right)^*, \qquad P_4 = \left(\partial_{R_{\text{iron}}} g_2\right)^*. \qquad (19)$$

$$c_{11} = -\alpha_{1,\text{public}}\,\epsilon/r, \qquad c_{12} = -\alpha_{2,\text{public}}\,\epsilon/r,$$
$$c_{21} = 1/(\alpha_{1,\text{growth}}\gamma r), \qquad c_{22} = 1/(\alpha_{2,\text{growth}}\gamma r).$$

385   Then it can be proven that under the setting of $\alpha_{1,\text{growth}} < \alpha_{2,\text{growth}}$, the criteria (2) $a_4 > 0$
386   requires that:

$$\alpha_{1,\text{public}} > \alpha_{2,\text{public}}. \tag{20}$$

387   In the iron-partition model, Eq. (7) and Eq. (20) suggest that, if two strains should stably
388   coexist, not only the strain that invests fewer resources in growth must invest more in
389   private siderophores, but also play a more cooperative role.
390
391   The remaining two criteria, (3) $a_3 > 0$ and (4) $a_1 a_2 a_3 - a_1^2 a_4 - a_3^2 > 0$, differentiate
392   between the classical model and the iron-partition model. By definitions in Eq. (12)-(15),
393   equations for these two criteria are the additive or multiplicative combination of a_4 with
394   other positive definite terms involving c_{11} and c_{12} (details in Supplementary Material,
395   Section 3):

$$a_3 = d\big(m_1^*(c_{11}P_1 + c_{21}dP_2) + m_2^*(c_{12}P_3 + c_{22}dP_4)\big) + \frac{a_4}{d}. \tag{21}$$

$$a_1 a_2 a_3 - a_1^2 a_4 - a_3^2 \tag{22}$$

$$\begin{aligned}
&= (2d^3 + a_3)\Big(m_1^*(c_{11}P_1 + dc_{21}P_2)(d + c_{21}P_2 m_1^*) \\
&\quad + m_2^*\big(c_{12}(dP_3 + c_{21}P_1 P_4 m_1^*) \\
&\quad + c_{22}(d^2 P_4 + P_2(c_{11}P_3 + 2dc_{21}P_4)m_1^*)\big) \\
&\quad + m_1^{*2}c_{22}P_4(c_{12}P_3 + dc_{22}P_4)\Big).
\end{aligned}$$

396
In the classical consumer resource model, c_{ij} describes the consumption of species i on
resource j, which is always positive. Consequently, once the criteria (2) $a_4 > 0$ holds,
criteria (3) $a_3 > 0$ and criteria (4) $a_1 a_2 a_3 - a_1^2 a_4 - a_3^2 > 0$ are satisfied automatically.
Actually, the determination of the fixed point stability only depends on the tangent slope of
growth contours $(\frac{P_1}{P_2}, \frac{P_3}{P_4})$ and the slope of the consumption vector $(\frac{c_{21}}{c_{11}}, \frac{c_{22}}{c_{12}})$. If the criteria in
Eq.17-18 are satisfied, the coexistence is stable as long as the supply vector locates
within the sector area formed by the reverse extension of the two consumption vectors.
However, in the iron-partition model, the fulfillment of criteria (2) no longer guarantees the
satisfaction of criteria (3) and (4). Even when Eq.(17)-(18) are satisfied and the supply
vector locates within the sector formed by consumption vectors, changes in the
iron-supply can still change the stability of the growth contour intersection, driving the
system into other forms of dynamics instead of stable coexistence. As shown in Figure 3A,
in the chemical space, the growth contours of a partial cooperator and the pure cheater
intersect. At the crossing point, the consumption vector of the partial cooperators points to
the upper-left direction, suggesting it consumes iron while creating public siderophores.
The consumption vector of the pure cheater horizontally points to the left, as it solely
consumes iron but does not contribute to public siderophore. These sectors formed by two
consumption vectors overlap a region on the iron-supply axis (x-axis). Nevertheless,
within this region of iron-supply, the stability of the fixed point changes: when the iron

supply is low, the system will still collapse to extinction even if the supply (black circles,
Fig. 3C) locates within the sector zone; oscillation begins as the iron supply increases
(yellow circles, Fig. 3D); and eventually, stable coexistence can be reached when the iron
supply is high (blue circles, Fig. 3E).
The collapse of the system in Figure 3C can be attributed to the changes in criteria (3)
$a_3 > 0$. Due to the negative signs of c_{11} and c_{12}, the first term of a_3, $d\big(m_1^*(c_{11}P_1 +$

$c_{21}dP_2) + m_2^*(c_{12}P_3 + c_{22}dP_4)\big)$ now has an uncertain sign. If $P_1 = \big(\partial_{R_1}g_1\big)^*$ or $P_3 =$

$\big(\partial_{R_1}g_2\big)^*$ are sufficiently large, indicating that strains are highly sensitive to free iron, the

first term might be negative, and is able to drive the whole equation of a_3 into the
negative regime. Moreover, the sign of the first term also relies on m_1^* and m_2^*, both of
which are dependent on the iron-supply concentration. As shown in Region 2 in Figure 3B,
the sign of a_3 remains negative when a_4 is positive, which contributes to the extinction
of both species even when the iron supply falls within the sector formed by the reverse
extension of consumption vectors (Fig. 3C).
Similarly (See Supplementary Materials, Section 3.2), with negative c_{11} and c_{12}, criteria
(4) in the iron-partition model now has an uncertain sign, with complex dependence on
m_1^* and m_2^*. When criteria (4) remains negative (Region 3) while the other criteria are
fulfilled(Region 3, Fig. 3B), the fixed point remains unstable, and the system oscillates.
In the classical model, due to the positive sign of c_{11} and c_{12}, Region 2 and Region 3 in
Figure 3F would not exist. As the existence of Region 2 and Region 3 is the result of
negative c_{11} and c_{12}, such extinction and oscillation dynamics are not exceptional. Two
partial cooperators with distinct allocation strategies may also undergo similar transitions,
with the dynamics experiencing extinction, oscillation, coexistence, and exclusion as the
iron supply increases (Fig. S3).

Comprehensive assessment of strategies under within-species and

between-species competitions.

In our model, distinct strains within the same species are distinguished by distinct
resource allocation strategies, and their public siderophores can be shared. Parameters
such as the cost of siderophores and the growth coefficient can vary between species. In
addition, we assumed that siderophores produced by different species can not be shared.
We systematically evaluated the competitiveness of strategies within and between
species based on these assumptions.
448
First, the competition between all potential strategies against the pure cheater of the same
species exhibits complex dynamics (as illustrated in the ternary plot of Fig. 4A): the pure
cheater does trigger extinction for highly cooperative strains (high in α_{public} and low in
α_{growth}). However, over a broad range of strategy space, partial cooperator strategies can
coexist stably or dynamically with pure cheater strategies. Moreover, this coexistence can
exist in a wide range of different parameters and public siderophore production and

recycling sets (Fig. S5, S6). Self-seeker strategies can even result in the exclusion of the
pure cheater. In terms of coexistence, increasing investment in public siderophores
increases the likelihood of oscillations with the cheater, whereas increasing investment in
private siderophores increases the tendency of stable coexistence.

We evaluated a species' "non-invasive strategy" by employing the invasion chain
method(Taillefumier, Posfai et al. 2017): starting with one of the arbitrarily chosen
strategies, we added strains with the highest growth rate in the existing chemical
environment into the environment until no species could be added. A typical invasion
chain is depicted in Figure 4B: the pure cheater has the highest growth rate in the
environment shaped by the initial partial cooperator, but in the environment co-created by
these two strains, the self-seekers ($\alpha_{\text{public}} = 0, \alpha_{\text{private}} > 0$) have a growth advantage,
allowing them to invade and create an environment devoid of public siderophores. In such
an environment, only self-seekers can survive, which brings the invasion chain to an end.
In summary, within the same species, the self-seeker strategy is the most resistant
against pure cheater strains and is "evolutionarily optimal", because it generates a
chemical environment that cannot be invaded by other strategies. Also, the self-seeker

strategy can establish a community with a lower initial population, which benefits the start

of colonization (Fig. S7).

However, when evaluating a species' overall competitiveness, the self-seeker strategy is
not always superior. For example, in evaluating the steady-state size of the population
composed of a single strain (Fig. 4C), strategies that are close to the pure cheater have a
larger overall population size. Meanwhile, in terms of resilience to harsher external
environments, pure cooperators are more resistant to increased dilution (Fig. 4D).

Regarding the competition between species using different public siderophores, strains
that invest more in public siderophores are more capable of invading another species (Fig.
S4A), whereas strains that invest less in private siderophores are more resistant to
invasion by another species (Fig. S4C). Self-seeker strategies, on the other hand, are not
effective both in invasion and resistance(Fig. S4B, D).

Discussion

Understanding diversity has long been a cornerstone of microbial ecology (Delmont, Robe
et al. 2011). While the number of species may be limited by chemical dimensions,
microorganisms are capable of expanding this limit by actively generating chemical
diversity in their microhabitat. In this work, we modeled the specific system of
siderophore-mediated iron competition in order to investigate more general questions: first,
how could cooperators responsible for chemical diversity resist extinction induced by
cheaters; and second, what is the ecological consequence of microbes creating new
chemical dimensions. For the first question, we suggested the privatization of
siderophores acting as a "game changer" to allow partial cooperators advantage over
pure cheaters, and derived that a necessary condition for coexistence is that the
cooperators who invest more in public siderophores also need to invest more in private
siderophores. Concerning the second question, we analytically compared the difference in
stability criteria between the traditional consumer resource models and the "resource
partition model," in which organisms not only consume but also produce resources. In
addition, the public siderophore does not become a "resource" until it forms an association
with ferric iron, the actual resource that microorganisms require. This is an additional layer
of meaning for the term "resource-partition models": externally supplied resources (iron)
and microbe-generated resources (siderophore) interact to form the actual resources
(iron-associated siderophores) taken up by the microorganisms.

Microbes interact via influencing their shared environment (McGill, Enquist et al. 2006).
Classical ecological models emphasize the "consumption" aspect of such influences
(Tikhonov and Monasson 2017, Altieri and Franz 2019). With the rapid development of
microbiology, it has become increasingly apparent that bacteria have enormous potential
for introducing new chemicals into their microhabitats(Gavriilidou, Kautsar et al. 2022).
Siderophore is just one of the many secondary metabolites that microbes actively
produce for their own benefit. Antibiotics, bacteriocins, signaling molecules, and even
bacterial vesicles can all be considered "chemical dimensions" generated by microbes
themselves (Bajić, Rebolleda-Gómez et al. 2021, Niehus, Oliveira et al. 2021). Regarding
these "self-generated dimensions," a general ecological framework has yet to be
established. The conventional consumer resource model provides intuitive coexistence
criteria (i.e. species should preferentially consume the resource that limits their growth),
with consumption vectors clearly segregating supply conditions into zones of
the same stability (Tilman 1982, Koffel, Daufresne et al. 2016). In this "resource partition
model", however, due to the reversed direction of consumption vectors, there are
additional parameter areas where the RH criterion can change signs. Consequently, in the
regions of steady-state equilibrium in the classical model, non-equilibrium dynamics
become possible. In general, our analysis suggested that an ecosystem with
"self-generated dimensions" tends to be more dynamic and complex.

Oscillation usually bridges two distinctive states and often plays critical roles in various
biological systems (Goldbeter 2008). Notably, the oscillation in our model differs from the
oscillation in the work of Huisman et al. where the oscillation bridges between the stable
equilibrium and the chaos with higher-than-CEP dynamical coexistence (Huisman and
Weissing 1999, Huisman and Weissing 2001). In our model, the parameter region of
oscillation locates between total extinction and stable coexistence between partial
cooperators and cheaters. Oscillation here is more of a "danger zone" indicating that the
system is on the verge of collapse, similar to the early warning signatures of ecosystems
(Carpenter, Cole et al. 2011). Indeed, our continuous equations assume that organisms
can recover from arbitrarily small values, but the troughs of oscillatory dynamics increase
the probability of stochastic extinction in real systems (Reichenbach, Mobilia et al. 2006,
Ovaskainen and Meerson 2010). Recent studies in the field of game theory, however,
have detected oscillations when the environment is made explicit and have shown that
oscillatory dynamics prevent the extinction of cooperators (Weitz, Eksin et al. 2016, Hilbe,
Šimsa et al. 2018, Tilman, Plotkin et al. 2020). Given the specificity of oscillatory dynamics
in ecology and the propensity of resource partition models to enter the oscillation zone, it
would be intriguing to explore the facilitative or destructive functions of oscillations in
diverse ecological systems with chemical innovations.
547
In the specific game of microbial iron competition, the pervasiveness of siderophore
production remains the subject of continuing investigation (Barber and Elde 2015, Kramer,
Özkaya et al. 2020, Lee, Eldakar et al. 2021). Our work emphasized that private
siderophores only provide possible solutions in some species, such as marine microbes
or mycobacterium, where the siderophores can be modified into membrane-attached
form.
Other kinds of siderophore privatization, such as keeping siderophores intracellularly for
relieving oxidative stress, have been proposed to confer cheater resistance (Jin, Li et al.
2018). Actually, we hypothesize that the division of the iron resource by siderophores
provides a universal mechanism for "resource privatization": numerous siderophores exist
in the natural world, each with their own specific receptors (Cornelis and Matthijs 2002, Jin,
Li et al. 2018). For a given type of siderophore, microorganisms with corresponding
receptors can share it as a public good, whereas microbes without comparable receptors
perceive it as inaccessible "private goods" (Leventhal, Ackermann et al. 2019).
Theoretical models suggested that the populations of different cooperators utilizing
distinct siderophores can be regulated by their cheaters, similar to how parasites impose
negative frequency selection (Lee, van Baalen et al. 2012). In addition, a model with
different types of siderophores suggested that coexistence between cooperators and
cheaters is possible if a "loner" uses a kind of relatively inefficient siderophores (Inglis,
Biernaskie et al. 2016). In the future, it would be exciting to investigate iron interactions in
a more realistic biological setting and systematically investigate the ecological
consequences of microbial chemical innovation.
570
571

Data Availability

The source code and parameters used are available in the supplementary material.

Acknowledgements

This work was supported by grants from Peking-Tsinghua Center for Life Sciences.

Funding

This work was supported by the National Natural Science Foundation of China (No. 32071255) and Clinical Medicine Plus X - Young Scholars Project, Peking University, the Fundamental Research Funds for the Central Universities (No. PKU2022LCXQ009).

Author contributions

Jiqi Shao performed the majority of computational and mathematical analysis in this research and drafted the manuscript. Nan Rong performed the preliminary computational analysis in this research and drafted the manuscript. Zhenchao Wu, Beibei Liu, and Ning Shen offered insightful comments and assisted in revising the manuscript. Zhiyuan Li conceptualized and oversaw the project and revised the manuscript. All authors gave final approval for publication and agreed to be held accountable for the work performed herein.

Competing interests

The authors declare no competing interests.

Figures

Figure 1. Privatization of siderophore enables the coexistence between partial

cooperators and pure cheaters.

(A) The schematic diagram of a chemostat model.
(B) The schematic diagram showing two iron uptake fluxes from the public siderophores
(left) and private siderophores (right).
(C) The strategy space (the strategic phase diagram) by ternary plot, showing all resource

allocation strategies $\vec{\alpha} = \left(\alpha_{\text{growth}}, \alpha_{\text{private}}, \alpha_{\text{public}}\right)$. The model has four typical strategies,

distinguished according to whether to produce public siderophores and whether to use

private siderophores. Pure cooperators (yellow) only produce and use public siderophores,

while the pure cheater (orange) only use public siderophores and put all resources into

α_{growth}; partial cooperators (green) produce and use both public and private siderophores,

while self-seekers (blue) only produce private siderophores and use both public and

private siderophores.

(D) The schematic illustration of the four typical strategies shown in (C).

(E-F) The growth contours in the chemical space, with the concentration of Fe^{3+} by the

x-axis and the concentration of public siderophore by the y-axis. (E) shows the growth

contours of a pure cooperator and the pure cheater, and (F) shows the growth contours of

a partial cooperator and the pure cheater (color scheme same as that in (C) and (D)). The

blue area represents a growth-advantageous zone in which the partial cooperator

outperforms the pure cheater. Inserts are the biomass time-courses of species competing

in the chemostat.

Figure 2. The partial cooperators and cheaters generate rich ecological dynamics

(A) The partial cooperators can oscillate with the cheater. In the chemical space, the
intersection between the growth contours of a partial cooperator (blue) and a pure cheater
(red) is represented by a black circle, and the dynamical trajectory beginning is
represented by a black line. The background color indicates the difference in growth rate
between partial cooperators and cheaters. Insert shows the competition dynamics of the
two species.

(B) The partial cooperators can exclude the cheater. Same as (A), but the partial
cooperators allocate more resources to private siderophores.

(C) The phase diagram illustrates the mechanism of oscillation shown in (A). In the upper
panel, the chemical space is divided into four regions with varying relative finesses
(growth rate relative to dilution rate) between the cheater and the partial cooperators,
denoted by the symbols + and -. The background color of the lower panel corresponds to
the regions in the chemical space depicted in the upper panel.

(D-E) Bifurcation diagrams for the system in (A) as the dilution rate (D) and the Fe^{3+}
supply changes (E). Empty circles represent the minimum and maximum of the oscillation,
while solid dots represent a steady state.

Figure 3. The self-created dimension enables oscillation and coexistence.

(A) The growth contours of a partial cooperator (blue) and the cheater (red), and their consumption vectors at the intersection. Different types of dynamics induced by different supplies of Fe^{3+} are indicated by the colors of circles along the x-axis. Black indicates the extinction of both species; Yellow indicates oscillatory dynamics; Blue indicates steady-state coexistence.

(B) The values of three stability criteria as the iron supply increases along the x-axis. The signs of three stability criteria divide the x-axis into four regions, as indicated by the text.

(C-E) Exemplary time-courses of the three types of dynamics induced by different supplies of Fe^{3+}, such as extinction (C), oscillation (D), and coexistence (E).

(F) The signs of all four stability criteria in four regions shown in (B).

Figure 4. Assessment of all possible strategies, by their interplay with cheaters, evolutionary stability, stable population, and critical dilution rate.

(A) How different strategies interact with the pure cheater. Each strategy's competition outcomes with the pure cheater are indicated by dot color in the ternary plot (light gray: non-viable by itself; deep gray: viable by itself but extinct with the cheater; yellow: oscillate with the cheater; blue: stably coexist with the cheater; red: exclude the cheater).

(B) The chain of invasion in the strategy space. For each arrow, the dot at the beginning of the arrow is the initial strategy that creates the steady-state environment, and the dot at the endpoint of the arrow indicates the strategy with the maximal growth rate in the formal environment. The endpoint of the whole invasion chain is the evolutionarily stable strategy that can not be invaded by any other strategies. The path starts with a partial cooperator strategy(green dot, $\vec{\alpha} = (0.6, 0.2, 0.2)$), then directs to the pure cheater strategy(orange dot), and ends at the self-seeker strategy(blue dots, $\vec{\alpha} = (0.65, 0.35, 0)$).

(C) The stable population for different strategies when existing alone.

(D) The maximal dilution rate for non-zero biomass in a steady-state chemostat.

678 **Reference**

679 Allison, S. D. (2005). "Cheaters, diffusion and nutrients constrain decomposition by
680 microbial enzymes in spatially structured environments." Ecology Letters **8**(6): 626-635.
681

682 Altieri, A. and S. Franz (2019). "Constraint satisfaction mechanisms for marginal stability
683 and criticality in large ecosystems." Physical Review E **99**(1): 010401.
684

685 Andrews, S. C., et al. (2003). "Bacterial iron homeostasis." FEMS Microbiology Reviews
686 **27**(2-3): 215–237.
687

688 Bajić, D., et al. (2021). "The macroevolutionary consequences of niche construction in
689 microbial metabolism." Frontiers in Microbiology: 2641.
690

691 Barber, M. F. and N. C. Elde (2015). "Buried Treasure: Evolutionary Perspectives on
692 Microbial Iron Piracy." Trends in Genetics **31**(11): 627–636.
693

694 Boyd, P. W. and M. J. Ellwood (2010). "The biogeochemical cycle of iron in the ocean."
695 Nature Geoscience.
696

697 Buckling, A., et al. (2010). "Siderophore-mediated cooperation and virulence in
698 Pseudomonas aeruginosa." Fems Microbiology Ecology **62**(2): 135-141.
699

700 Butaitė, E., et al. (2017). "Siderophore cheating and cheating resistance shape
701 competition for iron in soil and freshwater Pseudomonas communities." Nature
702 Communications **8**(1): 414.
703

704 Carpenter, S. R., et al. (2011). "Early warnings of regime shifts: a whole-ecosystem
705 experiment." Science **332**(6033): 1079-1082.
706

707 Cordero, O. X., et al. (2012). "Public good dynamics drive evolution of iron acquisition
708 strategies in natural bacterioplankton populations." Proceedings of the National Academy
709 of Sciences **109**(49): 20059–20064.
710

Cornelis, P. and S. Matthijs (2002). "Diversity of siderophore-mediated iron uptake
systems in fluorescent pseudomonads: not only pyoverdines." Environ Microbiol **4**(12):
787-798.
Fluorescent pseudomonads are gamma-proteobacteria known for their capacity to
colonize various ecological niches. This adaptability is reflected by their
sophisticated and diverse iron uptake systems. The majority of fluorescent
pseudomonads produce complex peptidic siderophores called pyoverdines or
pseudobactins, which are very efficient iron scavengers. A tremendous variety of
pyoverdines has been observed, each species producing a different pyoverdine.
This variety can be used as an interesting tool to study the diversity and taxonomy

of fluorescent pseudomonads. Other siderophores, including newly described ones, are also produced by pseudomonads, sometimes endowed with interesting properties in addition to iron scavenging, such as formation of complexes with other metals or antimicrobial activity. Factors other than iron limitation, and different regulatory proteins also seem to influence the production of siderophores in pseudomonads and are reviewed here as well. Another peculiarity of pseudomonads is their ability to use a large number of heterologous siderophores via different TonB-dependent receptors. A first genomic analysis of receptors in four different fluorescent pseudomonads suggests that their siderophore ligand repertoire is likely to overlap, and that not all receptors recognize siderophores as ligands.

Cremer, J., et al. (2019). "Cooperation in microbial populations: theory and experimental model systems." Journal of molecular biology **431**(23): 4599-4644.

Czaran, T. L., et al. (2002). "Chemical warfare between microbes promotes biodiversity." Proceedings of the National Academy of Sciences.

Delmont, T. O., et al. (2011). "Accessing the soil metagenome for studies of microbial diversity." Applied and environmental microbiology **77**(4): 1315-1324.

Dutkiewicz, S., et al. (2020). "Dimensions of marine phytoplankton diversity." Biogeosciences **17**(3): 609-634.

Emerson, D., et al. (2012). "The microbial ferrous wheel: iron cycling in terrestrial, freshwater, and marine environments." Frontiers in Microbiology **3**: 383.

Erez, A., et al. (2020). "Nutrient levels and trade-offs control diversity in a serial dilution ecosystem." Elife **9**: e57790.

Estrela, S., et al. (2022). "Diversity begets diversity under microbial niche construction." bioRxiv.

Fischbach, M. A. and J. A. Segre (2016). "Signaling in host-associated microbial communities." Cell **164**(6): 1288-1300.

Gavriilidou, A., et al. (2022). "Compendium of specialized metabolite biosynthetic diversity encoded in bacterial genomes." Nature Microbiology **7**(5): 726-735.

Goldbeter, A. (2008). "Biological rhythms: clocks for all times." Current biology **18**(17): R751-R753.

Goldford, J. E., et al. (2018). "Emergent simplicity in microbial community assembly." Science **361**(6401): 469-474.

Gore, J., et al. (2009). "Snowdrift game dynamics and facultative cheating in yeast." Nature **459**(7244): 253-256.

The origin of cooperation is a central challenge to our understanding of evolution. The fact that microbial interactions can be manipulated in ways that animal interactions cannot has led to a growing interest in microbial models of cooperation and competition. For the budding yeast Saccharomyces cerevisiae to grow on sucrose, the disaccharide must first be hydrolysed by the enzyme invertase. This hydrolysis reaction is performed outside the cytoplasm in the periplasmic space between the plasma membrane and the cell wall. Here we demonstrate that the vast majority (approximately 99 per cent) of the monosaccharides created by sucrose hydrolysis diffuse away before they can be imported into the cell, serving to make invertase production and secretion a cooperative behaviour. A mutant cheater strain that does not produce invertase is able to take advantage of and invade a population of wild-type cooperator cells. However, over a wide range of conditions, the wild-type cooperator can also invade a population of cheater cells. Therefore, we observe steady-state coexistence between the two strains in well-mixed culture resulting from the fact that rare strategies outperform common strategies-the defining features of what game theorists call the snowdrift game. A model of the cooperative interaction incorporating nonlinear benefits explains the origin of this coexistence. We are able to alter the outcome of the competition by varying either the cost of cooperation or the glucose concentration in the media. Finally, we note that glucose repression of invertase expression in wild-type cells produces a strategy that is optimal for the snowdrift game-wild-type cells cooperate only when competing against cheater cells.

Granato, E. T., et al. (2019). "The evolution and ecology of bacterial warfare." Current biology **29**(11): R521-R537.

Gu, S., et al. (2020). "Competition for iron drives phytopathogen control by natural rhizosphere microbiomes." Nature Microbiology **5**(8): 1002–1010.

Gupta, D., et al. (2021). "Effective Resource Competition Model for Species Coexistence." Physical review letters **127**(20): 208101.

Hagstrom, G. I. and S. A. Levin (2017). "Marine ecosystems as complex adaptive systems: emergent patterns, critical transitions, and public goods." Ecosystems **20**(3): 458-476.

Hardin and G. (1960). "The Competitive Exclusion Principle." Science **131**(3409): 1292-1297.

Hardin, G. (1968). "The Tragedy of the Commons." Science **162**(3859): 1243-1248.

809 Hilbe, C., et al. (2018). "Evolution of cooperation in stochastic games." <u>Nature</u> **559**(7713):
810 246-249.
811
812 Ho, P.-Y., et al. (2022). "Competition for fluctuating resources reproduces statistics of
813 species abundance over time across wide-ranging microbiotas." <u>Elife</u> **11**: e75168.
814
815 Huisman, J. and F. J. Weissing (1999). "Biodiversity of plankton by species oscillations
816 and chaos." <u>Nature</u> **402**(6760): 407–410.
817
818 Huisman, J. and F. J. Weissing (2001). "Biological conditions for oscillations and chaos
819 generated by multispecies competition." <u>Ecology</u> **82**(10): 2682-2695.
820
821 Inglis, R. F., et al. (2016). "Presence of a loner strain maintains cooperation and diversity
822 in well-mixed bacterial communities." <u>Proceedings. Biological sciences</u> **283**(1822).
823
824 Jin, Z., et al. (2018). "Conditional privatization of a public siderophore enables
825 Pseudomonas aeruginosa to resist cheater invasion." <u>Nature Communications</u> **9**(1): 1-11.
826
827 Julou, T., et al. (2013). "Cell-cell contacts confine public goods diffusion inside
828 Pseudomonas aeruginosa clonal microcolonies." <u>Proceedings of the National Academy of</u>
829 <u>Sciences</u> **110**(31): 12577–12582.
830
Koffel, T., et al. (2016). "Geometrical envelopes: Extending graphical contemporary niche
theory to communities and eco-evolutionary dynamics." <u>J Theor Biol</u> **407**: 271-289.
Contemporary niche theory is a powerful structuring framework in theoretical
ecology. First developed in the context of resource competition, it has been
extended to encompass other types of regulating factors such as shared
predators, parasites or inhibitors. A central component of contemporary niche
theory is a graphical approach popularized by Tilman that illustrates the different
outcomes of competition along environmental gradients, like coexistence and
competitive exclusion. These food web modules have been used to address
species sorting in community ecology, as well as adaptation and coexistence on
eco-evolutionary time scales in adaptive dynamics. Yet, the associated graphical
approach has been underused so far in the evolutionary context. In this paper, we
provide a rigorous approach to extend this graphical method to a continuum of
interacting strategies, using the geometrical concept of the envelope. Not only
does this approach provide community and eco-evolutionary bifurcation diagrams
along environmental gradients, it also sheds light on the similarities and
differences between those two perspectives. Adaptive dynamics naturally merges
with this ecological framework, with a close correspondence between the
classification of singular strategies and the geometrical properties of the envelope.
Finally, this approach provides an integrative tool to study adaptation between
levels of organization, from the individual to the ecosystem.
852

853 Kramer, J., et al. (2020). "Bacterial siderophores in community and host interactions."
854 Nature Reviews Microbiology **18**(3): 152–163.
855
856 Lafferty, K. D., et al. (2015). "A general consumer-resource population model." Science
857 **349**(6250): 854-857.
858
859 Lee, I. P. A., et al. (2021). "Bacterial cooperation through horizontal gene transfer." Trends
860 in ecology & evolution.
861
862 Lee, W., et al. (2012). "An evolutionary mechanism for diversity in siderophore-producing
863 bacteria." Ecology Letters **15 2**: 119-125.
864
865 Leventhal, G. E., et al. (2019). "Why microbes secrete molecules to modify their
866 environment: the case of iron-chelating siderophores." Journal of the Royal Society
867 Interface **16**(150): 20180674.
868
869 Ley, R. E., et al. (2006). "Ecological and evolutionary forces shaping microbial diversity in
870 the human intestine." Cell **124**(4): 837-848.
871
872 Li, Z., et al. (2020). "Modeling microbial metabolic trade-offs in a chemostat." PLoS
873 Computational Biology **16**(8): e1008156.
874
875 Lv, H., et al. (2014). "Metabolomic analysis of siderophore cheater mutants reveals
876 metabolic costs of expression in uropathogenic Escherichia coli." Journal of Proteome
877 Research **13**(3): 1397–1404.
878
879 Macarthur, R. and R. Levins (1964). "Competition, Habitat Selection, and Character
880 Displacement in a Patchy Environment." Proc Natl Acad Sci U S A **51**: 1207-1210.
881
882 Martinez, J. S., et al. (2003). "Structure and membrane affinity of a suite of amphiphilic
883 siderophores produced by a marine bacterium." Proceedings of the National Academy of
884 Sciences of the United States of America **100**(7): 3754–3759.
885
886 May, R. and P. Allen (1977). "Stability and Complexity in Model Ecosystems." Systems,
887 Man and Cybernetics, IEEE Transactions on **44**: 887-887.
888
889 McGill, B. J., et al. (2006). "Rebuilding community ecology from functional traits." Trends
890 in ecology & evolution **21**(4): 178-185.
891
892 Miransari, M. (2013). "Soil microbes and the availability of soil nutrients." Acta
893 physiologiae plantarum **35**(11): 3075-3084.
894
Niehus, R., et al. (2021). "The evolution of strategy in bacterial warfare via the regulation
of bacteriocins and antibiotics." Elife **10**: e69756.

Niehus, R., et al. (2017). "The evolution of siderophore production as a competitive trait."
Evolution; international journal of organic evolution **71**(6): 1443–1455.

Ovaskainen, O. and B. Meerson (2010). "Stochastic models of population extinction."
Trends in ecology & evolution **25**(11): 643-652.

Reichenbach, T., et al. (2006). "Coexistence versus extinction in the stochastic cyclic
Lotka-Volterra model." Physical Review E **74**(5): 051907.

Ross-Gillespie, A., et al. (2007). "Frequency dependence and cooperation: theory and a
test with bacteria." The American Naturalist **170**(3): 331–342.

Ross-Gillespie, A. and R. Kümmerli (2014). "Collective decision-making in microbes."
Frontiers in Microbiology **5**: 54.

Roy, S. and J. Chattopadhyay (2007). "The stability of ecosystems: a brief overview of the
paradox of enrichment." Journal of biosciences **32**(2): 421-428.

Schmidt, R., et al. (2019). "Microbe-driven chemical ecology: past, present and future."
The ISME journal **13**(11): 2656-2663.

Scholz, R. L. and E. P. Greenberg (2015). "Sociality in Escherichia coli: Enterochelin Is a
Private Good at Low Cell Density and Can Be Shared at High Cell Density." Journal of
Bacteriology **197**(13): 2122–2128.

Sexton, D. J. and M. Schuster (2017). "Nutrient limitation determines the fitness of
cheaters in bacterial siderophore cooperation." Nature Communications **8**(1): 230.

Smith, P. and M. Schuster (2019). "Public goods and cheating in microbes." Current
biology **29**(11): R442-R447.

Souza, M. O., et al. (2009). "Evolution of cooperation under N-person snowdrift games."
Journal of Theoretical Biology **260**(4): 581-588.

Taillefumier, T., et al. (2017). "Microbial consortia at steady supply." Elife **6**: e22644.

Tikhonov, M. and R. Monasson (2017). "Collective phase in resource competition in a
highly diverse ecosystem." Physical review letters **118**(4): 048103.

Tilman, A. R., et al. (2020). "Evolutionary games with environmental feedbacks." Nature
Communications **11**(1): 1-11.

Tilman, D. (1982). Resource competition and community structure. Princeton; Guildford,

Princeton University Press.

Weitz, J. S., et al. (2016). "An oscillating tragedy of the commons in replicator dynamics
with game-environment feedback." <u>Proc Natl Acad Sci U S A</u> **113**(47): E7518-E7525.

A tragedy of the commons occurs when individuals take actions to maximize their
payoffs even as their combined payoff is less than the global maximum had the
players coordinated. The originating example is that of overgrazing of common
pasture lands. In game-theoretic treatments of this example, there is rarely
consideration of how individual behavior subsequently modifies the commons and
associated payoffs. Here, we generalize evolutionary game theory by proposing a
class of replicator dynamics with feedback-evolving games in which
environment-dependent payoffs and strategies coevolve. We initially apply our
formulation to a system in which the payoffs favor unilateral defection and
cooperation, given replete and depleted environments, respectively. Using this
approach, we identify and characterize a class of dynamics: an oscillatory tragedy
of the commons in which the system cycles between deplete and replete
environmental states and cooperation and defection behavior states. We
generalize the approach to consider outcomes given all possible rational choices
of individual behavior in the depleted state when defection is favored in the replete
state. In so doing, we find that incentivizing cooperation when others defect in the
depleted state is necessary to avert the tragedy of the commons. In closing, we
propose directions for the study of control and influence in games in which
individual actions exert a substantive effect on the environmental state.

968   .

971