1

# ARTseq-FISH reveals position-dependent differences in gene expression of micropatterned mESCs

Xinyu Hu[1,2], Bob van Sluijs[1,2], Óscar García-Blay[1], Yury Stepanov[1], Koen Rietrae[1], Wilhelm T.S. Huck[1,*], Maike M. K. Hansen[1,*].

**Affiliations:**
[1]Institute for Molecules and Materials, Radboud University, Heyendaalseweg 135, 6525 AJ Nijmegen, the Netherlands.
[2]These authors contributed equally to this work: Xinyu Hu, Bob van Sluijs.
*Correspondence to: w.huck@science.ru.nl (W.T.S.H); maike.hansen@ru.nl (M.M.K.H)

**Abstract:** Differences in gene-expression profiles between individual cells can give rise to distinct cell fate decisions. Yet how localisation on a micropattern impacts initial changes in mRNA, protein, and phosphoprotein abundance remains unclear. To identify the effect of cellular position on gene expression, we developed a scalable antibody and mRNA targeting sequential fluorescence *in situ* hybridization (ARTseq-FISH) method capable of simultaneously profiling mRNAs, proteins, and phosphoproteins in single cells. We studied 67 (phospho-)protein and mRNA targets in individual mouse embryonic stem cells (mESCs) cultured on circular micropatterns. ARTseq-FISH reveals relative changes in both abundance and localization of mRNAs and (phospho-)proteins during the first 48 hours of exit from pluripotency. We confirm these changes by conventional immunofluorescence and time-lapse microscopy. Chemical labelling, immunofluorescence, and single-cell time-lapse microscopy further show that cells closer to the edge of the micropattern exhibit increased proliferation compared to cells at the centre. Together these data suggest that while gene expression is still highly heterogeneous position-dependent differences in mRNA and protein levels emerge as early as 12 hours after LIF withdrawal.

**Introduction**

A complex interplay of signalling pathways and transcriptional networks guides morphological transformations during mammalian development[1-3]. During early development the balance between key signalling pathways defines the cellular choice between differentiation and self-renewal[4-6]. Therefore, early changes in mRNA and protein expression levels can influence cell fate decisions[7-9]. Micropatterning has proven to be a powerful tool in unveiling position-dependent differentiation that recapitulates the spatial patterning of germ layers observed in early embryonic development[10-17]. Measuring mRNA and protein levels as mouse embryonic stem cells (mESCs) exit pluripotency will offer insights into how micropattern positioning affects early gene expression changes. This requires a scalable image-based approach capable of simultaneously detecting and quantifying mRNAs and proteins with a high spatial resolution across large areas.

Recently, next generation sequencing (NGS)-based spatial-omics methods have greatly increased the multiplexing capacities of RNA or protein detection, albeit with low efficiency due to *in situ* reverse transcription and limited resolution of the beads used to capture targets[18-22]. Techniques that resolve the resolution issue, predominantly for RNA, are image-based spatial-omics methods. Image-based transcriptomic methods have greatly expanded the number of transcripts that can be resolved, using sequential fluorescence *in situ* hybridization (FISH)[23-27] and barcoding approaches[24-27]. These methods have been combined with immunofluorescence (IF), but multiplexed protein detection remains a challenge[28]. For example, targeting and visualizing RNAs and proteins often require separate sequential experimental and imaging steps, doubling the experimental time[26]. Thus, scaling of these techniques in terms of multiplexing capabilities are bottlenecks, limiting the ability to simultaneously visualize mRNAs and proteins in high throughput[29-32]. Furthermore, the resolution of protein detection is often below the level of RNA detection proving a barrier to quantitative insights into protein location or spatial correlation of RNA and protein abundance[24, 26, 28-32]. Conversely, DNA-barcoding dependent methods (for example, CODEX[33], ImmunoSABER[34] and InSituPlex®[35]) have shown their multiplexing capacity in protein detection, but have thus far not been integrated with RNA detection[36, 37].

Here we develop and validate a multiplexed spatial profiling method: antibody-mRNA targeted sequential FISH (ARTseq-FISH). This method enables simultaneous quantification of mRNA, protein, and phosphoprotein levels in individual mouse embryonic stem cells (mESCs) with high spatial resolution. The gene expression profiles of 67 targets were analysed during the initial 48 hours of differentiation, allowing us to capture relative changes in mRNA and protein abundance as mESCs exit pluripotency. Similar to previous evidence[15, 16], ARTseq-FISH reveals distinct gene-expression profiles in cells located at different positions on the micropattern. Specifically, position-dependent differences in gene expression profiles emerge already during the first 12 hours of Leukemia inhibitory factor (LIF) withdrawal. Finally, we confirm that cells closer to the edge of the micropattern display increased proliferation compared to cells at the centre of the micropattern already during the first 12 hours of LIF withdrawal. In summary, here we describe and validate a method that quantifies relative differences in mRNA and (phospho)protein abundance.

## Results

### ARTseq-FISH enables simultaneous visualization of mRNAs, proteins, and phosphoproteins

The ARTseq-FISH workflow comprises a series of consecutive steps (Fig. 1a and d, Supplementary Notes 1, Supplementary Fig. 1-16). In short, fixed cells are stained with DNA-barcoded antibodies against (phospho-)protein targets of choice (Supplementary Table 1-2). The DNA-barcode of each antibody, known not to influence antibody binding efficeincy[38, 39], contains a 10-nt polyadenylation (poly A) linker and a specific 36-nt sequence that is complementary to a corresponding target-specific single stranded DNA (ssDNA) padlock probes (PLPs). On the other hand, cellular mRNAs are directly targeted by one specific ssDNA PLP with a target sequence of 36-nt (Supplementary Fig. 1a-b), resulting in DNA/RNA hybridization. Therefore, the circularizing of the PLPs that hybridize to either proteins or RNAs require T4 DNA ligase and SplintR ligase respectively[40-42] (Fig.1a, Supplementary Fig. 4-6). Next, rolling circle amplification (RCA) generates rolling circle products (RCPs) that amplify the original (phospho-)protein or mRNA target (Fig. 1a and Supplementary Fig. 8). To decode the original target amplified by the RCPs, we first hybridize target-specific bridge probes (BrPs), which are each recognized by unique combinations of fluorescently labelled readout probes (RoPs) (Fig. 1a, Supplementary Fig.1c-d, Supplementary Fig. 9, and Supplementary Table 1), allowing the detection of individual spots for each mRNA and protein (Fig. 1b-c, Supplementary Fig. 17-18). ARTseq-FISH provides similar relative protein abundance compared to classic immunofluorescence (Supplementary Fig. 19). We determined a detection efficiency of ~47% (for *Nanog* mRNA) by comparing mRNA quantification using ARTseq-FISH with conventional single molecule RNA FISH (smFISH)[27] (Supplementary Fig. 20, Supplementary Table 3). Furthermore, ARTseq-FISH enhances the signal ~4-fold with respect to conventional smFISH (Supplementary Fig. 20c-d), resulting in a higher signal-to-noise ratio. On average, ~0.03-0.5 spots per cell detected as false positives from unspecific PLP (protein detection only), BrP, and RoP hybridization (Supplementary Fig. 3), in particular in the presence of true hybridization signal (Supplementary Fig. 18b). Similar to existing methods, we are unable to quantify the degree of false positives coming from unspecific binding of the antibody or PLP and the target detection efficiency is highly antibody[33, 34] and PLP-dependent[40-43] (Supplementary Fig. 10). Lastly, here we specifically employ one PLP per mRNA target, since we are interested in relative mRNA differences. However, should more rigorous quantification of mRNA be necessary at least 3 PLPs per target are likely required[40-43].

Fig. 1: Experimental outline of ARTseq-FISH. a, The experimental workflow of ARTseq-FISH enables detection of different molecular species (mRNA, proteins, and phosphoproteins) simultaneously. Each antibody conjugated to oligonucleotides, targets its associated (phospho-)protein *in situ*. Target-specific PLPs bind to these oligonucleotides as well as cellular mRNA, respectively. Then the PLPs are circularized and amplified into RCPs. A set of target-specific BrPs hybridize to RCPs and then RoPs labelled with fluorophores bind to one of the four flanking sequences on the BrP. b, Representative raw data of simultaneously detecting *Beta-catenin* mRNA, protein, and phosphoprotein with same resolution. Scale bar, 20 µm, 1 µm. c, The relative mean pixel intensity of the hybridization signal's point spread functions (n = 2000) around the local maxima. Comparison of the average observed signal of ARTseq-FISH to smFISH and a 200 nm fluorescent bead. d, The sequential stripping and rehybridization of a new known RoPs allows subsequent decoding of stacked signals.

## Automated analysis and quantification of mRNAs and proteins in single cells

Multiplexed, simultaneous, detection and quantification of mRNAs and (phospho-)proteins over a large area is achieved through serial stripping and rehybridization of RoPs following a predetermined colour barcoding scheme[27] (Fig. 1d, Methods, Supplementary Notes 1-2, Supplementary Fig. 14-16, Supplementary Table 4). The barcode of each target is determined by the presence of a given colour (green, red, or far red) in a sequence of specific hybridization rounds. To correctly decode the detected signal and translate this into mRNA or protein counts for individual cells, we developed an automated image analysis pipeline that returns the abundance of each target per cell.

The pipeline (Fig. 2a, more detailed description in Supplementary Notes 2, SN2 Fig. 1-3, software packages used[44-49]) initiates the analysis by identifying the local maxima per colour channel[50] by denoising and sharpening the image in 6 steps (SN2 Fig. 3-4) and classifying the local maxima as a true hybridization signal. This is done by using a support vector machine

(SVM) model trained on data of true point spread functions, which allows for signal
deconvolution (SN2 Fig. 5-6). Single cell analysis requires the reconstruction of single cells in
3D from a collection of images (Fig. 2a and SN2 Fig. 7-15). However, consistent segmentation
is challenging because a segmentation algorithm that maps onto one nucleus might not map
onto others (SN2 Fig. 8), or a global image filter that cleanly separates the nuclei in one image
might not be optimal for another (SN2 Fig. 10). Therefore, an iterative approach was
implemented, repeatedly segmenting nuclei using different global image filters and
segmentation parameters[51] (SN2 Fig. 9 and 13). We collected the segmentation labels and used
an SVM model to filter out any failed segmentations. The remaining labels were subsequently
used to reconstruct the nuclei in 3D by clustering their centroids with a node-based graph (SN2
Fig. 14-15). After interpolating any missing nucleus segmentations across the z-axis, we
obtained a 3D bounding box. Another SVM model subsequently assessed if this bounding box
contains a single nucleus. Before assigning mRNA and protein counts to a cell, we adjusted for
any drift, by using the segmented nuclei as unique markers to align the images. Once a global
shift was found, we refined the alignment by moving all hybridization signals across a narrow
range of shifted pixels until the signals overlapped (SN2 Fig. 16-17). Ultimately, we decoded
the overlapping signals and assigned each hybridization signal to a particular protein or mRNA
in a specific cell (Fig. 2b, Supplementary Fig. 2, SN2 Fig.18-19). Notably, we assign each
target to a specific cell based on the shortest distance to the nucleus. Therefore, although the
cellular spot count correlates strongly with the nuclear spot count (SN2 Fig. 23), there is a
possibility of incorrect assignments if the distance from the nucleus to the cell membrane varies
significantly between neighbouring cells.

**a**

**1. Detect Hybridization Signals**

**2. AI Based Iterative Segmentation**

i. Filter Image

ii. Segment Nuclei

Change Segmentation Parameters

Change Image Filters

Assess Segmentation

**3. Cluster Segments and Reconstruct Single Cell in 3D**

Centroid

A

B

x

z

y

A    B

**4. Correct Image Drift and Decode Hybridization Signals**

**5. Single Cell Analysis**

z

y

x

**b**

● mRNA
● Protein
● Phosphorylated protein

20 μm

● mRNA

20 μm

● Protein

20 μm

● Phosphorylated protein

20 μm

**Fig. 2: Computational outline of ARTseq-FISH. a,** A condensed summary of the automated computational pipeline that converts the raw microscope images into a single cell dataset, providing the abundance of each target per cell. First the hybridization signal is detected, followed by an iterative AI based segmentation approach, the segments are clustered, and the cells are reconstructed in 3D. Next, an image drift correction is applied to align the hybridization between rounds. In the final step these hybridization signals are decoded and assigned to the targets as well as individual cells. **b,** Reconstructed images show all the detected molecules in the cells cultured on micropattern 48 hours after leukemia inhibitory factor (LIF) withdrawal. The spots are subdivided into three classes: mRNA (yellow), protein (blue) and phosphorylated protein (brown). Scale bar, 20 μm.

## ARTseq-FISH profiles the expression of mRNAs, proteins, and phosphoproteins in individual mESCs

We utilized ARTseq-FISH to explore how the expression of mRNAs and (phospho-)proteins is affected during the early stages of differentiation on micropatterns. LIF/serum cultured mESCs are heterogenous (Supplementary Fig. 22), due to their pluripotent nature, enabling them to make different cell fate decisions upon LIF withdrawal[52-54]. Therefore, we plated mESCs directly on a 750 μm diameter micropatterns and withdrew LIF from the serum medium for up to 48 hours to allow for cells to exit pluripotency. Notably, cells cultured under this condition are not the same as the cells primed to epiblast-like cells (EpiLCs)[12]. After fixation, we performed ARTseq-FISH on the micropatterned mESCs. Using our analysis pipeline, we visualized the locations of unique mRNA and (phospho-)protein targets (Fig. 3a) and quantified the abundance of 67 targets in individual cells (Fig. 3a-d). When discussing the protein targets detected using the respective antibodies (Supplementary Table 2), it is important to note that these measurements likely represent the total amount of protein detected, which includes both the unmodified and modified forms. Conversely, phosphorylated proteins were detected with specific antibodies (*protein* or phos-*protein* in figures).

Among protein targets, lineage-associated markers i.e., N-Cadherin, FOXA2, GATA6, showed higher abundance than pluripotency-related markers, i.e., NANOG, OCT4, and SOX2. Although we cannot rule out that these differences are due to dissimilarities in antibody efficiency, this is expected in mESCs that are exiting pluripotency (Fig. 3b)[10, 12, 15]. Notably, while RB shows low spot counts at a protein level, its mRNA and Phos-RB display higher spot counts (Fig. 3b-d), indicating that the difference in spot count between total RB and Phos-RB is caused by differences in antibody specificity. As mentioned previously, similar to other omics methods that rely on antibodies[33, 34], ARTseq-FISH is dependent on the binding efficiency of primary antibodies (for proteins) and PLPs (for mRNAs). Therefore, we cannot compare absolute counts between species. However, strong correlation of target counts between four different biological replicates (Fig. 3f) demonstrate the high reproducibility of ARTseq-FISH (Supplementary Fig. 21 and SN2 Fig. 20).

In addition, after filtering out the low abundant RNA targets in single-cell RNA sequencing (scRNAseq) data, ARTseq-FISH and scRNAseq mRNA quantification showed a high correlation ($r^2=0.73$) (Fig. 3g). Finally, to ensure true signals were detected with the image analysis software since no fixed intensity threshold is implemented, we tested the software on three samples: i) samples that did not include any readout probes, i.e., negative control (Fig. 3e, top panel); ii) samples that included readout probes, i.e., true signal; and iii) an *in silico* merged image which combined samples i and ii. For the merged image the number of false positives decreased six-fold for all channels (Fig. 3e, bottom panel). This demonstrates that in the presence of true signal the false positive rate for the noisiest channel is <1%. Together these data indicate that ARTseq-FISH is a reproducible technique that should be implemented when comparing relative spatial or temporal changes in expression levels of mRNA and proteins.

**Fig. 3: ARTseq-FISH profiles the distribution of mRNAs and proteins in individual mESCs on micropatterns. a,** Reconstructed images show the detected spots in Figure 2b assigned to individual cells and the identity of these targets at a specific location. Scale bar, 20 μm, 5 μm, and 0.5 μm. **b-d,** Quantification of the single cell count abundance of proteins (**b**), mRNAs (**c**), and phosphorylated proteins (**d**). 48 cells were analysed per target. **e,** Quantification of spots per cell for negative controls in three different channels (the three fluorophores are AT: Atto488, Tm: TAMRA, and CY5). Negative controls were performed without PLPs and 579 cells were analysed. When adding true signal to negative control images *in silico*, the detection of spots in the negative control decreases >6-fold for the green channel (bottom). **f,** The scatter plot shows the average correlation between the total count of individual targets detected within the image for 4 biological replicates at 48 hours differentiation of mESCs. **g,** The correlation of RNA targets per cell between ARTseq-FISH and scRNA-seq performed on serum/LIF cultured mESCs. scRNAseq data was analysed by Seurat[55]. 455, 503, 470, 482, 454, 466, 488, 408, 408, 408, 180, 241, 200, 200 and 233 non-micropatterned cells were analysed for *Smad2, Tcf3, E-cadherin, Cdk4, CyclinD, Smad1, Klf4, Oct4, Nanog, Sox2, p53, Beta-catenin, Stat3, CyclinE, Yap* respectively.

**ARTseq-FISH reveals gene expression gradients across micropatterned mESCs after 48 hours of LIF withdrawal**

ARTseq-FISH provides single cell quantification (Fig. 1-3) of each molecular species while retaining spatial information. As confirmed by immunofluorescence, ARTseq-FISH data show three patterns of molecular distribution across mESCs on micropatterns: i) increased expression in cells located at the centre (i.e., p53); ii) evenly distributed throughout (i.e., phosphorylated RNA Pol II); and iii) increased expressed in cells at the periphery (i.e., BRACHYURY) (Fig. 4a-f, Supplementary Fig. 23). Consistently, targets are classified into these three groups (Fig. 4e and Supplementary Fig. 24, purple, white, and orange respectively). Overall, lineage, pluripotency as well as cell cycle associated proteins are more highly expressed in cells localised towards the edge of the micropattern. Specifically, BRACHYURY and GATA6 are mostly enriched in cells at the edge of the micropattern, as are the NANOG and SOX2 (Fig. 4e and Supplementary Fig. 25a). Yet, BRACHYURY and GATA6 are not necessarily highly expressed in the same cells as NANOG and SOX2 (Supplementary Fig. 25e). Surprisingly, some mRNAs (i.e., *Stat3* and *Cdk2*) appear localised to the centre although their (phospho-)protein counterparts are localised to the edge. This could either be because cells at the edge are translationally more active and thereby require less mRNA to produce a higher amount of protein. Alternatively, cells could be in the process of up or down regulating their mRNA, and differences in mRNA and protein half-life could lead to a lack of correlation between the two species[56]. Lastly, when calculating the mean fold change in expression level in cells located in the centre versus edge, cells >125 μm from the edge were defined as being in the centre of the micropattern (Supplementary Fig. 24), which is possibly an oversimplification. For example, *Smad2* and *Rb* mRNA and protein are enriched in cells at the edge while our analysis suggests that the phosphorylated proteins are more homogeneously expressed across the micropattern (Fig. 4e). However, when quantifying the average expression profile of spots per cell at seven positions across the micropattern, a reduction in Phos-RB and Phos-SMAD2 protein at the centre (~375μm from the edge) of the micropattern becomes evident (Supplementary Fig. 25b-d). Consequently, the localisation analysis likely masks more subtle changes in gene expression between cells at the edge and centre of the micropattern.

To more closely examine the relationship between target expression levels and cellular location, we quantified the expression levels of some targets in individual cells and plotted these against cellular position on the micropattern. This analysis revealed that targets that are more highly expressed in cells at the edge (i.e., BRACHYURY), display more of an exponential trend (Fig. 4g), rather than linear. Conversely, some targets that are classified as localised towards the centre (i.e., Phos-RB) or the edge (i.e., *Smad1* mRNA) appear to be most highly expressed in cells located ~150 μm from the edge, rather than 375 μm from the edge, the latter being the true centre. Interestingly, the single-cell expression levels of individual targets are highly heterogeneous (Fig. 4g). While this could be a technical consequence of ARTseq-FISH, the heterogeneity of p53, Phos-PolII, and BRACHYURY is also visible with regular immunofluorescence (Fig. 4b), and mRNA levels have previously been reported to become more heterogeneous as cells exit pluripotency[57]. Furthermore, we have shown that the mean gene expression values are reproducible across replicates when quantified by ARTseq-FISH (Fig 3f). Therefore, this heterogeneity is likely biological and indicates that although cellular location on the micropattern does generate trends in gene expression profiles, both mRNA and protein synthesis is still heterogenous after 48 hours of LIF withdrawal.

269
**Fig. 4: ARTseq-FISH reveals gradients of mRNA and protein expression levels within micropattern-**
**differentiated mESCs.**
**a,** Schematic illustration of three representative spatial distributions of molecules on micropatterns. Created with
BioRender.com. **b,** Immunofluorescence results of p53, Phos-Pol II (Phosphorylated RNA Pol II), and
BRACHYURY within a micropattern-differentiated mESCs. Scale bar, 100 μm. **c,** Average expression profile of
spots per cell at different positions on the micropattern (edge to centre) for p53, Phos-Pol II, and BRACHYURY
proteins from 3 biological replicates (150, 125, and 158 cells and 1 micropattern were per replicate). **d,** Pearson
correlation between relative mean fluorescence in **(b)** and relative spots per cell with respect to their position on
the micropattern **(c)**. **e,** Mean fold change of the expression level of individual targets between the edge and centre

(purple and orange respectively) of the micropattern across 3 biological replicates (125-158 cells and 1
micropattern analysed per replicate). *: phosphorylated protein. Error bars represents SEM of 4 biological
replicates. **f,** UMAP of cells after 48 hours of LIF withdrawal, with the colour representing if cells are localised
towards the edge (orange) or centre (purple) of the micropattern (left). Same UMAP with single cell expression
level of p53, Phos-Pol II, and BRACHYURY proteins indicated in blue. 158 cells and 1 micropattern were
analysed. **g,** The spots per cell (y-axis) with respect to the position of the cell on the micropattern (x-axis) of
selected targets. The edge of the micropattern serves as the reference value for the relative position of the cells.
Cells are found at the edge at 0 μm and near the centre at 375 μm. 158 cells were analysed per target.

## Position-dependent mRNA and protein profiles emerge over time

To investigate how mRNA and protein abundance changes over time, we examined the spatial
distribution of molecules in micropatterned cells at different time points (0 hours, 12 hours, 24
hours, and 48 hours) following LIF withdrawal (Fig. 5a). To better understand the changes, we
grouped targets into different categories (Supplementary Table 5) and quantified the relative
changes in target abundance across the micropattern in 2D (Fig. 5b) over the 48-hour LIF
withdrawal period. At the initial time point (0 hours), both RNA and protein levels across the
different categories showed a relatively homogeneous distribution. Interestingly, we noticed a
specific region, approximately 150 μm from the edge of the micropattern, where mRNAs
related to cell cycle class 2 (genes that promote cell cycle progression) were elevated (Fig. 5c,
cell cycle class 2). Moreover, this region exhibited slightly increased levels of mRNAs and
proteins related to (LIF) signalling at 0 hours (Fig. 5c, signalling). While we considered that
this might be an artefact of initial uneven cell seeding, cells were seeded on the micropattern
24 hours before LIF withdrawal (i.e., at T = -24 hours). Therefore, these subtle initial
expression differences could be indicative of first changes in position-based expression
profiles. This is reinforced by two distinct areas on the micropattern emerging where both
mRNA and protein levels tended to increase over time: these were found at position 2 and 6 of
the micropattern, located approximately 150 μm from either side of the edge (Fig. 5b).

To further explore whether proteins are expressed differently based on cellular location on the
micropattern, we quantified gene expression profiles for individual cells situated either at the
true edge or at the true centre of the micropattern (Fig. 5b). Once again, we categorized targets
based on their functions and plotted the relative single-cell changes over the 0, 12, 24, and 48
hours of LIF withdrawal. At 0 hours, cells at the micropattern's edge contain higher levels of
proteins associated with cell cycle 1 (lengthening G1) than at the centre (Fig. 5d, 0h cell cycle
1 top compared to bottom). Subsequently, at the edge after 12 hours we observed a decrease in
proteins associated with cell cycle 1 (lengthening G1) and an increase in proteins related to cell
cycle 2 (cell cycle progression) (Fig. 5d). These findings suggested that within the first ~12
hours of LIF withdrawal, cells at the edge alter the expression of genes associated with cell
cycle progression, potentially leading to higher proliferation rates during this time. In contrast,
at the micropattern's centre, the levels of cell cycle 1 (lengthening G1) increases already from
0-12 hours, while cell cycle 2 genes (cell cycle progression) initially remain constant (Fig. 5d,
bottom). These results indicated that, the G1 phase of cells located at the centre might start
increasing 0-12 hours after LIF withdrawal.

When analysing genes associated with pluripotency and particular lineages, we only observed
slightly different gene expression profiles at the centre compared to the edge of the
micropattern. Proteins that exhibit the most significant differences in expression profiles are
SOX17, BRACHYURY, (Fig. 5d, right) and NANOG (Fig 5e, left). In particular, SOX17
levels increase mostly between 0-12 hours at the edge of the micropattern, while at the centre
there is a more gradual increase over the course of 48 hours of LIF withdrawal. BRACHYURY
levels increase gradually at the edge of the micropattern from 12-48 hours of LIF withdrawal

and remain relatively constant at the centre. Lastly, NANOG levels increase from 0-12 hours
and decrease again from 12-24 hours of LIF withdrawal at the edge of the micropattern. In
contrast, at the centre of the micropattern NANOG levels increase from 12-24 hours of LIF
withdrawal.
334
To validate these findings, we performed time-lapse microscopy on micropatterned NANOG-
GFP mESCs[58, 59] for 46.5 hours of LIF withdrawal (Fig 5e, right). When quantifying the mean
NANOG-GFP intensity for different positions at the edge and the centre of the micropattern,
we observed distinct NANOG-GFP expression changes. Specifically, cells at the edge of the
micropattern show an initial increase in NANOG-GFP expression (lasting ~12-15 hours)
followed by a gradual decrease in NANOG-GFP expression. Conversely, the centre of the
micropattern displays a decrease in NANOG-GFP expression from 15-20 hours post LIF
withdrawal, followed by a continuous increase in NANOG-GFP expression (Fig 5e, right).
While the overall trend at the edge is consistent with ARTseq-FISH NANOG data (Fig 5e, left,
orange), the initial decrease observed in the time-lapse experiment for cells at the centre of the
micropattern is not visible by ARTseq-FISH (Fig. 5e, left, purple). This could be because
ARTseq-FISH reveals expression levels of cells that are fixed after 0, 12, 24, and 48 hours post
LIF withdrawal and do not represent a true time-course, which might conceal rapid changes in
expression levels. Furthermore, the time-lapse experiment was not analysed at a single-cell
level. Alternatively, it is possible that the differences between the time-lapse experiment and
ARTseq-FISH are cell-type specific. Therefore, we sought to confirm the expression of another
pluripotency marker (OCT4) by immunofluorescence at 0 and 48 hours of LIF withdrawal,
which shows similar behaviour as ARTseq-FISH (Supplementary Fig. 26).
353
Interestingly, at the single-cell level, we found a negative correlation between pluripotency-
associated proteins (i.e., OCT4 and SOX2) and lineage-associated markers (i.e., FOXA2,
GATA6 and SOX17) (Fig. 5f) in cells at the edge of the micropattern. This indicates that cells
at the edge are not uniform in their protein expression levels. When we project the data on a
2D UMAP, we observe the most prominent separation of cellular expression profiles occurring
between 0-12 hours post LIF withdrawal (Fig. 5g). Furthermore, while there is some separation
in cells located at the edge and centre, this separation is subtle, and predominantly visible at 48
hours post LIF withdrawal (Fig. 5g).
362

**Fig. 5: Spatial-temporal quantification of mRNAs and proteins after LIF withdrawal from mESCs on micropatterns. a,** Schematic of the experiment where ARTseq-FISH was performed at 0, 12, 24 and 48 hours after LIF withdrawal. Created with BioRender.com. **b,** Schematic of the definition of centre and edge on the micropattern. Created with BioRender.com. **c,** Heatmap showing the abundance of different classes of targets (see Table S5 for full list of targets) across 1 micropattern at 0, 12, 24 and 48 hours after LIF withdrawal. The expression level of targets is normalised to the maximum abundance across all four time points. **d,** Relative single cell expression levels of the targets shown in **c**, at the edge (top) and the centre (bottom) of the micropattern at 0, 12, 24 and 48 hours after LIF withdrawal. 0–24-hour data includes 2 biological replicates; 48 hours includes 3 biological replicates (for the edge 393 cells and the centre 238 cells were analysed in total). For each replicate and

time point 1 micropattern was analysed. Shaded regions represent 95% confidence intervals. **e,** Single cell
NANOG protein expression at the edge and centre of micropattern at 0, 12, 24 and 48 hours after LIF withdrawal
detected by ARTseq-FISH (left). 0–24-hour data includes 2 biological replicates; 48 hours includes 3 biological
replicates (edge and centre include 393 and 238 cells in total respectively). For each replicate and time point 1
micropattern was analysed. Quantification of live cell time-lapse of NANOG-GFP positive mESCs for 5 different
positions at the edge (cells) and the centre (cells) of 1 micropattern over 46.5 hours of LIF withdrawal (right).
Shaded regions represent 95% confidence intervals. **f,** Pearson correlation and hierarchical clustering of
pluripotency and lineage related markers at the edge of the micropattern (3 biological replicates, 1 micropattern
each, 111 cells total) at 48 hours after LIF withdrawal. **g (i)** UMAP clustering of the individual cells within a
single micropattern experiment at different time points (0, 12, 24 and 48 hours) after LIF withdrawal. **(ii)** UMAP
clustering of single cells at different positions (centre, edge or between centre and edge) within the micropattern
(purple, orange, and white respectively).  1 biological replicate and 1 micropattern (0, 12, 24 and 48 hours include
269, 139, 141, 158 cells respectively).

## Cellular location on the micropattern impacts proliferation

In light of the gene expression profiles (Fig. 5c) indicating differences in protein signals
between the micropattern's edge and centre, we proceeded to investigate the translational
activity of cells across the micropattern. To do this, we exposed micropatterned mESCs to the
puromycin analogue O-propargyl-puromycin (OPP), which is incorporated into newly
synthesized proteins[60]. Through click chemistry, we fluorescently labelled the nascently
translated polypeptides (Fig. 6a). Our data revealed a gradual decrease in protein synthesis
across the entire micropattern over time (Fig. 6b). Notably, the signal in the micropattern's
centre decreased more readily, particularly after 24-48 hours (Fig. 6b-c, light and dark green).
These findings are in line with ART-seqFISH data, showing more prominent protein detection
at the micropattern's edge after 48 hours of LIF withdrawal (Fig. 6d). These data suggest that
spatial organization plays a role in the dynamics of translational activity in micropatterned
mESCs.

To gain insights into why translation might be reduced in the centre of the micropattern, we
next tried to determine if cells proliferate differently depending on their position on the
micropattern. To this end, we first quantified cell density across the micropattern as a function
of DAPI intensity. Interestingly, on our 750 µm diameter micropatterns, cell density shows two
radially symmetric peaks over time, approximately 100 µm from the edge of the micropattern
(Fig. 6e-f), indicating that the colonies are not flat but somewhat 'donut' shaped (Fig. 6g). To
determine if these differences in cell density were caused by increased proliferation, we
performed EdU staining after 0, 12, 24 and 48 hours of LIF withdrawal and quantified cell
proliferation from the incorporation of EdU into newly synthesized DNA[61] (Fig. 6h). The
results indicate that there are fewer cells in the S phase at the centre of the micropattern than
towards the edge and therefore likely display reduced proliferations already after 12 hours of
LIF withdrawal (Fig. 6i). Interestingly, at the very edge of the micropattern there still appear
to be cells incorporating EdU after 48 hours of LIF withdrawal. To confirm that actively cycling
cells are more abundant at the edge of the micropattern after LIF withdrawal, we performed
Ki-67 staining[62] (Fig. 6j and Supplementary Fig. 27), an indirect marker for proliferating cells.
The data show that at 0 hours after LIF withdrawal Ki-67 staining is mostly homogenous across
the micropattern, indicating comparable cell cycling at the centre and edge of the micropattern
(Fig. 6i). Notably, at 0 hours there is reduced Ki-67 signal at the very centre of the micropattern
(micropattern position 1 and 8), consistent with quantified EdU signal (Fig. 6i, light blue).
From 0 to 12 hours, there appears a slight increase in Ki-67 intensity (i.e., cell cycling) at the
centre of the micropattern, which subsequently drops from 24 to 48 hours post LIF withdrawal,
demonstrating a decrease in cellular proliferation or a state of cell cycle arrest (Fig. 6j and
Supplementary Fig. 27). Caspase-3 and 7-AAD staining as well as nuclear morphology

analysis indicate that this lack of proliferation could be connected to increased apoptosis at
centre of the micropattern[63] (Supplementary Fig. 28). Conversely, at the edge of the
micropattern cells show more consistent Ki-67 intensities over time (Fig. 6j), although at 48
hours the cells at the very edge (micropattern position 1 and 8), do not show increased cycling
as would be expected from the EdU staining (Fig. 6j compared to Fig. 6i), this could be because
Ki-67 is less sensitive, in particular for cells with a short G1 phase[64]. Nevertheless, these
combined assays indicate that cells positioned more towards the edge of the micropattern
proliferate more readily than cells at the centre.
433

434
**Fig. 6: mESCs display position-dependent differences in protein synthesis and proliferation rates.**
**a,** Representative images showing the protein synthesis of the micropatterned mESCs at 0, 12, 24 and 48 hours
after LIF withdrawal. Images from different time points are set as the same maximum and minimum gray value.
Scale bar, 120 μm. **b,** Quantification of relative fluorescent intensity of the protein synthesis in mESCs across 2-
10 micropatterns at 0, 12, 24 and 48 hours after LIF withdrawal. **c,** Representative images showing the protein
synthesis of the micropatterned mESCs at 48 hours after LIF withdrawal (left). Scale bar, 120 μm. Quantification
of relative fluorescent intensity of the protein synthesis in mESCs across 10 micropatterns at 48 hours after LIF
withdrawal (right). **d,** Pie chart showing the percentage of mRNAs, proteins, and phosphoproteins of targets that
are more abundant at the centre (purple) and targets that are more abundant at the edge (orange) of the

micropattern. **e and f,** Heatmap of DAPI intensity across the micropattern and normalized DAPI intensity across the micropattern at 0, 12, 24 and 48 hours after LIF withdrawal. **g,** Schematic shows the "donut" shape of the micropatterned mESCs. **h,** Representative images showing EdU staining of the micropatterned mESCs at 0, 12, 24 and 48 hours after LIF withdrawal. Images from different time points are set as the same maximum and minimum grey value. Insets: image of EdU staining at 48 hours after LIF withdrawal with adjusted contrast. Scale bar, 120 μm. **i,** Quantification of relative fluorescent intensity of EdU staining in mESCs across 2-10 micropatterns at 0, 12, 24 and 48 hours after LIF withdrawal. **j,** Heatmap of Ki-67 expression level at z-projection across 1 micropattern at 0, 12, 24 and 48 hours after LIF withdrawal.

## Cellular location on the micropattern impacts cell cycling dynamics

Given the distinct expression profiles of cell cycle-related genes observed between cells at the micropattern's edge and centre using ARTseq-FISH (Fig. 5d), we aimed to quantitatively analyse cell cycle progression on a single-cell basis. To this end, we conducted time-lapse microscopy on micropatterned Fucci mESCs[65, 66], which enable the visualisation of cell cycle dynamics in live cells (Fig. 7a). Single-cell tracking and measurement of Azami Green (Az1) and Kusabira Orange-2 (KO-2) fluorescence over time delineates the individual stages of the cell cycle (Fig. 7b-c). We performed single-cell tracking of 35 cells across the micropattern (20 at the centre and 15 at the edge) over a 45-hour period following of LIF withdrawal. When assessing the length of the G1 phase at different time intervals post LIF withdrawal, intriguing differences emerge between cells at the micropattern's edge and those at the centre. Specifically, cells at the micropattern's centre that enter the G1 phase within the initial 5 hours of LIF withdrawal consistently exhibit a relatively short G1 phase of approximately 3 hours (Fig. 7d, top and Supplementary Fig. 31a, left). Conversely, cells that enter the G1 phase after this initial 5-hour period show a notably extended G1 phase (Fig. 7e). Markedly, cells at the micropattern's edge display a G1 phase duration similar to cells at the centre (Fig. 7d, bottom and Supplementary Fig. 31a, middle). However, a subset (about 25%) of cells at the edge already exhibit a prolonged G1 phase within the first 5 hours of LIF withdrawal (Supplementary Fig. 31a, right). Similarly, after the initial 5-hour period, just over half (approximately 60%) of cells closer to the edge continue to cycle, while the remaining portion (about 40%) display an extended G1 phase (Fig. 7f). These single-cell analyses show that the G1 phase of cells at the centre gradually elongates, while >half of the cells at the edge continue cycling until ~30 hours post LIF withdrawal. These data suggested that the cell cycle pace is more heterogeneous at the edge of the micropattern than at the centre.

We next performed a more comprehensive population-based analysis spanning the entire micropattern over the course of the 45-hour LIF withdrawal. By comparing the proportion of cells in G1 versus S/G2/M phases for cells at the micropattern's edge and centre over time, we again observe a subtle yet discernible difference (Fig. 7g). Specifically, in the centre there was a gradual decline in the fraction of cells in S/G2/M phase, beginning approximately 6 hours post LIF withdrawal (Fig. 7g, left). Instead, cells at the edge of the micropattern show a relatively unchanged distributions across the cell cycle stages over time, with the proportion of cells in the S/G2/M phase only declining after ~40 hours of LIF withdrawal (Fig. 7g. right). Since these data align with the single-cell analysis, the latter is likely representative of the full population. Furthermore, these findings are consistent with the Ki-67 staining results (Fig. 6j and Supplementary Fig. 27) and the cell cycle stages quantified from DAPI intensity differences[67-69] (Supplementary Fig. 29-30). Although differences in timing of these changes might arise because the Fucci mESCs are a separate cell line[65, 66], the fraction of cells in the late S/G2/M phase (Fucci mESCs) follow a similar trend to the EdU staining performed on mESCs (Fig. 7h). Lastly, single cell tracking of both NANOG-GFP and Fucci mESCs suggests that cell migration is minimal (Supplementary Fig. 31b and c, respectively). While the potential for rapid cell migration that elude cell tracking exists, the prevailing data indicates that cells

remain either at the centre or the edge of the micropattern (Supplementary Fig. 31c). Collectively, the data indicate that cells closer to the micropattern's edge exhibit increased proliferation compared to those at the centre, evident 5-24 hours after LIF withdrawal.

**Fig. 7: Live cell imaging reveals differences in cell-cycle progression.**
**a,** Schematic illustration of the Fucci mESCs demonstrating different colours at different cell cycle stages. Created with BioRender.com. **b-c,** Kusabira Orange-2 (KO-2) (G1) and Azami Green (Az1) (S/G2/M) intensity of a tracked cell in the centre (**b**) and edge (**c**) of the micropattern. **d,** Average KO-2 intensity throughout the G1 phase for cells at the centre (top) and edge (bottom) of the micropattern that enter G1 within 5 hours of LIF withdrawal. Average is of 12 cells for the centre and 6 cells for the edge. Error bars represent standard deviation. **e-f,** Single cell KO-2 (G1) traces of cells that enter G1 after 5 hours of LIF withdrawal at the centre (**e**) and edge (**f**) of the micropattern. 7 (centre) and 9 (edge) cells were analysed respectively. **g,** Quantification of the fraction of G1, early S, and late S/G2/M mESCs at the centre (left) and the edge (right) of the micropattern over time after 45 hours of LIF withdrawal. 1961 cells at the edge and 2190 cells at the centre were analysed in total. **h,** Quantification of mean EdU intensity of mESCs for 10 different positions at the edge and the centre of 1 micropattern at 0 hours, 12 hours, 24 hours, and 48 hours after LIF withdrawal. Shaded regions represent 95% confidence intervals.

## Discussion

Thus far, there have been two major areas of focus when studying micropatterned cells: i) implementation of immunohistochemistry (IHC) to determine spatial-temporal protein abundance in low throughput[10-15]; or ii) revealing the emergence of specific cell types by single-cell transcriptomic sequencing in high throughput[15]. Here we developed and exploit a

technique that provides quantitative spatial information on different functional proteins and
their respective mRNAs at same time.
ARTseq-FISH, enables simultaneous measurement and quantification of relative mRNA and
(phospho-)protein abundance in individual cells. The combined protocol of mRNA and protein
detection provides the same resolution of different molecular species and is scalable (Fig. 1).
Additionally, the deep learning-based image analysis pipeline in ARTseq-FISH (Fig. 2)
identifies the cellular nucleus and FISH signals, which allowed detection of 67 mRNAs,
proteins, and phosphoproteins simultaneously (Fig. 3). We applied ARTseq-FISH to identify
position-dependent gene expression profiles on micropatterned mESCs (Fig. 4). The analysis
revealed the expression levels of individual cells and their dependence on the respective
location on the micropattern during 48 hours of LIF withdrawal. Intriguingly, cells located at
the edge and centre of the micropattern obtain different gene expression profiles over time (Fig.
5) while maintaining a high degree of cell-to-cell variability in expression levels. We
demonstrate that depending on the micropattern location, cells display differences in protein
synthesis, cell cycling and proliferation (Fig. 6-7). Taken together, our findings indicted that
while gene expression is still highly heterogeneous differences in mRNA and protein levels
emerge as early as 12 hours after LIF withdrawal.

While other impressive spatial-omics methods that simultaneously measure mRNA and
proteins exist (DSP[18] and SMI[36], introduced by NanoString; $\pi$-FISH[70]), we focused on
developing a protocol that allows for a combined amplification step yielding comparable
resolution for mRNA and protein detection. This combined detection and analysis for mRNA
and protein offers the opportunity to circumvent the challenges in integration of different
datasets[71-73]. Since the sensitivity of ARTseq-FISH relies on the antibodies (for proteins) and
PLPs (for mRNAs), similar limitations apply compared to existing methods[33, 34, 41, 43]. Although
there is a risk that the affinity of the antibodies may be influenced by the conjugation method,
it is a widely used procedure in current proteomic technologies[33-35]. While we have shown that
the detection efficiency of the PLP-based method depends on the binding efficiency of the
chosen padlock probe sequence, we only used one PLP to target each mRNA in this study. This
is feasible, when comparing relative RNA abundance, either between individual cells (similar
to scRNA-seq) or as we do here between different locations or different time points during the
initial stages of mESCs differentiation.

In future, ARTseq-FISH has the potentials to detect thousands of targets simultaneously by
introducing more unique readout probe sequences and maximizing the hybridization rounds.
For instance, to detect 2048 targets, we will need 40 different readout probe sequences and at
least 10 rounds of hybridization. However, in theory sequential FISH methods are able to
perform 81 hybridization rounds, which would allow the detection of more than 10,000 targets
with ARTseq-FISH.  Yet, to achieve this, it will be critical to assign different hybridization
rounds to high abundance targets to avoid optical crowding. A future application of ARTseq-
FISH lies specifically in capturing the spatial information of different molecular species (i.e.,
mRNAs, proteins, phosphoproteins) and quantifying their relative abundance in particular
biological processes. One of the advantages of using padlock probes is the power of signal
amplification by RCA. Therefore, ARTseq-FISH has the potential to detect DNA (i.e., single
nucleotide polymorphisms (SNPs))[74], miRNAs[75, 76], viral RNA[77], and proximity of proteins[37,
78] or even proximity of RNA and protein. In addition, due to the signal amplification of RCA,
the results of ARTseq-FISH can also be read out by flow cytometry[79, 80], which makes this and
future techniques more broadly accessible.

## Methods
### Cell culture
mESC-E14 (mESCs) in this study were obtained from Hendrik Marks's group at Radboud University[81]. The culture plates were coated with 0.1% gelatin (Sigma Aldrich, 48723-500G). Undifferentiated mESCs were cultured and maintained in the medium consisting of high glucose-Dulbecco's Modified Eagles Medium (Gibco™, 41965039) supplemented with 15% fetal bovine serum (Gibco™, A3840002), 2 mM L-glutamine (Gibco™, A2916801), 1 mM sodium pyruvate (Gibco™, 11360039), 1% penicillin–streptomycin (Gibco™, 15140122), 0.1 mM beta-mercaptoethanol (Gibco™ 31350010) and 500 U/mL recombinant leukemia inhibitory factor (LIF; Millipore, ESG1107) at 37 °C in 5% $CO_2$. Cells were passaged every 2-3 days using 0.05% Trypsin/EDTA (Gibco™, 25300062). Cell density was 5x $10^4$ per well for the experiments in the μ-Slide (ibidi, 81501) and 1x $10^4/cm^2$ for the experiments in the μ-Slide 3D perfusion (ibidi, 80376). For differentiation assays, cells were cultured on micropatterns in LIF medium for 1 day and then changed to a medium without LIF (-LIF) for up to 2 days.

### Micropatterned Substrate Fabrication
Micropatterned islands with different diameters (750 μm) were produced on a silicon mask using standard photolithography [82]. Distances between the microfeatures were 100 μm and the height of the microfeatures was between 50 μm and 100 μm. The PDMS and the curing agent (SYLGARD 184, 1.1KG) was mixed with a ratio of 10:1 in a plastic beaker. The PDMS mixture was degassed to remove the air bubbles under vacuum. Then, the degassed PDMS mixture was poured onto the master in a petri dish. A gel was formed in a 60°C oven over night. Finally, the gel PDMS stamps was peeled off the master.

The μ-Slide 3D perfusion microchambers were plasma oxidized to improve protein printing efficiency and cell attachment on the glass bottoms [83]. The surface of the PDMS stamps was placed with a drop of 50 μg/mL fibronectin in PBS and incubated for 1 hour at room temperature. The PDMS stamps were washed twice with PBS and dried thoroughly. The glass bottoms of μ-Slide 3D perfusion were then coated with the fibronectin from the PDMS stamps, followed by the graft of the PLL (20)-g [3.5]- PEG (2) (0.1 mg/mL, SuSoS AG) in 10 mM HEPES (Gibco™, 15630080) for 30 minutes at room temperature. The PLL-g-PEG prevents cell attachment in the nonprinted areas [84].

### Immunostaining
Cells were fixed with 4% paraformaldehyde (PFA) (Sigma Aldrich, 1040021000-1L) for 10 minutes at room temperature and then permeabilized with 0.5% Triton in 1x PBS at room temperature for 10 minutes. Cells were washed with 1x PBS three times after fixation and permeabilization. After permeabilization, cells were incubated in blocking buffer (5% BSA in 1x PBS containing 0.5% Triton) for 1 hour at room temperature. The primary antibodies were diluted in blocking buffer and incubated at room temperature for 2-4 hours. Cells were then washed with 1x PBS three times and incubated in secondary antibody solution for 2 hours at room temperature. The secondary antibodies were diluted 1:1000 in 1x PBS. DAPI (10 μg/mL) (Thermo Scientific™ 62247) staining was performed together with secondary antibodies.

For Ki-67 (Cell Signaling, 9192), Caspase-3 (Cell Signaling, 94530), OCT4 staining, the primary antibodies were diluted as recommended ratio in blocking buffer. The cells were then incubated overnight at 4°C in the primary antibody solution. The other steps of the protocol remained the same as above.

### smFISH

Probes of *Nanog* were purchased from Stellaris (LGC Biosearch Technologies, Novato, CA) (http://www.singlemoleculefish.com/). Each set contains 28 probes and each probe is 20 nt long (Table S1). Probes are conjugated with TAMRA. Cells were fixed with 4% PFA (Sigma Aldrich, 1040021000-1L) for 10 minutes at room temperature and then permeabilized with pre-chilled 70% EtOH at 4°C for a minimum of 1 hour. Cells were washed three times with 1x PBS. Probes were diluted to a final concentration of 50 nM in 1 g/mL dextran sulfate (Sigma, catalog # 42867), 2x SSC and 10% formamide (Invitrogen, AM9342). The hybridization of the probes was performed overnight at 37°C. The next day, samples were washed with 2x SSC buffer at 37°C for 30 minutes and then incubated with DAPI solution (diluted in 2x SSC) at 37°C for 15-20 minutes. Cells were washed with 2x SSC and then switch to a photo-protective buffer containing 50% glycerol (Thermo Fisher Scientific, Waltham, MA), 75 μg/mL glucose oxidase (Sigma Aldrich, Darmstadt, Germany), 520 μg/mL catalase (Sigma Aldrich, Darmstadt, Germany), and 0.5 mg/mL Trolox (Sigma Aldrich, Darmstadt, Germany). Samples were imaged with a spinning disk confocal microscope (OLYMPUS PlanXApo 60x/1.40- N.A. oil-immersion objective, and Andor iXon camera). DAPI was excited by 405 nm, and TAMRA was excited by 540 nm lasers. The distance of Z-stacks was set to 0.5 μm. The exposure time and laser intensity were set to reach a maximum gray value of 16000 each image.

**Probe design of ARTseq-FISH**

The mRNA sequences were obtained from NCBI website (Mus musculus). Probes were designed according to the targeted regions by using online software (http://biosearchtech.com/stellaris-designer/), for individual probe sequences see Table S1. For targeting RNA directly, we selected 36 nt from the mRNA sequence (18 nt for 5' arm and 18 nt for 3'arm of PLP). Each PLP contains two arms (18 nt each) for binding to the targeting region, one unique 28 nt sequence identical to the bridge probe and one common RCA primer binding region. Besides the 28 nt sequences identical to paired PLPs, BrPs consist of four overhang sequences with 18 nt each.

Oligonucleotides used for antibody conjugation consist of a 10 nt linker (polyA) and a 36 nt sequences corresponding to the complementary PLP. The 36 nt sequences for antibody conjugation, 28 nt sequences of PLPs and BrPs, and 18nt readout probe sequences were identified using a previously published algorithm [24]. The readout probes are 18 nt in length and labeled with three different fluorophores (ATTO488, TAMRA, CY5). We used BLAST from NCBI to check that the 18 nt probe sequence is less than 13 nt (<70%) complementary to other Mus musculus transcripts. All the probes have GC content within the range 45%-65% and were purchased from Integrated DNA technologies (IDT).

| Oligonucleotides (46 nt) | 5'- Azide-AAAAAAAAAA- [36 nt unique sequence] -3' |
|---|---|
| Padlock probe (84 nt) | 5' -phosphorylated- [18 nt 5'-arm] - [28 nt unique sequence] - [20 nt RCA primer] - [18 nt 3'- arm] - 3' |
| Bridge probe (108 nt) | 5' - [readout 1] -AA- [readout 2] -AA- [28 nt unique sequence] -AA- [readout 3] -AA- [readout 4] - 3' |

**Antibody–DNA conjugation**

Antibodies (Abs) used in ARTseq-FISH were labelled with DNA oligonucleotides as previous described[39]. In brief, Abs were first buffer exchanged into 0.2 M NaHCO3 (pH 8.3) (Thermo Scientific Acros 217125000) by using Zeba™ Spin Desalting Columns, 40K MWCO (Thermo Scientific™ 87767). Abs were then reacted with 10x molar excess of dibenzocyclooctyne-S-S-N-hydroxysuccinimidyl (DBCO-S-S-NH) ester (Sigma Aldrich, 761532-1mg) for 2 hours at room temperature in the dark. Next, Abs were purified using Zeba™ Spin Desalting Columns,

40K MWCO to remove excess DBCO-s-s-NHS. 5' Azide modified DNA oligonucleotides
(IDT) were incubated with Abs (molar ratio of antibody to single-stranded DNA of 1:15)
overnight at 4°C in the dark. The conjugated Abs were buffer exchanged into PBS (Gibco™
20012019) containing 0.05% sodium azide (Sigma Aldrich, S8032-25G) and 0.1 mM EDTA
(Lonza, 51201), and excess oligos were removed by 100K Amicon centrifuge filters (Merck
UFC510096). The DNA-conjugated Abs were stored at 4°C. The list of antibodies and
corresponding DNA oligonucleotides is provided in Table S1 and S2. The dilution of
conjugated antibodies was optimized between ratios of 1:1000, 1:2000 and at 1:4000 and the
ratio of 1:4000 was used for ARTseq-FISH.
671
**ARTseq-FISH**
***Fixation and permeabilization*** Cells were fixed with 4% PFA (Sigma Aldrich, 1040021000-
1L) for 10 minutes at room temperature and then permeabilized with pre-chilled 100%
methanol at -20°C for 10 minutes. Cells were washed with 1x PBS three times after fixation
and permeabilization.
***Blocking and Primary antibody incubation*** After permeabilization, cells were incubated in
blocking buffer (5% BSA in PBS containing 0.5% Triton) for 1 hour at room temperature. The
DNA oligonucleotide labelled primary antibodies were diluted to 1:4000 in blocking buffer
and incubated at room temperature for 2-4 hours.
***RNA PLP hybridization and ligation*** Phosphorylated PLPs were hybridized to the mRNA
sequences directly at a final concentration of 100 nM/PLP. The reaction was performed
overnight at 37°C using 1x Ampligase ligase buffer (Lucigen, A1905B), 50 mM KCl
(Invitrogen, AM9640G), 20% Formamide (Invitrogen, AM9342), 0.2 µg/µL tRNA, 1U/ µL
RiboLock RNase Inhibitor (Thermo Scientific, EO0381) and 0.2 µg/µL of BSA in Nuclease-
free water (Invitrogen, AM9937). Next, PLP ligation was performed for 4 hours at room
temperature using 0.5 U/µl SplintR Ligase (NEB, M0375), 1x Ampligase ligase buffer, 10 µM
ATP (Invitrogen, 18330019) and 0.2 µg/µL of BSA in Nuclease-free water.
***Ab PLP hybridization and ligation*** The hybridization of PLPs to DNA oligonucleotides
conjugated to Abs was performed at a final concentration of 5 nM/PLP. The ligation of PLPs
was performed in the same reaction with T4 DNA ligase, 1x T4 DNA ligase buffer, 50 mM
KCl (Invitrogen, AM9640G), 20% Formamide (Invitrogen, AM9342), 0.2 µg/µL of BSA and
Nuclease-free water. This reaction was incubated at 37°C for 30 minutes followed by 45°C for
90 minutes.
***Rolling circle amplification (RCA)*** RCA was performed with 0.5 U/µL Phi29 polymerase
(NEB, M0269L), 1x Phi29 polymerase buffer, 5% Glycerol (Sigma, G5516), 0.25 mM each
dNTP (Thermo Scientific, R0192), 0.2 U/µL Exonuclease I (Thermo Scientific, EN0581), 0.2
µM RCA primer and 0.2 µg/µL of BSA in Nuclease-free water overnight at 30˚C.
***Bridge probe hybridization*** BrPs were hybridized at a final concentration of 100 nM per probe.
The reaction took place in the hybridization buffer which contains 2x SSC, 20% formamide,
0.05 g/mL dextran sulfate and nuclease-free water, for 1-2 hours at room temperature. An extra
fixation with 4% PFA was performed to stabilize the binding of bridge probes and rolling circle
amplification products.
***Readout probe hybridization*** RoPs were hybridized at a final concentration of 100 nM per
probe and in the same hybridization buffer as the BrPs. The reaction can be performed together
with DAPI (10 µg/mL) (Thermo Scientific™ 62247) staining for 1h at room temperature. Cells
were washed with 2x SSC and then switch to a photo-protective buffer containing 50% glycerol
(Thermo Fisher Scientific, Waltham, MA), 75 µg/mL glucose oxidase (Sigma Aldrich,
Darmstadt, Germany), 520 µg/mL catalase (Sigma Aldrich, Darmstadt, Germany), and 0.5
mg/mL Trolox (Sigma Aldrich, Darmstadt, Germany).

*__Image acquisition__* Images of sequential hybridization were acquired with a spinning disk confocal microscope (OLYMPUS PlanXApo 60x/1.40- N.A. oil-immersion objective, and Andor iXon camera). DAPI was excited by 405 nm, and ATTO488, TAMRA, Cy5 was excited by 475 nm, 540 nm and 638 nm lasers respectively. The distance of Z-stacks was set as 0.5 μm. The exposure time and laser intensity were set differently in different channels to obtain the same maximum intensity value of spots.

*__Stripping__* Samples were incubated in 60% formamide solution (60% formamide in 2x SSC) for 10 mins to remove the readout probes after image acquisition and then washed three times with 2x SSC.

*__Rehybridization__* A new set of RoPs were hybridized in the hybridization buffer for 1hour at room temperature in dark.

*__Image acquisition__* Same as described above. Repeat the cycle of image acquisition, stripping off readout probes and rehybridization.

**Single-cell RNA sequencing (scRNAseq) in mESCs**

mESC were cultured as previous described. Cells were grown for 30 hours in a 37 °C incubator with 5% $CO_2$. Cells were exposed to 0.007% EtOH for the last 6 hours. Cells were then washed once with PBS and detached by 0.05% trypsin-EDTA for 2 minutes at 37 °C. Detached cells were pelleted and resuspended in 1mL of freezing media consisting of 80% culture media, 10% of extra heat inactivated FBS and 10% of DMSO. Proper viability of the cells prior to freezing was quantified with propidium iodide/acridine orange staining in a LUNA-FL Dual Fluorescence Cell Counter. Frozen cells were delivered to the commercial company Single Cell Discoveries (https://www.scdiscoveries.com) in dry ice. Single cell barcoding was performed in a 10x microfluidic genomic chip in order to encapsulate individual cells in water droplets in oil, containing cell specific barcoded beads. Labelled RNA molecules were pooled and subjected to a poly-A specific reverse transcription. cDNA molecules where linearly amplified using an in vitro transcription reaction and the final sequencing library was obtained through a second step of reverse transcription. 3'-end sequencing was performed, followed by quality control and genomic mapping. Final read count per gene and per cell matrices where generated and used in posterior analysis steps.

**Flow cytometry analysis of cell cycle**

mESCs were seeded in 1.9 $cm^2$ wells (24-well format) with a density of 4.4 x $10^3$ cells/$cm^2$. The wells were previously coated with 0.1% gelatin. Cells were seeded in serum/LIF medium and cultured for 24 hours. Then, cells were fixed in pre-chilled 70% ethanol for 30 minutes at -20°C and washed by 1x PBS. Finally, cells were stained with 1 μg/mL propidium iodide (Sigma-Aldrich, P4864-10ML) containing 10 μg/mL RNaseA (VWR International B.V, 0675-250MG) for 15 minutes. After staining, cells were transferred to a flow cytometry tube and kept on ice until their immediate analysis in a BD FACS Calibur™ flow cytometer.

**Flow cytometry analysis of NANOG-GFP mESC cell line in serum/LIF conditions**

The NANOG-GFP mESC line was kindly provided by Leor Weinberger's lab (Gladstone Institutes)[58]. NANOG-GFP mESCs and non-labelled mESCs were seeded in 9.5 $cm^2$ wells (6-well format) with a density of 4x $10^4$ cells/$cm^2$. The wells were previously coated with 0.1% gelatin. Cells were seeded in serum/LIF medium. 24 hours post-seeding, cells were detached with 400 μL of 0.05% trypsin-EDTA for 2 minutes and resuspended in growing media. Cells were transferred to a flow cytometry tube and kept on ice until their immediate analysis in a BD FACS Calibur™ flow cytometer.

**Flow cytometry analysis of CD24 expression in mESC**

mESCs were seeded in 1.9 cm$^2$ wells (24-well format) with a density of 4.4x 10$^3$ cells/cm$^2$.
The wells were previously coated with 0.1% gelatin. Cells were seeded in serum/LIF medium
and cultured for 96 hours, with a medium renewal at 48 hours of culture. Cells were stained
with an anti-mouse CD24 antibody conjugated to alexa647 fluorophore (BioLegend, 101818)
or an alexa647 isotype control antibody (BioLegend, 400626) for 30 minutes in serum/LIF
media at 37˚C. After staining, cells were washed with 1x PBS and detached with accutase
(StemCell™ technologies, 07922) for 5 minutes at 37˚C, transferred to a flow cytometry tube
and kept on ice until their analysis in a BD FACSVerse™ flow cytometer.
769
**EdU proliferation assay**
mESCs were seeded onto the micropatterns, ensuring complete coverage, and cultured for 16
hours. Subsequently, the medium was changed to a LIF-free medium. To assess cell
proliferation at different time points, the culture medium was supplemented with 10 μM of
EdU two hours before fixation at each time point. The detection of EdU incorporation was
performed according to the protocol provided by the EdU proliferation kit (Abcam, ab219801,
iFluor 488). In brief, the cells were washed twice with Wash Buffer and then fixed with 4%
PFA for 10 minutes at room temperature, while avoiding exposure to light. After the fixation
step, the cells were washed twice and permeabilized with 1x Permeabilization Buffer for 20
minutes at room temperature. Subsequently, the cells were washed twice and incubated in the
dark for 30 minutes at room temperature in a Reaction mix containing TBS, 4 mM Copper
Sulfate, 1.2 μM iFluor 488 azide dye (500 μM in DMSO), and 1x EdU additive solution.
Following another round of washing, the cells were stained with Hoechst 33342 (5 μg/mL).
Finally, the cells were imaged using a spinning disk confocal microscope (OLYMPUS
UPlanFL N 10x/0.30 N.A. objective, and Andor iXon camera). The Hoechst 33342 and EdU
signals were excited at wavelengths of 405 nm and 475 nm, respectively.
786
**Protein synthesis assay**
mESCs were seeded onto the micropatterns, ensuring complete coverage, and cultured for 16
hours. Subsequently, the medium was changed to a LIF-free medium. To assess the protein
synthesis at different time points, the culture medium was supplemented with 20 μM of Click-
iT® OPP 30 minutes before fixation at each time point. The detection of protein synthesis was
performed according to the protocol provided by the Click-iT® Plus OPP Protein Synthesis
Assay Kits (Thermo Fisher, C10456). In brief, the cells were washed once with 1x PBS and
then fixed with 4% PFA for 10 minutes at room temperature. After the fixation step, the cells
were washed twice and permeabilized with 0.5% Triton-X-100 in PBS for 15 minutes at room
temperature. Subsequently, the cells were washed twice and incubated in the dark for 30
minutes at room temperature in a Click-iT® Plus OPP reaction cocktail containing 1x Click-
iT® OPP Reaction Buffer, Copper Protectant (Component D), Alexa Fluor® picolyl azide
(Component B) and 1x Click-iT® Reaction Buffer Additive. Following another round of
washing, the cells were stained with 1X HCS NuclearMask™ Blue Stain for 30 minutes at
room temperature, protected from light. Finally, the cells were imaged using a spinning disk
confocal microscope (OLYMPUS UPlanFL N 10x/0.30 N.A. objective, and Andor iXon
camera). The NuclearMask™ Blue Stain and protein synthesis signals were excited at
wavelengths of 405 nm and 638 nm, respectively.
805
**Apoptosis detection**
To detect apoptosis of mESCs, Cleaved Caspase-3 (Cell Signaling, 94530) and 7-AAD
(Thermier Fisher, 00-6993-50) were measured to assess the middle and late stage of apoptosis
respectively. mESCs were seeded onto the micropatterns, ensuring complete coverage, and
cultured for 16 hours. Subsequently, the medium was changed to a LIF-free medium. To assess

the middle stage of apoptosis after 48 hours differentiation, fixed cells were stained with
Caspase-3 (cleaved) as mentioned in the section of " Immunostaining". To assess the late stage
of apoptosis after 48 hours differentiation, live cells were stained with 7-AAD and Hoechst
33342 (Sigma-Aldrich, B2261-25MG) for 30 minutes at a 37°C incubator. Finally, the cells
were imaged using a spinning disk confocal microscope (OLYMPUS UPlanFL N 10x/0.30
N.A. objective, and Andor iXon camera). Quantification of apoptosis from nuclear morphology
was performed on DAPI images acquired by ART-seqFISH as previously described[63].
818
**Live cell imaging**
Fucci mESCs[65, 66] were kindly provided by Menno Ter Huurne and Hendrik Marks at Radboud
University. NANOG-GFP mESCs[58, 59] and Fucci mESCs were seeded onto the micropatterns,
ensuring complete coverage, and cultured for 16 hours. Subsequently, the medium was changed
to a LIF-free medium and live cell imaging was performed after LIF withdrawal. Imaging was
conducted using the SP8x AOBS-WLL confocal laser scanning microscope, equipped with a
Leica 40x/1.1 Water objective and a live-cell environmental chamber maintained at a
humidified temperature of 37°C and 5% $CO_2$. For NANOG-GFP mESCs, images were
acquired using a HyDs detector and a 488 nm laser. As for Fucci mESCs, images were acquired
using a HyDs detector, along with both 488 nm and 561 nm lasers. Z-series was obtained with
a step size of 1.5 μm. The time interval between image acquisitions was set to 1.5 hours. Single
cell tracking was performed manually using the Multi-point tool in Fiji[85].
831
832   **Image and data analysis**
833   Details described in Supplementary Information, Supplementary Note 2: Computational
834   pipelines development of ARTseq-FISH.
835
836   **Data availability**
837   All data is available in the main text or the supplementary information.

**References**

1.  Rivron, N.C. et al. Blastocyst-like structures generated solely from stem cells. *Nature* **557**, 106-111 (2018).
2.  Martyn, I., Kanno, T.Y., Ruzo, A., Siggia, E.D. & Brivanlou, A.H. Self-organization of a human organizer by combined Wnt and Nodal signalling. *Nature* **558**, 132-135 (2018).
3.  Girgin, M.U. et al. Bioengineered embryoids mimic post-implantation development in vitro. *Nat Commun* **12**, 5140 (2021).
4.  Lolas, M., Valenzuela, P.D.T., Tjian, R. & Liu, Z. Charting Brachyury-mediated developmental pathways during early mouse embryogenesis. *Proceedings of the National Academy of Sciences* **111**, 4478-4483 (2014).
5.  Li, Z. et al. BMP4 Signaling Acts via Dual-Specificity Phosphatase 9 to Control ERK Activity in Mouse Embryonic Stem Cells. *Cell Stem Cell* **10**, 171-182 (2012).
6.  Pauklin, S. & Vallier, L. The Cell-Cycle State of Stem Cells Determines Cell Fate Propensity. *Cell* **155**, 135-147 (2013).
7.  Lu, R. et al. Systems-level dynamic analyses of fate change in murine embryonic stem cells. *Nature* **462**, 358-362 (2009).
8.  Abranches, E. et al. Stochastic NANOG fluctuations allow mouse embryonic stem cells to explore pluripotency. *Development* **141**, 2770-2779 (2014).
9.  Hu, S. et al. Transcription factor antagonism regulates heterogeneity in embryonic stem cell states. *Mol Cell* **82**, 4410-4427.e4412 (2022).
10. Martyn, I., Brivanlou, A.H. & Siggia, E.D. A wave of WNT signaling balanced by secreted inhibitors controls primitive streak formation in micropattern colonies of human embryonic stem cells. *Development* **146** (2019).
11. Etoc, F. et al. A Balance between Secreted Inhibitors and Edge Sensing Controls Gastruloid Self-Organization. *Developmental Cell* **39**, 302-315 (2016).
12. Morgani, S.M., Metzger, J.J., Nichols, J., Siggia, E.D. & Hadjantonakis, A.K. Micropattern differentiation of mouse pluripotent stem cells recapitulates embryo regionalized cell fate patterning. *Elife* **7** (2018).
13. Chhabra, S., Liu, L., Goh, R., Kong, X. & Warmflash, A. Dissecting the dynamics of signaling events in the BMP, WNT, and NODAL cascade during self-organized fate patterning in human gastruloids. *PLoS Biol* **17**, e3000498 (2019).
14. Tewary, M. et al. A stepwise model of reaction-diffusion and positional information governs self-organized human peri-gastrulation-like patterning. *Development* **144**, 4298-4312 (2017).
15. Minn, K.T. et al. High-resolution transcriptional and morphogenetic profiling of cells from micropatterned human ESC gastruloid cultures. *Elife* **9**, e59445 (2020).
16. Blin, G. et al. Geometrical confinement controls the asymmetric patterning of brachyury in cultures of pluripotent cells. *Development* **145** (2018).
17. Plouhinec, J.-L., Simon, G., Vieira, M., Collignon, J. & Sorre, B. Dissecting signaling hierarchies in the patterning of the mouse primitive streak using micropatterned EpiLC colonies. *Stem Cell Rep* **17**, 1757-1771 (2022).
18. Merritt, C.R. et al. Multiplex digital spatial profiling of proteins and RNA in fixed tissue. *Nat Biotechnol* **38**, 586-599 (2020).
19. Vickovic, S. et al. High-definition spatial transcriptomics for in situ tissue profiling. *Nat Methods* **16**, 987-990 (2019).
20. Liu, Y. et al. High-Spatial-Resolution Multi-Omics Sequencing via Deterministic Barcoding in Tissue. *Cell* **183**, 1665-1681 e1618 (2020).
21. Stahl, P.L. et al. Visualization and analysis of gene expression in tissue sections by spatial transcriptomics. *Science* **353**, 78-82 (2016).

22.    Chen, A. et al. Spatiotemporal transcriptomic atlas of mouse organogenesis using
DNA nanoball-patterned arrays. *Cell* **185**, 1777-1792.e1721 (2022).
23.    Codeluppi, S. et al. Spatial organization of the somatosensory cortex revealed by
osmFISH. *Nature Methods* **15**, 932-935 (2018).
24.    Moffitt, J.R. et al. High-throughput single-cell gene-expression profiling with
multiplexed error-robust fluorescence in situ hybridization. *Proc Natl Acad Sci U S A*
**113**, 11046-11051 (2016).
25.    Chen, K.H., Boettiger, A.N., Moffitt, J.R., Wang, S. & Zhuang, X. Spatially resolved,
highly multiplexed RNA profiling in single cells. *Science* **348**, aaa6090 (2015).
26.    Takei, Y. et al. Integrated spatial genomics reveals global architecture of single
nuclei. *Nature* **590**, 344-350 (2021).
27.    Eng, C.-H.L. et al. Transcriptome-scale super-resolved imaging in tissues by RNA
seqFISH+. *Nature* **568**, 235-239 (2019).
28.    Wang, G., Moffitt, J.R. & Zhuang, X. Multiplexed imaging of high-density libraries
of RNAs with MERFISH and expansion microscopy. *Sci Rep-Uk* **8**, 4847 (2018).
29.    Rao, A., Barkley, D., Franca, G.S. & Yanai, I. Exploring tissue architecture using
spatial transcriptomics. *Nature* **596**, 211-220 (2021).
30.    Wang, F. et al. RNAscope: a novel in situ RNA analysis platform for formalin-fixed,
paraffin-embedded tissues. *J Mol Diagn* **14**, 22-29 (2012).
31.    Popovic, D., Koch, B., Kueblbeck, M., Ellenberg, J. & Pelkmans, L. Multivariate
Control of Transcript to Protein Variability in Single Mammalian Cells. *Cell Syst* **7**,
398-411 e396 (2018).
32.    Schulz, D. et al. Simultaneous Multiplexed Imaging of mRNA and Proteins with
Subcellular Resolution in Breast Cancer Tissue Samples by Mass Cytometry. *Cell*
*Syst* **6**, 25-36 e25 (2018).
33.    Goltsev, Y. et al. Deep Profiling of Mouse Splenic Architecture with CODEX
Multiplexed Imaging. *Cell* **174**, 968-981 e915 (2018).
34.    Saka, S.K. et al. Immuno-SABER enables highly multiplexed and amplified protein
imaging in tissues. *Nat Biotechnol* **37**, 1080-1090 (2019).
35.    Manesse, M., Patel, K.K., Bobrow, M. & Downing, S.R. in Biomarkers for
Immunotherapy of Cancer: Methods and Protocols. (eds. M. Thurin, A. Cesano &
F.M. Marincola) 585-592 (Springer New York, New York, NY; 2020).
36.    He, S. et al. High-plex imaging of RNA and proteins at subcellular resolution in fixed
tissue by spatial molecular imaging. *Nature Biotechnology* (2022).
37.    Vistain, L. et al. Quantification of extracellular proteins, protein complexes and
mRNAs in single cells by proximity sequencing. *Nature Methods* **19**, 1578-1589
(2022).
38.    van Buggenum, J.A.G. et al. Immuno-detection by sequencing enables large-scale
high-dimensional phenotyping in cells. *Nat Commun* **9**, 2384 (2018).
39.    Rivello, F. et al. Single-cell intracellular epitope and transcript detection reveals
signal transduction dynamics. *Cell Reports Methods* **1**, 100070 (2021).
40.    Sountoulidis, A. et al. SCRINSHOT enables spatial mapping of cell states in tissue
sections with single-cell resolution. *PLoS Biol* **18**, e3000675 (2020).
41.    Lee, H., Marco Salas, S., Gyllborg, D. & Nilsson, M. Direct RNA targeted in situ
sequencing for transcriptomic profiling in tissue. *Sci Rep* **12**, 7976 (2022).
42.    Alon, S. et al. Expansion sequencing: Spatially precise in situ transcriptomics in intact
biological systems. *Science* **371** (2021).
43.    Gyllborg, D. et al. Hybridization-based in situ sequencing (HybISS) for spatially
resolved transcriptomics in human and mouse brain tissue. *Nucleic Acids Res* **48**, e112
(2020).

44. Collaboration, T.A. et al. Astropy: A community Python package for astronomy. *A&A*
**558**, A33 (2013).

45. Zhou, Q.-Y., Park, J. & Koltun, V. Open3D: A modern library for 3D data processing.
*arXiv preprint arXiv:1801.09847* (2018).

46. Bradski, G. The openCV library. *Dr. Dobb's Journal: Software Tools for the*
*Professional Programmer* **25**, 120-123 (2000).

47. Waskom, M.L. Seaborn: statistical data visualization. *Journal of Open Source*
*Software* **6**, 3021 (2021).

48. McKinney, W. in Proceedings of the 9th Python in Science Conference, Vol. 445 51-
56 (Austin, TX, 2010).

49. Pedregosa, F. et al. Scikit-learn: Machine learning in Python. *the Journal of machine*
*Learning research* **12**, 2825-2830 (2011).

50. Hansen, M.M.K., Desai, R.V., Simpson, M.L. & Weinberger, L.S. Cytoplasmic
Amplification of Transcriptional Noise Generates Substantial Cell-to-Cell Variability.
*Cell Syst.* **7**, 384-397.e386 (2018).

51. van Sluijs, B., Maas, R.J.M., van der Linden, A.J., de Greef, T.F.A. & Huck, W.T.S.
A microfluidic optimal experimental design platform for forward design of cell-free
genetic networks. *Nature Communications* **13**, 3626 (2022).

52. Acampora, D. et al. Functional Antagonism between OTX2 and NANOG Specifies a
Spectrum of Heterogeneous Identities in Embryonic Stem Cells. *Stem Cell Reports* **9**,
1642-1659 (2017).

53. Guo, G. et al. Serum-Based Culture Conditions Provoke Gene Expression Variability
in Mouse Embryonic Stem Cells as Revealed by Single-Cell Analysis. *Cell Rep* **14**,
956-965 (2016).

54. Marks, H. et al. The Transcriptional and Epigenomic Foundations of Ground State
Pluripotency. *Cell* **149**, 590-604 (2012).

55. Hao, Y. et al. Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-
3587.e3529 (2021).

56. Raj, A., Peskin, C.S., Tranchina, D., Vargas, D.Y. & Tyagi, S. Stochastic mRNA
Synthesis in Mammalian Cells. *PLOS Biology* **4**, e309 (2006).

57. Semrau, S. et al. Dynamics of lineage commitment revealed by single-cell
transcriptomics of differentiating embryonic stem cells. *Nature Communications* **8**,
1096 (2017).

58. Desai, R.V. et al. A DNA repair pathway can regulate transcriptional noise to promote
cell fate transitions. *Science* **373**, eabc6506 (2021).

59. Sokolik, C. et al. Transcription Factor Competition Allows Embryonic Stem Cells to
Distinguish Authentic Signals from Noise. *Cell Syst* **1**, 117-129 (2015).

60. Altman, M.K. et al. Suppression of the GTPase-activating protein RGS10 increases
Rheb-GTP and mTOR signaling in ovarian cancer cells. *Cancer Lett* **369**, 175-183
(2015).

61. Salic, A. & Mitchison, T.J. A chemical method for fast and sensitive detection of
DNA synthesis *in vivo*. *Proceedings of the National Academy of Sciences*
**105**, 2415-2420 (2008).

62. Cai, C. et al. c-Myc regulates neural stem cell quiescence and activation by
coordinating the cell cycle and mitochondrial remodeling. *Signal Transduction and*
*Targeted Therapy* **6**, 306 (2021).

63. Filippi-Chiela, E.C. et al. Nuclear Morphometric Analysis (NMA): Screening of
Senescence, Apoptosis and Nuclear Irregularities. *PLOS ONE* **7**, e42522 (2012).

64. Miller, I. et al. Ki-67 is a Graded Rather than a Binary Marker of Proliferation versus
Quiescence. *Cell Rep* **24**, 1105-1112.e1105 (2018).

65. Ter Huurne, M., Chappell, J., Dalton, S. & Stunnenberg, H.G. Distinct Cell-Cycle Control in Two Different States of Mouse Pluripotency. *Cell Stem Cell* **21**, 449-455.e444 (2017).

66. ter Huurne, M. et al. Critical Role for P53 in Regulating the Cell Cycle of Ground State Embryonic Stem Cells. *Stem Cell Rep* **14**, 175-183 (2020).

67. Ferro, A. et al. Blue intensity matters for cell cycle profiling in fluorescence DAPI-stained images. *Laboratory Investigation* **97**, 615-625 (2017).

68. Olofsson, K., Carannante, V., Takai, M., Önfelt, B. & Wiklund, M. Single cell organization and cell cycle characterization of DNA stained multicellular tumor spheroids. *Sci Rep-Uk* **11**, 17076 (2021).

69. Roukos, V., Pegoraro, G., Voss, T.C. & Misteli, T. Cell cycle staging of individual cells by fluorescence microscopy. *Nat Protoc* **10**, 334-348 (2015).

70. Tao, Y. et al. Highly efficient and robust π-FISH rainbow for multiplexed in situ detection of diverse biomolecules. *Nature Communications* **14**, 443 (2023).

71. Palla, G., Fischer, D.S., Regev, A. & Theis, F.J. Spatial components of molecular tissue biology. *Nature Biotechnology* **40**, 308-318 (2022).

72. Stuart, T. & Satija, R. Integrative single-cell analysis. *Nature Reviews Genetics* **20**, 257-272 (2019).

73. Qin, Y. et al. A multi-scale map of cell structure fusing protein images and interactions. *Nature* **600**, 536-542 (2021).

74. Szemes, M. et al. Diagnostic application of padlock probes--multiplex detection of plant pathogens using universal microarrays. *Nucleic Acids Res* **33**, e70 (2005).

75. Fang, C. et al. MiRNA Detection Using a Rolling Circle Amplification and RNA-Cutting Allosteric Deoxyribozyme Dual Signal Amplification Strategy. *Biosensors (Basel)* **11** (2021).

76. Lin, C. et al. Visualization of individual microRNA molecules in fixed cells and tissues using target-primed padlock probe assay. *Biochemical and Biophysical Research Communications* **526**, 607-611 (2020).

77. Kumari, R. et al. A Novel Rolling Circle Amplification-Based Detection of SARS-CoV-2 with Multi-Region Padlock Hybridization. *Diagnostics* **12** (2022).

78. Söderberg, O. et al. Direct observation of individual endogenous protein complexes in situ by proximity ligation. *Nature Methods* **3**, 995-1000 (2006).

79. Jain, S., Dandy, D.S., Geiss, B.J. & Henry, C.S. Padlock probe-based rolling circle amplification lateral flow assay for point-of-need nucleic acid detection. *Analyst* **146**, 4340-4347 (2021).

80. Zhao, Y. et al. Single Cell RNA Expression Analysis Using Flow Cytometry Based on Specific Probe Ligation and Rolling Circle Amplification. *Acs Sensors* **5**, 3031-3036 (2020).

81. van Mierlo, G. et al. Integrative Proteomic Profiling Reveals PRC2-Dependent Epigenetic Crosstalk Maintains Ground-State Pluripotency. *Cell Stem Cell* **24**, 123-137.e128 (2019).

82. Théry, M. & Piel, M. Adhesive micropatterns for cells: a microcontact printing protocol. *Cold Spring Harb Protoc* **2009**, pdb.prot5255 (2009).

83. van Kooten, T.G., Spijker, H.T. & Busscher, H.J. Plasma-treated polystyrene surfaces: model surfaces for studying cell-biomaterial interactions. *Biomaterials* **25**, 1735-1747 (2004).

84. Bao, M., Xie, J., Piruska, A. & Huck, W.T.S. 3D microniches reveal the importance of cell size and shape. *Nat Commun* **8**, 1962 (2017).

85. Schindelin, J. et al. Fiji: an open-source platform for biological-image analysis. *Nature Methods* **9**, 676-682 (2012).

**Acknowledgments**

We thank members of the Huck and Hansen laboratories for the thoughtful discussion and suggestions; Aigars Piruska for making the silicon masters with micropatterns and technical assistance with CSU spinning disk confocal; Erik van Buijtenen and Kinga Matuła for sharing the antibody conjugation protocols, and fruitful discussions and suggestions from Erik van Buijtenen. We also thank Leor Weinberger for the NANOG-GFP mESC line as well as Hendrik Marks for the mESC-E14 and mESC Fucci cell line and thoughtful suggestions. We acknowledge generous support from Radboud University (W.T.S.H.), the Christine Mohrmann fellowship (M.M.K.H), the European Union (ERC, ChOICE, 101041939, M.M.K.H) and the Netherlands Organization for Scientific Research (NWO) (ENW-XS awards OCENW.XS3.055 and OCENW.XS21.2.050, both M.M.K.H., and a Spinoza grant, W.T.S.H.). X.H. acknowledges financial support from the China Scholarship Council (CSC).

**Author information**

Authors and Affiliations

**Institute for Molecules and Materials, Radboud University, Heyendaalseweg 135, 6525 AJ Nijmegen, the Netherlands**

Xinyu Hu, Bob van Sluijs, Óscar García-Blay, Yury Stepanov, Koen Rietrae, Wilhelm T.S. Huck, Maike M. K. Hansen.

Contributions

X.H., B.v.S., W.T.S.H., and M.M.K.H. conceived and designed the study. X.H. developed and optimized the experimental procedure for ARTseq-FISH. B.v.S. developed the software for ARTseq-FISH. Ó.G.B. helped with probe design and optimization of ARTseq-FISH workflow. Ó.G.B., also conducted supporting experiments. Y.S., and K.R., contributed to validation experiments of ARTseq-FISH. X.H., B.v.S., W.T.S.H., and M.M.K.H. wrote the manuscript and Ó.G.B. edited the manuscript.

Corresponding author

Correspondence to Wilhelm T.S. Huck and Maike M. K. Hansen.

**Ethics declarations**
Competing interests
Authors declare no competing interests

**Supplementary Information:**

Supplementary Notes 1 Experimental development of ARTseq-FISH

Supplementary Notes 2 Computational pipeline development of ARTseq-FISH

Supplementary Figs. 1-31

Supplementary Table Captions

References

**Supplementary Tables 1-5**