Rapid learning of neural circuitry from holographic ensemble stimulation enabled by model-based compressed sensing

Marcus A. Triplett[1,2,3,4,†,*], Marta Gajowa[5,*], Benjamin Antin[1,2,3,4], Masato Sadahiro[5], Hillel Adesnik[5], and Liam Paninski[1,2,3,4]

[1]Mortimer B. Zuckerman Mind Brain Behavior Institute, Columbia University, NY
[2]Grossman Center for the Statistics of Mind, Columbia University, NY
[3]Center for Theoretical Neuroscience, Columbia University, NY
[4]Department of Statistics, Columbia University, NY
[5]Department of Molecular and Cell Biology, University of California, Berkeley, CA
[†] Correspondence: marcus.triplett@columbia.edu
[*] These authors contributed equally

Abstract

Discovering how neural computations are implemented in the cortex at the level of monosynaptic connectivity requires probing for the existence of synapses from possibly thousands of presynaptic candidate neurons. Two-photon optogenetics has been shown to be a promising technology for mapping such monosynaptic connections via serial stimulation of neurons with single-cell resolution. However, this approach is limited in its ability to uncover connectivity at large scales because stimulating neurons one-by-one requires prohibitively long experiments. Here we developed novel computational tools that, when combined, enable learning of monosynaptic connectivity from high-speed holographic neural ensemble stimulation. First, we developed a model-based compressed sensing algorithm that identifies connections from postsynaptic responses evoked by stimulation of many neurons at once, considerably increasing the rate at which the existence and strength of synapses are screened. We show that this model-based approach, explicitly incorporating known biophysics of optogenetic mapping experiments, is critical for accurately determining synaptic connectivity using compressed sensing. Second, we developed a deep learning method that isolates the postsynaptic response evoked by each stimulus, allowing stimulation to rapidly switch between ensembles without waiting for the postsynaptic response to return to baseline. Together, our system increases the throughput of monosynaptic connectivity mapping by an order of magnitude over existing approaches, enabling the acquisition of connectivity maps at speeds needed to discover the synaptic circuitry implementing neural computations.

Introduction

The structure of synaptic connectivity is central to how the brain processes information, stores long-term memories, and mediates cognition. To uncover such connectivity, two-photon optogenetic stimulation has emerged as a promising technology due to its ability to flexibly probe neurons with single-cell resolution while monitoring postsynaptic currents (PSCs) using whole-cell recordings [Packer et al., 2012, Baker et al., 2016, Shemesh et al., 2017, Naka et al., 2019, Hage et al., 2022, Printz et al., 2023]. Yet existing optogenetic circuit mapping techniques have been limited to probing

connectivity from small numbers of neurons that must be slowly stimulated one-by-one, and therefore require aggregating small-scale maps across experiments to obtain large-scale maps of connectivity [Hage et al., 2022]. By contrast, an ideal monosynaptic connectivity mapping technique would enable large numbers of synaptic connections to be identified at high speed within a single experimental session. This would provide a crucial advantage in that each experiment would produce a more comprehensive and representative map of neural circuitry, rather than having to pool together smaller experiments that each provide just a partial view of connectivity.

Multiple strategies could serve to advance optogenetic circuit mapping towards this state, although each introduces experimental and computational challenges. The simplest strategy to improve the throughput of a connectivity mapping experiment is to increase the rate at which stimulation switches between neurons. This is primarily determined by the refresh rate of the holographic spatial light modulator (SLM) if using advanced light sculpting techniques, or, in the case of laser scanning approaches, just the time required for a neuron to integrate photocurrent until it elicits an action potential [Rickgauer and Tank, 2009]. However, while this approach allows a mapping experiment to be completed in a shorter time frame, naively stimulating too quickly confounds postsynaptic measurements because the membrane conductance of the postsynaptic neuron will not have sufficient time to return to baseline conditions before the next stimulus is applied. The speed at which a mapping experiment can feasibly take place therefore becomes limited by the ability to computationally demix PSC waveforms that overlap in time.

Simulation studies predict that connectivity mapping could also be greatly accelerated through the use of compressed sensing, where, in principle, sparse connectivity could be reconstructed from few measurements provided that stimulation is applied to ensembles of randomly selected neurons at once [Hu and Chklovskii, 2009, Fletcher et al., 2011, Mishchenko and Paninski, 2012, Shababo et al., 2013, Draelos and Pearson, 2020, Navarro and Oweiss, 2023]. However, the efficacy of existing compressed sensing algorithms is fundamentally limited outside of simplified simulations because they neglect the complex biophysics intrinsic to mapping experiments. For example, failures to elicit spikes in presynaptic neurons (due to inadequate laser power or physiological stochasticity) or for spikes to propagate across synapses are known to significantly impact the accuracy of compressed sensing [Hu and Chklovskii, 2009, Navarro and Oweiss, 2023]. Further, high rates of spontaneous synaptic currents could lead to both false-positive connections and mischaracterized synaptic weights [Shababo et al., 2013, Printz et al., 2023].

To overcome these limitations and enable high-speed, large-scale connectivity mapping experiments, we developed new computational tools for inferring connectivity from holographic stimulation of neural ensembles. While existing optogenetic circuit mapping techniques perform slow single-neuron stimulation, our tools allow experiments to proceed with rapid holographic ensemble stimulation, minimizing downtime of the holographic SLM since evoked PSCs are isolated in time, deconfounded of spontaneous PSCs, and cleaned of electrical noise by a computational demixing procedure post-hoc. To infer connectivity, we developed a model-based compressed sensing algorithm that simultaneously estimates synaptic weights, presynaptic spikes, and how such spikes depend on laser power, while accounting for biophysical constraints and any residual spontaneous PSCs not already eliminated by the demixing procedure. To validate these tools, we performed connectivity mapping experiments using the recently engineered family of fast, potent ChroME2.0 opsins [Sridharan et al., 2022] together with two-photon holographic light sculpting [Pégard et al., 2017]. We routinely mapped connectivity from hundreds of presynaptic candidates within 680x680x100 μm^3 volumes of cortex, together totalling more than 12,000 probed targets. By combining rapid ensemble stimulation with computational demixing and model-based compressed sensing, we reduced the stimulation time required to reconstruct connectivity by an order of magnitude over existing approaches, allowing large numbers of synapses to be quickly mapped within individual experimental sessions.

To facilitate the use of our techniques, we provide an open access software toolbox and web-based notebook implementation that researchers can straightforwardly use and validate in their own mapping experiments. While our analysis shows that the highest mapping throughput is obtained using holographic ensemble stimulation, our techniques also increase the accuracy and speed of more widely accessible mapping approaches based on conventional single-target stimulation, and therefore can enable fast and accurate connectivity mapping in a broad range of laboratories.

Results

Optogenetic circuit mapping framework

Accurately characterizing the presence and strength of a synaptic connection requires precise control over the initiation of presynaptic action potentials. Ideally, each stimulus would evoke just a single presynaptic spike since this provides a direct measurement of synaptic charge transfer. To this end, we combined the family of potent soma-targeted ChroME opsins [Mardinly et al., 2018, Sridharan et al., 2022] with a scanless computer-generated holography system named 3D-SHOT [Pégard et al., 2017] (Figure 1a). We first expressed ChroME2f in parvalbumin-expressing (PV) neurons by viral transfection with an adeno-associated virus [Sridharan et al., 2022]. We calibrated laser powers and illumination time such that brief (3-5 ms) periods of stimulation almost always led to either 0 or 1 action potential, with minimal instances of multiple action potentials resulting from a single pulse (Figure S1). Next, we confirmed through a similar process that we could reliably evoke action potentials when holographically stimulating ensembles of neurons at once (Figure S1, Figure S2). Finally, we combined holographic stimulation of ensembles of PV neurons with whole-cell recordings from pyramidal neurons. We focused our initial experiments on mapping connections of this kind for two reasons. First, in this preparation opsin expression is restricted to PV neurons, and thus any measurements of evoked PSCs in a pyramidal neuron would not be affected by direct photocurrents arising from stimulation at locations near the recording electrode. Second, PV neurons make strong synaptic connections onto pyramidal neurons [Packer and Yuste, 2011], and therefore were likely to be reliably identified computationally.

Having established the high spatiotemporal precision of our holographic stimulation technique here and in previous work [Pégard et al., 2017, Bounds et al., 2021, Sridharan et al., 2022], we then developed a computational system for processing optogenetic data and inferring synaptic connectivity. The first component of our system seeks to extract precise measurements of optogenetically-evoked PSCs that have been affected by three adverse factors intrinsic to mapping experiments: (1) electrical noise arising from the recording electrode, which increases the variability of the observed synaptic currents and can require additional repetitions of each presynaptic stimulus to overcome; (2) spontaneous currents that, depending on their timing relative to the optogenetic stimulus, can obscure optogenetically-evoked PSCs or increase the number of false-positive connections; and (3) postsynaptic responses from preceding or subsequent trials when the interstimulus interval is shorter than the PSC decay time, which are prevalent when attempting to perform mapping experiments at very high speeds. To simultaneously eliminate these factors, we developed a deep neural network architecture for demixing and denoising PSC waveforms (Figure 1b,c) in a process that we call neural waveform demixing (NWD). The NWD network attempts to isolate optogenetically evoked PSCs stimulus-by-stimulus, such that confounding synaptic currents not driven by the optogenetic stimulus are "subtracted out" by the network and the resulting currents cleaned of electrical noise. Consequently, the NWD network allows experiments to proceed with very short interstimulus intervals since evoked PSCs from previous and subsequent trials are subtracted out and the initial baseline current reset to zero (further details given below).

The second component of our system is a model-based compressed sensing algorithm (Figure 1d). The key challenge for a compressive approach to connectivity mapping is to robustly infer presynaptic spikes from PSCs evoked by ensemble stimulation despite a multitude of unobserved sources of biophysical variability. For example, the illumination intensity required to elicit a presynaptic spike varies unpredictably between neurons because opsin is expressed differentially throughout a neural population, regardless of whether the opsin construct is delivered using a viral or transgenic targeting strategy [Bounds et al., 2021]. In a mapping experiment, this relationship between light and presynaptic spike probability cannot be observed from the presynaptic neuron itself, but must be inferred through electrophysiological measurements from the postsynaptic neuron. This inference problem is in turn complicated by the fact that PSCs can occur spontaneously, that postsynaptic responses are corrupted by electrical noise, that synaptic transmission can fail despite successful elicitation of a presynaptic spike, and that potentially tens of neurons are stimulated at a time. Accounting for the power-dependence of presynaptic spike initiation for individual neurons when only observing postsynaptic responses

Figure 1: Overview of proposed high-throughput optogenetic circuit mapping framework. a, A fast, potent, and soma-targeted opsin (here ChroME2f) is expressed in a pool of candidate presynaptic neurons while intracellular currents are monitored from a voltage-clamped postsynaptic neuron. Two-photon holography is used to elicit spikes in ensembles of presynaptic candidates, increasing the number of putative synaptic connections tested compared to single-target stimulation. **b**, To further increase throughput, stimulation is applied at high speeds (e.g., $30-50$ Hz), such that the postsynaptic membrane conductance is not guaranteed to return to baseline conditions (i.e., trials are "overlapping" in time). This allows more synapses to be tested per minute, but complicates downstream identification of their existence and strength unless the evoked PSCs are subsequently demixed. **c**, A neural network is trained to isolate the components of the postsynaptic response that are plausibly due to the optogenetic stimulus, allowing high-fidelity measurements of postsynaptic current even at very high stimulation frequencies. **d**, The "demixed" postsynaptic responses evoked by holographic ensemble stimulation are used in a model-based compressed sensing algorithm (CAVIaR) to reconstruct the underlying connectivity. Unlike traditional compressed sensing techniques, compressive mapping of synaptic circuitry requires inference of any optogenetically elicited presynaptic spikes (left) as well as the synapses themselves (right). CAVIaR alternates between inference of presynaptic spikes and synaptic connectivity, in addition to a number of other pertinent biophysical variables (see main text). **e**, Presynaptic neurons (white circles) identified via the proposed pipeline in an example experiment mapping connectivity from presynaptic PV neurons to a postsynaptic pyramidal neuron. Probed targets considered unconnected shown as red circles.

to ensemble stimulation is therefore computationally challenging, and constitutes only one of the inference problems that must be solved to accurately reconstruct connectivity using a compressed sensing approach.

To resolve these challenges, we embedded the compressed sensing step in a hierarchical Bayesian statistical model. Our model relates patterns of ensemble stimulation to electrophysiological measurements from the postsynaptic cell while, at the same time, accounting for variability in photoactivation and synaptic transmission across the opsin-expressing population. We developed a variational inference technique called CAVIaR (coordinate-ascent variational inference and isotonic regularization) to learn posterior distributions over the model parameters (Methods). CAVIaR identifies the existence and strength of individual synaptic connections from ensemble stimulation, substantially increasing the rate at which neural circuits can be mapped compared to single-target stimulation and increasing the accuracy compared to existing compressed sensing techniques.

Neural waveform demixing allows mapping experiments to proceed rapidly

The NWD network has a sequential U-Net architecture [Ronneberger et al., 2015, Falk et al., 2019] that uses one dimensional convolutional filters to learn representations of the input signal at increasing levels of temporal compression. This allows the network to integrate information across the entire PSC trace, such that confounding synaptic currents preceding or following an admissible PSC "initiation zone" (typically selected to be a window of 3-12 ms following stimulation) can be accurately subtracted and the baseline current reset to zero (Figure 2a). Since the network is trained to generate noise-free, demixed PSCs from tens of thousands of simulated noisy inputs, NWD also markedly improves the overall signal quality (Figure 2). Furthermore, unlike computationally intensive algorithms for deconvolving intracellular currents [Merel et al., 2016], NWD can isolate the PSC evoked by optogenetic stimulation with just a simple forward pass through the network, with the resulting demixed PSCs usable in any connectivity inference algorithm and in real time.

Performing optogenetic connectivity mapping experiments at high speed poses a trade-off: the experiment can be completed much more quickly, but measurements of the postsynaptic response are much more confounded by currents from previous and subsequent trials. For this reason, a typical rate of stimulation for a connectivity mapping experiment is ~10 Hz (i.e., stimulating a neuron approximately every 100 ms [Hage et al., 2022, Printz et al., 2023, Chen et al., 2023], much longer than the decay time of a typical PSC), which we take as a baseline rate in later method comparisons. To characterize the improvement in experiment speed-up resulting from the application of NWD, we simulated connectivity mapping experiments at full 20 kHz sampling resolution with varying synaptic connection densities and stimulation frequencies (Figure 2b, Figure S3). In a sparsely connected circuit, the primary action of NWD is to eliminate spontaneous synaptic currents and reduce electrical noise, leading to just a marginal improvement in signal fidelity compared to the raw PSC traces (Figure 2b, 0.1 connection density example). However, as synapses become more prevalent the occurrence of postsynaptic events greatly increases, with many confounding synaptic currents elicited from previous trials. Application of NWD therefore leads to a substantial improvement in the signal fidelity of the postsynaptic response at high stimulation frequencies (Figure 2b, 0.2 to 0.4 connection density examples). Similarly, the fidelity of the postsynaptic response degrades as a function of the number of simultaneously stimulated neurons and the background rate of spontaneous synaptic currents (Figure 2b, middle and right panels). However, NWD largely eliminates this effect, and thus reduces experimental constraints on the duration of interstimulus intervals since confounding synaptic currents are significantly reduced. Stimulation can therefore be increased to rates closer to the refresh rate of the SLM. In practice, this enables us to stimulate much faster than the ~40 ms decay time typical of PSC traces in our preparations (e.g., Figure 2d).

We applied NWD to each of our experimental settings, including demixing of inhibitory PSCs from PV-pyramidal and SST-pyramidal mapping experiments under holographic single-target and ensemble stimulation (Figure 2d-e), as well as demixing of excitatory PSCs from pyramidal-pyramidal mapping experiments (Figure 2f). We found that we could improve the performance of the NWD network by tuning it to the time constants of either inhibitory or excitatory synaptic currents

Figure 2: Simultaneous demixing and denoising of optogenetically evoked postsynaptic currents. a, Neural waveform demixing overview. Raw electrophysiological traces (left) are confounded by synaptic currents from previous and subsequent trials during fast stimulation, in addition to spontaneous synaptic currents. The NWD network (middle) is trained to isolate the PSC waveform associated with the optogenetic stimulus, resulting in a demixed trace (right, red) with the confounding synaptic currents subtracted and baseline current (at 0 ms) adjusted to 0 nA. Note that a 5 ms "pre-stimulus" window is provided to the network as additional context. **b**, Performance of NWD in simulation as a function of stimulation frequency under increasing connection density (left), number of simultaneously targeted cells (middle) and rate of spontaneous activity (right). In all cases NWD leads to a substantial improvement in signal fidelity (measured by the mean square error of the raw vs demixed trace compared to the ground truth evoked PSC). Errors are evaluated across all conditions using a single trained network. **c, d**, Examples of NWD applied to optogenetically evoked inhibitory PSCs under single-target (c) and ten-target (d) holographic stimulation of PV neurons expressing ChroME2f. Panel (d) shows responses to stimulating ensembles containing the neuron in panel (c). **e**, Same as (d), but for stimulation of SST neurons. **f**, same as (c), but for stimulation of pyramidal neurons expressing ChroME2s [Sridharan et al., 2022] while recording from another pyramidal neuron.

(Figure S4). In each case the NWD network led to a dramatic reduction in confounding synaptic currents and electrical noise, in agreement with our simulated results (Figure 2b).

Ensemble stimulation can test for many synaptic connections simultaneously

We sought to capitalize on recent technological developments in opsin engineering [Baker et al., 2016, Shemesh et al., 2017, Mardinly et al., 2018, Sridharan et al., 2022], holographic light sculpting [Papagiakoumou et al., 2010, Pégard et al., 2017, Papagiakoumou et al., 2020, Adesnik and Abdeladim, 2021] and fundamental mathematical results on efficient signal reconstruction in sparse settings [Donoho, 2006, Candes and Tao, 2006, Candes et al., 2006] by developing a statistical method for inferring synaptic connectivity from holographic ensemble stimulation. This was based on the logic that pairing rapid stimulation and computational demixing with the ability to test for many potential connections with each hologram could dramatically speed up circuit mapping. We therefore created and tested a variety of connectivity inference algorithms, and ultimately found that a variational inference approach for a hierarchical Bayesian statistical model was able to overcome the limitations of existing compressed sensing approaches. Our model relates patterns of holographic stimulation to PSCs through a series of key latent variables: (1) optogenetic "power curves" that characterize the relationship between laser power and presynaptic spike probability due to variation in opsin expression and rheobase across neurons, (2) presynaptic spikes successfully elicited by photostimulation and transmitted to the postsynaptic

neuron, and (3) synaptic weights that determine the amplitude of the resulting PSCs (Figure 3a). We learn posterior distributions over the latent variables using CAVIaR, an algorithm that iteratively updates the parameters of a variational approximation to the posterior such that the inferred presynaptic spikes respect the biophysical plausibility constraint of having isotonically increasing spike probabilities as a function of laser power (i.e., such that the probability of evoking a presynaptic spike increases with laser power on average; see Methods).

A key component of our experimental design was to randomly switch between three or more laser powers while mapping. Since CAVIaR infers presynaptic spikes, we could then estimate optogenetic power curves to determine whether the plausibility constraint was satisfied. Any putative presynaptic neurons for which spike counts happen to implausibly decrease with increasing power are considered spurious, and immediately disconnected by CAVIaR. When using our software in practice, a user simply sets a parameter that imposes a minimum success rate (between 0 and 1) for evoking presynaptic spikes (given by the isotonically-constrained power curve at maximal power) and CAVIaR identifies putatively connected neurons meeting that criterion. However, PSCs can occur spontaneously, inflating the estimated presynaptic spike rate and increasing the number of false positives. We therefore simultaneously estimate the background rate of spontaneous currents and use this to adaptively adjust the plausibility criterion during inference.

We compared CAVIaR to two other approaches. (1) Standard compressed sensing. While there are a number of widely-used algorithms for compressed sensing [Rani et al., 2018], they each solve a common mathematical problem and therefore lead to closely related results. Compressive sampling matching pursuit (CoSaMP, [Needell and Tropp, 2009]) is a particularly scalable algorithm for solving the compressed sensing problem and was used in the first simulation study on compressive connectivity mapping [Hu and Chklovskii, 2009] which we therefore benchmark CAVIaR against and consider representative of the performance from related studies (e.g., ref. [Chen et al., 2023]). (2) A variational inference technique for a spike-and-slab connectivity model similar to CAVIaR but that, among other methodological differences, does not impose isotonic regularization or account for spontaneous synaptic currents (CAVI-SnS, [Shababo et al., 2013]). We evaluated the ability of each method to reconstruct connectivity in simulations where the sparsity of the underlying connectivity varied, since compressed sensing-based techniques are known to be particularly sensitive to this parameter [Donoho, 2006]. We found that CAVIaR greatly outperformed conventional compressed sensing (Figure 3b, left vs right panels), primarily because the latter does not have a model for stochastic, power-dependent presynaptic spikes. Importantly, in line with earlier results [Hu and Chklovskii, 2009], we found that algorithms failing to account for this stochasticity will yield strongly biased synaptic weight estimates since they will average over postsynaptic measurements when no presynaptic spikes occurred. CAVIaR also outperformed CAVI-SnS, which transiently demonstrated high accuracy, but ultimately generated increasingly worse estimates of connectivity with increasing amounts of data since it lacked mechanisms to prevent spontaneous PSCs from being misidentified as false positives. We found that CAVIaR consistently achieved state-of-the-art performance when repeating the analysis while varying the number of times each stimulus was repeated, the number of simultaneous targets, and the rate of background spontaneous activity as a function of the amount of experimental data collected (Figure S7, Figure S8).

Next, we estimated the speed-up obtained by combining rapid ensemble stimulation with NWD and CAVIaR, and compared this against previously established techniques. Our simulations showed that even with large population sizes (e.g., mapping 1000 potential presynaptic neurons), the speed of connectivity mapping was optimal when stimulating ensembles of 20 neurons at a time (Figure S8) beyond which the accuracy of presynaptic spike inference gradually degraded. Our speed-up simulations were therefore performed using 20-cell stimulation. We anticipated that the background rate of spontaneous PSCs would also be a key factor influencing the ultimate accuracy of each connectivity inference method, since a high number of spontaneous PSCs that happen to fall within the stimulation window could mimic a synapse and lead to false positives. Single-target mapping at 10 Hz with CAVI-SnS required more than 30 minutes of stimulation to cross an accuracy of 0.95 in simulations with 1000 neurons, 10% connectivity, and 1 Hz spontaneous activity (Figure 3c, left panel; accuracy measured as the R^2 between the true and estimated synaptic weights). By comparison, NWD-enabled 50 Hz stimulation of 20-cell ensembles recovered synaptic connectivity at an accuracy exceeding 0.95 with just

Figure 3: CAVIaR accurately maps synaptic connectivity at high speeds in simulation. **a**, Statistical model relating patterns of holographic stimulation to measurements of postsynaptic current via latent optogenetic power curves, presynaptic spikes, and synaptic weights. **b**, Comparison of CAVIaR (ours, left) with CAVI-SnS (middle) and CoSaMP (right) for varying connection densities as a function of increasing amounts of experimental data. Simulated presynaptic candidate population size of N=1000 (hence, e.g., 4500 trials corresponds to an average of just 4.5 stimuli per neuron). All PSCs are demixed using NWD. Note that CoSaMP requires the number of desired connections to be specified in advance; we therefore provided CoSaMP with the true number of connections, such that the reported performance represents an upper bound. R^2 represents the coefficient of determination between the ground-truth and inferred connectivity vectors. **c**, Order of magnitude improvement in mapping efficiency using NWD-accelerated circuit mapping with ensemble stimulation. Left, convergence speed of CAVIaR, CoSaMP, and CAVI-SnS with and without NWD, assuming a background rate of spontaneous PSCs of 1 Hz. Without NWD, the model-based methods CAVIaR and CAVI-SnS perform substantially more poorly than with NWD enabled. A higher rate of spontaneous activity (5 Hz, right panel) causes fast degradation of CAVI-SnS performance, but impacts CAVIaR and CoSaMP much less. Error bars show mean $\pm$ 1 s.d. over 20 simulations.

30 seconds of stimulation using CAVIaR, resulting in a rate of connectivity testing of more than 2000 neurons per minute in low spontaneous activity conditions. In the same time period CoSaMP obtained an accuracy of less than 0.6. Use of our computational tools in optimal conditions is therefore predicted to enable a speed-up of more than an order of magnitude.

Increasing the rate of spontaneous activity (to 5 Hz, a more challenging regime) revealed several important behaviors of CAVIaR and CAVI-SnS (Figure 3c, right panel). Namely, the accuracy of CAVI-SnS both with and without NWD behaved non-monotonically (similar to Figure 3b, final R^2 at 5 Hz with NWD, 0.74; without NWD, <0), with connectivity estimates ultimately getting worse with additional experiment time due to the accumulation of spontaneous PSCs. However, CAVIaR with NWD obtained nearly the same accuracy when spontaneous PSCs occurred at 5 Hz as it did when they occurred at 1 Hz (final R^2 with spontaneous rate 1 Hz, 0.98; 5 Hz, 0.91) due to CAVIaR's built-in mechanisms for estimating and adapting to spontaneous PSCs. These results thus indicate that our model-based compressed sensing approach should be much more robust in conditions with high spontaneous activity.

Validating CAVIaR using mapping experiments in cortical slices

Having established the accuracy and speed of our connectivity mapping system in simulations, we next sought to validate our computational tools experimentally. We first wanted to confirm that connectivity maps obtained using ensemble stimulation were closely aligned to those obtained using the existing approach of single-target stimulation. We therefore performed experiments where the same population of PV neurons was mapped using randomly interleaved trials of single-target and holographic ensemble stimulation (ensembles of 10 neurons, stimulation performed at 30 Hz). We separated trials into two sets depending on whether they were obtained using single-target or ensemble stimulation, demixed the optogenetically evoked PSCs using NWD, and detected putative synapses in each set independently using CAVIaR (Figure 4a-d). For the example shown in Figure 4a, model-based compressed sensing identified the same set of synaptically connected PV neurons as with single-target stimulation and almost identical synaptic strengths (varying from 57 to 228 pA, Figure 4c,d). Across 14 different experiments we probed 2619 total presynaptic candidates (mean number of targets probed, 187; minimum, 107; maximum, 269). CAVIaR identified very similar numbers of connections when using single-target stimulation compared to ensemble stimulation (Figure 4e,f; R^2, 0.81). Further, we confirmed that these were primarily the same set of connections when evaluated based on both their numerical synaptic strengths (Figure 4g, left; average R^2 between connectivity vectors estimated using single-target and ensemble stimulation, 0.89), as well their binary classification (i.e., whether an ROI was connected or not, without regard to synaptic strength; Figure 4g, right; average Jaccard index, 0.79). Additional examples are given in Figure S9. We note, however, that connectivity mapping using single-target stimulation does not necessarily represent a perfect ground truth as false positives can arise from spontaneous activity (Figure 3c), and so some discrepancy should be expected.

In some experimental preparations opsin-expressing neurons do not also express a separate nuclear fluorophore for straightforward identification of photosensitive cell nuclei [Bounds et al., 2021]. We therefore considered whether our connectivity mapping system could be used in the case where neuron locations are not known *a priori* (i.e., when mapping is performed "blind"). In this context, the field of view is divided into a three-dimensional grid and locations are stimulated one grid point at a time [Wang et al., 2007, Baker et al., 2016, Naka et al., 2019]. We performed both single-target and 10-target stimulation of grid points at 30 Hz and applied NWD and CAVIaR independently to the two sets of mapping data. Both single-target and ensemble stimulation identified a common ROI, and overlaying the stimulation grid on an image of the underlying tissue showed that this ROI aligned with a unique opsin-expressing neuron (Figure S12; see Figure S13 for full, multi-plane analysis of blind grid mapping experiment). This confirmed that NWD and CAVIaR could also facilitate rapid connectivity mapping in the blind stimulation regime.

We next wanted to confirm that CAVIaR inferences are well-calibrated. Specifically, since compressive approaches to connectivity mapping depend closely on the accuracy of the presynaptic spike inference step [Hu and Chklovskii,

a Single-target stimulation | Ensemble stimulation

Connected ROIs: 1, 2, 3, 4, 5, 6, 7
Unconnected ROIs: 8, 9, 10, ⋮, 167, 168, 169

30 mW   45 mW   60 mW

300 pA / 45 ms

■ detected presynaptic spike (from listed neuron)   ★ detected spontaneous

b Ensemble (detection) | Single-target (validation)

Current (pA): 0, 100, 200, 300, 400, 500
Time (ms): 0, 20, 40

Raw / Demixed

c Inferred synaptic weight (this example)

$R^2=0.99$

Ensemble stim. (pA): 0, 100, 200, 300
Single-target stim. (pA): 0, 100, 200, 300

f Number of inferred connections (all data)

Ensemble stimulation: 0, 5, 10, 15
Single-target stimulation: 0, 10, 20

g Score (all data)

1.0, 0.8, 0.6, 0.4, 0.2, 0.0

R^2 Jaccard index

d Inferred presynaptic neurons (z-projection, 5 planes)

single-target stimulation | compressed sensing $R^2=0.99$, Jaccard index=1

PV-expressing neurons
Pyramidal neuron
50 μm

e single-target stimulation | compressed sensing $R^2=0.96$, Jaccard index=0.8

h n=4 pairs
post   pre

i pre   post
50 μm

j Example ground-truth spiking scenarios

(i) single-target stim true positive
(ii) ensemble stim true positive
3ms stim
post
pre

(iii) ensemble stim false negative
(iv) ensemble stim synaptic failure

0.1 nA
0.2 nA
10 ms

k Presynaptic spike accuracy

30 mW   45 mW   60 mW
1.00, 0.75, 0.50, 0.25, 0.00

Single   Ensemble

Figure 4: Validation of model-based compressed sensing approach for mapping PV-pyramidal connections. a, "Checkerboard" visualization of CAVIaR inferences. Left column shows demixed PSCs evoked by single-target stimulation of the listed ROI. Right column shows PSCs evoked by holographic stimulation of ensembles containing the listed neuron. Trials separated by dashed lines and rearranged to be in order of increasing laser power (from 30 mW to 60 mW). While all single-target stimulation trials are shown on the left, for ease of visualization only a fraction of ensemble stimulation trials are shown on the right. Shaded trials indicate when CAVIaR detects that the observed PSC was evoked by successfully spiking the listed (presynaptic) ROI; opacity of shading represents the estimated posterior uncertainty of the spike (lighter, less certain; darker, more certain). Blue stars represent when CAVIaR determines that the PSC is spontaneous (due to e.g. an uncharacteristic amplitude). Out of 169 stimulated ROIs, the same set of 7 were identified by CAVIaR as being presynaptic by both single-target and ensemble stimulation. **b**, Example PSCs (corresponding to neuron marked by white arrow in panel d, below) evoked by holographic stimulation of ensembles containing a putatively connected neuron (left panel). Single-target stimulation of the same neuron validates the existence of the individual synapse (right panel). Gray traces show raw PSCs, colored traces show demixed PSCs. Traces colored at random. **c**, Synaptic weights estimated by CAVIaR independently from single-target stimulation and ensemble stimulation trials show strong agreement (R^2, 0.99). **d**, Z-projection (over 5 different planes) of stimulation FOV for experiment corresponding to panels a-c. White circles show identified presynaptic neurons. Left, presynaptic neurons identified using single-target stimulation; right, presynaptic neurons identified using ensemble stimulation. NWD and CAVIaR were used in both cases. **e**, Additional example of similarity between synaptic connectivity maps obtained using single-target (left) vs ensemble stimulation (right) in combination with NWD and CAVIaR. **f**, Number of connections identified by CAVIaR using both single-target and ensemble stimulation across 14 experiments. Dashed gray line shows identity. **g**, Agreement between connectivity maps obtained using single-target stimulation compared to model-based compressed sensing. Agreement measured using both the R^2 (which takes synaptic strength into account; mean $\pm$ 1 s.d., 0.89 $\pm$ 0.19) and the Jaccard index (which measures agreement using only a binary classification; mean $\pm$ 1 s.d., 0.79 $\pm$ 0.15). **h**, Simultaneous whole-cell and cell-attached recordings provide ground-truth presynaptic spikes associated with postsynaptic responses during single-target and ensemble stimulation. **i**, Example image showing locations of presynaptic and postsynaptic neurons, among other segmented ROIs. **j**, Example scenarios from a paired recording associated with successful photoelicitation of presynaptic spikes. While trials (i) and (ii) resulted in correct predictions of presynaptic spiking by CAVIaR (i.e., are true positives), trials (iii) and (iv) were declared trials on which the stimulated neuron did not spike, either due to an unusually low-amplitude PSC or synaptic failure. Underlaid gray traces show raw PSC without NWD, black traces show PSC with NWD. Scales at bottom left apply to all subpanels. **k**, Performance of presynaptic spike inference over $n = 4$ paired recordings, separated by single-target vs ensemble stimulation and laser power. Performance of presynaptic spike inference does not depend on power or single-target vs ensemble stimulation ($p > 0.05$ for all pairs, dependent t-test).

2009, Shababo et al., 2013], we wanted to determine whether the spikes inferred by CAVIaR were allocated to the correct presynaptic neurons. We therefore performed multiple (n=4) paired recordings to obtain ground-truth spikes. We first used single-target stimulation to identify a likely presynaptic neuron in real time, then established a cell-attached patch clamp on the presynaptic neuron, and then randomly alternated between single-target and ensemble stimulation across multiple power levels while recording presynaptic spikes (Figure 4h,i). We confirmed that each of the neurons identified as connected using single-target stimulation following establishment of the patch clamp were also considered connected when the population was subsequently mapped using ensemble stimulation.

Paired recordings revealed a range of different scenarios associated with ground-truth spiking (Figure 4j). This included true positives, where CAVIaR correctly inferred a presynaptic spike from a given PSC evoked by single-target or ensemble stimulation (scenarios (i) and (ii) in Figure 4j). This also included false negatives, however. For example, in some cases CAVIaR incorrectly estimated that an unusually small PSC did not arise by a stimulation-evoked spike of the patched presynaptic neuron but rather from another, more weakly connected neuron that was stimulated at the same time (scenario (iii)). False negatives also occurred due to synaptic failure, where the presynaptic neuron successfully elicited, but did not transmit, a spike (scenario (iv)). Despite the latter two challenging scenarios, CAVIaR achieved high accuracy in its overall estimates of presynaptic spikes with no significant degradation in its ability to infer presynaptic spikes for individual neurons, even when stimulating ten neurons at once and over a range of laser powers (Figure 4k; all pair-wise differences not significant, p > 0.05, dependent t-test).

CAVIaR estimates converge rapidly and accurately predict postsynaptic responses to novel holographic stimulation patterns

To go beyond single-target confirmation of the presence and magnitude of the synapses identified using CAVIaR (Figure 4f,g), we next asked whether the inferred connectivity could be used to predict the postsynaptic response to holograms

targeting combinations of neurons that the model had never seen before. We also wanted to know whether ensemble stimulation was systematically engaging polysynaptic network effects. For example, PV neurons are known to provide high levels of mutual inhibition [Pfeffer et al., 2013], which could be triggered when stimulating many PV neurons at once. In this case, other PV neurons would be inhibited, potentially reducing the number of driven PV neurons and resulting in smaller inhibitory PSCs than would be expected by CAVIaR.

To simultaneously answer these questions, we performed an out-of-sample testing method that we refer to as leave-one-hologram-out cross-validation (LOHO-CV, Figure 5a). Assuming that H different holograms are used in the experiment (where each hologram targets a different set of neurons), LOHO-CV works by running the CAVIaR algorithm on data corresponding to $H-1$ holograms, such that the algorithm has never observed postsynaptic responses from simultaneous stimulation of the neurons targeted in the Hth hologram. Then, samples from the posterior distribution over the model parameters are used to predict the mean postsynaptic response to hologram H at three different power levels, a process that includes sampling over the model's uncertainty about whether spikes will be elicited by stimulation at any given laser power. The predicted response is compared against the held-out response, and the process continues with a different hologram selected to be held-out until the response to every hologram has been predicted.

LOHO-CV confirmed that CAVIaR achieved high accuracy in its predicted responses to novel holograms (Figure 5b; example R^2, 0.92). This indicated that CAVIaR did not miss synapses for which ensemble stimulation evoked large and reliable PSCs. Such synapses would manifest as large observed test responses, but with predicted responses of 0 (i.e., the test data would lie along the $y = 0$ axis in Figure 5b). Additionally, high LOHO-CV performance implied that CAVIaR did not make systematic errors arising due to polysynaptic network effects, which would manifest as clear off-diagonal points in Figure 5b. We performed a similar cross-validation analysis for CAVI-SnS and CoSaMP, which achieved substantially lower performance compared to CAVIaR (Figure 5b, middle and right panels). Repeating the analysis across all 14 experiments showed that these two previously developed techniques exhibited systematic errors in their ability to predict responses to unseen holographic stimuli (Figure 5c, d), indicating that they did not learn connectivity consistent with the underlying physiology.

We next wanted to determine how efficient our protocol was for discovering synaptic connectivity as a function of continuous stimulation time (namely, without regard to experiment time devoted to pausing stimulation to measure access resistance, adjusting the seal of the patch pipette, etc.). To do so, we subsampled mapping data from the total set of available trials and, for each subset, applied NWD and CAVIaR to estimate connectivity. This allowed us to monitor the rate of convergence of CAVIaR to its final estimates in increments of tens of seconds. Note that because our principal aim was validation of the connectivity inferences themselves rather than a demonstration of speed-up specifically, each hologram was presented multiple (3) times at random throughout the experiment. This experimental design provided a means to characterize performance via LOHO-CV analysis, but could only establish a lower bound on mapping speed because simulations demonstrate that optimal speeds are obtained only when unique holograms are used on every trial (Figure S8b). Nevertheless, we found that our mapping protocol converged to an R^2 of 0.9 with less than just 200 ms of stimulation time per neuron at 30 Hz (Figure 5e, left plot; obtained by normalizing mapping time by number of neurons in experiment). By comparison, single-target stimulation using CAVIaR and NWD reached an R^2 of 0.9 in 800 ms per neuron (though with much higher variance; Figure 5e, middle plot), and single-target stimulation using the existing approach of CAVI-SnS required 1.1 seconds per neuron (Figure 5e, right plot), more than five times slower than model-based compressed sensing and (in accordance with our simulation and LOHO-CV results; Figure 3c, Figure 5c) with substantially lower predictive accuracy.

Inference of synaptic circuitry from ensemble stimulation across multiple cortical cell types

Finally, we wanted to characterize how model-based compressed sensing performed across a variety of cortical cell-type combinations. We used Emx-Cre or SST-Cre lines to express either ChroME2s or ChroME2f respectively in pyramidal

Figure 5: Predictive performance and convergence rates using model-based compressed sensing outperform existing methods. a, A "leave-one-hologram-out" cross-validation approach can be used to determine the predictive performance of connectivity mapping techniques. **b**, LOHO-CV R^2 scores for CAVIaR, CAVI-SnS, and CoSaMP on an example mapping experiment, showing that CAVIaR obtains the highest accuracy. No method suggests recruitment of polysynaptic effects, which would manifest as systematic off-diagonal data points. For these comparisons, CoSaMP was provided with the number of connections identified by CAVIaR. **c**, Summary of LOHO-CV R^2 scores across 14 different PV-pyramidal mapping experiments. Dashed gray line denotes an R^2 of 0; scores below this line indicate poorer predictive performance than simply predicting the mean of the true responses without regard to which neurons were stimulated. **d**, Direct comparison of CAVIaR LOHO-CV R^2 scores to CAVI-SnS and CoSaMP, showing that CAVIaR outperforms both methods on each dataset. **e**, Convergence time for proposed model-based compressed sensing technique (ensemble stimulation with CAVIaR and NWD) compared to conventional single-target mapping (with and without NWD). Stimulation performed at 30 Hz. Ensemble stimulation applied to 10 targets at once. Convergence time reported per neuron due to varying population sizes (larger populations take longer to map). Red/gray lines show convergence times for individual experiments. Each point is obtained by evaluating the R^2 between the estimated connectivity at that time point and at the end of the experiment. Black line with markers shows median across experiments. **f**, Z-projected map of connection strength across 14 pooled experiments. Colored circles indicate position of identified presynaptic neurons, open gray circles indicate positions of all probed ROIs. Individual maps aligned to have postsynaptic neuron at location (0, 0). **g**, Synaptic strength as a function of distance from 14 pooled experiments.

neurons and somatostatin-expressing (SST) inhibitory interneurons. Then, in addition to our prior experiments mapping PV-pyramidal connections, we also mapped pyramidal-pyramidal (n=12 experiments), SST-pyramidal (n=8), and pyramidal-PV connections (n=5, Figure 6). In each case we used our previous experimental design, where we randomly interleaved single-target and ensemble stimulation trials, and used NWD and CAVIaR to infer connectivity independently from each trial type.

Our experiments mapping pyramidal-pyramidal connections involved probing hundreds of potential presynaptic candidates per cortical slice (Figure 6a,b; mean number of targets, 486; minimum, 214; maximum, 687). This experimental preparation posed a particular challenge because the high opsin expression density made it difficult to establish a whole-cell recording on a pyramidal neuron that was opsin-negative, and consequently we typically patched an opsin-positive postsynaptic neuron. When probing nearby presynaptic candidates we thus frequently encountered direct photocurrent artifacts in our whole-cell recordings that could affect our ability to discriminate synaptic connectivity (i.e. direct photocurrents could increase the false-positive rate in the immediate vicinity of the patch-electrode). However, because NWD is trained to isolate PSCs occurring within a specified window (typically 3-12 ms after stimulation), confounding photocurrents (which begin rising immediately upon stimulation) were suppressed by the demixing process (Figure S10). Following the application of CAVIaR to infer connectivity, there was (as expected) some variability between connectivity maps obtained using single-target stimulation and model-based compressed sensing, but the identity, strength, and overall number of identified connections tended to be conserved across experiments (Figure 6a, c-d; mean R^2 between single-target and model-based compressed sensing connectivity vectors, 0.36; mean Jaccard index, 0.61). Though the mean R^2 was lower than in the PV-E condition, this was largely due to an outlier experiment (Figure S11), and was not representative of the distribution at large (median R^2, 0.57).

Mapping pyramidal-PV connections involved probing similar numbers of potential presynaptic candidates as in our pyramidal-pyramidal maps due to a similarly high opsin expression density and intrinsic cell-count (Figure 6e, mean number of targets across 5 experiments, 602; minimum, 562; maximum, 661). However, because in this preparation opsin is expressed specifically in pyramidal neurons, when recording from PV neurons we faced no risk of direct photocurrent artifacts. Across 5 experiments, connectivity maps obtained using single-target stimulation and model-based compressed sensing achieved a mean R^2 of 0.52, and a mean Jaccard index of 0.55 (Figure 6f-h).

Finally, we considered whether using compressed sensing to map connections from SST neurons to pyramidal neurons could pose an additional challenge because SST-pyramidal connections are largely axo-dendritic, potentially causing significant filtering and attenuation of PSCs through the dendrites of pyramidal neurons [Cristo et al., 2004, Wang et al., 2004, Dorsett et al., 2021], though prior work has shown the feasibility of using single-target stimulation (via two-photon glutamate uncaging) to map connections of this type [Fino and Yuste, 2011]. Across 8 experiments we found that, while some variability persisted, the connectivity maps obtained using single-target stimulation and model-based compressed sensing were highly conserved, despite this more challenging experimental setting (Figure 6i-l; mean R^2, 0.72; mean Jaccard index, 0.66; mean number of targets, 129; minimum 51; maximum 175).

By performing a relatively small number of experiments (5-14) for each of the mapped populations, we acquired connectivity maps that corresponded to the probing of thousands of presynaptic candidates within the local volume. These maps recapitulated the increased connection probability in the proximity of the postsynaptic neuron ($\leq$100 μm around the cell) characteristic of connectivity mapping studies (Figure 6b,f,j), which typically require pooling together many more smaller-scale experiments [Baker et al., 2016, Seeman et al., 2018, Hage et al., 2022, Campagnola et al., 2022]. While a detailed analysis of the spatial properties of individual maps of connectivity is beyond the scope of this study, our results indicate the possibility of investigating map-to-map variability (Figure 4d,e; supp figs) [Nikolenko et al., 2007]. Further, our results show the feasibility and utility of compressive connectivity mapping across a diversity of cortical cell-type combinations, despite challenges arising due to synaptic physiology or experimental preparation (i.e., direct photocurrent artifacts).

Figure 6: Model-based compressed sensing of synaptic connectivity applied to multiple cortical cell type pairs. a, Comparison of pyramidal-pyramidal connectivity maps obtained on the same population of neurons using single-target stimulation and model-based compressed sensing (10-target ensembles, 30 Hz stimulation, connections identified using CAVIaR with NWD). Experiment performed over 5 planes, image shows z-projection. Agreement between synaptic weights estimated by CAVIaR independently from single-target stimulation and ensemble stimulation trials. **b**, Map of synaptic connections identified using ensemble stimulation across 12 pooled experiments. Dashed cyan circle represents region with 30 μm radius where synaptic connections are most uncertain due to potential photocurrent contamination in opsin-positive postsynaptic neurons (however see Figure S10). **c**, Number of connections identified by CAVIaR using both single-target and ensemble stimulation across 12 experiments. Dashed gray line shows identity. **d**, Agreement between connectivity maps obtained using single-target stimulation compared to model-based compressed sensing. Mean R^2, 0.36 $\pm$ 0.62; median R^2, 0.57; mean Jaccard index, 0.61 $\pm$ 0.2. Gray arrow indicates outlier (c.f. Figure S11). **e–h**, same as a–d, but for mapping pyramidal-PV connections over 5 experiments. Blue circle represents postsynaptic PV neuron. Mean R^2 in panel h, 0.52 $\pm$ 0.21; mean Jaccard index, 0.55 $\pm$ 0.22. **i–l**, Same as a–d, but for mapping SST-pyramidal connections over 5 experiments. Mean R^2 in panel l, 0.72 $\pm$ 0.27; mean Jaccard index, 0.63 $\pm$ 0.15.

Discussion

Uncovering how neural computations are implemented in the cortex requires knowledge of the existence and strength of the underlying synaptic connections. We therefore developed a pair of complementary computational tools (NWD and CAVIaR) that together enable high-speed acquisition of large-scale synaptic connectivity maps using two-photon holographic optogenetics. Other studies that develop compressive connectivity mapping techniques either use generic compressed sensing algorithms or have only been demonstrated in simplified simulations [Hu and Chklovskii, 2009, Shababo et al., 2013, Draelos and Pearson, 2020, Navarro and Oweiss, 2023, Chen et al., 2023]. Here, however, we show that considerable methodological extensions over these previous algorithms are required to ensure that connectivity is accurately determined in light of the substantial sources of biophysical variability that are intrinsic to real holographic stimulation experiments. We found that use of NWD and CAVIaR provided a mapping speed-up of over an order of magnitude compared to existing approaches in realistic settings, granting the speed and scalability necessary to collect large-scale maps of connectivity within individual experimental sessions.

To implement and validate these tools in real mapping experiments we used cutting-edge optogenetic technologies that offered several key advantages. Namely, since the ChroME2f opsin is potent with fast off-kinetics, we could titrate the pulse width and range of stimulation laser powers such that with high probability stimulation generated either 0 or 1 presynaptic spike(s), with minimal instances of more than one spike (Figure S1), which facilitated tractable presynaptic spike inference. By comparison, potent opsins with slow off-kinetics (such as ChRmine [Marshel et al., 2019]) elicit larger and less predictable presynaptic spike counts [Sridharan et al., 2022] that make connectivity inference more computationally challenging as the combinatorial space of how presynaptic spikes can be allocated across the stimulated population grows rapidly.

We also relied on the fact that ChroME2f expression is targeted to the soma through a fusion with the Kv2.1 tag [Mardinly et al., 2018, Sridharan et al., 2022], reducing the likelihood of triggering spikes in untargeted neurons by off-target stimulation of nearby neurites [Baker et al., 2016, Shemesh et al., 2017]. Moreover, we used two-photon holographic light sculpting to illuminate entire somas at once [Papagiakoumou et al., 2010, Hernandez et al., 2016, Pégard et al., 2017] and randomly switched the stimulation laser between three or more powers, effectively probing for presynaptic neurons at multiple excitation levels. Stimulating with low power further reduces the risk of off-target stimulation, but may fail to elicit spikes in neurons with low opsin expression or high rheobases. Notably, it cannot in general be known ahead of time whether a lack of optogenetically evoked PSCs was due to the stimulated neuron not being connected or because of a failure to photoelicit spikes, an ambiguity that can only be resolved through high power stimulation. On the other hand, stimulating with high power increases the risk of off-target stimulation but elicits spikes with high probability. We therefore elected to use a randomized design that probed neurons at multiple power levels spanning a range of photosensitivies, ultimately allowing us to map multiple cortical cell-types with high precision using CAVIaR.

One potential concern with our proposed connectivity mapping approach is that it may engage mechanisms for short-term plasticity (i.e., synaptic facilitation or depression) given our suggested high speed of stimulation and the fact that entire ensembles of neurons are stimulated at once. To help calibrate mapping protocols, we derived a simple combinatorial expression for the expected interstimulus interval (ISI, in seconds) for any given neuron (Methods). Assuming connectivity from N total neurons is mapped by stimulating a random ensemble of size R on each trial at a speed of f Hz, the mean ISI is $\frac{N}{Rf}$. Thus, a typical experiment mapping ∼300 neurons with 10-target stimulation at 30 Hz yields an expected stimulation return time of 1 second. An additional concern associated with ensemble stimulation is the recruitment of potential polysynaptic network effects arising due to recurrent connectivity in the cortex [Oldenburg et al., 2022]. However, in slice experiments we do not expect this to be a major factor influencing the quality of our inferences, and we confirmed this using LOHO-CV (Figure 5b,c), though the impact of polysynaptic effects likely depends on brain region and number of simultaneously illuminated targets.

We demonstrated the applicability of NWD and CAVIaR by mapping connectivity between three different cell-types (PV, SST, and pyramidal neurons), using two ChroME variants (ChroME2f and ChroME2s), and using both targeted and blind experimental designs. In every case, application of our tools led to a substantial improvement in signal quality and speed of connectivity mapping. While in our experiments we typically mapped connectivity from several hundred neurons, this was only limited by factors outside of the influence of NWD and CAVIaR, such as the number of neurons targetable by our SLM or the rate at which phase masks could be computed. Our simulations predict that the scale and throughput of compressive connectivity mapping will only grow with the adoption of experimental steps to increase the targetable field of view (e.g., by translating the microscope or SLM stimulation field), with larger commercially available SLMs, with the use of holographic mesoscopes [Abdeladim et al., 2023], or with even more potent opsins enabling stimulation of more targets for the same laser power. Such advances would allow for precise, low-latency holographic control over larger neural ensembles or for mapping larger population sizes than those used here.

Our results indicate that NWD and CAVIaR can substantially reduce the cumulative time spent stimulating tissue. However, the speed of a mapping experiment can also be impacted by other factors, including the time required to identify and segment neurons, monitor the access resistance of the patch clamp, and compute hologram phase masks. Each of these factors similarly affects mapping performed using single-target stimulation and ensemble stimulation. While acquiring an image stack to identify presynaptic candidates is required for all experiments using targeted stimulation, measuring access resistance poses an additional trade-off between the speed of a mapping experiment and the desire to make regular quality checks. An ideal experimental design would therefore proceed with the longest periods of uninterrupted stimulation possible to yield the maximal benefits of NWD and CAVIaR, but with brief ($\sim$200 ms) automatic pauses included to check access resistance. The attending experimenter could then be prompted to address inadequate access only when necessary. In the case of phase mask generation, efficient computation is critical since otherwise experiment time could be dominated by this step. We expect that high-speed phase mask estimation methods, such as those based on deep learning [Eybposh et al., 2020, Eybposh et al., 2022], could play an increasingly important role in rapid connectivity mapping.

Combining NWD and CAVIaR with two-photon holographic optogenetics should enable new experiments where, with just a few minutes of rapid ensemble stimulation, one can obtain an extensive map of connectivity. This could allow for very high throughput collection and screening of synaptic connectivity maps from brain slices, or for future *in vivo* applications [Chen et al., 2023] where anatomical connectivity can be directly related to functional activity. Critically, *in vivo* experiments are impacted by higher background rates of spontaneous activity and more prevalent polysynaptic effects. While further validation must be performed to ensure accuracy *in vivo*, the impact of each of these impediments is demonstrably reduced by our computational methods, implying that NWD and CAVIaR may be important tools for precise and scalable mapping in this experimental regime.

Acknowledgements

The authors thank Kenneth Kay, Darcy Peterka, Ben Shababo, and Shizhe Chen for many insightful discussions and helpful suggestions. We thank Uday Jagadisan for technical assistance related to the optical setup and Karthika Gopakumar for helping with animal husbandry and the opsin expression procedure. This work was funded by NIH awards 1RF1MH120680 and 1U19NS107613-01 to HA and LP. MAT, BA, and LP were supported by the Gatsby Charitable Foundation and NSF NeuroNex award 1707398.

Author contributions

MAT, MG, HA, and LP conceived of the project. MG designed and performed all experiments, with assistance from MS during the initial project phase. MAT developed the computational methods together with LP, with contributions from BA.

MAT performed all simulations. MAT and MG performed all analyses. MAT and MG wrote the manuscript with input from all authors.

Code availability

CAVIaR is implemented in Python using JAX [Bradbury et al., 2018], can be run on GPU, and is freely available at https://github.com/marcustriplett/circuitmap together with a PyTorch Lightning implementation of NWD.

References

[Abdeladim et al., 2023] Abdeladim, L., Shin, H., Jagadisan, U. K., Ogando, M. B., and Adesnik, H. (2023). Probing inter-areal computations with a cellular resolution two-photon holographic mesoscope. *bioRxiv*, pages 2023–03.

[Adesnik and Abdeladim, 2021] Adesnik, H. and Abdeladim, L. (2021). Probing neural codes with two-photon holographic optogenetics. *Nature Neuroscience*, 24(10):1356–1366.

[Ayer et al., 1955] Ayer, M., Brunk, H. D., Ewing, G. M., Reid, W. T., and Silverman, E. (1955). An empirical distribution function for sampling with incomplete information. *The annals of mathematical statistics*, pages 641–647.

[Baker et al., 2016] Baker, C. A., Elyada, Y. M., Parra, A., and Bolton, M. M. (2016). Cellular resolution circuit mapping with temporal-focused excitation of soma-targeted channelrhodopsin. *Elife*, 5:e14193.

[Bishop, 2006] Bishop, C. M. (2006). *Pattern Recognition and Machine Learning*, volume 4. Springer.

[Blei et al., 2017] Blei, D. M., Kucukelbir, A., and McAuliffe, J. D. (2017). Variational inference: A review for statisticians. *Journal of the American statistical Association*, 112(518):859–877.

[Bounds et al., 2021] Bounds, H. A., Sadahiro, M., Hendricks, W. D., Gajowa, M., Oldenburg, I. A., Gopakumar, K., Quintana, D., Daigle, T., Zeng, H., and Adesnik, H. (2021). Multifunctional cre-dependent transgenic mice for high-precision all-optical interrogation of neural circuits. *bioRxiv*.

[Bradbury et al., 2018] Bradbury, J., Frostig, R., Hawkins, P., Johnson, M. J., Leary, C., Maclaurin, D., Necula, G., Paszke, A., VanderPlas, J., Wanderman-Milne, S., and Zhang, Q. (2018). JAX: composable transformations of Python+NumPy programs.

[Campagnola et al., 2022] Campagnola, L., Seeman, S. C., Chartrand, T., Kim, L., Hoggarth, A., Gamlin, C., Ito, S., Trinh, J., Davoudian, P., Radaelli, C., et al. (2022). Local connectivity and synaptic dynamics in mouse and human neocortex. *Science*, 375(6585):eabj5861.

[Candes et al., 2006] Candes, E. J., Romberg, J. K., and Tao, T. (2006). Stable signal recovery from incomplete and inaccurate measurements. *Communications on Pure and Applied Mathematics: A Journal Issued by the Courant Institute of Mathematical Sciences*, 59(8):1207–1223.

[Candes and Tao, 2006] Candes, E. J. and Tao, T. (2006). Near-optimal signal recovery from random projections: Universal encoding strategies? *IEEE transactions on information theory*, 52(12):5406–5425.

[Chen et al., 2023] Chen, I.-W., Chan, C. Y., Navarro, P., de Sars, V., Ronzitti, E., Oweiss, K., Tanese, D., and Emiliani, V. (2023). High-throughput in vivo synaptic connectivity mapping of neuronal micro-circuits using two-photon holographic optogenetics and compressive sensing. *bioRxiv*, pages 2023–09.

[Cristo et al., 2004] Cristo, G. D., Wu, C., Chattopadhyaya, B., Ango, F., Knott, G., Welker, E., Svoboda, K., and Huang, Z. J. (2004). Subcellular domain-restricted gabaergic innervation in primary visual cortex in the absence of sensory and thalamic inputs. *Nature neuroscience*, 7(11):1184–1186.

[Donoho, 2006] Donoho, D. L. (2006). Compressed sensing. *IEEE Transactions on information theory*, 52(4):1289–1306.

[Dorsett et al., 2021] Dorsett, C., Philpot, B. D., Smith, S. L., and Smith, I. T. (2021). The impact of sst and pv interneurons on nonlinear synaptic integration in the neocortex. *Eneuro*, 8(5).

[Draelos and Pearson, 2020] Draelos, A. and Pearson, J. (2020). Online neural connectivity estimation with noisy group testing. *Advances in Neural Information Processing Systems*, 33.

[Eybposh et al., 2020] Eybposh, M. H., Caira, N. W., Atisa, M., Chakravarthula, P., and Pégard, N. C. (2020). Deepcgh: 3d computer-generated holography using deep learning. *Optics Express*, 28(18):26636–26650.

[Eybposh et al., 2022] Eybposh, M. H., Curtis, V. R., Rodríguez-Romaguera, J., and Pégard, N. C. (2022). Advances in computer-generated holography for targeted neuronal modulation. *Neurophotonics*, 9(4):041409.

[Falk et al., 2019] Falk, T., Mai, D., Bensch, R., Çiçek, Ö., Abdulkadir, A., Marrakchi, Y., Böhm, A., Deubner, J., Jäckel, Z., Seiwald, K., et al. (2019). U-net: deep learning for cell counting, detection, and morphometry. *Nature methods*, 16(1):67–70.

[Fino and Yuste, 2011] Fino, E. and Yuste, R. (2011). Dense inhibitory connectivity in neocortex. *Neuron*, 69(6):1188–1203.

[Fletcher et al., 2011] Fletcher, A. K., Rangan, S., Varshney, L. R., and Bhargava, A. (2011). Neural reconstruction with approximate message passing (neuramp). In *NIPS*, pages 2555–2563.

[Gerchberg, 1972] Gerchberg, R. W. (1972). A practical algorithm for the determination of plane from image and diffraction pictures. *Optik*, 35(2):237–246.

[Hage et al., 2022] Hage, T. A., Bosma-Moody, A., Baker, C. A., Kratz, M. B., Campagnola, L., Jarsky, T., Zeng, H., and Murphy, G. J. (2022). Synaptic connectivity to l2/3 of primary visual cortex measured by two-photon optogenetic stimulation. *Elife*, 11:e71103.

[Hernandez et al., 2016] Hernandez, O., Papagiakoumou, E., Tanese, D., Fidelin, K., Wyart, C., and Emiliani, V. (2016). Three-dimensional spatiotemporal focusing of holographic patterns. *Nature communications*, 7(1):1–11.

[Hu and Chklovskii, 2009] Hu, T. and Chklovskii, D. (2009). Reconstruction of sparse circuits using multi-neuronal excitation (rescume). *Advances in Neural Information Processing Systems*, 22:790–798.

[Lee et al., 2020] Lee, J., Mitelut, C., Shokri, H., Kinsella, I., Dethe, N., Wu, S., Li, K., Reyes, E. B., Turcu, D., Batty, E., et al. (2020). Yass: Yet another spike sorter applied to large-scale multi-electrode array recordings in primate retina. *bioRxiv*.

[Mardinly et al., 2018] Mardinly, A. R., Oldenburg, I. A., Pégard, N. C., Sridharan, S., Lyall, E. H., Chesnov, K., Brohawn, S. G., Waller, L., and Adesnik, H. (2018). Precise multimodal optical control of neural ensemble activity. *Nature neuroscience*, 21(6):881–893.

[Marshel et al., 2019] Marshel, J. H., Kim, Y. S., Machado, T. A., Quirin, S., Benson, B., Kadmon, J., Raja, C., Chibukhchyan, A., Ramakrishnan, C., Inoue, M., et al. (2019). Cortical layer–specific critical dynamics triggering perception. *Science*, 365(6453):eaaw5202.

[Merel et al., 2016] Merel, J., Shababo, B., Naka, A., Adesnik, H., and Paninski, L. (2016). Bayesian methods for event analysis of intracellular currents. *Journal of Neuroscience Methods*, 269:21–32.

[Mishchenko and Paninski, 2012] Mishchenko, Y. and Paninski, L. (2012). A bayesian compressed-sensing approach for reconstructing neural connectivity from subsampled anatomical data. *Journal of computational neuroscience*, 33(2):371–388.

[Naka et al., 2019] Naka, A., Veit, J., Shababo, B., Chance, R. K., Risso, D., Stafford, D., Snyder, B., Egladyous, A., Chu, D., Sridharan, S., et al. (2019). Complementary networks of cortical somatostatin interneurons enforce layer specific control. *Elife*, 8:e43696.

[Navarro and Oweiss, 2023] Navarro, P. and Oweiss, K. (2023). Compressive sensing of functional connectivity maps from patterned optogenetic stimulation of neuronal ensembles. *Patterns*.

[Needell and Tropp, 2009] Needell, D. and Tropp, J. A. (2009). Cosamp: Iterative signal recovery from incomplete and inaccurate samples. *Applied and computational harmonic analysis*, 26(3):301–321.

[Nikolenko et al., 2007] Nikolenko, V., Poskanzer, K. E., and Yuste, R. (2007). Two-photon photostimulation and imaging of neural circuits. *Nature methods*, 4(11):943–950.

[Oldenburg et al., 2022] Oldenburg, I. A., Hendricks, W. D., Handy, G., Shamardani, K., Bounds, H. A., Doiron, B., and Adesnik, H. (2022). The logic of recurrent circuits in the primary visual cortex. *bioRxiv*, pages 2022–09.

[Packer et al., 2012] Packer, A. M., Peterka, D. S., Hirtz, J. J., Prakash, R., Deisseroth, K., and Yuste, R. (2012). Two-photon optogenetics of dendritic spines and neural circuits. *Nature methods*, 9(12):1202–1205.

[Packer and Yuste, 2011] Packer, A. M. and Yuste, R. (2011). Dense, unspecific connectivity of neocortical parvalbumin-positive interneurons: a canonical microcircuit for inhibition? *Journal of Neuroscience*, 31(37):13260–13271.

[Pakman and Paninski, 2014] Pakman, A. and Paninski, L. (2014). Exact hamiltonian monte carlo for truncated multi-variate gaussians. *Journal of Computational and Graphical Statistics*, 23(2):518–542.

[Papagiakoumou et al., 2010] Papagiakoumou, E., Anselmi, F., Bègue, A., De Sars, V., Glückstad, J., Isacoff, E. Y., and Emiliani, V. (2010). Scanless two-photon excitation of channelrhodopsin-2. *Nature methods*, 7(10):848–854.

[Papagiakoumou et al., 2020] Papagiakoumou, E., Ronzitti, E., and Emiliani, V. (2020). Scanless two-photon excitation with temporal focusing. *Nature Methods*, 17(6):571–581.

[Pégard et al., 2017] Pégard, N. C., Mardinly, A. R., Oldenburg, I. A., Sridharan, S., Waller, L., and Adesnik, H. (2017). Three-dimensional scanless holographic optogenetics with temporal focusing (3d-shot). *Nature communications*, 8(1):1–14.

[Pfeffer et al., 2013] Pfeffer, C. K., Xue, M., He, M., Huang, Z. J., and Scanziani, M. (2013). Inhibition of inhibition in visual cortex: the logic of connections between molecularly distinct interneurons. *Nature neuroscience*, 16(8):1068–1076.

[Printz et al., 2023] Printz, Y., Patil, P., Mahn, M., Benjamin, A., Litvin, A., Levy, R., Bringmann, M., and Yizhar, O. (2023). Determinants of functional synaptic connectivity among amygdala-projecting prefrontal cortical neurons in male mice. *Nature communications*, 14(1):1667.

[Rani et al., 2018] Rani, M., Dhok, S. B., and Deshmukh, R. B. (2018). A systematic review of compressive sensing: Concepts, implementations and applications. *IEEE access*, 6:4875–4894.

[Rickgauer and Tank, 2009] Rickgauer, J. P. and Tank, D. W. (2009). Two-photon excitation of channelrhodopsin-2 at saturation. *Proceedings of the National Academy of Sciences*, 106(35):15025–15030.

[Ronneberger et al., 2015] Ronneberger, O., Fischer, P., and Brox, T. (2015). U-net: Convolutional networks for biomedical image segmentation. In *International Conference on Medical image computing and computer-assisted intervention*, pages 234–241. Springer.

[Seeman et al., 2018] Seeman, S. C., Campagnola, L., Davoudian, P. A., Hoggarth, A., Hage, T. A., Bosma-Moody, A., Baker, C. A., Lee, J. H., Mihalas, S., Teeter, C., et al. (2018). Sparse recurrent excitatory connectivity in the microcircuit of the adult mouse and human cortex. *Elife*, 7:e37349.

[Shababo et al., 2013] Shababo, B., Paige, B., Pakman, A., and Paninski, L. (2013). Bayesian inference and online experimental design for mapping neural microcircuits. *Advances in Neural Information Processing Systems*, 26:1304–1312.

[Shemesh et al., 2017] Shemesh, O. A., Tanese, D., Zampini, V., Linghu, C., Piatkevich, K., Ronzitti, E., Papagiakoumou, E., Boyden, E. S., and Emiliani, V. (2017). Temporally precise single-cell-resolution optogenetics. *Nature neuroscience*, 20(12):1796–1806.

[Sridharan et al., 2022] Sridharan, S., Gajowa, M. A., Ogando, M. B., Jagadisan, U. K., Abdeladim, L., Sadahiro, M., Bounds, H. A., Hendricks, W. D., Turney, T. S., Tayler, I., et al. (2022). High-performance microbial opsins for spatially and temporally precise perturbations of large neuronal networks. *Neuron*.

[Wang et al., 2007] Wang, H., Peca, J., Matsuzaki, M., Matsuzaki, K., Noguchi, J., Qiu, L., Wang, D., Zhang, F., Boyden, E., Deisseroth, K., et al. (2007). High-speed mapping of synaptic connectivity using photostimulation in channelrhodopsin-2 transgenic mice. *Proceedings of the National Academy of Sciences*, 104(19):8143–8148.

[Wang et al., 2004] Wang, Y., Toledo-Rodriguez, M., Gupta, A., Wu, C., Silberberg, G., Luo, J., and Markram, H. (2004). Anatomical, physiological and molecular properties of martinotti cells in the somatosensory cortex of the juvenile rat. *The Journal of physiology*, 561(1):65–90.

Methods

Experimental methods

Animals

All experiments on animals were conducted with approval of the Animal Care and Use Committee of the University of California, Berkeley. In all experiments we attempted to use male and female mice equally. Mice used for experiments in this study were transgenic Emx-Cre, PV-Cre, or SST-Cre mice obtained by crossing the corresponding lines in-house with a wild type (CD-1 (ICR) white strain, obtained from Charles River). Mice were housed in cohorts of five or fewer in a reverse light:dark cycle of 12:12 hours, with experiments occurring during the dark phase.

Opsin expression method

We used a neonatal injection procedure to induce expression of ChroME2s (AAV9.CAG.DIO.ChroME-ST.P2A.H2B-mR uby6) in the visual cortex of Emx-Cre animals or ChroME2f (AAV9.CAG.DIO.Chrome2f.P2A.H2B-mRuby3.WPRE.SV43) in PV-Cre and SST-Cre animals. Both constructs expressed the mRuby3 fluorophore in opsin-positive cell nuclei. Young pups at P3 or P4 were anaesthetized by placing them on ice for approximately 3 minutes. Next, each animal was stabilized under the nanoliter-injector (WPI) and a small portion (30 nL/injection) of virus was injected directly in 3-5 places around V1 via the skin and skull and at 3-5 depths to target L2/3. After the procedure the animal was placed on a

heating pad until it recovered. At the end of the procedure the injected litter was returned to their cage with their parents and housed together until reaching approximately 21 days of age.

3D-SHOT holography setup

All experiments were performed using the 3D-SHOT multiphoton holography setup (see [Mardinly et al., 2018] for details). Briefly, the setup was custom built around a commercial Sutter MOM (movable object microscope) platform (Sutter Instruments) and combined a 3D photostimulation path, a fast resonant-galvo two-photon raster scanning imaging path and a wide-field one-photon epifluorescence/IR (infrared) transmitted light imaging path. The stimulation and imaging beams were merged together using a polarizing beamsplitter placed before the microscope tube lens.

A femtosecond fiber laser was used for two-photon photostimulation (Monaco 1035-80-60; 1040 nm, 1 MHz, 300 fs, Coherent). The stimulation laser was directed onto a blazed diffraction grating (600 l/mm, 1000 nm blaze, Edmund Optics 49-570) for temporal focusing. In order to be able to utilize the total available laser power (60 W laser output), the beam was enlarged by a factor of 2.5 to prevent heat damage of the grating surface. The spot on the grating was relayed onto a rotating diffuser where it formed a temporally focused spot. The rotating diffuser was used to both randomize the phase pattern imprinted on the temporally focused spot and to expand the beam in the direction orthogonal to the temporal focusing direction and fully fill the spatial light modulator (HSP 1920 1920x1152 pixels, Meadowlark Optics). The SLM plane was relayed through 4f systems to the back aperture of an Olympus 20x water immersion objective, resulting in a custom 3D distribution of temporally-focused spots at the focus of the objective. Holographic phase masks were calculated using the iterative, in-house written and GPU-optimized Gerchberg-Saxton algorithm [Gerchberg, 1972] and the intensity distribution was corrected to accommodate for diffraction efficiencies.

The two-photon imaging path relied on a Ti:sapphire laser, Mai Tai (Spectra Physics), with external power control via Pockels cell (Conoptics, Inc). For fast raster scanning, the system was equipped with conjugated 8 kHz resonant galvo-galvo systems. The imaging path hardware was controlled by ScanImage software and custom Matlab code was used to control the spatial light modulator for targeted photostimulation and synchronize with imaging.

Epifluorescent one-photon excitation was via a Spectra X (Lumencor) light source filtered by an appropriate excitation filter set. For slice transillumination we used a 750 nm and IR diffuser. The image was collected using an Olympus 20x magnification water-immersion objective and a CCD camera and displayed on a screen enabling targeted patch clamping.

Slice electrophysiology

In vitro slice recordings were performed on 300 μm-thick coronal slices coming from 4–6-week-old animals expressing opsin in L2/3 of V1. During slicing the level of opsin expression was checked using a simple laser light to visualize the targeting of the opsin to V1 and the general brightness of the mRuby3 nuclear marker.

Whole-cell patch-clamp protocols were performed in ACSF (artificial cerebrospinal fluid) perfusion solution (in mM: NaCL 119, NaHCO3 26, Glucose 20, KCl 2.5, CaCl 2.5, MgSO4 1.3, NaH2PO4 1.3) in temperature-controlled (33°C) conditions. Patch pipettes (4-7 MΩ) were pulled from borosilicate glass filaments (Sutter Instruments) and filled with a Cesium (Cs2+)-based internal solution (in mM: 135 CeMeSO4, 3 NaCl, 10 HEPES, 0.3 EGTA, 4 Mg-ATP, 0.3 Na-GTP, 1 QX-314, 5 TEA-Cl, 295 mOsm, pH=7.45) also containing 50 μM Alexa Fluor hydrazide 488 or 594 dye (ThermoFisher Scientific). For loose-patch recordings the pipettes were filled with standard ACSF. Data was recorded at 20 kHz using a 700b Multiclamp Axon Amplifier (Molecular Devices). The headstage with the electrode holder (G23 Instruments) was controlled by a motorized micromanipulator (MP285A, Sutter Instruments). All data was acquired and analyzed with custom code written in Matlab using the National Instruments Data Acquisition Toolbox.

Learning physiological point spread functions

An opsin-positive cell was loose-patched at various slice depths between 20-90 μm and a volume of tissue surrounding that cell was then probed using a dense grid of holograms. The total size of the grid was 65x65x75 μm, resulting in a 10x10x7 voxel grid. The interval between the centers of the hologram targets were 5 pixels (6.5 μm) in the x/y dimensions and 12.5 μm in the z dimension. The phase masks of the grid holograms were pre-computed and stored in memory before performing the measurement. Two versions of the physiological point spread function experiments were performed. One where each hologram was a single spot randomly selected from the grid (n=8 experiments), and another where each hologram contained 10 spots: 1 spot randomly taken from the grid and 9 spots (either fixed or randomly placed) outside the grid (n=7 experiments).

Evoked spiking in response to 3-5 ms laser stimulation (30 Hz) across powers between 10-60 mW/spot were collected and analysed using a custom written script in Matlab. Each trial was randomly repeated 5-7 times. Evoked spiking was calculated by simple thresholding and the mean spiking probability across each grid point and power was calculated for single and 10-spot ensemble experiments. A Gaussian fit to the per-cell normalized mean spiking probability data was applied to estimate the FWHM of the spiking physiological point spread function.

Whole-cell targeted mapping experimental protocol

First, a L2/3 opsin-negative cell of pyramidal shape was sealed on (below 40 μm of cortical depth, opsin absence judged by fluorophore presence). The tissue was then quickly imaged with approximately 40 frames at 4-5 planes spaced by 25 μm. The cell was always positioned to be in the second plane from the top within the collected stack. The position of the presynaptic candidates were automatically identified by an in-house algorithm detecting round shapes in a specified FOV due to the presence of mRuby3 fluorophore in the opsin-positive cell nuclei.

Next, we computed 20-25 different sets of holograms, where every set of holograms was designed so that every target was stimulated once. Each hologram either targeted an ensemble of presynaptic candidates, or targeted one cell at a time. Further, each set of holograms was organized into sweeps, where each sweep was composed of a baseline period, followed by a single set of holographic stimulation trials, and ended with a short baseline period. A single set of ensemble holograms was obtained by randomly partitioning all targets into 10-target ensembles (without repeating any targets). Additional sets of holograms were obtained by repeatedly performing this random partitioning.

For example, an experiment probing 100 presynaptic candidates would contain data from 45 sweeps of 100 single-target holograms across 3 powers (15 repetitions per power), and 20 sweeps representing 20 sets of 10-target holograms. Each such set of 10-target holograms would be comprised of 100 targets split into 10 different 10-target holograms.This would ultimately result in 200 unique ensemble holograms organized in 20 sweeps. Each sweep would be performed across 3 powers and repeated 3 times per power (leading to 180 ensemble sweeps). In total, this would result in 225 sweeps (45 single-target sweeps and 180 ensemble sweeps). Importantly, the order of all holograms and powers within each repetition and sweep was randomized.

Stable access to the cell was obtained during hologram computation. The cell properties (Rm and Rs) were checked and the response to one-photon photocurrent was characterized. Following the completion of hologram computation, presynaptic candidates were mapped using randomly interleaved experimental trials containing single-target holograms or samples from the set of 10-target holograms. Each single-target hologram was repeated repeated 7-15 times, while ensemble holograms were repeated up to 3 times. All stimulations were performed across 3-4 laser powers in both mapping regimes.

The Rm and Rs parameters were controlled and frequently logged during the mapping protocol. Every 10 trials (or 20-180 seconds, depending on the number of presynaptic targets and the power condition), a series of hyperpolarizing steps was applied and the Rm and Rs logged. During offline analysis, trials when Rs significantly changed or were above 30 MOhm were excluded from the data. Experiments where zero or one connection(s) were identified were also excluded

from analysis.

For pyramidal-pyramidal mapping, the postsynaptic neuron was targeted based on its characteristic pyramidal shape and the absence of an mRuby3 signal in the cell. In an area with high opsin expression it was difficult to identify a completely opsin-negative cell. Thus, for a large portion of putatively opsin-negative cells, stimulating near the electrode still elicited a small direct photocurrent artifact (with a typical amplitude of 50-300 pA). For mapping pyramidal-PV connections, the postsynaptic cell type was confirmed to be PV by observing a fast-spiking response type as a result of a typical current step protocol.

Power calibration for single and ensemble target

To ensure that the connectivity maps obtained using single-target and ensemble stimulation were comparable, we carefully characterized the power needed to evoke similar photocurrent amplitudes at each presynaptic candidate in both stimulation regimes. To do this, a set of whole-cell experiments was performed in several cell types (pyramidal, PV, and SST). In each such experiment, an opsin-positive cell was patched and a direct opsin-photocurrent was recorded while all other experimental parameters were kept the same as in the connectivity mapping experiments. We used a range of powers that were typical for our connectivity mapping protocols. The requested laser power was a multiple of the diffraction efficiency of a given hologram (either single-target or ensemble) and power per cell multiplied by a number of targets in a hologram for a given condition.

In these experiments (Figure S2), we compared the opsin-photocurrent amplitude evoked in response to stimulation of a single target (the patched opsin-positive cell) with the opsin-photocurrent amplitudes evoked by ensemble holograms containing the patched opsin-positive cell. This set of experiments provided a measurement of the difference in the power delivered to the patched-cell between these two stimulation regimes. Correcting this difference required ∼35% additional power to the ensemble holograms (Figure S2b-d). Applying the correction compensated for the necessary photocurrent and was implemented in all mapping experiments (Figure S2e-g).

Pair-patch experimental protocol

The pair-patch mapping protocol was similar to the single-patch protocol, but contained an additional "pre-mapping" step. After establishing stable access to the opsin-negative pyramidal cell, all presynaptic candidates were probed with 10 trials of single-power, single-target holograms. A fast online analysis based on the z-score provided the coordinates of cells that responded to this simple connection screening procedure. One such identified cell would be approached and loose-patched with a second electrode while monitoring the stability of the postsynaptic recording. The main prerequisite for deciding which cell would be loose-patched (from a handful of putatively connected cells obtained via our simple screening procedure) was the ability to identify the cell in the slice under IR illumination and its accessibility. While the loose patch was established, the surrounding area would be re-imaged to update the positions of the presynaptic candidates. Mapping would then be performed according to the above-described procedure, while recording both postsynaptic responses and presynaptic spiking.

Whole-cell grid mapping experimental protocol

Opsin-negative cells (or those expressing minimal amounts of opsin) were whole-cell patched at various slice depths between 30-90 μm. Cells were voltage-clamped at a holding potential of -70 mV. After establishing stable access to a cell, we acquired its response to one-photon pulses of increasing duration (1, 3, and 5 ms pulses at 10 Hz). A volume of tissue surrounding that cell would then be probed using a dense grid of holograms across 3-4 powers. The total size of the grid was ∼160x160x100 μm, resulting in a 25x25x4 voxel grid. The distance between the centers of the hologram targets was 5 pixels (6.5 μm) in the x/y dimensions and 25 μm in the z dimension. The phase masks of the grid holograms

were pre-computed and stored in memory before performing the measurement. Recording parameters were frequently checked and logged by pausing the mapping protocol and probing the cell with a series of hyperpolarizing pulses.

Computational methods

Neural waveform demixing network architecture and training

The NWD network is a sequential U-Net [Ronneberger et al., 2015] that forms compressed representations of 45 ms single-trial PSC traces via a "contraction path," and generates a PSC waveform at the original trial length but with confounding synaptic currents removed via an "expansion path." Each block in the contraction path consists of a 2x temporal decimation step, a 1d convolution, a batch-norm step, and a rectified linear activation function. Each block in the expansion path consists of a "transposed" 1d convolution (also known as a fractionally strided convolution, or as a pseudo deconvolution), a batch-norm step, a rectified linear activation function, and a 2x linear interpolation step. There are four contraction blocks and four expansion blocks, with the outputs of the contraction blocks submitted to the expansion blocks at the corresponding temporal resolution through skip connections as in the original U-Net architecture.

The NWD networks used in this study were trained using 50,000 simulated PSC traces. The simulated PSCs obeyed the following generative structure. First, a template PSC $\varrho(\,\cdot\,;\tau_r,\tau_d,\Delta)$ with parameters τ_r,τ_d,Δ took the form

$$\varrho(t;\tau_r,\tau_d,\Delta) = \left[\exp\left(\frac{t-\Delta}{\tau_d}\right) - \exp\left(\frac{t-\Delta}{\tau_r}\right)\right]\mathbb{1}_{[t\geq\Delta]}. \tag{1}$$

Every simulated PSC $c(t)$ was the sum of a random number of templates, accounting for the stimulation of multiple connected presynaptic cells,

$$c(t) = \sum_{j=1}^{J}\varrho(t;\tau_r^j,\tau_d^j,\Delta^j), \tag{2}$$

$$\tau_X^j \sim \mathsf{Uniform}(\tau_X^{\mathsf{min}},\tau_X^{\mathsf{max}}), \tag{3}$$

$$\Delta^j \sim \mathsf{Uniform}(\Delta^{\mathsf{min}},\Delta^{\mathsf{max}}), \tag{4}$$

$$J \sim p(J), \tag{5}$$

where $X \in \{r,\mathsf{diff}\}$ with $\tau_d = \tau_r + \tau_{\mathsf{diff}}$ to ensure $\tau_d > \tau_r$, and where $p(J)$ determined the probability of selecting $J \in \mathbb{N}$ (including the possibility of $J = 0$). The PSC c_k for training example k was then given as a function of time t by

$$c_k(t) = c_k^{\mathsf{prev}}(t) + c_k^{\mathsf{tar}}(t) + c_k^{\mathsf{next}}(t) + g_k(t) + \epsilon_k(t) \tag{6}$$

where c_k^{prev}, c_k^{tar} and c_k^{next} refer to PSC effects from the previous trial, the target trial, and the next trial(s). Note that c_k^{prev} is not the same as c_{k-1}^{tar} as each trial was simulated entirely independently. Here $g_k(t)$ represents temporally-correlated noise from a Gaussian process,

$$g_k \sim \mathsf{Normal}(\mathbf{0},\mathbf{K}), \tag{7}$$

with the covariance matrix $\mathbf{K}$ defined by the radial basis function kernel with noise variance σ_{scale} and characteristic lengthscale ℓ_{gp}

$$\mathbf{K}_{t_1,t_2} = \sigma_{\mathsf{scale}}\exp\left(\frac{-(t_1-t_2)^2}{2\ell_{\mathsf{gp}}^2}\right). \tag{8}$$

Finally, $\epsilon_k(t)$ represents uncorrelated noise sampled from a univariate Gaussian,

$$\epsilon_k(t) \sim \mathsf{Normal}(0,\sigma_{\mathsf{noise}}^2) \tag{9}$$

where $\sigma^2_{\text{noise}} \sim U(\sigma^2_{\text{noise,min}}, \sigma^2_{\text{noise,max}})$. The NWD network is trained to infer c_k^{tar} from c_k by minimizing the mean-squared error loss.

The time constants of the training data were selected to match the PSCs recorded in experiments via an interactive simulator that overlaid the template waveform on experimental traces. We found that the NWD network performed best when trained separately on training data matched to either excitatory or inhibitory synaptic currents depending on the data being processed.

The NWD network intrinsically performs a denoising of the evoked PSCs since each target PSC in the training data was uncorrupted by the noise terms g_k and ϵ_k. We found that we could obtain further improvement in PSC denoising by randomly incorporating 45 ms snippets of pure electrical noise taken from experiments during periods without stimulation. For such "negative" templates, the network was trained to produce an output of zeros.

Finally, we performed a minor correction step to guarantee that the output of the NWD network monotonically decayed after a predefined time [Lee et al., 2020]. Namely, after a user-specified time t_{monotone}, the network output N_t for all $t > t_{\text{monotone}}$ was adjusted by recursively setting $N_t = \min\{N_t, N_{t-1}\}$.

We implemented NWD using PyTorch Lightning. Networks were trained using stochastic gradient descent with a batch size of 64 and learning rate of 0.01 for 3000-6000 epochs, which was sufficiently long for the optimizer to converge in our training regime.

Statistical model

CAVIaR aims to fit a statistical model to the demixed optogenetic data that relates patterns of holographic stimulation to the resulting measurements of postsynaptic current. Let $\mathbf{c}_k \in \mathbb{R}^T$ represent the PSC trace on trial k. In accordance with the trial structure used for NWD, we assume that $T = 900$ frames (or 45 ms), with photostimulation beginning at $t = 100$ (or 5 ms). To make use of compressed sensing-style techniques, we collapse the demixed, denoised PSC trace into a single number by integrating over the duration of the trial,

$$y_k = \int_0^T c_k^{\text{NWD}}(t)dt, \quad k = 1, \ldots, K, \tag{10}$$

where c_k^{NWD} represents the demixed trace on trial k. The measurements y_k therefore represent the total synaptic charge transfer resulting from the transmission of presynaptic spikes. Since the postsynaptic neuron is held under voltage clamp, we may model these measurements as a simple sum of optogenetically-evoked presynaptic spikes weighted by the strengths of the corresponding synapses,

$$y_k \sim \text{Normal}\left(\mathbf{w}^\top \mathbf{s}_{:,k}, \sigma^2\right), \qquad \sigma^{-2} \sim \text{Gamma}\left(t_{\text{sh}}, t_{\text{ra}}\right). \tag{11}$$

Here $\mathbf{w}$ is a vector of synaptic weights encoding the charge transfer resulting from a single spike, and $\mathbf{s}_{:,k}$ is the vector of presynaptic spikes on trial k. Note that, as described below, spike generation is highly stochastic, and hence s_{nk} is a latent variable. The synaptic weights are regularized by imposing a Gaussian prior,

$$w_n \sim \text{Normal}(u, b^2), \tag{12}$$

with hyperparameters u and b^2, and the variance σ^2 is intended to account for electrical noise from the recording electrode and variability in the amplitude of the evoked PSCs. Model parameters are listed in Table 1.

The variability of spike generation in our experimental data shows that the probability of generating a spike is not only different between cells, but also across power levels, and is therefore effectively impossible to know ahead of time. We propose a structured presynaptic model that can flexibly adjust to varying spike probabilities on a cell-by-cell basis. In particular, presynaptic spikes are assumed to be driven by holographic photostimulation according to a linear-nonlinear-Bernoulli model,

$$s_{nk} \mid \phi_n \sim \text{Bernoulli}\left(f\left(\phi_n^0 I_{nk} - \phi_n^1\right)\right) \tag{13}$$

Symbol	Interpretation	Hyperparameters	Posterior
$\mathbf{w}$	Connection strengths	u, b^2	$\boldsymbol{\mu}, \boldsymbol{\Omega}$
s_{nk}	Presynaptic spike	–	λ_{nk}
ϕ_n^0, ϕ_n^1	Presynaptic sigmoid coefficients	$\mathbf{v}, \mathbf{L}$	$\boldsymbol{\nu}_n, \boldsymbol{\Sigma}_n$
σ^2	Observation noise variance	$t_{\mathsf{sh}}, t_{\mathsf{ra}}$	$\theta_{\mathsf{sh}}, \theta_{\mathsf{ra}}$
z_k	Spontaneous synaptic current		
y_k	Observed response (integral of the PSC)		
I_{nk}	Laser power applied to neuron n on trial k		
N (n)	Number of neurons (neuron index)		
K (k)	Number of trials (trial index)		
f	Sigmoid function		

Table 1: Table of model parameters and their interpretations.

where $f(x) = 1/(1 + \exp(-x))$ is the sigmoid nonlinearity, I_{nk} denotes the power of the stimulation laser focused on cell n on trial k, and the sigmoid coefficients follow bivariate normal distributions

$$\phi_n \sim \mathsf{Normal}(\mathbf{v}_n, \mathbf{L}_n). \tag{14}$$

During inference we thus effectively perform a Bayesian logistic regression of each cell's inferred presynaptic spikes on the laser targets. This presynaptic model allows spike probabilities to adapt to variation in opsin expression via the sigmoid coefficient term ϕ_n^0, as well as to variation in rheobase via the intercept term ϕ_n^1.

Posterior inference

Variational approximation

The goal of CAVIaR is to infer the posterior distribution over the hidden variables, including the synaptic weights and presynaptic spikes. The posterior describes the distribution of parameter values that are consistent with the observed data, given our prior assumptions. Importantly, this provides a description of uncertainty in the inferred synaptic weights. It is not computationally feasible to compute the exact posterior distribution, but standard methods have been developed to approximate the posterior [Blei et al., 2017]. Here we modify the conventional coordinate-ascent variational inference algorithm by augmenting it with several steps that account for spontaneous synaptic currents and biophysical plausibility of the inferred spikes. To this end, we note that the posterior distribution over the latent variables factorizes as

$$p(\mathbf{w}, \mathbf{s}, \boldsymbol{\phi}, \sigma^2 \mid \mathbf{y}, \mathcal{I}) \propto p(\sigma^2) \prod_{n=1}^{N} p(w_n) p(\phi_n) \prod_{k=1}^{K} p(s_{nk} \mid I_{nk}, \phi_n) p(y_k \mid \mathbf{w}, \mathbf{s}_{:,k}, \sigma^2). \tag{15}$$

We approximate the posterior distribution $p(\mathcal{Z} \mid \mathbf{y}, \mathcal{I})$ by a variational model $q(\mathcal{Z})$ (where $\mathcal{Z}$ represents the set of latent variables) that obeys a similar factorization,

$$q(\mathbf{w}, \mathbf{s}, \boldsymbol{\phi}, \sigma^2) = q(\sigma^2 \mid \theta_{\mathsf{sh}}, \theta_{\mathsf{ra}}) q(\mathbf{w} \mid \boldsymbol{\mu}, \boldsymbol{\Omega}) \prod_{n=1}^{N} q(\phi_n \mid \boldsymbol{\nu}_n, \boldsymbol{\Sigma}_n) \prod_{k=1}^{K} q(s_{nk} \mid \lambda_{nk}) \tag{16}$$

where the individual factors are $q(\mathbf{w} \mid \boldsymbol{\mu}, \boldsymbol{\Omega}) = \mathsf{Normal}(\mathbf{w} \mid \boldsymbol{\mu}, \boldsymbol{\Omega})$, $q(s_{nk} \mid \lambda_{nk}) = \mathsf{Bernoulli}(s_{nk} \mid \lambda_{nk})$, $q(\phi_n \mid \boldsymbol{\nu}_n, \boldsymbol{\Lambda}_n) = \mathsf{Normal}(\phi_n \mid \boldsymbol{\nu}_n, \boldsymbol{\Sigma}_n)$, and $q(\sigma^{-2} \mid \theta_{\mathsf{sh}}, \theta_{\mathsf{ra}}) = \mathsf{Gamma}(\sigma^{-2} \mid \theta_{\mathsf{sh}}, \theta_{\mathsf{ra}})$.

Given the above approximation, coordinate-ascent variational inference seeks to perform an update of each factor $q(\mathcal{Z}_i)$ one-by-one for all $\mathcal{Z}_i \in \mathcal{Z}$, with the idea being that each update moves the approximate posterior $q(\mathcal{Z})$ closer to the true posterior $p(\mathcal{Z} \mid \mathbf{y}, \mathcal{I})$ in the sense of the Kullback-Leibler divergence (KL-divergence) [Blei et al., 2017]. We take a

variety of approaches to updating each factor depending on the relative tractability of the update. The complete algorithm is given in Algorithm 1.

Inference of synaptic weights

First, it can be shown that the optimal variational update for the synaptic strengths $\mathbf{w}$, conditional on all other latent variables $\mathcal{Z} \setminus \mathbf{w}$, obeys the equation

$$q(\mathbf{w} \mid \boldsymbol{\mu}, \boldsymbol{\Omega}) \propto \exp \mathbb{E}_{q(\mathcal{Z} \setminus \mathbf{w})} \left[\ln p(\mathbf{y}, \mathcal{Z} \mid \mathcal{I}) \right], \tag{17}$$

see, e.g., [Bishop, 2006]. Since both the prior on $\mathbf{w}$ and the observations are Gaussian-distributed, one can evaluate Equation 17 analytically and solve for $\boldsymbol{\mu}$ and $\boldsymbol{\Omega}$ by "completing the square", yielding the block update

$$\boldsymbol{\Omega} = \left(\frac{\theta_{\mathsf{sh}}}{\theta_{\mathsf{ra}}} \sum_{k=1}^{K} (\mathbf{D}_k + \mathbf{M}_k) + \frac{1}{b^2} \mathbf{I}_n \right)^{-1}, \quad \boldsymbol{\mu} = \boldsymbol{\Omega} \left(\frac{\theta_{\mathsf{sh}}}{\theta_{\mathsf{ra}}} \sum_{k=1}^{K} y_k \boldsymbol{\lambda}_{:,k} + \frac{1}{b^2} u \right) \tag{18}$$

where $\mathbf{D}_k = \mathrm{diag}\left(\lambda_{1,k}(1 - \lambda_{1,k}), \ldots, \lambda_{Nk}(1 - \lambda_{Nk}) \right) \in \mathbb{R}^{N \times N}$ is diagonal and $\mathbf{M}_k = \boldsymbol{\lambda}_{:,k} \boldsymbol{\lambda}_{:,k}^{\top} \in \mathbb{R}^{N \times N}$. Inspecting the update for $\boldsymbol{\mu}$, one sees that a neuron's synaptic weight is only determined by the magnitude of the postsynaptic responses on trials for which that neuron actually elicited a spike. However, those spikes are not directly observed, and must themselves be inferred from the data.

Inference of presynaptic spikes

Given a measurement y_k resulting from the photostimulus $I_{:,k}$ and our current estimates of synaptic strengths, electrical noise, and *a priori* presynaptic spike probabilities, we must decide which of the targeted cells actually elicited a spike. In early trials, this depends more on the prior (i.e., the presynaptic model) rather than on the likelihood, and this balance shifts in favor of the likelihood the more experimental data that is collected. However, unlike in Equation 17, the optimal variational update for the presynaptic spike probabilities λ_{nk} cannot be solved for analytically. Instead, one can recognize ([Bishop, 2006]) that the KL-divergence from the approximate to the true posterior, $\mathrm{KL}(q\|p)$, can be decomposed into the sum

$$\mathrm{KL}(q\|p) = \ln p(\mathbf{y}) - \mathcal{L}(\mathcal{Z}) \tag{19}$$

where $\ln p(\mathbf{y}) = \ln \int p(\mathbf{y}, \mathcal{Z}) d\mathcal{Z}$ is the logarithmic model evidence and where

$$\mathcal{L}(\mathcal{Z}) = \mathbb{E}_q \left[\ln p(\mathbf{y}, \mathbf{w}, \mathbf{s}, \boldsymbol{\phi}, \sigma^2) \right] + \mathcal{H}_q \left[\mathbf{w}, \mathbf{s}, \boldsymbol{\phi}, \sigma^2 \right], \tag{20}$$

with $\mathcal{H}_q[x]$ the Shannon entropy of variable x under density q. Since the KL-divergence is always non-negative, the inequality $\mathcal{L}(\mathcal{Z}) \leq \ln p(\mathbf{y})$ always holds, and therefore $\mathcal{L}(\mathcal{Z})$ is a lower bound on the logarithmic model evidence (referred to as the "evidence lower bound"). By maximizing the evidence lower bound as a function of the variational model parameters, $\mathcal{L}(\mathcal{Z})$ approaches $\ln p(\mathbf{y})$, and hence drives $\mathrm{KL}(q\|p)$ towards zero. If the bound is tight, i.e., if $\mathcal{L}(\mathcal{Z}) = \ln p(\mathbf{y})$, the KL-divergence is zero and the variational approximation to the posterior becomes exact. To this end, differentiating $\mathcal{L}$ and solving for λ_{nk} yields the update

$$\ln \left(\frac{\lambda_{nk}}{1 - \lambda_{nk}} \right) = \mathbb{E}_{q(\boldsymbol{\phi})} \left[\ln \left(\frac{f_{nk}}{1 - f_{nk}} \right) \right] - \frac{\theta_{\mathsf{sh}}}{2\theta_{\mathsf{ra}}} \left\{ -2y_k \mu_n + 2\mu_n \sum_{j \neq n} \mu_j \lambda_{jk} - \left(\mu_n^2 + \beta_n^2 \right) \right\}$$

where we have defined $f_{nk} = f\left(\phi_n^0 I_{nk} - \phi_n^1 \right)$. The expectation in the above term is analytically intractable due to multiple nonlinearities. We thus make a Monte Carlo approximation to the expectation

$$\mathbb{E}_{q(\boldsymbol{\phi})} \left[\ln \left(\frac{f_{nk}}{1 - f_{nk}} \right) \right] \approx \frac{1}{M} \sum_{m=1}^{M} \ln \left(\frac{f_{nk}[m]}{1 - f_{nk}[m]} \right) \tag{21}$$

with $f_{nk}[m] = f\left(\phi_n^0[m]I_{nk} - \phi_n^1[m]\right)$, and for each pair $(\phi_n^0[m], \phi_n^1[m])^\top \sim q(\phi_n \mid \nu_n, \Lambda_n)$. As discussed below, the sigmoid coefficients are sampled from a *truncated* multivariate normal distribution. However, sampling from truncated multivariate normal distributions is computationally demanding and remains an open area of research [Pakman and Paninski, 2014]. Instead, the terms ϕ_n^0 and ϕ_n^1 are sampled independently from their truncated marginal distributions.

Inspecting the updates for λ_{nk}, one notices that each update depends on all other neurons in the population, but is independent of all other trials. Thus, we update the spike probabilities for all trials simultaneously, but perform this one neuron at a time (where the order of the neuron updates is randomized every iteration). Immediately following the inference of a neuron's spikes, we evaluate their biophysical plausibility as a function of laser power. It is well known from patch-clamp studies with opsin-expressing neurons that the probability of photoeliciting an action potential should increase monotonically with laser power [Sridharan et al., 2022]. We therefore perform an isotonic regression through the inferred spike rate using the pool adjacent violators algorithm (PAVA) [Ayer et al., 1955].

Concretely, let $K_n^p = \{k \in \{1, \ldots, K\} : I_{nk} = p\}$ be the set of all trials on which neuron n was stimulated with power p, and let

$$\bar{\lambda}_n^p = \frac{1}{|K_n^p|} \sum_{k \in K_n^p} \lambda_{nk} \tag{22}$$

denote the average inferred spike probability for neuron n at power p. PAVA computes an isotonic regression function $\hat{F}_n$ by solving the optimization problem

$$\text{minimize} \sum_{p \in \mathcal{P}} (\bar{\lambda}_n^p - \hat{F}_n(p))^2 \quad \text{such that } \hat{F}_n \text{ is non-decreasing,} \tag{23}$$

where $\mathcal{P}$ is the set of non-zero powers used in the experiment. This provides us with an isotonically-constrained optogenetic "power curve" $\hat{F}_n$ that we use to judge whether or not the putative synapse connecting neuron n to the patched neuron is sufficiently plausible. In particular, we require that at the maximal power used in the experiment the inferred probability of successfully initiating and transmitting a spike is greater than some threshold $\theta_{\text{PAVA}} \in [0, 1]$, which we refer to as the "min spike rate at max power" (MSRMP). However, some fraction of the inferred spikes may result from spontaneous synaptic currents, which can lead to false positives if left unaccounted for. We therefore infer the rate of spontaneous currents (λ_{spont}, described below) and add this to the user-defined threshold θ_{PAVA} to adaptively adjust the plausibility criterion. Neurons with $\hat{F}_n(\max_k I_{nk}) < \theta_{\text{PAVA}} + \lambda_{\text{spont}}$ (i.e., whose adjusted maximal spike rate is below θ_{PAVA}) are considered unconnected and have their synaptic weights and inferred spike probabilities set to zero. We used $\theta_{\text{PAVA}} = 0.4$ for mapping inhibitory presynaptic neurons, and $\theta_{\text{PAVA}} = 0.3$ for mapping excitatory presynaptic neurons.

We also use a "masking" procedure to prevent small, noise-driven postsynaptic measurements from causing spurious spike inferences. To do this, we evaluate a test statistic τ on each trace c_k. If $\tau(c_k) < \tau_{\text{min}}$ we force $\lambda_{nk} = 0$ for all $n = 1, \ldots, N$. The statistic τ is typically chosen to be the sample autocorrelation.

Inference of presynaptic sigmoid coefficients

The presynaptic sigmoid coefficients ϕ (governing the presynaptic models) are updated using Laplace's method. Namely, rather than solving for the optimal variational approximation or optimizing the evidence lower bound, we learn a posterior distribution over each ϕ_n by making a second-order Taylor approximation about the posterior mode, with the posterior covariance matrix given by the inverse of the Hessian matrix appearing in the Taylor series. In some cases we encountered pathological spike inference when the sigmoid coefficients ϕ_n became negative, and hence we enforce a positivity constraint in the mode-finding algorithm described below. Moreover, the Monte Carlo approximation described in Equation 21 depends closely on the non-negativity of the sampled variates. Thus we truncate the Laplace approximation to the non-negative real line,

$$q(\phi_n \mid \nu_n, \Sigma_n) = \text{TruncNorm}(\phi_n \mid \nu_n, \Sigma_n) \tag{24}$$

with $\nu_n^i > 0$ for $i \in \{0,1\}$ and supp $q(\phi_n \mid \nu_n, \Sigma_n) = \mathbb{R}_{>0}^2$.

Note that, while PAVA is a non-parametric estimator of the spike rate, its use differs from the presynaptic spike model which takes a parametric form. The presynaptic model is used to evaluate the prior probability of photoeliciting a spike, whereas PAVA is used to evaluate the plausibility of the inferred spikes.

We use Newton's method with a log barrier and backtracking line search to identify the posterior mode ν_n. To this end, define the objective function with barrier penalty t as

$$\Psi_n^t = -\mathbb{E}_{q(\mathbf{s}_n)} \left[\sum_{k=1}^K \ln p(s_{nk} \mid \phi_n, I_{nk}) + \ln p(\phi_n \mid \mathbf{v}, \mathbf{L}) \right] - \frac{1}{t} \sum_{i=0}^1 \ln(\phi_n^i),$$

where we average over the spike uncertainty. Note that the presynaptic models depend only on the spikes $\mathbf{s}$, and are independent conditional on $\mathbf{s}$, allowing us to compute Laplace approximations in parallel across cells.

The mode-finding algorithm starts with an initial t. We then solve for the mode by iterating the Newton steps

$$\nu_n^{(t)} \leftarrow \nu_n^{(t)} - \kappa \mathbf{H}_{nt}^{-1} \mathbf{J}_{nt} \tag{25}$$

until convergence, where $\mathbf{J}$ and $\mathbf{H}$ are respectively the Jacobian and Hessian of the presynaptic model for neuron n with barrier sharpness t

$$\mathbf{J}_{nt} = \nabla_{\phi_n} \Psi_n^t, \quad \mathbf{H}_{nt} = \nabla \nabla_{\phi_n} \Psi_n^t. \tag{26}$$

The stepsize κ is adaptively selected using a standard backtracking line search rule. We then increase t to sharpen the log barrier and repeat as required. Typically we only sharpen t two or three times as this proved sufficient for our data.

Inference of observation noise variance

To infer the observation noise variance, we use the same approach as with the update rule for the synaptic weights (Equation 17). In particular, noting that

$$q(\sigma^{-2} \mid \theta_{\mathsf{sh}}, \theta_{\mathsf{ra}}) \propto \exp \mathbb{E}_{q(\mathcal{Z} \backslash \sigma^{-2})} \left[\ln p(\mathbf{y}, \mathcal{Z} \mid \mathcal{I}) \right],$$

and recognizing that the gamma prior is conjugate to the Gaussian likelihood, one obtains the variational update

$$\theta_{\mathsf{sh}} = t_{\mathsf{sh}} + \frac{K}{2}, \quad \theta_{\mathsf{ra}} = t_{\mathsf{ra}} + \frac{1}{2} \sum_{k=1}^K \mathbb{E}_q \left[\left(y_k - \mathbf{w}^\top \mathbf{s}_{:,k} \right)^2 \right]$$

where the expectation above can be evaluated analytically, though we leave it in the above form for legibility.

Inference of spontaneous synaptic currents

Finally, we estimate spontaneous synaptic currents z_k using soft-thresholding. The idea is that, given a numerical tolerance ϵ (where typically $\epsilon = 0.05$) and positively rectified residuals

$$e_k := [y_k - \mu^\top \lambda_{:,k}]_+ \tag{27}$$

we apply a soft-thresholding function S with penalty γ defined by

$$S(e, \gamma) := \begin{cases} 0 & \text{if } e \leq \gamma \\ e - \gamma & \text{otherwise.} \end{cases} \tag{28}$$

We then iteratively shrink the penalty γ until the norm of the residual data comprises no more than ϵ of the norm of the observed data; i.e., until

$$\frac{\sum_{k=1}^{K}(y_k - \boldsymbol{\mu}^\top \boldsymbol{\Lambda}_{:,k} - z_k)^2}{\sum_{k=1}^{K} y_k^2} \leq \epsilon \tag{29}$$

where z_k is obtained via the soft-thresholded residuals. However, we also require that $(z_1, \ldots, z_K)$ be approximately orthogonal to $\boldsymbol{\lambda}_n$ for all n, and apply the masking procedure as noted in the spike inference step above, leading to the spontaneous synaptic current estimator

$$z_k = \begin{cases} 0 & \text{if } \sum_{n=1}^{N} \lambda_{nk} > \theta_{\text{orthog}} \text{ or } \tau(c_k) < \tau_{\text{min}} \\ S(e_k, \gamma) & \text{otherwise} \end{cases} \tag{30}$$

where $\theta_{\text{orthog}} \approx 0$, but is not exactly zero to allow for numerical imprecision. The spontaneous events are constrained to be approximately orthogonal to the inferred spikes to prevent variability in PSC amplitude causing false positives.

Post-hoc scan for false negatives

Occasionally the CAVIaR algorithm will erroneously declare a neuron to be disconnected, either due to a failure to meet the PAVA-based plausibility criterion in the earliest iterations of the algorithm, or due to the biological data violating the assumptions of the underlying statistical model. To correct for this, following inference we perform a post-hoc scan for potential false negatives. The idea is to reconnect neurons that CAVIaR originally declared to be disconnected by checking whether spontaneous PSCs coinciding with the stimulation of a selected neuron could constitute valid postsynaptic responses when the rigidity of the statistical model is relaxed. Since the overwhelming bulk of connections are already identified by the first CAVIaR pass, missed connections are comparatively rare and thus we found a simple greedy algorithm to be effective.

The false-negative scanning algorithm is given in Algorithm 3. Briefly, the algorithm begins by collecting all neurons S_{disc} declared to be disconnected by the first CAVIaR pass. It then selects the neuron $n^* \in S_{\text{disc}}$ with the greatest number of coincidental spontaneous PSCs, and checks if assigning these PSCs to neuron n^* would satisfy the PAVA criterion. If so, neuron n^* is declared connected, with the posterior distribution of the reconnected neuron's parameters determined by sample statistics of the corresponding spontaneous PSCs. Neuron n^* is then removed from S_{disc} (whether reconnected or not), and the algorithm repeats until $S_{\text{disc}} = \emptyset$.

Inference of canonical postsynaptic current waveforms

Once we have inferred the presynaptic spike matrix $\boldsymbol{\Lambda}$, we can obtain accurate PSC waveforms $\mathbf{r}_n \in \mathbb{R}^T$ using ridge regression. Collecting the waveforms in the rows of a matrix $\mathbf{R}$, they can be obtained simultaneously by solving the non-negative L_2 problem

$$\hat{\mathbf{R}} = \underset{\mathbf{R} \geq \mathbf{0}}{\operatorname{argmin}} \left\{ \|\mathbf{C} - \boldsymbol{\Lambda}^\top \mathbf{R}\|_F + \gamma \|\mathbf{R}\|_F \right\} \tag{31}$$

where $\mathbf{C} \in \mathbb{R}^{K \times T}$ is the matrix of PSC traces $(\mathbf{c}_1, \ldots, \mathbf{c}_K)$, $\gamma > 0$ is the ridge penalty, and $\| \cdot \|_F$ is the Frobenius norm. Note that by using the spike matrix $\boldsymbol{\Lambda}$ instead of the optogenetic "design matrix" (where each element of the matrix determines which neurons were merely stimulated as in [Hu and Chklovskii, 2009], rather than which spiked as a result of photostimulation), the estimated waveforms are much less biased by trials in which neurons were not photoactivated. We used this ridge regression approach to determine synaptic weights from PSCs throughout the paper.

Algorithm 1: Coordinate-ascent variational inference and isotonic regularization (CAVIaR)

---

    input: PSC traces $\mathbf{c}$, stimulus information $\mathcal{I}$, PAVA threshold θ_{PAVA}, spontaneous penalty backtracking scalar α,
        soft orthogonality threshold θ_{orthog}, minimal test statistic τ_{min}, number of iterations *iters*

1   initialise $\lambda_{nk} \leftarrow 1$ for all n, k such that $I_{nk} > 0$ and $\tau_{\mathsf{test}}(\mathbf{c}_k) \geq \tau_{\mathsf{min}}$

2 $\lambda_{\mathsf{spont}} \leftarrow 0$ // initialize spontaneous rate to 0

3 $i \leftarrow 1$

4   **while** $i \leq iters$ **do**

5   update $q(\mathbf{w} \mid \boldsymbol{\mu}, \boldsymbol{\Omega}) \propto \exp \mathbb{E}_{q(\mathcal{Z} \backslash \mathbf{w})}\left[\ln p(\mathbf{y}, \mathcal{Z} \mid \mathcal{I})\right]$ // variational solution for synaptic weights

6   **for** $n = 1, \ldots, N$ **do**   // infer spikes via Monte Carlo ELBO solution

7   sample $\phi_n[m] \sim q(\phi_n \mid \boldsymbol{\nu}, \boldsymbol{\Sigma})$ for $m = 1, \ldots, M$

8   **for** $k = 1, \ldots, K$ **do**

9 $\lambda_{nk} \leftarrow \mathrm{argmax}_{\lambda_{nk}} \, \mathsf{ELBO}(\lambda_{nk} \mid \{\phi_n[m]\}_{m=1}^{M})$

10   **end**

11 $\hat{F}_n \leftarrow \mathsf{PAVA}(\mathcal{I}, \boldsymbol{\lambda}_n)$ // estimate optogenetic power curve

12   **if** $\hat{F}_n(\max_k I_{nk}) < \theta_{PAVA} + \lambda_{spont}$ **then**   // check plausibility criterion

13 $\mu_n \leftarrow 0, \boldsymbol{\lambda}_n \leftarrow \mathbf{0}$

14   **end**

15   **end**

16   **for** $n = 1, \ldots, N$ **do**   // Laplace approx of receptive field posterior

17 $\boldsymbol{\nu}_n, \boldsymbol{\Sigma}_n \leftarrow \mathrm{R{\small ECEPTIVE}F{\small IELD}L{\small APLACE}}(\mathbf{s}_n, \phi_n, \mathbf{v}, \mathbf{L})$

18   **end**

19   update $q(\sigma^{-2} \mid \theta_{\mathsf{sh}}, \theta_{\mathsf{ra}}) \propto \exp \mathbb{E}_{q(\mathcal{Z} \backslash \sigma^{-2})}\left[\ln p(\mathbf{y}, \mathcal{Z} \mid \mathcal{I})\right]$ // variational solution for noise precision

20   **while** $\|\mathbf{y} - \boldsymbol{\mu}^\top \boldsymbol{\Lambda} - \mathbf{z}\|_2^2 / \|\mathbf{y}\|_2^2 \geq \epsilon$ **do**   // begin spontaneous PSC inference

21   **for** $k = 1, \ldots, K$ **do**

22   **if** $\sum_{n=1}^N \lambda_{nk} \leq \theta_{orthog}$ *and* $\tau_{test}(\mathbf{c}_k) \geq \tau_{min}$ **then**

23 $z_k \leftarrow S([y_k - \boldsymbol{\mu}^\top \boldsymbol{\Lambda}_{:,k}]_+, \gamma)$ // soft-threshold residual with penalty γ

24   **end**

25   **end**

26 $\gamma \leftarrow \alpha\gamma$ // shrink penalty

27   **end**

28 $\lambda_{\mathsf{spont}} \leftarrow \frac{1}{K} \sum_{k=1}^K \mathbb{1}_{[z_k \neq 0]}$ // update spontaneous rate

29 $i \leftarrow i + 1$

30   **end**

31 $\boldsymbol{\mu}, \boldsymbol{\Omega}, \boldsymbol{\Lambda}, \phi, \mathbf{z} \leftarrow \mathsf{FNS{\small CAN}}(\boldsymbol{\mu}, \mathbf{z})$ // scan resulting spontaneous PSCs for potential false negatives

---

Algorithm 2: RECEPTIVEFIELDLAPLACE

---

for $n = 1, \ldots, N$ **do**

for $t = 1, \ldots, T_{max}$ **do**

$\kappa \leftarrow 1$ // Newton stepsize

$\Psi_n(\boldsymbol{\phi}_n) = -\mathbb{E}_{q(s_n \mid \boldsymbol{\lambda}_n)} \left[\sum_{k=1}^{K} \ln p(\mathbf{s}_{nk} \mid \boldsymbol{\phi}_n, I_{nk}) + \ln p(\boldsymbol{\phi}_n \mid \mathbf{v}, \mathbf{L}) \right] - \frac{1}{\alpha_{\text{barrier}}} \sum_{i=0}^{1} \ln(\phi_n^i)$

$\mathbf{J} = \nabla_{\boldsymbol{\phi}_n} \Psi_n, \, \mathbf{H} = \nabla\nabla_{\boldsymbol{\phi}_n} \Psi_n$

$\mathbf{d} = \mathbf{H}_n^{-1} \mathbf{J}_n$ // search direction

while $\Psi_n(\boldsymbol{\phi}_n + \kappa \mathbf{d}) > \Psi_n(\boldsymbol{\phi}_n) + \alpha_{\text{backtrack}} \, \kappa \, \mathbf{J}_n^{\top} \mathbf{d}$ **do**   // backtrack

8             $\kappa \leftarrow \beta_{\text{backtrack}} \, \kappa$

end

$\boldsymbol{\phi}_n \leftarrow \boldsymbol{\phi}_n - \kappa \mathbf{d}$ // make step

end

$\boldsymbol{\nu}_n \leftarrow \boldsymbol{\phi}_n$

$\boldsymbol{\Sigma}_n \leftarrow \mathbf{H}_n^{-1}$

end

$q(\boldsymbol{\phi} \mid \boldsymbol{\nu}_n, \boldsymbol{\Sigma}_n) = \text{TruncNormal}(\boldsymbol{\nu}_n, \boldsymbol{\Sigma}, 0, \infty)$ // truncate support to $(0, \infty)$

Algorithm 3: FNSCAN

---

input: Synaptic weights $\boldsymbol{\mu}$, spontaneous synaptic currents $\mathbf{z}$, stimulus information $\mathcal{I}$, PAVA threshold θ_{PAVA}

$S_{\text{disc}} \leftarrow \{n \in \{1, \ldots, N\} : \mu_n = 0\}$ // initialize pool of candidate neurons

while $|S_{disc}| > 0$ **do**

for $n = 1, \ldots, N$ **do**   // collect spontaneous PSC indices aligning with neuron stim times

$\text{spont}_n \leftarrow \{k \in \{1, \ldots, K\} : z_k \neq 0 \text{ and } I_{nk} > 0\}$

end

$n^* \leftarrow \text{argmax}_n |\text{spont}_n|$ // select neuron with most coincidental spontaneous PSCs

$\hat{F}_{n^*}^{\text{spont}} \leftarrow \text{PAVA}\left((I_{n^*k})_{k \in \text{spont}_n}, (z_k)_{k \in \text{spont}_n} \right)$ // estimate putative power curve

if $\hat{F}_{n^*}^{\text{spont}}(\max_k I_{n^*k}) \geq \theta_{PAVA}$ **then**

$\mu_{n^*} \leftarrow \text{mean}(\{z_k : k \in \text{spont}_{n^*}\})$ // neuron passes PAVA criterion, declare connected

$\beta_{n^*} \leftarrow \text{s.e.m.}(\{z_k : k \in \text{spont}_{n^*}\})$

for $k \in \text{spont}_{n^*}$ **do**

12             $\lambda_{n^*k} \leftarrow 1$ // declare spontaneous PSC to be spike from neuron n^*

13             $z_k \leftarrow 0$ // remove spontaneous PSC from vector $\mathbf{z}$

14   **end**

15   **end**

16      $S_{\text{disc}} \leftarrow S_{\text{disc}} \setminus \{n^*\}$ // remove n^* from pool of disconnected neurons

17 **end**

Comparing estimated connectivity vectors

To quantify the similarity between connectivity vectors obtained using single-target stimulation and ensemble stimulation, we used the coefficient of determination (R^2) and the Jaccard index (both from the scikit-learn Python package). For sets of connections $\mathcal{C}_{\text{single}}, \mathcal{C}_{\text{ensemble}}$ identified respectively using single-target and ensemble stimulation, the Jaccard index is given by

$$J(\mathcal{C}_{\text{single}}, \mathcal{C}_{\text{ensemble}}) = \frac{\mathcal{C}_{\text{single}} \cap \mathcal{C}_{\text{ensemble}}}{\mathcal{C}_{\text{single}} \cup \mathcal{C}_{\text{ensemble}}}. \tag{32}$$

While the R^2 accounts for the magnitude of synaptic strengths, the Jaccard index only accounts for whether neurons are classified as connected or not (i.e., their binary status). For example, the Jaccard index takes the value 1 if single-target and ensemble stimulation yield connections that agree perfectly, and takes the value 0 if they do not share any common elements.

Leave-one-hologram-out cross-validation

We assess the accuracy of the CAVIaR inferences using leave-one-hologram-out cross-validation (LOHO-CV, Algorithm 4). Let $\mathcal{H}$ represent the complete set of hologram designs; i.e., $h \in \mathcal{H}$ determines which neurons will be targeted for stimulation, regardless of laser power. LOHO-CV proceeds by selecting $h \in \mathcal{H}$, fitting CAVIaR to the PSC traces and stimuli corresponding to all other holograms $\mathcal{H} \setminus \{h\}$, and then averaging samples from the posterior predictive distribution over responses to hologram h to obtain an estimate of the postsynaptic response (at multiple power levels).

Algorithm 4: LOHO-CV

input: PSCs $\mathbf{c}$, stimulus information $\mathcal{I}$, holograms $\mathcal{H}$, number of posterior samples J

1 **for** $h \in \mathcal{H}$ **do**
2   estimate variational parameters $\boldsymbol{\mu}, \boldsymbol{\beta}, \boldsymbol{\nu}, \boldsymbol{\Sigma}$ from $\{(\mathbf{c}_k, I_{:,k}) : I_{:,k} \text{ is not an instance of } h\}$ with CAVIaR
3   **for** $j = 1, \ldots, J$ **do**
4   **for** $n = 1, \ldots, N$ **do**   // begin ancestral sampling from posterior predictive distribution
5   sample $(\phi_n^0, \phi_n^1) \sim \text{TruncNorm}(\boldsymbol{\nu}_n, \boldsymbol{\Sigma}_n)$
6   sample $w_n \sim \text{Normal}(\mu_n, \beta_n^2)\mathbb{1}_{[\mu_n \neq 0]} + \delta_0 \mathbb{1}_{[\mu_n = 0]}$
7   **for** *each laser power p* **do**
8   sample $s_n^p \sim \text{Bernoulli}(\sigma(\phi_n^0 h_n^p - \phi_n^1))$, where h_n^p is the power delivered to neuron n assuming hologram h is applied with power p
9   **end**
10   **end**
11   **for** *each laser power p* **do**
12   set $y^p[j] \leftarrow \mathbf{w}^\top \mathbf{s}^p$
13   **end**
14   **end**
15   predict response to hologram h at power p as $\frac{1}{J} \sum_{j=1}^{J} y^p[j]$
16 **end**

Simulated circuit mapping experiments

We used simulated data to characterize the performance of the techniques we tested. To ensure the accuracy of this characterization, rather than sampling directly from the generative model that we propose, we added several layers of

biophysical realism. In particular, we sampled noisy synaptic currents that were first demixed using NWD before being supplied as input to the connectivity inference algorithms. This way we could test for the combined accuracy of NWD and connectivity inference.

The simulated data was generated as follows. First, sigmoid parameters ϕ_n^X were sampled uniformly from $U(\phi_{\min}^X, \phi_{\max}^X)$ for $X \in \{0, 1\}$. Then, presynaptic spikes s_{nk} were sampled from a linear-nonlinear-Bernoulli model

$$s_{nk} \sim \text{Bernoulli}\left(\sigma\left(\phi_n^0 I_{nk} - \phi_n^1\right)\right) \tag{33}$$

where the laser power on each trial was randomly selected from a discrete set matched to the experimental data. Each neuron had a canonical PSC transient $h(t, \Delta_{nk}; \tau_r^n, \tau_d^n)$ that took a form similar to that used for training the NWD networks. The unnormalized transient $\tilde{h}_n$ was defined by

$$\tilde{h}(t, \Delta_{nk}; \tau_r^n, \tau_d^n) = \left[\exp\left(-\frac{t - \Delta_{nk}}{\tau_d^n}\right) - \exp\left(-\frac{t - \Delta_{nk}}{\tau_r^n}\right)\right] \mathbb{1}_{[t \geq \Delta_{nk}]} \tag{34}$$

where τ_r^n and τ_d^n are rise and decay time constants and Δ_{nk} represents the spike delay (i.e., the combined spike photoelicitation and transmission latencies) for neuron n on trial k. The transient was then normalized to take integral 1,

$$h(t, \Delta_{nk}; \tau_r^n, \tau_d^n) = \tilde{h}(t, \Delta_{nk}; \tau_r^n, \tau_d^n) / \int \tilde{h}_n(t', \Delta_{nk}; \tau_r^n, \tau_d^n) dt', \tag{35}$$

such that the synaptic weight for that neuron (w_n, defined as the synaptic charge transfer below) was preserved when multiplying the weight by the PSC.

The spike delays were laser power-dependent so that, in accordance with experimental data for the opsins used in this study [Sridharan et al., 2022], neurons initiated and propagated spikes faster if stimulated at higher powers. Concretely, for laser power I_{nk} targeted at neuron n on trial k, the spike time was sampled from a right-shifted gamma distribution,

$$\Delta_{nk} \sim \text{Gamma}\left(\frac{\alpha}{I_{nk}^2}, \beta, \Delta_{\min}\right) \tag{36}$$

where

$$\text{Gamma}(x \mid a, b, t) = \frac{b^a}{\Gamma(a)}(x - t)^{a-1} \exp(-b(x - t)) \mathbb{1}_{[x > t]}. \tag{37}$$

Note that since the mean of the shifted gamma distribution is $t + a/b$, the optogenetically evoked spikes followed, in expectation, an inverse-square dependence of time on power, consistent with the physics of two-photon absorption [Papagiakoumou et al., 2020]:

$$\mathbb{E}[\Delta_{nk}] = \Delta_{\min} + \frac{\alpha}{\beta I_{nk}^2}. \tag{38}$$

The synaptic weights w_n are intended to model the total synaptic charge transfer resulting from the transmission of a presynaptic spike to the postsynaptic neuron. We sampled the weights in a way that reflects the typical observations from our experiments. Namely, of the neurons that were chosen to be synaptically connected, a small number of them were strongly connected (in our simulations this was 20% of the connected population, though the precise value did not notably impact our results) and a large number of them were more weakly connected (80%). Specifically, for a given connectivity rate $\alpha \in (0, 1)$, a subset of $\lceil \alpha N \rceil$ neurons were randomly selected as being connected to the postsynaptic neuron. Then, if a neuron n was chosen to be strongly connected, its weights were sampled as

$$w_n \sim U(w_{\min}^{\text{strong}}, w_{\max}^{\text{strong}}) \tag{39}$$

where $w_{\min}^{\text{strong}}$ and $w_{\max}^{\text{strong}}$ respectively represent the lower and upper bounds of the uniform distribution. If n was weakly connected then its weights were sampled as

$$w_n \sim \text{Exp}(w_{\text{mean}}^{\text{weak}}, w_{\min}^{\text{weak}}) \tag{40}$$

where the exponential distribution above is in its two-parameter, right-shifted form

$$\text{Exp}(x \mid \lambda, \Delta) = \frac{1}{\lambda} \exp\left(-\frac{1}{\lambda}(x - \Delta)\right) \mathbb{1}_{[x \geq \Delta]} \tag{41}$$

with $\Delta \geq 0$.

We used two approaches to generating PSC traces. Either we directly simulated individual 45 ms trials for each $k = 1, \ldots, K$, or we simulated continuous circuit mapping experiments at 20 kHz sampling resolution (matched to the experimental data) that lasted for tens of minutes. Using the latter approach we could very closely mimic the exact contribution of confounding synaptic currents arising from stimulation at very high frequencies. Simulations performed in this continuous manner were subsequently restructured into the usual 45 ms snippets of activity.

In the first "trial-wise" approach to simulating data, the presynaptic spikes, synaptic weights, and PSC kernels were used to generate the postsynaptic responses y_k as

$$y_k = \int \left(w_n m_{nk} s_{nk} h_n(t) + h_k^{\text{spont}}(t) + g_k(t) + \epsilon_k(t)\right) dt. \tag{42}$$

Here m_{nk} is a multiplicative noise term, $h_k^{\text{spont}}(t)$ is a spontaneous PSC term, $g_k(t)$ is a temporally correlated noise term, and $\epsilon_k(t)$ is an additive noise term.

The multiplicative noise term m_{nk} accounted for the fact that, in our experimental data, the precise amplitudes of PSC transients were observed to vary from trial to trial. We sampled m_{nk} from a log-normal distribution,

$$m_{nk} \sim \text{LogNormal}(0, \sigma_{\text{mult}}^2), \tag{43}$$

such that the median postsynaptic charge transfer following a presynaptic spike still took the value w_n. The spontaneous term h_k^{spont} was either a PSC as in Equation 35 with random time constants, amplitudes, and spike times, or the zero vector, depending on the probability of spontaneous events. Temporally correlated noise, $\mathbf{g}_k \in \mathbb{R}^T$, was sampled from a Gaussian process,

$$\mathbf{g}_k \sim \text{Normal}(\mathbf{0}, \mathbf{K}) \tag{44}$$

where $\mathbf{K}$ was defined by the radial basis function kernel,

$$\mathbf{K}_{t_1, t_2} = \sigma_{\text{scale}} \exp\left(\frac{-(t_1 - t_2)^2}{2\ell_{\text{gp}}^2}\right) \tag{45}$$

for $t_1, t_2 = 1, \ldots, T$. Finally, the additive noise $\epsilon_k(t)$ was sampled independently and identically from a zero-mean Gaussian, $\epsilon_k(t) \sim \text{Normal}(0, \sigma_{\text{noise}}^2)$. Due to high computational tractability this trial-wise approach was used to generate the heatmaps in Figure 3.

The second "continuous experiment" approach to simulation was much more computationally demanding. Simulating a 30 minute experiment at 20 KHz requires 36,000,000 time points, making, for example, the use of Gaussian processes prohibitive. Our approach was to generate vectors encoding spike times for each neuron and convolve this with the corresponding neuron's PSC kernel while batching over blocks of time. To this end, an experiment of length T was evenly subdivided into trials according to the stimulation frequency f. For each such trial, we sampled spikes, spike latencies, and multiplicative noise for each neuron as described above. Then, using this collection of variables, we defined spike vectors $\boldsymbol{\zeta}_n \in \mathbb{R}^T$ for $n = 1, \ldots, N$ by setting

$$\zeta_n(\lceil t_k + \Delta_{nk} \rceil) = w_n m_{nk} s_{nk}, \quad \text{for each } k = 1, \ldots, K, \tag{46}$$

where t_k is the timebin at which the kth trial begins, and $\zeta_n(t) = 0$ at all other timebins. Similarly, we generated spontaneous spike vectors $\{\boldsymbol{\zeta}_j^{\text{spont}}\}_{j=1}^J$ at a specified rate λ_{spont} (in Hz), where each ζ_j^{spont} encodes the activation of a single spontaneous PSC with waveforms $\mathbf{h}_j^{\text{spont}}$ defined by randomly sampled time constants, amplitudes, and onset times. The

full-length postsynaptic measurement vector was then obtained by convolving the spike vectors with the PSC waveforms and summing across neurons and spontaneous inputs,

$$\mathbf{c} = \sum_{n=1}^{N} \mathbf{h}_n * \boldsymbol{\zeta}_n + \sum_{j=1}^{J} \mathbf{h}_j^{\text{spont}} * \boldsymbol{\zeta}_j^{\text{spont}} + \boldsymbol{\epsilon}. \tag{47}$$

Here the noise follows a first-order autoregressive process

$$\epsilon(t) \sim \text{Normal}(\gamma \epsilon(t-1), \sigma_{\text{noise}}^2) \tag{48}$$

with autoregressive coefficient $\gamma \in (0,1)$, which was suitably scalable for continuous simulated experiments.

The postsynaptic responses y_k used by the connectivity inference algorithms were extracted from $\mathbf{c}$ by setting

$$y_k := \int_{t_k - t_{\text{pre}}}^{t_k + t_{\text{post}}} c^{\text{NWD}}(t) dt \tag{49}$$

where t_k represents the beginning of the k-th trial, and $t_{\text{pre}} = 100$ and $t_{\text{post}} = 800$. Note that as the rate of stimulation increases, the interstimulus-interval decreases, and the trial windows increasingly overlap in time. This leads to a confounding of the observed PSCs as in Figure 2, which the NWD network must suppress.

Combinatorics of expected interstimulus intervals

Let N represent the total number of potential presynaptic neurons, R the size of the ensemble, and f the rate of stimulation (in Hz). Assuming neurons are chosen uniformly at random, the probability of selecting neuron $n \in \{1, \dots, N\}$ is

$$\frac{\binom{N-1}{R-1}}{\binom{N}{R}} = \frac{R}{N}. \tag{50}$$

Hence, one must stimulate N/R times on average to return to the same neuron. If stimulation occurs at f Hz, this takes N/Rf seconds.

Supplementary information: Rapid learning of neural circuitry from holographic ensemble stimulation enabled by Bayesian model-based compressed sensing

Symbol	Interpretation	Value
$\tau_r^{\min}$	Rise time constant minimum	10
$\tau_r^{\max}$	Rise time constant maximum	40
$\tau_{\text{diff}}^{\min}$	Difference between rise and decay time constants minimum	150
$\tau_{\text{diff}}^{\max}$	Difference between rise and decay time constants maximum	340
$\Delta_{\text{tar}}^{\min}$	PSC onset time minimum (target trial)	160
$\Delta_{\text{tar}}^{\max}$	PSC onset time maximum (target trial)	400
$\Delta_{\text{prev}}^{\min}$	PSC onset time minimum (previous trial effects)	-400
$\Delta_{\text{prev}}^{\max}$	PSC onset time maximum (previous trial effects)	159
$\Delta_{\text{next}}^{\min}$	PSC onset time minimum (next trial effects)	400
$\Delta_{\text{next}}^{\max}$	PSC onset time maximum (next trial effects)	899
σ_{scale}	GP variance	0.045
ℓ_{gp}	GP lengthscale	45
$\sigma_{\text{noise,min}}^2$	Noise variance minimum	0.001
$\sigma_{\text{noise,max}}^2$	Noise variance maximum	0.02

Table S1: Parameters used for training an NWD network for inhibitory-to-excitatory mapping experiments. Assumes timesteps measured at 20 kHz.

Symbol	Interpretation	Value
$\tau_r^{\min}$	Rise time constant minimum	10
$\tau_r^{\max}$	Rise time constant maximum	40
$\tau_{\text{diff}}^{\min}$	Difference between rise and decay time constants minimum	60
$\tau_{\text{diff}}^{\max}$	Difference between rise and decay time constants maximum	120
$\Delta_{\text{tar}}^{\min}$	PSC onset time minimum (target trial)	160
$\Delta_{\text{tar}}^{\max}$	PSC onset time maximum (target trial)	400
$\Delta_{\text{prev}}^{\min}$	PSC onset time minimum (previous trial effects)	-400
$\Delta_{\text{prev}}^{\max}$	PSC onset time maximum (previous trial effects)	159
$\Delta_{\text{next}}^{\min}$	PSC onset time minimum (next trial effects)	400
$\Delta_{\text{next}}^{\max}$	PSC onset time maximum (next trial effects)	899
σ_{scale}	GP variance	0.045
ℓ_{gp}	GP lengthscale	45
$\sigma_{\text{noise,min}}^2$	Noise variance minimum	0.001
$\sigma_{\text{noise,max}}^2$	Noise variance maximum	0.02

Table S2: Parameters used for training an NWD network for excitatory-to-excitatory mapping experiments. Assumes timesteps measured at 20 kHz.

Symbol	Interpretation	Default value
ϕ_{min}^0	Presynaptic sigmoid coefficient (0) minimum	0.2
ϕ_{max}^0	Presynaptic sigmoid coefficient (0) maximum	0.25
ϕ_{min}^1	Presynaptic sigmoid coefficient (1) minimum	10
ϕ_{max}^1	Presynaptic sigmoid coefficient (1) maximum	15
$\tau_{r,\mathsf{min}}$	Rise time constant minimum	10
$\tau_{r,\mathsf{max}}$	Rise time constant maximum	40
$\tau_{\Delta,\mathsf{min}}$	Difference between rise and decay time constants minimum	250
$\tau_{\Delta,\mathsf{max}}$	Difference between rise and decay time constants maximum	300
α	Rate parameter for gamma-distributed spike times	10^4
β	Shape parameter for gamma-distributed spike times	15
Δ_{min}	Minimum spike time	60
$w_{\mathsf{min}}^{\mathsf{strong}}$	Strong synaptic weight minimum	20
$w_{\mathsf{max}}^{\mathsf{strong}}$	Strong synaptic weight maximum	40
$w_{\mathsf{mean}}^{\mathsf{weak}}$	Weak synaptic weight mean (of the unshifted distribution)	4
$w_{\mathsf{min}}^{\mathsf{weak}}$	Weak synaptic weight minimum	5
ℓ_{gp}	GP lengthscale	50
σ_{scale}	GP noise variance	4^{-3}

Table S3: Table of default parameters for simulation studies.

Figure S1: Physiological point spread functions (PPSFs) for holographic single-target and ensemble stimulation. **a**, Illustration of multi-plane grid of stimulation sites used to obtain single-target PPSFs. A loose-patched, opsin-positive cell was positioned to be in the center plane of the grid (37.5 μm). Planes separated by 12.5 μm. **b**, Single-cell example of mean spike probability when stimulating individual points on grid (7 repetitions per power). **c**, Mean radial (left panel) and axial (right panel) PPSF for a population of opsin-expressing PV cells. Error bars indicate standard deviation over n=8 cells. **d**, Illustration of multi-plane grid of stimulation sites used to obtain 10-target ensemble PPSFs. In these experiments, one point on the grid is targeted by the hologram while the remaining 9 "control" targets are outside of the grid area and kept fixed (i.e., do not change from stimulus to stimulus). **e**, Single-cell example of mean spike probability when stimulating using a 10-target hologram (with one target being the opsin-positive cell; 7 repetitions per power). Panels (a) and (c) display results from the same cell. **f**, Mean radial (left panel) and axial (right panel) PPSF for a population of opsin-positive PV cells when stimulating 10 targets at once. Error bars indicate standard deviation over n=7 cells. **g**, Incidence of multiple action potentials (i.e., doublets) in response to a single laser pulse. Trials shown are those where the patched cell spiked at least once. Postsynaptic neuron same as in panel (b). 10 ms time windows from the start of a 3 ms laser pulse are displayed. Exemplary traces displayed on the right are marked by arrows.

Figure S2: "Power gap" between single-target and ensemble holograms. a, Illustration of the two compared stimulation conditions. An opsin-positive cell was patched and photocurrents were recorded in whole-cell configuration while the regular connectivity mapping protocol was used. Trials when single-target holograms were targeting the patched cell (left) were compared with trials when the patched cell was a part of a 10-target ensemble hologram (right). **b**, Photocurrents from an example experiment. Red stars indicate the photocurrent amplitude across increasing stimulation power for a single-target hologram applied to the patched cell, blue circles represent the response amplitude across power for different ensemble holograms containing the patched cell. **c**, Same data as in b, but showing the mean ± s.e.m. for the different ensemble hologram sets (left), and normalized to the corresponding single-target photocurrents (right, magenta triangles). **d**, Mean photocurrents evoked by ensemble holograms, normalized to photocurrents evoked by single-target holograms at the corresponding laser powers. Error bars show one standard deviation. Data collected across a population of cells; n=6 experiments performed on 1 pyramidal cell (Emx-Cre), 1 PV cell (PV-Cre), 4 SST cells (SST-Cre). **e-g**, same as b-d but with laser power for ensemble holograms increased by 35%. For g, data collected across n=2 experiments (1 pyramidal cell, 1 SST cell).

Figure S3: Components of an example simulated connectivity mapping experiment. **a**, Presynaptic spike probabilities as a function of laser power. Connected neurons shown in red, disconnected neurons shown in gray. **b**, Example histograms (shown as normalized densities) of presynaptic spike latencies for three typical laser powers. Average spike and transmission times decrease quadratically with increasing power. **c**, Example log-normal distribution of multiplicative noise terms that induce trial-to-trial variability in PSC amplitude. **d**, Distribution of synaptic weights, shown on log scale. **e**, Example PSC templates. Disconnected neuron templates represented by the zero vector. **f**, Three example noise processes sampled from a first-order autoregressive process. **g**, Example connectivity mapping experiment under increasing rates of stimulation. Simulation has 300 neurons with 10% connection probability and spontaneous synaptic currents occuring at 5 Hz.

Figure S4: a, NWD network performance on simulated test data. Solid black line shows raw data given as input to the neural network. Orange line shows target trace. Dashed purple line shows prediction from the NWD network. **b**, Performance of NWD trained on simulated data matched to IE mapping experiments (solid, blue). For comparison, NWD trained on simulated data matched to EE mapping experiments shown as dashed red line. **c**, Performance of NWD trained on simulated data matched to EE mapping experiments (solid, red). For comparison, NWD trained on simulated data matched to IE mapping experiments shown as dashed blue line.

Figure S5: Example application of NWD to individual PSCs evoked by holographic ensemble stimulation in PV (**a**) and SST (**b**) to pyramidal mapping experiments. PSCs selected uniformly at random and sorted by magnitude of postsynaptic response.

Figure S6: Example application of NWD to individual PSCs evoked by holographic ensemble stimulation in a pyramidal-pyramidal mapping experiment. PSCs selected uniformly at random and sorted by magnitude of postsynaptic response.

Figure S7: Large-scale classification of presynaptic spikes, spontaneous synaptic currents, and synaptic connectivity in a 3 minute simulated connectivity mapping experiment with 300 neurons mapped using 10-target ensemble stimulation at 10 Hz. Spontaneous PSCs occur at a rate of 5 Hz. **a**, "Checkerboard" visualization of CAVIaR model inference. Each row shows sample of PSCs evoked by stimulating the listed neuron across multiple powers. In this example, 9 neurons in addition to the listed neuron are stimulated on each trial. Shaded red cells indicate detected presynaptic spike for listed neuron. Shaded blue cells indicate detected spontaneous PSC. Triangles indicate ground-truth presynaptic spikes. Accuracy of inferred presynaptic spikes for this simulation, 94.3%. Traces shown in gray indicate neurons that the model declared disconnected. Out of 300 neurons mapped, one putative connection was a false negative (neuron 22) due to CAVIaR incorrectly assigning its PSCs to other stimulated neurons or to spontaneous activity. **b**, Comparison of connectivity inference accuracy between CAVIaR, CAVI-SnS, and CoSaMP.

Figure S8: Performance and phase transitions for three connectivity inference techniques for a population of 1000 candidate presynaptic neurons. **a**, Performance as a function of underlying density of synaptic connectivity (connection probability). **b**, Performance as a function of stimulus diversity (hologram repetitions). **c**, Performance as a function of the number of simultaneously targeted neurons. **d**, Performance as a function of probability of spontaneous PSC lying within admissible PSC initiation window.

Figure S9: Additional comparisons between connectivity maps obtained using single-target and ensemble stimulation. Opacity of circles represents synaptic strength, normalized by largest synaptic strength across both single-target and ensemble connections (fainter connections are weaker). Arrows in panel c show connections found using single-target stimulation that were not identified using ensemble stimulation.

Figure S10: Analysis of the effect of NWD on direct photocurrent artifacts in pyramidal-pyramidal (Emx-Cre; AAV-st-ChroME2s-mRuby3) mapping experiments (corresponds to Figure 6a). a-e, Photocurrents and postsynaptic responses for multiple conditions (single-target stimulation, ensemble stimulation), connection statuses (connected, not connected), and orderings (sorted by largest photocurrents, proximity to electrode). Raw traces shown in gray, demixed traces shown in color. Numbers inside parentheses show distance from electrode tip. **f-h**, Stimulation FOVs showing all targets (f), connections inferred using single-target stimulation (g), and connections inferred using ensemble stimulation (h). **i, j**, Spatial representation of the magnitude of evoked currents during stimulation period before (i) and after (j) applying NWD. Magnitude given as the average current (in picoamperes) during the stimulation window (shaded red regions in a-e) across all stimulation trials and powers. Heightened currents near the electrode tip correspond to direct photocurrents on the patched cell.

Figure S11: Outlier experiment corresponding to marked datapoint in Figure 6d. **a**, Checkerboard visualization of synaptic currents and photocurrents evoked by single-target (left) and holographic ensemble (right) stimulation. Solid grey traces correspond to demixed PSCs from ROIs considered unconnected; dashed faint grey traces are the same but without demixing. Note the presence of photocurrents in row 6 largely suppressed by NWD. **b**, Z-projected stimulation FOV and comparison between connections identified using single-target stimulation (left) and holographic ensemble stimulation (right). Given the R^2 and Jaccard scores, this experiment is the worst performing in the study, yet the disagreement between the two maps is only moderate.

Figure S12: Demixing and connectivity inference for "blind" grid mapping experiments. a, Grid mapping of PV-pyramidal connectivity using single-target stimulation over four powers (30-75 mW). One out of five planes are shown as an example. Note that maps of synaptic connectivity obtained by CAVIaR "merge" the multi-power demixed maps into a single map. **b,c**, Grid mapping of PV-pyramidal connectivity using both single-target (on plane 0 μm) and holographic ensemble stimulation (on five planes over 25 to -75 μm, separated by 25 μm each; only plane 0 μm shown for comparison) over four powers. Agreement between the two maps validates the use of CAVIaR in this regime. Each pixel in the raw maps shown in (c) obtained by averaging across all PSCs evoked by stimulation of an ensemble containing that pixel. **d**, Overlay of connected pixels (red circles) on underlying anatomy (obtained by imaging expression of mRuby) confirms that grid mapping using both single-target and ensemble stimulation correctly identifies an opsin-expressing neuron. Note that no such anatomical image is available for the Vglut1-Cre; st-ChroME-GCaMP7 mice (panel (a)) since the opsin construct does not include a constitutively fluorescent protein (e.g., mRuby).

Figure S13: Demixing and denoising of optogenetic grid mapping data using NWD and CAVIaR. Example shows PV-pyramidal connectivity mapping (PV-Cre ChroME2f). Postsynaptic cell is on plane 0 μm.