1   **Effects of social presence on behavioural, neural and physiological aspects of empathy for**

2   **pain**

4   Abbreviated title: Effects of social presence on empathy for pain

6   Pauline Petereit[1,2] *, Ronja Weiblen[1,2,3], Anat Perry[4], Ulrike M. Krämer[1,2,5] *

[1] Department of Neurology, University of Lübeck, 23562 Lübeck, Germany

[2] Center of Brain, Behavior and Metabolism (CBBM), University of Lübeck, 23562 Lübeck, Germany

[3] Department of Psychiatry and Psychotherapy, University of Lübeck, 23562 Lübeck, Germany

[4] Psychology Department, Hebrew University of Jerusalem, Jerusalem, Israel

[5] Department of Psychology, University of Lübeck, 23562 Lübeck, Germany

* Corresponding authors: Pauline Petereit, pauline.petereit@neuro.uni-luebeck.de; Ulrike M. Krämer, ulrike.kra-

emer@uni-luebeck.de

31   **Summary**

In mediated interactions (e.g. video calls), less information is available about the other. To investigate how this

affects our empathy for one another, we conducted an EEG study, in which thirty human participants observed

one of 5 targets undergoing painful electric stimulation, once in a direct interaction and once in a live, video-

mediated interaction. We found that observers were as accurate in judging others' pain and showed as much

affective empathy via video as in a direct encounter. While mu suppression, a common neural marker of empa-

thy, was not sensitive to others' pain, theta responses to others' pain as well as skin conductance coupling

between participants were reduced in the video-mediated condition. We conclude that physical proximity with

its rich social cues is important for nuanced physiological resonance with the other's experience. More studies

are warranted to confirm these results and to understand their behavioural significance for remote social inter-

actions.

Key words: social presence, pain empathy, empathic accuracy, mu suppression, physiological coupling

Introduction

Over the last decades, many social interactions in private life and at work, including medical and psychotherapeutic contexts, have shifted from personal encounters to mediated interactions such as video calls. These mediums provide less detailed social cues and information channels and limit opportunities for immediate, reciprocal interaction compared to personal interactions. These factors are sometimes referred to as intimacy and immediacy, and together they contribute to the degree of social presence offered within social interactions (Cui et al., 2013; Short et al., 1976; see Biocca et al., 2003, for alternative accounts of social presence). Given the reduced social presence of mediated interactions, the question arises of how this changes our ability to share and understand others' affective states, i.e., our ability to empathize (Decety & Jackson, 2004; Shamay-Tsoory, 2011). Focusing on intimacy, we asked how the reduction of social cues – using a video call versus a direct interaction – affects empathy. We used empathy for pain as a well-established model to investigate the behavioural, neural and physiological aspects of empathy (Singer & Lamm, 2009).

Empathy for pain is a multifaceted process, including an affective response, feelings of distress and empathic care towards the person suffering (Goubert et al., 2009; Lamm et al., 2007; Singer & Lamm, 2009), as well as cognitive processes. The latter are sometimes measured as empathic accuracy, which is the accuracy of one's perception of the other's pain (Laursen et al., 2014; Zaki et al., 2009b).

On the neural level, empathy for pain has been associated with mu suppression in the EEG: reduced power between 8 and 13 Hz over the somatosensory cortex (Cheng et al., 2008; Gallo et al., 2018; Peled-Avron et al., 2018; Peng et al., 2021; Perry et al., 2010a). Mu suppression has been linked to higher empathic accuracy (Goldstein et al., 2018) and might aid empathy by representing the other's bodily state in one's own somatosensory system (Riečanský & Lamm, 2019). Other studies examined mid-frontal theta activity (4–8 Hz) as a possible electrophysiological component of empathy (Mu et al., 2008; Peng et al., 2021) and own pain experience (Misra et al., 2017; Peng et al., 2021; Ploner et al., 2017). Mid-frontal theta is thought to indicate activity in the anterior cingulate cortex (ACC, Mitchell et al., 2008; van der Molen et al., 2017), which is often reported in fMRI studies on empathy for pain and is associated with the negative affect during own and others' pain (Fallon et al., 2020).

Finally, several studies suggest that physiological "coupling" (in cardiac activity or skin conductance), i.e., aligning to the physiological state of someone in pain, might reflect empathic sharing and facilitate understanding of the other (Goldstein et al., 2017; Jospe et al., 2020; Reddan et al., 2020; Zerwas et al., 2021).

Humans respond to social cues such as facial expressions or eye contact by sharing the other's emotional state (Ensenberg et al., 2017; Hess, 2021; Perry et al., 2010b). We thus expected that reduced availability

of social cues and thereby reduced intimacy of the interaction would reduce affective resonance and impair

understanding of others. To test the effect of intimacy on empathy for pain in the current study, pairs of partici-

pants underwent an empathy-for-pain paradigm in two conditions: one direct, face-to-face interaction and one

interaction mediated via real-time video transfer. In both conditions, one participant (the target) received painful

electrical stimulation to the back of the right hand while the other (the observer) was watching. As both observer

and target were able to see each other in both conditions, their behaviours could subtly influence each other.

Therefore, the task can be considered an interactive empathy task (Cañigueral et al., 2022; Cui et al., 2013).

The task of the observer was to judge the pain intensity experienced by the target and her/his own feelings of

unpleasantness while watching the others' pain. As targets themselves also rated their experienced pain inten-

sity, the correspondence between targets' and observers pain provided a behavioural read-out for empathic

accuracy for pain. In addition, we measured EEG from the observer and cardiac and skin conductance re-

sponses from both members of the dyad. Contrary to our hypothesis, we found that the reduced level of intimacy

in mediated compared to direct interactions did not diminish empathic accuracy (Agahi & Wanic, 2020; Jospe et

al., 2020; Zaki et al., 2009a). While pain empathy did not modulate mu suppression, theta responses to another's

pain were reduced in mediated interactions. Finally, physiological responses to another's pain were not affected

by social presence, but physiological coupling within the dyad was reduced in the mediated compared to the

direct interaction.

Results

All dyads experienced both conditions, the direct condition, sitting face-to-face in the same room, and the medi-

ated condition, where they sat in different rooms and saw each other via a real-time video transfer. In each

condition, observers watched the target receiving 80 painful electrical shocks of varying intensity (20 levels from

"perceivable" to "extremely painful") and were instructed to judge how painful these shocks were for the target.

The targets rated how painful the shocks were for themselves. Before the two interactive conditions, observers

received 80 painful electrical shocks themselves (own pain condition), to become familiar with the experience

of electrical shocks, and to asses EEG responses to own pain (see Fig. 1). Dyads consisted of one of five

targets, who came to the lab six times to interact with a different observer each time, and one out of 30 observ-

ers.The results are thus based on data from 30 dyads. For all response variables, we hypothesized that they

would be predicted by others' pain intensity, and that this effect would show an interaction with the condition, if

social presence modulated pain empathy.

Figure 1. (A) Schematic overview of experimental conditions. (B) Overview of the analysed outcomes.

Social presence does not influence empathic accuracy

To assess effects of social presence on empathic accuracy, observers' ratings of targets' pain were predicted

from shock intensity, condition ("direct" vs. "mediated") and their interaction using generalized linear mixed mod-

els with single trials nested in observers and observers nested in targets. The best model had a random-effects

structure containing random slopes for *condition* and *intensity* (AIC difference to the next best model during

fitting of random effects: -63). Moreover, it contained a fixed effect of *intensity* (b(SE) = 0.07(0.01), z = 10.24, p

< 0.0001, AIC difference to a model without the fixed effect of intensity: -14). The Bayes factor for the hypothe-

sized interaction of condition and intensity was 0.075, indicating strong evidence for the model without the inter-

action (Lee & Wagenmakers, 2014). This shows that the intensity of the shocks received by the targets predicted

the observers' pain ratings (Fig. 2A & B), indicating meaningful empathic accuracy. However, this effect did not

differ between conditions.

Social presence increases unpleasantness ratings

In addition to the pain intensity ratings, observers also rated how unpleasant it was for them to watch

the target receiving the painful shocks. We used a generalized linear mixed model to assess whether shock

intensity and condition ("direct" vs. "mediated") predicted the unpleasantness ratings. The best model had a

random-effects structure containing random slopes for the interaction between *condition* and *intensity* (AIC dif-

ference to the next best model during fitting of random effects: -29). It contained a fixed effect of *intensity* (b(SE)

= 0.07(0.01), *z* = 8.07, *p* < 0.0001) and a fixed effect of *condition* (b(SE) = -0.14(0.05), *z* = -2.63, *p* = 0.009, AIC

difference to a model without the fixed effect of *condition*: -4). The Bayes factor for the hypothesized interaction

of condition and intensity was 0.021, indicating very strong evidence for the model without the interaction (Lee

& Wagenmakers, 2014). This shows that the intensity of the shocks received by the targets predicted the ob-

servers' unpleasantness ratings (Fig. 2C), but equally so in both conditions. However, the observers rated the

pain of the target as slightly more unpleasant in the direct than in the mediated interaction (mean difference (SD)

= 1.47(6.45)).

Figure 2. (A) Example pain rating data over trials from one randomly chosen sample dyad. (B) Predicted observer pain ratings for direct- and mediated-interaction conditions. The black thick line represents the fixed effect of shock intensity; single coloured lines represent predicted data from single participants. The colour shading from blue to red represents the value of the random slope for intensity of each participant, and the solid and dotted lines show predicted ratings for direct and mediated condition, respectively. (C) Predicted observer valence ratings in direct and mediated interaction. The black thick line represents the fixed effect of shock intensity; single coloured lines represent predicted data from single participants. The colour shading from blue to red represents the value of the random slope for the interaction between intensity and condition of each participant. The boxplots represent summary statistics for each condition. * = $p < 0.01$.

Mu suppression is not sensitive to observed pain

Using a permutation test, we analysed whether mu suppression was sensitive to the observed pain intensity, and whether it was reduced in the mediated condition. For each participant, we chose an individualized frequency (in the range of 8 – 13 Hz) and electrode (of central electrodes), which showed the strongest desynchronization in the own pain condition. Data from 29 dyads were included in these analyses, as data from the own

pain condition of one participant were not analysable due to technical issues during data collection. The analyses

of peak mu suppression in the "direct" and "mediated interaction" conditions showed no effect of *intensity* or

condition, nor an interaction that was significant after maximum value correction (see Fig. 3A & 3B middle and

right). This shows that there was no mu suppression which was sensitive to others' pain intensity in either con-

dition. Similarly, analysing averaged power over the canonical mu band (8–13 Hz) yielded no significant effect

of either factor (*intensity*: b(SE) = 0.34(0.30), *t*(df) =   1.13(87), *p* = 0.26; *condition*: b(SE) = -0.04(0.30), t(df) =

-0.14(87), *p* = 0.88; *interaction intensity*condition*: b(SE) = -0.02(0.42), *t*(df) = -0.04(87), *p* = 0.97). However,

peak mu significantly differed between pain levels when participants experienced pain themselves (Fig. 3A left).

Figure 3. (A) Shown is peak mu power averaged over participants, dichotomized into low- and high-intensity shocks for display purposes only. Blue solid lines indicate time windows where the main effect of intensity reached significance (uncorrected level); blue dotted lines indicate time windows where the intensity effect survived maximum value correction. The main effect of condition and the interaction of condition x intensity did not survive maximum value correction, at any time-point. On the left side, the clear effect of pain intensity on mu power in the own-pain condition can be assessed. (B) Shown is the topography of averaged peak mu power differences between high- and low-intensity shocks in the three conditions. (C) Shown are boxplots for theta power averaged over 4–8 Hz and 0–500 ms after shock onset at electrode Fz for the direct and mediated conditions. Grey lines depict means from single participants. Significance asterisks refer to post-hoc tests from linear mixed models. * = $p < 0.05$, ** = $p < 0.001$. (D) Shown is the time course of averaged theta power (4–8 Hz) after stimulus onset. (E) Shown is the topography of averaged theta power differences between high- and low-intensity shocks in the two conditions.

Sensitivity of theta power to others' pain is modulated by social presence

We then tested whether theta responses were sensitive to observed pain intensity, and whether this was modulated by social presence. Data from 30 dyads were included in these analyses. The best-fitting linear mixed model for the power averaged over the canonical theta band (4–8 Hz) yielded a significant main effect of *intensity* (b(SE) = 0.57(0.14), t(df) = 3.99(90), $p < 0.001$) and a significant interaction between *condition* and *intensity* (b(SE) = -0.46(0.20), t(df) = -2.30(90), $p = 0.024$, AIC difference to next best model: 2.9). Follow-up models on the significant interaction showed a significant effect of *intensity* in the direct condition (b(SE) = 0.57(0.13), t(df) = 4.46(28.99), $p < 0.001$), but not in the mediated condition (b(SE) = 0.10(0.15), t(df) = 0.70(28.99), $p = 0.49$). This indicated that frontal theta was more responsive to the other's shock intensity in the direct condition than in the mediated condition (see Fig. 3C, D, E).

Exploratory analyses of whole time-frequency-electrode space

To test for modulations of neural activity by observed shock intensity or social presence beyond mu and theta band, we ran a cluster based permutation test on the whole time-frequency-electrode data space. We used a cluster size correction and a maximum value correction to correct for multiple comparisons. Data from 30 dyads were included in these analyses. During both direct and mediated interaction, others' pain *intensity* was positively related to 2 to 6 Hz power between 56 and 840 ms after shock onset at the frontal and central electrodes (cluster 1, maximal at F4, see Fig. 4A left, B and E top left). Others' pain *intensity* was also negatively related to 12 to 20 Hz power between 680 and 1040 ms after shock onset over parietal and occipital electrodes (cluster 2, maximal at P2, see Fig. 4A left, B and E bottom left).

Neither a permutation test on the *condition* effect (mediated vs. direct) nor the interaction between *condition* and *intensity* yielded any cluster that survived the cluster size correction or the maximum value correction.

However, due to the exploratory nature of the analyses, we further examined the biggest cluster, which was

significant on an uncorrected level. In direct compared to mediated interactions (main effect of condition; see

Fig. 4A middle, C and E middle), theta/alpha (5–12 Hz) power at the frontal electrodes and lower beta (13–20

Hz) power at the frontal, central and centro-parietal electrodes was enhanced in an early time window (-256–

644 ms after shock onset). In a later time window (511–1500 ms after shock onset), alpha/lower beta (8–20 Hz)

power at the centro-parietal electrodes was enhanced during direct interactions. For the interaction between

condition and *intensity*, the biggest uncorrected significant cluster showed a stronger positive effect of *intensity*

in the "direct" than in the "mediated" condition. This cluster spanned 3–20 Hz between -220 and 1380 ms after

shock onset and included most left-hemisphere and central electrodes (see Fig. 4A right, D). Its time course is

displayed separately for low (3–9 Hz, electrode with maximal interaction: Cz, Fig. 4E top right) and high frequen-

cies (10–20 Hz, electrode with maximal interaction: CP1, Fig. 4E bottom right).

Figure 4: (A) Clusters found in the permutation tests on the correlation between shock intensity and EEG power across both conditions (left), on the main effect of condition (middle) and on the interaction between shock intensity (dichotomized for display purposes only) and condition (right), averaged over electrodes. (B) Results of the permutation test on the correlation between shock intensity and EEG power across direct and mediated

conditions. Data points that were significant after cluster size correction are displayed in colour; data points that
were significant after maximum value correction are marked in white. (C) Results of the permutation tests on the
condition effect. Data points of the largest cluster (uncorrected significant) are displayed in colour. (D) Results
of the permutation tests on the interaction between condition and intensity (dichotomized for display purposes
only). Data points of the largest cluster (uncorrected significant) are displayed in colour. (E) Power time course
of cluster 1 (top left) and cluster 2 (bottom left), and power time course for the early cluster (5–20 Hz, F3, middle
top) and the late cluster (8–20 Hz, TP8, middle bottom) showing a condition effect and power time course of
lower frequencies (3–9 Hz; top, right), and higher frequencies (10–20 Hz; bottom, right) for interaction effects
between condition and intensity. Data are dichotomized into low and high intensities for display purposes only.
241

Social presence does not modulate physiological responses to other's pain

Next, we tested whether observers showed stronger cardiac responses (interbeat interval, IBI, responses) to

observed pain in the "direct" than in the "mediated" condition, again using linear mixed models on single trials.

The best model contained random slopes for *condition* and *intensity* but no fixed effects. The Bayes factor for

the hypothesized effect of intensity was 0.027, indicating very strong evidence for the model without the effect.

The Bayes factor for the hypothesized interaction of condition and intensity was 0.088, indicating strong evi-

dence for the model without the interaction (Lee & Wagenmakers, 2014). These results indicate that observers'

IBI responses were not sensitive to the observed shock intensity on the sample level (see Fig. 5A).

Moreover, we tested whether observers' skin conductance responses (SCRs) were more responsive to

observed pain intensity in the "direct" than in the "mediated" condition. Observers' SCRs were greatest in the

time window from 1 to 4 seconds after shock onset, hence this time window was used for all further analyses.

The best single-trial linear mixed model on observers' SCR to the observed shocks contained random slopes

for *condition* and *intensity* and a fixed positive effect of *intensity*. The Bayes factor for the hypothesized interac-

tion of condition and intensity was 0.036, indicating strong evidence for the model without the interaction (Lee &

Wagenmakers, 2014). These results indicate that observers' SCRs were sensitive to the shock intensity, but

equally so in direct and mediated conditions (Fig. 5C and 5D). All model parameters are listed in Table 1. Models

with dichotomized intensities yielded the same results.

Social presence enhances skin conductance coupling between target and observer

Finally, we used a single-trial linear mixed model to test whether the direct interaction enhanced the coupling

between targets and observers IBI responses. To this end, the model tested whether observers' IBI responses

could be predicted from targets' IBI responses and whether this effect was modulated by the condition ("direct"

vs. "mediated" interaction). The best model contained a random slope for *condition* but no fixed effects. The

Bayes factor for the hypothesized effect of targets' IBI was 0.059, indicating strong evidence for the model

without the effect. The Bayes factor for the hypothesized interaction of condition and targets' IBI was 0.076,

indicating strong evidence for the model without the interaction (Lee & Wagenmakers, 2014). These results

indicate that there was no coupling between targets' and observers' IBI responses (Fig. 5B). Comparing the

correlations between the continuous IBI traces in the two conditions showed the same results (correlations were

highest for a lag of 2 seconds).

The same rationale was used to test for effects of condition on coupling between targets' and observers'

skin conductance responses to pain. The best single-trial linear mixed model on SCR coupling – predicting

observers' SCR from targets' SCR – contained a random slope for *condition* and *target SCR* and fixed effects

of *target SCR*, *condition* and their interaction (for model parameters, see Table 1). Follow-up models on the

interaction between *condition* and *target SCR* showed a significant positive effect of *target SCR* on observers'

SCR in the "direct" condition (b(SE) = 0.12(0.03), t = 4.15, p < 0.0001), but not in the "mediated" condition (b(SE)

= 0.03(0.02), t = 1.64, p = 0.1). When comparing the correlation coefficients of targets' and observers' SCRs

between conditions, there was only a marginally significant effect of *condition* (b(SE) = -0.07(0.04), t(df) = -

1.81(26), p = 0.082). These results indicate that coupling between targets' and observers' SCRs was greater in

the direct than the mediated interaction, but that the effect was rather small (see Fig. 5E).

281   - insert table 1 here -

Figure 5: (A) Grand averages of observers' IBI responses in direct and mediated interactions, dichotomized into low- and high-intensity trials for display purposes only. (B) Observer IBI responses predicted from the single-trial model on physiological coupling. The thick black line represents the fixed effect of target IBI; the thin lines represent predicted data for single participants. (C) Grand averages of observers' SCRs in direct and mediated interactions, dichotomized into low- and high-intensity trials for display purposes only. (D) Observer SCRs predicted from shock intensity in the generalized linear mixed model. The black thick line represents the fixed effect of intensity; the coloured lines display predicted values for single participants. The colour shading from blue to red represents the value of the random slopes of shock intensity for each participant. Solid and dotted lines show predicted values for direct and mediated conditions, respectively. (E) Observer SCRs predicted from target SCRs in the generalized linear mixed model on physiological coupling. The black thick line represents the fixed effect of targets' SCRs; the coloured lines display predicted values for single participants. The colour shading from blue to red represents the value of the random slopes of target SCR for each participant.

Targets do not express their pain differently in the two conditions

To test whether possible differences in empathic responses towards the targets' pain in the two conditions were

due to different expressivity of the targets' in the two conditions, we conducted a control experiment. We showed

100 video segments of targets' experiencing pain from the "direct" and "mediated" conditions to 31 naïve partic-

ipants, who rated the pain intensity of the shocks displayed in the video. These participants were unaware of

the condition manipulation in the videos. If targets' expressed their pain e.g. systematically stronger in the "direct"

than in the "mediated" condition, the new pain ratings should be higher for videos from the "direct" than from the

"mediated" condition. This was not the case. The means of the pain ratings from the video control experiment

did not differ significantly between conditions (mean difference = 0.79, $t(30)$ = 1.66, p = 0.11, Cohen's d = 0.3).

The mean Spearman's correlations between video observers' ratings and shock intensity also did not differ

between videos from direct and mediated interaction (mean$_{direct}$(SD) = 0.36(0.16), mean$_{mediated}$(SD) = 0.32(0.13),

mean difference = 0.040, $t(30)$ = 1.36, p = 0.18). These results indicate that the targets did not express their

pain significantly differently in the direct versus mediated interaction.

311
312
313
Discussion

Although mediated social interactions through video calls are becoming the new norm, the impacts on under-

standing others and their feelings have not yet been researched thoroughly (Grondin et al., 2019), especially in

social neuroscience. In the current study, we explored how a video-mediated interaction affects empathy for

pain on behavioural, physiological and neural levels. We expected that less availability of social cues in a medi-

ated interaction would hamper empathizing with the other. However, we found that observers were just as ac-

curate in judging the other's pain in the mediated interaction as in the direct interaction. Moreover, participants

experienced the other's suffering as only slightly less unpleasant in the mediated interaction. On the neural level,

we did not find mu suppression over the somatosensory cortex that was sensitive to the other's pain in either

condition. However, mid-frontal theta tracked the other's pain intensity more in the direct than in the mediated

interaction. Exploratory analyses of the whole time-frequency-electrode space showed no additional differences

between direct and mediated conditions after correcting for multiple comparisons. On a physiological level, ob-

servers' SCRs were coupled to targets' SCRs to a stronger degree in the direct compared to the mediated

interaction. In sum, behavioural empathy was not reduced in the mediated interaction, whereas some neural

and physiological aspects of empathy were dampened.

Effects of social presence on behavioural aspects of empathy for pain

Surprisingly, among the many studies on empathy for pain, hardly any measured empathic accuracy, and none explored which type of information is necessary or sufficient to judge others' pain accurately (Gauthier et al., 2008; Laursen et al., 2014; Leonard et al., 2013). In story-based empathy studies, empathic accuracy for emotion was reduced when auditory linguistic information was completely removed, whereas missing visual information did not impact empathic accuracy (Jospe et al., 2020; Zaki et al., 2009a). Similarly, we reveal that participants could judge the targets' pain quite well, and this ability did not decline in the mediated interaction. In contrast to the story-based paradigm, our results imply that visual information (apparent in both the direct interaction and the video calls) is sufficient for empathic accuracy for others' pain. As our participants did not experience severe pain (expressed by moaning or crying), auditory information might have been less important than for example in empathic responses to the pain of hospital patients (Agahi & Wanic, 2020). As our control analysis showed no condition differences in target expressivity, we can be assured that these did not mask true condition differences in empathic accuracy.

Although observers showed more affective empathy in the direct than in the mediated condition, the effect size was so small that it was practically negligible. One reason for this might be that many of our participants remembered their own recent pain experience and so were empathic to the similar experience of another. This was also stated by many in the debriefing questionnaires. This strategy might have led to imagination of others' pain independent of the medium, causing similar affective empathy (Goubert et al., 2005).

Effects of social presence on neural aspects of empathy for pain

As most former studies on empathy for pain used abstract cues or static pictures, we expected to find even stronger mu suppression in our paradigm using real stimuli and focusing on individual peak-mu frequency (i.e., Perry et al., 2010a; Riečanský & Lamm, 2019; Zebarjadi et al., 2021). Instead, we did not find any mu suppression in response to others' pain. Speculatively, mu suppression is a compensatory mechanism that aids empathy for pain via somatosensory representation of others' pain if insufficient sensory cues are available. Alternatively, it may be a weak signal that requires many repetitions – and stronger pain signals – to find a significant effect. However, if mu suppression is a general empathy mechanism, our analysis should have been sensitive enough to detect it. Therefore, our null findings on mu suppression align with recent criticism of its robustness and validity as a mechanism underlying empathy in general (Hobson & Bishop, 2016).

The mid-frontal theta/delta response constitutes another neural component of empathy for pain that has

so far been rarely examined in EEG studies (but see Mu et al., 2008; Peng et al., 2021). Mid-frontal theta has

been related to the salience, unexpectedness and aversiveness of many different types of stimuli (Cavanagh &

Shackman, 2015; González-Roldan et al., 2011; Güntekin & Başar, 2014). Therefore, the heightened sensitivity

of theta to the other's pain level might indicate that the other's pain elicits more arousal and negative affect in

the direct interaction (Balconi et al., 2009), although this did not result in measurable behavioural differences.

Mid-frontal theta might stem from the ACC, which shows reliable activity to both own and others' pain in fMRI

studies (Cavanagh & Frank, 2014; Fallon et al., 2020). Confirming this assumption with source analyses was

beyond the scope of this paper but could be an important step in future studies. The exploratory single-trial

permutation test confirmed on a trend level the interaction between others' pain intensity and direct versus me-

diated interaction.

Lastly, our exploratory analysis revealed stronger parietal beta suppression relating to stronger ob-

served pain irrespective of the condition. Parietal beta decrease has been linked to attention to affective touch

(von Mohr et al., 2018). Future EEG studies should clarify its role in empathy for pain.

Effects of social presence on physiological coupling

Observers' cardiac activity was not sensitive to others' pain and showed no coupling with targets' cardiac activity.

In contrast, previous empathy studies found cardiac coupling in emotional empathy (Zerwas et al., 2021), espe-

cially when semantic and auditory information was missing (Jospe et al., 2020). The shocks elicited a strong

cardiac acceleration in the targets in our paradigm (data not shown in this paper). Possibly this was not easily

mimicked by observers cardiac activity, which might be one reason for the discrepancies with former studies

(Goldstein et al., 2017).

In contrast, we found coupling in skin conductance responses, and this was the one aspect of pain

empathy that was markedly reduced in the mediated interaction. This indicates that the physiological coupling

component of empathy, specifically in SCR, might rely on physical proximity (Chatel-Goldman et al., 2014; Mu-

rata et al., 2020) and possibly olfactory cues that are missing in mediated interactions (Calvi et al., 2020; de

Groot et al., 2014).

Conclusions and future directions

Many recent studies examined direct interactions between participants, claiming that this is necessary for understanding social cognition (Fan et al., 2021; Levy et al., 2021; Redcay & Schilbach, 2019). However, few studies have explicitly compared these new paradigms to similar tasks using mediated interactions (but see e.g. Hietanen et al., 2020). Hence, it remains unclear whether the degree of social presence affects social cognition and if these effects are due to the interactivity (here called "immediacy") or to the amount of information transferred and the shared physical space (called "intimacy") (Cui et al., 2013; Grondin et al., 2019). Therefore, by examining the impact of social presence on empathy for the first time, we add a potentially important dimension to the study of social cognition. We show that the effects are nuanced: Only the immediate mid-frontal theta response to others' pain, presumably relating to emotional arousal (Balconi et al., 2009), and SCR coupling were affected by the reduced intimacy. This could indicate that intimacy is especially important for more automatic, stimulus-driven empathy components. Future studies should address whether the immediacy within the interaction might have a stronger impact on all components of empathy (Hamilton & Lind, 2016; Hietanen et al., 2020).

        To conclude, we do not find evidence that empathy for pain is markedly impaired in video-mediated interactions, although physiological and neural resonance with the other's pain was reduced, which implies that some level of synchronization with the other is impaired. This suggests that empathic abilities might be preserved in everyday mediated social interactions, which are becoming more common. By showing specific changes in empathy components in a mediated interaction, we start to fill the gap in knowledge about social presence in social neuroscience.

Limitations of the study

The main limitation of this study is the small sample size, which was due to the complexity of the design. This is especially prominent when reporting mostly null results, as one might argue that our small sample size prevented us from detecting subtle effects of social presence. However, as power analyses showed, by analysing single trials and using a within-subject design, we had sufficient power to detect meaningful effects of social presence. Moreover, the small number of men in the sample prevented us from assessing effects of gender. However, even if there are differences in empathy between female-female and female-male dyads, effects of social presence on it should still be detectable in our within-subjects design. Another limitation is the comparably low standardization of our laboratory task. Using a task with real live people, we aimed to capture real-life empathy for pain in the best way possible in the EEG laboratory. At the same time, by using a standardized pain-

419   stimulation protocol, we maintained a high degree of standardization compared to studies using unstructured

420   interactive paradigms (i.e., Levy et al., 2017). Our study therefore answers recent calls for more interactive and

421   contextual experimental methods for researching social interaction (Dumas, 2011; Przyrembel et al., 2012; Son-

422   kusare et al., 2019).

Acknowledgements:

UMK is supported by the German Science Foundation (grant number KR3691/8-1). We thank Lou Maria Lütjo-

hann, Celina Mävers, Marthe Mieling, Leah Reinicke, Ellyn Sänger and Jasmin Thurley for help with data col-

lection and pre-processing, Charlotte Petereit for help with the figures, and two anonymous reviewers for their

helpful comments.

Author contributions: Conceptualization: P.P. and U.M.K; Methodology: P.P. and U.M.K; Software: P.P; For-

mal Analysis: P.P.; Investigation: P.P.; Data Curation: P.P.; Writing- Original Draft: P.P; Writing- Review & Edit-

ing: P.P., R.W., A.P and U.M.K.; Visualization: P.P.; Supervision: U.M.K.; Project Administration: P.P. and

U.M.K; Funding Acquisition: U.M.K

Declaration of interests: "The authors declare no competing interests."

479   **References**

480   Abler, B., & Kessler, H. (2009). Emotion Regulation Questionnaire – Eine deutschsprachige Fassung des ERQ

481   von Gross und John. *Diagnostica*, *55*(3), 144–152. https://doi.org/10.1026/0012-1924.55.3.144

482   Agahi, S., & Wanic, R. (2020). Supremacy of auditory versus visual Input in somatic Empathy and perceived

483   Pain Level. *Pain Management Nursing: Official Journal of the American Society of Pain Management*

484   *Nurses*, *21*(2), 201–206. https://doi.org/10.1016/j.pmn.2019.06.013

485   Akaike, H. (1998). Information Theory and an Extension of the Maximum Likelihood Principle. In E. Parzen, K.

486   Tanabe, & G. Kitagawa (Eds.), *Selected Papers of Hirotugu Akaike* (pp. 199–213). Springer.

487   https://doi.org/10.1007/978-1-4612-1694-0_15

488   Balconi, M., Brambilla, E., & Falbo, L. (2009). Appetitive vs. Defensive responses to emotional cues. Autonomic

489   measures   and   brain   oscillation   modulation. *Brain   Research*, *1296*, 72–84.

490   https://doi.org/10.1016/j.brainres.2009.08.056

491   Bates, D., Maechler, M., Bolker, B., Walker, S., Bojesen Christensen, R. H., Singmann, H., Da, B., Scheipl, F.,

492   Grothendieck, G., Green, P., Fox, J., Bauer, A., & Krivitsky, P. N. (2020). *Linear Mixed-Effects Models*

493   *using "Eigen" and S4* (1.1-25) [R]. https://github.com/lme4/lme4/

494   Benedek, M., & Kaernbach, C. (2010). Decomposition of skin conductance data by means of nonnegative de-

495   convolution. *Psychophysiology*, *47*(4), Article 4. https://doi.org/10.1111/j.1469-8986.2009.00972.x

496   Biocca, F., Harms, C., & Burgoon, J. K. (2003). Toward a More Robust Theory and Measure of Social Presence:

497   Review and Suggested Criteria. *Presence: Teleoperators and Virtual Environments*, *12*(5), Article 5.

498   https://doi.org/10.1162/105474603322761270

499   Burnham, K. P., & Anderson, D. R. (2004). Multimodel Inference: Understanding AIC and BIC in Model Selec-

500   tion. *Sociological Methods & Research*, *33*(2), Article 2. https://doi.org/10.1177/0049124104268644

501   Calvi, E., Quassolo, U., Massaia, M., Scandurra, A., D'Aniello, B., & D'Amelio, P. (2020). The scent of emotions:

502   A systematic review of human intra- and interspecific chemical communication of emotions. *Brain and*

503   *Behavior*, *10*(5), e01585. https://doi.org/10.1002/brb3.1585

504   Cañigueral, R., Krishnan-Barman, S., & Hamilton, A. F. de C. (2022). Social signalling as a framework for sec-

505   ond-person neuroscience. *Psychonomic Bulletin & Review*. https://doi.org/10.3758/s13423-022-02103-

506   2

507   Cavanagh, J. F., & Frank, M. J. (2014). Frontal theta as a mechanism for cognitive control. *Trends in Cognitive*

508   *Sciences*, *18*(8), 414–421. https://doi.org/10.1016/j.tics.2014.04.012

Cavanagh, J. F., & Shackman, A. J. (2015). Frontal Midline Theta reflects Anxiety and cognitive Control: Meta-

analytic   Evidence.   *Journal   of   Physiology,   Paris*,   *109*(0),   3–15.

https://doi.org/10.1016/j.jphysparis.2014.04.003

Chatel-Goldman, J., Congedo, M., Jutten, C., & Schwartz, J.-L. (2014). Touch increases autonomic coupling

between   romantic   partners.   *Frontiers   in   Behavioral   Neuroscience*,   *8*.

https://doi.org/10.3389/fnbeh.2014.00095

Cheng, Y., Yang, C.-Y., Lin, C.-P., Lee, P.-L., & Decety, J. (2008). The perception of pain in others suppresses

somatosensory oscillations: A magnetoencephalography study. *NeuroImage*, *40*(4), 1833–1840.

https://doi.org/10.1016/j.neuroimage.2008.01.064

Cohen, M. X. (2014). *Analyzing Neural Time Series Data: Theory and Practice* (1st ed.). The MIT Press.

Corcoran, A. W., Alday, P. M., Schlesewsky, M., & Bornkessel-Schlesewsky, I. (2018). Toward a reliable, auto-

mated method of individual alpha frequency (IAF) quantification. *Psychophysiology*, *55*(7), e13064.

https://doi.org/10.1111/psyp.13064

Cui, G., Lockee, B., & Meng, C. (2013). Building modern online social presence: A review of social presence

theory and its instructional design implications for future trends. *Education and Information Technolo-

gies*, *18*(4), 661–685. https://doi.org/10.1007/s10639-012-9192-1

de Groot, J. H. B., Semin, G. R., & Smeets, M. A. M. (2014). I can see, hear, and smell your fear: Comparing

olfactory and audiovisual media in fear communication. *Journal of Experimental Psychology: General*,

143(2), 825–834. https://doi.org/10.1037/a0033731

Decety, J., & Jackson, P. L. (2004). The Functional Architecture of Human Empathy. *Behavioral and Cognitive

Neuroscience Reviews*, *3*(2), 31. https://doi.org/10.1177/1534582304267187

Delorme, A., & Makeig, S. (2004). EEGLAB: An open source toolbox for analysis of single-trial EEG dynamics

including independent component analysis. *Journal of Neuroscience Methods*, *134*(1), 9–21.

https://doi.org/10.1016/j.jneumeth.2003.10.009

Dumas, G. (2011). Towards a two-body neuroscience. *Communicative & Integrative Biology*, *4*(3), 349–352.

https://doi.org/10.4161/cib.4.3.15110

Einthoven, W., Fahr, G., & De Waart, A. (1950). On the direction and manifest size of the variations of potential

in the human heart and on the influence of the position of the heart on the form of the electrocardiogram.

American Heart Journal, *40*(2), 163–211. https://doi.org/10.1016/0002-8703(50)90165-7

Ensenberg, N. S., Perry, A., & Aviezer, H. (2017). Are you looking at me? Mu suppression modulation by facial

539   expression direction. *Cognitive, Affective & Behavioral Neuroscience*, *17*(1), 174–184.

540   https://doi.org/10.3758/s13415-016-0470-z

541   Fallon, N., Roberts, C., & Stancak, A. (2020). Shared and distinct functional networks for empathy and pain

542   processing: A systematic review and meta-analysis of fMRI studies. *Social Cognitive and Affective Neu-*

543   *roscience*, *15*(7), 709–723. https://doi.org/10.1093/scan/nsaa090

544   Fan, S., Dal Monte, O., & Chang, S. W. C. (2021). Levels of naturalism in social neuroscience research. *ISci-*

545   *ence*, *24*(7), 102702. https://doi.org/10.1016/j.isci.2021.102702

546   Gallo, S., Paracampo, R., Müller-Pinzler, L., Severo, M. C., Blömer, L., Fernandes-Henriques, C., Henschel, A.,

547   Lammes, B. K., Maskaljunas, T., Suttrup, J., Avenanti, A., Keysers, C., & Gazzola, V. (2018). The causal

548   role   of   the   somatosensory   cortex   in   prosocial   behaviour.   *ELife*,   *7*,   e32740.

549   https://doi.org/10.7554/eLife.32740

550   Gauthier, N., Thibault, P., & Sullivan, M. J. L. (2008). Individual and relational correlates of pain-related empathic

551   accuracy   in   spouses   of   chronic   pain   patients.   *The   Clinical   Journal   of   Pain*,   *24*(8),   669–677.

552   https://doi.org/10.1097/AJP.0b013e318173c28f

553   Goldstein, P., Weissman-Fogel, I., Dumas, G., & Shamay-Tsoory, S. G. (2018). Brain-to-brain coupling during

554   handholding is associated with pain reduction. *Proceedings of the National Academy of Sciences*,

555   *115*(11), E2528–E2537. https://doi.org/10.1073/pnas.1703643115

556   Goldstein, P., Weissman-Fogel, I., & Shamay-Tsoory, S. G. (2017). The role of touch in regulating inter-partner

557   physiological coupling during empathy for pain. *Scientific Reports*, *7*. https://doi.org/10.1038/s41598-

558   017-03627-7

559   González-Roldan, A. M., Martínez-Jauand, M., Muñoz-García, M. A., Sitges, C., Cifre, I., & Montoya, P. (2011).

560   Temporal dissociation in the brain processing of pain and anger faces with different intensities of emo-

561   tional expression. *PAIN*, *152*(4), 853–859. https://doi.org/10.1016/j.pain.2010.12.037

562   Goubert, L., Craig, K. D., & Buysse, A. (2009). Perceiving Others in Pain: Experimental and Clinical Evidence

563   on the Role of Empathy. In J. Decety & W. Ickes (Eds.), *The Social Neuroscience of Empathy* (pp. 153–

564   166). The MIT Press. https://doi.org/10.7551/mitpress/9780262012973.003.0013

565   Goubert, L., Craig, K. D., Vervoort, T., Morley, S., Sullivan, M. J. L., Williams, de C. A. C., Cano, A., & Crombez,

566   G.   (2005).   Facing   others   in   pain: The   effects   of   empathy: *Pain*,   *118*(3),   285–288.

567   https://doi.org/10.1016/j.pain.2005.10.025

568   Green, P., & MacLeod, C. J. (2016). SIMR: An R package for power analysis of generalized linear mixed models

569   by simulation. *Methods in Ecology and Evolution*, *7*(4), Article 4. https://doi.org/10.1111/2041-

570   210X.12504

571   Grondin, F., Lomanowska, A. M., & Jackson, P. L. (2019). Empathy in computer-mediated interactions: A con-

572   ceptual framework for research and clinical practice. *Clinical Psychology: Science and Practice*, *26*(4),

573   17–17. https://doi.org/10.1111/cpsp.12298

574   Gross, J. J., & John, O. P. (2003). Individual differences in two emotion regulation processes: Implications for

575   affect, relationships, and well-being. *Journal of Personality and Social Psychology*, *85*(2), 348–362.

576   https://doi.org/10.1037/0022-3514.85.2.348

577   Güntekin, B., & Başar, E. (2014). A review of brain oscillations in perception of faces and emotional pictures.

578   *Neuropsychologia*, *58*, 33–51. https://doi.org/10.1016/j.neuropsychologia.2014.03.014

579   Hamilton, A. F. de C., & Lind, F. (2016). Audience effects: What can they tell us about social neuroscience,

580   theory of mind and autism? *Culture and Brain*, *4*(2), 159–177. https://doi.org/10.1007/s40167-016-0044-

581   5

582   Hess, U. (2021). Who to whom and why: The social nature of emotional mimicry. *Psychophysiology*, *58*(1),

583   e13675. https://doi.org/10.1111/psyp.13675

584   Hietanen, J. O., Peltola, M. J., & Hietanen, J. K. (2020). Psychophysiological responses to eye contact in a live

585   interaction and in video call. *Psychophysiology*, *57*(6), e13587. https://doi.org/10.1111/psyp.13587

586   Hobson, H. M., & Bishop, D. V. M. (2016). Mu suppression – A good measure of the human mirror neuron

587   system? *Cortex*, *82*, 290–310. https://doi.org/10.1016/j.cortex.2016.03.019

588   Jospe, K., Genzer, S., Klein Selle, N., Ong, D., Zaki, J., & Perry, A. (2020). The contribution of linguistic and

589   visual cues to physiological synchrony and empathic accuracy. *Cortex; a Journal Devoted to the Study*

590   *of the Nervous System and Behavior*, *132*, 296–308. https://doi.org/10.1016/j.cortex.2020.09.001

591   Lamm, C., Nusbaum, H. C., Meltzoff, A. N., & Decety, J. (2007). What Are You Feeling? Using Functional Mag-

592   netic Resonance Imaging to Assess the Modulation of Sensory and Affective Responses during Empa-

593   thy for Pain. *PLoS ONE*, *2*(12), e1292. https://doi.org/10.1371/journal.pone.0001292

594   Laursen, H. R., Siebner, H. R., Haren, T., Madsen, K., Grønlund, R., Hulme, O., & Henningsson, S. (2014).

595   Variation in the oxytocin receptor gene is associated with behavioral and neural correlates of empathic

596   accuracy. *Frontiers in Behavioral Neuroscience*, *8*. https://doi.org/10.3389/fnbeh.2014.00423

597   Lawless, J. F. (1987). Negative binomial and mixed poisson regression. *Canadian Journal of Statistics*, *15*(3),

598   209–225. https://doi.org/10.2307/3314912

599   Lee, M. D., & Wagenmakers, E.-J. (2014). *Bayesian Cognitive Modeling: A Practical Course*. Cambridge Uni-

600   versity Press.

601   Leonard, M. T., Issner, J. H., Cano, A., & Williams, A. M. (2013). Correlates of Spousal Empathic Accuracy for

602   Pain-related   Thoughts   and   Feelings.   *The   Clinical   Journal   of   Pain*,   *29*(4),   324–333.

603   https://doi.org/10.1097/AJP.0b013e3182527bfd

604   Levy, J., Goldstein, A., & Feldman, R. (2017). Perception of social synchrony induces mother-child gamma

605   coupling in the social brain. *Social Cognitive and Affective Neuroscience*, *12*(7), 1036–1046.

606   https://doi.org/10.1093/scan/nsx032

607   Levy, J., Lankinen, K., Hakonen, M., & Feldman, R. (2021). The integration of social and neural synchrony: A

608   case for ecologically valid research using MEG neuroimaging. *Social Cognitive and Affective Neurosci-*

609   *ence*, *16*(1–2), 143–152. https://doi.org/10.1093/scan/nsaa061

610   Lopez-Calderon, J., & Luck, S. J. (2014). ERPLAB: An open-source toolbox for the analysis of event-related

611   potentials. *Frontiers in Human Neuroscience*, *8*. https://doi.org/10.3389/fnhum.2014.00213

612   Misra, G., Wang, W., Archer, D. B., Roy, A., & Coombes, S. A. (2017). Automated classification of pain percep-

613   tion using high-density electroencephalography data. *Journal of Neurophysiology*, *117*(2), 786–795.

614   https://doi.org/10.1152/jn.00650.2016

615   Mitchell, D. J., McNaughton, N., Flanagan, D., & Kirk, I. J. (2008). Frontal-midline theta from the perspective of

616   hippocampal "theta." *Progress in Neurobiology*, *86*(3), 156–185. https://doi.org/10.1016/j.pneuro-

617   bio.2008.09.005

618   Mu, Y., Fan, Y., Mao, L., & Han, S. (2008). Event-related theta and alpha oscillations mediate empathy for pain.

619   *Brain Research*, *1234*, 128–136. https://doi.org/10.1016/j.brainres.2008.07.113

620   Murata, A., Nishida, H., Watanabe, K., & Kameda, T. (2020). Convergence of physiological responses to pain

621   during face-to-face interaction. *Scientific Reports*, *10*(1), Article 1. https://doi.org/10.1038/s41598-019-

622   57375-x

623   Peled-Avron, L., Goldstein, P., Yellinek, S., Weissman-Fogel, I., & Shamay-Tsoory, S. G. (2018). Empathy dur-

624   ing consoling touch is modulated by mu-rhythm: An EEG study. *Neuropsychologia*, *116*, 68–74.

625   https://doi.org/10.1016/j.neuropsychologia.2017.04.026

Peng, W., Lou, W., Huang, X., Ye, Q., Tong, R. K.-Y., & Cui, F. (2021). Suffer together, bond together: Brain-

to-brain synchronization and mutual affective empathy when sharing painful experiences. *NeuroImage*,

238, 118249. https://doi.org/10.1016/j.neuroimage.2021.118249

Perry, A., Bentin, S., Bartal, I. B.-A., Lamm, C., & Decety, J. (2010). "Feeling" the pain of those who are different

from us: Modulation of EEG in the mu/alpha range. *Cognitive, Affective, & Behavioral Neuroscience*,

10(4), 493–504. https://doi.org/10.3758/CABN.10.4.493

Perry, A., Troje, N. F., & Bentin, S. (2010). Exploring motor system contributions to the perception of social

information: Evidence from EEG activity in the mu/alpha frequency range. *Social Neuroscience*, *5*(3),

272–284. https://doi.org/10.1080/17470910903395767

Ploner, M., Sorg, C., & Gross, J. (2017). Brain Rhythms of Pain. *Trends in Cognitive Sciences*, *21*(2), 100–110.

https://doi.org/10.1016/j.tics.2016.12.001

Przyrembel, M., Smallwood, J., Pauen, M., & Singer, T. (2012). Illuminating the dark matter of social neurosci-

ence: Considering the problem of social interaction from philosophical, psychological, and neuroscien-

tific perspectives. *Frontiers in Human Neuroscience*, *6*. https://doi.org/10.3389/fnhum.2012.00190

Redcay, E., & Schilbach, L. (2019). Using second-person neuroscience to elucidate the mechanisms of social

interaction. *Nature Reviews Neuroscience*, *20*(8), 495–505. https://doi.org/10.1038/s41583-019-0179-4

Reddan, M. C., Young, H., Falkner, J., López-Solà, M., & Wager, T. D. (2020). Touch and social support influ-

ence   interpersonal   synchrony   and   pain.   *Social   Cognitive   and   Affective   Neuroscience*.

https://doi.org/10.1093/scan/nsaa048

Riečanský, I., & Lamm, C. (2019). The Role of Sensorimotor Processes in Pain Empathy. *Brain Topography*,

32(6), 965–976. https://doi.org/10.1007/s10548-019-00738-4

Rutgen, M., Seidel, E.-M., Rie ansky, I., & Lamm, C. (2015). Reduction of Empathy for Pain by Placebo Anal-

gesia Suggests Functional Equivalence of Empathy and First-Hand Emotion Experience. *Journal of

Neuroscience*, *35*(23), 8938–8947. https://doi.org/10.1523/JNEUROSCI.3936-14.2015

Shamay-Tsoory, S. G. (2011). The Neural Bases for Empathy. *The Neuroscientist*, *17*(1), 7.

Short, J., Williams, E., & Christie, B. (1976). *The social psychology of telecommunications*. Wiley.

Singer, T., & Lamm, C. (2009). The Social Neuroscience of Empathy. *Annals of the New York Academy of

Sciences*, *1156*(1), 81–96. https://doi.org/10.1111/j.1749-6632.2009.04418.x

Sonkusare, S., Breakspear, M., & Guo, C. (2019). Naturalistic Stimuli in Neuroscience: Critically Acclaimed.

Trends in Cognitive Sciences, *23*(8), Article 8. https://doi.org/10.1016/j.tics.2019.05.004

Sperl, M. F. J., Panitz, C., Hermann, C., & Mueller, E. M. (2016). A pragmatic comparison of noise burst and

electric shock unconditioned stimuli for fear conditioning research with many trials. *Psychophysiology*,

53(9), 1352–1365. https://doi.org/10.1111/psyp.12677

Stropahl, M., Bauer, A.-K. R., Debener, S., & Bleichner, M. G. (2018). Source-Modeling Auditory Processes of

EEG Data Using EEGLAB and Brainstorm. *Frontiers in Neuroscience*, *12*. https://www.frontiersin.org/ar-

ticle/10.3389/fnins.2018.00309

van der Molen, M. J. W., Dekkers, L. M. S., Westenberg, P. M., van der Veen, F. M., & van der Molen, M. W.

(2017). Why don't you like me? Midfrontal theta power in response to unexpected peer rejection feed-

back. *NeuroImage*, *146*, 474–483. https://doi.org/10.1016/j.neuroimage.2016.08.045

von Mohr, M., Crowley, M. J., Walthall, J., Mayes, L. C., Pelphrey, K. A., & Rutherford, H. J. V. (2018). EEG

captures affective touch: CT-optimal touch and neural oscillations. *Cognitive, Affective, & Behavioral*

Neuroscience, *18*(1), 155–166. https://doi.org/10.3758/s13415-017-0560-6

Zaki, J., Bolger, N., & Ochsner, K. (2008). It Takes Two: The Interpersonal Nature of Empathic Accuracy. *Psy-*

chological Science, *19*(4), 399–404. https://doi.org/10.1111/j.1467-9280.2008.02099.x

Zaki, J., Bolger, N., & Ochsner, K. (2009). Unpacking the informational bases of empathic accuracy. *Emotion*,

9(4), 478–487. https://doi.org/10.1037/a0016551

Zaki, J., Weber, J., Bolger, N., & Ochsner, K. (2009). The neural bases of empathic accuracy. *Proceedings of*

the National Academy of Sciences, *106*(27), Article 27. https://doi.org/10.1073/pnas.0902666106

Zebarjadi, N., Adler, E., Kluge, A., Jääskeläinen, I. P., Sams, M., & Levy, J. (2021). Rhythmic Neural Patterns

During Empathy to Vicarious Pain: Beyond the Affective-Cognitive Empathy Dichotomy. *Frontiers in*

Human Neuroscience, *15*, 708107. https://doi.org/10.3389/fnhum.2021.708107

Zerwas, F. K., Springstein, T., Karnilowicz, H. R., Lam, P., Butler, E. A., John, O. P., & Mauss, I. B. (2021). "I

feel you": Greater linkage between friends' physiological responses and emotional experience is asso-

ciated with greater empathic accuracy. *Biological Psychology*, *161*, 108079. https://doi.org/10.1016/j.bi-

opsycho.2021.108079

681
682

687   **Figures**
688
689   **Figure 1.** (A) Schematic overview of experimental conditions. (B) Overview of the analysed outcomes.
690
Figure 2. Social presence does not influence empathic accuracy but increases unpleasantness ratings (A) Ex-
ample pain rating data over trials from one randomly chosen sample dyad. (B) Predicted observer pain ratings
for direct- and mediated-interaction conditions. The black thick line represents the fixed effect of shock intensity;
single coloured lines represent predicted data from single participants. The colour shading from blue to red
represents the value of the random slope for intensity of each participant, and the solid and dotted lines show
predicted ratings for direct and mediated condition, respectively. (C) Predicted observer valence ratings in direct
and mediated interaction. The black thick line represents the fixed effect of shock intensity; single coloured lines
represent predicted data from single participants. The colour shading from blue to red represents the value of
the random slope for the interaction between intensity and condition of each participant. The boxplots represent
summary statistics for each condition. * = $p < 0.01$.

701
Figure 3. Mu suppression is not sensitive to observed pain while sensitivity of theta power to others' pain is
modulated by social presence(A) Shown is peak mu power averaged over participants, dichotomized into low-
and high-intensity shocks for display purposes only. Blue solid lines indicate time windows where the main effect
of intensity reached significance (uncorrected level); blue dotted lines indicate time windows where the intensity
effect survived maximum value correction. The main effect of condition and the interaction of condition x intensity
did not survive maximum value correction, at any time-point. On the left side, the clear effect of pain intensity on
mu power in the own-pain condition can be assessed. (B) Shown is the topography of averaged peak mu power
differences between high- and low-intensity shocks in the three conditions. (C) Shown are boxplots for theta
power averaged over 4–8 Hz and 0–500 ms after shock onset at electrode Fz for the direct and mediated con-
ditions. Grey lines depict means from single participants. Significance asterisks refer to post-hoc tests from linear
mixed models. * = $p < 0.001$. (D) Shown is the time course of averaged theta power (4–8 Hz) after stimulus
onset. (E) Shown is the topography of averaged theta power differences between high- and low-intensity shocks
in the two conditions.

715
Figure 4: Exploratory analyses of whole time-frequency-electrode space (A) Clusters found in the permutation
tests on the correlation between shock intensity and EEG power across both conditions (left), on the main effect
of condition (middle) and on the interaction between shock intensity (dichotomized for display purposes only)
and condition (right), averaged over electrodes. (B) Results of the permutation test on the correlation between
shock intensity and EEG power across direct and mediated conditions. Data points that were significant after
cluster size correction are displayed in colour; data points that were significant after maximum value correction
are marked in white. (C) Results of the permutation tests on the condition effect. Data points of the largest cluster
(uncorrected significant) are displayed in colour. (D) Results of the permutation tests on the interaction between
condition and intensity (dichotomized for display purposes only). Data points of the largest cluster (uncorrected
significant) are displayed in colour. (E) Power time course of cluster 1 (top left) and cluster 2 (bottom left), and
power time course for the early cluster (5–20 Hz, F3, middle top) and the late cluster (8–20 Hz, TP8, middle
bottom) showing a condition effect and power time course of lower frequencies (3–9 Hz; top, right), and higher
frequencies (10–20 Hz; bottom, right) for interaction effects between condition and intensity. Data are dichoto-
mized into low and high intensities for display purposes only.

730
Figure 5: Social presence does not modulate physiological responses to other's pain but enhances skin con-
ductance coupling between target and observer (A) Grand averages of observers' IBI responses in direct and
mediated interactions, dichotomized into low- and high-intensity trials for display purposes only. (B) Observer
IBI responses predicted from the single-trial model on physiological coupling. The thick black line represents the
fixed effect of target IBI; the thin lines represent predicted data for single participants. (C) Grand averages of
observers' SCRs in direct and mediated interactions, dichotomized into low- and high-intensity trials for display
purposes only. (D) Observer SCRs predicted from shock intensity in the generalized linear mixed model. The
black thick line represents the fixed effect of intensity; the coloured lines display predicted values for single
participants. The colour shading from blue to red represents the value of the random slopes of shock intensity
for each participant. Solid and dotted lines show predicted values for direct and mediated conditions, respec-
tively. (E) Observer SCRs predicted from target SCRs in the generalized linear mixed model on physiological
coupling. The black thick line represents the fixed effect of targets' SCRs; the coloured lines display predicted
values for single participants. The colour shading from blue to red represents the value of the random slopes of
shock intensity for each participant.

Tables

Table 1: Results of the single-trial (generalized) linear mixed models on IBI responses and SCRs

model	random slopes	SD	fixed effects	b(SE)	t(df)	p	AIC Δ
observers' IBI							-13
(n=29)	intercept/obs	25.01	intercept	4.33(2.16)	2.00(27.6)	0.055	
	condition /obs	13.00	-				
	intensity/obs	0.80					
	intercept/targ	7.9					
	condition/targ	0.13					
	intensity/targ	0.36					
observers' IBI (coupling with target IBI)							-26
(n=29)	intercept/obs	24.2	intercept	4.696 (2.154)	2.18 (28.9)	<0.038	
	condition/obs	12.93					
	intercept/targ	<0.000001					
	condition /targ	<0.000001					
observers' SCR							-3
	intercept/obs	0.16	intercept	0.34(0.03)	11.8	<0.0001	
(n=27)	condition/obs	0.21	intensity	0.01(0.002)	3.22	0.001	
	intensity/obs	0.007					
	intercept/targ	0.03					
	condition /targ	0.03					
	intensity/targ	0.0009					

observers' SCR (coupling with target SCR)						-28
	intercept/obs	0.10	intercept	0.17(0.05)	3.34	0.0008
	condition /obs	0.19	target SCR	0.12(0.02)	5.85	<0.0001
(n=27)	targ-SCR/obs	0.06	condition	0.18(0.07)	2.81	0.005
	intercept/targ	0.0000	target SCR * condition	-0.08(0.02)	-5.72	<0.0001
	condition/targ	0.05				
	targ SCR/targ	0.02				

748 *Notes: IBI = interbeat interval, SCR = skin conductance response, AIC Δ = difference in Akaike information*
749 *criterion to the next best model. Models on IBI responses are linear mixed models, and models on SCRs are*
750 *generalized linear mixed models (note that the latter do not provide degrees of freedom for fixed effects).*

STAR Methods

Resource availability

Lead contact

Further information and requests for resources should be directed to and will be

fulfilled by the lead contact, Pauline Petereit (pauline.petereit@neuro.uni-luebeck.de).

Materials availability

This study did not create any new unique materials.

Data and code availability

• Behavioral and physiological raw data have been deposited at

https://osf.io/pqmra/?view_only=df6b8e7cd19743d481d86ef7cb83cb83 and are publicly available at the

date of publication. The DOI is listed in the key resources table. Video recording data of the experiments

cannot be shared to guarantee participant anonymity.

• All original code has been deposited at

https://osf.io/pqmra/?view_only=df6b8e7cd19743d481d86ef7cb83cb83 and is publicly available as of

the date of publication. DOIs are listed in the key resources table.

• Any additional information required to reanalyze the data reported in this paper is available

from the lead contact upon request.

Experimental model and subject details

Five female psychology students were recruited as targets. Only females were recruited for the targets to reduce

possible gender effects over the dyads. They were on average 19.8 years old (SD = 0.75). Thirty psychology

students (7 males, 23 females, mean age(SD) = 24.07(4.68) years) were recruited as observers. A priori power

analyses with data simulations using the *simr* package (Green & MacLeod, 2016) in R showed that this sample

size is sufficient to detect a small effect (f_2 = 0.02) of condition on empathic accuracy (measured via an interac-

tion effect between shock intensity and direct vs. mediated condition in a generalized linear mixed model on

single trials; see below for statistical analyses) with a power of at least 0.98 (depending on exact model struc-

ture). Data were simulated using sample parameters (means and variance parameters as well as correlations

between conditions) from a pilot study. We did not conduct a separate power analysis for the analysis of neural

data because effect size estimations for neural effects are difficult. Moreover, the neural analysis was conducted

with the full amount of trials (80 trials) in contrast to the 40 trials used for the behavioural analysis. Hence, the

power calculation for the behavioural analyses is more conservative. For both targets and observers, exclusion

criteria were current psychiatric or cardiovascular and past or current neurological disorders, current or chronic

pain conditions or current pain-medication intake. For one observer, all physiological data from the own-pain

condition had to be excluded because of missing stimulus triggers. Skin conductance and electrocardiogram

(ECG) data from one target (mediated-interaction condition) were missing due to a technical error. Skin con-

ductance data from two observers could not be analysed due to poor data quality. These dyads were excluded

from all analyses of the missing outcome variable. Targets received €10 per hour; observers received course

credit or €10 per hour. All participants provided written informed consent prior to taking part in the study, includ-

ing consent for video-recording them and showing the videos to other work group members, and in case of the

targets, showing the videos to other participants in future studies. The experiment was carried out according to

the Declaration of Helsinki and was approved by the Ethics Committee of the University of Lübeck.

Method details

Experimental design

Each target interacted with six different observers on six different study days. Observers came to the lab once

for one session (two interactions) with one target. For the observers, there were three conditions, which were

implemented in separate blocks (within-subject design, see Fig. 1A). In the "own pain" condition, the observer

was alone in the laboratory and received electric shocks. This ensured that observers knew what the electric

shocks felt like. In the "direct interaction" condition, observer and target sat opposite each other at a table, and

the target received electric shocks while the observer watched. In the "mediated interaction" condition, target

and observer sat in adjacent rooms and saw each other over a real-time video transmission. The video showed

the face, upper body, arms and hands of the target, and targets were instructed to place the hand with the shock

electrode clearly visible in front of them on the table in both conditions. In both conditions, a blue board was

placed behind the target to make the background as similar as possible in the two conditions. Observers watched

the same target in the latter two conditions and rated the observed pain experience of the target. Targets rated

their own pain experience. In all conditions, skin conductance and ECG were recorded from both participants,

and EEG was recorded from the observer. The "own pain" condition was always carried out first, and the order

of "direct interaction" and "mediated interaction" conditions was pseudo-randomized over participants. In the

latter two conditions, targets' and observers' behaviour was video-recorded.

Stimulus calibration

Prior to the pain task, pain stimuli were calibrated to the subjective pain thresholds of the participants (the ob-

servers in the "own pain" condition, and the targets prior to the first interaction condition). To this end, participants

received electric shocks starting from 0 mA, increasing in amplitude in steps of 0.5 mA. They were required to

rate each stimulus on a scale from 0 ("not perceivable") to 8 ("strongest pain imaginable"). As soon as they rated

a stimulus with "7" ("unbearably painful"), stimuli were decreased in amplitude (again in steps of 0.5 mA) until

participants rated the stimulus as "0" or an amplitude of 0 mA was reached. The procedure was then repeated

once more with increasing stimulus intensity. The stimulus intensity that was rated as 1 ("noticeable") in this last

round was used as the lower limit for the stimuli presented during the task, with the stimulus intensity rated as 6

("extremely painful") used as the upper limit. For the pain task itself, the range from the lower limit to the upper

limit was evenly divided into 20 levels, to have a finer variation in pain intensity levels.

Pain task

The pain task itself was adapted from Rutgen and colleagues (2015). Electric shocks were delivered using a DS

5 isolated bipolar constant current stimulator (Digitimer) and a bar electrode (Digitimer, two electrodes with 9-

mm diameter, 30 mm apart) attached to the back of the right hand. The skin under the electrode was treated

with an abrasive paste and conductive gel to reduce the electric resistance of the skin.

Each trial started with an auditory cue lasting 500 ms that did not predict the shock stimulus intensity.

At 1000 ms after the cue, the electric shock was delivered for 500 ms (series of 2-ms electric pulses, interspersed

with approximately 20-ms breaks). After a randomly varying interval (6000–9000 ms), the next trial or the rating

followed. In 50% of the trials, participants were prompted to rate the stimulus by a vocal recording saying,

"Please rate". We included the rating only in 50% of the trials to keep the duration of the experiment feasible.

The rating was given on a tablet computer. Targets rated how painful the electric shock was for themselves on

a visual analogue scale ranging from "not at all painful" to "extremely painful". Observers rated how painful they

thought the electric shock was for the target (on the same visual analogue scale) as well as how unpleasant it

was for them to watch the target receive the electric shock (on a visual analogue scale ranging from "not at all

unpleasant" to "extremely unpleasant"). The latter rating served as a measure of affective empathy. Electric

shocks varied in intensity in 20 steps from the intensity the targets had rated as "noticeable" (intensity level 1)

to the intensity they had rated as "extremely painful" (intensity level 20) during the calibration. There were 80

trials in each condition, and each intensity occurred four times. The order of intensities was pseudo-random

(with no more than four shocks with intensity level higher than 10 or lower than 11 in a row) but fixed for all

participants and conditions. The order of trials that had to be rated was fixed as well. In the "own pain" condition,

the task was the same except that observers rated their own pain experience on the visual analogue scale.

Targets and observers were instructed not to talk or move excessively during the task but were otherwise al-

lowed to express their emotions freely. Observers were instructed to rate the pain of the other as accurately as

possible. Both observers and targets were told who would receive shocks in the task, hence the observers knew

that they would receive shocks in the "own pain" condition, but would not receive any shocks in the other two

conditions.

Experimental procedure

Target selection

Before targets interacted with observers, they came to the laboratory alone to familiarize themselves with the

procedures and the stimuli. In this first session, they did the same pain task as in the main experiment, but with

no other person in the room. As in the main experiment, skin conductance and ECG were recorded. After the

first session, targets decided whether they wanted to participate in the main experiment. Moreover, we used this

first session for target selection, as we aimed to recruit only targets who set the intensity limits of the stimuli

during stimulus calibration to a level that was actually painful. This was important to ensure that we measured

actual pain empathy during the main experiment. We therefore defined the minimum upper intensity limit that

targets had to reach during the first session to be eligible for the full experiment as within +/-1 standard deviation

of the mean upper intensity limit of a pilot study (resulting in a minimum upper intensity limit of 2.5 mA). We

invited seven potential targets to this first session. Due to the criteria, we had to exclude one target, and one

participant dropped out after the first session, which left five targets for the main experiment.

Main experiment

For the main experiment, the observer arrived first in the laboratory. After the informed-consent form was signed,

the EEG, skin conductance and ECG measurement equipment as well as the stimulus electrode were prepared.

The pain-calibration procedure was carried out. After five practice trials to familiarize themselves with the ratings

on the tablet computer, participants did the pain task in the "own pain" condition. Meanwhile, the target arrived

in a different room, responded to questionnaires and was equipped with the electrodes for physiological meas-

urement. As soon as the "own pain" condition was finished, the stimulus electrode was attached to the target's

right hand, and the target underwent the pain-calibration procedure. Meanwhile, the observer completed ques-

tionnaires. When both were finished, the experimental tasks started (see "pain task" above; either "direct inter-

action" or "mediated interaction" first). Between conditions, there was a short break in which the target moved

between rooms (either into our out of the room in which the observer was seated). ECG and skin conductance

cables of the observer were unplugged from the amplifier during moving, and plugged in again before the next

condition started. Data quality was checked again before the second condition started. Afterwards, target and

observer were seated in different rooms again and replied to post-experimental questionnaires. Finally, observ-

ers were debriefed about the aim of the study. Targets were debriefed only after completing all six sessions.

Questionnaires

We assessed participants' age, gender, body weight and height, educational degree, and habits regarding smok-

ing, caffeine consumption and physical activity. After the experiment, we obtained participants' subjective eval-

uation of the experiment and observers' evaluation of the target. They also filled out two personality question-

naires: the Interpersonal Reactivity Index (Davis, 1983; German version: Paulus, 2009) and the Emotion Regu-

lation Questionnaire (Gross & John, 2003; German version: Abler & Kessler, 2009). These data are not further

evaluated here.

Physiological data acquisition

Participants were asked to refrain from smoking, exercise, alcohol and caffeine for at least six hours before the

experiment to prevent these factors from impacting the physiological measurements. EEG data were recorded

with 59 Ag/AgCl electrodes placed on an elastic cap according to the international 10-20-system (using a Brain-

Amp MR plus amplifier, BrainProducts GmbH,). An online reference electrode was placed on the left earlobe,

while an offline reference electrode was placed on the right earlobe. Horizontal and vertical EOG were recorded

with four electrodes placed next to the outer corners of the eyes and above and below the left eye, respectively.

Sampling rate was 500 Hz, and data were recorded with an online high pass filter of 0.016 Hz, a low pass filter

of 48 Hz and a notch filter at 50 Hz. Impedances were kept below 5kΩ.

911   ECG data were recorded with bipolar Ag/AgCl recording electrodes and one reference electrode, using

a 50-Hz notch filter. One of the recording electrodes was placed on the right forearm, the other one on the left

lower calf of the participant (following Einthoven lead II configuration, Einthoven et al., 1950). Skin conductance

was measured with two electrodes placed on the thenar and hypothenar of the left hand, using a 50-Hz notch

filter. An electrode attached to the left forearm served as ground for both ECG and skin conductance, which

were recorded with the same amplifier (BrainAmp ExG, BrainProducts *GmbH*). In the "direct interaction" and

"mediated interaction" conditions, data from observer and target were recorded synchronously by connecting all

amplifiers to the same USB adapter feeding the data into BrainVisionRecorder (version 1.21.0102, Brain-

Products GmbH).

Quantification and statistical analysis

Physiological data processing

EEG data

All pre-processing was done in EEGLAB, version 2020.0 (Delorme & Makeig, 2004), implemented in MATLAB

R2019b (The Mathworks*)*. Data were re-referenced to the right earlobe, and bipolar horizontal and vertical EOG

channels were computed. Consistently bad channels (mean number = 2.01, range = 0 to 8 channels per partic-

ipant and condition) and data segments with large artefacts were removed from the data (resulting in on average

1.6% of removed trials per participant and condition in the final epoched data). A bandpass filter was applied

(finite impulse response filter, lower limit: 1 Hz, upper limit: 40 Hz, filterorder: 16500). Next, independent com-

ponent analysis (ICA; implemented with the *runica* function in EEGLAB) was used for ocular artefact correction.

Independent components that were clearly related to eye blinks or horizontal eye movements based on topog-

raphy and time course were visually detected and removed (ranging from 2 to 6 components per participant).

Afterwards, the weights of the remaining components were projected onto the original unfiltered data (Stropahl

et al., 2018). Channels that had been removed before the ICA were interpolated (spherical interpolation); For

some participants additional bad channels (mean number = 0.44, range = 0 to 3 channels per participant and

condition) had to be interpolated. Data were then filtered with a bandpass filter with a lower limit of 0.2 Hz and

an upper limit of 40 Hz (Finite impulse response filter, filter order = 16500, hamming window). Afterwards, data

were segmented into stimulus-locked epochs of 4500-ms lengths (1000 ms before and 3500 ms after stimulus

onset) and baseline-corrected to 1000 ms before stimulus onset. A voltage threshold (between -70/70 μV and -

100/100 μV) was manually set for each participant in a way that all trials with non-ocular artefacts were removed.

The number of rejected trials varied from 0 to 31% per participant and condition and did not differ between

conditions (M = 13% in all conditions).

For the time-frequency analysis, single-trial data of all electrodes were convolved with a complex Morlet

wavelet as implemented in MATLAB (function *cwt* with parameter specification 'cmor1-1.5'):

945
$$\omega(t) = (\pi f_b)^{-0.5} e^{-2\pi i f_c t} e^{-\frac{t^2}{f_b}},$$

where f_b = 1 is the bandwidth parameter, and f_c = 1.5 is the wavelet center frequency. Specifically, for each

participant, changes in time-varying energy were computed (square of the convolution between wavelet and

signal) in the frequencies (1–40 Hz, linear increase) for the 1500 ms after shock onset. Power values were

converted to decibels with respect to an average baseline from 500 to 50 ms before stimulus onset (Cohen,

2014). For analyses of peak-frequency power (see next paragraph), we subtracted the averaged data from 500

to 50 ms before stimulus onset as a baseline correction.

To analyse mu suppression, we determined the individual peak frequency of mu power for each partic-

ipant by using the *restingIAF* toolbox (Corcoran et al., 2018). Frequency peaks in the range from 8 to 13 Hz

were detected in the baseline data from 1000 to 0 ms before shock onset in the "own pain" condition at central

electrodes (C1, C2, C3, C4, C5, C6, CP1, CP2, CP3, CP4, CP5, CP6, FC1, FC2, FC3, FC4, FC5, FC6). For

each detected frequency peak, the difference in average power between baseline (-1000 to 0 ms) and stimula-

tion (0 to 1000 ms after shock onset) was calculated. The electrode and corresponding peak frequency with the

strongest shock-related desynchronization for each participant was chosen for all further analyses. In 16 partic-

ipants, a clear peak frequency was detectable. Peaks occurred at all possible frequencies except 9 Hz, and at

the following electrodes: C1, C4, C3, CP2, CP3, C6, CP6, FC6, and FC2. The remaining 13 participants did not

show a peak frequency, and for them data from the most frequent peak frequency and electrode were used (11

Hz at C4).

ECG data

ECG data were loaded into EEGLAB, filtered with a bandpass filter (lower: 1 Hz, upper: 30 Hz, finite impulse

response filter, filterorder = 8250) and segmented into epochs of 2 s before and 8 s after stimulus onset for the

stimulus-locked analyses. The MATLAB function *findpeaks* was used to detect the r-peaks in the segmented as

well as the continuous data (for additional analyses of physiological coupling). Afterwards, data were visually

screened for wrongly assigned or missing r-peaks. Data sections containing extrasystoles or otherwise unde-

tectable R-peaks were treated as missing values. The interbeat interval (IBI) in ms was calculated for every pair

of heartbeats and used as IBI value for each original data point in between the two heartbeats. In this way, the

IBI trace had the same time resolution as the original data. For the analysis of condition differences in IBI re-

sponses to shocks, the mean IBI from 2 to 5 s after cue onset (Sperl et al., 2016) was computed and baseline-

corrected to the mean of the 2 s before cue onset.

Skin conductance data

Skin conductance responses (SCRs) were analysed using the Ledalab-Toolbox (Benedek & Kaernbach, 2010,

version 3.4.8) in Matlab. Data were downsampled to 50 Hz, smoothed with an adaptive gaussian window and

visually screened for strong artefacts, which were spline-interpolated. Afterwards, a continuous deconvolution

analysis was conducted to separate phasic from tonic skin conductance activity (Benedek & Kaernbach, 2010).

In the following, the mean phasic driver activity 1–4 s after shock onset was used for analyses (for targets' SCR

in physiological coupling). For the observer data (condition differences and physiological coupling), four different

time lags (1–4 s) were considered to account for a time lag in the observer's response to the target's pain

expression.

Statistical analyses

We first outline the general statistical analysis approach before specifying the details. One set of statistical anal-

yses examined the observers' responses to the targets' pain (empathic accuracy, unpleasantness ratings, neural

and physiological responses; see Fig. 1B, left side). In these analyses, the predictors, shock intensity, condition

(direct vs. mediated) and their interaction were tested. A significant effect of shock intensity indicates that the

observer's responses are influenced by the other's pain and are therefore interpreted as empathic. A significant

effect of condition indicates that social presence generally changes the observers' behaviour and physiology,

whereas an interaction between shock intensity and condition indicates that social presence alters the sensitivity

to another's pain. The second set of analyses examined physiological coupling between observer and target

responses (see Fig. 1B, bottom). In these analyses, the targets' responses, the condition (direct vs. mediated)

and their interaction served as predictors. A significant effect of targets' responses indicates that observers' and

targets' responses are generally coupled, whereas an interaction with condition indicates stronger coupling in

one condition compared to the other.

In both sets of analyses, (generalized) linear mixed models with single trials (Level 1) nested in observ-

ers (Level 2) and observers nested in targets (Level 3) were calculated. For theta activity, responses were

averaged over trials. In cases where the hypothesized effects were not significant, Bayes Factors were calcu-

lated by comparing the Bayesian Information Criterion between models with and without the predictor in ques-

tion. This was done to confirm evidence for the null-hypothesis (Lee & Wagenmakers, 2014). For the peak-mu

analysis, permutation tests over the whole time course were conducted, as there was no predefined time win-

dow. To test for effects of intensity, condition and their interaction beyond mu suppression, exploratory permu-

tation tests on the whole EEG data space were conducted.

To assess the robustness of the findings, the analyses of mu, IBI and SCRs were repeated with data

averaged over trials. In these analyses, the factor intensity was dichotomized into low and high intensity (low: 1

to 10; high: 11 to 20). The results of these analyses are not reported, as they did not differ from the single-trial

results. In the figures, data are dichotomized into low and high intensities for display purposes only. Permutation

tests were carried out in MATLAB (version R2019b, The Mathworks*), and all other statistical analyses were

carried out in R (version 4.0.2, *R core team*).

1012

Behavioural data

To test for condition differences in empathic accuracy, we conducted a negative binomial generalized linear

mixed model (using the function *glmer.nb* in the *lme4* package in R, Bates et al., 2020) on observers' single-trial

ratings of the other's pain (Lawless, 1987). To find the best random slopes structure, first models with the full

fixed-effects structure and different random slopes were compared using the Akaike Information Criterion (AIC,

Akaike, 1998). An AIC difference greater than 2 was set as the threshold for a significant difference (Burnham

& Anderson, 2004). Then one fixed predictor after the other was added to the model with the optimized random-

effects structure, and only predictors that significantly improved the model were kept. The same procedure was

used to test for differences in unpleasantness ratings between the conditions.

1022

EEG data

To test whether mu suppression was modulated by shock intensity (20 levels), condition (direct vs. mediated)

or their interaction, permutation tests were conducted on the mu peak-frequency power within the window from

-500 to 1500 ms after stimulus onset (see Cohen, 2014). To test for effects of shock intensity on mu suppression,

Spearman correlations between normalized single-trial power and single-trial shock intensity were calculated for

each data point and across participants. This yielded a time course of correlation coefficients between peak-

frequency power and shock intensity. Correlation values were z-transformed by comparing them to a permuta-

tion-based null-hypothesis distribution (based on randomly shuffling over trials 1000 times). To correct for mul-

tiple comparisons, a maximum value correction was used (Cohen, 2014).

To test for condition differences in mu suppression regardless of shock intensity, mu power was aver-

aged over trials and compared between conditions. For the null-hypothesis distribution, the assignment of con-

dition was randomly shuffled over participants (1000 permutations). To test for condition differences in the cor-

relation between shock intensity and power (whether power tracked shock intensity to a larger degree in the

direct condition), the condition difference between correlation coefficients was calculated for each data point and

participant and compared to a random distribution (shuffled between conditions in 1000 permutations). For a

comparison, we also analysed power in the traditional mu band (8–13 Hz, electrode C4, 500–1000 ms after

shock onset). We chose the time window according to previous literature and to be comparable to the time

window in which the response to own pain occurred (Zebarjadi et al., 2021).

For the effects of condition, shock intensity and their interaction on theta responses, we analysed power

in the traditional theta band (4–8 Hz, electrode Fz, 0–500 ms after shock onset). We chose the time window

according to previous literature (Mu et al., 2008; Peng et al., 2021). In the latter two analyses, we used a linear

mixed model with condition averages nested in observers and observers nested in targets and dichotomized

shock intensities.

Finally, in an exploratory analysis, using the same permutation test method as described above for

peak-frequency mu, we looked for main effects of shock intensity (20 levels), condition and their interaction on

power across the whole time-frequency-electrode space (1–20 Hz, all electrodes). For the effect of shock inten-

sity, a time window from 0 to 1500 ms was used, for the other analyses a time window from -500 to 1500 ms.

For the effect of shock intensity data were downsampled to 125 Hz to reduce computation time. Again, a maxi-

mum value correction and additionally a cluster size correction (Cohen, 2014) were used to correct for multiple

comparisons.

1053

IBI and SCRs

To test for condition and shock intensity (20 levels) effects on the observers' IBI, linear mixed models on the

single-trial IBI averages were conducted (using the *lmer* function in *lme4* in R). The same procedure was used

for the skin conductance data. Here, first the time lag with the greatest SCR over all conditions was selected for

further analyses. To account for interindividual variation in SCR levels, SCR means were normalized by dividing

them by participants' individual standard deviation. As SCR data were not normally distributed, generalized

linear mixed models (using the *glmer* function in *lme4*, Gamma family, log-link function) were used for these

data. To assess the robustness of the findings, all analyses were also carried out with data averaged over trials

(with dichotomized shock intensities).

To test for coupling between targets' and observers' IBI and SCRs, similar (generalized) linear mixed

models were used, but this time the targets' IBI response or SCR was entered as a fixed predictor instead of the

stimulus intensity. For all (generalized) linear mixed models, fitting of random- and fixed-effects structure was

carried out in the same way as for the single-trial behavioural data. To assess the robustness of the findings,

skin conductance coupling was also analysed by calculating Spearman's correlation coefficients between tar-

gets' and observers' SCRs and comparing them between conditions using a linear mixed model. Correlations

were calculated for the four time lags, and the time lag with the highest correlation coefficients across both

conditions was chosen for the linear mixed model. For the robustness analysis of the IBI coupling, the Spear-

man's correlation between the targets' and the observers' continuous IBI over the whole task was calculated.

For this, IBI data were smoothed with a moving average function of 2 seconds to reduce the influence of strong

outliers. The correlation coefficients were calculated for 20 different time lags (steps of 0.5 s) between target

and observer data. For testing condition effects, the lag with the highest correlation over all conditions was used.

Correlation coefficients were transformed using the Fisher's z-transformation to obtain normally distributed data.

They were then compared between conditions using linear mixed models with correlation coefficients from dif-

ferent conditions (Level 1) nested in observers (Level 2) and targets (Level 3). The results of these analyses are

reported in the results section when they diverge from the results of the single-trial analyses.

1079

Control analysis of target expressivity

Empathic accuracy depends on the expressivity of the other (Zaki et al., 2008), and differences between direct

and mediated interactions might result from altered expressivity of the targets in either condition. To test whether

the targets show systematic differences in their pain expression between the conditions, a control experiment

with a different sample was conducted. Thirty-one participants (25 females, 6 males, mean age(SD) =

23.51(4.70)) were shown 100 segments from the video recordings of both direct and mediated interaction without

being aware of that manipulation. Video segments included four seconds before and four seconds after an

electric shock and were chosen such that there was an equal number of videos from each original session,

condition and target pain rating (summarized in 5 bins of 20 rating points each). Videos were shown in random

order. After each video the participants had 10 seconds to rate how painful the stimulus was for the target in the

video on a visual analogue scale ranging from "not painful at all" to "extremely painful". If targets expressed their

pain differently in the two conditions, we would expect condition differences in the mean pain ratings or in the

empathic accuracy of the control participants. Mean pain ratings and mean empathic accuracy scores (Spear-

man's correlations) were then compared between the conditions using t-tests.

Pre-registration

A pilot study using a similar design was pre-registered at OSF: https://osf.io/gcyqs.

1096