1   **Effects of social presence on behavioural, neural and physiological aspects of empathy for pain**

3   Abbreviated title: Effects of social presence on empathy for pain

5   Pauline Petereit[1,2] *, Ronja Weiblen[1,2,3], Anat Perry[4], Ulrike M. Krämer[1,2,5] *

[1] Department of Neurology, University of Lübeck, 23562 Lübeck, Germany

[2] Center of Brain, Behavior and Metabolism (CBBM), University of Lübeck, 23562 Lübeck, Germany

[3] Department of Psychiatry and Psychotherapy, University of Lübeck, 23562 Lübeck, Germany

[4] Psychology Department, Hebrew University of Jerusalem, Jerusalem, Israel

[5] Department of Psychology, University of Lübeck, 23562 Lübeck, Germany

* Corresponding authors:

Pauline Petereit,

University of Lübeck, Ratzeburger Allee 160, 23562 Lübeck, Germany

E-Mail: pauline.petereit@neuro.uni-luebeck.de

Phone: 004945131017422

Ulrike M. Krämer,

University of Lübeck, Ratzeburger Allee 160, 23562 Lübeck, Germany

E-Mail: ulrike.kraemer@uni-luebeck.de

Phone: 004945131017421

Abstract

In mediated interactions (e.g. video calls), less information is available about the other. To investigate how this affects

our empathy for one another, we conducted an EEG study, in which thirty human participants observed one of 5

targets undergoing painful electric stimulation, once in a direct interaction and once in a live, video-mediated interac-

tion. We found that observers were as accurate in judging others' pain and showed as much affective empathy via

video as in a direct encounter. While mu suppression, a common neural marker of empathy, was not sensitive to

others' pain, theta responses to others' pain as well as skin conductance coupling between participants were reduced

in the video-mediated condition. We conclude that physical proximity with its rich social cues is important for nuanced

physiological resonance with the other's experience. More studies are warranted to confirm these results and to

understand their behavioural significance for remote social interactions.

Key words: adults, electroencephalography, empathic accuracy, neural oscillations, physiological coupling

Introduction

Over the last decades, many social interactions in private life and at work, including medical and psychotherapeutic

contexts, have shifted from personal encounters to mediated interactions. In contrast to a direct interaction, a medi-

ated interaction is not happening face-to-face in the same space, but instead mediated via any means of communi-

cation, for example a letter, a phone call, or a video call. These interactions differ in two aspects from personal

encounters: First, they provide less detailed social cues and information channels, a feature often termed intimacy.

Second, they limit opportunities for immediate, reciprocal exchange compared to personal encounters, a feature

often termed immediacy. These factors contribute together to the degree of social presence offered within social

interactions. Note that focusing on them extends the concept of social presence from the dichotomous presence or

absence of another person to a feature of the medium that can vary continuously (Short et al. 1976; Cui et al. 2013;

but see Biocca et al. 2003 for alternative accounts of social presence). Given the reduced social presence of medi-

ated interactions, the question arises of how they change our ability to share and understand others' affective states,

i.e., our ability to empathize (Decety and Jackson 2004; Shamay-Tsoory 2011). Focusing on intimacy, we asked how

the reduction of social cues – using a video call versus a direct, personal interaction – affects empathy. While visual

and auditory cues are usually still present in a video mediated interaction, visual cues are limited as people can only

see a cutting of the scene and not the whole body and environment of the other. Moreover, the resolution and other

quality aspects of visual and auditory data might be reduced in a video call. In addition, olfactory and tactile infor-

mation, which are thought to be important for empathy, are completely missing in a video call (Goldstein et al. 2018;

Calvi et al. 2020). To investigate the effects of changed social presence on behavioural, neural and physiological

aspects of empathy, we used empathy for pain as a well-established model (Singer and Lamm 2009).

Empathy for pain is a multifaceted process, including an affective response, feelings of distress and empathic care

towards the person suffering (Lamm et al. 2007; Goubert et al. 2009; Singer and Lamm 2009), as well as cognitive

processes. The latter are sometimes measured as empathic accuracy, which is the accuracy of one's perception of

the other's pain (J. Zaki et al. 2009; Laursen et al. 2014).

On the neural level, empathy for pain has been hypothesized to be supported by representing the other's bodily state

in one's own somatosensory system and thus with neural activations in that system while observing others in pain

(Riečanský and Lamm 2019). These neural activations are associated with mu suppression in the EEG: reduced

power between 8 and 13 Hz over the somatosensory cortex (Cheng et al. 2008; Perry, Bentin, et al. 2010; Peled-

Avron et al. 2018; Peng et al. 2021). Mu suppression has been linked to higher empathic accuracy (Goldstein et al.

2018). Although one study has shown that activity in the somatosensory cortex is sensitive to others' painful facial

expressions (Gallo et al. 2018), most studies have used static pictures or videos of injured body limbs (Whitmarsh et

al. 2011; Motoyama et al. 2017; Levy et al. 2018; Zebarjadi et al. 2021) and it is therefore not clear whether this

phenomenon is specific to the perception of others' injured bodies. It is therefore useful to additionally examine neural

activity at other frequencies or areas of the cortex that might be linked to pain empathy. Some studies examined mid-

frontal theta activity (4–8 Hz) as a possible electrophysiological shared component of empathy (Mu et al., 2008; Peng

et al., 2021) and own pain experience (Misra et al. 2017; Ploner et al. 2017; Peng et al. 2021). Mid-frontal theta is

thought to indicate activity in the anterior cingulate cortex (ACC, Mitchell et al. 2008; van der Molen et al. 2017),

which is often reported in fMRI studies on empathy for pain and is associated with the negative affect during own and

others' pain (Fallon et al. 2020). ACC activation is also sensitive to facial expressions of different pain intensities

(Saarela et al. 2007).

Finally, several studies suggest that physiological "coupling" (in cardiac activity or skin conductance), i.e., aligning to

the physiological state of someone in pain, might reflect empathic sharing and facilitate understanding of the other

(Goldstein et al. 2017; Jospe et al. 2020; Reddan et al. 2020 Apr 17; Zerwas et al. 2021).

Humans share others' emotional state especially in response to social cues such as facial expressions or eye contact

(Ensenberg et al., 2017; Hess, 2021; Perry et al., 2010b). We thus expected that reduced availability of social cues

and thereby reduced intimacy of the interaction would dampen affective responses and impair understanding of

others. Additionally, if people do not share the same space, with the potential of physical interaction, feelings of

closeness and the salience of the other person and with that also empathic responses might be diminished (Melloni

et al. 2014; Järvelä et al. 2016; Weisz and Zaki 2018; Bogdanova et al. 2022). To test the effect of intimacy on

empathy for pain in the current study, pairs of participants underwent an empathy-for-pain paradigm in two conditions:

one direct, face-to-face interaction and one interaction mediated via real-time video transfer. In both conditions, one

participant (the target) received painful electrical stimulation to the back of the right hand while the other (the ob-

server) was watching. As both observer and target were able to see each other in both conditions, their behaviours

could subtly influence each other. Therefore, the task can be considered an interactive empathy task (Cui et al. 2013;

Cañigueral et al. 2022 Jun 1). The task of the observer was to judge the pain intensity experienced by the target and

her/his own feelings of unpleasantness while watching the others' pain. As targets themselves also rated their expe-

rienced pain intensity, the correspondence between targets' and observers pain provided a behavioural read-out for

empathic accuracy for pain. In addition, we measured EEG from the observer and cardiac and skin conductance

responses from both members of the dyad.

Methods

Participants

Five female psychology students were recruited as targets. Only females were sampled for the targets to reduce

possible gender effects over the dyads. They were on average 19.8 years old (SD = 0.75). Thirty psychology students

(7 males, 23 females, mean age(SD) = 24.07(4.68) years) were recruited as observers. As all the students were from

the same university and most of them from the same year, some of them had already met before the experiment (n

=10), while others had not (n =20). A priori power analyses with data simulations using the *simr* package (Green and

MacLeod 2016) in R showed that this sample size is sufficient to detect a small effect (f_2 = 0.02) of condition on

empathic accuracy (measured via an interaction effect between shock intensity and direct vs. mediated condition in

a generalized linear mixed model on single trials; see below for statistical analyses) with a power of at least 0.98

(depending on exact model structure). Data were simulated using sample parameters (means and variance param-

eters as well as correlations between conditions) from a pilot study. We did not conduct a separate power analysis

for the neural data because effect size estimations for neural effects are difficult. Moreover, the neural analysis was

conducted with the full amount of trials (80 trials) in contrast to the 40 trials used for the behavioural analysis. Hence,

the power calculation for the behavioural analyses is more conservative. For both targets and observers, exclusion

criteria were current psychiatric or cardiovascular and past or current neurological disorders, current or chronic pain

conditions or current pain-medication intake. For one observer, all physiological data from the own-pain condition

had to be excluded because of missing stimulus triggers. Skin conductance and electrocardiogram (ECG) data from

one target (mediated-interaction condition) were missing due to a technical error. Skin conductance data from two

observers could not be analysed due to poor data quality. These dyads were excluded from all analyses of the

missing outcome variable. Targets received €10 per hour; observers received course credit or €10 per hour. All

participants provided written informed consent prior to taking part in the study, including consent for video-recording

them and showing the videos to other work group members, and in case of the targets, showing the videos to other

participants in future studies. The experiment was carried out according to the Declaration of Helsinki and was ap-

proved by the Ethics Committee of the University of Lübeck.

Experimental design

Each target interacted with six different observers on six different study days. Observers came to the lab once for

one session (two interactions) with one target. For the observers, there were three conditions, which were imple-

mented in separate blocks (each observer went through all three conditions, within-subject design, see Fig. 1A). In

the "own pain" condition, the observer was alone in the laboratory and received electric shocks. This procedure

ensured that observers knew what the electric shocks felt like. In the "direct interaction" condition, observer and target

sat opposite each other at a table, and the target received electric shocks while the observer watched. Thus, in this

condition the observers were able to see the whole body of the target apart from the legs, which were placed under

the table. In the "mediated interaction" condition, target and observer sat in adjacent rooms and saw each other over

a real-time video transmission. The video showed the face, upper body, arms and hands of the target, and targets

were instructed to place the hand with the shock electrode clearly visible in front of them on the table in both condi-

tions. As the camera filming the target had to be placed above the screen the target was watching, the visual per-

spective on the target was slightly different between the "mediated" and the "direct" condition. In both conditions, a

blue board was placed behind the target to make the background as similar as possible. Observers watched the

same target in the latter two conditions and rated the observed pain experience of the target. Targets rated their own

pain experience. In all conditions, skin conductance and ECG were recorded from both participants, and EEG was

recorded from the observer. The "own pain" condition was always carried out first, and the order of "direct interaction"

and "mediated interaction" conditions was pseudo-randomized over participants such that each order ("direct inter-

action" first, or "mediated interaction" first) occurred an equal number of times. In the latter two conditions, targets'

and observers' behaviour was video-recorded.

A

B

Figure 1. (A) Schematic overview of experimental conditions. (B) Overview of the analysed outcomes.

Stimulus calibration

Prior to the pain task, pain stimuli were calibrated to the subjective pain thresholds of the participants (the observers in the "own pain" condition, and the targets prior to the first interaction condition). To this end, participants received electric shocks starting from 0 mA, increasing in amplitude in steps of 0.5 mA. They were required to rate each stimulus on a 9-point Likert scale from 0 ("not perceivable") to 8 ("strongest pain imaginable"). The intermediate steps on the scale were likewise labelled (1 = "noticeable", 2 = "unpleasant", 3 = "slightly painful", 4 = "medium painful", 5 = "very painful", 6 = "extremely painful", 7 = "unbearably painful"). We increased the stimulus intensity until participants rated a stimulus with "7" (Rutgen et al. 2015). After that, stimuli were decreased in amplitude (again in steps of 0.5 mA) until participants rated the stimulus as "0" or an amplitude of 0 mA was reached. The procedure was then repeated once more with increasing stimulus intensity. The stimulus intensity that was rated as 1 ("noticeable") in this last round was used as the lower limit for the stimuli presented during the task, with the stimulus intensity rated

as 6 ("extremely painful") used as the upper limit. For the pain task itself, the range from the lower limit to the upper

limit was evenly divided into 20 levels, to have a finer variation in shock intensity levels. As the physical intensity of

the stimuli was increased until participants rated them as "extremely painful", we could make sure that participants

really experienced pain during the experiment.

Pain task

The pain task itself was adapted from Rutgen and colleagues (2015). Electric shocks were delivered using a DS 5

isolated bipolar constant current stimulator (Digitimer) and a bar electrode (Digitimer, two electrodes with 9-mm di-

ameter, 30 mm apart) attached to the back of the right hand. The skin under the electrode was treated with an

abrasive paste and conductive gel to reduce the electric resistance of the skin.

Each trial started with an auditory cue lasting 500 ms that did not predict the shock stimulus intensity. At 1000 ms

after the cue, the electric shock was delivered for 500 ms (series of 2-ms electric pulses, interspersed with approxi-

mately 20-ms breaks). After a randomly varying interval (6000–9000 ms), the next trial or the rating followed. In 50%

of the trials, participants were prompted to rate the stimulus by a vocal recording saying, "Please rate". We included

the rating only in 50% of the trials to keep the duration of the experiment feasible. The rating was given on a tablet

computer. Targets rated how painful the electric shock was for themselves on a visual analogue scale ranging from

"not at all painful" to "extremely painful". Observers rated how painful they thought the electric shock was for the

target (on the same visual analogue scale) as well as how unpleasant it was for them to watch the target receive the

electric shock (on a visual analogue scale ranging from "not at all unpleasant" to "extremely unpleasant"). The latter

rating served as a measure of affective empathy. Electric shocks varied in intensity in 20 steps from the intensity the

targets had rated as "noticeable" (intensity level 1) to the intensity they had rated as "extremely painful" (intensity

level 20) during the calibration. There were 80 trials in each condition, and each intensity occurred four times. The

number of trials was similar to other EEG studies on pain empathy with a similar paradigm (Rutgen et al. 2015).

However, compared to former studies, we enhanced the number of rated trials from one third to half of the trials,

because our power analysis had shown that this was necessary for a reliable assessment of empathic accuracy. The

order of intensities was pseudo-random (with no more than four shocks with intensity level higher than 10 or lower

than 11 in a row) but fixed for all participants and conditions. The order of trials that had to be rated was fixed as well.

In the "own pain" condition, the task was the same except that observers rated their own pain experience on the

visual analogue scale. Targets and observers were instructed not to talk or move excessively during the task but

were otherwise allowed to express their emotions freely. Observers were instructed to rate the pain of the other as

accurately as possible. We assumed that it would be possible for the observer to infer the intensity of targets' pain

from facial expressions, gestures, para-lingual expression and body movement. Observers were free to use all these

cues as they wished. Both observers and targets were told who would receive shocks in the task, hence the observers

knew that they would receive shocks in the "own pain" condition, but would not receive any shocks in the other two

conditions.

Experimental procedure

Target selection

Before targets interacted with observers, they came to the laboratory alone to familiarize themselves with the proce-

dures and the stimuli. In this first session, they did the same pain task as in the main experiment, but with no other

person in the room. As in the main experiment, skin conductance and ECG were recorded. After the first session,

targets decided whether they wanted to participate in the main experiment. Moreover, we used the first session for

target selection, as we aimed to recruit only targets who set the intensity limits of the stimuli during stimulus calibration

to a level that was actually painful. This constraint was important to ensure that we measured actual pain empathy

during the main experiment. We therefore defined the minimum upper intensity limit that targets had to reach during

the first session to be eligible for the full experiment as within +/-1 standard deviation of the mean upper intensity

limit of a pilot study (resulting in a minimum upper intensity limit of 2.5 mA). We invited seven potential targets to the

first session. Due to the criteria, we had to exclude one target, and one participant dropped out after the first session,

which left five targets for the main experiment.

Main experiment

For the main experiment, the observer arrived first in the laboratory. After the informed-consent form was signed, the

EEG, skin conductance and ECG measurement equipment as well as the stimulus electrode were prepared. The

pain-calibration procedure was carried out. After five practice trials to familiarize themselves with the ratings on the

tablet computer, participants did the pain task in the "own pain" condition. Meanwhile, the target arrived in a different

room, responded to questionnaires and was equipped with the electrodes for physiological measurement. As soon

as the "own pain" condition was finished, the stimulus electrode was attached to the target's right hand, and the target

underwent the pain-calibration procedure. Meanwhile, the observer completed questionnaires. When both were fin-

ished, the experimental tasks started (see "pain task" above; either "direct interaction" or "mediated interaction" first).

Between conditions, there was a short break in which the target moved between rooms (either into our out of the

room in which the observer was seated). ECG and skin conductance cables of the observer were unplugged from

the amplifier during moving, and plugged in again before the next condition started. Data quality was checked again

before the second condition started. Afterwards, target and observer were seated in different rooms again and replied

to post-experimental questionnaires. Finally, observers were debriefed about the aim of the study. Targets were

debriefed only after completing all six sessions.

Questionnaires

We assessed participants' age, gender, body weight and height, educational degree, and habits regarding smoking,

caffeine consumption and physical activity. After the experiment, we obtained participants' subjective evaluation of

the experiment and observers' evaluation of the target. They also filled out two personality questionnaires: the Inter-

personal Reactivity Index (Davis, 1983; German version: Paulus, 2009) and the Emotion Regulation Questionnaire

(Gross & John, 2003; German version: Abler & Kessler, 2009). These data are not further evaluated here.

Physiological data acquisition

Participants were asked to refrain from smoking, exercise, alcohol and caffeine for at least six hours before the

experiment to prevent these factors from impacting the physiological measurements. EEG data were recorded with

59 Ag/AgCl electrodes placed on an elastic cap according to the international 10-20-system (using a BrainAmp MR

plus amplifier, BrainProducts GmbH,). An online reference electrode was placed on the left earlobe, while an offline

reference electrode was placed on the right earlobe. Horizontal and vertical EOG were recorded with four electrodes

placed next to the outer corners of the eyes and above and below the left eye, respectively. Sampling rate was 500

Hz, and data were recorded with an online high pass filter of 0.016 Hz, a low pass filter of 48 Hz and a notch filter at

50 Hz. Impedances were kept below 5kΩ.

ECG data were recorded with bipolar Ag/AgCl recording electrodes and one reference electrode, using a 50-Hz notch

filter. One of the recording electrodes was placed on the right forearm, the other one on the left lower calf of the

participant (following Einthoven lead II configuration, Einthoven et al., 1950). Skin conductance was measured with

two electrodes placed on the thenar and hypothenar of the left hand, using a 50-Hz notch filter. An electrode attached

to the left forearm served as ground for both ECG and skin conductance, which were recorded with the same amplifier

(BrainAmp ExG, BrainProducts *GmbH*). In the "direct interaction" and "mediated interaction" conditions, data from

observer and target were recorded synchronously by connecting all amplifiers to the same USB adapter feeding the

data into BrainVisionRecorder (version 1.21.0102, BrainProducts GmbH).

Quantification and statistical analysis

Physiological data processing

EEG data

All pre-processing was done in EEGLAB, version 2020.0 (Delorme & Makeig, 2004), implemented in MATLAB

R2019b (The Mathworks*).* Data were re-referenced to the right earlobe, and bipolar horizontal and vertical EOG

channels were computed. Consistently bad channels (mean number = 2.01, range = 0 to 8 channels per participant

and condition) and data segments with large artefacts were removed from the data (resulting in on average 1.6% of

removed trials per participant and condition in the final epoched data). A bandpass filter was applied (finite impulse

response filter, lower limit: 1 Hz, upper limit: 40 Hz, filterorder: 16500). Next, independent component analysis (ICA;

implemented with the *runica* function in EEGLAB) was used for ocular artefact correction. Independent components

that were clearly related to eye blinks or horizontal eye movements based on topography and time course were

visually detected and removed (ranging from 2 to 6 components per participant). Afterwards, the weights of the

remaining components were projected onto the original unfiltered data (Stropahl et al., 2018). Channels that had

been removed before the ICA were interpolated (spherical interpolation); For some participants additional bad chan-

nels (mean number = 0.44, range = 0 to 3 channels per participant and condition) had to be interpolated. Data were

then filtered with a bandpass filter with a lower limit of 0.2 Hz and an upper limit of 40 Hz (Finite impulse response

filter, filter order = 16500, hamming window). Afterwards, data were segmented into stimulus-locked epochs of 4500-

ms lengths (1000 ms before and 3500 ms after stimulus onset) and baseline-corrected to 1000 ms before stimulus

onset. A voltage threshold (between -70/70 μV and -100/100 μV) was manually set for each participant in a way that

all trials with non-ocular artefacts were removed. The number of rejected trials varied from 0 to 31% per participant

and condition and did not differ between conditions (M = 13% in all conditions).

For the time-frequency analysis, single-trial data of all electrodes were convolved with a complex Morlet wavelet as

implemented in MATLAB (function *cwt* with parameter specification 'cmor1-1.5'):

$$\omega(t) = (\pi f_b)^{-0.5} e^{-2\pi i f_c t} e^{-\frac{t^2}{f_b}},$$

where f_b = 1 is the bandwidth parameter, and f_c = 1.5 is the wavelet center frequency. Specifically, for each participant,

changes in time-varying energy were computed (square of the convolution between wavelet and signal) in the fre-

quencies (1–40 Hz, linear increase) for the 1500 ms after shock onset. Power values were converted to decibels with

respect to an average baseline from 500 to 50 ms before stimulus onset (Cohen, 2014). For analyses of peak-

frequency power (see next paragraph), we subtracted the averaged data from 500 to 50 ms before stimulus onset

as a baseline correction.

To analyse mu suppression, we determined the individual peak frequency of mu power for each participant by using

the *restingIAF* toolbox (Corcoran et al. 2018). Although mu power has often been analysed as the mean of the power

in the frequency range from 8-13 Hz, resting state oscillations over the somatosensory cortex often show a distinct

peak in one specific frequency, which is suppressed upon somatosensory activation. The peak frequency differs

between individuals (Berchicci et al. 2011; Gundlach et al. 2020). It is therefore useful to analyse desynchronization

in individual peak frequencies to target oscillations functionally related to somatosensory activation, instead of aver-

aging over a whole frequency band. Frequency peaks in the range from 8 to 13 Hz were detected in the baseline

data from 1000 to 0 ms before shock onset in the "own pain" condition at central electrodes (C1, C2, C3, C4, C5, C6,

CP1, CP2, CP3, CP4, CP5, CP6, FC1, FC2, FC3, FC4, FC5, FC6). For each detected frequency peak, the difference

in average power between baseline (-1000 to 0 ms) and stimulation (0 to 1000 ms after shock onset) was calculated.

The electrode and corresponding peak frequency with the strongest shock-related desynchronization for each par-

ticipant was chosen for all further analyses. In 16 participants, a clear peak frequency was detectable. Peaks occurred

at all possible frequencies except 9 Hz, and at the following electrodes: C1, C4, C3, CP2, CP3, C6, CP6, FC6, and

FC2. The remaining 13 participants did not show a peak frequency, and for them data from the most frequent peak

frequency and electrode were used (11 Hz at C4).

ECG data

ECG data were loaded into EEGLAB, filtered with a bandpass filter (lower: 1 Hz, upper: 30 Hz, finite impulse response

filter, filterorder = 8250) and segmented into epochs of 2 s before and 8 s after stimulus onset for the stimulus-locked

analyses. The MATLAB function *findpeaks* was used to detect the r-peaks in the segmented as well as the continuous

data (for additional analyses of physiological coupling). Afterwards, data were visually screened for wrongly assigned

or missing r-peaks. Data sections containing extrasystoles or otherwise undetectable R-peaks were treated as miss-

ing values. The interbeat interval (IBI) in ms was calculated for every pair of heartbeats and used as IBI value for

each original data point in between the two heartbeats. In this way, the IBI trace had the same time resolution as the

original data. For the analysis of condition differences in IBI responses to shocks, the mean IBI from 2 to 5 s after

cue onset (Sperl et al. 2016) was computed and baseline-corrected to the mean of the 2 s before cue onset.

Skin conductance data

Skin conductance responses (SCRs) were analysed using the Ledalab-Toolbox (Benedek & Kaernbach, 2010, ver-

sion 3.4.8) in Matlab. Data were downsampled to 50 Hz, smoothed with an adaptive gaussian window and visually

screened for strong artefacts, which were spline-interpolated. Afterwards, a continuous deconvolution analysis was

conducted to separate phasic from tonic skin conductance activity (Benedek & Kaernbach, 2010). In the following,

the mean phasic driver activity 1–4 s after shock onset was used for analyses (for targets' SCR in physiological

coupling). For the observer data (condition differences and physiological coupling), four different time lags (1–4 s)

were considered to account for a time lag in the observer's response to the target's pain expression.

Statistical analyses

We first outline the general statistical analysis approach before specifying the details. One set of statistical analyses

examined the observers' responses to the targets' pain (empathic accuracy, unpleasantness ratings, neural and

physiological responses; see Fig. 1B, left side). In these analyses, the predictors, shock intensity, condition (direct

vs. mediated) and their interaction were tested. A significant effect of shock intensity indicates that the observer's

responses are influenced by the other's pain and are therefore interpreted as empathic. A significant effect of condi-

tion indicates that social presence generally changes the observers' behaviour and physiology, whereas an interac-

tion between shock intensity and condition indicates that social presence alters the sensitivity to another's pain. The

second set of analyses examined physiological coupling between observer and target responses (see Fig. 1B, bot-

tom). In these analyses, the targets' responses, the condition (direct vs. mediated) and their interaction served as

predictors. A significant effect of targets' responses indicates that observers' and targets' responses are generally

coupled, whereas an interaction with condition indicates stronger coupling in one condition compared to the other.

In both sets of analyses, (generalized) linear mixed models with single trials (Level 1) nested in observers (Level 2)

and observers nested in targets (Level 3) were calculated. For theta activity, responses were averaged over trials. In

cases where the hypothesized effects were not significant, Bayes Factors were calculated by comparing the Bayesian

Information Criterion between models with and without the predictor in question. This was done to confirm evidence

for the null-hypothesis (Lee and Wagenmakers 2014). For the peak-mu analysis, permutation tests over the whole

time course were conducted, as there was no predefined time window. To test for effects of intensity, condition and

their interaction beyond mu suppression, exploratory permutation tests on the whole EEG data space were con-

ducted.

To assess the robustness of the findings, the analyses of mu, IBI and SCRs were repeated with data averaged over

trials. In these analyses, the factor intensity was dichotomized into low and high intensity (low: 1 to 10; high: 11 to

20). The results of these analyses are not reported, as they did not differ from the single-trial results. In the figures,

data are dichotomized into low and high intensities for display purposes only. Permutation tests were carried out in

MATLAB (version R2019b, The Mathworks*)*, and all other statistical analyses were carried out in R (version 4.0.2, *R*

core team).

Behavioural data

To test for condition differences in empathic accuracy, we conducted a negative binomial generalized linear mixed

model (using the function *glmer.nb* in the *lme4* package in R, Bates et al., 2020) on observers' single-trial ratings of

the other's pain (Lawless 1987). To find the best random slopes structure, first models with the full fixed-effects

structure and different random slopes were compared using the Akaike Information Criterion (AIC, Akaike, 1998). An

AIC difference greater than 2 was set as the threshold for a significant difference (Burnham and Anderson 2004).

Then one fixed predictor after the other was added to the model with the optimized random-effects structure, and

only predictors that significantly improved the model were kept. The same procedure was used to test for differences

in unpleasantness ratings between the conditions.

EEG data

To test whether mu suppression was modulated by shock intensity (20 levels), condition (direct vs. mediated) or their

interaction, permutation tests were conducted on the mu peak-frequency power within the window from -500 to 1500

ms after stimulus onset (see Cohen, 2014). To test for effects of shock intensity on mu suppression, Spearman

correlations between normalized single-trial power and single-trial shock intensity were calculated for each data point

and across participants. This procedure yielded a time course of correlation coefficients between peak-frequency

power and shock intensity. Correlation values were z-transformed by comparing them to a permutation-based null-

hypothesis distribution (based on randomly shuffling over trials 1000 times). To correct for multiple comparisons, a

maximum value correction was used (Cohen, 2014).

To test for condition differences in mu suppression regardless of shock intensity, mu power was averaged over trials

and compared between conditions. For the null-hypothesis distribution, the assignment of condition was randomly

shuffled over participants (1000 permutations). To test for condition differences in the correlation between shock

intensity and power (whether power tracked shock intensity to a larger degree in the direct condition), the condition

difference between correlation coefficients was calculated for each data point and participant and compared to a

random distribution (shuffled between conditions in 1000 permutations). For a comparison, we also analysed power

in the traditional mu band (8–13 Hz, electrode C4, 500–1000 ms after shock onset). We chose the time window

according to previous literature and to be comparable to the time window in which the response to own pain occurred

(Zebarjadi et al. 2021).

For the effects of condition, shock intensity and their interaction on theta responses, we analysed power in the tradi-

tional theta band (4–8 Hz, electrode Fz, 0–500 ms after shock onset). We chose the time window according to pre-

vious literature (Mu et al. 2008; Peng et al. 2021). In the latter two analyses, we used a linear mixed model with

condition averages nested in observers and observers nested in targets and dichotomized shock intensities.

Finally, in an exploratory analysis, using the same permutation test method as described above for peak-frequency

mu, we looked for main effects of shock intensity (20 levels), condition and their interaction on power across the

whole time-frequency-electrode space (1–20 Hz, all electrodes). For the effect of shock intensity, a time window from

0 to 1500 ms was used, for the other analyses a time window from -500 to 1500 ms. For the effect of shock intensity

data were downsampled to 125 Hz to reduce computation time. Again, a maximum value correction and additionally

a cluster size correction (Cohen, 2014) were used to correct for multiple comparisons.

IBI and SCRs

To test for condition and shock intensity (20 levels) effects on the observers' IBI, linear mixed models on the single-

trial IBI averages were conducted (using the *lmer* function in *lme4* in R). The same procedure was used for the skin

conductance data. Here, first the time lag with the greatest SCR over all conditions was selected for further analyses.

To account for interindividual variation in SCR levels, SCR means were normalized by dividing them by participants'

individual standard deviation. As SCR data were not normally distributed, generalized linear mixed models (using the

glmer function in *lme4*, Gamma family, log-link function) were used for these data. To assess the robustness of the

findings, all analyses were also carried out with data averaged over trials (with dichotomized shock intensities).

To test for coupling between targets' and observers' IBI and SCRs, similar (generalized) linear mixed models were

used, but this time the targets' IBI response or SCR was entered as a fixed predictor instead of the stimulus intensity.

For all (generalized) linear mixed models, fitting of random- and fixed-effects structure was carried out in the same

way as for the single-trial behavioural data. To assess the robustness of the findings, skin conductance coupling was

also analysed by calculating Spearman's correlation coefficients between targets' and observers' SCRs and com-

paring them between conditions using a linear mixed model. Correlations were calculated for the four time lags, and

the time lag with the highest correlation coefficients across both conditions was chosen for the linear mixed model.

For the robustness analysis of the IBI coupling, the Spearman's correlation between the targets' and the observers'

continuous IBI over the whole task was calculated. In this analysis, IBI data were smoothed with a moving average

function of 2 seconds to reduce the influence of strong outliers. The correlation coefficients were calculated for 20

different time lags (steps of 0.5 s) between target and observer data. For testing condition effects, the lag with the

highest correlation over all conditions was used. Correlation coefficients were transformed using the Fisher's z-trans-

formation to obtain normally distributed data. They were then compared between conditions using linear mixed mod-

els with correlation coefficients from different conditions (Level 1) nested in observers (Level 2) and targets (Level

3). The results of these analyses are reported in the results section when they diverge from the results of the single-

trial analyses.

Control analysis of target expressivity

Empathic accuracy depends on the expressivity of the other (Zaki et al. 2008), and differences between direct and

mediated interactions might result from altered expressivity of the targets in either condition. To test whether the

targets show systematic differences in their pain expression between the conditions, a control experiment with a

different sample was conducted. Thirty-one participants (25 females, 6 males, mean age(SD) = 23.51(4.70)) were

shown 100 segments from the video recordings of both direct and mediated interaction without being aware of that

manipulation. Video segments included four seconds before and four seconds after an electric shock and were cho-

sen such that there was an equal number of videos from each original session, condition and target pain rating

(summarized in 5 bins of 20 rating points each). Videos were shown in random order. After each video the participants

had 10 seconds to rate how painful the stimulus was for the target in the video on a visual analogue scale ranging

from "not painful at all" to "extremely painful". If targets expressed their pain differently in the two conditions, we would

expect condition differences in the mean pain ratings or in the empathic accuracy of the control participants. Mean

pain ratings and mean empathic accuracy scores (Spearman's correlations) were then compared between the con-

ditions using t-tests.

Pre-registration

A pilot study using a similar design was pre-registered at OSF: https://osf.io/gcyqs.

Behavioral and physiological raw data as well as the main analysis code have been deposited at

https://osf.io/pqmra/?view_only=df6b8e7cd19743d481d86ef7cb83cb83 and are publicly available at the date of

publication. Further data and code are available upon reasonable request from the first author.

Results

Social presence does not influence empathic accuracy

To assess effects of social presence on empathic accuracy, observers' ratings of targets' pain were predicted from shock intensity, condition ("direct" vs. "mediated") and their interaction using generalized linear mixed models with single trials nested in observers and observers nested in targets. The best model had a random-effects structure containing random slopes for *condition* and *intensity* (AIC difference to the next best model during fitting of random effects: -63). Moreover, it contained a fixed effect of *intensity* (b(SE) = 0.07(0.01), z = 10.24, p < 0.0001, AIC difference to a model without the fixed effect of intensity: -14). The Bayes factor for the hypothesized interaction of condition and intensity was 0.075, indicating strong evidence for the model without the interaction (Lee and Wagenmakers 2014). These results show that the intensity of the shocks received by the targets predicted the observers' pain ratings (Fig. 2A & B), indicating meaningful empathic accuracy. However, this effect did not differ between conditions. To control for possible learning effects, we assured that participants were not more accurate in the second condition (interaction between time and intensity on observer ratings: b(SE) = 0.001(0.005), z = 0.21, p = 0.84), and that the order of conditions did not moderate any effects of social presence on empathic accuracy (interaction between intensity, condition, and order condition: b(SE) = -0.002(0.024), z = -0.073, p = 0.942).

As the familiarity within the dyads varied slightly, we did additional control analyses to assure that the varying familiarity between targets and observers in our sample did not masks effects of social presence on empathic accuracy (Ma et al. 2011; Meyer et al. 2013; Motomura et al. 2015). Although familiarity might modulate the effect of social presence on empathy (see supplementary materials for detailed analyses), the results show that it is unlikely that the inclusion of familiar dyads masked any effects of social presence on empathic accuracy, rather the contrary. However, it has to be noted that the number of dyads who had met more than twice was very small (four dyads) and therefore results of these analyses have to be interpreted with caution.

Social presence increases unpleasantness ratings

We used a generalized linear mixed model to assess whether shock intensity and condition ("direct" vs. "mediated") predicted the unpleasantness ratings. The best model had a random-effects structure containing random slopes for the interaction between *condition* and *intensity* (AIC difference to the next best model during fitting of random effects: -29). It contained a fixed effect of *intensity* (b(SE) = 0.07(0.01), z = 8.07, p < 0.0001) and a fixed effect of *condition* (b(SE) = -0.14(0.05), z = -2.63, p = 0.009, AIC difference to a model without the fixed effect of *condition*: -4). The Bayes factor for the hypothesized interaction of condition and intensity was 0.021, indicating very strong evidence

487   for the model without the interaction (Lee and Wagenmakers 2014). This shows that the intensity of the shocks

488   received by the targets predicted the observers' unpleasantness ratings (Fig. 2C), but equally so in both conditions.

489   However, the observers rated the pain of the target as slightly more unpleasant in the direct than in the mediated

490   interaction (mean difference (SD) = 1.47(6.45)).

491
492
493

A

B

C

494
Figure 2. (A) Example pain rating data over trials from one randomly chosen sample dyad. (B) Predicted observer
pain ratings averaged over direct- and mediated-interaction conditions. The black thick line represents the fixed effect
of shock intensity; single coloured lines represent predicted data from single participants. The colour shading from
blue to red represents the value of the random slope for intensity of each participant (left). Predicted observer pain
ratings depending on condition (direct vs. mediated). Lines represent values for single participants. The colour shad-
ing from blue to red represents the value of the random slope for condition of each participant. The boxplots represent
summary statistics for each condition (C) Predicted observer unpleasantness ratings in direct and mediated interac-
tion. The black thick line represents the fixed effect of shock intensity; single coloured lines represent predicted data
from single participants. The colour shading from blue to red represents the value of the random slope for the inter-
action between intensity and condition of each participant. The boxplots represent summary statistics for each con-
dition. * = $p < 0.01$, n.s. = not significant.

Mu suppression is not sensitive to observed pain

Using a permutation test, we analysed whether mu suppression was sensitive to the observed pain intensity, and

whether it was reduced in the mediated condition. For each participant, we chose an individualized frequency (in the

range of 8 – 13 Hz) and electrode (of central electrodes), which showed the strongest desynchronization in the own

pain condition. Data from 29 dyads were included in these analyses, as data from the own pain condition of one

participant were not analysable due to technical issues during data collection. The analyses of peak mu suppression

in the "direct" and "mediated interaction" conditions showed no effect of *intensity* or *condition,* nor an interaction that

was significant after maximum value correction (see Fig. 3A & 3B middle and right). These results show that there

was no mu suppression which was sensitive to others' pain intensity in either condition. Similarly, analysing averaged

power over the canonical mu band (8–13 Hz) yielded no significant effect of either factor (*intensity*: b(SE) = 0.34(0.30),

t(df) =   1.13(87), *p* = 0.26; *condition*: b(SE) = -0.04(0.30), t(df) =   -0.14(87), *p* = 0.88; *interaction intensity*condition*:

b(SE) = -0.02(0.42), *t*(df) = -0.04(87), *p* = 0.97). However, peak mu significantly differed between pain levels when

participants experienced pain themselves (Fig. 3A left).

Figure 3. (A) Shown is peak mu power averaged over participants, dichotomized into low- and high-intensity shocks for display purposes only. Blue solid lines indicate time windows where the main effect of intensity reached significance (uncorrected level); blue dotted lines indicate time windows where the intensity effect survived maximum value correction. The main effect of condition and the interaction of condition x intensity did not survive maximum value correction, at any time-point. On the left side, the clear effect of pain intensity on mu power in the own-pain condition can be assessed. (B) Shown is the topography of averaged peak mu power differences between high- and low-intensity shocks in the three conditions. (C) Shown are boxplots for theta power averaged over 4–8 Hz and 0–500 ms after shock onset at electrode Fz for the direct and mediated conditions. Grey lines depict means from single participants. Significance asterisks refer to post-hoc tests from linear mixed models. * = $p < 0.001$.(D) Shown is the

533   time course of averaged theta power (4–8 Hz) after stimulus onset. (E) Shown is the topography of averaged theta
534   power differences between high- and low-intensity shocks in the two conditions.
535

Sensitivity of theta power to others' pain is modulated by social presence

We then tested whether theta responses were sensitive to observed pain intensity, and whether their sensitivity was

modulated by social presence. Data from 30 dyads were included in these analyses. The best-fitting linear mixed

model for the power averaged over the canonical theta band (4–8 Hz) yielded a significant main effect of *intensity*

(b(SE) = 0.57(0.14), *t*(df) = 3.99(90), *p* < 0.001) and a significant interaction between *condition* and *intensity* (b(SE)

= -0.46(0.20), *t*(df) = -2.30(90), *p* = 0.024, AIC difference to next best model: 2.9). Follow-up models on the significant

interaction showed a significant effect of *intensity* in the direct condition (b(SE) = 0.57(0.13), *t*(df) = 4.46(28.99), *p* <

0.001), but not in the mediated condition (b(SE) = 0.10(0.15), *t*(df) = 0.70(28.99), *p* = 0.49). This indicated that frontal

theta was more responsive to the other's shock intensity in the direct condition than in the mediated condition (see

Fig. 3C, D, E).

Exploratory analyses of whole time-frequency-electrode space

Grand Averages of EEG power in the different conditions can be found in supplementary figure S1-S6. To test for

modulations of neural activity by observed shock intensity or social presence beyond mu and theta band, we ran a

cluster based permutation test on the whole time-frequency-electrode data space. We used a cluster size correction

and a maximum value correction to correct for multiple comparisons. Data from 30 dyads were included in these

analyses. During both direct and mediated interaction, others' pain *intensity* was positively related to 2 to 6 Hz power

between 56 and 840 ms after shock onset at the frontal and central electrodes (cluster 1, maximal at F4, see Fig. 4A

left, B and E top left). Others' pain *intensity* was also negatively related to 12 to 20 Hz power between 680 and 1040

ms after shock onset over parietal and occipital electrodes (cluster 2, maximal at P2, see Fig. 4A left, B and E bottom

left).

Neither a permutation test on the *condition* effect (mediated vs. direct) nor the interaction between *condition* and

intensity yielded any cluster that survived the cluster size correction or the maximum value correction. However, due

to the exploratory nature of the analyses, we further examined the biggest cluster, which was significant on an un-

corrected level. In direct compared to mediated interactions (main effect of condition; see Fig. 4A middle, C and E

middle), theta/alpha (5–12 Hz) power at the frontal electrodes and lower beta (13–20 Hz) power at the frontal, central

and centro-parietal electrodes was enhanced in an early time window (-256–644 ms after shock onset). In a later

time window (511–1500 ms after shock onset), alpha/lower beta (8–20 Hz) power at the centro-parietal electrodes

was enhanced during direct interactions. For the interaction between *condition* and *intensity*, the biggest uncorrected

significant cluster showed a stronger positive effect of *intensity* in the "direct" than in the "mediated" condition. This

cluster spanned 3–20 Hz between -220 and 1380 ms after shock onset and included most left-hemisphere and central

electrodes (see Fig. 4A right, D). Its time course is displayed separately for low (3–9 Hz, electrode with maximal

interaction: Cz, Fig. 4E top right) and high frequencies (10–20 Hz, electrode with maximal interaction: CP1, Fig. 4E

bottom right).

572
Figure 4: (A) Clusters found in the permutation tests on the correlation between shock intensity and EEG power
across both conditions (left), on the main effect of condition (middle) and on the interaction between shock intensity

(dichotomized for display purposes only) and condition (right), averaged over electrodes. (B) Results of the permu-
tation test on the correlation between shock intensity and EEG power across direct and mediated conditions. Data
points that were significant after cluster size correction are displayed in colour; data points that were significant after
maximum value correction are marked in white. (C) Results of the permutation tests on the condition effect. Data
points of the largest cluster (uncorrected significant) are displayed in colour. (D) Results of the permutation tests on
the interaction between condition and intensity (dichotomized for display purposes only). Data points of the largest
cluster (uncorrected significant) are displayed in colour. (E) Power time course of cluster 1 (top left) and cluster 2
(bottom left), and power time course for the early cluster (5–20 Hz, F3, middle top) and the late cluster (8–20 Hz,
TP8, middle bottom) showing a condition effect and power time course of lower frequencies (3–9 Hz; top, right), and
higher frequencies (10–20 Hz; bottom, right) for interaction effects between condition and intensity. Data are dichot-
omized into low and high intensities for display purposes only.
586

Social presence does not modulate physiological responses to other's pain

Next, we tested whether observers showed stronger cardiac responses (interbeat interval, IBI, responses) to ob-

served pain in the "direct" than in the "mediated" condition, again using linear mixed models on single trials. The best

model contained random slopes for *condition* and *intensity* but no fixed effects. The Bayes factor for the hypothesized

effect of intensity was 0.027, indicating very strong evidence for the model without the effect. The Bayes factor for

the hypothesized interaction of condition and intensity was 0.088, indicating strong evidence for the model without

the interaction (Lee and Wagenmakers 2014). These results indicate that observers' IBI responses were not sensitive

to the observed shock intensity on the sample level (see Fig. 5A).

Moreover, we tested whether observers' skin conductance responses (SCRs) were more responsive to observed

pain intensity in the "direct" than in the "mediated" condition. Observers' SCRs were greatest in the time window from

1 to 4 seconds after shock onset, hence this time window was used for all further analyses. The best single-trial linear

mixed model on observers' SCR to the observed shocks contained random slopes for *condition* and *intensity* and a

fixed positive effect of *intensity*. The Bayes factor for the hypothesized interaction of condition and intensity was

0.036, indicating strong evidence for the model without the interaction (Lee and Wagenmakers 2014). These results

indicate that observers' SCRs were sensitive to the shock intensity, but equally so in direct and mediated conditions

(Fig. 5C and 5D). All model parameters are listed in Table 1. Models with dichotomized intensities yielded the same

results.

Social presence enhances skin conductance coupling between target and observer

Finally, we used a single-trial linear mixed model to test whether the direct interaction enhanced the coupling between

targets and observers IBI responses. To this end, the model tested whether observers' IBI responses could be pre-

dicted from targets' IBI responses and whether this effect was modulated by the condition ("direct" vs. "mediated"

interaction). The best model contained a random slope for *condition* but no fixed effects. The Bayes factor for the

hypothesized effect of targets' IBI was 0.059, indicating strong evidence for the model without the effect. The Bayes

factor for the hypothesized interaction of condition and targets' IBI was 0.076, indicating strong evidence for the

model without the interaction (Lee and Wagenmakers 2014). These results indicate that there was no coupling be-

tween targets' and observers' IBI responses (Fig. 5B). Comparing the correlations between the continuous IBI traces

in the two conditions showed the same results (correlations were highest for a lag of 2 seconds).

The same rationale was used to test for effects of condition on coupling between targets' and observers' skin con-

ductance responses to pain. The best single-trial linear mixed model on SCR coupling – predicting observers' SCR

from targets' SCR – contained a random slope for *condition* and *target SCR* and fixed effects of *target SCR*, *condition*

and their interaction (for model parameters, see Table 1). Follow-up models on the interaction between *condition* and

target SCR showed a significant positive effect of *target SCR* on observers' SCR in the "direct" condition (b(SE) =

0.12(0.03), *t* = 4.15, *p* < 0.0001), but not in the "mediated" condition (b(SE) = 0.03(0.02), *t* = 1.64, *p* = 0.1). When

comparing the correlation coefficients of targets' and observers' SCRs between conditions, there was only a margin-

ally significant effect of *condition* (b(SE) = -0.07(0.04), *t*(df) = -1.81(26), *p* = 0.082). These results indicate that cou-

pling between targets' and observers' SCRs was greater in the direct than the mediated interaction, but that the effect

was rather small (see Fig. 5E).

Table 1: Results of the single-trial (generalized) linear mixed models on IBI responses and SCRs

model	random slopes	SD	fixed effects	b(SE)	t(df)	p	AIC Δ
observers' IBI							-13
(n=29)	intercept/obs	25.01	intercept	4.33(2.16)	2.00(27.6)	0.055	
	condition /obs	13.00	-				
	intensity/obs	0.80					
	intercept/targ	7.9					
	condition/targ	0.13					
	intensity/targ	0.36					
observers' IBI (coupling							-26

						AIC Δ
with target IBI)						
(n=29)	intercept/obs	24.2	intercept	4.696 (2.154)	2.18 (28.9)	<0.038
	condition/obs	12.93				
	intercept/targ	<0.000001				
	condition /targ	<0.000001				
observers' SCR						-3
	intercept/obs	0.16	intercept	0.34(0.03)	11.8	<0.0001
(n=27)	condition/obs	0.21	intensity	0.01(0.002)	3.22	0.001
	intensity/obs	0.007				
	intercept/targ	0.03				
	condition /targ	0.03				
	intensity/targ	0.0009				
observers' SCR (coupling with target SCR)						-28
	intercept/obs	0.10	intercept	0.17(0.05)	3.34	0.0008
	condition /obs	0.19	target SCR	0.12(0.02)	5.85	<0.0001
(n=27)	targ-SCR/obs	0.06	condition	0.18(0.07)	2.81	0.005
	intercept/targ	0.0000	target SCR * condition	-0.08(0.02)	-5.72	<0.0001
	condition/targ	0.05				
	targ SCR/targ	0.02				

627 *Notes: IBI = interbeat interval, SCR = skin conductance response, AIC Δ = difference in Akaike information crite-*
628 *rion to the next best model. Models on IBI responses are linear mixed models, and models on SCRs are general-*
629 *ized linear mixed models (note that the latter do not provide degrees of freedom for fixed effects).*

Figure 5: (A) Grand averages of observers' IBI responses in direct and mediated interactions, dichotomized into low- and high-intensity trials for display purposes only. (B) Observer IBI responses predicted from the single-trial model on physiological coupling. The thick black line represents the fixed effect of target IBI; the thin lines represent predicted data for single participants. (C) Grand averages of observers' SCRs in direct and mediated interactions, dichotomized into low- and high-intensity trials for display purposes only. (D) Observer SCRs predicted from shock intensity in the generalized linear mixed model. The black thick line represents the fixed effect of intensity; the coloured lines display predicted values for single participants. The colour shading from blue to red represents the value of the random slopes of shock intensity for each participant (left). Observer SCrs predicted from condition. The coloured lines represent values for single participants, the colour shading from blue to red depicts the random slope of condition for the single participants (right). (E) Observer SCRs predicted from target SCRs in the generalized linear

642 mixed model on physiological coupling. The black thick line represents the fixed effect of targets' SCRs; the coloured
643 lines display predicted values for single participants. The colour shading from blue to red represents the value of the
644 random slopes of target SCR for each participant. * = p < .001, n.s. = not significant.
645

Targets do not express their pain differently in the two conditions

To test whether possible differences in empathic responses towards the targets' pain in the two conditions were due

to different expressivity of the targets' in the two conditions, we conducted a control experiment. We showed 100

video segments of targets' experiencing pain from the "direct" and "mediated" conditions to 31 naïve participants,

who rated the pain intensity of the shocks displayed in the video. These participants were unaware of the condition

manipulation in the videos. If targets' expressed their pain e.g. systematically stronger in the "direct" than in the

"mediated" condition, the new pain ratings should be higher for videos from the "direct" than from the "mediated"

condition. This was not the case. The means of the pain ratings from the video control experiment did not differ

significantly between conditions (mean difference = 0.79, $t(30)$ = 1.66, p = 0.11, Cohen's d = 0.3). The mean Spear-

man's correlations between video observers' ratings and shock intensity also did not differ between videos from direct

and mediated interaction (mean$_{direct}$(SD) = 0.36(0.16), mean$_{mediated}$(SD) = 0.32(0.13), mean difference = 0.040, $t(30)$

= 1.36, p = 0.18). These results indicate that the targets did not express their pain significantly differently in the direct

versus mediated interaction.

660
Discussion

Although mediated social interactions through video calls are becoming the new norm, the impacts on understanding

others and their feelings have not yet been researched thoroughly (Grondin et al. 2019), especially in social neuro-

science. In the current study, we explored how a video-mediated interaction affects empathy for pain on behavioural,

physiological and neural levels. We expected that less availability of social cues in a mediated interaction would

hamper empathizing with the other. However, we found that observers were just as accurate in judging the other's

pain in the mediated as in the direct interaction. Moreover, participants experienced the other's suffering as only

slightly less unpleasant in the mediated interaction. On the neural level, we did not find mu suppression over the

somatosensory cortex that was sensitive to the other's pain in either condition. However, mid-frontal theta tracked

the other's pain intensity more in the direct than in the mediated interaction. Exploratory analyses of the whole time-

frequency-electrode space showed no additional differences between direct and mediated conditions after correcting

for multiple comparisons. On a physiological level, observers' SCRs were coupled to targets' SCRs to a stronger

degree in the direct compared to the mediated interaction. In sum, behavioural empathy was not reduced in the

mediated interaction, whereas some neural and physiological aspects of empathy were dampened.

Effects of social presence on behavioural aspects of empathy for pain

Surprisingly, among the many studies on empathy for pain, hardly any measured empathic accuracy, and none

explored which type of information is necessary or sufficient to judge others' pain accurately (Gauthier et al. 2008;

Leonard et al. 2013; Laursen et al. 2014). In story-based empathy studies, empathic accuracy for emotion was re-

duced when auditory linguistic information was completely removed, whereas missing visual information did not im-

pact empathic accuracy (Zaki et al. 2009; Jospe et al. 2020). Similarly, we reveal that participants could judge the

targets' pain quite well, and this ability did not decline in the mediated interaction. In contrast to the story-based

paradigm, our results imply that visual information (apparent in both the direct interaction and the video calls) is

sufficient for empathic accuracy for others' pain. As our participants did not experience severe pain (expressed by

moaning or crying), auditory information might have been less important than for example in empathic responses to

the pain of hospital patients (Agahi and Wanic 2020). As our control analysis showed no condition differences in

target expressivity, we can be assured that these did not mask true condition differences in empathic accuracy.

Although observers showed more affective empathy in the direct than in the mediated condition, the effect size was

so small that it was practically negligible. One reason for that might be that many of our participants remembered

their own recent pain experience to help them feel with the similar experience of the other, as they reported in the

debriefing questionnaires. This strategy might have led to imagination of others' pain independent of the medium,

causing similar affective empathy (Goubert et al. 2005).

Effects of social presence on neural aspects of empathy for pain

As most former studies on empathy for pain used abstract cues or static pictures, we expected to find even stronger

mu suppression in our paradigm using real stimuli and focusing on individual peak-mu frequency (i.e., Perry, Bentin,

et al. 2010; Riečanský and Lamm 2019; Zebarjadi et al. 2021). Instead, we did not find any mu suppression in

response to others' pain. Speculatively, mu suppression is a compensatory mechanism that aids empathy for pain

via somatosensory representation of others' pain if insufficient sensory cues are available. Alternatively, it might be

specific for empathy for pain that is caused by visually perceivable physical injury, and might not be sensitive to facial

expressions of pain. Otherwise, it may be a weak signal that requires many repetitions – and stronger pain signals –

to find a significant effect. However, if mu suppression is a general empathy mechanism, our analysis should have

been sensitive enough to detect it. Therefore, our null findings on mu suppression align with recent criticism of its

robustness and validity as a mechanism underlying empathy in general (Hobson and Bishop 2016).

The mid-frontal theta/delta response constitutes another neural component of empathy for pain that has so far been

rarely examined in EEG studies (but see Mu et al. 2008; Peng et al. 2021). Mid-frontal theta has been related to the

salience, unexpectedness and aversiveness of many different types of stimuli (González-Roldan et al. 2011; Gün-

tekin and Başar 2014; Cavanagh and Shackman 2015). Therefore, the heightened sensitivity of theta to the other's

pain level might indicate that the other's pain elicits more arousal and negative affect in the direct interaction (Balconi

et al. 2009), although without resulting in measurable behavioural differences. Mid-frontal theta might stem from the

ACC, which shows reliable activity to both own and others' pain in fMRI studies (Cavanagh and Frank 2014; Fallon

et al. 2020). Confirming this assumption with source analyses was beyond the scope of the current paper but could

be an important step in future studies. The exploratory single-trial permutation test confirmed on a trend level the

interaction between others' pain intensity and direct versus mediated interaction.

Lastly, our exploratory analysis revealed stronger parietal beta suppression relating to stronger observed pain irre-

spective of the condition. Parietal beta decrease has been linked to attention to affective touch (von Mohr et al. 2018).

Future EEG studies should clarify its role in empathy for pain.

Effects of social presence on physiological coupling

Observers' cardiac activity was not sensitive to others' pain and showed no coupling with targets' cardiac activity. In

contrast, previous empathy studies found cardiac coupling in emotional empathy (Zerwas et al., 2021), especially

when semantic and auditory information was missing (Jospe et al., 2020). The shocks elicited a strong cardiac ac-

celeration in the targets in our paradigm (data not shown in this paper). Possibly these responses were not easily

mimicked by observers cardiac activity, which might be one reason for the discrepancies with former studies (Gold-

stein et al. 2017).

In contrast, we found coupling in skin conductance responses, which was the one aspect of pain empathy that was

markedly reduced in the mediated interaction. This finding indicates that the physiological coupling component of

empathy, specifically in SCR, might rely on physical proximity (Chatel-Goldman et al. 2014; Murata et al. 2020) and

possibly olfactory cues that are missing in mediated interactions (de Groot et al. 2014; Calvi et al. 2020).

Conclusions and future directions

Many recent studies examined direct interactions between participants, claiming that it is necessary for understanding social cognition (Redcay and Schilbach 2019; Fan et al. 2021; Levy et al. 2021). However, few studies have explicitly compared these new paradigms to similar tasks using mediated interactions (but see e.g. Hietanen et al., 2020). Hence, it remains unclear whether the degree of social presence affects social cognition and if these effects are due to the interactivity (here called "immediacy") or to the amount of information transferred and the shared physical space (called "intimacy") (Cui et al. 2013; Grondin et al. 2019). Therefore, by examining the impact of social presence on empathy for the first time, we add a potentially important dimension to the study of social cognition. We show that the effects are nuanced: Only the immediate mid-frontal theta response to others' pain, presumably relating to emotional arousal (Balconi et al. 2009), and SCR coupling were affected by the reduced intimacy. These findings could indicate that intimacy is especially important for more automatic, stimulus-driven empathy components. Future studies should address whether the immediacy within the interaction might have a stronger impact on all components of empathy (Hamilton and Lind 2016; Hietanen et al. 2020).

To conclude, we do not find evidence that empathy for pain is markedly impaired in video-mediated interactions, although physiological and neural resonance with the other's pain was reduced, which implies that some level of synchronization with the other is impaired. This suggests that empathic abilities might be preserved in everyday mediated social interactions, which are becoming more common. By showing specific changes in empathy components in a mediated interaction, we start to fill the gap in knowledge about social presence in social neuroscience.

Limitations of the study

The main limitation of the current study is the small sample size, which was due to the complexity of the design. This constraint is especially prominent when reporting mostly null results, as one might argue that our small sample size prevented us from detecting subtle effects of social presence. However, as power analyses showed, by analysing single trials and using a within-subject design, we had sufficient power to detect meaningful effects of social presence. Moreover, the small number of men in the sample prevented us from assessing effects of gender. However, even if there are differences in empathy between female-female and female-male dyads, effects of social presence on it should still be detectable in our within-subjects design. Nevertheless, as the literature suggests possible gender differences in empathy (Christov-Moore et al. 2014), the overrepresentation of women in our sample might prevent a generalization to men. Future studies should therefore assess the effects of social presence on empathy in gender-balanced samples. As we assessed only young students, who were used to video calls in their daily lives, our results

might not be generalizable to populations who are less familiar with computer mediated communication. The effects

of social presence might be stronger in these populations and future studies should therefore include more diverse

samples. Our sample was fairly homogenous in terms of age and sociodemographic background, as we sampled

only young psychology students. Homogeneity between participants might influence perceived similarity within a

dyad, which might influence empathic accuracy or even its modulation by social presence (Majdandžić et al. 2016;

Han 2018). We did not explicitly assess perceived similarity between participants within a dyad and can assume that

this varied slightly between dyads, so that we cannot rule out that this slightly influenced our results. However, we

also aimed to test the effects of social presence in a generalizable sample, to be sure the effects do not only hold

under very specific conditions. Similar to the effects of familiarity between target and observer, future studies will

have to address modulatory effects of similarity and other participant characteristics. Another limitation is the com-

parably low standardization of our laboratory task. Using a task with real live people, we aimed to capture real-life

empathy for pain in the best way possible in the EEG laboratory. At the same time, by using a standardized pain-

stimulation protocol, we maintained a high degree of standardization compared to studies using unstructured inter-

active paradigms (i.e., Levy et al., 2017). Our study therefore answers recent calls for more interactive and contextual

experimental methods for researching social interaction (Dumas 2011; Przyrembel et al. 2012; Sonkusare et al.

2019).

Acknowledgements:

UMK is supported by the German Science Foundation (grant number KR3691/8-1). We thank Lou Maria Lütjohann,

Celina Mävers, Marthe Mieling, Leah Reinicke, Ellyn Sänger and Jasmin Thurley for help with data collection and

pre-processing, Charlotte Petereit for help with the figures, and two anonymous reviewers for their helpful comments.

Author contributions: Conceptualization: P.P. and U.M.K; Methodology: P.P. and U.M.K; Software: P.P; Formal

Analysis: P.P.; Investigation: P.P.; Data Curation: P.P.; Writing- Original Draft: P.P; Writing- Review & Editing: P.P.,

R.W., A.P and U.M.K.; Visualization: P.P.; Supervision: U.M.K.; Project Administration: P.P. and U.M.K; Funding

Acquisition: U.M.K

Declaration of interests: "The authors declare no competing interests."

References

Abler B, Kessler H. 2009. Emotion Regulation Questionnaire – Eine deutschsprachige Fassung des ERQ von Gross und John. Diagnostica. 55(3):144–152. doi:10.1026/0012-1924.55.3.144.

Agahi S, Wanic R. 2020. Supremacy of auditory versus visual Input in somatic Empathy and perceived Pain Level. Pain Manag Nurs Off J Am Soc Pain Manag Nurses. 21(2):201–206. doi:10.1016/j.pmn.2019.06.013.

Akaike H. 1998. Information Theory and an Extension of the Maximum Likelihood Principle. In: Parzen E, Tanabe K, Kitagawa G, editors. Selected Papers of Hirotugu Akaike. New York, NY: Springer. (Springer Series in Statistics). p. 199–213. [accessed 2022 Apr 27]. https://doi.org/10.1007/978-1-4612-1694-0_15.

Balconi M, Brambilla E, Falbo L. 2009. Appetitive vs. defensive responses to emotional cues. Autonomic measures and brain oscillation modulation. Brain Res. 1296:72–84. doi:10.1016/j.brainres.2009.08.056.

Bates D, Maechler M, Bolker B, Walker S, Bojesen Christensen RH, Singmann H, Da B, Scheipl F, Grothendieck G, Green P, et al. 2020. Linear Mixed-Effects Models using "Eigen" and S4. https://github.com/lme4/lme4/.

Benedek M, Kaernbach C. 2010. Decomposition of skin conductance data by means of nonnegative deconvolution. Psychophysiology. 47(4):647–658. doi:10.1111/j.1469-8986.2009.00972.x.

Berchicci M, Zhang T, Romero L, Peters A, Annett R, Teuscher U, Bertollo M, Okada Y, Stephen J, Comani S. 2011. Development of Mu Rhythm in Infants and Preschool Children. Dev Neurosci. 33(2):130–143. doi:10.1159/000329095.

Biocca F, Harms C, Burgoon JK. 2003. Toward a More Robust Theory and Measure of Social Presence: Review and Suggested Criteria. Presence Teleoperators Virtual Environ. 12(5):456–480. doi:10.1162/105474603322761270.

Bogdanova OV, Bogdanov VB, Miller LE, Hadj-Bouziane F. 2022. Simulated proximity enhances perceptual and physiological responses to emotional facial expressions. Sci Rep. 12(1):109. doi:10.1038/s41598-021-03587-z.

Burnham KP, Anderson DR. 2004. Multimodel Inference: Understanding AIC and BIC in Model Selection. Sociol Methods Res. 33(2):261–304. doi:10.1177/0049124104268644.

Calvi E, Quassolo U, Massaia M, Scandurra A, D'Aniello B, D'Amelio P. 2020. The scent of emotions: A systematic review of human intra- and interspecific chemical communication of emotions. Brain Behav. 10(5):e01585. doi:10.1002/brb3.1585.

Cañigueral R, Krishnan-Barman S, Hamilton AF de C. 2022 Jun 1. Social signalling as a framework for second-person neuroscience. Psychon Bull Rev. doi:10.3758/s13423-022-02103-2.

Cavanagh JF, Frank MJ. 2014. Frontal theta as a mechanism for cognitive control. Trends Cogn Sci. 18(8):414–421. doi:10.1016/j.tics.2014.04.012.

Cavanagh JF, Shackman AJ. 2015. Frontal Midline Theta reflects Anxiety and cognitive Control: Meta-analytic Evidence. J Physiol Paris. 109(0):3–15. doi:10.1016/j.jphysparis.2014.04.003.

Chatel-Goldman J, Congedo M, Jutten C, Schwartz J-L. 2014. Touch increases autonomic coupling between romantic partners. Front Behav Neurosci. 8:1. doi:10.3389/fnbeh.2014.00095.

Cheng Y, Yang C-Y, Lin C-P, Lee P-L, Decety J. 2008. The perception of pain in others suppresses somatosensory oscillations: A magnetoencephalography study. NeuroImage. 40(4):1833–1840. doi:10.1016/j.neuroimage.2008.01.064.

Christov-Moore L, Simpson EA, Coudé G, Grigaityte K, Iacoboni M, Ferrari PF. 2014. Empathy: Gender effects in brain and behavior. Neurosci Biobehav Rev. 46(Pt 4):604. doi:10.1016/j.neubiorev.2014.09.001.

Cohen MX. 2014. Analyzing Neural Time Series Data: Theory and Practice. 1st ed. Cambridge, Massachusetts: The MIT Press.

Corcoran AW, Alday PM, Schlesewsky M, Bornkessel-Schlesewsky I. 2018. Toward a reliable, automated method of individual alpha frequency (IAF) quantification. Psychophysiology. 55(7):e13064. doi:10.1111/psyp.13064.

Cui G, Lockee B, Meng C. 2013. Building modern online social presence: A review of social presence theory and its instructional design implications for future trends. Educ Inf Technol. 18(4):661–685. doi:10.1007/s10639-012-9192-1.

Decety J, Jackson PL. 2004. The Functional Architecture of Human Empathy. Behav Cogn Neurosci Rev. 3(2):31. doi:10.1177/1534582304267187.

Delorme A, Makeig S. 2004. EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. J Neurosci Methods. 134(1):9–21. doi:10.1016/j.jneumeth.2003.10.009.

Dumas G. 2011. Towards a two-body neuroscience. Commun Integr Biol. 4(3):349–352. doi:10.4161/cib.4.3.15110.

Einthoven W, Fahr G, De Waart A. 1950. On the direction and manifest size of the variations of potential in the human heart and on the influence of the position of the heart on the form of the electrocardiogram. Am Heart J. 40(2):163–211. doi:10.1016/0002-8703(50)90165-7.

Ensenberg NS, Perry A, Aviezer H. 2017. Are you looking at me? Mu suppression modulation by facial expression direction. Cogn Affect Behav Neurosci. 17(1):174–184. doi:10.3758/s13415-016-0470-z.

Fallon N, Roberts C, Stancak A. 2020. Shared and distinct functional networks for empathy and pain processing: a systematic review and meta-analysis of fMRI studies. Soc Cogn Affect Neurosci. 15(7):709–723. doi:10.1093/scan/nsaa090.

Fan S, Dal Monte O, Chang SWC. 2021. Levels of naturalism in social neuroscience research. iScience.
24(7):102702. doi:10.1016/j.isci.2021.102702.
Gallo S, Paracampo R, Müller-Pinzler L, Severo MC, Blömer L, Fernandes-Henriques C, Henschel A, Lammes BK,
Maskaljunas T, Suttrup J, et al. 2018. The causal role of the somatosensory cortex in prosocial behaviour. eLife.
7:e32740. doi:10.7554/eLife.32740.
Gauthier N, Thibault P, Sullivan MJL. 2008. Individual and relational correlates of pain-related empathic accuracy in
spouses of chronic pain patients. Clin J Pain. 24(8):669–677. doi:10.1097/AJP.0b013e318173c28f.
Goldstein P, Weissman-Fogel I, Dumas G, Shamay-Tsoory SG. 2018. Brain-to-brain coupling during handholding is
associated with pain reduction. Proc Natl Acad Sci. 115(11):E2528–E2537. doi:10.1073/pnas.1703643115.
Goldstein P, Weissman-Fogel I, Shamay-Tsoory SG. 2017. The role of touch in regulating inter-partner physiological
coupling during empathy for pain. Sci Rep. 7:3252. doi:10.1038/s41598-017-03627-7.
González-Roldan AM, Martínez-Jauand M, Muñoz-García MA, Sitges C, Cifre I, Montoya P. 2011. Temporal disso-
ciation in the brain processing of pain and anger faces with different intensities of emotional expression. PAIN.
152(4):853–859. doi:10.1016/j.pain.2010.12.037.
Goubert L, Craig KD, Buysse A. 2009. Perceiving Others in Pain: Experimental and Clinical Evidence on the Role of
Empathy. In: Decety J, Ickes W, editors. The Social Neuroscience of Empathy. The MIT Press. p. 153–166. [accessed
2022 Apr 1]. http://mitpress.universitypressscholarship.com/view/10.7551/mitpress/9780262012973.001.0001/upso-
9780262012973-chapter-13.
Goubert L, Craig KD, Vervoort T, Morley S, Sullivan MJL, Williams de CAC, Cano A, Crombez G. 2005. Facing
others in pain: the effects of empathy: Pain. 118(3):285–288. doi:10.1016/j.pain.2005.10.025.
Green P, MacLeod CJ. 2016. SIMR: an R package for power analysis of generalized linear mixed models by simu-
lation. Methods Ecol Evol. 7(4):493–498. doi:10.1111/2041-210X.12504.
Grondin F, Lomanowska AM, Jackson PL. 2019. Empathy in computer-mediated interactions: A conceptual frame-
work for research and clinical practice. Clin Psychol Sci Pract. 26(4):17–17. doi:10.1111/cpsp.12298.
de Groot JHB, Semin GR, Smeets MAM. 2014. I can see, hear, and smell your fear: Comparing olfactory and audi-
ovisual media in fear communication. J Exp Psychol Gen. 143(2):825–834. doi:10.1037/a0033731.
Gross JJ, John OP. 2003. Individual differences in two emotion regulation processes: implications for affect, relation-
ships, and well-being. J Pers Soc Psychol. 85(2):348–362. doi:10.1037/0022-3514.85.2.348.
Gundlach C, Müller MM, Hoff M, Ragert P, Nierhaus T, Villringer A, Sehm B. 2020. Reduction of somatosensory
functional connectivity by transcranial alternating current stimulation at endogenous mu-frequency. NeuroImage.
221:117175. doi:10.1016/j.neuroimage.2020.117175.
Güntekin B, Başar E. 2014. A review of brain oscillations in perception of faces and emotional pictures. Neuropsy-
chologia. 58:33–51. doi:10.1016/j.neuropsychologia.2014.03.014.
Hamilton AF de C, Lind F. 2016. Audience effects: what can they tell us about social neuroscience, theory of mind
and autism? Cult Brain. 4(2):159–177. doi:10.1007/s40167-016-0044-5.
Han S. 2018. Neurocognitive Basis of Racial Ingroup Bias in Empathy. Trends Cogn Sci. 22(5):400–421.
doi:10.1016/j.tics.2018.02.013.
Hess U. 2021. Who to whom and why: The social nature of emotional mimicry. Psychophysiology. 58(1):e13675.
doi:10.1111/psyp.13675.
Hietanen JO, Peltola MJ, Hietanen JK. 2020. Psychophysiological responses to eye contact in a live interaction and
in video call. Psychophysiology. 57(6):e13587. doi:10.1111/psyp.13587.
Hobson HM, Bishop DVM. 2016. Mu suppression – A good measure of the human mirror neuron system? Cortex.
82:290–310. doi:10.1016/j.cortex.2016.03.019.
Järvelä S, Kätsyri J, Ravaja N, Chanel G, Henttonen P. 2016. Intragroup Emotions: Physiological Linkage and Social
Presence. Front Psychol. 7:105. doi:10.3389/fpsyg.2016.00105.
Jospe K, Genzer S, Klein Selle N, Ong D, Zaki J, Perry A. 2020. The contribution of linguistic and visual cues to
physiological synchrony and empathic accuracy. Cortex J Devoted Study Nerv Syst Behav. 132:296–308.
doi:10.1016/j.cortex.2020.09.001.
Lamm C, Nusbaum HC, Meltzoff AN, Decety J. 2007. What Are You Feeling? Using Functional Magnetic Resonance
Imaging to Assess the Modulation of Sensory and Affective Responses during Empathy for Pain. Warrant E, editor.
PLoS ONE. 2(12):e1292. doi:10.1371/journal.pone.0001292.
Laursen HR, Siebner HR, Haren T, Madsen K, Grønlund R, Hulme O, Henningsson S. 2014. Variation in the oxytocin
receptor gene is associated with behavioral and neural correlates of empathic accuracy. Front Behav Neurosci. 8.
doi:10.3389/fnbeh.2014.00423.
Lawless JF. 1987. Negative binomial and mixed poisson regression. Can J Stat. 15(3):209–225.
doi:10.2307/3314912.
Lee MD, Wagenmakers E-J. 2014. Bayesian Cognitive Modeling: A Practical Course. Cambridge University Press.
Leonard MT, Issner JH, Cano A, Williams AM. 2013. Correlates of Spousal Empathic Accuracy for Pain-related
Thoughts and Feelings. Clin J Pain. 29(4):324–333. doi:10.1097/AJP.0b013e3182527bfd.

Levy J, Goldstein A, Feldman R. 2017. Perception of social synchrony induces mother-child gamma coupling in the
social brain. Soc Cogn Affect Neurosci. 12(7):1036–1046. doi:10.1093/scan/nsx032.
Levy J, Goldstein A, Pratt M, Feldman R. 2018. Maturation of Pain Empathy from Child to Adult Shifts from Single to
Multiple Neural Rhythms to Support Interoceptive Representations. Sci Rep. 8(1):1810. doi:10.1038/s41598-018-
19810-3.
Levy J, Lankinen K, Hakonen M, Feldman R. 2021. The integration of social and neural synchrony: a case for eco-
logically   valid   research   using   MEG   neuroimaging.   Soc   Cogn   Affect   Neurosci.   16(1–2):143–152.
doi:10.1093/scan/nsaa061.
Lopez-Calderon J, Luck SJ. 2014. ERPLAB: an open-source toolbox for the analysis of event-related potentials.
Front Hum Neurosci. 8:213. doi:10.3389/fnhum.2014.00213.
Ma Q, Shen Q, Xu Q, Li D, Shu L, Weber B. 2011. Empathic responses to others' gains and losses: An electrophys-
iological investigation. NeuroImage. 54(3):2472–2480. doi:10.1016/j.neuroimage.2010.10.045.
Majdandžić J, Amashaufer S, Hummer A, Windischberger C, Lamm C. 2016. The selfless mind: How prefrontal
involvement in mentalizing with similar and dissimilar others shapes empathy and prosocial behavior. Cognition.
157:24–38. doi:10.1016/j.cognition.2016.08.003.
Melloni M, Lopez V, Ibanez A. 2014. Empathy and contextual social cognition. Cogn Affect Behav Neurosci.
14(1):407–425. doi:10.3758/s13415-013-0205-3.
Meyer ML, Masten CL, Ma Y, Wang C, Shi Z, Eisenberger NI, Han S. 2013. Empathy for the social suffering of friends
and   strangers   recruits   distinct   patterns   of   brain   activation.   Soc   Cogn   Affect   Neurosci.   8(4):446–454.
doi:10.1093/scan/nss019.
Misra G, Wang W, Archer DB, Roy A, Coombes SA. 2017. Automated classification of pain perception using high-
density electroencephalography data. J Neurophysiol. 117(2):786–795. doi:10.1152/jn.00650.2016.
Mitchell DJ, McNaughton N, Flanagan D, Kirk IJ. 2008. Frontal-midline theta from the perspective of hippocampal
"theta." Prog Neurobiol. 86(3):156–185. doi:10.1016/j.pneurobio.2008.09.005.
von Mohr M, Crowley MJ, Walthall J, Mayes LC, Pelphrey KA, Rutherford HJV. 2018. EEG captures affective touch:
CT-optimal touch and neural oscillations. Cogn Affect Behav Neurosci. 18(1):155–166. doi:10.3758/s13415-017-
0560-6.
van der Molen MJW, Dekkers LMS, Westenberg PM, van der Veen FM, van der Molen MW. 2017. Why don't you
like me? Midfrontal theta power in response to unexpected peer rejection feedback. NeuroImage. 146:474–483.
doi:10.1016/j.neuroimage.2016.08.045.
Motomura Y, Takeshita A, Egashira Y, Nishimura T, Kim Y-K, Watanuki S. 2015. Interaction between valence of
empathy and familiarity: is it difficult to empathize with the positive events of a stranger? J Physiol Anthropol. 34(1):13.
doi:10.1186/s40101-015-0049-3.
Motoyama Y, Ogata K, Hoka S, Tobimatsu S. 2017. Frequency-dependent changes in sensorimotor and pain affec-
tive systems induced by empathy for pain. J Pain Res. Volume 10:1317–1326. doi:10.2147/JPR.S129791.
Mu Y, Fan Y, Mao L, Han S. 2008. Event-related theta and alpha oscillations mediate empathy for pain. Brain Res.
1234:128–136. doi:10.1016/j.brainres.2008.07.113.
Murata A, Nishida H, Watanabe K, Kameda T. 2020. Convergence of physiological responses to pain during face-
to-face interaction. Sci Rep. 10(1):450. doi:10.1038/s41598-019-57375-x.
Peled-Avron L, Goldstein P, Yellinek S, Weissman-Fogel I, Shamay-Tsoory SG. 2018. Empathy during consoling
touch is modulated by mu-rhythm: An EEG study. Neuropsychologia. 116:68–74. doi:10.1016/j.neuropsycholo-
gia.2017.04.026.
Peng W, Lou W, Huang X, Ye Q, Tong RK-Y, Cui F. 2021. Suffer together, bond together: Brain-to-brain synchroni-
zation and mutual affective empathy when sharing painful experiences. NeuroImage. 238:118249. doi:10.1016/j.neu-
roimage.2021.118249.
Perry A, Bentin S, Bartal IB-A, Lamm C, Decety J. 2010. "Feeling" the pain of those who are different from us:
Modulation of EEG in the mu/alpha range. Cogn Affect Behav Neurosci. 10(4):493–504. doi:10.3758/CABN.10.4.493.
Perry A, Troje NF, Bentin S. 2010. Exploring motor system contributions to the perception of social information:
Evidence   from   EEG   activity   in   the   mu/alpha   frequency   range.   Soc   Neurosci.   5(3):272–284.
doi:10.1080/17470910903395767.
Ploner   M,   Sorg   C,   Gross   J.   2017.   Brain   Rhythms   of   Pain.   Trends   Cogn   Sci.   21(2):100–110.
doi:10.1016/j.tics.2016.12.001.
Przyrembel M, Smallwood J, Pauen M, Singer T. 2012. Illuminating the dark matter of social neuroscience: Consid-
ering the problem of social interaction from philosophical, psychological, and neuroscientific perspectives. Front Hum
Neurosci. 6:190. doi:10.3389/fnhum.2012.00190.
Redcay E, Schilbach L. 2019. Using second-person neuroscience to elucidate the mechanisms of social interaction.
Nat Rev Neurosci. 20(8):495–505. doi:10.1038/s41583-019-0179-4.
Reddan MC, Young H, Falkner J, López-Solà M, Wager TD. 2020 Apr 17. Touch and social support influence inter-
personal synchrony and pain. Soc Cogn Affect Neurosci. doi:10.1093/scan/nsaa048.

Riečanský I, Lamm C. 2019. The Role of Sensorimotor Processes in Pain Empathy. Brain Topogr. 32(6):965–976.
doi:10.1007/s10548-019-00738-4.
Rutgen M, Seidel E-M, Riečanský I, Lamm C. 2015. Reduction of Empathy for Pain by Placebo Analgesia Suggests
Functional Equivalence of Empathy and First-Hand Emotion Experience. J Neurosci. 35(23):8938–8947.
doi:10.1523/JNEUROSCI.3936-14.2015.
Saarela MV, Hlushchuk Y, Williams AC de C, Schürmann M, Kalso E, Hari R. 2007. The Compassionate Brain:
Humans Detect Intensity of Pain from Another's Face. Cereb Cortex. 17(1):230–237. doi:10.1093/cercor/bhj141.
Shamay-Tsoory SG. 2011. The Neural Bases for Empathy. The Neuroscientist. 17(1):7.
Short J, Williams E, Christie B. 1976. The social psychology of telecommunications. London; New York: Wiley.
Singer T, Lamm C. 2009. The Social Neuroscience of Empathy. Ann N Y Acad Sci. 1156(1):81–96.
doi:10.1111/j.1749-6632.2009.04418.x.
Sonkusare S, Breakspear M, Guo C. 2019. Naturalistic Stimuli in Neuroscience: Critically Acclaimed. Trends Cogn
Sci. 23(8):699–714. doi:10.1016/j.tics.2019.05.004.
Sperl MFJ, Panitz C, Hermann C, Mueller EM. 2016. A pragmatic comparison of noise burst and electric shock
unconditioned stimuli for fear conditioning research with many trials. Psychophysiology. 53(9):1352–1365.
doi:10.1111/psyp.12677.
Stropahl M, Bauer A-KR, Debener S, Bleichner MG. 2018. Source-Modeling Auditory Processes of EEG Data Using
EEGLAB and Brainstorm. Front Neurosci. 12:309. doi:10.3389/fnins.2018.00309.
Weisz E, Zaki J. 2018. Motivated empathy: a social neuroscience perspective. Curr Opin Psychol. 24:67–71.
doi:10.1016/j.copsyc.2018.05.005.
Whitmarsh S, Nieuwenhuis I, Barendregt H, Jensen O. 2011. Sensorimotor Alpha Activity is Modulated in Response
to the Observation of Pain in Others. Front Hum Neurosci. 5:91. doi:doi: 10.3389/fnhum.2011.00091.
Zaki J, Bolger N, Ochsner K. 2008. It Takes Two: The Interpersonal Nature of Empathic Accuracy. Psychol Sci.
19(4):399–404. doi:10.1111/j.1467-9280.2008.02099.x.
Zaki Jamil, Bolger N, Ochsner K. 2009. Unpacking the informational bases of empathic accuracy. Emotion. 9(4):478–
487. doi:10.1037/a0016551.
Zaki J., Weber J, Bolger N, Ochsner K. 2009. The neural bases of empathic accuracy. Proc Natl Acad Sci.
106(27):11382–11387. doi:10.1073/pnas.0902666106.
Zebarjadi N, Adler E, Kluge A, Jääskeläinen IP, Sams M, Levy J. 2021. Rhythmic Neural Patterns During Empathy
to Vicarious Pain: Beyond the Affective-Cognitive Empathy Dichotomy. Front Hum Neurosci. 15:708107.
doi:10.3389/fnhum.2021.708107.
Zerwas FK, Springstein T, Karnilowicz HR, Lam P, Butler EA, John OP, Mauss IB. 2021. "I feel you": Greater linkage
between friends' physiological responses and emotional experience is associated with greater empathic accuracy.
Biol Psychol. 161:108079. doi:10.1016/j.biopsycho.2021.108079.

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014

1015

1016

Figures

Figure 1. (A) Schematic overview of experimental conditions. (B) Overview of the analysed outcomes.

Figure 2. (A) Example pain rating data over trials from one randomly chosen sample dyad. (B) Predicted observer pain ratings averaged over direct- and mediated-interaction conditions. The black thick line represents the fixed effect of shock intensity; single coloured lines represent predicted data from single participants. The colour shading from blue to red represents the value of the random slope for intensity of each participant (left). Predicted observer pain ratings depending on condition (direct vs. mediated). Lines represent values for single participants. The colour shading from blue to red represents the value of the random slope for condition of each participant. The boxplots represent summary statistics for each condition (C) Predicted observer unpleasantness ratings in direct and mediated interaction. The black thick line represents the fixed effect of shock intensity; single coloured lines represent predicted data from single participants. The colour shading from blue to red represents the value of the random slope for the interaction between intensity and condition of each participant. The boxplots represent summary statistics for each condition. * = p < 0.01, n.s. = not significant.

Figure 3. (A) Shown is peak mu power averaged over participants, dichotomized into low- and high-intensity shocks for display purposes only. Blue solid lines indicate time windows where the main effect of intensity reached significance (uncorrected level); blue dotted lines indicate time windows where the intensity effect survived maximum value correction. The main effect of condition and the interaction of condition x intensity did not survive maximum value correction, at any time-point. On the left side, the clear effect of pain intensity on mu power in the own-pain condition can be assessed. (B) Shown is the topography of averaged peak mu power differences between high- and low-intensity shocks in the three conditions. (C) Shown are boxplots for theta power averaged over 4–8 Hz and 0–500 ms after shock onset at electrode Fz for the direct and mediated conditions. Grey lines depict means from single participants. Significance asterisks refer to post-hoc tests from linear mixed models. * = p < 0.001.(D) Shown is the time course of averaged theta power (4–8 Hz) after stimulus onset. (E) Shown is the topography of averaged theta power differences between high- and low-intensity shocks in the two conditions.

Figure 4: (A) Clusters found in the permutation tests on the correlation between shock intensity and EEG power across both conditions (left), on the main effect of condition (middle) and on the interaction between shock intensity (dichotomized for display purposes only) and condition (right), averaged over electrodes. (B) Results of the permutation test on the correlation between shock intensity and EEG power across direct and mediated conditions. Data points that were significant after cluster size correction are displayed in colour; data points that were significant after maximum value correction are marked in white. (C) Results of the permutation tests on the condition effect. Data points of the largest cluster (uncorrected significant) are displayed in colour. (D) Results of the permutation tests on the interaction between condition and intensity (dichotomized for display purposes only). Data points of the largest cluster (uncorrected significant) are displayed in colour. (E) Power time course of cluster 1 (top left) and cluster 2 (bottom left), and power time course for the early cluster (5–20 Hz, F3, middle top) and the late cluster (8–20 Hz, TP8, middle bottom) showing a condition effect and power time course of lower frequencies (3–9 Hz; top, right), and higher frequencies (10–20 Hz; bottom, right) for interaction effects between condition and intensity. Data are dichotomized into low and high intensities for display purposes only.

Figure 5: (A) Grand averages of observers' IBI responses in direct and mediated interactions, dichotomized into low- and high-intensity trials for display purposes only. (B) Observer IBI responses predicted from the single-trial model on physiological coupling. The thick black line represents the fixed effect of target IBI; the thin lines represent predicted data for single participants. (C) Grand averages of observers' SCRs in direct and mediated interactions, dichotomized into low- and high-intensity trials for display purposes only. (D) Observer SCRs predicted from shock intensity in the generalized linear mixed model. The black thick line represents the fixed effect of intensity; the coloured lines display predicted values for single participants. The colour shading from blue to red represents the value of the random slopes of shock intensity for each participant (left). Observer SCrs predicted from condition. The coloured lines represent values for single participants, the colour shading from blue to red depicts the random slope of condition for the single participants (right). (E) Observer SCRs predicted from target SCRs in the generalized linear mixed model on physiological coupling. The black thick line represents the fixed effect of targets' SCRs; the coloured lines display predicted values for single participants. The colour shading from blue to red represents the value of the random slopes of target SCR for each participant. * = p < .001, n.s. = not significant.