Hippocampal damage causes retrograde amnesia for objects' visual, but not odour, properties

Running title: Odour memory is not hippocampus dependent

Sean G. Lacoursiere[1], Brendan B. McAllister[1], Crystal Hadikin[1,3], Wayne W. Tschetter[1,4], Hugo Lehmann[2], Robert J. Sutherland[1]

[1] Department of Neuroscience, Canadian Centre for Behavioural Neuroscience, The University of Lethbridge, Lethbridge, Alberta, Canada

[2] Department of Psychology, Trent University, Peterborough, Ontario, Canada

[3] Canadian School of Natural Nutrition, Sooke, British Columbia, Canada

[4] Department of Ophthalmology, Oregon Health and Science University, Portland, Oregon, United States of America

Corresponding author:

  Robert J. Sutherland; email: robert.sutherland@uleth.ca

ABSTRACT

Damage to the hippocampus produces profound retrograde amnesia, but odour and object discrimination memories can be spared in the retrograde direction following hippocampal damage. Prior lesion studies testing retrograde amnesia for object/odour discriminations are problematic due to sparing of large parts of the hippocampus, which may support memory recall, and/or the presence of uncontrolled, distinctive odours that may support object discrimination. To address these issues, we used a simple object discrimination test to assess memory in rats. Two visually distinct objects, paired with distinct odour cues, were presented to the rats. One object was associated with a reward. Following training, neurotoxic hippocampal lesions were made using *N-methyl-D-aspartate* (NMDA). The rats were then tested on the preoperatively learned object discrimination problem, with and without the availability of odour or visual cues during testing. The rats were also postoperatively trained on a new object discrimination problem. Lesion sizes ranged from 67-97% of the hippocampus, with an average of 87%. On the preoperatively learned discrimination problem, the rats with hippocampal lesions showed preserved object discrimination memory when tested in the dark (i.e., without visual cues) but not when the explicit odour cues were removed from the objects. Hippocampal lesions did not prevent rats from solving a postoperatively learned object discrimination problem. Our results support the idea that long-term memories for odours, unlike recall of visual properties of objects, does not depend on the hippocampus in rats, consistent with previous observations that hippocampal damage does not cause retrograde amnesia for odour memories.

KEYWORDS: Hippocampal Formation; Discrimination Learning; Memory, Long-Term; Olfaction; Amnesia, Retrograde; Excitatory Amino Acid Agonists

1. INTRODUCTION

Studies in humans and laboratory animals show that damage to the hippocampus produces profound retrograde amnesia. The magnitude of the loss of previously learned information is related to the extent of the damage to the hippocampus (Epp et al., 2008; Lehmann et al., 2010; Mumby et al., 1999; Sutherland et al., 2001). Hippocampal damage in rats causes retrograde amnesia for a wide range of memory types, including place (Morris et al., 1982), context (Broadbent & Clark, 2013), object recognition (Gaskin et al., 2003), shock-probe conditioning (Lehmann et al., 2006), and visual discrimination (Epp et al., 2008) memories (see Lee et al., 2016 for review). However, olfactory and object discrimination memories have been reported to be spared in the retrograde direction by similar hippocampal damage (Jonasson et al., 2004; Mumby et al., 1999). Olfaction is a highly conserved sensory system and may be the main sensory modality rodents use to interact with an environment (Ache & Young, 2005). The responses from different olfactory receptors are segregated in different cells in the olfactory epithelium, allowing discrimination between odorants (Ache & Young, 2005; Aqrabawi & Kim, 2020), perhaps explaining how the olfactory system in rodents can function independently of the hippocampus in storing and retrieving odour memories acquired during simple odour discrimination tasks.

Two-choice discrimination problems are commonly used to assess memory in rodents. The typical task involves presenting a rat with two cues, which could be objects, visual cues, odours, and so on, depending on the nature of the task. The rat is trained to criteria to select the rewarded cue (S+) and not the other (S-). In the testing phase, rats that have learned the task properly will persist in selecting S+ despite no longer receiving a reward. Hippocampal damage after visual discrimination training has been found to cause retrograde amnesia (Driscoll et al.,

2005; Epp et al., 2008), suggesting the hippocampus is important for this type of memory. In odour discrimination tasks, however, rats with extensive hippocampal damage show postoperative performance similar to their preoperative performance. For example, in a study by Jonasson et al. (2004), rats were presented with a forced decision on a simple odour discrimination. Following training, the rats were subjected to NMDA lesions and, after recovery, were tested on the preoperatively learned task. The rats with hippocampus damage showed equivalent recall to the intact rats. These results suggest that, unlike other types of memory, the memories used to solve odour discrimination problems may not be dependent on the hippocampus for recall.

Prior lesion studies conducted to test retrograde amnesia for object/odour discriminations are problematic because the objects used may have allowed for inadvertent use of odours to enable discrimination (Lehmann et al., 2007; Mumby et al., 1999; Wible et al., 1992), leading to ambiguity in the interpretation of the role of the hippocampus in object discrimination memory. Therefore, in the present study we trained rats to discriminate between objects constructed of identical materials, to eliminate the use of intrinsic odour cues to differentiate between the objects. The objects were visually distinct and, during training, were paired with explicit odours that could be removed during testing, allowing the experimenter to control whether the rats could use odour cues to solve the problem.

Following nearly-complete hippocampal damage, the ability of the rats to solve the pre-operative object discrimination problem was tested with and without odour and visual cues. The rats showed preserved memory when the use of visual cues was eliminated, but retrograde amnesia for the same object discrimination problem when the explicit odour cues were removed. These results indicate that rats with hippocampal lesions can use odour cues to solve a

preoperatively acquired discrimination problem, supporting the idea that long-term memory for odours, unlike recall of visual cues, does not depend on hippocampal circuitry in rats.

2. METHODS

2.1. Subjects

The University of Lethbridge Animal Care Committee, which follows the guidelines set by the Canadian Council on Animal Care, approved all procedures. In total, 29 male Long-Evans hooded rats (CCBN vivarium, Lethbridge, AB, Canada), weighing 250-350 g, were used. The rats were housed individually in standard laboratory cages on a 12:12 light-dark cycle (lights on at 7:00 AM), and all testing was done during the light phase. The rats had free access to water and were fed ~25 g of rat chow per day throughout the experiment, except during the first week of recovery from surgery, during which food was given *ad libitum.*

2.2. Apparatus

The apparatus for the object discrimination task has been described in detail elsewhere (Mumby et al., 1990). Briefly, it consisted of an elevated runway, separated from identical goal areas at each end by opaque guillotine doors. Each goal area contained two food wells, into which food pellets (45 mg Bio-Serv, Inc., Frenchtown, NJ, USA) could be delivered by hand through plastic tubes that were mounted on the outside of the apparatus. A short divider wall protruded from the center of the end wall and separated the two food wells.

The objects used as discriminanda (see Fig. 1) were large enough to cover a food well in the apparatus but small enough to be easily displaced by a rat. Specifically, the first pair of objects was a small house constructed out of red and blue Lego and a small step stool constructed of yellow and green Lego. The objects were of equal weight and contained openings in which scented cotton balls could be placed. The openings were fashioned so that that the rats

were unable to touch or lick the cotton balls. The second pair of objects was a plastic egg holder

and a centrifuge tube. The objects were washed after every session with a solution of diluted

chlorine bleach to remove any extraneous scents acquired during displacement by the rats or

handling by the experimenter.

A.　　　　　　　　　　　　　　　　　　**B.**

Figure 1. Discriminanda used for two object discrimination problems: Discrimination 1 (**A**), which
was learned prior to hippocampal lesion surgery and tested postoperatively, and Discrimination 2
(**B**), which was learned and tested postoperatively.

The odours consisted of vanilla and peppermint concentrates manufactured by LorAnn

Oils®. To present the odours, clean cotton balls containing 0.3 ml of oil were placed inside the

objects each day. Based on previous reports (Scott et al., 2013), the odours used were chosen so

that their subjective olfactory characteristics were distinct enough to discriminate between the

two.

2.3. Preoperative Object Discrimination Training

The rats were initially habituated to the apparatus for several days, and their behavior was

shaped to retrieve food pellets from the food wells (see Mumby et al., 1990). At least 24 h after

their last habituation/shaping session, the rats were trained on the first discrimination problem

(Discrimination 1) in a single short session. The object pair consisted of the vanilla-scented Lego house and the peppermint-scented Lego stool (Fig. 1). One of the objects was designated S+ (rewarded) and the other was designated S- (not rewarded), counterbalanced within groups. The spatial position of S+ and S- was pseudorandomly counterbalanced across trials. Each rat was given a minimum of 40 trials, and the session ended when the rat reached criterion (7 consecutive choices of S+). The rats were then matched according to the number of trials they required to reach criterion and assigned to either the SHAM or HPC group.

2.4. Surgery

Approximately 72 h after training on Discrimination 1, the rats underwent surgery. Hippocampal lesions were completed following the protocol described by Lehmann et al. (2010). Briefly, the rats were anaesthetized with isoflurane (Janssen, Toronto, ON, Canada) in 1.0 L/min oxygen at 21 °C (Benson Medical Industries, Markham, ON, Canada) and administered buprenorphine (0.07 cc, 0.3 mg/mL, i.p.; Reckitt & Colman, Richmond, VA, USA) as an analgesic. The rats were then transferred to a stereotaxic frame (Kopf Instruments, Tujunga, CA, USA), and the scalp was cleaned with 4% stanhexidine followed by 70% isopropyl alcohol. A midline incision was made to expose the skull, and the periosteum was excised. The rats in the HPC group received a complete bilateral lesion of the hippocampal formation made by intrahippocampal injections of N-methyl-D-Aspartic acid (NMDA; 7.5 µg/µL in 0.9% saline; Sigma Chemical Co., St. Louis, MO, USA) at ten sites bilaterally, following coordinates from Lehmann et al. (2010). The infusions were done sequentially through a 30-gauge injection cannula attached to a 10 µL Hamilton syringe by polyethylene tubing (PE-50). At each site, 0.4 µL was infused at a rate of 0.15 µL/min. The cannula was left in place for 2.5 min following injection to facilitate diffusion. The scalp was closed using wound clips following the injections.

Diazepam (0.2-0.6 cc, 10 mg/mL, i.p.; Sabex, Boucherville, QC, Canada) was given to the rats as a prophylactic against seizures as they awakened. The same surgical procedure was given to the rats in the SHAM group, but no damage was done to the skull or brain of these rats. This sham surgical procedure (not drilling into the skull and/or not lowering the infusion cannula into the cortex overlying the hippocampus) is now common across many labs investigating the effects of hippocampal damage on memory, including retrograde memory (Broadbent et al., 2007; Epp et al., 2008; Gaskin et al., 2003; Gidyk et al., 2021; McDonald et al., 2018; Quinn et al., 2008; Tse et al., 2007; Winocur et al., 2005, 2013).

2.5. Postoperative object discrimination training

Fourteen days after surgery, the rats were trained on a new object discrimination problem (Discrimination 2). The purpose of Discrimination 2 was, first, to motivate the rats to continue displacing the objects on the Discrimination 1 extinction trials during retention testing and, second, to correct any procedural/motor deficits in the object discrimination performance that may have resulted from the lesions. Training the rats postoperatively on a second discrimination also allowed for assessment of the effects of the lesion on object discrimination acquisition, and it provided a way to distinguish the anterograde and retrograde effects of the lesions on memory.

For Discrimination 2, a new pair of objects (egg holder and centrifuge tube), with no odour added to either object, served as the discriminanda (Fig. 1). As before, one of the objects was designated S+ and the other one was designated S-, counterbalanced within groups. The rats were given 20 training trials per day for a minimum of 2 days. The training continued for a given rat until it reached a criterion of at least 16 out of 20 choices of S+ in a session.

2.6. Object Discrimination Retention Testing

The rats' retention for the object discrimination problems was assessed 24 h after criterion was reached for Discrimination 2. The rats were tested over 2 days to determine the retention of Discrimination 1. On one day, the rats were tested for Discrimination 1 with the explicit peppermint and vanilla odour cues and under the original visual/lighting conditions (Standard test). On the other day, the rats assigned to the No-Odour condition were tested without explicit odour cues (i.e., with the peppermint and vanilla odour cues removed), increasing the need for the rats to rely on the visual cues to succeed on the discrimination test. In contrast, the rats assigned to the Dark condition were tested in complete darkness. In this instance, the rats needed to rely on olfactory cues to complete the discrimination, and the experimenter used night vision goggles to conduct the testing.

The order of the testing days (Standard vs. No-Odour or Dark) was counterbalanced. For each testing condition, the rats were given 2 trials of Discrimination 1, and their selections of S+ were not rewarded (i.e., extinction trials). These trials were then followed by 2 trials of Discrimination 2, in which selection of S+ was rewarded. This sequence was continued until 10 trials for each problem were completed. Subjecting rats to extinction trials for Discrimination 1 allowed confidence that any selection bias in the retention test could be attributed to the rat's memory of the preoperative acquisition, with no influence from post-surgery reinforcement. The interleaving of the rewarded Discrimination 2 trials served to maintain the rat's motivation to displace the objects.

2.7. Histology

Upon completing the behavioural testing, the rats were given an overdose of sodium pentobarbital (0.5 cc; 320 mg/ml, i.p.; Schering Canada Inc., Pointe-Claire, QC, Canada) and

intracardially perfused with 0.9% saline followed by 10% formalin. Their brains were collected

and stored in 10% formalin-30% sucrose for at least 48 h. The brains were sectioned at 40 μm

thickness using a freezing sliding microtome. Every twelfth section of the hippocampus was

mounted on gelatin-coated glass slides, and the sections were stained with Cresyl violet and

cover slipped. Coronal sections spanning the entire hippocampus were imaged and digitized

using a slide scanner (NanoZoomer-RS, Hamamatsu).

Hippocampal damage was estimated using principles of the Cavalieri method, following

the protocol described by Scott et al. (2016), with minor modifications. A systematic sampling

grid (with an area per square of 0.0144 mm^2) was randomly laid over each image, and the

number of points hitting the dentate gyrus granule cell layer or the CA1/CA3 pyramidal cell

layers was counted. Grids were generated using ImageJ software (https://imagej.nih.gov/ij). For

each HPC rat, the total number of hits was divided by the average number of hits obtained from

three SHAM rats (i.e., 1209 hits), resulting in the proportion of remaining HPC tissue. The

complement proportion was used as an estimate of the percentage of HPC damage.

2.8. Statistics

SPSS Statistics (Version 27, IBM) and GraphPad Prism 8 (Version 8.4.3; macOS) were

used for statistical analysis and figure production, respectively. An alpha level of 0.05 was used.

To test whether the rate of S+ responses was significantly greater than would be predicted by

chance (i.e., greater than 50%) within each group, one-tailed one-sample t-tests were used.

Comparisons between two groups were made using two-tailed independent-samples t-tests. A

two-way ANOVA was used to determine if S+ responses in the Discrimination 1 retention test

were affected by lesion (SHAM vs. HPC) or testing condition (Standard vs. No-Odour vs. Dark).

Equal variances were not assumed if Levene's test was significant. Means reported in the text below are presented ± the standard error of the mean (SEM).

3. RESULTS

3.1. HPC Lesions

The NMDA injections produced extensive cell loss across all principal subfields of the HPC (see Fig. 2 for a representative lesion). Estimates of HPC damage in the HPC rats from the No-Odour condition ranged from 84%-95% with a mean of 90% ± 2%. Estimates of HPC damage in the HPC rats from the Dark condition ranged from 67%-97%, with a mean of 87% ± 5%.

Figure 2. Photomicrographs of Nissl-stained sections from a rat brain with a representative lesion, estimated at 88% of the hippocampus. Scale bar represents 5 mm.

3.2. Object Discrimination

3.2.1. Discrimination 1 – Preoperative training.

The SHAM and HPC rats, whether from the No-Odour or Dark condition, learned the preoperative object discrimination problem (Discrimination 1) at a similar rate. The SHAM No-Odour group ($n = 8$) required 96.4 ± 10.6 trials to reach criterion, the SHAM Dark group ($n = 7$) required 88.0 ± 8.7 trials, the HPC No-Odour group ($n = 7$) required 99.9 ± 13.6 trials, and the HPC Dark group ($n = 7$) required 90.0 ± 10.9 trials. A two-way ANOVA revealed no main effect of lesion (SHAM vs. HPC) [$F_{1,25} = 0.06$, $p = 0.807$], no main effect of cue (No-Odour vs. Dark) [$F_{1,25} = 0.68$, $p = 0.418$], and no interaction [$F_{1,25} < 0.01$, $p = 0.947$]. Moreover, the time required to learn the discrimination did not differ across the lesion groups [$F_{1,23} < 0.01$, $p = 0.951$] or cue [$F_{1,23} = 0.43$, $p = 0.516$], with a range of 50-63 min per group. Two time values were missing due to a recording problem – one from the SHAM No-Odour group and one from the HPC Dark group – but the missing times were not noted to be out of range. These results confirm that the rats assigned to each group were properly matched for their acquisition performance.

3.2.2. Discrimination 1 - Postoperative retention.

Figure 3 illustrates the retention performance of the SHAM and HPC rats on Discrimination 1 when tested in the Standard (visual and odour cues present), No-Odour (discrete odour cues removed), and Dark (complete darkness) conditions. The data from the two SHAM groups were pooled for the Standard test because they did not statistically differ [77.5% vs. 74.3% S+ choices; $t_{13} = 0.27$, $p = 0.789$]. The same was done for the data from the two HPC groups on the Standard test, again because of similar performance [81.4% vs. 67.1% S+ choices; $t_{12} = 1.01$, $p = 0.333$]. This pooling simplified presentation of the data and overall analysis.

In the postoperative retention test for Discrimination 1 (Fig. 3), a two-way ANOVA showed that there was no significant main effect of testing condition [$F_{2,52} = 1.22$, $p = 0.303$] or lesion [$F_{1,52} = 0.73$, $p = 0.397$], nor was there an interaction between the two factors [$F_{2,52} = 0.49$, $p = 0.615$]. Comparing the performance of the rats in each condition to chance, one-sample t-tests showed that the SHAM rats could solve the problem under all the conditions, as indicated by selecting S+ at a rate significantly greater than 50% [Standard: $t_{14} = 4.58$, $p < 0.001$; No-Odour: $t_7 = 2.96$, $p = 0.011$; Dark: $t_6 = 2.38$, $p = 0.027$]. In contrast, the HPC rats performed no better than chance under the No-Odour condition, selecting S+ on $55.7 \pm 10.2\%$ of the trials [$t_6 = 0.56$, $p = 0.298$]. However, the HPC rats were able to select S+ at a rate significantly greater than chance when they could rely on odour cues, both in the Standard and Dark conditions [Standard: $t_{13} = 3.43$, $p = 0.002$; Dark: $t_6 = 2.80$, $p = 0.016$].

Figure 3. Postoperative performance of rats on Discrimination 1, tested with or without the explicit odour or visual cues that were present during preoperative acquisition of the problem. For each group, performance was assessed by comparing the rate of S+ selections to the rate predicated by chance (i.e., 50%, dotted line on figure). Whereas the SHAM rats could solve the problem with

odour and visual cues intact (Standard), without the explicit odour cues (No-Odour), and without visual cues (Dark), the HPC rats were selectively impaired in the No-Odour condition. Error bars = SEM.

3.2.3. Discrimination 2 – Postoperative training and retention.

Following surgery and recovery, the SHAM and HPC rats were trained to criterion on a new discrimination problem (Discrimination 2). The HPC rats required significantly more trials to reach criterion than did the SHAM rats [$t_{27} = 2.83$, $p = 0.009$; Fig. 4A]. Despite this difference in acquisition, the performance of the SHAM and HPC rats on the final training session, as assessed by the percentage of S+ selections out of the 20 trials, was very similar [$t_{27} = 0.34$, $p = 0.735$; Fig. 4B]. Furthermore, on the Discrimination 2 retention test, performance did not significantly differ between the HPC and SHAM rats [$t_{27} = 1.87$, $p = 0.072$; Fig. 4C]. These results show that, while the HPC rats required more trials to learn the anterograde discrimination task compared to the SHAM rats, by the end of training and during retention testing the SHAM and HPC rats showed similar performance.

Figure 4. Performance of rats on an object discrimination problem (Discrimination 2) acquired after the rats had received either SHAM or HPC lesions. **A**. Trials to criterion during acquisition of Discrimination 2. The HPC rats required significantly more acquisition trials to reach criterion than did the SHAM rats. **B.** The S+ choice rate during the final session of Discrimination 2 acquisition was similar between the SHAM and HPC rats. **C.** The S+ choice rate during Discrimination 2 retention testing did not differ significantly between groups. Error bars = SEM.

4. DISCUSSION

The purpose of this study was to test whether memory for discrimination problems acquired before receiving a complete, bilateral hippocampal lesion is affected by the presence or absence of odour and visual cues. To this end, rats were trained on an object discrimination problem (Discrimination 1), which could be solved using visual cues, explicit odour cues (i.e., strong, distinct odours applied to the objects by the experimenter), or tactile cues. In principle, the use of implicit odour cues to solve the problem is a possibility, but it is unlikely in the present experiment, because the objects were constructed out of identical materials and so would very likely not differ in scent. The Discrimination 1 problem was successfully acquired by the rats in a single, short training session. Retention for the problem was then tested after hippocampal lesions. In one condition (No-Odour), retention was tested without the explicit odour cues that

were associated with the objects during acquisition. In another condition (Dark), the contribution of visual cues was assessed by testing the rats' retention performance with the lights off and no other light source present, making the room completely dark.

Our key finding was that NMDA lesions encompassing almost the entirety of the hippocampus did not cause retrograde amnesia for the object discrimination problem when rats could use odour cues to make a choice. This was the case even when the rats were tested in the dark, rendering them unable to use visual cues. With the explicit odours present in the retention test, the HPC rats selected S+ at a rate significantly greater than would be predicted by chance. However, when the explicit odour cues were removed during the retention test, forcing the rats to rely on visual or other cues to solve the problem, the HPC damaged rats failed to select S+ at a rate greater than chance, suggesting retrograde amnesia for the object/reward association. The SHAM rats, in contrast, were able to solve the problem without using the explicit odour cues. It is unlikely that visual impairments caused by hippocampal damage contributed to the failure of the HPC rats to solve the discrimination problem in the No-odour condition, because rats with similar hippocampal lesions were not impaired at learning simple visual discriminations in a visual-only task (Driscoll et al., 2005).

Our results suggest that there may be something unique about how the brain stores odour memories, relative to memories derived from vision and perhaps other sensory modalities. Whatever these unique neurobiological features may be, they allowed the odour features of an object/reward association memory, learned with an intact hippocampus, to be successfully recalled without the hippocampus. In contrast, memory for the non-odour features (e.g., colour and shape of the objects) did not survive a complete lesion of the hippocampus. This idea is consistent with previous findings that complete hippocampal lesions cause retrograde object

recognition memory deficits when rats are tested in the novel object preference test using objects without explicit odours (Broadbent et al., 2010; Gaskin et al., 2003). It is also consistent with the observation that hippocampal lesions cause retrograde amnesia for the association between a visual cue (e.g., a picture) and an escape platform in a visual discrimination test (Driscoll et al., 2005; Epp et al., 2008; Lehmann et al., 2021; Sutherland et al., 2001). Finally, it is consistent with previous findings that hippocampal lesions do not cause retrograde amnesia for odour recognition memories (Scott et al., 2013) or in an odour discrimination task (Broadbent et al., 2007).

In several studies, rats with complete hippocampal lesions did not show retrograde amnesia in object discrimination tasks similar to the one used in the present study (Lehmann et al., 2007; Mumby et al., 1999; Wible et al., 1992). In another study, rats with hippocampal lesions performed worse than control rats, but did not show complete retrograde amnesia, as evidenced by the ability to solve the discrimination problem at a rate greater than chance (Broadbent et al., 2007). Importantly, none of these previous studies reported pairing objects with explicit odour cues. According to the interpretation presented above, without explicit odour cues, the rats in these studies should not have been able remember the object/reward association after hippocampal damage. An alternate explanation is required, then, that addresses the question of why the HPC rats in the present study failed to recall the object/reward association when they could not rely on explicit odour cues, whereas rats in previous studies that did not use explicit odour cues performed the task adequately. One possibility is that, in at least some previous studies, the rats with hippocampal lesions may not have needed explicit odour cues because they could rely on implicit ones instead. In the study by Wible et al. (1999), objects made of varying materials (e.g., metal, rubber) were used as discriminanda, potentially making it so that the rats

could discriminate between the objects based on odour. In other studies, the authors reported

using objects with no obvious scents (Mumby et al., 1999) or consistently used objects made of

plastic (Broadbent et al., 2007; Lehmann et al., 2007). However, in these studies it was not

reported that objects made of identical materials were used, as was done for Discrimination 1 in

the present study to minimize the ability of rats to use implicit odour cues to solve the

discrimination problem. Notably, the postoperatively acquired Discrimination 2 used two

different plastic objects, and the rats were able to discriminate between them in complete

darkness, though whether the rats relied on implicit odour cues, tactile cues, or both to solve this

problem is not known.

      Cue overshadowing may offer another explanation. The rats trained on the discrimination

problem without explicit odour cues are, by default, forced to rely on other information, such as

visual and tactile cues, to solve the problem. The rats therefore pay close attention to these cues,

allowing the rats to form a strong memory of the object that is resilient to hippocampal damage.

When rats are trained on the same task *with* explicit odour cues, the rats may rely heavily on

these odours to solve the problem – perhaps because odour cues are particularly salient to a rat –

and pay less attention to the non-odour features of the object. This might result in a weaker

memory of these features, which is sufficiently strong for a normal rat to recall the object/reward

association, but not strong enough for a rat without a hippocampus to do so. In other words, it is

not that rats with hippocampal lesions are incapable of recalling object/reward associations

formed pre-lesion, but rather that the explicit odour cues overshadow the non-odour cues,

preventing the rats from forming a hippocampal-independent memory of the object/reward

association that they *would* have been able to form had the explicit odours not been present.

Regardless of which interpretation is correct, the totality of the evidence suggests that odour memories are less susceptible to hippocampal damage than are similar memories involving non-odour information. Previous studies have shown that complete hippocampal lesions do not affect the ability to form new odour recognition memories and maintain these memories over long (i.e., 5 week) retention intervals, nor do lesions disrupt the ability to recall odour recognition memories acquired preoperatively (Feinberg et al., 2012; Hunsaker et al., 2008; Scott et al., 2013). Similarly, hippocampal damage does not cause retrograde or anterograde amnesia for odour/reward associations in odour discrimination memory tasks (Broadbent et al., 2007; Jonasson et al., 2004; Wood et al., 2004). These findings, in conjunction with the present results, indicate that extrahippocampal systems in the brain are responsible for the long-term maintenance of odour memories. The striatal memory system may be particularly important for remembering odour/reward associations, as lesions of dorsal striatum have been found to cause retrograde amnesia and impair, though not abolish, memory in the anterograde direction on an odour discrimination task (Broadbent et al., 2007).

This is not to say that the hippocampus has no role in odour processing or in odour memory tasks. For instance, lesioning the ventral dentate gyrus with colchicine impairs performance in a delayed-matching-to-sample odour discrimination working memory test (Weeden et al., 2014). The hippocampal damage did not impair ability to differentiate between odours, as indicated by normal performance when the delay between sample and test was short (15 s). However, at a longer delay (60 s), the rats with lesions were impaired, particularly when they were required to distinguish between similar odours. With larger lesions to the ventral hippocampus, rats were impaired even at the 15 s delay, though they could still discriminate between odours in a simple discrimination task (Kesner et al., 2011). Moreover, some studies

have shown retrograde amnesia for odour/flavour memories acquired very shortly (i.e., a couple

of days) before hippocampal lesions are sustained (Tse et al., 2007; Winocur, 1990; Winocur et

al., 2001), suggesting that odour recognition memories may be transiently dependent on the

hippocampus. The possibility should not be dismissed, however, that this short period of

retrograde amnesia may result from disrupted cellular consolidation in structures other than the

hippocampus, as lesions targeted to the hippocampus may temporarily disrupt cellular

consolidation even in off-target structures (Glenn et al., 2005; Rudy & Sutherland, 2008). In any

case, none of these studies contradicts the idea that long-term storage of odour memories,

particularly relatively simple odour recognition or odour discrimination memories, is

independent of the hippocampus.

The story is similar for other brain structures directly upstream of the hippocampus, in

that the bulk of the evidence indicates that these structures are involved in performing certain

odour-based tasks, but each structure on its own is not essential for simple odour memories. The

lateral entorhinal cortex (LEC) is a primary source of input to the hippocampus and receives

projections from olfactory areas like the olfactory bulbs and the piriform cortex (Kerr et al.,

2007). Reversibly inactivating the LEC impairs the performance of rats in an odour

discrimination task when a fine discrimination is required, demonstrating a role for the LEC in

processing odour information (Chapuis et al., 2013). The task used by Chapuis and colleagues

involved making an immediate response after sampling the odours, and so it does not reveal

much about the role of LEC in odour memory; but, aspiration of the LEC does not impair the

ability to form new odour recognition memories or to retain these memories for intervals up to

48 h (Wirth et al., 1998). Transection of the lateral olfactory tract - a major source of olfactory

input to the entorhinal cortex and, therefore, also to the hippocampus - had no effect on

odour/reward association memory in an odour discrimination task (Slotnick & Risser, 1990).

However, when this transection was combined with electrolytic lesions of the mediodorsal

thalamic nucleus, part of the olfactory thalamocortical system, rats did show anterograde

amnesia in this task. In the case of the perirhinal cortex, which also receives input from the

piriform cortex and projects directly and indirectly to the hippocampus (Furtak et al., 2007),

excitotoxic lesions cause anterograde impairments of recognition memory for social odours at

relatively long retention intervals (1-48 h), but not of recognition memory for more distinct non-

social odours (household spices) at the same intervals (Feinberg et al., 2012).

    A secondary finding of the present study was that complete hippocampal lesions do not

prevent learning and memory of new object/reward associations, nor do they impair the ability to

discriminate between distinct objects. This was observed when rats were tested on the

Discrimination 2 problem, which was acquired post-lesion. HPC lesions may have had some

effect on the ability to learn and remember Discrimination 2. Specifically, relative to the SHAM

rats, the HPC rats required more trials to learn this discrimination. However, with additional

training, the HPC rats did eventually reach a level of performance that was very similar to the

level achieved by the SHAM rats. Moreover, on the Discrimination 2 retention test, the HPC rats

did not perform significantly worse than the SHAM rats, and both groups performed well above

chance levels. Thus, complete HPC lesions may cause some mild anterograde object

discrimination learning impairments, but clearly do not cause complete anterograde amnesia for

object/reward association memories. This is consistent with previous findings that rats with

hippocampal lesions are still able to form new object recognition memories, as assessed using

the novel object preference test (Forwood et al., 2005; Gaskin et al., 2003; Mumby et al., 2002,

2005), and new object/reward associations, as assessed using object discrimination tasks

(Broadbent et al., 2007; Lehmann et al., 2007; Mumby et al., 1999). Rats with hippocampal damage have been shown to exhibit small decreases in novel object preference relative to sham-lesioned animals, but object recognition memory was still intact in these lesioned animals, as indicated by preference for the novel object significantly greater than chance (Broadbent et al., 2010). Based on our present findings, and on previous work using the same object discrimination apparatus (Lehmann et al., 2007), there is no reason to believe that learning Discrimination 2 interfered with Discrimination 1.

In summary, we observed that complete hippocampal lesions in rats caused retrograde, but not anterograde, amnesia in an object discrimination task, so long as the task could not be solved using odour cues. When the rats were able to use odour cues to solve the discrimination problem, they were able to perform the task successfully, demonstrating that memory for the odour features of an object/reward association memory is spared following hippocampal damage. When considered alongside previous findings, our results support the idea that object/reward association memories that would normally be hippocampal-independent can be made dependent by pairing the objects with explicit odour cues during acquisition and then removing these odour cues during retention testing.

Acknowledgements: The authors thank Valérie Lapointe for assistance with histology. Funding was provided by NSERC Discovery Grants to HL and RJS.

Conflict of interest statement: The authors declare no conflict of interest.

REFERENCES

Ache, B., & Young, J. (2005). Olfaction: Diverse Species, Conserved Principles. *Neuron*, *48*(3), 417–430. https://doi.org/10.1016/j.neuron.2005.10.022

Aqrabawi, A. J., & Kim, J. C. (2020). Olfactory memory representations are stored in the anterior olfactory nucleus. *Nature Communications*, *11*(1). https://doi.org/10.1038/s41467-020-15032-2

Broadbent, N., & Clark, R. E. (2013). Remote context fear conditioning remains hippocampus-dependent irrespective of training protocol, training-surgery interval, lesion size, and lesion method. *Neurobiology of Learning and Memory*, *106*, 300–308. https://doi.org/10.1016/j.nlm.2013.08.008

Broadbent, N., Gaskin, S., Squire, L., & Clark, R. (2010). Object recognition memory and the rodent hippocampus. *Learning & Memory*, *17*, 5–11. https://doi.org/10.1101/lm.1650110

Broadbent, N., Squire, L., & Clark, R. (2007). Rats depend on habit memory for discrimination learning and retention. *Learning & Memory*, *14*, 145–151. https://doi.org/10.1101/lm.455607

Chapuis, J., Cohen, Y., He, X., Zhang, Z., Jin, S., Xu, F., & Wilson, D. (2013). Lateral entorhinal modulation of piriform cortical activity and fine odor discrimination. *The Journal of Neuroscience*, *33*(33), 13449–13459. https://doi.org/10.1523/JNEUROSCI.1387-13.2013

Driscoll, I., Howard, S., Prusky, G., Rudy, J., & Sutherland, R. (2005). Seahorse wins all races: Hippocampus participates in both linear and non-linear visual discrimination learning. *Behavioural Brain Research*, *164*(1), 29–35. https://doi.org/10.1016/j.bbr.2005.05.006

Epp, J., Keith, J., Spanswick, S., Stone, J., Prusky, G., & Sutherland, R. (2008). Retrograde amnesia for visual memories after hippocampal damage in rats. *Learning & Memory*, *15*, 214–221. https://doi.org/10.1101/ lm.788008

Feinberg, L., Allen, T., Ly, D., & Fortin, N. (2012). Recognition memory for social and non-social odors: Differential effects of neurotoxic lesions to the hippocampus and perirhinal cortex. *Neurobiology of Learning and Memory*, *97*(1), 7–16. https://doi.org/10.1016/j.nlm.2011.08.008

Forwood, S., Winters, B., & Bussey, T. (2005). Hippocampal lesions that abolish spatial maze performance spare object recognition memory at delays of up to 48 hours. *Hippocampus*, *15*(3), 347–355. https://doi.org/10.1002/hipo.20059

Furtak, S., Wei, S.-M., Agster, K., & Burwell, R. (2007). Functional neuroanatomy of the parahippocampal region in the rat: The perirhinal and postrhinal cortices. *Hippocampus*, *17*(9), 709–722.

Gaskin, S., Tremblay, A., & Mumby, D. (2003). Retrograde and anterograde object recognition in rats with hippocampal lesions. *Hippocampus*, *13*(8), 962–969. https://doi.org/10.1002/hipo.10154

Gidyk, D. C., McDonald, R. J., & Sutherland, R. J. (2021). Intact behavioral expression of contextual fear, context discrimination, and object discrimination memories acquired in the absence of the hippocampus. *Journal of Neuroscience, 41*(11), 2437–2446. https://doi.org/10.1523/JNEUROSCI.0546-20.2020

Glenn, M., Lehmann, H., Mumby, D., & Woodside, B. (2005). Differential Fos Expression Following Aspiration, Electrolytic, or Excitotoxic Lesions of the Perirhinal Cortex in Rats. *Neuroscience*, *119*(3), 806–813. https://doi.org/10.1037/0735-7044.119.3.806

Hunsaker, M., Fieldsted, P., Rosenberg, J., & Kesner, R. (2008). Dissociating the roles of dorsal and ventral CA1 for the temporal processing of spatial locations, visual objects, and odors. *Behavioral Neuroscience*, *122*(3), 643–650. https://doi.org/10.1037/0735-7044.122.3.643

Jonasson, Z., Ballantyne, J., & Baxter, M. (2004). Preserved anterograde and retrograde memory of rapidly acquired olfactory discrminations after neurotoxic hippocampal lesions. *Hippocampus*, *14*(1), 28–39. https://doi.org/10.1002/hipo.10146

Kerr, K., Agster, K., Furtak, S., & Burwell, R. (2007). Functional neuroanatomy of the parahippocampal region: The lateral and medial entorhinal areas. *Hippocampus*, *17*(9), 697–708. https://doi.org/10.1002/hipo.20315

Kesner, R., Hunsaker, M., & Ziegler, W. (2011). The role of the dorsal and ventral hippocampus in olfactory working memory. *Neurobiology of Learning and Memory*, *96*(2), 361–366. https://doi.org/10.1016/j.nlm.2011.06.011

Lee, J., Zelinski, E., McDonald, R., & Sutherland, R. (2016). Heterarchic reinstatement of long-term memory: A concept on hippocampal amnesia in rodent memory research. *Neuroscience & Biobehavioral Reviews*, *71*, 154–166.

Lehmann, H., Glenn, M., & Mumby, D. (2007). Consolidation of object-discrimination memory is independent of the hippocampus in rats. *Experimental Brain Research*, *180*, 755–764. https://doi.org/10.1007/s00221-007-0895-2

Lehmann, H., Lecluse, V., Houle, A., & Mumby, D. (2006). Retrograde amnesia following hippocampal lesions in the shock-probe conditioning test. *Hippocampus*, *16*(4), 379–387.

Lehmann, H., Sparks, F., O'Brien, J., McDonald, R., & Sutherland, R. (2010). Retrograde amnesia for fear-potentiated startle in rats after complete, but not partial, hippocampal

damage. *Neuroscience*, *167*(4), 974–984.

https://doi.org/10.1016/j.neuroscience.2010.03.005

Lehmann, H., Stykel, M. G., & Glenn, M. J. (2021). Overtraining Strengthens the Visual

Discrimination Memory Trace Outside the Hippocampus in Male Rats. *Frontiers in

Behavioral Neuroscience*, *15*, 768552. https://doi.org/10.3389/fnbeh.2021.768552

McDonald, R. J., Balog, R. J., Lee, J. Q., Stuart, E. E., Carrels, B. B., & Hong, N. S. (2018). Rats

with ventral hippocampal damage are impaired at various forms of learning including

conditioned inhibition, spatial navigation, and discriminative fear conditioning to similar

contexts. *Behavioural Brain Research, 351*, 138–151.

https://doi.org/10.1016/j.bbr.2018.06.003

Morris, R. G. M., Garrud, P., Rawlins, J. N. P., & O'Keefe, J. (1982). Place navigation impaired

in rats with hippocampal lesions. *Nature*, *297*(5868), 681–683.

https://doi.org/10.1038/297681a0

Mumby, D., Astur, R., Weisend, M., & Sutherland, R. (1999). Retrograde amnesia and selective

damage to the hippocampal formation: Memory for places and object discriminations.

Behavioural Brain Research, *106*, 97–107.

Mumby, D., Gaskin, S., Glenn, M., Schramek, T., & Lehmann, H. (2002). Hippocampal Damage

and Exploratory Preferences in Rats: Memory for Objects, Places, and Contexts.

Learning & Memory, *9*, 49–57. https://doi.org/10.1101/lm.41302

Mumby, D., Pinel, J., & Wood, E. (1990). Nonrecurring-items delayed nonmatching-to-sample

in rats: A new paradigm for testing nonspatial working memory. *Psychobiology*, *18*, 321–

326.

Mumby, D., Tremblay, A., Lecluse, V., & Lehmann, H. (2005). Hippocampal damage and anterograde object-recognition in rats after long retention intervals. *Hippocampus*, *15*(8), 1050–1056. https://doi.org/10.1002/hipo.20122

Quinn, J. J., Ma, Q. D., Tinsley, M. R., Koch, C., & Fanselow, M. S. (2008). Inverse temporal contributions of the dorsal hippocampus and medial prefrontal cortex to the expression of long-term fear memories. *Learning & Memory, 15*(5), 368-372. https://doi.org/10.1101/lm.813608

Rudy, J., & Sutherland, R. (2008). Is it systems or cellular consolidation? Time will tell. An alternative interpretation of the Morris group's recent science paper. *Neurobiology of Learning and Memory*, *89*(4), 366–369. https://doi.org/10.1016/j.nlm.2007.07.017

Scott, G., Mtetwa, M., & Lehmann, H. (2013). Novel odour recognition memory is independent of the hippocampus in rats. *Experimental Brain Research*, *224*, 199–209. https://doi.org/10.1007/s00221-012-3304-4

Scott, G., Saucier, D., & Lehmann, H. (2016). Contrasting the Amnesic Effects of Temporary Inactivation with Lesions of the Hippocampus on Context Memory. *Journal of Behavioral and Brain Science*, *6*(4), 184–198.

Slotnick, B., & Risser, J. (1990). Odor memory and odor learning in rats with lesions of the lateral olfactory tract and mediodorsal thalamic nucleus. *Brain Research*, *529*(1–2), 23–29. https://doi.org/10.1016/0006-8993(90)90807-N

Sutherland, R., Weisend, M., Mumby, D., Astur, R., Hanlon, F., Koerner, A., Thomas, M., Wu, Y., Moses, S., Cole, C., Hamilton, D., & Hoesing, J. (2001). Retrograde Amnesia After Hippocampal Damage: Recent vs. Remote Memories in Two Tasks. *Hippocampus*, *11*, 27–42.

Tse, D., Langston, R., Kakeyama, M., Bethus, I., Spooner, P., Wood, E., Witter, M., & Morris, R. (2007). Schemas and Memory Consolidation. *Science*, *316*(5821), 76–82. https://doi.org/10.1126/science.1135935

Weeden, C., Hu, N., Ho, L., & Kesner, R. (2014). The role of the ventral dentate gyrus in olfactory pattern separation. *Hippocampus*, *24*(5), 553–559. https://doi.org/10.1002/hipo.22248

Wible, C., Shiber, J., & Olton, D. (1992). Hippocampus, fimbria-fornix, amygdala, and memory: Object discriminations in rats. *Behavioral Neuroscience*, *106*(5), 751–761. https://doi.org/10.1037/0735-7044.106.5.751

Winocur, G. (1990). Anterograde and retrograde amnesia in rats with dorsal hippocampal or dorsomedial thalamic lesions. *Behavioural Brain Research*, *38*(2), 145–154. https://doi.org/10.1016/0166-4328(90)90012-4

Winocur, G., McDonald, R., & Moscovitch, M. (2001). Anterograde and retrograde amnesia in rats with large hippocampal lesions. *Hippocampus*, *11*(1), 18–26. https://doi.org/10.1002/1098-1063(2001)11:1<18::AID-HIPO1016>3.0.CO;2-5

Winocur, G., Moscovitch, M., Caruana, D. A., & Binns, M. A. (2005). Retrograde amnesia in rats with lesions to the hippocampus on a test of spatial memory. *Neuropsychologia, 43*(11), 1580-1590.

Winocur, G., Sekeres, M. J., Binns, M. A., & Moscovitch, M. (2013). Hippocampal lesions produce both nongraded and temporally graded retrograde amnesia in the same rat. *Hippocampus*. https://doi.org/10.1002/hipo.22093

Wirth, S., Ferry, B., & Di Scala, G. (1998). Facilitation of olfactory recognition by lateral entorhinal cortex lesion in rats. *Behavioural Brain Research*, *91*(1–2), 49–59. https://doi.org/10.1016/S0166-4328(97)00102-2

Wood, E., Agster, K., & Eichenbaum, H. (2004). One-Trial Odor-Reward Association: A Form of Event Memory Not Dependent on Hippocampal Function. *Behavioral Neuroscience*, *118*(3), 526–539. https://doi.org/10.1037/0735-7044.118.3.526