# Parthenogenetic Panagrolaimus species evolve at lower mutation rates, but show increased nucleotide diversity

Laura I. Villegas[1,], Luca Ferretti[2], Thomas Wiehe[3], Ann-Marie Waldvogel[1,*], and Philipp H. Schiffer[1,*] *Both authors had equal contributions

[1]Institute for Zoology, University of Cologne, 50674 Köln, Germany
[2]Big Data Institute, University of Oxford, Oxford, United Kingdom
[3]Institute for Genetics, University of Cologne, Köln, 50674 Germany
Correspondence: lvilleg1@uni-koeln.de

May 1, 2023

**Abstract**

Asexual reproduction is assumed to lead to the accumulation of deleterious mutations, and reduced heterozygosity due to the absence of recombination. Panagrolaimid nematodes display different modes of reproduction. Sexual reproduction through distinct males and females, asexual reproduction through parthenogenesis found in the genus *Panagrolaimus*, and hermaphroditism, found in *Propanagrolaimus*. Here, we compared genomic features of free-living nematode populations in strains and species isolated from geographically distant regions to study genomic diversity and genome-wide differentiation under different modes of reproduction. We firstly estimated genome-wide spontaneous mutation rates per genome for a polyploid parthenogenetic *Panagrolaimus* strain and a diploid hermaphroditic *Propanagrolaimus* species via mutation-accumulation-lines. Secondly, we calculated population genomic parameters including nucleotide diversity and fixation index ($F_{ST}$) between populations of asexually and sexually reproducing nematodes. Thirdly, we used phylogenetic network methods on sexually and asexually reproducing *Panagrolaimus* strains to understand evolutionary relationships between them. The estimated mutation rate was slightly lower for the asexual strain, as expected for taxa with this reproductive mode. Natural polyploid asexual strains revealed higher nucleotide diversity. Despite their common ancestor, a gene network revealed a high level of genetic differentiation among asexual strains. The elevated heterozygosity found in the triploid parthenogens could be explained by the third genome copy. Given their tendentially lower mutation rates it can be hypothesized that this is part of the mechanism to evade Muller's ratchet. Our findings in parthenogenetic triploid nematode populations seem to challenge common expectations of evolution under asexuality.

**Keywords: Nematodes, parthenogenesis, mutation rate, diversification, population genomics, asexuality.**

*Dedicated to the memory of Einhard Schierenberg.*

# 1   Introduction

Parthenogenetic organisms reproduce asexually, without paying the cost of sex, i.e. production of males, courtship or mate finding. Consequently, parthenogenetic reproduction should result in an evolutionary advantage (Otto & Lenormand, 2002), since asexually reproducing organisms are capable of generating more offspring from the same resources, than their sexual sister species. The Red Queen Hypothesis (RQH), originally focused on coevolution in a host-parasite context, has been extended to propose that sexually reproducing organisms have an evolutionary advantage in habitats with many biotic interactions, while asexually reproducing taxa are expected to be more frequent in habitats with challenging environmental conditions and lowered biotic preassure (e.g. higher altitudes) (Hartmann et al., 2017; Van Valen, 1973).

However, sexual reproduction is the predominant reproductive mode among Metazoans, leading to a paradox that was coined as the "Queen of Evolutionary questions" by Graham Bell (Bell, 1982). To explain the apparent "lack" of parthenogenetic taxa, evolutionary theory posits that the absence of recombination inhibits the efficient purging of deleterious mutations which accordingly accumulate over time. This effect, called Muller's ratchet (or mutational meltdown e.g. for RNA viruses) (Gabriel, Lynch, & Burger, 1993) (Felsenstein, 1974) counteracts the long-term persistence of obligate asexual lineages lacking recombination. Additionally in parthenogens, due to the lack of outcrossing, alleles can't be recombined and the linkage between sites in finite populations would reduce the overall effectiveness of selection. This phenomenon, described as Hills-Robertson interference, (Hill & Robertson, 1966) suggests that asexuality hinders local adaptation (Barton, 1994) (Engelstädter, 2008).

Mutation rates, the rate at which mutations arise *de novo* in a genome, are also proposed to vary between organisms with different reproductive modes. As a result of the strong selection that favours DNA repair and replication mechanisms and, on shorter time scales, the purging of deleterious mutations as is the majority of non-synonymous mutations, mutation rates are generally low throughout the animal kingdom but vary between species (Lynch, 2010). It has been proposed that asexually reproducing organisms have lower mutation rates than sexually reproducing relatives (Sloan & Panjeti, 2010).

Empirical studies paint a different, more complex picture: Bdelloid rotifers, oribatid mites and Darwinulid ostracods are examples of the so called 'evolutionary scandals' that prove the long-term persistence of asexual taxa (Smith, 1978). Studies on asexual populations of monkey flowers and parasitoid wasps show how asexual taxa can locally adapt to their environment (Amat, van Alphen, Kacelnik, Desouhant, & Bernstein, 2017)(Gibson, 2019) (Simón-Porcar, Silva, & Vallejo-Marín, 2021). Despite the expected reduced genetic diversity in parthenogenetic taxa, a study on the triploid asexual crayfish *Procambarus virginalis*, has shown that this species is highly heterozygous even when the expected outcome of its reproductive mechanism would be homozygosity (Schwarz, 2017). It might thus be possible for natural populations of asexually reproducing organisms to genetically diverge due to local adaptation, genetic drift, geographic isolation, and the generation of distinct mutations in different populations.

Nematoda are one of the most species-rich groups within Metazoa. The phylum is highly diverse in terms of ecological niches inhabited, size ranges of different groups and reproductive strategies (Blaxter, 2011). The wide distribution of species in the phylum opens the opportunity to perform studies not only in terrestrial ecosystems, but also in fresh waters and the marine environment. The short generation times in many species, and relative ease of establishing large cultures in the laboratory allows researchers to perform experiments to answer ecological and evolutionary questions. Here we focused on studying members of the Panagrolaimidae family, where gonochoristic (sexual), and parthenogenetic (asexual), as well as hermaphroditic (sexual, but selfing) reproduction can be found (Kiontke et al., 2004). This system allows to compare closely related taxa with different reproduction modes providing information on how their genomic features may differ accordingly. In this study we wanted to understand how mutation rates vary in free-living closely related species with different reproduction modes (sexual, but selfing, and asexual) and their genomic traits, such as nucleotide diversity and population divergence between the different mutation accumulation lines (MAL) and parental state. We furthermore extended the study of these genomic traits to natural populations of diploid sexual and triploid parthenogenetic nematodes of the genus *Panagrolaimus*. We tested whether the results obtained in bottle-necked laboratory strains held true for natural populations from different geographical locations, and if these patterns of genome evolution in parthenogens are consistent with theoretical expectations or challenge them.

## 2   Materials and Methods

### 2.1   Sampling, sequencing and data pre-processing

This study includes natural *Panagrolaimus* strains from different geographic locations that were previously described in McGill et al. 2015 (McGill et al., 2015) (table 1). Asexual nematode strains used here are triploid (allopolyloids) and sexual strains are diploid selfing hermaphrodites, in which no males have been observed so far. Strains were kept as laboratory cultures (at 15°C on low nutrient agar plates inoculated with OP50 *(E. coli)* from

which DNA was extracted from several plates. Adult nematodes, as well as larvae and eggs were washed off from the plates and cleaned in 3 washing steps. After three rounds of freeze-thaw cycles on lysis buffer, genomic DNA was extracted following a salt-out protocol or using Qiagen's genomic tip. Whole genome sequencing of pooled specimens was performed on Illumina HiSeq2000 and NovaSeq platforms, sequencing data was deposited in SRA and is available under the Bioproject PRJNA374706 ([Dataset]Schiffer, 2017). After standard quality filtering and trimming of raw reads using fastp (versions 0.23.0 and 0.20.1) (Chen, Zhou, Chen, & Gu, 2018), paired-end reads were mapped to the respectively closest related reference assembly (available for strains PS1159, JU765, and ES5 (*P. sp.* 'bornheim'), table1) using bwa-mem2 (version 2.2.1) (Vasimuddin, Misra, Li, & Aluru, 2019). For strains where reads were too short, mapping was done using NextGenMap (version 0.5.5) (Sedlazeck, Rescheneder, & von Haeseler, 2013). For strains where the insert size was smaller than double the read length, pear (version 0.9.8) (Zhang, Kobert, Flouri, & Stamatakis, 2014) was used before mapping with bwa-mem2. The alignments were filtered to remove duplicates using PICARD tools (MarkDuplicatesWithMateCigar)(version 2.26.8) (Institute, 2018), and low-quality reads (<30) were removed using samtools view (version 1.13) (Danecek et al., 2021). Asexual strain PS1159 and sexual strain JU765 were used in the mutation accumulation lines experiment for estimation of mutation rates. All other sexual and asexual strains were used in the population analysis and phylogenetic network. A list of the commands implemented in the following segments of Materials and Methods is provided here: https://github.com/lauraivillegasr/parthenogenomics.

## 2.2 Estimation of mutation rates from a MAL experiment

For the estimation of mutation rates, an asexual strain *Panagrolaimus* sp. PS1159 and a sexual selfing strain *Propanagrolaimus* sp. JU765 were subjected to a mutation accumulation lines (MALs) experiment for 30 to 52 generations of continuous inbreeding. Whole genome sequencing data was then generated from the starting point of the experiment and the end point of each MAL (Figure 1). DNA extraction was performed as described above. To allow for sufficient DNA content, several libraries of the respective strain at the start of the MAL experiment were pooled into one, hereafter referred to as the *RefPool*. To ensure all quality scores were in Sanger encoding seqret (Madeira et al., 2019) and fastQC (version 0.11.9) (Andrews, 2010) were used. Mutations were called using a probabilistic approach (accuMUlate version 0.2.1)(Winter et al., 2018), following authors recommendations. Alignments from MAL were merged using samtools merge along with the RefPool. The analysis was performed only on a reduced set of positions commonly covered between all the lines and parental state per reproductive mode, respectively. Putative mutations were filtered by coverage ranges, number of reads supporting mutations, absence of mutant allele in other samples, absence of mutation in RefPool, apparent mutation being caused by mismapped reads (Anderson-Darling test statistic), read pair successfully mapped to the reference genome (Fisher's exact test)(specific values used for filtering can be found under: https://github.com/lauraivillegasr/parthenogenomics). Resulting candidates were manually curated using the Integrative Genomics Viewer (IGV version 2.16.1) (Thorvaldsdottir, Robinson, & Mesirov, 2013). The number of callable sites was estimated for each MAL as the number of positions along the assembly within the depth coverage of 10 and 50. The mutation rates were obtained by dividing the number curated *de novo* mutations by the total number of callable sites using

$$\mu = \frac{\text{called mutations}}{\text{generations} \times \text{callable sites}}. \tag{1}$$

High-credibility intervals (HCI) for the estimated mutation rates were obtained using the Bayesian First Aid R package (Bååth, 2014). Mutation rates estimated for each of the MAL, as well as the average mutation rate for both reproductive modes along with the average number of callable site per strain, were provided as input for this.

## 2.3 Analysis of populations

### 2.3.1 Population parameters

Population genomic parameters $\theta_w$, nucleotide diversity ($\pi$), effective population size $N_e$, and fixation index $F_{ST}$ were estimated in two different settings (A) intrinsic estimation of $F_{ST}$ in MA lines between each other and correspondent parental state and (B) between natural populations from different geographic origins (sexual populations were compared against asexual populations) (table1). Both approaches aimed to understand if patterns in found in long kept laboratory strains are similar to those found in natural populations as well as understanding how populations parameters vary between deferentially reproducing taxa.

To account for ploidy in the triploid asexuals, coverage ranges per genome copy were selected for all further analysis based on coverage distribution of the reads mapped against any of the reference genomes (supplementary figures 1-3). For (A) MAL data a coverage range of 15X per genome copy was selected (10X-55X for asexual triploid lines and 10-40 for sexual diploid lines). For (B) natural populations a coverage range of 23X per genome copy was selected (10X-80X for asexual triploid populations and 10-56 for sexual diploid populations). Pileup files were created individually for each genome data set of MALs and natural populations, respectively, using samtools mpileup (Danecek et al., 2021). A sync file was obtained using mpileup2sync.jar from Popoolation2 (version 1.201) (Kofler, Pandey, & Schlotterer, 2011) from all data sets together to maintain only all common positions, which served as input for the estimation of the fixation index $F_{ST}$ on non-overlapping 1kb windows using Fst-sliding.pl. An approximate size of 3000 individuals per population was used.

$F_{ST}$ was also estimated for 5 MAL of population size N=100 of *C. elegans* (Konrad, Brady, Bergthorsson, & Katju, 2019) from the Bioproject PRJNA448413 ([Dataset]Konrad, 2018) to compare our hermaphroditic sexual strain to the model organism.

Individual pileup files were used as input for Variance-sliding.pl from Popoolation (Kofler, Orozco-terWengel, et al., 2011), using the options –measure theta and –measure pi respectively. The effective population size was then obtained as $N_e = \theta/2n\mu$ for the sexual populations and asexual populations, where $\mu$ is the mutation rate estimated as described above and n corresponds to the ploidy of the studied system (n=2 for sexuals and n=3 for asexuals).

Theta of homologous genome copies (diploid part of the triploid genome) was estimated using Tetmer (version 2.2.1) (Becher et al., 2020). A k-mer spectrum of the Illumina reads per population was obtained using the K-mer analysis toolkit (KAT version 2.4.2) (Mapleson, Garcia Accinelli, Kettleborough, Wright, & Clavijo, 2016) (K=27) and provided to Tetmer as input. The manual fitting mode was used for both reproductive modes, for asexuals the triploid allopolyploid model (AAB) was selected, the diploid (AA) model was selected for sexual populations.

Plots for visualizing $\theta_w$, $\pi$, $F_{ST}$ and effective population size results were obtained using the R package ggplot2 (Wickham, 2016). Statistical analyses to test for significance between the results obtained from each of the populations were done using Welch's t-test. Welch's t-test is used since sample sizes are unequal (more asexual populations than sexual populations) to compare the mean values of $F_s t$, $\pi$, $\theta_w$ for the different reproductive modes.

### 2.3.2 Phylogenetic network construction

A phylogenetic network was constructed for the parthenogenetic strains to test whether they can be defined as phylogenetically distinct species due to their genetic divergence. A phylogenetic network was also constructed for the sexual species as a reference. Orthologues present in all asexual and in all sexual strains were used for this analysis, each reproductive mode was assessed separately. Coordinates of the orthologue genes were obtained for each reference genome using Benchmarking Universal Single-Copy Orthologs (BUSCO) (Simão, Waterhouse, Ioannidis, Kriventseva, & Zdobnov, 2015). The coordinates of the genes on the reference genome extract single gene bam files, from alignments produced as previously described, for each strain using samtools. To obtain a consensus sequence for each gene, bcftools (version 1.13) mpileup and bcftools call were used for variant calling (Danecek et al., 2021). These served as input for GATK (FastaAlternateReferenceMake)(version 4.2.3.0))(Auwera & O'Connor, 2020) to obtain a consensus sequence per strain for each gene, a multiple sequence alignment was obtained using MAFFT (version 7.47.1) (Katoh, 2002). A gene network was obtained for the consensus gene sequences of orthologues present in all the populations per reproductive mode using the median network algorithm from Splitstree4 (Huson, 1998).

## 3 Results

### 3.1 The spontaneous mutation rate is marginally lower in parthenogens

We conducted a mutation accumulation line (MAL) experiment with an asexual (triploid) and a sexual strain (diploid) (Figure 1). MALs were maintained for up to 40 generations, with many lines being lost over the course of the experiment. Starting with 100 and 71 lines respectively, 30 (30%) sexual lines and 15 (21%) parthenogenetic lines survived. Of these, six lines per reproduction mode were randomly chosen for whole genome sequencing and mutation calling. High-confidence mutations were called from 31,735 and 25,812 common positions in asexual and sexual lines, respectively. For the asexual lines, 72375 candidate mutations were called, whereas 7852 candidates were found for the sexual lines. After filtering for coverage ranges, read support, unique mutations per line and miss-mapped regions, and manual curation, 11 DNMs (De novo mutations) with high support remained for the asexual lines and 10 DNMs for the sexual ones (supplementary table 3).

Per reproductive mode five MAL showed at least one DNM. For the asexual lines an average of $7.5 \times 10^{10}$ callable sites were found and $5.6 \times 10^{10}$ for the sexual lines (supplementary table 3). This resulted in a total mutation rate of $5.828 \times 10^{-10}$ (Bayesian HCI $1.2 \times 10^{-17}$, $5.1 \times 10^{-9}$) for asexual Panagrolaimus and a respective mutation rate of $8.923 \times 10^{-10}$ (Bayesian HCI $2.9 \times 10^{-17}$, $6.9 \times 10^{-9}$) for the sexual Panagrolaimus. The mean rate for asexuals is lower than that found for sexuals, however, due to wide credibility intervals, mutation rates between asexual and sexual nematodes do not differ significantly. The majority of DNMs were transitions, with a transition/transversion ratio of 4.5 (9:2) in the asexual lines and 1.5 (6:4) in the sexual lines (supplementary table 5).

### 3.2 Population genetic patterns differ between asexual and sexual MAL

Standing nucleotide diversity, i.e. $\pi$ of the respective parental lines, is lower in the sexual srefpool than in the asexual arefpool (Figure 2).

Estimating population genetic parameters for MALs of both reproduction modes revealed significantly lower $\theta_w$ ($p < 0.0005$) in sexual lines (range 0.0022-0.0029 - lowest and highest $\theta_w$ values found in the different sexual lines) when compared to the asexual lines (range 0.0029-

0.0037 - lowest and highest $\theta_w$ values found in the different asexual lines). Sexual lines are also significantly lower in nucleotide diversity (range 0.0024-0.0036)($p$value $= 0.001315$) with the lowest value found in the sexual parental strain , when compared to asexual lines. Nucleotide diversity of asexual lines ranges between 0.0032 and 0.0046, with the lowest $\pi$ found in one of the asexual MAL (al4), which is also the asexual line with the lowest $\theta_w$.

Comparisons of population differentiation among parental and descendant lines per reproduction mode revealed distinct differences. The sexual lines showed a significantly higher $F_{ST}$ between parental and descendant lines when compared to the $F_{ST}$ among descendant lines ($p$-value$== 6e{-}7$). For the asexual lines, this comparison was not significant. In general, genome-wide population differentiation was found to be more pronounced among asexual lines (including parental arefpool - PS1159parental) with $F_{ST}$ ranging between 0.1312 and 0.1755 in comparison to the $F_{ST}$ range in sexual lines (0.0304-0.0796). Population differentiation was significantly higher in asexual lines than in sexual lines ($p$-value $=7.5 \times 10^{24}$). $F_{ST}$ values found on the sexual descendant lines were similar to those estimated for *C. elegans* descendant lines (population size 100)(supplementary table 9) from a MAL experiment performed by Konrad et al. (2019).

## 3.3 High genetic diversity but low differentiation in natural populations of parthenogens

Nucleotide diversity $\pi$ for asexual populations ranged from 0.0402963 (Raleigh, USA; apop2) to 0.206227 (Beijing, China; apop3). For sexual populations $\pi$ ranged from 0.00241247 (Guangxi, China; spop4) to 0.0545159 (Cologne, Germany; spop3). Nucleotide diversity was on average higher for asexual populations ($p$-value$= 0.0145$) (Figure 3) (supplementary table 10).

Genome-wide estimator $\theta_w$ for asexual populations ranged from 0.0306732 (Diourbel, Senegal; apop1) to 0.192175 (Beijing, China; apop3). For sexual populations $\theta_w$ ranged from 0.0027009 (Guangxi, China; spop4) to 0.0452624 (Cologne, Germany; spop3). Asexual populations showed a significantly higher $\theta_w$ when compared to sexual populations ($p$-value $= 0.0242$).

To test whether the higher diversity found on asexuals could be caused by the third genome copy, estimator $\theta_w$ was obtained for homologous genome copies (asexual triploid *Panagrolaimus* have a hybrid origin). For triploid asexual populations, the mean $\theta_w$ obtained for homologous copies (2 genome gopies) was generally lower than genome wide $\theta_w$ estimations (3 genome copies), but not significantly different ($p$value $= 0.0747$).

### 3.3.1 Population differentiation of natural populations

In natural populations, population differentiation $F_{ST}$ between asexual Diourbel strain, Senegal (apop1) and asexual Corvallis strain, USA (apop7) was highest with an $F_{ST}$ of 0.468 for asexual populations. For sexual populations, the highest level of population differentiation was found between the Santiago Island strain, Cape Verde (spop1) and the Cologne strain, Germany (spop3) with an $F_{ST}$ of 0.851948. Population differentiation was more pronounced among sexual populations as compared to among asexual populations ($p$value$=0.00002471$).

## 3.4 Asexual *Panagrolaimus* strains are genetically distinct

To analyse divergence between the monophyletic asexual *Panagrolaimus* strains under phylogenetic species concept using a network approach in Splits Tree (Huson, 1998). In the asexual

reference genome (*Panagrolaimus sp.* PS1159) we idendtified 2173 universal single-copy or-
thologues. For each strain, reads corresponding to the coordinates of the orthologues were
extracted in order to generate consensus sequences for genes with sufficient read support: 375
orthologues were shared on the asexual populations, these genes were aligned and used on
the gene network. We conducted a corresponding analysis for the sexual species as proof of
principle. For the sexual reference (*Panagrolaimus sp.* bornheim) genome 1983 orthologues
were found and reads corresponding to these genes' coordinates were extracted; a total of 213
orthologues were used for the gene network. The Median network calculated in Splits Tree
showed a total of 23 splits between the asexual strains, while there were 3 splits between the
sexual species, i.e. the minimum possible number of splits (Figure 4).

# 4    Discussion

In this study we aimed to analyze divergence patterns in the evolutionary context of asexual
animals using parthenogenetic *Panagrolaimus* nematodes as a system, and sexual, in this
case fully selfing hermaphroditic *Propanagrolaimus* strains, as a comparator(Figure 5). By
running long-term MAL we conducted a classical experiment to measure mutation rates
under nearly neutral evolution (Halligan & Keightley, 2009). We also estimated standard
population genomic parameters from natural populations of widely-spread geographic origin
(Figure 3)(gonochoristic sexual *Panagrolaimus* and parthenogenetic asexual *Panagrolaimus*),
and used phylogenetic networks (Figure 4) to narrow the complexity of population genomic
patterns under parthenogenesis.

## 4.1    Decreased mutation rates of parthenogens could aid to diminish the effect of Muller's ratchet

Parthenogenesis is assumed to have low evolutionary potential due to the accumulation of
deleterious mutations (Muller's Ratchet) (Muller, 1932), lack of recombination and low ge-
netic diversity (Simon et al. 2003). In *Panagrolaimus*, the parthenogenetic strains analysed
here appear to be monophyletic, originating 1.3–8.5 Mya ago through hybridisation involv-
ing triploidization (Schiffer et al., 2019). Whereas sexual strains and species in our analysis
undergo normal meiosis, eggs of the nematode *Panagrolaimus sp.* PS1159 (arefpool and
apop2) develop without fertilization and the offspring is exclusively female. Asexual meiosis
(presence of polar body) occurs without recombination [pers. comment, Caroline Blanc and
Marie Delattre, Lyon] and as a result, offspring are clones of their mother. The majority
of non-neutral mutations that occur are deleterious (Sloan & Panjeti, 2009). To explain
long-term persistence of asexuals it can be assumed that they have lower mutation rates,
thus the process of mutation accumulation is slowed down. Although the estimated mutation
rates did not differ significantly between sexual and asexual *Panagrolaimus* strains, genetic
diversity in the MAL was found to be much higher in asexual *Panagrolaimus* than in sexual
*Propanagrolaimus*.

If mutation rates are low and recombination is absent, clonal interference can be reduced
when beneficial mutations in a population arise. With higher mutation rates, if many bene-
ficial mutations are carried in different clones, fixation time of these mutations is increased
(De Visser & Rozen, 2005). In our study, mutation rates found for both reproductive modes
are one order of magnitude smaller than that reported for the distantly-related nematode *C.
elegans* (Denver et al., 2009) and for the arthropod model *Drosophila melanogaster* (Keight-
ley, Ness, Halligan, & Haddrill, 2014). This not only shows that findings in model organisms
cannot be generalized across phyla, but also indicates divergent evolutionary mechanisms in

both panagrolaimid nematodes analysed here. To better understand these, it will be necessary to estimate mutation rates with additional MALs, run for shorter accumulation periods to reduce variation of the data and thus allowing for better statistical validation (Oppold & Pfenninger, 2017).

## 4.2 Reproduction modes impacts on patterns of population differentiation

Asexual reproduction may affect population differentiation in different ways and accordingly patterns are expected to differ between asexual and sexual species (Balloux, Lehmann, & de Meeûs, 2003; Prugnolle & De Meeûs, 2008). For a theoretical example of a clonal (=asexual) diploid parasitic species, it is expected that populations are less differentiated to their parents (preceeding generations) than to other populations (Prugnolle & De Meeûs, 2008). This is because clonal reproduction tends to decrease the average expected heterozygosity within populations but does not change, in average, the expected heterozygosity overall populations. In our study the triploid asexual MAL did not produce a distinct pattern of differentiation that would differ between a MA line and the parental state or among MA lines (Figure 2). For sexual reproduction, highly inbred laboratory selfing nematode strains previously described, such as *C. briggsae* and *C. elegans*, show low heterozygosity and a higher differentation between populations and the parental state than among populations, which is a result of alternative alleles being fixed separately in each lineage (Barrière et al., 2009; Teotónio, Estes, Phillips, & Baer, 2017). This pattern, being opposite to the theoretically expected pattern of differentiation in asexuals, aligns with reduced heterozygosity, which we observed in the hermaphroditic (=sexual).

The lower rate of differentiation among sexual lines from the MAL experiment in comparison to asexual lines could be explained by (i) lower standing genetic variation in the hermaphroditic parent at the beginning of the experiment and (ii) increased heterozygosity for asexuals arising from the third genome copy. In asexual lines "inbreeding" does not change the genetic structure of a population (Bengtsson, 2003). The pattern of population differentiation we found for asexual MA lines could show the differentiation pattern of natural populations: maintained heterozygosity within offspring due to the lack of allele segregation (Stoeckel & Masson, 2014). It has been proposed that under asexuality, heterozygosity can even increase since alleles of a same gene could independently accumulate mutations over generations, the so-called Meselson effect. This has recently been observed in obligate asexual oribatid mites (Brandt et al., 2021), but could not be tested in the nematodes with our current data.

## 4.3 Heterozygosity is maintained in natural asexual populations

Studies on populations of the sexual parasitic nematode *Baylisascaris schroederi* isolated from different mountain ranges, showed very low and non-significant $F_{ST}$ (range: 0.01911 to 0.02875) between three populations (Zhou et al., 2013). Natural populations of sexual *C. brenneri* also show low values of differentiation between populations, in this case, from geographically distant regions eastern India and French Guiana (0.092)(Dey, Chan, Thomas, & Cutter, 2013). Our results reveal contrasting patterns of on average higher differentiation among natural sexual populations. In our study, both sexual and sexual populations have two-fold higher $F_{ST}$ values than those reported for *Caenorhabditis* nematodes and *Baylisascaris schroederi*. These higher $F_{ST}$ values could be an initial indication for local adaptation (Hirase et al., 2021) to the distinct regions where the populations were isolated from. It has been previously shown that asexual taxa can in fact adapt to their local environment, as is the case of paristoid wasps: asexual parasitoid wasps have a higher investment on production of

eggs on food-depleted environments for adults, whereas the sexual counterpart often found in food-rich environments with patchy distribution of hosts show a higher investment in flight capacity and have a higher longevity(Amat et al., 2017).

## 4.4 Natural populations of asexual nematodes show higher genomic diversity than their sexual counterparts

Population genomic parameters obtained for asexual natural populations showed a mean nucleotide diversity higher than that found for sexual populations. This pattern contradicts the expectation that the lack of recombination is assumed to result in reduced genetic diversity in parthenogenetic taxa compared to sexually reproducing counterparts (Charlesworth & Willis, 2009). However, the same conflictual pattern of increased nucleotide diversity has also been found for the amazon molly, an asexual fish of hybrid origin (Jaron et al., 2021), as well as in the asexual bark lice (*Echmepteryx hageni*) (Shreve, Mockford, & Johnson, 2011) when compared to respective sexual counterparts.

$\theta_w$ was also found to be higher in the asexual populations genome-wide estimations. However, when only homologous genome copies were compared (accounted for by allopolyploid model using Tetmer), $\theta_w$ estimations were generally lower, revealing the third genome copy as a major source for genetic diversity due to the hybrid origin. Increased genetic diversity in asexuals could also be explained by a larger effective population size $N_e$ (Soulé, 1976). Under the neutral theory, genetic diversity is expected to increase with $N_e$ and this has been seen in taxa such as chordates, annelids, arthropods and molluscs among others (Buffalo, 2021; Leffler et al., 2012). Larger, more stable populations are then expected to maintain greater levels of neutral genetic diversity than populations with lower effective population sizes $N_e$ (Leffler et al., 2012).It is predicted that large population sizes in asexual taxa aid the avoidance of extinction since the effectiveness of natural selection is increased and the lack of recombination can be compensated in such lineages Ross, Hardy, Okusu, and Normark (2013). Consequently, studies have found larger populations for asexual oribatid mites than in sexual ones in temperate and boreal forests (Brandt et al., 2017).

Heterozygosity is heavily affected by effective population size. This could differ between asexual and sexual strains because of differences in actual population size, reproduction mode and the different impact of purifying selection. For example, an increased mutational load in asexuals could reduce their effective population size due to purifying selection, but this effect could be compensated by larger population sizes.

Asexual reproduction accompanied by high genetic diversity has shown to allow for rapid adaptive responses on parasitic nematodes of *Meloidogyne* species to their hosts (Castagnone-Sereno, 2006). Nucleotide diversity of natural *C. elegans* populations from Hawaii has been found to be lower than that found here for asexual and sexual populations, these Hawaiian isolates are known to harbor a higher genetic diversity than all other known *C. elegans* populations. However, the average genome-wide diversity found for these *C. elegans* populations ($\pi$ 0.00109) is very similar to what was obtained for JU765 (spop4 - 0.00129). Both *C. elegans* and our *Propanagrolaimus* populations are diploid and selfing hermaphrodites (Crombie et al., 2019). The variable proportion of nucleotide diversity found on different genomic regions, is consistent with the assumed pattern across the genomic landscape. Some regions have very low diversity and could correspond to coding regions, whereas introns are expected to be more diverse (Tatarinova et al., 2016). Follow-up studies, including the annotation of reference genomes, will allow for more precise estimations of $\pi$ across the genomic landscape.

## 4.5 Genomic distinctiveness of *Panagrolaimus* strains

Split networks have been successfully applied to compare distantly related taxa (yeast, mammals, *Drosophila* and *C. briggsae*)(Huson & Bryant, 2006), as well as populations of hyperdiverse nematodes (Dey et al., 2013). Our split network analysis based on 375 orthologues in the asexual strains and 213 orthologues in the sexual species, appears to indicate that the former are as genetically distinct as the latter. Notably, the network for the sexual species is tree-like, while it shows more splits for the parthenogens. This is expected in a triploid system, where homeologs are not resolved (or phased in the genome assemblies) and thus recapitulate a pattern usually seen in recombining populations.

The classical biological species concept, defines species as groups of (potentially) interbreeding populations which are reproductively isolated from other similar groups (Mayr, 1999). Naturally the concept is restricted to organisms with sexual reproduction. Asexual organisms can undergo speciation and form distinct clusters due to adaptation to different niches or to physical separation (Birky, Adams, Gemmel, & Perry, 2010) (William Birky & Barraclough, 2009). Distinct species have been found for asexual organisms such as bdelloid rotifers, oribatid mites and oligochaete worms by using genomic data (Birky et al., 2010), i.e. applying a phylogenetic or phylogenomic species concept. Similar to these taxa the monophyletic asexual *Panagrolaimus* nematodes in our study might be seen as species under a phylogenomic concept of the term.

## 4.6 Studying genome evolution in parthenogens requires improved analysis frameworks

In our study it became obvious that studying "parthenogenomic" complexity is currently limited on two very different levels. Firstly, the availability of genomic resources is limited (phased reference genomes from tiny invertebrate organisms are not available for enough representatives of the Phyla). Secondly, population genetic theory as well as its implementation in analysis tools is not geared towards accounting for variation in ploidy and mode of reproduction in non-model organisms.

### 4.6.1 Limitations in sequencing technology still affect inferences of population genomic parameters

The accuracy of population genomic analyses depends on the quality of the underlying genomic data obtained through sequencing. Short-read data, such as the one used here, can provide ample information about a genome, but it is insufficient to distinguish repetitive regions of the genome and cannot be used to properly to study structural variants (Adewale, 2020). The maximum resolution that can be derived from short-read data ultimately depends on the quality of the reference genome that is used to map and compare the data to (Waldvogel et al., 2020). Polyploid genomes, as those in the asexual strains studied here, are technically challenging as they are likely to involve divergent evolutionary trajectories between homeologs, which carry different patterns of variation (Hörandl et al., 2020). To better infer measures such as $\pi$ phased reference genomes should be used and with the current advances in long-read sequencing methods these should be available in the future (The Darwin Tree of Life Project Consortium et al., 2022) even for tiny invertebrate taxa, as nematodes or rotifers.

Genome sequencing has seen another drastic advancement in the last years with long-read methods becoming available for single individuals with sufficient amounts of DNA (e.g. vertebrates), thus potentially allowing for higher resolution in population genomic assays

(De Coster, Weissensteiner, & Sedlazeck, 2021). In our study we still depend on short-read sequencing data of pools of individuals, as single individual long-read sequencing from tiny organisms remains challenging. Consequently, our study remains reference-based with limited resolution on aspects such as structural genomic variation (Adewale, 2020). At the same time, short-read (Illumina) sequencing still outperforms long-read methods in terms of the cost-coverage-ratio when aiming for deep sequencing studies, which is inevitable for the here conducted MAL experiment for the identification of *de novo* mutations.

### 4.6.2 Population genetic theory and software implementation are not well suited to study complex genomes

Asexual reproduction adds another level of complexity to our understanding of mutational evolution, selection, and adaptation. Many expectations for genome evolutionary processes, as e.g. the mode of reproduction violates the classical population assumption of random mating (Fisher, 1923) (Wright, 1931).

Even when phased reference genomes become available for tiny invertebrate taxa more appropriate software tools to disentangle different genotypes (e.g. autopolyploids or allopolyploids) are needed in a population genomic analysis framework. Tools that account for both ploidy and asexuality simultaneously in population genomic analysis are limited, making current estimations and analysis from data of organisms with these characteristics non precise despite adjustments to reduce bias (Jighly, Abbott, & Daetwyler, 2019).

In this study we aimed to account for potential bias by adjusting coverage ranges to ensure high enough representation of the respective genotypes in the read data, and thus allow for each position on each genome copy to be called equally likely. However, to study such genotypes on a single individual level in polyploid systems it will be necessary to develop more sensitive software tools.

### 4.6.3 Theoretical expectations about mutation rate evolution need to be adapted to diverse and complex systems

Mutations are the ground source of genetic variation, even if most non-synonymous mutations that occur are deleterious. While both mutation and recombination rates are variable along the genome, in obligate asexual taxa recombination can be completely absent, possibly leading to the accumulation of mildly deleterious mutations. We have found indications that natural populations of asexual *Panagrolaimus* show an elevated level of heterozygosity, potentially due to their third genome copy and the lack of allele segregation. The mutation rate in these organisms appears to be low, thus maybe delaying the effect of Muller's ratchet. The differentiation found within sets of natural populations, as well as the genomic differentiation between strains seems to show the potential for evolution in parthenogenetic animals. Asexual reproduction does not occur in the same way across the tree of life. In some taxa like asexual angiosperms, asexual reproduction occurs with meiosis still present and where gene conversion can occur. In other taxa, such as *Panagrolaimus* nematodes, asexual meiosis happens without recombination, whereas in other nematode species (Castagnone-Sereno & Danchin, 2014) and stick insects (Schwander, Henry, & Crespi, 2011), mitotic parthenogenesis without meiosis can take place. This diversity necessarily will be reflected in at least equally diverse mechanisms of genome evolutionary processes which ultimately cannot be resolved as long as we aim to apply one unifying theory.

## 5 Outlook

Due to the ability of parthenogentic taxa to save resources that would otherwise be allocated to finding a mating partner as well as the potential to found new populations by the dislocation of single individuals, taxa with this reproductive mode could be favored in extreme challenging environments (geographical parthenogenesis)(Tilquin & Kokko, 2016). This pattern has been observed in plants, flatworms (Lorch, Zeuss, Brandl, & Brändle, 2016) and ostracods (Symonová et al., 2018), and ambivalent data has been reported on salamanders (Greenwald, Denton, & Gibbs, 2016). Given our findings that genomic diversity is high in asexual populations, we would like to extend our research to further analyse a large set of populations isolated from distinct extreme environments to test geographical parthenogenesis in nematodes.

## 6 Acknowledgments

The authors gratefully acknowledge the help of Michael Kroiher and Einhard Schierenberg in conceiving the original MAL experiment. The authors are thankful to Tanja Schwander for comments on analyses methods, and to Caroline Blanc and Marie Delattre for a cytological analysis in two *Panagrolaimus* nematodes. This project was supported by, initially, a grant from the Volkswagen Foundation to P. Schiffer. The project was then supported through a DFG ENP Projekt [grant number 434028868] to P. Schiffer, and the DFG funded project B08 in the CRC1211 [grant number 268236062] to P. Schiffer and A-M Waldvogel, in which L. Villegas is employed. We furthermore thank the Regional Computing Center of the University of Cologne (RRZK) for providing computing time on the DFG-funded [Funding number INST 216/512/1FUGG] High Performance Computing (HPC) system CHEOPS as well as support.

## References

Adewale, B. A. (2020, November). Will long-read sequencing technologies replace short-read sequencing technologies in the next 10 years? *African Journal of Laboratory Medicine*, *9*(1). Retrieved 2022-08-10, from http://www.ajlmonline.org/index.php/AJLM/article/view/1340 doi: 10.4102/ajlm.v9i1.1340

Amat, I., van Alphen, J. J., Kacelnik, A., Desouhant, E., & Bernstein, C. (2017, September). Adaptations to different habitats in sexual and asexual populations of parasitoid wasps: a meta-analysis. *PeerJ*, *5*, e3699. doi: 10.7717/peerj.3699

Andrews. (2010, Jun). Fastqc. *https://qubeshub.org/resources/fastqc*.

Auwera, G. v. d., & O'Connor, B. D. (2020). *Genomics in the cloud: using Docker, GATK, and WDL in Terra* (First edition ed.). Sebastopol, CA: O'Reilly Media. (OCLC: on1148137471)

Bååth, R. (2014). Bayesian first aid: A package that implements bayesian alternatives to the classical *.test functions in r. In *User! 2014 - the international r user conference.*

Balloux, F., Lehmann, L., & de Meeûs, T. (2003, August). The population genetics of clonal and partially clonal diploids. *Genetics*, *164*(4), 1635–1644. doi: 10.1093/genetics/164.4.1635

Barrière, A., Yang, S.-P., Pekarek, E., Thomas, C. G., Haag, E. S., & Ruvinsky, I. (2009, March). Detecting heterozygosity in shotgun genome assemblies: Lessons from obligately outcrossing nematodes. *Genome Research*, *19*(3), 470–480. doi: 10.1101/gr.081851.108

Barton, N. H. (1994, December). The reduction in fixation probability caused by substitutions at linked loci. *Genetical Research*, *64*(3), 199–208. Retrieved 2023-02-20, from `https://www.cambridge.org/core/product/identifier/S0016672300032857/type/journal_article` doi: 10.1017/S0016672300032857

Becher, H., Brown, M. R., Powell, G., Metherell, C., Riddiford, N. J., & Twyford, A. D. (2020, November). Maintenance of Species Differences in Closely Related Tetraploid Parasitic Euphrasia (Orobanchaceae) on an Isolated Island. *Plant Communications*, *1*(6), 100105. Retrieved 2023-02-24, from `https://linkinghub.elsevier.com/retrieve/pii/S2590346220301334` doi: 10.1016/j.xplc.2020.100105

Bell, G. (1982). *The masterpiece of nature: the evolution and genetics of sexuality*. Berkeley: University of California Press.

Bengtsson, B. O. (2003, February). Genetic variation in organisms with sexual and asexual reproduction: Genetic variation in partially asexual organisms. *Journal of Evolutionary Biology*, *16*(2), 189–199. doi: 10.1046/j.1420-9101.2003.00523.x

Birky, C. W., Adams, J., Gemmel, M., & Perry, J. (2010, May). Using Population Genetic Theory and DNA Sequences for Species Detection and Identification in Asexual Organisms. *PLoS ONE*, *5*(5), e10609. doi: 10.1371/journal.pone.0010609

Blaxter, M. (2011, April). Nematodes: The Worm and Its Relatives. *PLoS Biology*, *9*(4), e1001050. Retrieved 2023-02-20, from `https://dx.plos.org/10.1371/journal.pbio.1001050` doi: 10.1371/journal.pbio.1001050

Brandt, A., Schaefer, I., Glanz, J., Schwander, T., Maraun, M., Scheu, S., & Bast, J. (2017, December). Effective purifying selection in ancient asexual oribatid mites. *Nature Communications*, *8*(1), 873. doi: 10.1038/s41467-017-01002-8

Brandt, A., Tran Van, P., Bluhm, C., Anselmetti, Y., Dumas, Z., Figuet, E., ... Bast, J. (2021, September). Haplotype divergence supports long-term asexuality in the oribatid mite *Oppiella nova*. *Proceedings of the National Academy of Sciences*, *118*(38), e2101485118. doi: 10.1073/pnas.2101485118

Buffalo, V. (2021, August). Quantifying the relationship between genetic diversity and population size suggests natural selection cannot explain Lewontin's Paradox. *eLife*, *10*, e67509. Retrieved 2023-04-28, from `https://elifesciences.org/articles/67509` doi: 10.7554/eLife.67509

Castagnone-Sereno, P. (2006, April). Genetic variability and adaptive evolution in parthenogenetic root-knot nematodes. *Heredity*, *96*(4), 282–289. doi: 10.1038/sj.hdy.6800794

Castagnone-Sereno, P., & Danchin, E. G. J. (2014, July). Parasitic success without sex - the nematode experience. *Journal of Evolutionary Biology*, *27*(7), 1323–1333. Retrieved 2022-08-19, from `https://onlinelibrary.wiley.com/doi/10.1111/jeb.12337` doi: 10.1111/jeb.12337

Charlesworth, D., & Willis, J. H. (2009, November). The genetics of inbreeding depression. *Nature Reviews Genetics*, *10*(11), 783–796. Retrieved 2023-03-08, from `https://www.nature.com/articles/nrg2664` doi: 10.1038/nrg2664

Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018, September). fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics*, *34*(17), i884–i890. Retrieved 2022-08-09, from `https://academic.oup.com/bioinformatics/article/34/17/i884/5093234` doi: 10.1093/bioinformatics/bty560

Crombie, T. A., Zdraljevic, S., Cook, D. E., Tanny, R. E., Brady, S. C., Wang, Y., ... Andersen, E. C. (2019, December). Deep sampling of Hawaiian Caenorhabditis elegans reveals high genetic diversity and admixture with global populations. *eLife*, *8*, e50465. doi: 10.7554/eLife.50465

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., ... Li,

H. (2021, January). Twelve years of SAMtools and BCFtools. *GigaScience*, *10*(2), giab008. doi: 10.1093/gigascience/giab008

[Dataset]Konrad, A. (2018, April). Experimental mutation accumulation in Caenorhabditis elegans under differing population size. *SRA - Bioproject PRJNA448413*. Retrieved from `https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA448413`

[Dataset]Schiffer, P. (2017, July). Evolution of parthenogenesis in Panagrolaimus nematodes. *SRA - Bioproject PRJNA374706*. Retrieved from `https://www.ncbi.nlm.nih.gov/bioproject/374706`

De Coster, W., Weissensteiner, M. H., & Sedlazeck, F. J. (2021, September). Towards population-scale long-read sequencing. *Nature Reviews Genetics*, *22*(9), 572–587. Retrieved 2023-03-08, from `https://www.nature.com/articles/s41576-021-00367-3` doi: 10.1038/s41576-021-00367-3

Denver, D. R., Dolan, P. C., Wilhelm, L. J., Sung, W., Lucas-Lledo, J. I., Howe, D. K., ... Baer, C. F. (2009, September). A genome-wide view of Caenorhabditis elegans base-substitution mutation processes. *Proceedings of the National Academy of Sciences*, *106*(38), 16310–16314. doi: 10.1073/pnas.0904895106

De Visser, J. A. G. M., & Rozen, D. E. (2005, July). Limits to adaptation in asexual populations. *Journal of Evolutionary Biology*, *18*(4), 779–788. doi: 10.1111/j.1420-9101.2005.00879.x

Dey, A., Chan, C. K. W., Thomas, C. G., & Cutter, A. D. (2013, July). Molecular hyperdiversity defines populations of the nematode Caenorhabditis brenneri. *Proceedings of the National Academy of Sciences*, *110*(27), 11056–11060. doi: 10.1073/pnas.1303057110

Engelstädter, J. (2008, November). Constraints on the evolution of asexual reproduction. *BioEssays*, *30*(11-12), 1138–1150. doi: 10.1002/bies.20833

Felsenstein, J. (1974, October). The evolutionary advantage of recombination. *Genetics*, *78*(2), 737–756. Retrieved 2023-04-05, from `https://academic.oup.com/genetics/article/78/2/737/5991150` doi: 10.1093/genetics/78.2.737

Fisher, R. A. (1923). Xxi.—on the dominance ratio. *Proceedings of the royal society of Edinburgh*, *42*, 321–341.

Gabriel, W., Lynch, M., & Burger, R. (1993, December). Muller's Ratchet and Mutational Meltdowns. *Evolution*, *47*(6), 1744. Retrieved 2022-08-09, from `https://www.jstor.org/stable/2410218?origin=crossref` doi: 10.2307/2410218

Gibson, A. K. (2019, December). Asexual parasites and their extraordinary host ranges. *Integrative and Comparative Biology*, *59*(6), 1463–1484. doi: 10.1093/icb/icz075

Greenwald, K. R., Denton, R. D., & Gibbs, H. L. (2016, November). Niche partitioning among sexual and unisexual *Ambystoma* salamanders. *Ecosphere*, *7*(11). Retrieved 2022-08-10, from `https://onlinelibrary.wiley.com/doi/10.1002/ecs2.1579` doi: 10.1002/ecs2.1579

Halligan, D. L., & Keightley, P. D. (2009, December). Spontaneous Mutation Accumulation Studies in Evolutionary Genetics. *Annual Review of Ecology, Evolution, and Systematics*, *40*(1), 151–172. Retrieved 2022-08-31, from `https://www.annualreviews.org/doi/10.1146/annurev.ecolsys.39.110707.173437` doi: 10.1146/annurev.ecolsys.39.110707.173437

Hartmann, M., Štefánek, M., Zdvořák, P., Heřman, P., Chrtek, J., & Mráz, P. (2017). The red queen hypothesis and geographical parthenogenesis in the alpine hawkweed hieracium alpinum (asteraceae). *Biological Journal of the Linnean Society*, *122*(4), 681–696. Retrieved 2023-02-22, from `http://academic.oup.com/biolinnean/article/122/4/681/4564920` doi: 10.1093/biolinnean/blx105

Hill, W. G., & Robertson, A. (1966, December). The effect of linkage on limits to artificial

selection. *Genetical Research*, *8*(3), 269–294.

Hirase, S., Yamasaki, Y. Y., Sekino, M., Nishisako, M., Ikeda, M., Hara, M., ... Kikuchi, K. (2021, October). Genomic Evidence for Speciation with Gene Flow in Broadcast Spawning Marine Invertebrates. *Molecular Biology and Evolution*, *38*(11), 4683–4699. doi: 10.1093/molbev/msab194

Huson, D. H. (1998, February). SplitsTree: analyzing and visualizing evolutionary data. *Bioinformatics*, *14*(1), 68–73. Retrieved 2022-08-24, from `https://academic.oup.com/bioinformatics/article-lookup/doi/10.1093/bioinformatics/14.1.68` doi: 10.1093/bioinformatics/14.1.68

Huson, D. H., & Bryant, D. (2006, February). Application of Phylogenetic Networks in Evolutionary Studies. *Molecular Biology and Evolution*, *23*(2), 254–267. doi: 10.1093/molbev/msj030

Hörandl, E., Bast, J., Brandt, A., Scheu, S., Bleidorn, C., Cordellier, M., ... Dunthorn, M. (2020). Genome Evolution of Asexual Organisms and the Paradox of Sex in Eukaryotes. In P. Pontarotti (Ed.), *Evolutionary Biology—A Transdisciplinary Approach* (pp. 133–167). Cham: Springer International Publishing. Retrieved 2022-09-07, from `http://link.springer.com/10.1007/978-3-030-57246-4_7` doi: 10.1007/978-3-030-57246-4_7

Institute, B. (2018). *Picard Toolkit*. Retrieved from `http://broadinstitute.github.io/picard/`

Jaron, K. S., Bast, J., Nowell, R. W., Ranallo-Benavidez, T. R., Robinson-Rechavi, M., & Schwander, T. (2021, March). Genomic Features of Parthenogenetic Animals. *Journal of Heredity*, *112*(1), 19–33. doi: 10.1093/jhered/esaa031

Jighly, A., Abbott, R. J., & Daetwyler, H. D. (2019, November). Editorial: Polyploid Population Genetics and Evolution—From Theory to Practice. *Frontiers in Ecology and Evolution*, *7*, 460. Retrieved 2022-08-10, from `https://www.frontiersin.org/article/10.3389/fevo.2019.00460/full` doi: 10.3389/fevo.2019.00460

Katoh, K. (2002, July). MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research*, *30*(14), 3059–3066. doi: 10.1093/nar/gkf436

Keightley, P. D., Ness, R. W., Halligan, D. L., & Haddrill, P. R. (2014, January). Estimation of the Spontaneous Mutation Rate per Nucleotide Site in a *Drosophila melanogaster* Full-Sib Family. *Genetics*, *196*(1), 313–320. Retrieved 2023-04-19, from `https://academic.oup.com/genetics/article/196/1/313/5935607` doi: 10.1534/genetics.113.158758

Kiontke, K., Gavin, N. P., Raynes, Y., Roehrig, C., Piano, F., & Fitch, D. H. A. (2004, June). *Caenorhabditis* phylogeny predicts convergence of hermaphroditism and extensive intron loss. *Proceedings of the National Academy of Sciences*, *101*(24), 9003–9008. Retrieved 2022-08-09, from `https://pnas.org/doi/full/10.1073/pnas.0403094101` doi: 10.1073/pnas.0403094101

Kofler, R., Orozco-terWengel, P., De Maio, N., Pandey, R. V., Nolte, V., Futschik, A., ... Schlötterer, C. (2011, January). PoPoolation: A Toolbox for Population Genetic Analysis of Next Generation Sequencing Data from Pooled Individuals. *PLoS ONE*, *6*(1), e15925. doi: 10.1371/journal.pone.0015925

Kofler, R., Pandey, R. V., & Schlotterer, C. (2011, December). PoPoolation2: identifying differentiation between populations using sequencing of pooled DNA samples (PoolSeq). *Bioinformatics*, *27*(24), 3435–3436. doi: 10.1093/bioinformatics/btr589

Konrad, A., Brady, M. J., Bergthorsson, U., & Katju, V. (2019, July). Mutational Landscape of Spontaneous Base Substitutions and Small Indels in Experimental *Caenorhabditis*

*elegans* Populations of Differing Size. *Genetics*, *212*(3), 837–854. Retrieved 2023-02-20, from `https://academic.oup.com/genetics/article/212/3/837/5931235` doi: 10.1534/genetics.119.302054

Leffler, E. M., Bullaughey, K., Matute, D. R., Meyer, W. K., Ségurel, L., Venkat, A., ... Przeworski, M. (2012, July). Revisiting an Old Riddle: What Determines Genetic Diversity Levels within Species? *PLoS Biology*. doi: 10.1371/journal.pbio.1001388

Lorch, S., Zeuss, D., Brandl, R., & Brändle, M. (2016, March). Chromosome numbers in three species groups of freshwater flatworms increase with increasing latitude. *Ecology and Evolution*, *6*(5), 1420–1429. Retrieved 2022-08-10, from `https://onlinelibrary.wiley.com/doi/10.1002/ece3.1969` doi: 10.1002/ece3.1969

Lynch, M. (2010, August). Evolution of the mutation rate. *Trends in Genetics*, *26*(8), 345–352. Retrieved 2022-08-09, from `https://linkinghub.elsevier.com/retrieve/pii/S0168952510001034` doi: 10.1016/j.tig.2010.05.003

Madeira, F., Park, Y. m., Lee, J., Buso, N., Gur, T., Madhusoodanan, N., ... Lopez, R. (2019, July). The EMBL-EBI search and sequence analysis tools APIs in 2019. *Nucleic Acids Research*, *47*(W1), W636–W641. Retrieved 2022-08-24, from `https://academic.oup.com/nar/article/47/W1/W636/5446251` doi: 10.1093/nar/gkz268

Mapleson, D., Garcia Accinelli, G., Kettleborough, G., Wright, J., & Clavijo, B. J. (2016, October). KAT: a K-mer analysis toolkit to quality control NGS datasets and genome assemblies. *Bioinformatics*, btw663. doi: 10.1093/bioinformatics/btw663

Mayr, E. (1999). *Systematics and the origin of species, from the viewpoint of a zoologist.* Harvard University Press.

McGill, L. M., Shannon, A. J., Pisani, D., Félix, M.-A., Ramløv, H., Dix, I., ... Burnell, A. M. (2015, March). Anhydrobiosis and Freezing-Tolerance: Adaptations That Facilitate the Establishment of Panagrolaimus Nematodes in Polar Habitats. *PLOS ONE*, *10*(3), e0116084. Retrieved 2023-02-20, from `https://dx.plos.org/10.1371/journal.pone.0116084` doi: 10.1371/journal.pone.0116084

Muller, H. J. (1932, March). Some Genetic Aspects of Sex. *The American Naturalist*, *66*(703), 118–138. Retrieved 2022-08-29, from `https://www.journals.uchicago.edu/doi/10.1086/280418` doi: 10.1086/280418

Oppold, A.-M., & Pfenninger, M. (2017, June). Direct estimation of the spontaneous mutation rate by short-term mutation accumulation lines in *Chironomus riparius*: MUTATION RATE OF *C. RIPARIUS*. *Evolution Letters*, *1*(2), 86–92. Retrieved 2023-03-08, from `https://academic.oup.com/evlett/article/1/2/86/6697412` doi: 10.1002/evl3.8

Otto, S. P., & Lenormand, T. (2002, April). Resolving the paradox of sex and recombination. *Nature Reviews Genetics*, *3*(4), 252–261. Retrieved 2022-08-12, from `http://www.nature.com/articles/nrg761` doi: 10.1038/nrg761

Prugnolle, F., & De Meeûs, T. (2008, September). The impact of clonality on parasite population genetic structure. *Parasite*, *15*(3), 455–457. doi: 10.1051/parasite/2008153p455

Ross, L., Hardy, N. B., Okusu, A., & Normark, B. B. (2013, January). LARGE POPULATION SIZE PREDICTS THE DISTRIBUTION OF ASEXUALITY IN SCALE INSECTS: DISTRIBUTION OF ASEXUALITY IN SCALE INSECTS. *Evolution*, *67*(1), 196–206. doi: 10.1111/j.1558-5646.2012.01784.x

Schiffer, P. H., Danchin, E. G., Burnell, A. M., Creevey, C. J., Wong, S., Dix, I., ... Blaxter, M. (2019, November). Signatures of the Evolution of Parthenogenesis and Cryptobiosis in the Genomes of Panagrolaimid Nematodes. *iScience*, *21*, 587–602. doi: 10.1016/j.isci.2019.10.039

Schwander, T., Henry, L., & Crespi, B. (2011, July). Molecular Evidence for Ancient Asex-

uality in Timema Stick Insects. *Current Biology*, *21*(13), 1129–1134. Retrieved 2022-08-24, from https://linkinghub.elsevier.com/retrieve/pii/S0960982211005537 doi: 10.1016/j.cub.2011.05.026

Schwarz, E. M. (2017, October). Evolution: A Parthenogenetic Nematode Shows How Animals Become Sexless. *Current Biology*, *27*(19), R1064–R1066. doi: 10.1016/j.cub .2017.08.040

Sedlazeck, F. J., Rescheneder, P., & von Haeseler, A. (2013, November). NextGen-Map: fast and accurate read mapping in highly polymorphic genomes. *Bioinformatics*, *29*(21), 2790–2791. Retrieved 2022-08-09, from https://academic.oup.com/bioinformatics/article-lookup/doi/10.1093/bioinformatics/btt468 doi: 10 .1093/bioinformatics/btt468

Shreve, S. M., Mockford, E. L., & Johnson, K. P. (2011, November). Elevated genetic diversity of mitochondrial genes in asexual populations of Bark Lice ('Psocoptera': Echmepteryx hageni): Genetic diversity in asexual Psocoptera. *Molecular Ecology*, *20*(21), 4433–4451. doi: 10.1111/j.1365-294X.2011.05298.x

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015, October). BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*, *31*(19), 3210–3212. doi: 10.1093/bioinformatics/btv351

Simón-Porcar, V. I., Silva, J. L., & Vallejo-Marín, M. (2021, April). Rapid local adaptation in both sexual and asexual invasive populations of monkeyflowers ( *Mimulus* spp.). *Annals of Botany*, *127*(5), 655–668. doi: 10.1093/aob/mcab004

Sloan, D. B., & Panjeti, V. G. (2009, October). Evolutionary feedbacks between reproductive mode and mutation rate exacerbate the paradox of sex. *Evolution*, *64*(4), 1129–1135. doi: 10.1111/j.1558-5646.2009.00869.x

Sloan, D. B., & Panjeti, V. G. (2010). Evolutionary feedbacks between reproductive mode and mutation rate exacerbate the paradox of sex. *Evolution*, *64*(4), 1129–1135. Retrieved 2023-02-20, from http://www.jstor.org/stable/27784076

Smith, J. M. (1978, November). Optimization Theory in Evolution. *Annual Review of Ecology and Systematics*, *9*(1), 31–56. Retrieved 2022-08-29, from https://www .annualreviews.org/doi/10.1146/annurev.es.09.110178.000335 doi: 10.1146/ annurev.es.09.110178.000335

Soulé, M. (1976). Allozyme variation: Its determinants in space and time. *Molecular Evolution*, 60–70.

Stoeckel, S., & Masson, J.-P. (2014, January). The Exact Distributions of FIS under Partial Asexuality in Small Finite Populations with Mutation. *PLoS ONE*, *9*(1), e85228. doi: 10.1371/journal.pone.0085228

Symonová, R., Vrbová, I., Lamatsch, D., Paar, J., Matzke-Karasz, R., Schmit, O., . . . Müller, S. (2018, March). Karyotype Variability and Inter-Population Genomic Differences in Freshwater Ostracods (Crustacea) Showing Geographical Parthenogenesis. *Genes*, *9*(3), 150. Retrieved 2022-08-10, from http://www.mdpi.com/2073-4425/9/3/150 doi: 10.3390/genes9030150

Tatarinova, T. V., Chekalin, E., Nikolsky, Y., Bruskin, S., Chebotarov, D., McNally, K. L., & Alexandrov, N. (2016, December). Nucleotide diversity analysis highlights functionally important genomic regions. *Scientific Reports*, *6*(1), 35730. doi: 10.1038/srep35730

Teotónio, H., Estes, S., Phillips, P. C., & Baer, C. F. (2017, June). Experimental Evolution with *Caenorhabditis* Nematodes. *Genetics*, *206*(2), 691–716. doi: 10.1534/genetics.115 .186288

The Darwin Tree of Life Project Consortium, Blaxter, M., Mieszkowska, N., Di Palma,

F., Holland, P., Durbin, R., ... Barnes, I. (2022, January). Sequence locally, think globally: The Darwin Tree of Life Project. *Proceedings of the National Academy of Sciences*, *119*(4), e2115642118. Retrieved 2022-08-19, from `https://pnas.org/doi/full/10.1073/pnas.2115642118` doi: 10.1073/pnas.2115642118

Thorvaldsdottir, H., Robinson, J. T., & Mesirov, J. P. (2013, March). Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. *Briefings in Bioinformatics*, *14*(2), 178–192. doi: 10.1093/bib/bbs017

Tilquin, A., & Kokko, H. (2016, October). What does the geography of parthenogenesis teach us about sex? *Philosophical Transactions of the Royal Society B: Biological Sciences*, *371*(1706), 20150538. Retrieved 2022-08-10, from `https://royalsocietypublishing.org/doi/10.1098/rstb.2015.0538` doi: 10.1098/rstb.2015.0538

Van Valen, L. (1973). A new evolutionary theory. *Evolutionary Theory*. Retrieved from `https://scholar.google.com/scholar_lookup?title=A\%20new\%20evolutionary\%20law&journal=Evol\%20Theory&volume=1&pages=1-30&publication_year=1973&author=Valen\%2CL`

Vasimuddin, M., Misra, S., Li, H., & Aluru, S. (2019, May). Efficient Architecture-Aware Acceleration of BWA-MEM for Multicore Systems. In *2019 IEEE International Parallel and Distributed Processing Symposium (IPDPS)* (pp. 314–324). Rio de Janeiro, Brazil: IEEE. Retrieved from `https://ieeexplore.ieee.org/document/8820962/` (Backup Publisher: 2019 IEEE International Parallel and Distributed Processing Symposium (IPDPS)) doi: 10.1109/IPDPS.2019.00041

Waldvogel, A., Feldmeyer, B., Rolshausen, G., Exposito‐Alonso, M., Rellstab, C., Kofler, R., ... Pfenninger, M. (2020, February). Evolutionary genomics can improve prediction of species' responses to climate change. *Evolution Letters*, *4*(1), 4–18. Retrieved 2022-08-24, from `https://onlinelibrary.wiley.com/doi/10.1002/evl3.154` doi: 10.1002/evl3.154

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis* (2nd ed. 2016 ed.). Cham: Springer International Publishing : Imprint: Springer.

William Birky, C., & Barraclough, T. G. (2009). Asexual Speciation. In I. Schön, K. Martens, & P. Dijk (Eds.), *Lost Sex* (pp. 201–216). Dordrecht: Springer Netherlands. Retrieved from `http://link.springer.com/10.1007/978-90-481-2770-2_10`

Winter, D. J., Wu, S. H., Howell, A. A., Azevedo, R. B., Zufall, R. A., & Cartwright, R. A. (2018). accuMUlate: a mutation caller designed for mutation accumulation experiments. *Bioinformatics (Oxford, England)*, *34*(15), 2659–2660. doi: 10.1093/bioinformatics/bty165

Wright, S. (1931, March). Evolution in Mendelian populations. *Genetics*, *16*(2), 97–159. Retrieved 2022-08-10, from `https://academic.oup.com/genetics/article/16/2/97/6045152` doi: 10.1093/genetics/16.2.97

Zhang, J., Kobert, K., Flouri, T., & Stamatakis, A. (2014, March). PEAR: a fast and accurate Illumina Paired-End reAd mergeR. *Bioinformatics*, *30*(5), 614–620. doi: 10.1093/bioinformatics/btt593

Zhou, X., Xie, Y., Zhang, Z.-h., Wang, C.-d., Sun, Y., Gu, X.-b., ... Yang, G.-y. (2013, December). Analysis of the genetic diversity of the nematode parasite Baylisascaris schroederi from wild giant pandas in different mountain ranges in China. *Parasites & Vectors*, *6*(1), 233. doi: 10.1186/1756-3305-6-233

# 7 Data Accessibility and Benefit-Sharing

Genome assemblies and sequencing data are deposited under Bioproject PRJNA374706 and are available through Wormbase.

# 8 Tables and figures

Table 1: Strains used, geographic origin, and mode of reproduction are specified. All species/strains apart from *P. davidi* are not described, i.e. *Panagrolaimus sp.* or Propanagrolaimus sp. in the case of JU765

| Original name | Renaming | Geographical origin | Analysis | Reference for mapping |
|---|---|---|---|---|
| JU765_parental | srefpool | Guangxi, China | | JU765 |
| JU765_100 | sl1 | | | JU765 |
| JU765_47 | sl2 | | Hermaphroditic | JU765 |
| JU765_60 | sl3 | | MAL | JU765 |
| JU765_71 | sl4 | Lines established from JU765_parental | | JU765 |
| JU765_73 | sl5 | | | JU765 |
| JU765_88 | sl6 | | | JU765 |
| PS1159_parental | arefpool | Raleigh, USA | | PS1159 |
| PS1159_82 | al1 | | | PS1159 |
| PS1159_22 | al2 | | Parthenogenetic | PS1159 |
| PS1159_8 | al3 | Lines established from PS1159_parental | (asexual) | PS1159 |
| PS1159_84 | al4 | | MAL | PS1159 |
| PS1159_86 | al5 | | | PS1159 |
| PS1159_19 | al6 | | | PS1159 |
| JU1646 | spop1 | Santiago Island, Cape Verde | Sexual | P. sp. bornheim |
| P. sp. bornheim | spop2 | Bonn, Germany | natural | P. sp. bornheim |
| P. sp. brombeer | spop3 | Cologne, Germany | populations | P. sp. bornheim |
| JU765 | spop4 | Guangxi, China | | JU765 |
| JB051 | apop1 | Diourbel, Senegal | Parthenogenetic | PS1159 |
| PS1159 | apop2 | Raleigh, USA | (asexual) | PS1159 |
| PS1162 | apop3 | Beijing, China | natural | PS1159 |
| PS1579 | apop4 | Pasadena, USA | populations | PS1159 |
| PS1806 | apop5 | Huntington, USA | | PS1159 |
| P. davidi | apop6 | McMurdo station, Antactic | | PS1159 |
| DL137G2 | apop7 | Corvallis, USA | | PS1159 |

Figure 1: Experimental setup of mutation accumulation. A total of 100 asexual lines (PS1159) and 71 sexual lines (JU765) were established at the beginning of the experiment; the total of final lines was 15 for PS1159 and 30 for JU765.

Figure 2: Nucleotide diversity (A) and population differentiation of (B) asexual and (C) sexual mutation accumulation lines. On average, $\theta_w$ and nucleotide diversity $\pi$ are higher in asexual lines. A pattern of differentiation can only be seen among sexual lines. Note that the range of Fst valued differs between asexuals and sexuals.

Figure 3: Nucleotide diversity $\pi$ on natural populations of nematodes isolated from distant geographical areas. Nucleotide diversity is higher on natural asexual populations when compared to natural sexual populations

Figure 4: Population differentiation of (A) natural sexual populations and (B) natural asexual populations. Orthologue gene network of (C) sexual and (D) asexual populations of *Panagrolaimus* nematodes. Sexual populations show higher differentiation than asexual ones. The gene network shows that asexual nematodes analyzed are genetically distinct from each other.

Figure 5: (A) Workflow of this work and (B) summary of results for asexual and sexual nematode populations. Mutation rate ($\mu$) was found to be tendentially lower on asexuals, nucleotide diversity ($\pi$) and Watterson's $\theta$ ($\theta_w$) were found to be larger on asexual population. Population differentiation ($F_st$)) was found to be higher in sexual natural populations and effective population size $N_e$ was higher for the asexual populations