# Glucocorticoids negatively relate to body mass on the short-term in a free-ranging ungulate

Lucas D. Lalande[1], Emmanuelle Gilot-Fromont[1,2], Jeffrey Carbillet[1,3], François Débias[1], Jeanne Duhayer[1,4], Jean-Michel Gaillard[1], Jean-François Lemaître[1], Rupert Palme[5], Sylvia Pardonnet[1,6], Maryline Pellerin[7], Benjamin Rey[1], Pauline Vuarin[1]

[1] Université de Lyon, Université Lyon 1, CNRS, Laboratoire de Biométrie et Biologie Evolutive UMR 5558, F-69622 Villeurbanne, France

[2] Université de Lyon, VetAgro Sup, Marcy l'Etoile, France

[3] Current institution: Institute of Ecology and Earth Sciences, University of Tartu, Tartu, 51014, Estonia

[4] Current institution: Office Français de la Biodiversité, Direction de la Recherche et de l'Appui Scientifique, Service Départemental de la Haute-Garonne, Villeneuve-de-Rivière, 31800, France

[5] Unit of Physiology, Pathophysiology and Experimental Endocrinology, Department of Biomedical Sciences, University of Veterinary Medicine, Veterinärplatz 1, 1210 Vienna, Austria

[6] Current institution: Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France

[7] Office Français de la Biodiversité, Direction de la Recherche et de l'Appui Scientifique, Service Conservation et Gestion Durable des Espèces Exploités, Châteauvillain, 52210, France

## Abstract

Environmental fluctuations force animals to adjust glucocorticoids (GCs) secretion and release to current conditions. GCs are a widely used proxy of an individual stress level. While short-term elevation in GCs is arguably beneficial for fitness components, previous studies have documented that the relationship between long-term baseline GCs elevation and fitness components can vary according to ecological and individual factors and according to the life-history of the species studied. Using longitudinal data on roe deer (*Capreolus capreolus*) from two populations facing markedly different environmental contexts, we tested whether baseline GC levels negatively correlate with body mass – a trait positively associated with demographic individual performance – on the short- to long-term. In support, higher baseline GC concentrations were associated to lighter body mass, both measured during the same capture event, in juvenile and adults of both populations. We also found evidence for an opposite relationship between FGMs and body mass gain during the early- and late-growth period, with no population or sex differences. Overall, we showed that despite the marked environmental and demographic differences between populations and despite the between-sex differences in life history (*i.e.* reproductive tactics), the relationship between body mass and GCs is consistent across environmental contexts, but differ mostly according to the life history stage of an individual. This work opens promising perspectives to further explore the relationship between GC and fitness-related traits according to life history stages in free-ranging mammals across seasonal and environmental contexts. The timing and context-dependence of GC levels highlight the complexity of studying stress responses in the wild.

**Keywords:** stress response, body condition, Faecal Cortisol Metabolites, individual performance, ungulates

## Introduction

Throughout their life, individuals have to adjust their morphology, physiology and/or behaviour to cope with environmental variation. Part of this variation is predictable as individuals can rely on environmental cues (*e.g.* photoperiod or temperature at the daily or seasonal scales) to anticipate future conditions (Wingfield 2003, 2008). Whereas predictable variation is not considered as stressful *per se* (*i.e.* eliciting a stress response), an unpredictable perturbation can disrupt life cycles and trigger a stress response through the activation of the hypothalamic-pituitary-adrenal (HPA) axis and the release of glucocorticoids (GCs), departing them from the baseline level (Reeder and Kramer 2005). The primary role of these hormones (namely cortisol and corticosterone) is to maintain an organism's energy balance according to its requirements in its current environment (Wingfield 2013, Hau et al. 2016), meaning that GCs promote allostasis (*i.e.* achieving stability through change) by ensuring energy homeostasis despite predictable and unpredictable environmental and life history variation (McEwen and Wingfield 2003, 2010, Romero et al. 2009). On the short-term, an elevation of GC concentration in an individual facing a challenging situation is beneficial because it promotes processes enhancing survival through reallocation of stored energy from non-immediately essential functions (e.g. growth, reproduction) towards cardiovascular functions or locomotion and foraging activities (Sapolsky et al. 2000). After a temporary perturbation, the GC concentration often quickly returns to the baseline level (Reeder and Kramer 2005). However, baseline GC concentration may remain elevated on the long-term (*i.e.* days to weeks) as a result from exposure to repeated or long-lasting perturbations such as prolonged inclement weather or food shortage (McEwen and Wingfield 2010, Hau et al. 2016). Such elevated GC concentration in the long-term cause 'allostatic overload' (McEwen and Wingfield 2010) and can have adverse consequences on individual performance through disruption of the immune function (Dhabhar 2014), inhibition of growth, or decreased body

condition (Reeder and Kramer 2005), with an intensity depending on the species (Boonstra

2013). Although variations in GC concentration does not reflect only stress (MacDougall-

Shackleton et al. 2019), GC levels are a widely used proxy for evaluating an individual

physiological stress response (Palme 2019).

While many studies analysed the effect of an increase of GC beyond the baseline level (*i.e.*

acute stress-induced GC concentration) on individual performance, fewer have considered

how GCs at baseline concentration (*i.e.* homeostatic levels of hormones) affect the

performance of animals in the wild. Results from these rare studies remain equivocal

(Rogovin et al. 2003, 2008, Altmann et al. 2004, Ethan Pride 2005, Pauli and Buskirk 2007,

Cabezas et al. 2007, Pedersen and Greives 2008, Pinho et al. 2019), which could be explained

by the complex influence of environmental conditions (Henderson et al. 2017). When

resources are limited, increased energy demands can be detrimental by resulting in high levels

of GC and allostatic overload. Nevertheless, this detrimental effect may not be observed when

resources are abundant, as high energy input can contribute reducing allostatic load (McEwen

and Wingfield 2003, 2010, Romero et al. 2009). Food availability (*i.e.* energy availability) or

behavioural flexibility stimulating foraging activity can thus compensate for elevated GC

levels and associated higher energy requirements (Henderson et al. 2017). Animals inhabiting

suboptimal habitats (*e.g.* exposed to human disturbance, predation, intra- or interspecific

competition or pathogens) experience continuous stressful conditions and generally show a

higher GC concentration compared to conspecifics inhabiting optimal areas (Dantzer et al.

2013, 2014, Formenti et al. 2018, Boudreau et al. 2019, Dulude-de Broin et al. 2020, Carbillet

et al. 2020).

Additionally, there is evidence that the direction of the relationship between GCs and

performance can also depend on individual attributes, such as body condition, sex or

reproductive status and reproductive tactics (Tilbrook et al. 2000, Ricklefs and Wikelski 2002,

Wey et al. 2015, Blumstein et al. 2016, Vuarin et al. 2019). Under high energy demands

triggered by elevated GC concentration, high quality individuals are more likely to prevent

themselves from reaching allostatic overload and may perform better than poor quality ones.

In African striped mice (*Rhabdomys pumilio*), elevated baseline GC concentration was related

to the selective disappearance of light individuals (Vuarin et al. 2019). Since the relationship

between GCs and performance is complex and involves individual attributes and

environmental context, both sources of variation have to be investigated to disentangle their

respective roles (Crespi et al. 2013, Dantzer et al. 2014).

Focusing on body mass provides a relevant approach to investigate the detrimental effects of

prolonged exposure to a high GC concentration. Indeed, body mass is often positively linked

to fitness components (*e.g.* Gaillard et al., 2000; Ronget et al., 2018) and is a trait easily

measured in the field that constitutes a reliable proxy of body condition in many species,

especially income breeders (Hewison et al. 1996, Andersen et al. 1998). High levels of GCs

are known to have a crucial role on body mass by increasing protein and amino-acid

catabolism and mobilising lipids (promoting gluconeogenesis), consequently decreasing body

mass (Rabasa and Dickson 2016).

However, the relationship between baseline GC concentration and body mass is complex.

Indeed, within seasonal baseline levels, GCs actually result in increased feeding behaviours

potentially promoting fat storing and body mass gain, as opposed to the previously-mentioned

GC actions on protein and amino-acid catabolism and lipid mobilisation (Landys et al. 2006).

Case studies have shown that the relationship between baseline GCs and body mass or

condition can be positive, negative or null (*e.g.* George et al. 2014, Wey et al. 2015, Hennin et

al. 2016, Boudreau et al. 2019). For instance, low body mass and poor body condition were

associated with higher baseline cortisol concentrations in Eurasian badgers (*Meles meles*,

George et al., 2014). On the other hand, experimental increase in GC concentration through

predation risk manipulation resulted in no changes in body condition in snowshoe hares

(*Lepus americanus*, Boudreau et al., 2019).

In this study, we aimed to contribute to a better understanding of how high baseline GC

concentration relates to body mass in roe deer (*Capreolus capreolus*), an income breeder for

which body mass is a good proxy for body condition (Hewison et al. 1996, Andersen et al.

2000, Pettorelli et al. 2006). We took advantage of the detailed and intensive monitoring data

collected through capture-mark-recapture for more than a decade at the individual level in two

populations subject to markedly different ecological contexts (*i.e.* one inhabiting a good

quality habitat and one population being strongly food-limited; Pettorelli et al., 2006). To

avoid measuring the capture-induced GC response (Romero & Reed, 2005), longer-term,

indirect measures of GC concentration can be obtained from biological matrices other than

blood (Sheriff et al. 2011). Indeed, blood GC concentration is a point in time measure and

circulating GCs concentration typically increases drastically within 3 minutes following an

individual's capture (Sapolsky et al. 2000, Romero and Reed 2005), which makes it difficult

to obtain reliable measures of the HPA baseline activity in free-ranging animals (but see

Sheriff et al. 2011, Lavergne et al. 2021). In this context, levels of faecal glucocorticoid

metabolites (FGMs) are particularly relevant as they represent an integrative measure of GC

concentration several hours before the capture event (*i.e.* baseline stress; Palme, 2019). Thus,

we measured FGMs, as arguably representative of baseline adrenocortical activity over a few

hours, with a time delay of 12h on average in roe deer (ranging from 6 to 23 hours, Dehnhard

et al. 2001).

As chronic stress can have both immediate and long-term consequences on fitness

components (Monaghan and Haussmann 2015), we tested the immediate relationship between

concomitant body mass and FGMs (*i.e.* short-term relationship), the relationship between

FGMs in a given year and the change of body mass between two consecutive years (*i.e.*

medium-term relationship) and the relationship between FGM measured on juveniles (*i.e.*

individuals in their first year of life) and body mass during the prime-age and senescent life

stages (*i.e.* long-term relationship). We first analysed the relationship between FGMs and

body mass measured at the same capture event. We expected individuals with higher FGM

concentration to be lighter than individuals with low FGM concentration. As roe deer are

income breeders that do not store fat (Hewison et al. 1996, Andersen et al. 1998), we expected

the catabolic actions of GCs to directly impact body mass and condition and to overcome the

effects of increased feeding behaviours promoted by GCs. Then, since chronically elevated

FGMs are expected to be deleterious on the medium to long-term, we tested whether FGMs

measured at a given capture influenced body mass changes across lifespan. Thus, we first

assessed the relationship between FGMs measured in a given year and a given body mass

change between two consecutive years, separately for growing individuals and prime-aged

adults having reached their full mass (*i.e.* between 4 and 10 years old; Douhard et al., 2017).

We expected that higher FGM concentration should result in a smaller mass gain for growing

individuals, and in a decrease in body mass for prime-aged adults, between two consecutive

years, as chronic stress can be linked to inhibited growth or decreased body condition (Reeder

and Kramer 2005). Finally, we tested for the effect of the stress experienced during the first

year of life on the individual body mass relative to the population and sex at a given age, from

the second year of life onwards. We predicted that individuals with higher FGMs during the

first year of life should be lighter later in life, as late-life performance in roe deer is affected

by early-life environmental conditions (Gaillard et al., 2003). Lastly, in all analyses we

expected the negative relationship between FGMs and body mass to be steeper in the food-

limited population due to higher allostatic load (McEwen and Wingfield 2003, Henderson et

al. 2017), and we expected sexes to respond differently to higher FGMs levels due to the

different reproductive tactics of males and females (Ricklefs and Wikelski 2002).

## Materials and Method

**Study population and sample collection**

We studied roe deer in two populations inhabiting closed forests at Trois-Fontaines (TF – 1360 ha) in north-eastern France (48°43'N, 4°55'E) and at Chizé (CH – 2614 ha) in western France (46°05'N, 0°25'W). Roe deer are medium-sized ungulates, weighing around 25 kg and common in lowland woodlands throughout most of Europe. Roe deer display weak sexual selection, with adult males only 10% heavier than females and party size less than three females per buck. They are income breeders (Andersen et al. 2000), and to meet the markedly increased energy needs during the late gestation and early lactation periods, adult females rely on food resources rather than body reserves. Both sexes allocate high energy expenditure to reproduction. Females produce twins every year from 2 to 12 years old (Andersen et al. 1998), and males allocate heavily in territory defence for 5-6 months during the rut period (Johansson 1996). The TF population is characterised by a homogeneous habitat of good quality on a broad spatial scale, with high forest productivity due to rich soils. On the other hand, the CH population inhabits a less suitable heterogeneous habitat comprising three types of varying quality (Pettorelli et al. 2001), with low forest productivity due to poor soils and frequent summer droughts (Pettorelli et al. 2006). In both study sites, large carnivores are absent, but hunting occurs occasionally to control population growth. Since 1975 in TF and 1977 in CH, a Capture-Mark-Recapture (CMR) program has taken place every winter, and CMR analyses show that both populations were quite below carrying capacity during the present study period (2010-2021 in TF, 2013-2021 in CH, unpublished data). Captures take place during 10-12 days each year, which are spread across December (at TF) or January (at CH) and March (Gaillard et al. 1993). Given that captures took place each year over 3 months, we might expect endocrine activity to vary throughout the capture season according to environmental conditions (*i.e.* populations) and sex (Dantzer et al. 2010, Sheriff et al.

2012). Therefore, we tested whether FGM levels varied as a function of the Julian date,

population and sex and found that FGMs increased throughout the season in a population-

specific manner (Figure S1). Thus, all FGM measurements were standardised for the median

Julian date of capture (9[th] of February) and corrected for the population effect. Successive

capture events within a capture season may also have consequences on FGM measurements

during the following captures. However, most capture days were more than 48 hours apart

while in roe deer, FGMs peak from 6 to 23 hours after an ATCH challenge and return to

baseline levels between 28 and 31 hours after treatment (Dehnhard et al. 2001). In some

cases, captures took place during two consecutive days, but were then conducted on opposite

areas of the forest to minimise disturbances. Likewise, morning and afternoon captures of a

given day took place in different areas within both forests. We thus considered that FGM

levels were not influenced by previous captures. Concerning the potential impact of the

capture event itself, since roe deer were captured by drive-netting and most animal

manipulations occurred between 1 and 4 hours after capture, we tested for a possible impact

of this delay on FGM levels. We did not detect any increase or decrease in FGMs (log-

transformed) along with the time between capture and sampling (linear regression accounting

for repeated measurements of a given individual: $0.006$ hour$^{-1} \pm 0.01$ SE, t = 0.44, p = 0.7),

even when the delay was > 6h (mean FGMs 6h onwards after capture compared to mean

FGMs measured within 6h following capture, accounting for repeated measurement of a given

individual, $-0.06 \pm 0.06$ SE, t = $-1.05$, p = 0.3) (Dehnhard et al. 2001). Individuals of known

age (*i.e.* captured within their first year of life) were weighed, and faecal matter has been

collected since 2010. During the capture period, diet is similar and mostly composed of

brambles (*Rubus* sp.) and ivy (*Hedera helix*) in the two deciduous forests (Tixier and Duncan

1996), so FGM measurements between populations should not be biased according to diet

composition. During this period, females can be gestating, and differences in female

reproductive status could result in FGM variability (Brunton et al. 2008, Dantzer et al. 2010).

Nevertheless, we did not account for females' reproductive status in our models, since i) we

have information about reproductive status (*i.e.* pregnant or not and number of foetuses) only

at CH where ultrasounds are performed on captured females, ii) most of them (*i.e.* 92 %) were

gestating at that time and iii) due to the delayed embryonic implantation (Aitken 1974),

gestation is at a very early stage during the capture season. Body mass measured on juveniles

(*i.e.* at about 8 months of age) depends on the date of capture. Juveniles gained on average 12

g/day and 24 g/day in CH and TF, respectively, during the capture period (Douhard et al.

2017). Therefore, we standardised juvenile body mass with a linear regression, using the

above-mentioned Julian date-body mass relationship. Thus, we computed the individual body

mass expected on the 9$^{th}$ of February, the median date of captures at CH and TF. In these two

populations, cohort quality is reliably measured by the cohort-specific average juvenile mass

corrected for the Julian date of capture (Gaillard et al. 1996). This proxy of environmental

quality from the birth of an individual (typically in May) to its capture as juvenile in winter

was obtained for all years, and will be referred to as 'cohort quality' hereafter.

**Faecal glucocorticoid metabolites (FGMs)**

Baseline adrenocortical activity was estimated through FGMs (Palme 2019). We collected

faecal matter rectally at capture from 2010 to 2021. Faeces were frozen at -20 °C within 24

hours prior to 2013 in CH and 2017 in TF, and immediately frozen at -80 °C after collection

since then. The time between faecal sampling and freezing, as well as freezing temperature,

may impact FGM values, with longer delays and higher temperature likely to result in lower

FGM levels due to bacterial activity (Hadinger et al., 2015; Lexen et al., 2008). In CH, the

first FGM values available in our dataset are post-2013, so all samples were immediately

frozen at -80 °C. However, in TF, FGM values are effectively higher when samples were

immediately frozen (0.43 ± 0.06 SE, t = 7.62, p < 0.0001). This is accounted for in our models

by adding the year of capture as a random effect. Extraction of FGM followed a methanol-

based procedure and the analysis was performed using a group-specific 11-

oxoaetiocholanolone enzyme immunoassay (EIA), a method previously described in details

(Möstl et al. 2002) and validated for roe deer (Zbyryt et al. 2018). In brief, 0.5 g (± 0.005) of

faeces were vortexed in 5 mL of 80% methanol before being centrifuged for 15 minutes at

2500 $g$ (Palme et al. 2013). The amount of FGM was determined in an aliquot of the

supernatant diluted 10 times in assay buffer. Measurements were done in duplicate with intra-

and inter-assay coefficients lower than 10% and 15%, respectively. FGMs are expressed as

nanograms per grams of wet faeces (ng/g). The data were then log-transformed for the

statistical analyses (henceforth called FGMs).

**Statistical analyses**

All analyses were performed using R version 4.0.5 (R Core Team 2021). Two raw FGM

measures were unusually high: 5192 and 6194 ng/g, for a female in CH aged 10, and a female

in TF aged 1, respectively (see Figure S2). Results exclude these extreme values (analyses

including them are reported in Table S1), but we specify whether adding them yielded

different conclusions or not.

*FGM repeatability*

Individual FGM repeatability was calculated for all individuals, and separately for each

population, using the 'rptR' package (Stoffel et al. 2017). Repeatability analysis included

individuals sampled only once to improve estimates of the within-individual variance (Martin

et al. 2011). Within-individual FGM repeatability was detectable but weak in both CH (r =

0.14, 95% CI = [0.03, 0.26]) and TF (r = 0.15 [0.01, 0.29]).

*Model structure*

We tested the hypothesis that body mass is related to FGM levels, and that this relationship

may depend on individual (*i.e.* sex) and environmental (*i.e.* population and condition at birth)

characteristics, using linear mixed effect models (LMMs) with a normal error distribution. All

continuous covariates were mean-centred so that the intercept is the relative individual body

mass for the mean cohort quality and/or mean FGMs. Visual inspection of model residuals

confirmed that they fulfilled the assumptions of Gaussian distribution and homoscedasticity.

*Short-term relationships between FGMs and body mass measured at the same capture event*

We first assessed the relationship between FGMs and body mass measured the same year (*i.e.*

at the same capture) to test the hypothesis that higher baseline FGMs result in short-term

adverse consequences on body mass. Individuals in their first year of life (*i.e* juveniles) were

analysed separately from individuals in their second year of life onwards, because juveniles

have to allocate to growth and are much more susceptible to any environmental harshness

than adults, making the first year of life is the critical period of roe deer population dynamics

(Hamel et al. 2009, Gaillard et al. 2013). In addition, metabolic rate is generally higher in

juveniles than in adults due to the costs of growth (Glazier 2005) and metabolic rates can alter

hormones metabolization and excretion, making it difficult to compare juvenile and adult

FGM concentrations (Goymann 2012). Since juveniles are not yet reproductively active, they

also likely display different endocrine profiles than adults (Dantzer et al. 2010). Finally, adult

roe deer habituate faster than juveniles to stress, which can in turn affect the relationship

between FGM and fitness-related trait (Bonnot et al. 2018). Body mass varies with age, and at

a given age, varies between sexes and according to environmental conditions at birth (*i.e.*

cohort effects, Hamel et al. 2016). We thus tested the relationship between FGMs measured in

a given year and individual body mass relative to the average body mass of all individuals of

the same age, population and sex, considering only ages for which we had data from at least 3

individuals. Therefore, the average body mass was calculated up to 10 and 15 years old in

males and females at CH, respectively, and up to 12 and 13 years old in males and females at TF, respectively. For juveniles, we analysed the relationship between FGMs and body mass measured the same year on 368 juveniles (78 females and 96 males in CH, 91 females and 103 males in TF). The response variable was the mass corrected for the date of capture (see above) and fixed explanatory variables included FGMs (corrected for the date of capture and the population), cohort quality, population, sex and all *4*-way, *3*-way and *2*-way interactions. Cohort quality was expressed as a relative cohort quality (*i.e.* the difference between mean cohort quality in each population and an individual cohort quality), so there is no redundancy with the population variable. The year of birth of the individuals (*i.e.* cohort) was included as a random effect. For individuals aged 2 years or older, the dataset included 655 observations on 377 individuals: 104 females (218 observations) and 83 males (136 obs.) in CH, and 106 females (164 obs.) and 84 males (137 obs.) in TF. The response variable used to test for the FGMs-mass relationship was the individual body mass minus the average mass of roe deer of the same population, same sex and same age, regardless of their year of capture (hereafter 'relative body mass). Fixed effects included continuous variation in FGMs, cohort quality (expressed as a relative cohort quality as described above), population, sex and all *4*-way, *3*-way and *2*-way interactions. Random effects of the cohort and of the year of capture were also included. Random effect of the individual identity (ID) was also included to account for repeated measurements on the same individuals.

*Medium-term relationships between FGMs and body mass change between two consecutive years*

Baseline FGMs can reflect medium-term consequences of GCs on body mass. The relationship between FGMs and the change of body mass between two consecutive years was analysed separately for early-growing individuals (*i.e.* between age 1 and 2), late-growing individuals (*i.e.* between the second year of life and adulthood at 4 years old, Hewison et al.

2011) and for prime-aged adults which had reached their full body mass (*i.e.* from 4 to 10

years old, Hewison et al., 2011). The response variable was the change in relative body mass.

Briefly, the change in relative mass expresses the change in body mass of a given individual

between two ages in relation to the change in body mass of all individuals of the same sex and

population between the same ages. The relative mass was calculated as the difference between

the mass of an individual at age $x$, and the mean mass of all individuals of age $x$ of the same

sex and the same population. The change in relative body mass was calculated as the

difference between the relative mass at age $x + 1$ and the relative mass at age $x$. Fixed effects

included relative cohort quality, population, sex, FGMs (at age $x$ or average of the FGM

values measured at age $x$ and age $x + 1$) and all *4*-way, *3*-way and *2*-way interactions, as well

as body mass at age $x$ (*i.e.* initial body mass) and its interaction with FGMs. Random effect of

the cohort was included, and for late-growing and prime-aged individuals, random effects of

the year of capture and individual ID were also included. For growing individuals, the dataset

included 99 individuals with FGMs measured as juveniles, and 67 individuals when including

FGMs measured during the first and second years of life to calculate the mean value. For both

late-growing individuals and adults, we kept a single observation per individual since the

inclusion of individual ID as a random effect created singularities. For individuals with

several observations, we kept the one for which the individual's age was the closest to the

mean age of all individuals with a unique observation according to sex and population, among

observations with complete data. The final dataset comprised 85 individuals in their late-

growth period and 90 prime-aged adults with FGMs measured in year *t*, and 63 late-growing

individuals and 71 prime-aged adults for which we had measurements in years *t* and *t+1*.

*Long-term relationships between FGMs during early life and body mass later in life*

As elevated baseline FGMs are expected to have negative effects on body mass on the long-

term, and as environmental conditions during the first year of life have carry-over effects on

performance later in life in roe deer (Gaillard et al. 2003), we tested whether FGMs during development are associated with adult body mass. Only individuals for which we measured FGMs as juvenile and body mass beyond the second year of life were analysed. We used 345 observations on 159 individuals for which we measured FGMs as juvenile and body mass as adult: 34 females (78 obs.) and 44 males (87 obs.) in CH, and 42 females (99 obs.) and 39 males (81 obs.) in TF. The response variable was the relative individual body mass as defined above. Fixed effects included FGMs measured when juvenile, relative cohort quality, population, sex and their *4*-way, *3*-way and *2*-way interactions. Random effects of the cohort and the year of capture were included, as well as individual ID to account for repeated measurements on the same individuals.

*Model selection*

Final models were selected based on AICc (Akaike Information Criterion corrected for small sample sizes). Among competitive models (assessed as the set of models within 2 ΔAICc, Burnham and Anderson 2002) we performed conditional model averaging based on the nesting selection rule (Richards 2008, Harrison et al. 2018). Briefly, model averaging was performed on the model with the lowest AICc value and simpler models nested within it, as well as models that were non-nested within the top-model as long as they did not include uninformative variables (*i.e.* an added parameter that did not substantially improve the model fit) (Richards 2008, Arnold 2010, Harrison et al. 2018). However, if the null model was present among the set of models within 2 ΔAICc, it was retained as the most plausible model. This model selection procedure allowed us to estimate the relative effects of the different variables on body mass while eliminating uninformative parameters to reach parsimony, and to alleviate the criticism that AIC tends to select over-complexified models (Link and Barker 2006). For each variable we estimated its effect size (β) with 85% confidence intervals (85%

CI, following the recommendations of Arnold 2010), and calculated marginal and conditional

$R^2$. We provide full model selection tables in supplementary information (Table S2).

**Results**

FGM concentrations ranged from 8 to 3428 ng/g with a median value of 689 ng/g. The

median FGMs was 674 ng/g (range: 34-3275 ng/g) in CH, and 715 ng/g (range: 8-3428 ng/g)

in TF.

**Short-term relationships between FGMs and body mass measured at the same capture**

**event**

*Juveniles*

The retained model highlighted a negative relationship between FGMs and body mass (both

corrected for the date of capture, $\beta_{FGM}$ = -0.23 [-0.4, -0.04], Figure 1(a), Table 1), as well as

the expected positive relationship between cohort quality and body mass ($\beta_{Cohort\ quality}$ = 1.08

[0.9, 1.3], Table 1). Estimates resulted from the conditional averaging of the model including

the additive effects of FGMs and cohort quality, and the model including cohort quality

(Table S2). Results were similar when including the high FGM value measured on a female at

TF (Table S1). Note that since the random effect 'cohort' created singularities, it was

removed and models therefore consisted of linear models.

*Adults*

We also found support for a short-term negative relationship between individual relative body

mass and FGMs in adults ($\beta_{FGM}$ = -0.32 [-0.4, -0.2], Figure 1(b), Table 1) as well as a positive

relationship between cohort quality and relative body mass ($\beta_{Cohort\ quality}$ = 0.26 [0.1, 0.5],

Table 1). Estimates resulted from the conditional averaging of the model including the

additive effects of FGMs and cohort quality, and the model including cohort quality (Table

S2). Results were similar when including the high FGM value measured on a female of CH

(Table S1). Since the random effect 'cohort' created singularities, the models only included

individual ID and year of capture as random effects.

394
**Figure 1.** Relationship between faecal glucocorticoid metabolites (log-transformed FGMs,
corrected for the date of capture and population) and individual relative body mass (individual
body mass (corrected for the date of capture for juveniles) minus the average body mass of
all individuals of the same age, population and sex) for (a) juveniles (N = 368 individuals) and
for (b) adults aged 2 years old onwards (N = 655 observations on 377 individuals). Points are
raw data and lines are predictions from the averaged selected models with 85% confidence
intervals.

**Medium-term relationships between FGMs and change in body mass between two**

**consecutive years**

*Early-growing individuals*

We did not find any evidence for a relationship between FGMs measured in juveniles and the

change in relative body mass for growing individuals. However, the change in relative body

mass was negatively related to the body mass of juveniles ($\beta_{mass}$ = -0.19 [-0.3, -0.1], Table 1).

Because body mass gain was lower for heavier juveniles, and because juvenile mass corrected

for the date of capture defines the cohort quality, we evidenced that the change in relative

body mass decreased with environmental quality at birth. This decrease was steeper in CH

than in TF ($\beta_{pop(CH)}$ = -1.49 [-2.9, -0.1], $\beta_{cohort}$ = -0.28 [-0.7, 0.1], $\beta_{pop(CH):cohort}$ = -0.73 [-1.4, -

0.1], Table 1). Estimates resulted from the conditional averaging of the model including the

additive effects of the population and initial body mass, and the model including the additive

effects and the two-way interaction between population and cohort quality (Table S2). Results

were similar when including the female of CH with the highest FGM value (Table S1). When

accounting for the mean FGM value between the first and second years of life, the change in

relative body mass was negatively related to the mean FGMs as well as to the juvenile mass

($\beta_{meanFGM}$ = -0.85 [-1.3, -0.4], $\beta_{mass}$ = -0.27 [-0.4, -0.1], Figure 2(a), Table 1). The change of

relative body mass was also lower in CH than TF ($\beta_{pop(CH)}$ = -1.57 [-2.3, -0.8], Figure 2(a),

Table 1). No model averaging was performed as this was the only model supported (Table

S2). However, when including the highest FGM value measured on a female at CH, we could

not detect any effect of FGMs, cohort quality, population, sex or initial body mass on the

change in relative body mass between the first and second years of life (Table S1). All models

included cohort as random effect.

*Late-growing individuals*

We found a positive relationship between the gain of body mass between two consecutive

years during late growth and FGMs measured year *t* ($\beta_{FGM}$ = 0.35 [0.1, 0.6], Figure 2(b),

Table 1) as well as a negative relationship between body mass gain and initial mass ($\beta_{mass}$ = -

0.15 [-0.2, -0.1], Table 1). Estimates resulted from the conditional averaging of the model

including the additive effects of FGMs and initial body mass, and the model including initial

body mass (Table S2). When accounting for the mean FGM value between FGMs measured

at year *t* and year *t+1*, we retained only a negative relationship between late growth and body

mass measured during year *t* ($\beta_{mass}$ = -0.17 [-0.2, -0.1], Table 1). This was the only retained

model so that no model averaging was performed (Table S2). Models resulted in singularities

when the random effect of the cohort was included so that only the random effect of the year

of capture was included in the models.

*Prime-aged adults*

In no cases (*i.e.* neither when accounting for FGM measured year $t$ or for the mean FGM

value between year $t$ and year $t+1$) was a relationship found between FGMs and change in

relative body mass (Figure 2(c), Table 1). Similarly, we found no evidence that mass

measured during year $t$, cohort quality, population or sex were related to changes in relative

body mass (Table 1, Table S2). Models resulted in singularities when the random effect of the

cohort was included. Thus, only the random effect of the year of capture was included in the

models.

447
**Figure 2.** Relationship between faecal glucocorticoid metabolites (log-transformed FGMs,
corrected for the date of capture and population) and change in relative body mass between
two consecutive years for (a) early-growing individuals (between the first and second years
of life), with FGMs either measured the first year of life (N = 99, pink points) or considered as
the mean FGMs between the first and second years (N = 67, green and blue points and lines
for TF and CH, respectively); for (b) late-growing individuals (between 2 and 4 years old) with
FGMs either measured during year $t$ (N = 85, pink points and line) or considered as the mean
FGMs between years $t$ and $t+1$ (N = 63, green points) and for (c) prime-aged adults which
had reached their full body mass (from 4 years old to 10 years old), with FGMs either
measured in a given year $t$ (N = 90, pink points) or considered as the mean FGM between
year $t$ and year $t+1$ (N = 71, green points). Points are raw data and lines are predictions from
the averaged selected models with 85% confidence intervals.

**Long-term relationships between FGMs during early-life and body mass later in life**

We found no evidence for a relationship between FGM measured in juveniles, cohort quality,

population or sex, and adult relative body mass (Table 1, Figure 3). Results were similar when

including the three observations of the adult female of CH with the highest FGM value

measured as juvenile (Table S1). We only included individual ID and year of capture as

random effects, because including 'cohort' created singularities.

**Figure 3.** Relationship between faecal glucocorticoid metabolites (log-transformed FGMs,
corrected for the date of capture and population) measured in juveniles and individual
relative body mass (*i.e.* individual body mass minus the average body mass of all individuals
of the same age, sex and population) (N = 345 observations on 160 individuals). Points are
raw data.

**Table 1.** Linear and linear mixed effect models selected for the short-, medium- and long-term
relationships between relative body mass and faecal glucocorticoid metabolites (FGMs). Models
accounted for sex (M: males), population (CH: Chizé) and for cohort quality (Qcoh). Models were
selected through model selection based on AICc and parameters estimated through model averaging.
85%CI: 85% confidence interval, V: variance, SD: standard-deviation.

| SHORT-TERM | | | | | |
|---|---|---|---|---|---|
| **Juveniles** | | | **Adults** | | |
| **Random effects** | **V** | **SD** | **Random effects** | **V** | **SD** |
| | | | Individual ID | 3.40 | 1.84 |
| | | | Year of capture | 0.25 | 0.50 |
| **Fixed effects** | **Estimate** | **85% CI** | **Fixed effects** | **Estimate** | **85% CI** |
| Intercept | -0.064 | [-0.22, 0.09] | Intercept | -0.010 | [-0.27, 0.25] |
| FGM | -0.231 | [-0,42, -0,04] | FGM | -0.318 | [-0.41, 0.46] |
| Qcoh | 1.083 | [0.91, 1.26] | Qcoh | 0.262 | [0.07, 0.46] |
| Marginal R² | 0.18 | | Marginal R² | 0.02 | |
| Conditional R² | | | Conditional R² | 0.81 | |

| MEDIUM-TERM | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| **FGM measured year *t*** | | | | | | | | |
| **Early-growth** | | | **Late-growth** | | | **Prime-aged** | | |
| **Random effects** | **V** | **SD** | **Random effects** | **V** | **SD** | **Random effects** | **V** | **SD** |
| Cohort | 0.35 | 0.59 | Year of capture | 0.41 | 0.64 | Year of capture | 0.24 | 0.49 |
| **Fixed effects** | **Estimate** | **85% CI** | **Fixed effects** | **Estimate** | **85% CI** | **Fixed effects** | **Estimate** | **85% CI** |
| Intercept | 0.132 | [-0.40, 0.66] | Intercept | -0.54 | [-0.88, -0.20] | Intercept | -0.193 | [0.50, 0.11] |
| Initial mass | -0.192 | [-0.31, -0.07] | FGM | 0.35 | [0.08, 0.61] | | | |
| Pop(CH) | -1.491 | [-2.86, -0.12] | Initial mass | -0.15 | [-0.22, -0.08] | | | |
| Qcoh | -0.277 | [-0.70, 0.15] | | | | | | |
| Pop(CH):Qcoh | -0.734 | [-1.35, -0.11] | | | | | | |
| Marginal R² | 0.15 | | Marginal R² | 0.12 | | Marginal R² | - | |
| Conditional R² | 0.26 | | Conditional R² | 0.35 | | Conditional R² | 0.15 | |

| **Mean FGM (FGM*t* and FGM*t+1*)** | | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| **Early-growth** | | | **Late-growth** | | | **Prime-aged** | | |
| **Random effects** | **V** | **SD** | **Random effects** | **V** | **SD** | **Random effects** | **V** | **SD** |
| Cohort | 0.24 | 0.49 | Year of capture | 0.32 | 0.56 | Year of capture | 0.18 | 0.42 |
| **Fixed effects** | **Estimate** | **85% CI** | **Fixed effects** | **Estimate** | **85% CI** | **Fixed effects** | **Estimate** | **85% CI** |
| Intercept | 0.322 | [-0.16, 0.81] | Intercept | -0.467 | [-0.81, -0.12] | Intercept | -0.285 | [-0.60, 0.03] |
| FGM | -0.845 | [-1.34, -0.35] | Initial mass | -0.167 | [-0.24, -0.09] | | | |
| Pop(CH) | -1.567 | [-2.34, -0.80] | | | | | | |
| Initial mass | -0.271 | [-0.41, -0.13] | | | | | | |
| Marginal R² | 0.18 | | Marginal R² | 0.13 | | Marginal R² | - | |
| Conditional R² | 0.26 | | Conditional R² | 0.32 | | Conditional R² | 0.10 | |

| LONG-TERM | | | | | | | | |
|---|---|---|---|---|---|---|---|---|

| Random effects | V | SD |
|---|---|---|
| Individual ID | 3.67 | 1.91 |
| Year of capture | 0.69 | 0.83 |
| **Fixed effects** | **Estimate** | **85% CI** |
| Intercept | -0.060 | [-0.48, 0.36] |
| Marginal R² | - | |
| Conditional R² | 0.83 | |

## Discussion

The relationship between GCs and fitness-related traits such as body mass is yet to be fully deciphered. Recent reviews indicate that studies in the wild have led to contrasting results by showing either negative, positive or no relationships between measures of stress and studied traits (Bonier et al. 2009, Crespi et al. 2013, Dantzer et al. 2014). A salient conclusion from these studies is that the relationship depends on an individual's ecological context and internal state. In the present study, the longitudinal monitoring of individual roe deer in two populations exposed to markedly different environmental contexts – one facing limiting resources and the other benefiting from a higher availability and quality of resources – allowed us to analyse the relationship between FGMs and body mass, while accounting for ecological factors and individual attributes such as sex and age. On the short-term, we found clear evidence for a negative relationship between baseline FGMs and body mass, an important driver of fitness in this species (Gaillard et al. 2000). Similarly, FGMs were negatively related to body mass gain between the first and second years of life. However, the relation was positive during late growth (*i.e.* individuals between 2 and 4 years old), and inexistant for prime-aged adults. Likewise, we detected no relationship between juvenile FGM levels and adult body mass.

Roe deer body mass correlated negatively with FGM levels on the short-term with no detectable population, age, or sex differences. This result suggests that the relationship between FGM and body mass is consistent across individuals and between environmental contexts in this species. Roe deer are income breeders, relying mostly on acquired resources

rather than on stored energy (Andersen et al. 2000). As GCs stimulate activity through the

mobilisation of energy stored (Sapolsky et al. 2000, Reeder and Kramer 2005), the

physiological costs of elevated GC levels are expected to directly impair body condition as

individuals do not have body reserves to could buffer elevated energy requirements

(Henderson et al. 2017). However, given marked between-population differences in

environmental context we would have expected individuals at TF to be able to mitigate the

physiological costs of high GC levels by reducing their allostatic load (McEwen and

Wingfield 2010, Henderson et al. 2017), compared to individuals at CH which were expected

to enter allostatic overload. Red deer (*Cervus elaphus*) showed different allostatic loads across

populations with different densities and environmental conditions, with individuals facing

high density or poor-quality habitat having higher FGM levels and lower body mass (Caslini

et al. 2016). Our results rather suggest that both roe deer populations are in a state of allostatic

overload. During the winter season, roe deer face more severe winters at TF than at CH,

which can in turn mitigate, during this period, the overall better resource quality in TF. This

could partly explain why no population differences were detected. It also suggests that

allostatic load can change across seasons, as shown in wild grey mouse lemurs (*Microcebus*

*murinus*, Hämäläinen et al. 2015), and that this variation could operate in a population-

specific way. It is therefore of great interest to assess the GC-fitness relationship across

different environmental contexts, including different years or climatic conditions, and through

various life history stages.

The change in body mass between two consecutive years was related to FGM levels

differently according to the individual life-stage. We found a negative relationship between

the change in body mass between the first and second years of life and FGMs. The fact that

only mean FGM levels (*i.e.* mean of the FGMs measured during the first and second years of

life) were related to body mass gain emphasizes that FGMs and body mass associate on the

short-term. Surprisingly, we found a positive relationship for individuals within their late-

growth phase. We rather expected a weaker negative relationship between FGM levels and

body mass gain, compared to early-growing individuals, or no relationship, as for prime-aged

adults. Such a positive relationship is intriguing and could suggest the existence of

compensatory growth (Metcalfe and Monaghan 2001), with individuals experiencing more

stress allocating more towards growth up to their full body mass, as shown in house sparrows

(*Passer domesticus*, Grace et al. 2017). However, for both early-growing and late-growing

individuals, we found that initial body mass (*i.e.* measured year *t*) was negatively related to

body mass gain. This suggests that compensatory growth does occur during the whole growth

period, which was accounted for in our models. We could not detect any relationship between

early-life FGM levels and adult body mass. As we found that higher FGMs at the juvenile

stage was related to lower juvenile body mass, this absence of carry-over effects on adult

body mass once again suggests the existence of compensatory growth (Metcalfe and

Monaghan 2001, Grace et al. 2017). Compensatory growth has adverse consequences on the

rate of body mass senescence in these populations (Douhard et al. 2017) and can ultimately

result in costs for survival or reproductive success. Whether GCs can modulate the process of

catching-up body mass remains to be investigated and would provide valuable insights into

the proximate mechanisms underpinning the associated costs.

Despite the numerous differences between our two study sites, we could not detect any

difference in the way individuals cope with stressors between populations, although we

expected different allostatic load in the two populations (McEwen and Wingfield 2010). Both

populations differ in terms of overall habitat quality, with CH being characterised by a low

forest productivity due to poor soils, mild winters, but frequent droughts during summer, and

TF by a high forest productivity due to rich soils and relatively wet summers (Pettorelli et al.

2006). Accordingly, life history parameters differ between populations. This is especially the

case for juvenile survival (Gaillard et al. 1993), body condition (Gilot-Fromont et al. 2012),

senescence patterns (haematological parameters: Jégo et al. 2014, body mass: Douhard et al.

2017, immune system: Cheynel et al. 2017), and the relation between FGMs and immune

traits (Carbillet et al. 2023). Thus, differences in the relationship between FGMs and body

mass was expected to occur between populations, with the strongest negative association in

CH, the poorer habitat in which individuals would be more at risk of allostatic overload

(McEwen and Wingfield 2010, Henderson et al. 2017). The absence of large predators and the

weak hunting pressure in both population might at least partly account for the absence of

support for a steeper relationship at CH. Studies of roe deer populations facing varying

conditions in terms of presence/absence of large predators and hunting pressure, in Poland,

showed that hunted populations without large predators had highest FGM levels with largest

variation (Zbyryt et al. 2018). In support, observed variation in FGM levels was high in both

populations (CV = 59.7 % and 55.3 % in CH and TF, respectively), which could mitigate the

relationship between body mass and FGMs and differences between populations.

Similarly, we could not detect any sex difference, despite the marked differences in the stress

response in males and females reported in mammals, due in part to the interaction between the

HPA and reproductive axes (Viau 2002, Wilson et al. 2005, and see Toufexis et al. 2014,

Novais et al. 2016 for reviews). This absence of sex-differences also suggests that during this

period of the year, female reproductive status does not influence the relationship between

FGMs and body mass. In reindeer (*Rangifer tarandus*), cortisol levels did not differ between

males, non-pregnant or pregnant females, but showed seasonal variation (Bubenik et al.

1998). Accordingly, red deer showed no sex-differences in their levels of FGMs but also

showed seasonal variation (Huber et al. 2003). Thus, it would be interesting to collect FGM

data of roe deer during different seasons as the stress response can vary accordingly (Bubenik

et al. 1998, Vera et al. 2011). It would also provide an opportunity to investigate whether the

different reproductive tactics of males and females can affect the relationship between GC and

fitness-related traits. Females of an iteroparous, long-lived species such as roe deer are

expected to trade reproduction for survival when environmental conditions are poor

(Hirshfield and Tinkle 1975). Thus, females are expected to be stress-responsive during their

reproductive period to be able to switch between alternative physiological states according to

whether they reproduce or not, due to their future opportunities of reproducing (Ricklefs and

Wikelski 2002). On the other hand, males allocate heavily to reproduction by defending

territories for half of the year (Johansson 1996). Territory defence results in elevated

physiological damages, whatever their reproductive success. Therefore, females could show a

larger variability in the way they respond to GC than males which are more constrained. This

provides promising perspectives to better understand the relationship between GC levels and

fitness-related traits according to sex-specific life history stages.

FGMs have been previously used as proxies to evaluate the homeostatic level or allostatic

load in wild populations, but the results should be interpreted with caution. FGMs do not fully

reflect an individual's stress response because GCs only represent one part of the complex

endocrine pathway involved in this response (MacDougall-Shackleton et al. 2019). Moreover,

although GCs are often referred to as "stress hormones", at baseline levels, their primary role

is to acquire, deposit and mobilise energy (Busch and Hayward 2009). Thus, unlike acute GC

concentrations, baseline GCs do not only provide information on how an individual responds

to stressors, but also on its physiology and activity (Reeder and Kramer 2005). It has been

reported that the acute GC response tends to be attenuated in individuals repeatedly exposed

to stressors (*e.g.* Ader et al. 1968, Kant et al. 1983, Vera et al. 2011). However, baseline GC

levels can also, but not always, be lower in individuals chronically exposed to environmental

stressors, making it to interpret FGM levels in the wild. Indeed, whether low baseline GCs

actually represent a non-stressed individual rather than an individual chronically exposed to

stressors is difficult to tell (Davis and Maney 2018).

In conclusion, we found that high baseline FGMs may have immediate adverse consequences

on body mass and that the relationship between body mass and FGM levels seems to depend

on an individual life-stage, rather than on the environmental context or the sex. Through this

work, we therefore emphasise the need to account for both environmental and individual

factors, including life history traits, to better capture the relationship between the stress

response and fitness-related traits. We also put emphasis on the idea to collect FGM across

different temporal and environmental contexts to evaluate how the GC-fitness relationship can

be modulated according to the seasonal and environmental conditions. To do so, it appears

critical to have access to longitudinal data. Measuring proxies of stress together with fitness-

related traits throughout the life of individuals is likely to be the only way to properly refine

our understanding of the implications of stress for fitness.

**Conflict of interest**

The authors declare having no conflict of interest.

**Ethics**

The research presented in this manuscript was done according to all institutional and/or

national guidelines. For both populations (Trois-Fontaines and Chizé), the protocol of capture

and blood sampling of roe deer under the authority of the Office Français de la Biodiversité

(OFB) was approved by the Director of Food, Agriculture and Forest (Prefectoral order 2009-

14 from Paris). All procedures were approved by the Ethical Committee of Lyon 1 University

(project DR2014-09, June 5, 2014).

**Funding**

This project was funded by the Université de Lyon, VetAgro Sup and the Office Français de

la Biodiversité (OFB, project CNV-REC-2019-08).

## Acknowledgements

This work was conducted as part of Ph.D. funded by the Université de Lyon and the Office

Français de la Biodiversité. We are grateful to all technicians, researchers and volunteers

participating in the collect of data on both sites (Trois-Fontaines and Chizé).

## References

Ader, R., Friedman, S. B., Grota, L. J. and Schaefer, A. 1968. Attenuation of the plasma
corticosterone response to handling and electric shock stimulation in the infant rat.
*Physiol. Behav.* **3**(2): 327–331. 10.1016/0031-9384(68)90109-1

Aitken, R. J. 1974. Delayed implantation in roe deer (*Capreolus capreolus*). *Reproduction*
**39**(1): 225–233. 10.1530/jrf.0390225

Altmann, J., Lynch, J. W., Nguyen, N., Alberts, S. C. and Gesquiere, L. R. 2004. Life-history
correlates of steroid concentrations in wild peripartum baboons. *Am. J. Primatol.*
**64**(1): 95–106. 10.1002/ajp.20064

Andersen, R., Duncan, P. and Linnell, J. D. C. 1998. *The European roe deer: The biology of
success*. Scandinavian University Press, Oslo.

Andersen, R., Gaillard, J.-M., Linnell, J. D. C. and Duncan, P. 2000. Factors affecting
maternal care in an income breeder, the European roe deer. *J. Anim. Ecol.* **69**(4): 672–
682. 10.1046/j.1365-2656.2000.00425.x

Arnold, T. W. 2010. Uninformative parameters and model selection using Akaike's
Information Criterion. *J. Wildl. Manag.* **74**(6): 1175–1178. 10.2193/2009-367

Blumstein, D. T., Keeley, K. N. and Smith, J. E. 2016. Fitness and hormonal correlates of
social and ecological stressors of female yellow-bellied marmots. *Anim. Behav.* **112**:
1–11. 10.1016/j.anbehav.2015.11.002

Bonier, F., Martin, P., Moore, I. and Wingfield, J. 2009. Do baseline glucocorticoids predict
fitness? *Trends Ecol. Evol.* **24**(11): 634–42. 10.1016/j.tree.2009.04.013

Bonnot, N. C., Bergvall, U. A., Jarnemo, A. and Kjellander, P. 2018. Who's afraid of the big
bad wolf? Variation in the stress response among personalities and populations in a
large wild herbivore. *Oecologia* **188**(1): 85–95. 10.1007/s00442-018-4174-7

Boonstra, R. 2013. Reality as the leading cause of stress: Rethinking the impact of chronic
stress in nature. *Funct. Ecol.* 27: 11–23. 10.1111/1365-2435.12008

Boudreau, M. R., Seguin, J. L., Boonstra, R., Palme, R., Boutin, S., Krebs, C. J. and Murray,
D. L. 2019. Experimental increase in predation risk causes a cascading stress response
in free-ranging snowshoe hares. *Oecologia* **191**(2): 311–323. 10.1007/s00442-019-
04500-2

Brunton, P. J., Russell, J. A. and Douglas, A. J. 2008. Adaptive responses of the maternal
hypothalamic-pituitary-adrenal axis during pregnancy and lactation. *J.*
*Neuroendocrinol.* **20**(6): 764–776. 10.1111/j.1365-2826.2008.01735.x

Bubenik, G. A., Schams, D., White, R. G., Rowell, J., Blake, J. and Bartos, L. 1998. Seasonal
levels of metabolic hormones and substrates in male and female reindeer (*Rangifer*
*tarandus*). *Comp. Biochem. Physiol. C Pharmacol. Toxicol. Endocrinol.* **120**(2): 307–
315. 10.1016/S0742-8413(98)10010-5

Burnham, K. P. and Anderson, D. R. 2002. *Model selection and multimodel inference: A*
*practical information-theoretic approach*. Springer, New York.

Busch, D. S. and Hayward, L. S. 2009. Stress in a conservation context: A discussion of
glucocorticoid actions and how levels change with conservation-relevant variables.
*Biol. Conserv.* **142**(12): 2844–2853. 10.1016/j.biocon.2009.08.013

Cabezas, S., Blas, J., Marchant, T. A. and Moreno, S. 2007. Physiological stress levels predict
survival probabilities in wild rabbits. *Horm. Behav.* **51**(3): 313–320.
10.1016/j.yhbeh.2006.11.004

Carbillet, J., Rey, B., Palme, R., Morellet, N., Bonnot, N., Chaval, Y., Cargnelutti, B.,
Hewison, A. J. M., Gilot-Fromont, E. and Verheyden, H. 2020. Under cover of the
night: Context-dependency of anthropogenic disturbance on stress levels of wild roe
deer *Capreolus capreolus*. *Conserv. Physiol.* **8**(1): coaa086. 10.1093/conphys/coaa086

Carbillet, J., Hollain, M., Rey, B., Palme, R., Pellerin, M., Regis, C., Geffré, A., Duhayer, J.,
Pardonnet, S., Debias, F., Merlet, J., Lemaître, J.-F., Verheyden, H. and Gilot-
Fromont, E. 2023. Age and spatio-temporal variations in food resources modulate
stress-immunity relationships in three populations of wild roe deer. *Gen. Comp.*
*Endocrinol.* **330**: 114141. 10.1016/j.ygcen.2022.114141

Caslini, C., Comin, A., Peric, T., Prandi, A., Pedrotti, L. and Mattiello, S. 2016. Use of hair
cortisol analysis for comparing population status in wild red deer (*Cervus elaphus*)
living in areas with different characteristics. *Eur. J. Wildl. Res.* **62**(6): 713–723.
10.1007/s10344-016-1049-2

Cheynel, L., Lemaître, J.-F., Gaillard, J.-M., Rey, B., Bourgoin, G., Ferté, H., Jégo, M.,
Débias, F., Pellerin, M., Jacob, L. and Gilot-Fromont, E. 2017. Immunosenescence
patterns differ between populations but not between sexes in a long-lived mammal.
*Sci. Rep.* **7**(1): 1–11. 10.1038/s41598-017-13686-5

Crespi, E. J., Williams, T. D., Jessop, T. S. and Delehanty, B. 2013. Life history and the
ecology of stress: how do glucocorticoid hormones influence life-history variation in
animals? *Funct. Ecol.* **27**(1): 93–106. 10.1111/1365-2435.12009

Dantzer, B., McAdam, A. G., Palme, R., Fletcher, Q. E., Boutin, S., Humphries, M. M. and
Boonstra, R. 2010. Fecal cortisol metabolite levels in free-ranging North American red

694 squirrels: Assay validation and the effects of reproductive condition. *Gen. Comp.*
695  *Endocrinol.* **167**(2): 279–286. 10.1016/j.ygcen.2010.03.024

696 Dantzer, B., Newman, A. E. M., Boonstra, R., Palme, R., Boutin, S., Humphries, M. M. and
697  McAdam, A. G. 2013. Density triggers maternal hormones that increase adaptive
698  offspring growth in a wild mammal. *Science* **340**(6137): 1215–1217.
699  10.1126/science.1235765

700 Dantzer, B., Fletcher, Q. E., Boonstra, R. and Sheriff, M. J. 2014. Measures of physiological
701  stress: A transparent or opaque window into the status, management and conservation
702  of species? *Conserv. Physiol.* **2**(1): cou023. 10.1093/conphys/cou023

703 Davis, A. K. and Maney, D. L. 2018. The use of glucocorticoid hormones or leucocyte
704  profiles to measure stress in vertebrates: What's the difference? *Methods Ecol. Evol.*
705  **9**(6): 1556–1568. 10.1111/2041-210X.13020

706 Dehnhard, M., Clauss, M., Lechner-Doll, M., Meyer, H. H. D. and Palme, R. 2001.
707  Noninvasive monitoring of adrenocortical activity in roe deer (*Capreolus capreolus*)
708  by measurement of fecal cortisol metabolites. *Gen. Comp. Endocrinol.* **123**(1): 111–
709  120. 10.1006/gcen.2001.7656

710 Dhabhar, F. S. 2014. Effects of stress on immune function: The good, the bad, and the
711  beautiful. *Immunol. Res.* **58**(2): 193–210. 10.1007/s12026-014-8517-0

712 Douhard, F., Gaillard, J.-M., Pellerin, M., Jacob, L. and Lemaître, J.-F. 2017. The cost of
713  growing large: Costs of post-weaning growth on body mass senescence in a wild
714  mammal. *Oikos* **126**(9): 1329–1338. 10.1111/oik.04421

715 Dulude-de Broin, F., Hamel, S., Mastromonaco, G. F. and Côté, S. D. 2020. Predation risk
716  and mountain goat reproduction: Evidence for stress-induced breeding suppression in
717  a wild ungulate. *Funct. Ecol.* **34**(5): 1003–1014. 10.1111/1365-2435.13514

718 Ethan Pride, R. 2005. High faecal glucocorticoid levels predict mortality in ring-tailed lemurs
719  (*Lemur catta*). *Biol. Lett.* **1**(1): 60–63. 10.1098/rsbl.2004.0245

720 Formenti, N., Viganó, R., Fraquelli, C., Trogu, T., Bonfanti, M., Lanfranchi, P., Palme, R. and
721  Ferrari, N. 2018. Increased hormonal stress response of Apennine chamois induced by
722  interspecific interactions and anthropogenic disturbance. *Eur. J. Wildl. Res.* **64**(6): 68.
723  10.1007/s10344-018-1228-4

724 Gaillard, J.-M., Delorme, D., Boutin, J.-M., Van Laere, G., Boisaubert, B. and Pradel, R.
725  1993. Roe deer survival patterns: A comparative analysis of contrasting populations. *J.*
726  *Anim. Ecol.* **62**(4): 778–791. 10.2307/5396

727 Gaillard, J.-M., Delorme, D., Boutin, J.-M., Van Laere, G. and Boisaubert, B. 1996. Body
728  mass of roe deer fawns during winter in 2 contrasting populations. *J. Wildl. Manag.*
729  **60**(1): 29–36. 10.2307/3802036

730 Gaillard, J.-M., Festa-Bianchet, M., Delorme, D. and Jorgenson, J. 2000. Body mass and
731  individual fitness in female ungulates: Bigger is not always better. *Proc. R. Soc. Lond.*
732  *B Biol. Sci.* **267**(1442): 471–477. 10.1098/rspb.2000.1024

Gaillard, J.-M., Loison, A., ToÏgo, C., Delorme, D. and Van Laere, G. 2003. Cohort effects
and deer population dynamics. *Écoscience* **10**(4): 412–420.
10.1080/11956860.2003.11682789

Gaillard, J.-M., Mark Hewison, A. J., Klein, F., Plard, F., Douhard, M., Davison, R. and
Bonenfant, C. 2013. How does climate change influence demographic processes of
widespread species? Lessons from the comparative analysis of contrasted populations
of roe deer. *Ecol. Lett.* **16**(s1): 48–57. 10.1111/ele.12059

George, S. C., Smith, T. E., Mac Cana, P. S. S., Coleman, R. and Montgomery, W. I. 2014.
Physiological stress in the Eurasian badger (*Meles meles*): Effects of host, disease and
environment. *Gen. Comp. Endocrinol.* **200**: 54–60. 10.1016/j.ygcen.2014.02.017

Gilot-Fromont, E., Jégo, M., Bonenfant, C., Gibert, P., Rannou, B., Klein, F. and Gaillard, J.-
M. 2012. Immune phenotype and body condition in roe deer: Individuals with high
body condition have different, not stronger immunity. *PLoS ONE* **7**(9): e45576.
10.1371/journal.pone.0045576

Glazier, D. S. 2005. Beyond the '3/4-power law': Variation in the intra- and interspecific
scaling of metabolic rate in animals. *Biol. Rev.* **80**(4): 611–662.
10.1017/S1464793105006834

Goymann, W. 2012. On the use of non-invasive hormone research in uncontrolled, natural
environments: the problem with sex, diet, metabolic rate and the individual. *Methods
Ecol. Evol.* **3**(4): 757–765. 10.1111/j.2041-210X.2012.00203.x

Grace, J. K., Froud, L., Meillère, A. and Angelier, F. 2017. House sparrows mitigate growth
effects of post-natal glucocorticoid exposure at the expense of longevity. *Gen. Comp.
Endocrinol.* **253**: 1–12. 10.1016/j.ygcen.2017.08.011

Hämäläinen, A., Heistermann, M. and Kraus, C. 2015. The stress of growing old: Sex- and
season-specific effects of age on allostatic load in wild grey mouse lemurs. *Oecologia*
**178**(4): 1063–1075. 10.1007/s00442-015-3297-3

Hamel, S., Gaillard, J.-M., Festa-Bianchet, M. and Côté, S. D. 2009. Individual quality, early-
life conditions, and reproductive success in contrasted populations of large herbivores.
*Ecology* **90**(7): 1981–1995. 10.1890/08-0596.1

Hamel, S., Gaillard, J.-M., Yoccoz, N. G., Albon, S., Côté, S. D., Craine, J. M., Festa-
Bianchet, M., Garel, M., Lee, P., Moss, C., Nussey, D. H., Pelletier, F., Stien, A. and
Tveraa, T. 2016. Cohort variation in individual body mass dissipates with age in large
herbivores. *Ecol. Monogr.* **86**(4): 517–543. 10.1002/ecm.1232

Harrison, X. A., Donaldson, L., Correa-Cano, M. E., Evans, J., Fisher, D. N., Goodwin, C. E.
D., Robinson, B. S., Hodgson, D. J. and Inger, R. 2018. A brief introduction to mixed
effects modelling and multi-model inference in ecology. *PeerJ* **6**: e4794.
10.7717/peerj.4794

Hau, M., Casagrande, S., Ouyang, J. and Baugh, A. 2016. Glucocorticoid-mediated
phenotypes in vertebrates: Multilevel variation and evolution. *Adv. Study Behav.* **48**:
41–115. 10.1016/bs.asb.2016.01.002

Henderson, L. J., Evans, N. P., Heidinger, B. J., Herborn, K. A. and Arnold, K. E. 2017. Do
glucocorticoids predict fitness? Linking environmental conditions, corticosterone and
reproductive success in the blue tit, *Cyanistes caeruleus*. *R. Soc. Open Sci.* **4**(10):
170875. 10.1098/rsos.170875

Hennin, H., Wells-Berlin, A. and Love, O. 2016. Baseline glucocorticoids are drivers of body
mass gain in a diving seabird. *Ecol. Evol.* **6**(6): 1702–1711. 10.1002/ece3.1999

Hewison, A. J. M., Angibault, J. M., Bideau, E., Vincent, J. P., Boutin, J. and Sempéré, A.
1996. Annual variation in body composition of roe deer (*Capreolus capreolus*) in
moderate environmental conditions. *Can. J. Zool.* **74**(2): 245–253. 10.1139/z96-031

Hewison, A. J. M., Gaillard, J.-M., Delorme, D., Van Laere, G., Amblard, T. and Klein, F.
2011. Reproductive constraints, not environmental conditions, shape the ontogeny of
sex-specific mass-size allometry in roe deer. *Oikos* **120**(8): 1217–1226.
10.1111/j.1600-0706.2011.19316.x

Hirshfield, M. F. and Tinkle, D. W. 1975. Natural selection and the evolution of reproductive
effort. *Proc. Natl. Acad. Sci.* **72**(6): 2227–2231. 10.1073/pnas.72.6.2227

Huber, S., Palme, R. and Arnold, W. 2003. Effects of season, sex, and sample collection on
concentrations of fecal cortisol metabolites in red deer (*Cervus elaphus*). *Gen. Comp.*
*Endocrinol.* **130**(1): 48–54. 10.1016/s0016-6480(02)00535-x

Jégo, M., Lemaître, J.-F., Bourgoin, G., Capron, G., Warnant, C., Klein, F., Gilot☐Fromont,
E. and Gaillard, J.-M. 2014. Haematological parameters do senesce in the wild:
Evidence from different populations of a long-lived mammal. *J. Evol. Biol.* **27**(12):
2745–2752. 10.1111/jeb.12535

Johansson, A. 1996. Territory establishment and antler cycle in male roe deer. *Ethology*
**102**(4): 549–559. 10.1111/j.1439-0310.1996.tb01147.x

Kant, G. J., Bunnell, B. N., Mougey, E. H., Pennington, L. L. and Meyerhoff, J. L. 1983.
Effects of repeated stress on pituitary cyclic AMP, and plasma prolactin,
corticosterone and growth hormone in male rats. *Pharmacol. Biochem. Behav.* **18**(6):
967–971. 10.1016/s0091-3057(83)80022-7

Landys, M. M., Ramenofsky, M. and Wingfield, J. C. 2006. Actions of glucocorticoids at a
seasonal baseline as compared to stress-related levels in the regulation of periodic life
processes. *Gen. Comp. Endocrinol.* **148**(2): 132–149. 10.1016/j.ygcen.2006.02.013

Lavergne, S. G., Krebs, C. J., Kenney, A. J., Boutin, S., Murray, D., Palme, R. and Boonstra,
R. 2021. The impact of variable predation risk on stress in snowshoe hares over the
cycle in North America's boreal forest: Adjusting to change. *Oecologia* **197**(1): 71–
88. 10.1007/s00442-021-05019-1

Link, W. A. and Barker, R. J. 2006. Model Weights and the Foundations of Multimodel
Inference. *Ecology* **87**(10): 2626–2635. 10.1890/0012-
9658(2006)87[2626:MWATFO]2.0.CO;2

MacDougall-Shackleton, S. A., Bonier, F., Romero, L. M. and Moore, I. T. 2019.
Glucocorticoids and "stress" are not synonymous. *Integr. Org. Biol.* **1**(1): obz017.
10.1093/iob/obz017

Martin, J. G. A., Nussey, D. H., Wilson, A. J. and Réale, D. 2011. Measuring individual
differences in reaction norms in field and experimental studies: A power analysis of
random regression models. *Methods Ecol. Evol.* **2**(4): 362–374. 10.1111/j.2041-
210X.2010.00084.x

McEwen, B. S. and Wingfield, J. C. 2003. The concept of allostasis in biology and
biomedicine. *Horm. Behav.* **43**(1): 2–15. 10.1016/S0018-506X(02)00024-7

McEwen, B. S. and Wingfield, J. C. 2010. What is in a name? Integrating homeostasis,
allostasis and stress. *Horm. Behav.* **57**(2): 105–111. 10.1016/j.yhbeh.2009.09.011

Metcalfe, N. B. and Monaghan, P. 2001. Compensation for a bad start: Grow now, pay later?
*Trends Ecol. Evol.* **16**(5): 254–260. 10.1016/s0169-5347(01)02124-3

Monaghan, P. and Haussmann, M. F. 2015. The positive and negative consequences of
stressors during early life. *Early Hum. Dev.* **91**(11): 643–647.
10.1016/j.earlhumdev.2015.08.008

Möstl, E., Maggs, J. L., Schrötter, G., Besenfelder, U. and Palme, R. 2002. Measurement of
cortisol metabolites in faeces of ruminants. *Vet. Res. Commun.* **26**(2): 127–139.
10.1023/A:1014095618125

Novais, A., Monteiro, S., Roque, S., Correia-Neves, M. and Sousa, N. 2016. How age, sex
and genotype shape the stress response. *Neurobiol. Stress* **6**: 44–56.
10.1016/j.ynstr.2016.11.004

Palme, R., Touma, C., Arias, N., Dominchin, M. F. and Lepschy, M. 2013. Steroid extraction:
Get the best out of faecal samples. *Wien. Tierärztl. Monatsschrift* **100**(9/10): 238–246.

Palme, R. 2019. Non-invasive measurement of glucocorticoids: Advances and problems.
*Physiol. Behav.* **199**: 229–243. 10.1016/j.physbeh.2018.11.021

Pauli, J. N. and Buskirk, S. W. 2007. Risk-disturbance overrides density dependence in a
hunted colonial rodent, the black-tailed prairie dog *Cynomys ludovicianus*. *J. Appl.*
*Ecol.* **44**(6): 1219–1230. 10.1111/j.1365-2664.2007.01337.x

Pedersen, A. B. and Greives, T. J. 2008. The interaction of parasites and resources cause
crashes in a wild mouse population. *J. Anim. Ecol.* **77**(2): 370–377. 10.1111/j.1365-
2656.2007.01321.x

Pettorelli, N., Gaillard, J.-M., Duncan, P., Ouellet, J.-P. and Van Laere, G. 2001. Population
density and small-scale variation in habitat quality affect phenotypic quality in roe
deer. *Oecologia* **128**(3): 400–405. 10.1007/s004420100682

Pettorelli, N., Gaillard, J.-M., Mysterud, A., Duncan, P., Stenseth, N. C., Delorme, D., Laere,
G. V., Toïgo, C. and Klein, F. 2006. Using a proxy of plant productivity (NDVI) to
find key periods for animal performance: The case of roe deer. *Oikos* **112**(3): 565–
572. https://doi.org/10.1111/j.0030-1299.2006.14447.x

Pinho, G. M., Ortiz-Ross, X., Reese, A. N. and Blumstein, D. T. 2019. Correlates of maternal
glucocorticoid levels in a socially flexible rodent. *Horm. Behav.* **116**: 104577.
10.1016/j.yhbeh.2019.104577

R Core Team 2021. R: A language and environment for statistical computing.https://www.R-
project.org

Rabasa, C. and Dickson, S. L. 2016. Impact of stress on metabolism and energy balance.
*Curr. Opin. Behav. Sci.* **9**: 71–77. 10.1016/j.cobeha.2016.01.011

Reeder, D. M. and Kramer, K. M. 2005. Stress in free-ranging mammals: Integrating
physiology, ecology, and natural history. *J. Mammal.* **86**(2): 225–235. 10.1644/BHE-
003.1

Richards, S. A. 2008. Dealing with overdispersed count data in applied ecology. *J. Appl. Ecol.*
**45**(1): 218–227. 10.1111/j.1365-2664.2007.01377.x

Ricklefs, R. E. and Wikelski, M. 2002. The physiology/life-history nexus. *Trends Ecol. Evol.*
**17**(10): 462–468. 10.1016/S0169-5347(02)02578-8

Rogovin, K. onstantin, Randall, J. A., Kolosova, I. and Moshkin, M. 2003. Social correlates
of stress in adult males of the great gerbil, *Rhombomys opimus*, in years of high and
low population densities. *Horm. Behav.* **43**(1): 132–139. 10.1016/s0018-
506x(02)00028-4

Rogovin, K. A., Randall, J. A., Kolosova, I. E. and Moshkin, M. P. 2008. Long-term
dynamics of fecal corticosterone in male great gerbils (*Rhombomys opimus Licht.*):
Effects of environment and social demography. *Physiol. Biochem. Zool.* **81**(5): 612–
626. 10.1086/588757

Romero, L. M., Dickens, M. J. and Cyr, N. E. 2009. The Reactive Scope Model - A new
model integrating homeostasis, allostasis, and stress. *Horm. Behav.* **55**(3): 375–389.
10.1016/j.yhbeh.2008.12.009

Romero, L. M. and Reed, J. M. 2005. Collecting baseline corticosterone samples in the field:
Is under 3 min good enough? *Comp. Biochem. Physiol. A. Mol. Integr. Physiol.*
**140**(1): 73–79. 10.1016/j.cbpb.2004.11.004

Ronget, V., Gaillard, J.-M., Coulson, T., Garratt, M., Gueyffier, F., Lega, J.-C. and Lemaître,
J.-F. 2018. Causes and consequences of variation in offspring body mass: Meta-
analyses in birds and mammals. *Biol. Rev.* **93**(1): 1–27. 10.1111/brv.12329

Sapolsky, R. M., Romero, L. M. and Munck, A. U. 2000. How do glucocorticoids influence
stress responses? Integrating permissive, suppressive, stimulatory, and preparative
actions. *Endocr. Rev.* **21**(1): 55–89. 10.1210/edrv.21.1.0389

Sheriff, M. J., Dantzer, B., Delehanty, B., Palme, R. and Boonstra, R. 2011. Measuring stress
in wildlife: Techniques for quantifying glucocorticoids. *Oecologia* **166**(4): 869–887.
10.1007/s00442-011-1943-y

Sheriff, M. J., Wheeler, H., Donker, S. A., Krebs, C. J., Palme, R., Hik, D. S. and Boonstra,
R. 2012. Mountain-top and valley-bottom experiences: the stress axis as an integrator

of environmental variability in arctic ground squirrel populations. *J. Zool.* **287**(1): 65–75. 10.1111/j.1469-7998.2011.00888.x

Stoffel, M. A., Nakagawa, S. and Schielzeth, H. 2017. rptR: repeatability estimation and variance decomposition by generalized linear mixed-effects models. *Methods Ecol. Evol.* **8**(11): 1639–1644. 10.1111/2041-210X.12797

Tilbrook, A. J., Turner, A. I. and Clarke, I. J. 2000. Effects of stress on reproduction in non-rodent mammals: The role of glucocorticoids and sex differences. *Rev. Reprod.* **5**(2): 105–113. 10.1530/ror.0.0050105

Tixier, H. and Duncan, P. 1996. Are European roe deer browsers? A review of variations in the composition of their diets. *Rev. Ecol. Terre Vie* **51**(1): 3–17. 10.3406/revec.1996.2189

Toufexis, D., Rivarola, M. A., Lara, H. and Viau, V. 2014. Stress and the reproductive axis. *J. Neuroendocrinol.* **26**(9): 573–586. 10.1111/jne.12179

Vera, F., Antenucci, C. D. and Zenuto, R. R. 2011. Cortisol and corticosterone exhibit different seasonal variation and responses to acute stress and captivity in tuco-tucos (*Ctenomys talarum*). *Gen. Comp. Endocrinol.* **170**(3): 550–557. 10.1016/j.ygcen.2010.11.012

Viau, V. 2002. Functional cross-talk between the hypothalamic-pituitary-gonadal and -adrenal axes. *J. Neuroendocrinol.* **14**(6): 506–513. 10.1046/j.1365-2826.2002.00798.x

Vuarin, P., Pillay, N. and Schradin, C. 2019. Elevated basal corticosterone levels increase disappearance risk of light but not heavy individuals in a long-term monitored rodent population. *Horm. Behav.* **113**: 95–102. 10.1016/j.yhbeh.2019.05.001

Wey, T. W., Lin, L., Patton, M. L. and Blumstein, D. T. 2015. Stress hormone metabolites predict overwinter survival in yellow-bellied marmots. *Acta Ethologica* **18**(2): 181–185. 10.1007/s10211-014-0204-6

Wilson, M. E., Legendre, A., Pazol, K., Fisher, J. and Chikazawa, K. 2005. Gonadal steroid modulation of the limbic-hypothalamic- pituitary-adrenal (LHPA) axis is influenced by social status in female rhesus monkeys. *Endocrine* **26**(2): 89–97. 10.1385/ENDO:26:2:089

Wingfield, J. C. 2003. Control of behavioural strategies for capricious environments. *Anim. Behav.* **66**(5): 807–816. 10.1006/anbe.2003.2298

Wingfield, J. C. 2008. Comparative endocrinology, environment and global change. *Gen. Comp. Endocrinol.* **157**(3): 207–216. 10.1016/j.ygcen.2008.04.017

Wingfield, J. C. 2013. Ecological processes and the ecology of stress: The impacts of abiotic environmental factors. *Funct. Ecol.* **27**(1): 37–44. 10.1111/1365-2435.12039

Zbyryt, A., Bubnicki, J. W., Kuijper, D. P. J., Dehnhard, M., Churski, M., Schmidt, K. and Editor, B. W. 2018. Do wild ungulates experience higher stress with humans than with large carnivores? *Behav. Ecol.* **29**(1): 19–30. 10.1093/beheco/arx142