O-antigen diversification masks identification of highly pathogenic STEC O104:H4-like strains

Christina Lang[1], Angelika Fruth[1], Ian W. Campbell[2], Claire Jenkins[3], Peyton Smith[4], Nancy Strockbine[4], François-Xavier Weill[5], Ulrich Nübel[6, 7, 8], Yonatan H. Grad[9], Matthew K. Waldor[2,9,10], and Antje Flieger[1#]

[1] Division of Enteropathogenic Bacteria and *Legionella*, National Reference Centre for *Salmonella* and other Enteric Bacterial Pathogens, Robert Koch-Institut, Wernigerode, Germany

[2] Department of Microbiology, Division of Infectious Diseases, Brigham & Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA

[3] Gastro and Food Safety (One Health) Division, Health Security Agency, London, UK

[4] Division of Foodborne, Waterborne and Environmental Diseases, National Center for Emerging and Zoonotic Infectious Diseases, Centers for Disease Control and Prevention, Atlanta, Georgia, USA

[5] Institut Pasteur, Université Paris Cité, Unité des bactéries pathogènes entériques, Paris, France

[6] Leibniz Institute DSMZ - German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany

[7] German Center for Infection Research (DZIF), Partner Site Braunschweig-Hannover, Germany

[8] Braunschweig Integrated Center of Systems Biology (BRICS), Technical University, Braunschweig, Germany

[9] Department of Immunology and Infectious Diseases, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA

27   [10] Howard Hughes Medical Institute, Boston, Massachusetts, USA

29   Running Head: **O-antigen diversification in STEC**

31   [#] Address correspondence to Antje Flieger, fliegera@rki.de

Abstract

Shiga toxin-producing *E. coli* (STEC) can give rise to a range of clinical outcomes from diarrhea to the life-threatening systemic condition, hemolytic uremic syndrome (HUS). Although STEC O157:H7 is the serotype most frequently associated with HUS, a major outbreak of HUS occurred in 2011 in Germany, and was caused by a rare serotype, STEC O104:H4. Prior to 2011 and since the outbreak, STEC O104:H4 strains have only rarely been associated with human infections. From 2012 to 2020 intensified STEC surveillance was performed in Germany where subtyping of ~8,000 clinical isolates by molecular methods including whole genome sequencing was carried out. A rare STEC serotype O181:H4 associated with HUS was identified and, like the STEC O104:H4 outbreak strain, this strain belongs to sequence type (ST) 678. Genomic and virulence comparisons revealed that the two strains are phylogenetically related and differ principally in the gene cluster encoding their respective lipopolysaccharide O-antigens but exhibit similar virulence phenotypes. In addition, five other serotypes belonging to ST678 from human clinical infection, such as OX13:H4, O127:H4, OgN-RKI9:H4, O131:H4, and O69:H4, were identified from diverse locations worldwide.

Importance

Our data suggest the high virulence ensemble of the STEC O104:H4 outbreak strain remains a global threat because genomically similar strains cause disease worldwide, but that horizontal acquisition of O-antigen gene clusters has diversified the O-antigens of strains belonging to ST678. Thus, identification of these highly pathogenic strains is masked by diverse and rare O-antigens, thereby confounding the interpretation of their potential risk.

Introduction

Shiga toxin-producing *E. coli* (STEC) are food-borne pathogens responsible for a range of clinical syndromes from diarrhea to the life-threatening systemic condition, hemolytic uremic syndrome (HUS), a triad of thrombotic microangiopathy, thrombocytopenia, and acute renal injury (1). Classical STEC include the pathovar enterohemorrhagic *E. coli* (EHEC), such as strains of serotype O157:H7, which in addition to the Shiga toxin gene, contain the locus of enterocyte effacement (LEE) pathogenicity island coding for a virulence-associated type III secretion system and effectors (1). Historically, the classification of STEC strains into different serotypes has proven invaluable for epidemiology and risk profiling (2). In *E. coli*, the serotype is determined by a combination of O- and H-antigen types (see below) whereas the O group solely denotes the O-antigen. Globally, O157:H7 strains are most frequently associated with HUS, but furthermore strains belonging to O groups O26, O103, O111, and O145, have also been regularly linked to HUS development (2, 3). In addition, a very rare serotype gave rise to a major HUS outbreak in early summer 2011, when an O104:H4 strain caused more than 3,000 cases of diarrhea and 800 cases of HUS including 54 fatalities, predominantly in Germany (4-6).

The O104:H4 outbreak strain encodes an exceptional set of virulence features (5-9). Like other HUS-associated strains, the strain produces Shiga toxin (Stx), specifically the Stx2a variant. But unlike most *E. coli* strains causing HUS, this strain belongs to enteroaggregative *E. coli* (EAEC), acquired a *stx2a*-encoding phage, and lacks LEE. EAEC strains characteristically harbor a plasmid (pAA) encoding aggregative adhesion fimbriae (AAF); in this case AAF of type I (AAF/I) (5, 7). AAF are responsible for bacterial autoaggregation, the stacked-brick adhesion phenotype to host cells, and contribute to the inflammatory response (10, 11). Furthermore, the outbreak strain encodes the virulence-linked serine-protease autotransporters (SPATE), SepA, SigA, and Pic and harbors an additional plasmid encoding

an extended spectrum beta-lactamase (ESBL) of the type CTX-M-15 (5, 11, 12). The

O104:H4 outbreak strain along with other O104:H4 strains all belong to multi-locus sequence

typing (MLST) sequence type (ST) ST678 and some possess *stx*, forming a distinct clade

among EAEC (5, 7, 11). Despite the extensive outbreak in May/June of 2011 and associated

wide distribution of the strain in affected regions, intensified molecular surveillance only

uncovered relatively few O104:H4 cases in Germany after the outbreak dissipated by July

2011.

In *E. coli*, serotypes are determined by the composition of the lipopolysaccharide (LPS) O-

antigen and the flagellar H-antigen, both important surface features of microorganisms that

shape pathogen host interactions (13, 14). LPS forms a major structural component of the

Gram-negative cell's outer membrane and its most distal part is the O-antigen. The O-antigen

is subject to strong selection pressure and is one of the most variable components of the

bacterial cell (13). Typically, in *E. coli* the O-antigen consists of chains of repeating

oligosaccharide subunits, usually composed of two to seven sugars often with additional

chemical modifications (15); currently, 182 O groups and 53 flagellar antigen types have been

described by phenotypic identification (14, 16). The genes encoding O-antigen biosynthesis

are organized in clusters that are flanked by colanic acid biosynthesis gene cluster (*wca*

genes) and histidine (*his*) biosynthesis operon (15). O-antigen biosynthesis gene clusters

typically have a lower GC content (often <40%) than that of the backbone of the *E. coli*

chromosome, which is ~50% GC content (13, 17, 18). These differences suggest that O-

antigen biosynthesis gene clusters are exchanged by lateral gene transfer and are under

diversifying selection, and therefore a hot spot of recombination (15, 19).

Although its wide distribution in the affected areas, the near disappearance of *E. coli*

O104:H4 in Germany after the large outbreak in 2011 was unanticipated. Here, we show that

the high virulence ensemble of the O104:H4 outbreak strain remains a threat, but that O-antigen gene exchange has cloaked the pathogen with several new O-antigens.

Results

The HUS-associated STEC O181:H4 strain 17-07187 shares a close phylogenetic relationship and similar virulence traits with the O104:H4 outbreak strain

After the large STEC O104:H4 outbreak in 2011, the German National Reference Centre for *Salmonella* and other Bacterial Enteric Pathogens intensified STEC surveillance and analyzed ~ 8,000 clinical isolates primarily from diarrhea and HUS patients from 2012 to 2020. This strain collection included a stool sample isolate (17-07187) from a 6-year-old girl who had bloody diarrhea and HUS in December 2017. She had not travelled outside her home in Northwest Germany before becoming ill. Serotyping and whole genome sequencing revealed that the strain belonged to an unusual serotype, O181:H4, that had not been previously associated with HUS. The strain had *stx2a* but lacked the LEE pathogenicity island (marker gene *eae*). Furthermore, the strain encoded characteristic EAEC markers including *aatA, aggR,* AAF/I genes and autotransporter protease genes *pic, sigA,* and *sepA* (Fig. 1A, Tab. S1).

MLST demonstrated that the 17-07187 isolate belonged to the same sequence type (ST678) as the O104:H4 outbreak strain (Table S2) (5, 7, 9). The EnteroBase core genome MLST (cgMLST) scheme (2,513 genes) (20) confirmed the close genomic relationship of this isolate with a panel of O104:H4 strains (Fig. 1B, Fig. S1A). There were only 30-34 allelic distances between this O181:H4 isolate and the O104:H4 outbreak strain (e.g. strains FWSEC0009, C227-11, or 11-02027), and O104:H4 clinical isolates from the Republic of Georgia in 2009 (2009EL-2071) and France in 2012 (Ec12-0465). In contrast, there was considerable allelic distance (AD) to STEC serotypes O181:H16 (ST6274), O181:H49 (ST173), O104:H21 (ST672), and O104:H7 (ST10075) isolated between 2012 and 2019 (AD >1500) (Fig. 1B).

Comparison of the virulence gene repertoire of the O181:H4 and the O104:H4 outbreak strain

also strongly suggested that they rely on very similar virulence mechanisms (Fig. 1A). Indeed,

both strains had comparable Stx-related cytotoxicity (Fig. 1C), exhibited a characteristic

enteroaggregative adherence pattern (Fig. 1D), and colonized intestinal tissues, particularly

the cecum and colon, similarly during *in vivo* infant rabbit infections (Fig. 1E). Thus, STEC

O181:H4 and O104:H4 isolates share marked genomic similarity and virulence-associated

genomic and phenotypic traits.

The genomes of the STEC O181:H4 strain 17-07187 and the O104:H4 outbreak strain

mainly differ in their O-antigen gene clusters and mobile genetic elements

The chromosomes (without plasmids) of the O181:H4 isolate 17-07187 and the O104:H4

outbreak strain FWSEC0009 were very similar (~99.7% nucleotide identity in the core

genome that is shared between these strains). The most striking difference between them were

their respective O-antigen gene clusters (OAGC) (Fig. 2A, 2B). Although these two clusters

were both situated in the same location in the chromosome, between *galF* and *hisI* (Fig. 2B),

their gene contents and organization were very different. Furthermore, their respective GC

contents, 36.8% for O181 and 37% for O104, differed from the chromosome GC content

(~50.7%), highlighting the likely role of lateral gene transfer in driving OAGC exchange.

Fourteen potential prophage regions are present in the O181:H4 isolate and 16 in the

O104:H4 outbreak strain (Fig. 2A, Table S3). Eleven of the fourteen prophages exhibited

substantial sequence identity (83-99.9%) with their O104:H4 counterparts and importantly,

the *stx2*-encoding prophages were nearly identical (~99.9% nucleotide identity) (Fig. 2C,

Table S4). Both *stx* phages are inserted into the tryptophan repressor binding protein gene

wrbA.

The genome of the O181:H4 isolate included three plasmids of ~81kb, ~76kb, and ~63kb (Table S2). The largest O181:H4 plasmid (pEc17-07187-1) was an incompatibility group I1 (IncI1) plasmid that showed only partial homology to the ESBL resistance plasmid of the O104:H4 outbreak strain (~75% nucleotide identity) but was very similar (98% nucleotide identity) to pHUSEC41-1 of STEC O104:H4 HUSEC41 from 2001 (92kb) (Fig. S1B) (3, 21). Both of these plasmids encode the *pilI-V* genes for thin pili. Unlike pHUSEC041-1, the O181:H4 pEc17-07187-1 did not contain antibiotic resistance genes (Fig. S1B, Tab. S5). O181:H4 plasmid (pEc17-07187-2/pAA) (76 kb) was very similar (99.8% nucleotide identity) to the O104:H4 outbreak strain pAA-EA11 and harbored virulence associated loci including *aggA/B/C/D*, which encode the AAF/I (Fig. 1A, Fig. S1C, Tab. S6) (7, 11). O181:H4 plasmid pEc17-07187-3 (63 kb) was not found in the O104:H4 outbreak strain; instead, it showed similarity to DHA plasmids of several enterobacteria coding for AmpC β-lactamase (22). However, unlike the DHA plasmids, resistance determinants were not present in the O181:H4 isolate (Fig. S1D, Tab. S7). Together these observations reveal the striking similarity of the chromosomes of the O181:H4 isolate and the O104:H4 outbreak strains and that their chief differences are confined to hot spots of recombination, i.e. their O-antigen gene clusters, and to mobile genetic elements, particularly their plasmids.

Additional recent global isolates of serotype O181:H4 and five other O groups belong to ST678

Next, we identified 158 genomes belonging to ST678 in EnteroBase (20), which contains ~202,200 *E. coli* genomes (as of April 11th, 2022). For a subset, serotype identification was not available, and for these cases, serotype was predicted based on the available genomic information (16). 123 of the ST678 strains were O104:H4, however, 18 additional O181:H4 genomes of ST678 were found. Furthermore, seven O127:H4, three O131:H4, and one each of O69:H4 and OX13:H4 were identified (Fig. 3A, Tab. S1). We categorized these as non-

O104:H4 ST678 strains. Additionally, based on a close phylogenetic relationship and single

difference in MLST alleles, we added three non-ST678 strains to the non-O104:H4 ST678

category: two O181:H4 strains (1472912 and 1472968) and one strain of the new and

provisionally assigned genoserotype OgN-RKI9:H4 (strain 608450) (Fig. 3A, Tab. S1).

cgMLST confirmed the close genetic relationship (AD ~50) between AAF/I gene positive

O104:H4 strains including the outbreak strain (FWSEC0009), and the 21 O181:H4 strains, the

OX13:H4 strain, and three of the seven O127:H4 strains (Fig. 3A, 3B). All of these strains

showed AAF/I which were also found in the 2011 outbreak strain (5, 7). Nine strains (four

O127:H4, three O131:H4, one each of O69:H4 and OgN-RKI9:H4) had higher allelic

distances up to 189. In contrast, these nine strains encoded AAF/III genes (Fig. 3A, 3B). The

25 AAF/I positive non-O104:H4 ST678 strains were all isolated in or after 2011 and the

majority was associated with diarrheal disease and a subset of six O181:H4 strains harbored

stx2a (Tab. S1, Fig. 3A, 3B, Fig. S2A). Interestingly, four of these six genomes (including 17-

07187 from Germany) shared a very similar *stx* phage with the O104:H4 outbreak strain, but

two of the six prophages were more distinct (Fig. 3A). AAF/I-positive strains shared regions

with a higher relatedness than those positive for AAF/III with the pAA plasmid of the 2011

outbreak strain (Fig. 3A).

The virulence gene profiles of the 34 non-O104 ST678 strains were generally similar to the

O104:H4 outbreak strain; however, there were a few differences. Specifically, *sepA* was

exclusively found in AAF/I-positive strains and EAST1 was present in all AAF/III-positive

strains and in only three of the AAF/I-positive isolates (Tab. S8).

The 34 non-O104:H4 ST678 strains were isolated in countries of Europe, Africa, and North

and South America. In addition, several were from individuals with a travel history that might

link these to East Asia (Fig. 3A, S2B and C). Together, these observations show that ST678

E. coli strains are found among seven different *E. coli* serotypes that have been linked to

diarrheal disease on several continents.

Phylogenomic analyses of ST678 *E. coli* suggest occurrence of multiple O-antigen gene

exchange events

The O-antigen gene clusters encoding the six O groups in the 34 non-O104:H4 ST678 strains

are found in the same chromosomal location as the 2011 O104:H4 outbreak strain but are

composed of largely disparate genes (Fig. 4A). These clusters also have a distinct GC content

(36.8-42.1%) from the backbone genome (50.5-50.7%), suggesting that they were acquired by

horizontal gene exchange. To explore the phylogenetic relationships among the 34 non-

O104:H4 ST678 strains and a set of O104:H4 strains, their shared single nucleotide

polymorphisms (SNPs) were analyzed using the O104:H4 outbreak strain FWSEC0009 as a

reference. A positive correlation (R=0.81; R^2=0,66) was found between isolation time and

genetic divergence (Fig. S2D) which supports that mutations have accumulated in a clock-like

fashion without notable outliers. A phylogenetic analysis shows that AAF/I positive strains

within ST678 are closely related to each other, clustering in clade I, whereas AAF/III positive

strains are more diverse and form more basal branches in a rooted maximum-likelihood

phylogenetic tree (Fig. 4B). This structure of the phylogeny suggests that clade I strains were

derived from an AAF/III-positive precursor. Within clade I, subclades Ia and Ib contain some

non-O104:H4 strains. Clade Ia is composed of O104:H4 non-outbreak strains isolated from

2015-2021 in the United Kingdom and in Kenya and the 2018 OX13:H4 strain that was

associated with travel to Ethiopia. Clade Ib contains all 21 O181:H4 strains and three

O127:H4 strains isolated from 2011 to 2021 from diverse continents. Since the majority of

ST678 strains are serotype O104:H4, including basal phylogenetic branches and most isolates

in clade I (Fig. 4B), it is parsimonious to assume that subclade Ib emerged from a O104:H4

predecessor, by replacing O104 antigen genes by O181 antigen genes. Similarly, our

phylogenetic analysis suggests that O127:H4 strains within clade Ib arose from an O181:H4

precursor. Together these observations indicate that O-antigen gene cluster exchange has

occurred repeatedly within ST678 strains.

Discussion

The *E. coli* O104:H4 outbreak in Germany in the early summer of 2011 was a public health

emergency; however, this serotype was rarely isolated as a cause of HUS after the epidemic

subsided. Nevertheless, our findings suggest that the unusual set of virulence factors that

characterized this Shiga toxin 2-producing EAEC strain remains a threat to human health.

Serotype conversion has cloaked this highly virulent genotype with several O-antigens,

including O181, O127, and OX13, which are present both in *stx* positive and *stx* negative

disease-linked isolates closely related to the O104:H4 outbreak strain. We found non-

O104:H4 ST678 strains from a variety of countries in Europe, Africa, and the Americas and

several of the cases were associated with travel to Africa and Asia, suggesting these

virulence-associated strains are globally distributed. Like the O104:H4 outbreak strain, the

AAF/I positive ST678 strains had similar chromosomes, similar pAA-linked virulence genes,

as well as virulence factors including the SPATEs (5, 7, 8, 12). The most salient difference of

the chromosomes of these strains with that of the outbreak strain were their respective O-

antigen gene clusters and other mobile genetic elements. Thus, horizontal exchange of these

clusters appears to have been a critical step in the evolution of these new pathogenic

serotypes, some of which were linked to HUS or bloody diarrhea.

The prime example uncovered here is the likely derivation of O181:H4 pathogens from an

O104:H4 precursor via O-antigen gene exchange. Among the 21 O181:H4 strains, six

harbored a *stx2a*-encoding prophage, including HUS-linked strain 17-07187. In four of these

strains, the *stx2a* prophage was very similar to the *stx2a* prophage of the 2011 outbreak strain

(Fig 3A), suggesting that the O-antigen exchange was a more recent event in their evolution

compared to the acquisition of the Shiga toxin-encoding prophage. The absence of *stx*

prophages in 15 of the O181:H4 isolates may be due to a lack of *stx* prophage acquisition or

have resulted from loss of their *stx* prophages, which is well documented in STEC in other

serotypes (23). Thus, in addition to O-antigen exchange, on-going horizontal transmission of

mobile genetic elements, such as *stx* phages, has contributed to diversification of

diarrheagenic ST678.

O-antigen gene clusters of Gram-negative bacteria are hot spots for diversifying selection and

recombination events (15, 19). Serotype conversion by lateral exchange of OAGCs has played

an important role in the evolution of enteric pathogens. For example, *Vibrio cholerae*

serogroup O139, which arose by exchange of O1 and O139 OAGCs, transiently replaced the

dominant O1 group as the cause of cholera in 1992-93 (13, 24). Our discovery of seven

distinct O-groups that share the flagellar H4 antigen and a highly similar virulence-linked

genetic makeup provides a compelling example of the role of O-antigen exchange in the

diversification of diarrheagenic *E. coli*. In addition, STEC O157:H7 is thought to have arisen

from an enteropathogenic *E. coli* (EPEC)-like O55:H7 strain which initially acquired *stx* via

phage transduction and subsequently acquired the O157 O group by exchange of the O55 with

the O157 OAGC (26). Also, an O182-O156 switch is thought to have occurred in STEC

strains persistently infecting cattle (18). Consequently, different OAGs may be found in

highly related genotypes (27, 28).

We can only speculate about the conditions driving OAGC exchange among ST678 strains.

Shiga toxin-producing O104:H4 EAEC strains, such as the 2011 outbreak strain, are

considered human-restricted pathogens and have not been isolated from animals, such as

cattle (29). However, interestingly a recent description highlighted the detection of an STEC

O104:H4 strain in pork (30). It is possible that the OAGC exchange occurred in a human host,

where the human intestinal microbiome may contain *E. coli* strains of O groups, such as O181

and O127, that could have donated their OAGC to an O104:H4 recipient by some means of

horizontal exchange. Also, epidemiological investigations suggest that fenugreek sprouts were

the food source that initiated the STEC O104:H4 epidemic; therefore plants colonized by

microbiota may be another possible site for OAGC exchange (4).

Conclusion

In conclusion, our study highlights how lateral exchange of O-antigen gene clusters can lead

to the rapid diversification of a globally important pathogen. Further, an important clinical

implication of these findings is that serotype surveillance cannot be used as a simple proxy for

strain virulence and needs to be complemented by virulence gene or genome analysis.

Surveillance to uncover how highly virulent strains may reemerge and spread in new O-

antigen outfits is warranted.

Material and Methods

Study strains

In the context of intensified STEC surveillance in Germany, clinical isolates were collected at

the National Reference Center for *Salmonella* and other Bacterial Enteric Pathogens and

analyzed for serotype, *stx* and subtypes, *eaeA*, *hlyA* and *aatA* as described (31). Strains were

grown on nutrient agar (Oxoid GmbH, Germany), Luria Bertani (LB) broth, or

enterohemolysin agar (Sifin GmbH, Germany). Genome sequencing was carried out on a

subset of strains and further genome sequence data of strains was gathered from EnteroBase

and NCBI (Table S1).

Whole genome sequencing (WGS)

Long read whole genome sequencing of O181:H4 strain 17-07187 was performed by GATC
Biotech (Konstanz, Germany) using a PacBio RS II sequencer (Pacific Biosciences, USA)
and short read genome sequencing of strains 12-04810, 14-03615, 14-01288, 16-00596, 16-
01499, 16-05332, 17-01774, 17-00416, 17-07187 and 19-02696 (Table S1) on an Illumina
MiSeq and HiSeq 1500 benchtop sequencer.

Bioinformatics analyses

De novo assembly of the PacBio sequence data (103-fold average coverage) was performed
by GATC utilizing HGAP3 (Pacific Biosciences, USA). Polishing of the assembled genome
and plasmids was performed with Illumina short reads by Pilon (version 1.22) (32). Quality
control and trimming of MiSeq raw reads with subsequent detection of serotype and virulence
genes was performed as described (16). Genomic comparisons were carried out using
MAUVE (version: 1.1.1) and MAFFT (version 1.3.7) as plugin in Geneious (version: 11.1.5;
Biomatters Ltd) (33, 34). Gaps within the MAFFT-alignment were excluded using the
Geneious "mask alignment" function. Ridom SeqSphere+ (version:7.2.0, Ridom GmbH,
Germany) was used to determine MLST Warwick sequence types and to create minimal
spanning trees based on 2513 allele targets (here genes) from the *E. coli* cgMLST EnteroBase
scheme with pairwise ignoring missing values from genome assemblies (20). Different
variants of the cgMLST genes among different genomes are represented as allelic distances
(AD). Phage prediction was carried out by analysing the genome sequences with PHASTER
(35). RAST was used for CDS annotation (36).

SNP-based alignment and maximum-likelihood based phylogenetic tree

Mapping of sequencing reads, generation of consensus sequences, and alignment calculation
was performed using the BatchMap pipeline with FWSEC0009 genome sequence as reference

(37). Gubbins (version 3.2.1) was used to identify loci containing elevated densities of base

substitutions (putative recombinations) within the alignment while concurrently constructing

an alignment and phylogeny (RAxML-tree) based on the putative point mutations (SNPs)

outside of these regions (38). Alignment generated by Gubbins was used to create a

maximum-likelihood based phylogenetic tree by PhyML 3.3.20180214 (Geneious plugin,

substitution model: HKY85, 100 bootstraps) (39).

Temporal signal and 'clocklikeness' of molecular phylogenies

TempEst was used to analyze the RAxML tree generated by Gubbins in conjunction with

collection year data to validate the molecular-clock assumption (40). Best-fitting root was

chosen for linear regression analyses.

Cytotoxicity, adherence, and infection assays

Viability of Vero cells after inoculation with diluted bacterial culture supernatants (1:200)

was examined using 3-[4,5-dimethylthiazole-2-yl]-2,5-diphenyltetrazolium bromide (MTT)

(5). Bacteria and Vero cells were prepared as described (31). Adherence to HEp-2 cells was

performed as described (37). For infant rabbit infection assays, litters of mixed gender 2-day-

old New Zealand White infant rabbits with the lactating doe were acquired from Charles

River (Canada, strain code 052). Infant rabbits were orogastrically inoculated on the day of

arrival with 10^9 CFU of Streptomycin-resistant strains O104:H4 C227-11 and O181:H4 17-

07187 suspended in 500µl 2.5% sodium bicarbonate (pH9) using a size 4 French catheter as

described previously, except that no ranitidine was administered (12). No antibiotics were

used prior or during infection. Infant rabbits were monitored for signs of illness and

euthanized three days (68-72 hours) post infection. Tissue samples taken from the stomach,

small intestine, cecum, and colon were homogenized in sterile PBS using a minibeadbeater-16

(Biospec Products, Inc.) and CFU determined by serial dilution, and plating on LB media

containing 200 µg/ml streptomycin (12).

Ethics statement

Rabbit studies were conducted according to protocols reviewed and approved by Brigham and

Women's Hospital Committee on Animals (IACUC protocol 2016N000334) and Animal

Welfare Assurance of Compliance (number A4752-01) in accordance with recommendations

in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health

and the Animal Welfare Act of the U.S. Department of Agriculture.

Data availability

The generated sequences were uploaded to NCBI project: PRJNA833419

(https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA833419).

Acknowledgements

We thank Ute Strutz, Thomas Garn, Tobias Größl, and Karsten Großhennig for excellent

technical assistance (Robert Koch Institute, Wernigerode, Germany). We also acknowledge

the MF1 bioinformatics and MF2 genome sequencing unit of the Robert Koch Institute

(Berlin) for the support of bioinformatic analysis and Illumina MiSeq and MinION

sequencing. We thank members of the Waldor and Flieger groups for helpful comments and

discussions during the study. AFl received funding from the German Ministry of Health (IGS-

ZOO grant number 2518FSB706) and AFl and AFr from the intensified molecular

surveillance initiative of the Robert Koch- Institute. IWC received funding from the National

Institute of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health

(Award Number 5 T32DK007477-37).

Author Contributions

conceptualized the study: C.L., A.Fl.

supervised the study: A.Fl.

conceived and designed the experiments: C.L., A.Fr., I.W.C., U.N., Y.H.G., M.K.W., A.Fl.

acquired data: C.L., A.Fr., I.W.C., C.J., P.S., N.S., F.-X.W., A.Fl.

conducted the experiments: C.L., I.W.C, A.Fl.

analyzed the data: C.L., A.Fr., I.W.C., C.J., P.S., N.S., F.-X.W., U.N., Y.H.G., A.Fl.

interpreted the results: C.L., A.Fr., I.W.C., C.J., P.S., N.S., F.-X.W., U.N., Y.H.G., M.K.W., A.Fl.

drafted the manuscript: C.L., M.K.W., A.Fl.

revised the manuscript: C.L., A.Fr., I.W.C., C.J., P.S., N.S., F.-X.W., U.N., Y.H.G., M.K.W., A.Fl.

approved the final version: C.L., A.Fr., I.W.C., C.J., P.S., N.S., F.-X.W., U.N., Y.H.G., M.K.W., A.Fl.

References

1.   Croxen MA, Law RJ, Scholz R, Keeney KM, Wlodarska M, Finlay BB. 2013. Recent advances in understanding enteric pathogenic *Escherichia coli*. Clin Microbiol Rev 26:822-80.

2.   Caprioli A, Scavia G, Morabito S. 2014. Public Health Microbiology of Shiga Toxin-Producing *Escherichia coli*. Microbiol Spectr 2:EHEC-0014-2013.

3.   Mellmann A, Bielaszewska M, Kock R, Friedrich AW, Fruth A, Middendorf B, Harmsen D, Schmidt MA, Karch H. 2008. Analysis of collection of hemolytic uremic syndrome-associated enterohemorrhagic *Escherichia coli*. Emerg Infect Dis 14:1287-90.

4.   Werber D, Krause G, Frank C, Fruth A, Flieger A, Mielke M, Schaade L, Stark K. 2012. Outbreaks of virulent diarrheagenic *Escherichia coli*--are we in control? BMC Med 10:11.

5.   Bielaszewska M, Mellmann A, Zhang W, Kock R, Fruth A, Bauwens A, Peters G, Karch H. 2011. Characterisation of the *Escherichia coli* strain associated with an outbreak of haemolytic uraemic syndrome in Germany, 2011: a microbiological study. Lancet Infect Dis 11:671-6.

6.   Frank C, Werber D, Cramer JP, Askar M, Faber M, an der Heiden M, Bernard H, Fruth A, Prager R, Spode A, Wadl M, Zoufaly A, Jordan S, Kemper MJ, Follin P, Müller L, King LA, Rosner B, Buchholz U, Stark K, Krause G, Team HI. 2011. Epidemic profile of Shiga-toxin-producing *Escherichia coli* O104:H4 outbreak in Germany. N Engl J Med 365:1771-1780.

7.   Rasko DA, Webster DR, Sahl JW, Bashir A, Boisen N, Scheutz F, Paxinos EE, Sebra R, Chin CS, Iliopoulos D, Klammer A, Peluso P, Lee L, Kislyuk AO, Bullard J, Kasarskis A, Wang S, Eid J, Rank D, Redman JC, Steyert SR, Frimodt-Moller J, Struve C, Petersen AM, Krogfelt KA, Nataro JP, Schadt EE, Waldor MK. 2011.

Origins of the *E. coli* strain causing an outbreak of hemolytic-uremic syndrome in

Germany. N Engl J Med 365:709-17.

8.   Rohde H, Qin J, Cui Y, Li D, Loman NJ, Hentschke M, Chen W, Pu F, Peng Y, Li J,

Xi F, Li S, Li Y, Zhang Z, Yang X, Zhao M, Wang P, Guan Y, Cen Z, Zhao X,

Christner M, Kobbe R, Loos S, Oh J, Yang L, Danchin A, Gao GF, Song Y, Li Y,

Yang H, Wang J, Xu J, Pallen MJ, Wang J, Aepfelbacher M, Yang R, Consortium

EcOHGAC-S. 2011. Open-source genomic analysis of Shiga-toxin-producing *E. coli*

O104:H4. N Engl J Med 365:718-24.

9.   Mellmann A, Harmsen D, Cummings CA, Zentz EB, Leopold SR, Rico A, Prior K,

Szczepanowski R, Ji Y, Zhang W, McLaughlin SF, Henkhaus JK, Leopold B,

Bielaszewska M, Prager R, Brzoska PM, Moore RL, Guenther S, Rothberg JM, Karch

H. 2011. Prospective genomic characterization of the German enterohemorrhagic

Escherichia coli O104:H4 outbreak by rapid next generation sequencing technology.

PLoS One 6:e22751.

10.   Schiller P, Knodler M, Berger P, Greune L, Fruth A, Mellmann A, Dersch P, Berger

M, Dobrindt U. 2021. The Superior Adherence Phenotype of *E. coli* O104:H4 is

Directly Mediated by the Aggregative Adherence Fimbriae Type I. Virulence 12:346-

359.

11.   Boisen N, Melton-Celsa AR, Scheutz F, O'Brien AD, Nataro JP. 2015. Shiga toxin 2a

and Enteroaggregative *Escherichia coli*--a deadly combination. Gut Microbes 6:272-8.

12.   Munera D, Ritchie JM, Hatzios SK, Bronson R, Fang G, Schadt EE, Davis BM,

Waldor MK. 2014. Autotransporters but not pAA are critical for rabbit colonization by

Shiga toxin-producing *Escherichia coli* O104:H4. Nat Commun 5:3080.

13.   Lerouge I, Vanderleyden J. 2002. O-antigen structural variation: mechanisms and

possible roles in animal/plant-microbe interactions. FEMS Microbiol Rev 26:17-47.

14.   Orskov F, Orskov I. 1992. *Escherichia coli* serotyping and disease in man and

461   animals. Can J Microbiol 38:699-704.

462   15.   Iguchi A, Iyoda S, Kikuchi T, Ogura Y, Katsura K, Ohnishi M, Hayashi T, Thomson

463   NR. 2015. A complete view of the genetic diversity of the *Escherichia coli* O-antigen

464   biosynthesis gene cluster. DNA Res 22:101-7.

465   16.   Lang C, Hiller M, Konrad R, Fruth A, Flieger A. 2019. Whole-Genome-Based Public

466   Health Surveillance of Less Common Shiga Toxin-Producing *Escherichia coli*

467   Serovars and Untypeable Strains Identifies Four Novel O Genotypes. J Clin Microbiol

468   57.

469   17.   Samuel G, Hogbin JP, Wang L, Reeves PR. 2004. Relationships of the *Escherichia*

470   *coli* O157, O111, and O55 O-antigen gene clusters with those of *Salmonella enterica*

471   and Citrobacter freundii, which express identical O antigens. J Bacteriol 186:6536-43.

472   18.   Geue L, Menge C, Eichhorn I, Semmler T, Wieler LH, Pickard D, Berens C, Barth

473   SA. 2017. Evidence for Contemporary Switching of the O-Antigen Gene Cluster

474   between Shiga Toxin-Producing *Escherichia coli* Strains Colonizing Cattle. Front

475   Microbiol 8:424.

476   19.   Milkman R, Jaeger E, McBride RD. 2003. Molecular evolution of the *Escherichia coli*

477   chromosome. VI. Two regions of high effective recombination. Genetics 163:475-83.

478   20.   Zhou Z, Alikhan NF, Mohamed K, Fan Y, Agama Study G, Achtman M. 2020. The

479   EnteroBase user's guide, with case studies on Salmonella transmissions, *Yersinia*

480   *pestis* phylogeny, and Escherichia core genomic diversity. Genome Res 30:138-152.

481   21.   Kunne C, Billion A, Mshana SE, Schmiedel J, Domann E, Hossain H, Hain T,

482   Imirzalioglu C, Chakraborty T. 2012. Complete sequences of plasmids from the

483   hemolytic-uremic syndrome-associated *Escherichia coli* strain HUSEC41. J Bacteriol

484   194:532-3.

485   22.   Gaillot O, Clement C, Simonet M, Philippon A. 1997. Novel transferable beta-lactam

486   resistance with cephalosporinase characteristics in *Salmonella enteritidis*. J

487   Antimicrob Chemother 39:85-7.

488   23.   Bielaszewska M, Prager R, Kock R, Mellmann A, Zhang W, Tschape H, Tarr PI,

489   Karch H. 2007. Shiga toxin gene loss and transfer in vitro and in vivo during

490   enterohemorrhagic *Escherichia coli* O26 infection in humans. Appl Environ Microbiol

491   73:3144-50.

492   24.   Faruque SM, Sack DA, Sack RB, Colwell RR, Takeda Y, Nair GB. 2003. Emergence

493   and evolution of *Vibrio cholerae* O139. Proc Natl Acad Sci U S A 100:1304-9.

494   25.   Sugiyama T, Kido N, Kato Y, Koide N, Yoshida T, Yokochi T. 1997. Evolutionary

495   relationship among rfb gene clusters synthesizing mannose homopolymer as O-

496   specific polysaccharides in *Escherichia coli* and Klebsiella. Gene 198:111-3.

497   26.   Feng P, Lempel KA, Karch H, Whittam TS. 1998. Genotypic and phenotypic changes

498   in the emergence of *Escherichia coli* O157:H7. J Infect Dis 177:1750-1753.

499   27.   Ingle DJ, Valcanis M, Kuzevski A, Tauschek M, Inouye M, Stinear T, Levine MM,

500   Robins-Browne RM, Holt KE. 2016. In silico serotyping of *E. coli* from short read

501   data identifies limited novel O-loci but extensive diversity of O:H serotype

502   combinations within and between pathogenic lineages. Microb Genom 2:e000064.

503   28.   Ochman H, Selander RK. 1984. Evidence for clonal population structure in

504   *Escherichia coli*. Proc Natl Acad Sci U S A 81:198-201.

505   29.   Wieler LH, Semmler T, Eichhorn I, Antao EM, Kinnemann B, Geue L, Karch H,

506   Guenther S, Bethe A. 2011. No evidence of the Shiga toxin-producing *E. coli*

507   O104:H4 outbreak strain or enteroaggregative *E. coli* (EAEC) found in cattle faeces in

508   northern Germany, the hotspot of the 2011 HUS outbreak area. Gut Pathog 3:17.

509   30.   Coipan CE, Friesema IH, van den Beld MJC, Bosch T, Schlager S, van der Voort M,

510   Frank C, Lang C, Fruth A, Franz E. 2022. Sporadic Occurrence of Enteroaggregative

Shiga Toxin-Producing *Escherichia coli* O104:H4 Similar to 2011 Outbreak Strain.

Emerg Infect Dis 28:1890-1894.

31. Prager R, Lang C, Aurass P, Fruth A, Tietze E, Flieger A. 2014. Two novel

EHEC/EAEC hybrid strains isolated from human infections. PLoS One 9:e95379.

32. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA,

Zeng Q, Wortman J, Young SK, Earl AM. 2014. Pilon: an integrated tool for

comprehensive microbial variant detection and genome assembly improvement. PLoS

One 9:e112963.

33. Darling AE, Mau B, Perna NT. 2010. progressiveMauve: multiple genome alignment

with gene gain, loss and rearrangement. PLoS One 5:e11147.

34. Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version

7: improvements in performance and usability. Mol Biol Evol 30:772-80.

35. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016.

PHASTER: a better, faster version of the PHAST phage search tool. Nucleic Acids

Res 44:W16-21.

36. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes

S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA,

McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R,

Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST Server: rapid

annotations using subsystems technology. BMC Genomics 9:75.

37. Lang C, Fruth A, Holland G, Laue M, Muhlen S, Dersch P, Flieger A. 2018. Novel

type of pilus associated with a Shiga-toxigenic *E. coli* hybrid pathovar conveys

aggregative adherence and bacterial virulence. Emerg Microbes Infect 7:203.

38. Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, Parkhill J,

Harris SR. 2015. Rapid phylogenetic analysis of large samples of recombinant

bacterial whole genome sequences using Gubbins. Nucleic Acids Res 43:e15.

39. Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010. New

algorithms and methods to estimate maximum-likelihood phylogenies: assessing the

performance of PhyML 3.0. Syst Biol 59:307-21.

40. Rambaut A, Lam TT, Max Carvalho L, Pybus OG. 2016. Exploring the temporal

structure of heterochronous sequences using TempEst (formerly Path-O-Gen). Virus

Evol 2:vew007.

 Figure-Legends:

 Figure 1: STEC O181:H4 17-07187 shares close phylogenetic relationship and similar

 virulence traits with the O104:H4 outbreak strain. (A) Overlapping virulence gene profiles

 of the O104:H4 outbreak strain (FWSEC0009) and the O181:H4 strain 17-07187. (B)

 Minimal spanning tree of O181:H4 17-07187 and selected O104:H4 strains based on

 cgMLST involving 2513 alleles confirms their close phylogenetic relationship. Strains

 representing phylogenetic diversity of O104:H4 isolates from humans with respect to isolation

 time (from 1998 to 2022) and location were selected. Only one cluster representative was

 used, except of the 2011 outbreak. O181:H4 17-07187 and O104:H4 strains 2009EL-2071,

 Ec12-0465, and outbreak strains from 2011 (including strains FWSEC0009, C227-11, and 11-

 02027) differ by only 30-34 alleles. An allele represents a variant form of a gene. Numbers in

 blue indicate allelic distances. Serotypes and AAF type are indicated in the key. For complete

 strain designations see Fig. S1A. (C) Cytotoxicity of O181:H4 17-07187 towards Vero cells

 and (D) adherence pattern to HEp-2 cells are comparable to the O104:H4 outbreak strain 11-

 02027. Results are representative of at least two additional experiments. EHEC O157:H7

 EDL933 producing Stx1 and Stx2 served as reference in the cytotoxicity assay and was set to

 100%. The cytotoxicity results represent the means and standard deviations of triplicate

 samples (n=3) and are representative of at least two additional experiments. E) Intestinal

 colonization in infant rabbits inoculated with O104:H4 outbreak strain C227-11 (n=6) or

 O181:H4 strain 17-07187 (n=5). CFU/g (colony forming units per gram of tissue) denotes

 concentration of bacteria recovered three days post inoculation from homogenated intestinal

 tissues. Data points represent individual rabbits from two independent litters split between the

 two strains. Bars show the geometric mean, SI is small intestine, ND is not detected.

Figure 2: O181:H4 17-07187 and O104:H4 outbreak strain genomes mainly differ in

OAGCs and mobile genetic elements. (A) Whole chromosome MAUVE alignment of

O181:H4 17-07187 and the O104:H4 outbreak strain FWSEC0009 highlighting mobile

genetic elements and differences in phage regions, IS-/ transposon elements, and OAGC

regions. (B) OAGCs of O181:H4 17-07187 and the O104:H4 FWSEC0009 outbreak strain

are flanked by homologous upstream (*galF*) and downstream (*gnd* to *hisI*) regions. MAFFT

alignment shows that the regions in between *galF* and *gnd* are very different. (C) MAFFT

alignment of the *stx2a*-encoding phages in O181:H4 17-07187 and the outbreak strain

O104:H4 outbreak strain FWSEC0009 shows that they are very similar (99,9% nucleotide

identity).

Figure 3: Recent global isolates of serotype O181:H4 and five other O groups belong to

ST678. (A) Summary of serotypes, isolation dates, along with relatedness of genomes

(cgMLST-based allelic distances) and mobile genetic elements (plasmids and *stx* phage) of

the 34 non-O104:H4 ST678 (ST12598, ST12610) clinical strains (found on EnteroBase)

compared to the O104:H4 outbreak strain FSWEC0009. (B) Serotypes of the non-O104:H4

ST678 strains and the O104:H4 outbreak strain FSWEC0009 in a minimal spanning tree

based on cgMLST. Numbers in blue indicate allelic distances. Although, phylogenetically

very close to ST678 strains, O181:H4 strains 1472912 and 1472968 in fact belong to

ST12610 due to a point mutation in *icd* and OgN-RKI9:H4 strain 608450 to ST12598 due to a

point mutation in *recA*.

Figure 4: O-antigen gene clusters and phylogeny within *E. coli* ST678. (A) O-antigen

gene clusters in ST678 strains differ in their gene contents and organization, yet they are

flanked by homologous regions starting upstream by *galF* and downstream by *gnd* to *hisI*,

respectively. (B) Maximum-likelihood phylogenetic tree generated by PhyML based on a

recombination-corrected alignment of genome-wide polymorphic sites. Strains selected for

representation in Fig. 1B and Fig. 3 were combined and O104:H4 2011 outbreak strain

FWSEC0009 served as reference for read mapping. The tree was rooted with an outgroup

consisting of *E. coli* strains K12 c600, Ec042, EDL933, O157:H7 strain Sakai, O26:H11

strain 11368, and O103:H2 strain 12009. Genome sequences from outgroup *E. coli* isolates

had been included in the phylogenetic analysis and for clarity are not shown in Fig. 4B (large

phylogenetic distance and associated limited resolution of the clades of interest) but in Fig.

4C. Bootstrap values >90% are indicated as black dots. Serotypes are as indicated in the key.

Heatmap on the right indicates the presence (filled box) or absence (white box) of genes for

AFF/I, AFF/III, and *stx2a*. Scale bar refers to a phylogenetic distance of 0.0002 nucleotide

substitutions per site. Grey boxes indicate clades. The box in light red highlights 2011

outbreak strains. (C) Maximum-likelihood phylogenetic tree with additional outgroup *E. coli*

isolates and strains represented in Fig. 4B as collapsed tree branch designated as ST678

strains.

A

		EHEC virulence genes													EAEC virulence genes											
		chromosome							plasmid						chromosome				pAA							
serotype	strain	stx	eae	lpf_{O26}	lpf_{O113}	terA	irp2	fyuA	ehxA	subAB	espP	iha	saa	set1	pic	sigA	aaiC	aatA,B,C,D,P	AAFI	aap	aggR	EAST1	iucA	sepA	
O104:H4	outbreak 2011	2a		+	+	+	+	+				+		+	+	+		+	+	+	+		+	+	
O181:H4	17-07187	2a		+	+	+	+	+				+		+	+	+		+	+	+	+	+	+	+	

B

C

Vero cell cytoxicity

D

Hep-2 cell adherence

O181:H4 17-07187
O104:H4 outbreak 2011
O157:H7 EDL933
O16:H48 K12 c600
not infected

E

Infant rabbit colonization

A — Chromosomes

O181:H4
17-07187 — regions of difference — phages — *stx2*-phage

O104:H4
2011 outbreak
FWSEC0009 — regions of difference — phages — *stx2*-phage

- ▮ phage intact
- ▯ phage questionable
- ▮ phage incomplete
- ▼ is-element/ transposon lacking in FWSEC0009
- ▽ is-element/ transposon lacking in 17-07187
- ⯅ OAGC genes and genes around differing between both strains
- ▽ genes lacking in FWSEC0009
- ▼ genes lacking in 17-07187

B — O-antigen gene regions

100% identity   30% to <100% identity   <30% identity   identity

O181
O104

38% identity   43% identity

- UTP-glucose-1-phosphate uridylyltransferase (*galF*)
- phosphoribosyl-AMP cyclohydrolase (*hisI*)
- O-antigen polymerase (*wzy*)
- O-antigen flippase (*wzx*)
- O-antigen chain length determinant protein (*wzz*)
- UDP-glucose 6-dehydrogenase (*ugd*)
- 6-phosphogluconate dehydrogenase (*gnd*)
- glycosyl transferase
- hypothetical protein
- capsular polysaccharide phosphotransferase (*cps12A*)
- UDP-N-acetylglucosamine 4,6-dehydratase (*capD*)
- dTDP-4-dehydrorhamnose reductase (*rmlD*)
- UDP-2,3-diacetamido-2,3-dideoxy-D-glucuronate 2-epimerase (*wpbI*)
- UDP-glucose 4-epimerase (*galE*)
- acetyl transferase (*nnaD*)
- sialic acid synthase (*nnaB*)
- N-acylneuraminate cytidylyltransferase (*nnaC*)
- UDP-N-acetylglucosamine 2-epimerase (*nnaA*)

C — *stx* phages

O181:H4
17-07187 — 99% identity

O104:H4
2011 outbreak
FWSEC0009 — *stx2*a

100% identity   30% to <100% identity   <30% identity

strain	serotype	country of isolation	travel association	year of isolation	allele distance to strain FWSEC0009	FWSEC0009 O104:H4				
						pl1	pl2 pAA		pl3	stx2a-phage
						cov.	cov.	AAF	cov.	cov.
FWSEC0009	O104:H4	Germany		2011	0	100	100	AAF/I	100	100
MEX-14	O181:H4	Mexico		2011	26	5.7	100	AAF/I	100	
497801	O181:H4	United Kingdom	Morocco	2013	27	1.7	100	AAF/I	100	
147828	O181:H4	United Kingdom	Morocco	2015	28	90.7	100	AAF/I	100	
B141_2	O181:H4	Ecuador		2015	28	2.9	95.5	AAF/I	100	
143508	O181:H4	United Kingdom		2015	30	17.5	100	AAF/I	100	
PNUSAE010936	O181:H4	United States		2017	30	1.1	100	AAF/I	100	
644185	O181:H4	United Kingdom	Mexico	2018	31	8.4	100	AAF/I	100	
PNUSAE023691	O181:H4	United States		2019	31	1.1	100	AAF/I	99.9	
285561	O181:H4	United Kingdom	Egypt	2016	32	32.1	100	AAF/I	100	
470195	O181:H4	United Kingdom	India	2018	32	16.1	100	AAF/I	20.3	100
PNUSAE010931	O181:H4	United States		2017	32	1.6	100	AAF/I	100	
201510555	O181:H4	France		2015	34	9	100	AAF/I	0	100
1078123	O181:H4	United Kingdom		2021	35	1.9	97.1	AAF/I	100	
1472912	O181:H4	United Kingdom		2021	35	4.7	100	AAF/I	100	
17-07187	O181:H4	Germany		2017	35	37.3	100	AAF/I	0	100
1472968	O181:H4	United Kingdom		2021	36	4.3	100	AAF/I	100	
SCPM-O-B-9428	O181:H4	Russia		2018	36	36.8	100	AAF/I	87.8	98.9
822586	O181:H4	United Kingdom	Morocco	2019	38	3.3	99.7	AAF/I	22.1	
896001	O181:H4	United Kingdom		2020	38	32.7	99.6	AAF/I	100	
PNUSAE005891	O181:H4	United States		2016	38	35.6	100	AAF/I	100	58.1
PNUSAE005897	O181:H4	United States		2017	39	35	99.9	AAF/I	100	58.4
649913	OX13:H4	United Kingdom	Ethiopia	2018	46	5.4	99.4	AAF/I	100	
524333	O127:H4	United Kingdom	India	2018	53	3.6	100	AAF/I	18.9	
524335	O127:H4	United Kingdom	India	2018	53	3	100	AAF/I	53.6	
535309	O127:H4	United Kingdom		2018	54	14.8	100	AAF/I	19.4	
789931	O127:H4	United Kingdom	Kenya	2019	106	17.5	70.8	AAF/III	13	
608450	OgN-RKI9:H4	United Kingdom		2018	112	7.3	62.1	AAF/III	30.5	
ALQ019547	O127:H4	Kenya		2015	113	16.2	69.6	AAF/III	0	
ALQ019554	O127:H4	Kenya		2015	113	16.4	69.3	AAF/III	0	
TMP018942	O127:H4	Kenya		2015	113	16.1	69.5	AAF/III	0	
103016	O131:H4	Gambia		2015	121	3.5	60.4	AAF/III	5.9	
178926	O131:H4	United Kingdom		2019	182	31.2	62.9	AAF/III	13	
102906	O69:H4	Gambia		2009	187	15	70.2	AAF/III	5.7	
808295	O131:H4	United Kingdom	India	2019	189	4.8	60.9	AAF/III	52.6	

AAF/I (dark grey) AAF/III (light grey)

0 ————— 100% coverage of reference

A   O-antigen gene regions

O104
O181
O127
O131
O69
OX13
OgN-RKI9

- O-antigen polymerase (*wzy*)
- UTP-glucose-1-phosphate uridylyltransferase (*galF*)
- O-antigen chain length determinant protein (*wzz*)
- acetyl transferase
- GDP-L-fucose synthetase (*fcl*)
- UDP-N-acetylglucosamine 2-epimerase (*nnaA*)
- UDP-N-acetylglucosamine 4,6-dehydratase (*capD*)
- CatB-related O-acetyltransferase
- hypothetical protein
- dTDP-4-dehydrorhamnose 3,5-epimerase (*rmlC*)
- acyltransferase

- O-antigen flippase (*wzx*)
- 6-phosphogluconate dehydrogenase (*gnd*)
- phosphoribosyl-AMP cyclohydrolase (*hisI*)
- sialic acid synthase (*nnaB*)
- capsular polysaccharide phosphotransferase (*cps12A*)
- mannose-1-phosphate guanylyltransferase (*manC*)
- GDP-mannose mannosyl hydrolase (*gmm*)
- UDP-glucose 4-epimerase (*galE*)
- dTDP-glucose 4,6 dehydratase (*rmlB*)
- coenzyme F420 hydrogenase
- NAD-dependent epimerase (*wbnF*)

- glycosyl transferase
- UDP-glucose 6-dehydrogenase (*ugd*)
- N-acylneuraminate cytidylyltransferase (*nnaC*)
- Phosphomannomutase (*manB*)
- dTDP-4-dehydrorhamnose reductase (*rmlD*)
- GDP-mannose 4,6-dehydratase (*gmd*)
- UDP-2,3-diacetamido-2,3-dideoxy-D-glucuronate 2-epimerase (*wpbI*)
- haloacid dehalogenase-like hydrolase
- glucose-1-phosphate thymidylyltransferase (*rmlA*)
- pyruvyl transferase

B   *E. coli* ST678 phylogeny

0.0002

AAF/I   AAF/III   stx2a

Legend:
- O104:H4 (pink)
- O127:H4 (yellow)
- O131:H4 (light green)
- O181:H4 (blue)
- OX13:H4 (olive green)
- O69:H4 (purple)
- OgN-RKI9:H4 (red)

I
Ia
Ib

outbreak 2011

C

Ec042
O157:H7 str. Sakai
EDL933
K12 c600
O26:H11 str. 11368
O103:H2 str. 12009
ST678 strains

0.2