Zinc-finger protein 518 plays a crucial role in pericentromeric heterochromatin formation by linking satellite DNA to heterochromatin

Shinya Ohta[1,2]*, Jun-Ichirou Ohzeki[3], Nobuko Sato[2], Ken-Ichi Noma[1,4], and Hiroshi Masumoto[3]*

[1]Institute for Genetic Medicine, Hokkaido University, Kita-15, Nishi-7, Kita-Ku, Sapporo 060-0815, Japan

[2]Department of Biochemistry, Medical School, Kochi University, Kohasu, Oko-cho, Nankoku, Kochi 783-8505, Japan

[3]Laboratory of Chromosome Engineering, Department of Frontier Research and Development, Kazusa DNA Research Institute, 2-6-7 Kazusa-Kamatari, Kisarazu 292-0818, Japan

[4]Institute of Molecular Biology, University of Oregon, 1370 Franklin Blvd, Eugene, OR, 97403, USA

*Corresponding authors:

shinya.ohta@igm.hokudai.ac.jp

masumoto@kazusa.or.jp

Summary

Aneuploidy is caused by chromosomal missegregation and is frequently observed in

cancers and hematological diseases. Pericentromeric heterochromatin formation is

essential for proper chromosome segregation; however, it is unclear how

heterochromatin is targeted to pericentromeres. In this study, we investigated the

involvement of two homologous zinc-finger proteins (ZNF518A and ZNF518B) in

heterochromatin formation at pericentromeres by determining the cellular localization

of ZNF518, centromeric proteins such as CENP-A and CENP-B, and heterochromatin

factors, including HP1 and histone H3K9 trimethylation. The results showed that

various segments in ZNF518 interact with CENP-B, HP1, and G9a histone H3K9

methyltransferase. In conclusion, we identified a novel mechanism underlying

pericentromeric heterochromatin formation mediated by ZNF518.

Keywords

centromere, pericentromere, heterochromatin, ZNF518, CENP-B

Introduction

Defects in centromere or kinetochore functions are related to chromosomal abnormalities, and anomalous chromosome segregation can lead to cancer or hematological diseases.[1,2] Chromosomal abnormalities can occur during mitosis, where replicated chromosomes are normally equally distributed between two daughter cells. During mitosis, compacted chromosomes are segregated when the spindle attaches to and pulls the kinetochore, a proteinaceous structure on the centromere. Centromere protein (CENP)-A, a centromere-specific histone H3 variant, is essential for kinetochore formation and directs the assembly of a complex containing at least 15 other proteins, including CENP-C.[3] CENP-A, CENP-B, and CENP-C are evolutionarily conserved from fission yeast to vertebrates and show distinct association patterns within the centromere.[4,5]

In normal human chromosomes, centromeres assemble on α-satellite DNA (alphoid DNA), a giant repetitive DNA locus consisting of 171-bp repeat units.[6,7] CENP-A assembles on a portion of α-satellite DNA, forming an epigenetic mark of kinetochore assembly.[8–11] Some α-satellite DNA subclasses contain a 17-nucleotide CENP-B box, bound by CENP-B.[8,12,13] Thus, CENP-B is distributed in both CENP-A chromatin (core centromere) and pericentromeric heterochromatin. CENP-B is involved in the assembly of CENP-A and CENP-C in centromeres and is crucial for functional centromere formation.[9,10] Furthermore, CENP-B and CENP-B boxes are required for human artificial chromosome formation and *de novo* centromeric CENP-A chromatin assembly in cells with synthetic alphoid DNA.[14–16] CENP-B also induces

heterochromatin formation in alphoid DNA integrated into chromosomal arms.[16]

Interestingly, CENP-B recruits numerous factors, including the open chromatin

modifier ASH1L (H3K36 methylase) and heterochromatin factor HP1, in a mutually

exclusive manner.[17] Heterochromatin plays important roles in chromosome stability

and the maintenance of sister chromatid cohesion during chromosome

segregation.[18,19] Therefore, both centromeric chromatin and heterochromatin are

important for accurate chromosome segregation.

Pericentromeric heterochromatin insulates centromeres from the rest of the

genome and is required for proper mitotic chromosome segregation.[19,20]

Heterochromatin has a more condensed structure than euchromatin and carries

specific post-translational histone modifications. Nucleosomal surfaces with charge

differences contribute to the condensed structure of heterochromatin.[21–23] Histone H3

lysine 9 trimethylation (H3K9me3) is recognized and bound by the chromodomain of

the heterochromatin protein HP1.[24] HP1 in turn recruits the H3K9 methyltransferase,

suppressor of variegation 3-9 homolog (SUV39H).[18,25] The HP1-mediated loading of

SUV39H to chromatin is required for the maintenance of heterochromatin at

pericentromeres.[26,27] As SUV39H localizes to pericentromeres in addition to other

constitutive heterochromatin regions, heterochromatin formation at pericentromeres

involves SUV39H. The mechanisms by which heterochromatin factors are initially

targeted to pericentromeres, however, remain elusive.[28]

H3K9 methyltransferases other than SUV39H have been identified. SET

domain bifurcated 1 (SETDB1) catalyzes H3K9me3 formation, whereas euchromatic

histone lysine methyltransferases 1 and 2 (GLP and G9a) are responsible for

H3K9me1 and H3K9me2, respectively. G9a can also methylate Lys9 and Lys27 of

histone H3.[29] These H3K9 methyltransferases, including SUV39H, are selectively

loaded onto distinct genomic regions through protein interactions with other factors.[30]

G9a and GLP bind to the widely interspaced zinc-finger (WIZ) protein.[29,31,32] H3K9

methylation at chromosomal arm regions is mostly lost in G9a-deficient cells, whereas

H3K9 methylation at pericentromeres is largely lost in SUV39H-deficient cells.[29]

Moreover, G9a interacts with zinc-finger protein 518B (ZNF518B) as recently indicated

by a proteomics analysis.[32] Our previous proteomics analysis revealed a possible role

for ZNF518B as a functional protein on mitotic chromosomes.[33,34] The present study

demonstrated that G9a, which acts as a H3K9 methyltransferase for chromosomal

arm regions, is also involved in pericentromeric heterochromatin formation via

interaction with ZNF518B recruited by CENP-B.

Results

Human ZNF518A/B is recruited to centromeres by CENP-B

We first examined the localization of green fluorescent protein (GFP)-tagged

ZNF518B and CENP-A, CENP-B, and CENP-C in human U2OS cells. Overexpression

of GFP fusion proteins allows for visualization of even weakly expressed proteins;

using this technique, we were able to successfully visualize the localization of ZNF518.

GFP-ZNF518B co-localized proximally to the centromeric factors (Fig. 1A–F).

ZNF518B localized to the centromeres during mitosis (Fig. S1A), suggesting

constitutive centromeric localization of ZNF518B throughout the cell cycle. GFP-

ZNF518B foci were slightly shifted away from CENP-A and CENP-C foci, but coincided

with CENP-B foci, indicating that ZNF518B localizes to the pericentromeric region,

where it is bound by CENP-B (Fig. 1A–F).

To investigate the localization dependency between ZNF518B and CENP-B,

we examined the localization of GFP-ZNF518B in *CENPB*-knockout (KO) HeLa

cells.[17] Centromeric/pericentromeric localization of GFP-ZNF518B was lost in

CENPB-KO cells, but was restored upon exogeneous CENP-B expression (Fig. 1G–

I), indicating that the centromeric localization of ZNF518B depends on CENP-B.

ZNF518A is a paralogue of ZNF518B present in humans (Fig. S1B). To

determine the localization of ZNF518A, GFP-ZNF518B and Halo-ZNF518A were

transiently co-expressed in U2OS cells (Fig. 1J). GFP-ZNF518B and Halo-ZNF518A

foci co-localized at the centromeres. Again, the co-localized signals were slightly

shifted away from CENP-A foci, suggesting that both ZNF518A and ZNF518B localize

at centromeres (Fig. 1J). Centromeric localization of GFP-ZNF518A was lost in

CENPB-KO cells, but restored upon exogenous CENP-B expression (Fig. S1C),

indicating that the centromeric localization of ZNF518A depends on CENP-B. Taken

together, the results suggested that both ZNF518A and ZNF518B localize to the

pericentromere via a common molecular mechanism involving CENP-B (Fig. S1D).

8   The C-terminus of ZNF518A/B is required for centromeric localization

To unravel the molecular mechanisms underlying ZNF518 recruitment to centromeres,

we sought to identify the segments in the 1,074-amino-acid (aa) ZNF518B protein that

are required for centromeric localization. We examined the centromeric localization of

15 ZNF518B derivatives expressed as GFP fusion proteins (Fig. 2). ZNF518B

derivatives containing regions from the N-terminus to 628 aa (ZNF518B$^{1-398}$,

ZNF518B$^{344-628}$, and ZNF518B$^{1-628}$) did not show centromeric localization (Fig. 2,

S2A). Therefore, the N-terminus of ZNF518B does not form sufficient interactions to

localize the protein to centromeres. Among four ZNF518B derivatives with

incrementing C-terminal truncations (ZNF518B$^{344-1074}$, ZNF518B$^{344-987}$, ZNF518B^{344-}

^{877}, and ZNF518B$^{344-769}$), ZNF518B$^{344-1074}$, ZNF518B$^{344-987}$, and ZNF518B$^{344-877}$

showed centromeric localization in 70.7%, 58.9%, and 57.1% of transfected cells,

respectively, whereas ZNF518B$^{344-769}$ did not show centromeric localization (Fig. 2).

Similarly, among ZNF518B$^{555-1074}$, ZNF518B$^{555-877}$, and ZNF518B$^{555-769}$, only

ZNF518B$^{555-769}$, which had the same C-terminal truncation as ZNF518B$^{344-769}$, did not

show centromeric localization (Figs. 2, S2A). These results suggested that residues

770–877 of ZNF518B are involved in its centromeric localization. However, only 2.2% of transfected cells showed centromeric localization of ZNF518B$^{770-877}$ (Fig. 2). Therefore, residues 770–877 are important, but insufficient for centromeric localization.

Given the large difference in centromeric localization among ZNF518B$^{555-1074}$, ZNF518B$^{555-877}$, and ZNF518B$^{555-769}$ (in 64.5%, 31.8%, and 0% of transfected cells, respectively), we examined the centromeric localization of ZNF518B derivatives carrying residues 770–1074 (Figs. 2, S2A). ZNF518B$^{878-1074}$, ZNF518B$^{878-987}$, and ZNF518B$^{770-987}$ showed centromeric localization in 53.6%, 56.7%, and 89.3% of cells, respectively (Figs. 2, S2A). ZNF518B$^{770-987}$ showed the highest frequency of centromeric localization among all ZNF518B derivatives. These results suggested that residues 770–987 are involved in the centromeric localization of ZNF518B.

We next examined the centromeric localization of ZNF518A derivatives (Fig. S2B). Wild-type (WT) ZNF518A (ZNF518A^{WT}) and two derivatives containing the C-terminal 1317–1483-aa segment (ZNF518A$^{971-1483}$ and ZNF518A$^{1317-1483}$) showed high frequencies of centromeric localization, suggesting that the C-terminal portion of ZNF518A harbors crucial segment(s) for centromeric localization (Fig. S2B). Notably, the C-terminal segments of ZNF518A and ZNF518B required for centromeric localization carry PxV(I)xL HP1-interaction motifs, which are widely conserved among higher vertebrates, including fish (Fig. S3), implying that their role is evolutionarily conserved potentially through their localization at centromeres.

The C-terminal PxIxL motif is required for the centromeric localization of ZNF518B

To elucidate the functional significance of the PxV(I)xL motifs in ZNF518B, we constructed the GFP-ZNF518B$^{KSS/C}$ derivative in which the conserved hydrophobic residues PxIxL in the C-terminus were substituted with the hydrophilic residues KxSxS (Fig. 2). ZNF518B$^{KSS/C}$ exhibited less frequent centromeric localization (14.3%) than ZNF518^{WT} (77.5%; Fig. 2). In contrast, the centromeric localization of ZNF518B$^{KSS/N}$, which carried a substitution of the PxVxL motif with KxSxS in the N-terminus, was comparable to that of ZNF518^{WT} (Fig. 2). ZNF518B$^{KSS/NC}$, which carried KxSxS substitutions of the PxV(I)xL motifs in both the N- and C-termini, showed a similar centromeric localization frequency (21.4%) as ZNF518B$^{KSS/C}$ (Fig. 2). Therefore, the PxIxL motif present in the C-terminus is involved in the centromeric localization of ZNF518B. ZNF518A$^{KSS/C}$ also showed reduced centromeric localization (Fig. S2B). These results demonstrated that the conserved PxV(I)xL motif in the C-terminus of ZNF518 is important for centromeric localization.

Although the C-terminal PxIxL motif is important for centromeric localization, ZNF518B$^{KSS/C}$ and ZNF518B$^{KSS/NC}$ localized to the centromeres in 14.3% and 21.4% of cells, respectively, implying the involvement of other interactions in centromeric localization. To test this, we substituted the PxVxL motif with KxSxS in the ZNF518B$^{770–987}$ derivative that localized to centromeres with the highest frequency (89.3%), yielding ZNF518B$^{770–987 \ KSS/C}$. ZNF518B$^{770–987 \ KSS/C}$ showed centromeric localization in 60.6% of transfected cells (Fig. 2, S2B), indicating that both the PxIxL

motif and the C-terminal segment (770–987 aa) participate in the centromeric

localization of ZNF518B.

ZNF518B interacts with HP1α and HP1β

The localization of full-length ZNF518B^{WT} to centromeres and pericentromeres

requires CENP-B. The ZNF518B$^{770-987}$ derivative with the C-terminal segment (770–

987 aa) including the PxIxL motif also localized to centromeres (Fig. 2). In *CENPB*-

KO cells, ZNF518B$^{770-987}$ showed a focal localization that was not observed for

ZNF518B^{WT} (Fig. S4A). However, the ZNF518B$^{770-987}$ foci were distant from

centromeres and rather co-localized with heterochromatin foci in *CENPB*-KO cells (Fig.

S4B). Thus, ZNF518B$^{770-987}$ can be recruited to non-centromeric heterochromatin

domains through its interaction with heterochromatic factors in the absence of CENP-

B (Fig. S4C).

Given its similarity to the canonical HP1-interacting motif PxVxL, we

hypothesized that the PxIxL motif in the C-terminus of ZNF518B interacts with HP1,

resulting in the accumulation of ZNF518B at heterochromatin regions in *CENPB*-KO

cells. Three isozymes of human HP1 have been identified: HP1α, HP1β, and HP1γ.[35]

We evaluated the interactions between ZNF518B and the HP1 isoforms using the

fluorescence microscopy-based interaction-trap (FMIT) assay,[36] which can be used to

examine protein interactions on chromatin *in vivo*. Given the overabundance of target

proteins compared to their endogenous counterparts, this technique allows for the

detection of direct and indirect protein–protein interactions using fluorescence microscopy.[17,20,36–40]

The FMIT assay was performed using HeLa-Int-03 cells, in which approximately 15,000 repeats of a synthetic alphoid DNA dimer unit (derived from the human chromosome 21 centromere) containing the *tet*O sequence (alphoid[*tetO*]) were ectopically integrated into a chromosome arm region, and *CENPB* was knocked out (Fig. 3A).[17] TetR-EYFP-ZNF518B localized to the alphoid[*tetO*] repeats, but did not exhibit centromeric localization due to the absence of CENP-B. We next examined whether Halo-HP1 co-expressed with TetR-EYFP-ZNF518B would be recruited to the alphoid[*tetO*] array (Fig. 3B). Halo-HP1α, -HP1β, and -HP1γ were recruited to alphoid[*tetO*] in 48%, 58%, and 14% of cells, respectively (Fig. 3C). In negative control cells expressing TetR-EYFP not fused to ZNF518B, Halo-HP1α, -HP1β, and -HP1γ were barely recruited to alphoid[*tetO*] (Fig. 3C). These results suggested that ZNF518B interacts with HP1.

ZNF518B interacts with HP1 isoforms via the C-terminal PxIxL motif

We next investigated which segments in ZNF518B are required for interaction with the HP1 isoforms by subjecting the ZNF518B derivatives described in Figure 2 to the FMIT assay (Fig. 3C). Upon co-expression with TetR-EYFP-fused ZNF518B derivatives (ZNF518B$^{1–398}$, ZNF518B$^{1–628}$, ZNF518B$^{344–628}$, ZNF518B$^{344–877}$, ZNF518B$^{555–877}$, and ZNF518B$^{770–877}$), Halo-HP1 was scarcely or not recruited to alphoid[*tetO*]. As these

derivatives lacked the PxIxL motif, this result indicated the importance of the PxIxL

motif in the interactions between ZNF518B and the HP1 isoforms (Fig. 3C).

We further examined the significance of the PxV(I)xL motifs in the ZNF518B-

HP1 interactions using TetR-EYFP-fused ZNF518B derivatives (ZNF518B$^{KSS/N}$,

ZNF518B$^{KSS/C}$, ZNF518B$^{KSS/NC}$, and ZNF518B$^{770-987\ KSS/C}$) carrying PxV(I)xL

substitutions with KxSxS, as depicted in Figure 2. Halo-HP1 was scarcely recruited to

alphoid^{tetO} by the ZNF518B derivatives, except for ZNF518B$^{KSS/N}$. Therefore, the PxIxL

motif in the C-terminus, not the PxVxL motif in the N-terminus, participates in the

interaction with HP1 (Fig. 3C).

Next, the TetR-EYFP-fused ZNF518B derivatives ZNF518B$^{344-1074}$,

ZNF518B$^{344-987}$, ZNF518B$^{555-1074}$, ZNF518B$^{878-1074}$, ZNF518B$^{878-987}$, and

ZNF518B$^{770-987}$, carrying the PxIxL motif in the ZNF518B C-terminus, were subjected

to the FMIT assay. Halo-HP1 was recruited to alphoid^{tetO} in more than 75% of cells

(Fig. 3C). Remarkably, these N-terminal-deleted derivatives were able to recruit the

three HP1 proteins with similar frequencies. Furthermore, ZNF518B$^{344-1074}$ was more

efficient at recruiting HP1 than ZNF518B^{WT}. Therefore, the N-terminal segment (1–

398) is inhibitory for ZNF518B interactions with HP1 isoforms (Fig. 3C).

The ZNF518B PxIxL motif interacts with the chromo-shadow domain of HP1

HP1 comprises a central linker flanked by chromo and chromo-shadow domains, the

latter of which interacts with the PxVxL motif in CAF1/p150.[41] As the PxIxL motif in

ZNF518B is crucial for interaction with HP1 isoforms, we predicted that ZNF518B

interacts with the chromo-shadow domain rather than the chromo domain of the HP1

isoforms. We examined the interactions between ZNF518B and the chromo (HP1β^{1-}

^{92}) and chromo-shadow (HP1β^{93-185}) domains of HP1β using the FMIT assay. As

expected, only the chromo-shadow domain was recruited to alphoid^{tetO} by ZNF518B

derivatives carrying the PxIxL motif in the C-terminus. This confirmed the interaction

between the PxIxL motif of ZNF518B and the chromo-shadow domain of HP1β (Fig.

3C).

ZNF518B$^{344-769}$ and ZNF518B$^{555-769}$ weakly bound HP1β despite the absence

of a PxIxL motif. The ZNF518B PxIxL motif had similar binding affinities for all three

HP1 isoforms. However, ZNF518A residues 971–1321, which correspond to ZNF518B

residues 555–769, did not show such specificity for the HP1 isoforms (Fig. S5B).

Therefore, the 555–769-aa segment in ZNF518B is a subsidiary segment for HP1

interactions that determine its specificity for HP1β.

ZNF518B maintains HP1 at alphoid^{tetO}

We next examined whether HP1β recruited to alphoid^{tetO} by TetR-EYFP-ZNF518B is

maintained at the alphoid^{tetO} repeat locus when ZNF518B is released from the

alphoid^{tetO} DNA following the addition of doxycycline (Dox) (Fig. 3D). TetR-EYFP-

ZNF518 was rapidly released from alphoid^{tetO} after the addition of Dox. EYFP foci

disappeared in 96% of cells 120 min after Dox addition (Fig. S5C). We quantified Halo-

HP1β foci at alphoid^{tetO} 150 min after Dox addition (Fig. 3E). The Halo-HP1β foci

disappeared from the alphoid^{tetO} DNA in all assays employing ZNF518B^{WT},

ZNF518B$^{878–987}$, and ZNF518B$^{770–987}$ (Fig. 3E). Therefore, ZNF518B is required to

maintain HP1β at the alphoid^{tetO}.

ZNF518B interacts directly with HP1 isoforms *in vitro*

The FMIT results strongly suggested that ZNF518B interacts with the HP1 isoforms,

corroborating previously published results regarding the interaction of ZNF518B with

the HP1 isoforms.[42] To further examine their direct interactions, we purified the

ZNF518B$^{770–987}$ derivative containing the PxIxL motif as a maltose-binding protein

(MBP) fusion protein and the HP1 isoforms as glutathione S-transferase (GST) fusion

proteins, respectively, using an *Escherichia coli* overexpression system. The two

purified proteins were mixed, and the MBP-fused ZNF518B$^{770–987}$ proteins were pulled

down using magnetic amylose beads. The GST-HP1 isoforms were co-purified with

ZNF518B$^{770–987}$ and were more abundant in pull-down (PD) samples with MBP-

ZNF518B$^{770–987}$ than in those with MBP alone (Fig. 3F), suggesting that the PxIxL motif

in the ZNF518B C-terminus directly interacts with HP1α, HP1β, and HP1γ (Fig. 3G).

ZNF518B localization cannot be maintained at centromeres without CENP-B

As CENP-B is required for centromeric localization of ZNF518B (Fig. 1), we next

investigated whether it is required to maintain ZNF518B at the centromeres. We

created a *CENPB* conditional KO (cKO) system using U2OS cells (Fig. S6A). The

Oryza sativa TIR1 gene, encoding an auxin receptor, was inserted into an adeno-

associated virus. In virus-infected cells, endogenous CENP-B fused with a mini auxin-

inducible degron (mAID) tag was rapidly targeted for degradation via the ubiquitin-

proteasome pathway following indole-3-acetic acid (IAA) treatment (Fig. S6A).[43]

The localization of GFP-ZNF518B was analyzed under three conditions (Fig.

4A): CENP-B:mAID available at all times (CENP-B^{ON}); CENP-B:mAID removed 24 h

before ZNF518B transfection (CENP-B^{OFF}); and CENP-B:mAID removed 4 h before

fixation (CENP-B^{4h-OFF}). As expected, GFP-ZNF518B localized at the centromeres in

CENP-B^{ON} cells (Fig. 4B–E). In CENP-B^{OFF} cells, CENP-B signals at the centromeres

were below the detection limit, and GFP-ZNF518B dissociated from the centromeres

(Fig. 4B–E). In CENP-B^{4h-OFF} cells, GFP-ZNF518B also dissociated from the

centromeres (Fig. 4B–E). These results demonstrated that the recruitment and

maintenance of ZNF518B at centromeres require CENP-B.

Interaction between ZNF518B and CENP-B

We further examined the interaction between ZNF518B and CENP-B using the FMIT

assay. For this, we constructed Halo-CENP-B$^{133–599}$, which is devoid of the helix-turn-

helix domain and thus cannot bind to α-satellites at centromeres.[17] Halo-CENP-B^{133-}

^{599} and TetR-EYFP-ZNF518B (WT and derivatives) were co-expressed in HeLa-03-Int

cells (Fig. 4F). As expected, Halo-CENP-B$^{133–599}$ did not localize to the centromeres.

Therefore, CENP-B$^{133–599}$ foci reflected the CENP-B recruitment to alphoid^{tetO} by TetR-

EYFP-ZNF518B.

CENP-B$^{133-599}$ was recruited to alphoid^{tetO} in 26% of cells expressing

ZNF518B^{WT}, but not in cells expressing TetR-EYFP and lacking ZNF518B. Moreover,

ZNF518B^{WT} was recruited to alphoid^{tetO} in cells expressing TetR:EYFP:CENP-B$^{133-599}$

(Fig. S6B). These results were consistent with the interaction between ZNF518B and

CENP-B (Fig. 4G). CENP-B$^{133-599}$ was barely recruited to alphoid^{tetO} by ZNF518B$^{1-398}$,

ZNF518B$^{1-628}$, and ZNF518B$^{344-628}$, suggesting that the ZNF518B segment from aa

629 to the C-terminus is important for the interaction. CENP-B$^{133-599}$ was recruited to

alphoid^{tetO} at high frequencies in cells expressing ZNF518B$^{344-1074}$, ZNF518B$^{344-987}$,

ZNF518B$^{344-877}$, ZNF518B$^{555-1074}$, ZNF518B$^{555-877}$, ZNF518B$^{770-987}$, and ZNF518B$^{770-877}$,

^{877}, all of which contain residues 770–877 (Fig. 4G). The recruitment frequency of

CENP-B$^{133-599}$ was reduced by 20% in cells expressing ZNF518$^{344-769}$ versus those

expressing ZNF518$^{344-877}$ and by 39% in cells expressing ZNF518$^{555-769}$ versus those

expressing ZNF518$^{555-877}$. ZNF518B$^{878-1074}$ and ZNF518B$^{878-987}$ did not exhibit CENP-

B$^{133-599}$ recruitment. These results support the importance of residues 770–877 of

ZNF518B for interaction with CENP-B (Fig. 4H).

We next examined the recruitment of CENP-B to alphoid^{tetO} in cells expressing

ZNF518B$^{KSS/N}$, ZNF518B$^{KSS/C}$, ZNF518B$^{KSS/NC}$, and ZNF518B$^{770-987\ KSS/C}$ and carrying

substitutions of the PxV(I)xL motifs with KxSxS (Fig. 4G). The PxV(I)xL substitutions

generally increased CENP-B recruitment to the *tetO* repeats, implying an inhibitory

role of the PxV(I)xL motifs in the interaction between ZNF518B and CENP-B. The

derivatives with the N-terminal region deleted showed considerably stronger

interactions with both HP1 and CENP-B than full-length ZNF518B (Figs. 3C and 4G).

Therefore, we speculate that the interaction domain in some ZNF518 segments is

exposed on the protein surface, allowing for easier interactions with the binding partner.

The C-terminus of ZNF518B interacts with G9a

Proteomics data have revealed an interaction between ZNF518B and G9a (EHMT2).[32]

We attempted to identify the ZNF518B segments that interact with G9a and vice versa

using the FMIT assay with cells co-expressing EYFP-TetR-ZNF518 and Halo-G9a.

G9a was recruited to *tetO* repeats in control cells expressing ZNF518B^{WT} (Fig. S6D

and 5A). G9a was recruited to alphoid*^{tetO}* at high frequencies in cells expressing

ZNF518B$^{KSS/C}$, ZNF518B$^{KSS/NC}$, ZNF518B$^{344-1074}$, ZNF518B$^{555-1074}$, and ZNF518B^{878-}

^{1074}, all of which carried the C-terminus, but barely in cells expressing the other

ZNF518B derivatives (Fig. 5A). Therefore, the C-terminus (residues 988–1074) of

ZNF518B is crucial for interaction with G9a.

We also examined the interaction between ZNF518A and G9a using the FMIT

assay. G9a was recruited to alphoid*^{tetO}* in cells expressing ZNF518A^{WT}, ZNF518A^{KSS},

ZNF518A$^{971-1483}$, ZNF518A$^{971-1483\ KSS}$, ZNF518A$^{1317-1487}$, and ZNF518A$^{1317-1483\ KSS}$, all

of which included the C-terminus, but less efficiently in cells expressing ZNF518A$^{1-970}$

and ZNF518A$^{971-1321}$ (Fig. 5B). These results suggested that the C-terminus (residues

1317–1483) of ZNF518A is important for interaction with G9a.

The segment downstream of the TAD in G9a interacts with ZNF518A/B

G9a and the C2H2-type zinc-finger motifs in ZNF644 and ZNF803, known as WIZ-

containing proteins, can interact.[31,44] The zinc-finger motifs in WIZ-containing proteins

also interact with the transcriptional regulatory CCAAT/enhancer-binding protein-β

transactivation domain (TAD) in the N-termini of G9a and GLP.[45] Because zinc-finger

motifs that interact with G9a are highly conserved in the ZNF518A/B C-termini in all

species, except *Xenopus laevis* (Fig. S6E), we hypothesized that the C-termini of

ZNF518A/B may interact with the G9a TAD.

To test this hypothesis, we conducted FMIT assays with several G9a

derivatives (Fig. 5C). In contrast to our hypothesis, G9a$^{1-290}$ was not recruited to

alphoid^{tetO} by ZNF518A/B, whereas G9a$^{291-1210}$, lacking the TAD, was. Moreover, two

291–545-aa segment-containing G9a derivatives, G9a$^{1-545}$ and G9a$^{291-545}$, were

recruited to alphoid^{tetO} by ZNF518A/B. These results indicated that residues 291–545

of G9a, not the TAD, are sufficient for interactions with ZNF518A/B.

ZNF518B recruits G9a H3K9 methyltransferase to centromeres

Finally, we assessed whether ZNF518 recruits G9a to centromeres. Plasmids

expressing GFP-ZNF518B or ZNF518B derivatives and Halo-G9a were co-transfected

into U2OS cells, and the localization of Halo-G9a was investigated using the FMIT

assay (Fig. 5D). Halo-G9a localized to the centromeres only when expressed with WT

or ZNF518B$^{344-1074}$ carrying the C-terminus, not when expressed with ZNF518B$^{344-987}$

lacking the C-terminal zinc-finger (Fig. 5D). Halo-G9a did not localize to the

centromeres when GFP-ZNF518B and Halo-G9a were expressed in *CENPB*-KO cells

(Fig. 5E). Therefore, CENP-B recruits ZNF518B to centromeres, and ZNF518B in turn

recruits G9a H3K9 methyltransferase (Fig. 5F).

Discussion

This study showed that the 770–877-aa segment in ZNF518B interacts with CENP-B and that CENP-B recruits ZNF518B to centromeres and pericentromeres. Furthermore, the 555–765 segment and PxIxL motif in ZNF518B are required for interaction with the chromo-shadow domain of HP1β. The pericentromere forms constitutive heterochromatin, promoting HP1 accumulation.[8,9,11] Therefore, among the ZNF518 proteins recruited to the centromere/pericentromere by CENP-B, those assembled in the pericentromere are likely to maintain a stable association with the chromatin through interaction with HP1. The ZNF518B C-terminus carrying the zinc-finger motif interacts with the segment downstream of the TAD in G9a, and CENP-B is required for the centromeric localization of G9a. Based on these findings, we propose the following mechanisms: 1) CENP-B binds to α-satellite DNA to initiate heterochromatin formation; 2) CENP-B recruits ZNF518B to centromeres and pericentromeres; 3) HP1β stabilizes ZNF518 only at pericentromeres; and 4) ZNF518B interacts with G9a for H3K9 methylation at the pericentromeres.

These conclusions do not disagree with a previously proposed mechanism involving SUV39H1 histone H3K9 methyltransferase.[26,27] HP1-dependent SUV39H1 recruitment and short RNA-mediated heterochromatinization are important for pericentromeric heterochromatin maintenance.[26,28] We suggest that G9a recruited by CENP-B and ZNF518B is required for pericentromeric heterochromatin establishment and that the CENP-B/ZNF518B/HP1β/G9a axis targets heterochromatin factors to

pericentromeres, where specificity is achieved by the binding of CENP-B to CENP-B box sequences in pericentromeric alphoid repeats.

ZNF518A is also recruited to pericentromeres in a CENP-B-dependent manner (Fig. S1) and interacts with HP1 and G9a (Figs. 5, S5). ZNF518A and ZNF518B are evolutionarily conserved in all vertebrates higher than amphibians. Therefore, we speculate that pericentromeric heterochromatin formation by ZNF518 is conserved among many vertebrates. Although further investigation is needed, our study provided insights into the potential mechanisms underlying the targeting of heterochromatin to human pericentromeres and indicated that a similar mechanism may exist in many species.

Material and Methods

DNA constructs

DNA fragments encoding human ZNF518A, ZNF518B, CENP-B$^{133-599}$, HP1α, HP1β,

HP1γ, and G9a were PCR-amplified (primers are listed in Table S1) from a cDNA

library generated by the Kazusa ORFeome Project and Genome Network Project

Clones by RIKEN BRC. The fragments were inserted into an entry vector (pENTR™

4 Dual Selection Vector; #A10465, Thermo Fisher Scientific) between the *Xmn*I

(#R0194S, New England BioLabs [NEB]) and *EcoR*V (#R0195S, NEB) sites using the

In-Fusion HD Cloning Kit (#Z9648N, Takara Bio), yielding entry plasmids

pENTR4XE153, pENTR4XE127, pENTR4ΔCENPB, pENTR4HP1α, pENTR4HP1β,

pENTR4HP1γ, and pENTR4G9a, which comprised the open reading frames encoding

the above respective proteins.[46–50]

DNA fragments encoding ZNF518A derivatives (ZNF518A^{KSS}, ZNF518A$^{1-970}$,

ZNF518A$^{971-1483}$, ZNF518A$^{971-1483KSS}$, ZNF518A$^{971-1321}$, ZNF518A$^{1317-1483}$, and

ZNF518A$^{1317-1483KSS}$), ZNF518B derivatives (ZNF518B$^{KSS/N}$, ZNF518B$^{KSS/C}$,

ZNF518B$^{KSS/NC}$, ZNF518B$^{1-398}$, ZNF518B$^{1-628}$, ZNF518B$^{344-628}$, ZNF518B$^{344-1074}$,

ZNF518B$^{344-987}$, ZNF518B$^{344-877}$, ZNF518B$^{344-769}$, ZNF518B$^{555-985}$, ZNF518B$^{555-877}$,

ZNF518B$^{555-769}$, ZNF518B^{878v1074}, ZNF518B$^{878-987}$, ZNF518B$^{770-987}$, ZNF518B^{770-}

$^{987KSS/C}$, and ZNF518B$^{770-877}$), HP1β derivatives (HP1$\beta^{N(1-92)}$ and HP1$\beta^{C(93-185)}$), and

G9a derivatives (G9a$^{1-92}$ and G9a$^{93-185}$) were PCR-amplified from pENTR4XN153,

pENTR4XN127, pENTR4HP1β, and pENTR4G9a, respectively, using the primers

listed in Table S1.

To overexpress the target proteins in human or *E. coli* cells, plasmids were

generated by DNA recombination between the above-mentioned entry plasmids and

pDEST131NGFP (for EGFP fusion proteins),[51] pDEST152JTY (for EYFP-TetR fusion

proteins), pDEST155Halo (for Halo-tagged proteins), pDEST174HATMBP (for HAT

[KDHLIHNVHKEEHAHNK]-MBP fusion proteins),[52] or pDEST172HGST (for His_6-GST

fusion proteins) using Gateway LR Clonase II Enzyme mix (#11791020, Thermo

Fisher Scientific)*.

For CRISPR-mediated double-strand break site formation, 54-bp DNA

fragments were generated by annealing two complementary single-stranded

oligonucleotides. The fragments were inserted into the *Age*I (#R3552S, NEB) site of

pTORA14 using the In-Fusion HD Cloning Kit,[53] yielding pTORA14AAVS1,

pTORA14CENP-B_C, pTORA14HA4, pTORA14HB7, and pTORA14HG1, which were

used to induce DNA cleavage at adeno-associated virus integration site 1 in *CENPB,*

HP1ALPHA, HP1BETA, and *HP1GAMMA*, respectively. The DNA sequences targeted

by the guide RNAs are listed in Table S2.

To construct the plasmid directing homologous recombination (HR) at the 3′

end of *CENPB*, a DNA fragment was PCR-amplified from human genomic DNA using

Tks Gflex DNA Polymerase (#R060A, Takara Bio) and the primers listed in Table S1.

This yielded a 584-bp DNA fragment containing the 267-bp left homological arm (HRL)

and 314-bp right homological arm (HRR) regions identical to the target locus. The

amplified DNA fragment comprising the HRL and HRR regions was cloned into pCR-

Blant (#K275020, Thermo Fisher Scientific). To construct HR donor plasmids

(pKANA1CENPB and pKANA2CENPB), mAID cassettes containing the mAID-

encoding region and a hygromycin or blasticidin resistance gene were PCR-amplified

from pMK287 or pMK288 and inserted between the HRL and HRR regions using the

In-Fusion HD Cloning Kit.[43] The resultant plasmids were used to generate a human

cell line expressing the CENP-B:mAID fusion protein.

Cell lines

HeLa-Int-03 cells containing approximately 15,000 repeats of the *tetO* sequence in

chromosome 21 with and without *CENPB* or *HP1* KO were established as previously

described.[17,37,54] A U2OS-*os*TIR1 cell line, which is the parental cell line of *CENPB-*

cKO cells, was generated by transfecting U2OS cells with pTORA14AAVS1 and

pMK243 using Lipofectamine LTX Reagent (#A12621, Thermo Fisher Scientific)[43]

followed by puromycin selection (1 µg/mL, #10-2100, Focus Biomolecules). *CENPB-*

cKO cells were generated by transfecting the U2OS-*os*TIR1 cells with

pTORA14CENP-B_C, pKANA1CENPB, and pKANA2CENPB followed by selection on

hygromycin B (1 mg/mL, #089-06151, Wako) and blasticidin S (5 µg/mL, #ant-bl-05,

InvivoGen).

Cell culture

U2OS and HeLa cells were cultured in Dulbecco's modified Eagle's medium (DMEM;

#044-29765, FUJIFILM Wako) supplemented with 10% (v/v) fetal bovine serum

(Bovogen Biologicals), 100 U/mL of penicillin, and 100 µg/mL of streptomycin (#168-

23191, Wako) at 37 °C in a humidified incubator with 5% CO_2. To generate CENP-

B[OFF] cells, *CENPB*-cKO cells were cultured with 0.2 µg/mL Dox (#D3447, Sigma-

Aldrich) for 24 h and subsequently with 0.5 mM IAA (#094-07121, Wako) for 24 h

before use.

Indirect immunofluorescence and microscopy

U2OS and HeLa cells in the exponential growth phase were seeded onto coverslips

at 30–40% confluence and allowed to grow overnight. To overexpress EGFP-

ZNF518A and EGFP-ZNF518B for indirect immunofluorescence experiments,

pDEST131NEGFP153 and pDEST131NEGFP127, respectively, were transfected into

the cells using Lipofectamine LTX. Twenty-four hours later, the cells were fixed with

4% (v/v) paraformaldehyde (#163-20145, Wako) in phosphate-buffered saline (PBS)

for 10 min and permeabilized with 0.15% (v/v) Triton X-100 (#1610407, Bio-Rad) in

PBS for 2 min. The cells were treated with 1% (v/v) bovine serum albumin (BSA;

#A7284, Sigma-Aldrich) in PBS and subsequently incubated with primary antibodies,

including mouse anti-CENP-A (1:200 dilution, #ab13939, Abcam), mouse anti-CENP-

B (1:50,000, 5E6C1),[16] rabbit anti-CENP-B (1:200, #ab25734, Abcam), 1:600-diluted

rabbit anti-CENP-C (#554),[55] mouse anti-HP1α (1:1,000, #MAB3584, Merck Millipore),

rabbit anti-HP1β (1:800, #8676 D2F2, Cell Signaling Technology), mouse anti-HP1γ

(1:3,000, #MAB3450 2MOD-1G6, Merck Millipore), or rabbit anti-histone H3K9me3

(1:1,000, #ab8898, Abcam). The cells were washed with PBS thrice for 5 min and

incubated with Alexa-conjugated secondary antibodies (1:600). DNA was stained with

0.1 µg/mL DAPI (#H-1200, Vector Laboratories).

The cells were observed using an Olympus FV1000 confocal microscope with

a UPlanSApo 60×/1.35 oil-immersion objective lens (Olympus) and FV10-SAW2.1

software (Olympus). Images were also acquired using a Nikon Confocal A1Rsi

microscope with a CFI Plan Apochromat VC 60×/1.40 oil-immersion objective lens

(Nikon) and FNIS-Elements AR software (Nikon). The images were acquired at 0.5-

µm intervals in the z-axis and Kalman-filtered to suppress background noise.

FMIT

pDEST172JTY and pDEST155Halo encoding EYFP-TetR and Halo fusion proteins,

respectively, were transfected into HeLa cells carrying the *tetO* array at 60%

confluence using Lipofectamine 3000 (#L3000001, Thermo Fisher Scientific). Twenty-

four hours later, the cells were incubated with 1 nM Halo Tag TMR Ligand (G825A,

Promega) at 37 °C for 2 h. The labeled cells were washed with DMEM and incubated

in DMEM at 37 °C for 30 min. Then, the cells were fixed with 4% (v/v)

paraformaldehyde in PBS for 10 min and DNA was stained with 0.1 µg/mL DAPI.

EYFP and TMR signals were observed using an Olympus BX53 fluorescence

microscope with a UPlanSApo 60×/1.35 oil-immersion objective lens and cellSens

software (Olympus). More than 100 cells were analyzed for each sample.

Quantification of microscopic data

Fluorescence quantitation and line plot analyses of the fluorescence signals were

performed using ImageJ 1.50e.[56] To quantitate fluorescence, z-stack images were

captured at every 500 nm between the lower and upper edges of nuclei and merged

using the maximum intensity z-projection function in ImageJ. In projected images,

nuclei and centromeres were defined by DAPI and CENP-A signals, respectively.

Nucleus- and centromere-associated GFP signals in the cells were estimated, and

their ratio was calculated to determine whether GFP-fused proteins were enriched at

centromeres. Statistical analyses were performed using R v.3.5.1 (R Foundation for

Statistical Computing, http://www.r-project.org).

For the line plot analyses, the slice showing the strongest signal of a

centromeric protein (CENP-A or CENP-B) and GFP-ZNF518 was selected from the z-

stack images. Using the RGB plot profile function in ImageJ, signal intensities of the

centromeric protein and GFP-ZNF518 were quantified for each pixel along the

indicated lines.

Protein purification

His$_6$-GST-HP1 and HAT-MBP-ZNF518B$^{770-987}$ were purified from BL21 *E. coli* cells.

To express the target proteins, *E. coli* cells carrying pDEST173HGSTHP1 for His$_6$-

GST-HP1 or pDEST174HATMBP127 for HAT-MBP-ZNF518B$^{770-987}$ were cultured in

L broth supplemented with 0.1 mM IPTG (#096-05143, Wako) at 30 °C for 2 h. His$_6$-

GST-HP1 proteins were extracted using a buffer containing 50 mM Tris-HCl (pH 8.0;

#201-06273, Wako), 500 mM NaCl (#191-01665, Wako), 10 mM β-mercaptoethanol

(#131-14571, Wako), and 10% glycerol (#075-00611, Wako). HAT-MBP-ZNF518B^{770-}

^{987} was extracted using above buffer including 1% Tween 20 (#H5152, Promega). The

target proteins were affinity-purified using Ni-NTA agarose (#143-09763, Wako).

MBP PD

All MBP PD procedures were performed using PD buffer comprising 25 mM Tris-HCl

(pH 8.0), 100 mM NaCl, 1 mM EDTA (#E5134, Sigma-Aldrich), 1 µM DTT (#GE17-

1318-02, Sigma-Aldrich), 1% Tween 20, 10% glycerol, and 0.02% BSA. First, 2 µg

HAT-MBP-ZNF518B$^{770-987}$ protein and 5 µg His$_6$-GST-HP1 protein, purified as

described above, were mixed and incubated under rotation at 4 °C for 4 h. Next, 50

µg amylose magnetic beads (#E8035S, NEB) was added to the protein mixture, which

was further incubated under rotation at 4 °C for 1 h. After washing the beads six times

with 1 mL of PD buffer, the sample was boiled with 1× SDS loading buffer at 95 °C for

5 min. The eluate was subjected to immunoblotting.

Immunoblotting

Immunoblotting was carried out using mouse anti-MBP (1:200, #sc-271524, Santa

Cruz Biotechnology), mouse anti-GST (1:200, #sc-138, Santa Cruz Biotechnology),

rabbit anti-HP1α (1:1,000, #2616, CST), mouse anti-HP1α (#MAB3584, Merck

Millipore), rabbit anti-HP1β (1:1,000,#8676, CST), rabbit anti-HP1γ (1:1,000,#2619,

CST), mouse anti-HP1γ (1:1,000, #MAB3450, Merck Millipore), and mouse anti-

GAPDH (1:10,000, #5A12,016-25523, Wako) primary antibodies. The target proteins

were detected using IRDye 800CW donkey anti-rabbit IgG and anti-mouse IgG

(1:15,000, #926-32213, #926-32210, Li-COR Biosciences) secondary antibodies.

Amino acid sequence analysis

The Basic Local Alignment Search Tool was used to identify amino acid similarities

between query sequences and sequences deposited in DDBJ/GenBank/EMBL.[57] A

neighbor-joining phylogenetic tree was constructed using MEGA X software.[58]

Acknowledgments

The authors thank Iyo Toramoto, Kana Yamazoe, and Fusako Kawasaki (Kochi University) for technical assistance; Koichiro Otake (Kazusa DNA Research Institute) for providing preliminary data; Koichi Honke, Hideaki Kuge, Kaoru Miyahara, Arisa Miyagawa-Yamaguchi, and Tatsuyuki Yamashita (Kochi University) for scientific advice; and William Earnshaw (University of Edinburgh) for critical manuscript reading. This work was supported by grants from JSPS KAKENHI (grant No. 18K06061), the Kato Memorial Bioscience Foundation, and the Takeda Science Foundation to SO, by grants from JST, CREST (grant No. 18070874), and MEXT KAKENHI (grant No. 20H00473), and by a grant from the Kazusa DNA Research Institute Foundation to HM.

Author contributions

SO, JO, and HM designed the study. SO, JO, and NS performed the experiments. SO, JO, and HM analyzed the data. SO, JO, KN, and HM wrote the manuscript.

Disclosure and competing interests statement

The authors declare no conflicts of interest.

Figure legends

Figure 1. ZNF518B and ZNF518A localize to centromeres in a CENP-B-

dependent manner.

(A, C, and E) Co-localization of GFP-ZNF518B (green) and the centromeric factors

(red) CENP-A **(A)**, CENP-B **(C)**, and CENP-C **(E)**. Areas marked by dashed boxes in

the left images are magnified in the right panel. Scale bars, 5 μm.

(B, D, and F) Line plot analyses along the white lines in panels A, C, and E for

quantification of GFP-ZNF518B (green) and the centromeric factors (red) CENP-A **(B)**

CENP-B **(D)**, and CENP-C **(F)**.

(G) Localization of GFP-ZNF518B (green) and CENP-A (red) in WT (upper) and

CENPB-KO cells (lower). Areas marked by dashed boxes in the left panels are

magnified in the right panels.

(H) Quantification of the fluorescence intensities of GFP-ZNF518B signals at

centromeres in WT, *CENPB*-KO [see panel **(G)**], and *CENPB*-KO cells overexpressing

CENP-B-Flag$_3$ [*CENPB*-KO + CENP-B-FL$_3$ OX; see panel **(I)**]. The y-axis shows the

relative fluorescence intensity of GFP-ZNF518B signals associated with centromeres

to total GFP signals in the nucleus. Circles represent measurements from the

individual cells shown in panels G and I. Red bars indicate medians. ****p* < 0.005.

(I) Co-localization of GFP-ZNF518B (green), CENP-A (red), and CENP-B (blue) in

CENPB-KO cells overexpressing CENP-B-Flag$_3$ (*CENPB*-KO + CENP-B-FL$_3$ OX).

The area marked by the dashed box in the left panel is magnified in the right panel.

(J) Co-localization of Halo-ZNF518A (red), GFP-ZNF518B (green), and CENP-A

(blue). The area marked by the dashed box in the left panel is magnified in the right

panel.

Figure 2. Identification of ZNF518B segments required for centromeric

localization.

Localization of the ZNF518B derivatives at centromeres. The schematic

representation of the respective ZNF518B derivatives is drawn approximately to scale.

Yellow boxes indicate zinc-finger motifs. The two PxV(I)xL motifs (purple boxes) were

substituted with KxSxS (light purple boxes), shown as KSS/N and KSS/C. KSS/NC

indicates that both PxV(I)xLs motifs were substituted with KxSxS. GFP-fused

ZNF518B signals (WT and 18 derivatives; green) were co-visualized with the

centromeric marker CENP-A or CENP-B (left images). The two images

(ZNF518B$^{KSS/NC}$ and ZNF518B$^{878-987}$) labeled with "A" show CENP-A foci, all other

images show CENP-B signals. Areas marked by dashed boxes in the left images are

magnified in the right panel. Numbers and colors (left) reflect % frequencies of cells

with GFP-ZNF518B signals at pericentromeres. Scale bar, 5 µm.

Figure 3. The PxIxL motif in ZNF518B is required for interaction with the chromo-

shadow domain of HP1.

(A) Schematic diagram of the FMIT assay for investigating the interaction between

ZNF518B and HP1. *CENPB*-KO cells were used.

(B) Tethering of YFP-ZNF518B (green) to the *tetO* repeats results in the accumulation of Halo-HP1β (top; red), but not Halo-HP1γ (bottom; red) at the locus.

(C) FMIT assay to examine the interaction between ZNF518B derivatives and HP1. Cells expressing TetR-EYFP-ZNF518B (WT or derivatives) and Halo-HP1 were used. The ZNF518B segment predicted to interact with HP1 is indicated by shaded boxes. Numbers and colors reflect % frequencies of the co-localization of TetR-EYFP-ZNF518B with full-length HP1 (HP1α, HP1β, and HP1γ) and the chromo domain (CD) and chromo-shadow domain (CSD) of HP1β.

(D) Experimental design to examine whether ZNF518B is required to maintain HP1β localization at the *tetO* repeats. TetR-EYFP-ZNF518 quickly dissociated from the *tetO* repeats in the presence of Dox. *CENPB* KO cells were used in this assay.

(E) Cells carrying TetR-EYFP-ZNF518 (WT or derivatives) and Halo-HP1β were subjected to the FMIT assay as explained in panel D.

(F) Pull-down (PD) experiments to examine the physical interaction between ZNF518B and HP1. Purified recombinant GST-HP1 (input) was incubated with MBP-ZNF518B$^{770-987}$, and MBP PD samples were subjected to immunoblotting using anti-GST. Signal intensities in the indicated samples were normalized to those in samples with MBP alone.

(G) Model depicting the central roles of ZNF518A and ZNF518B in pericentromeric heterochromatin formation via their interactions with CENP-B, HP1, and G9a. The red ellipse and arrow highlight the reaction step focused on in this figure.

Figure 4. Dissection of the interaction of ZNF518B with CENP-B.

(A) Schematic representation of the GFP-ZNF518B localization assay, in which

indole-3-acetic acid (IAA) was added at different time points to deplete CENP-B. Data

are shown in panel B.

(B) Representative images of GFP-ZNF518B-transfected cells quantified in panels C

and D**.** Localization of GFP-ZNF518B (green) relative to centromeres in CENP-B^{ON},

CENP-B^{OFF}, and CENP-B$^{4h\text{-}OFF}$ cells. CENP-A (red), CENP-B (blue), and DAPI

staining are shown. Areas marked by dashed boxes in the left images are magnified

in the right panel.

(C) Quantification of the fluorescence intensities of nuclear CENP-B signals in CENP-

B^{ON}, CENP-B^{OFF}, and CENP-B$^{4h\text{-}OFF}$ cells based on the images in panel B. ***$p < 0.005$,

red bars indicate medians.

(D) Quantification of the fluorescence intensities of GFP-ZNF518B signals at the

pericentromeres in CENP-B^{ON}, CENP-B^{OFF}, and CENP-B$^{4h\text{-}OFF}$ cells based on the

images shown in panel B. The y-axis shows the relative fluorescence intensity of GFP-

ZNF518B signals associated with pericentromeres to total GFP signals in the nucleus.

Circles represent measurements from the individual cells shown in panel B, red bars

indicate medians. **$p < 0.05$.

(E) Percentage of GFP-ZNF518B foci co-localized with centromeres (CENP-A foci) in

CENP-B^{ON}, CENP-B^{OFF}, and CENP-B$^{4h\text{-}OFF}$ cells based on the images in panel B.

(F) Schematic diagram of the FMIT assay for investigating the interaction between

ZNF518B and CENP-B. The CENP-B$^{133-599}$ derivative was used as it cannot bind to

the α-satellite repeats at pericentromeres by itself.

(G) FMIT assay to quantify the co-localization between TetR-EYFP-ZNF518 and Halo-CENP-B$^{133–599}$. Numbers and colors reflect % frequencies of Halo-CENP-B$^{133–599}$ recruitment onto TetR-EYFP-ZNF518B foci. Red shaded boxes represent the ZNF518B segment predicted to be involved in the interaction with CENP-B$^{133–599}$.

(H) Model depicting the central roles of ZNF518A and ZNF518B in pericentromeric heterochromatin formation via their interactions with CENP-B, HP1, and G9a. The red ellipse and arrow highlight the reaction step focused on in the present figure.

Figure 5. Dissection of ZNF518A and ZNF518B in terms of their requirement for the loading of G9a to centromeres.

(A) FMIT assay to quantify the co-localization between TetR-EYFP-ZNF518 (WT and 16 derivatives) and Halo-G9a. Green shaded boxes represent the ZNF518B segment potentially involved in its interaction with G9a.

(B) FMIT assay to quantify the co-localization between TetR-EYFP-ZNF518A (WT and seven derivatives) and Halo-G9a. Green shaded boxes represent the ZNF518A segment potentially involved in its interaction with G9a.

(C) FMIT assay to quantify the co-localization between Halo-G9a (WT and four derivatives) and TetR-EYFP-ZNF518A or TetR-EYFP-ZNF518B.

(D) Co-visualization of Halo-G9a (red) and GFP-ZNF518B (WT and two derivatives) or GFP alone (green) in WT cells. Areas marked by dashed boxes in the left images are magnified in the right panel. Scale bar, 5 μm.

(E) Localization of Halo-G9a (red) and GFP-ZNF518B^{WT} (green) in *CENPB*-KO cells. Scale bar, 5 μm.

1   **(F)** Model depicting the central roles of ZNF518A and ZNF518B in pericentromeric

2   heterochromatin formation via their interactions with CENP-B, HP1, and G9a (see

3   Discussion for details).

REFERENCES

1. Molina, O., Abad, M.A., Solé, F., and Menéndez, P. (2021). Aneuploidy in cancer:

lessons from acute lymphoblastic leukemia. Trends Cancer *7*, 37–47.

10.1016/j.trecan.2020.08.008.

2. Barra, V., and Fachinetti, D. (2018). The dark side of centromeres: types, causes

and consequences of structural abnormalities implicating centromeric DNA. Nat

Commun *9*, 4340. 10.1038/s41467-018-06545-y.

3. Cheeseman, I.M., and Desai, A. (2008). Molecular architecture of the kinetochore-

microtubule interface. Nat. Rev. Mol Cell Biol *9*, 33–46. 10.1038/nrm2310.

4. McKinley, K.L., and Cheeseman, I.M. (2016). The molecular basis for centromere

identity and function. Nat Rev Mol Cell Bio *17*, 16–29. 10.1038/nrm.2015.5.

5. Amor, D.J., Kalitsis, P., Sumer, H., and Choo, K.H.A. (2004). Building the

centromere: from foundation proteins to 3D organization. Trends Cell Biol *14*, 359–

368. 10.1016/j.tcb.2004.05.009.

6. Waye, J.S., and Willard, H.F. (1987). Nucleotide sequence heterogeneity of alpha

satellite repetitive DNA: a survey of alphoid sequences from different human

chromosomes. Nucleic Acids Res *15*, 7549–7569. 10.1093/nar/15.18.7549.

7. Schueler, M.G., Higgins, A.W., Rudd, M.K., Gustashaw, K., and Willard, H.F.

(2001). Genomic and genetic definition of a functional human centromere. Science

294, 109–115. 10.1126/science.1065042.

8. Schueler, M.G., and Sullivan, B.A. (2006). Structural and functional dynamics of

human centromeric chromatin. Annu Rev Genom Hum *7*, 301–313.

10.1146/annurev.genom.7.080505.115613.

9. Black, B.E., and Cleveland, D.W. (2011). Epigenetic centromere propagation and the nature of CENP-A nucleosomes. Cell *144*, 471–479. 10.1016/j.cell.2011.02.002.

10. Cleveland, D.W., Mao, Y., and Sullivan, K.F. (2003). Centromeres and kinetochores from epigenetics to mitotic checkpoint signaling. Cell *112*, 407–421. 10.1016/s0092-8674(03)00115-6.

11. Sullivan, B.A., and Karpen, G.H. (2004). Centromeric chromatin exhibits a histone modification pattern that is distinct from both euchromatin and heterochromatin. Nat Struct Mol Biol *11*, 1076–1083. 10.1038/nsmb845.

12. Masumoto, H., Nakano, M., and Ohzeki, J. (2004). The role of CENP-B and α-satellite DNA: de novo assembly and epigenetic maintenance of human centromeres. Chromosome Res *12*, 543–556. 10.1023/b:chro.0000036593.72788.99.

13. Earnshaw, W.C., Sullivan, K.F., Machlin, P.S., Cooke, C.A., Kaiser, D.A., Pollard, T.D., Rothfield, N.F., and Cleveland, D.W. (1987). Molecular cloning of cDNA for CENP-B, the major human centromere autoantigen. J Cell Biol *104*, 817–829. 10.1083/jcb.104.4.817.

14. Ohzeki, J., Nakano, M., Okada, T., and Masumoto, H. (2002). CENP-B box is required for de novo centromere chromatin assembly on human alphoid DNA. J Cell Biol *159*, 765–775. 10.1083/jcb.200207112.

15. Okamoto, Y., Nakano, M., Ohzeki, J., Larionov, V., and Masumoto, H. (2007). A minimal CENP-A core is required for nucleation and maintenance of a functional human centromere. EMBO J *26*, 1279–1291. 10.1038/sj.emboj.7601584.

16. Okada, T., Ohzeki, J., Nakano, M., Yoda, K., Brinkley, W.R., Larionov, V., and

Masumoto, H. (2007). CENP-B controls centromere formation depending on the

chromatin context. Cell *131*, 1287–1300. 10.1016/j.cell.2007.10.045.

17. Otake, K., Ohzeki, J., Shono, N., Kugou, K., Okazaki, K., Nagase, T.,

Yamakawa, H., Kouprina, N., Larionov, V., Kimura, H., et al. (2020). CENP-B creates

alternative epigenetic chromatin states permissive for CENP-A or heterochromatin

assembly. J Cell Sci *133*, jcs243303. 10.1242/jcs.243303.

18. Peters, A.H.F.M., O'Carroll, D., Scherthan, H., Mechtler, K., Sauer, S., Schöfer,

C., Weipoltshammer, K., Pagani, M., Lachner, M., Kohlmaier, A., et al. (2001). Loss

of the Suv39h histone methyltransferases impairs mammalian heterochromatin and

genome stability. Cell *107*, 323–337. 10.1016/s0092-8674(01)00542-6.

19. Grewal, S.I.S., and Jia, S. (2007). Heterochromatin revisited. Nat Rev Genet *8*,

35–46. 10.1038/nrg2008.

20. Molina, O., Vargiu, G., Abad, M.A., Zhiteneva, A., Jeyaprakash, A.A., Masumoto,

H., Kouprina, N., Larionov, V., and Earnshaw, W.C. (2016). Epigenetic engineering

reveals a balance between histone modifications and transcription in kinetochore

maintenance. Nat Commun *7*, 13334. 10.1038/ncomms13334.

21. Du, J., and Patel, D.J. (2014). Structural biology-based insights into

combinatorial readout and crosstalk among epigenetic marks. Biochim Biophys Acta-

Gene Regul Mech *1839*, 719–727. 10.1016/j.bbagrm.2014.04.011.

22. Hyun, K., Jeon, J., Park, K., and Kim, J. (2017). Writing, erasing and reading

histone lysine methylations. Exp Mol Med *49*, e324–e324. 10.1038/emm.2017.11.

23. Martin, C., and Zhang, Y. (2005). The diverse functions of histone lysine

methylation. Nat Rev Mol Cell Bio *6*, 838–849. 10.1038/nrm1761.

24. Canzio, D., Chang, E.Y., Shankar, S., Kuchenbecker, K.M., Simon, M.D.,

Madhani, H.D., Narlikar, G.J., and Al-Sady, B. (2011). Chromodomain-mediated

oligomerization of HP1 suggests a nucleosome-bridging mechanism for

heterochromatin assembly. Mol Cell *41*, 67–81. 10.1016/j.molcel.2010.12.016.

25. Rea, S., Eisenhaber, F., O'Carroll, D., Strahl, B.D., Sun, Z.-W., Schmid, M.,

Opravil, S., Mechtler, K., Ponting, C.P., Allis, C.D., et al. (2000). Regulation of

chromatin structure by site-specific histone H3 methyltransferases. Nature *406*, 593–

599. 10.1038/35020506.

26. Aagaard, L., Laible, G., Selenko, P., Schmid, M., Dorn, R., Schotta, G., Kuhfittig,

S., Wolf, A., Lebersorger, A., Singh, P.B., et al. (1999). Functional mammalian

homologues of the Drosophila PEV-modifier Su(var)3-9 encode centromere-

associated proteins which complex with the heterochromatin component M31.

EMBO J *18*, 1923–1938. 10.1093/emboj/18.7.1923.

27. Muramatsu, D., Kimura, H., Kotoshiba, K., Tachibana, M., and Shinkai, Y.

(2016). Pericentric H3K9me3 formation by HP1 interaction-defective histone

methyltransferase Suv39h1. Cell Struct Funct *41*, 145–152. 10.1247/csf.16013.

28. Shirai, A., Kawaguchi, T., Shimojo, H., Muramatsu, D., Ishida-Yonetani, M.,

Nishimura, Y., Kimura, H., Nakayama, J.-I., and Shinkai, Y. (2017). Impact of nucleic

acid and methylated H3K9 binding activities of Suv39h1 on its heterochromatin

assembly. eLife *6,* e25317. 10.7554/eLife.25317.

29. Tachibana, M., Sugimoto, K., Fukushima, T., and Shinkai, Y. (2001). SET

domain-containing protein, G9a, is a novel lysine-preferring mammalian histone

methyltransferase with hyperactivity and specific selectivity to lysines 9 and 27 of

histone H3. J Biol Chem *276*, 25309–25317. 10.1074/jbc.M101914200.

30. Fritsch, L., Robin, P., Mathieu, J.R.R., Souidi, M., Hinaux, H., Rougeulle, C.,

Harel-Bellan, A., Ameyar-Zazoua, M., and Ait-Si-Ali, S. (2010). A subset of the

histone H3 lysine 9 methyltransferases Suv39h1, G9a, GLP, and SETDB1

participate in a multimeric complex. Mol Cell *37*, 46–56.

10.1016/j.molcel.2009.12.017.

31. Ueda, J., Tachibana, M., Ikura, T., and Shinkai, Y. (2006). Zinc finger protein

WIZ links G9a/GLP histone methyltransferases to the co-repressor molecule CtBP. J

Biol Chem *281*, 20120–20128. 10.1074/jbc.M603087200.

32. Maier, V.K., Feeney, C.M., Taylor, J.E., Creech, A.L., Qiao, J.W., Szanto, A.,

Das, P.P., Chevrier, N., Cifuentes-Rojas, C., Orkin, S.H., et al. (2015). Functional

proteomic analysis of repressive histone methyltransferase complexes reveals

ZNF518B as a G9A regulator. Mol Cell Proteom *14*, 1435–1446.

10.1074/mcp.M114.044586.

33. Ohta, S., Bukowski-Wills, J.-C., Sanchez-Pulido, L., Alves, F. de L., Wood, L.,

Chen, Z.A., Platani, M., Fischer, L., Hudson, D.F., Ponting, C.P., et al. (2010). The

protein composition of mitotic chromosomes determined using multiclassifier

combinatorial proteomics. Cell *142*, 810–821. 10.1016/j.cell.2010.07.047.

34. Ohta, S., Montaňo-Gutierrez, L.F., Alves, F. de L., Ogawa, H., Toramoto, I., Sato,

N., Morrison, C.G., Takeda, S., Hudson, D.F., Rappsilber, J., et al. (2016).

Proteomics analysis with a nano random Forest approach reveals novel functional

interactions regulated by SMC complexes on mitotic chromosomes. Mol Cell

Proteom *15*, 2802–2818. 10.1074/mcp.M116.057885.

35. Canzio, D., Larson, A., and Narlikar, G.J. (2014). Mechanisms of functional

promiscuity by HP1 proteins. Trends Cell Biol *24*, 377–386.

10.1016/j.tcb.2014.01.002.

36. Ohzeki, J., Shono, N., Otake, K., Martins, N.M.C., Kugou, K., Kimura, H.,

Nagase, T., Larionov, V., Earnshaw, W.C., and Masumoto, H. (2016).

KAT7/HBO1/MYST2 regulates CENP-A chromatin assembly by antagonizing

Suv39h1-mediated centromere inactivation. Dev Cell *37*, 413–427.

10.1016/j.devcel.2016.05.006.

37. Ohzeki, J., Bergmann, J.H., Kouprina, N., Noskov, V.N., Nakano, M., Kimura, H.,

Earnshaw, W.C., Larionov, V., and Masumoto, H. (2012). Breaking the HAC barrier:

histone H3K9 acetyl/methyl balance regulates CENP-A assembly. EMBO J *31*,

2391–2402. 10.1038/emboj.2012.82.

38. Bergmann, J.H., Rodríguez, M.G., Martins, N.M.C., Kimura, H., Kelly, D.A.,

Masumoto, H., Larionov, V., Jansen, L.E.T., and Earnshaw, W.C. (2011). Epigenetic

engineering shows H3K4me2 is required for HJURP targeting and CENP-A

assembly on a synthetic human kinetochore. EMBO J *30*, 328–340.

10.1038/emboj.2010.329.

39. Martins, N.M.C., Bergmann, J.H., Shono, N., Kimura, H., Larionov, V.,

Masumoto, H., and Earnshaw, W.C. (2016). Epigenetic engineering shows that a

human centromere resists silencing mediated by H3K27me3/K9me3. Mol Biol Cell

27, 177–196. 10.1091/mbc.e15-08-0605.

40. Shono, N., Ohzeki, J., Otake, K., Martins, N.M.C., Nagase, T., Kimura, H.,

Larionov, V., Earnshaw, W.C., and Masumoto, H. (2015). CENP-C and CENP-I are

key connecting factors for kinetochore and CENP-A assembly. J Cell Sci *128*, 4572–

4587. 10.1242/jcs.180786.

41. Thiru, A., Nietlispach, D., Mott, H.R., Okuwaki, M., Lyon, D., Nielsen, P.R.,

Hirshberg, M., Verreault, A., Murzina, N.V., and Laue, E.D. (2004). Structural basis

of HP1/PXVXL motif peptide interactions and HP1 localisation to heterochromatin.

EMBO J *23*, 489–499. 10.1038/sj.emboj.7600088.

42. Nozawa, R.-S., Nagao, K., Masuda, H.-T., Iwasaki, O., Hirota, T., Nozaki, N.,

Kimura, H., and Obuse, C. (2010). Human POGZ modulates dissociation of HP1α

from mitotic chromosome arms through Aurora B activation. Nat Cell Biol *12*, 719–

727. 10.1038/ncb2075.

43. Natsume, T., Kiyomitsu, T., Saga, Y., and Kanemaki, M.T. (2016). Rapid protein

depletion in human cells by auxin-inducible degron tagging with short homology

donors. Cell Rep *15*, 210–218. 10.1016/j.celrep.2016.03.001.

44. Simon, J.M., Parker, J.S., Liu, F., Rothbart, S.B., Ait-Si-Ali, S., Strahl, B.D., Jin,

J., Davis, I.J., Mosley, A.L., and Pattenden, S.G. (2015). A role for widely

interspaced zinc finger (WIZ) in retention of the G9a methyltransferase on chromatin.

J Biol Chem *290*, 26088–26102. 10.1074/jbc.M115.654459.

45. Bian, C., Chen, Q., and Yu, X. (2015). The zinc finger proteins ZNF644 and WIZ regulate the G9a/GLP complex for gene repression. eLife *4*, e05606. 10.7554/eLife.05606.

46. Itoh, M., Yasunishi, A., Imamura, K., Kanamori-Katayama, M., Suzuki, H., Suzuki, M., Carninci, P., Kawai, J., and Hayashizaki, Y. (2006). Constructing ORFeome resources with removable termination codons. BioTechniques *41*, 44–50. 10.2144/000112209.

47. Nagase, T., Yamakawa, H., Tadokoro, S., Nakajima, D., Inoue, S., Yamaguchi, K., Itokawa, Y., Kikuno, R.F., Koga, H., and Ohara, O. (2008). Exploration of human ORFeome: high-throughput preparation of ORF clones and efficient characterization of their protein products. DNA Res *15*, 137–149. 10.1093/dnares/dsn004.

48. Kimura, K., Wakamatsu, A., Suzuki, Y., Ota, T., Nishikawa, T., Yamashita, R., Yamamoto, J., Sekine, M., Tsuritani, K., Wakaguri, H., et al. (2006). Diversification of transcriptional modulation: large-scale identification and characterization of putative alternative promoters of human genes. Gen Res *16*, 55–65. 10.1101/gr.4039406.

49. Otsuki, T., Ota, T., Nishikawa, T., Hayashi, K., Suzuki, Y., Yamamoto, J., Wakamatsu, A., Kimura, K., Sakamoto, K., Hatano, N., et al. (2005). Signal sequence and keyword trap in silico for selection of full-length human cDNAs encoding secretion or membrane proteins from oligo-capped cDNA libraries. DNA Res *12*, 117–126. 10.1093/dnares/12.2.117.

50. Ota, T., Suzuki, Y., Nishikawa, T., Otsuki, T., Sugiyama, T., Irie, R., Wakamatsu, A., Hayashi, K., Sato, H., Nagai, K., et al. (2004). Complete sequencing and

characterization of 21,243 full-length human cDNAs. Nat Genet *36*, 40–45.

10.1038/ng1285.

51. Ohta, S., Hamada, M., Sato, N., and Toramoto, I. (2015). Polyglutamylated

tubulin binding protein C1orf96/CSAP is involved in microtubule stabilization in

mitotic spindles. PloS One *10*, e0142798. 10.1371/journal.pone.0142798.

52. Cħaga, G., Bochkariov, D.E., Jokhadze, G.G., Hopp, J., and Nelson, P. (1999).

Natural poly-histidine affinity tag for purification of recombinant proteins on cobalt(II)-

carboxymethylaspartate crosslinked agarose. J Chromatogr A *864*, 247–256.

10.1016/s0021-9673(99)01008-0.

53. Ohta, S., Taniguchi, T., Sato, N., Hamada, M., Taniguchi, H., and Rappsilber, J.

(2019). Quantitative proteomics of the mitotic chromosome scaffold reveals the

association of baz1b with chromosomal axes. Mol Cell Proteom *18*, 169–181.

10.1074/mcp.ra118.000923.

54. Ruppert, J.G., Samejima, K., Platani, M., Molina, O., Kimura, H., Jeyaprakash,

A.A., Ohta, S., and Earnshaw, W.C. (2018). HP1α targets the chromosomal

passenger complex for activation at heterochromatin before mitotic entry. EMBO J

37, e97677. 10.15252/embj.201797677.

55. Ohta, S., Bukowski-Wills, J.-C., Wood, L., de Lima Alves, F., Chen, Z.,

Rappsilber, J., and Earnshaw, W.C. (2010). Proteomics of isolated mitotic

chromosomes identifies the kinetochore protein Ska3/Rama1. Cold Spring Harb

Symp Quant Biol *75*, 433–438. 10.1101/sqb.2010.75.022.

56. Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ:

years of image analysis. Nat Methods *9*, 671–675. 10.1038/nmeth.2089.

57. Altschul, S.F., Gish, W., Miller, W., Myers, E.W., and Lipman, D.J. (1990). Basic

local alignment search tool. J Mol Biol *215*, 403–410. 10.1016/S0022-

2836(05)80360-2.

58. Kumar, S., Stecher, G., Li, M., Knyaz, C., and Tamura, K. (2018). MEGA X:

molecular evolutionary genetics analysis across computing platforms. Mol Biol Evol

35, 1547–1549. 10.1093/molbev/msy096.