Formation of pericentromeric heterochromatin via ZNF518 that links satellite

DNA to heterochromatin

Shinya Ohta[1,2]*, Jun-Ichirou Ohzeki[3], Nobuko Sato[2], Ken-Ichi Noma[1,4], and Hiroshi

Masumoto[3]*

[1]Institute for Genetic Medicine, Hokkaido University, Kita-15, Nishi-7, Kita-Ku,

Sapporo 060-0815, Japan

[2]Department of Biochemistry, Medical School, Kochi University, Kohasu, Oko-cho,

Nankoku, Kochi 783-8505, Japan

[3]Laboratory of Chromosome Engineering, Department of Frontier Research and

Development, Kazusa DNA Research Institute, 2-6-7 Kazusa-Kamatari, Kisarazu 292-

0818, Japan

[4]Institute of Molecular Biology, University of Oregon, 1370 Franklin Blvd, Eugene,

OR, 97403, USA

*Corresponding authors:

shinya.ohta@igm.hokudai.ac.jp

masumoto@kazusa.or.jp

Abstract

Aneuploidy is caused by chromosomal missegregation and is frequently observed in

cancers and hematological diseases. The intricate structure formed at the centromere

or kinetochore plays an integral role in the segregation of chromosomes. For instance,

heterochromatin at pericentromere is essential for proper chromosomal segregation;

however, it is unclear how heterochromatin is targeted to pericentromeres. However

the connection between Heterochromatin and centromere still remains unresolved.

Here, we demonstrate a novel mechanism by which two homologous zinc-finger

proteins (ZNF518A and ZNF518B) play central roles in heterochromatin formation at

pericentromeres. We investigate this mechanism by determining the cellular

localization of ZNF518, centromeric proteins such as CENP-A and CENP-B, and

heterochromatin factors, including HP1 and histone H3K9 trimethylation. The results

show that multiple segments within ZNF518 contribute to its interactions with CENP-

B, HP1, and G9a histone H3K9 methyltransferase. This study identifies a novel

mechanism underlying pericentromeric heterochromatin formation mediated by the

central component ZNF518.

KEYWORDS

centromere, pericentromere, heterochromatin, ZNF518, CENP-B

1 Introduction

Defects in centromere or kinetochore functions are directly related to chromosomal

abnormalities, and the anomalous segregation of chromosomes can lead to cancer or

hematological diseases.[1,2] Chromosomal abnormalities can occur during mitosis,

where replicated chromosomes are normally distributed equally between two daughter

cells. During mitosis, compacted chromosomes are segregated when the spindle

attaches to and pulls the kinetochore, a proteinaceous structure on the chromosome.

The region of the chromosome that forms the kinetochore is the centromere.

Centromere protein A (CENP-A), a centromere-specific histone H3 variant, is essential

for kinetochore formation on centromeres and directs the assembly of a complex

containing at least 15 other proteins, including centromere protein C (CENP-C).[3] The

centromeric factors CENP-A, CENP-B, and CENP-C are evolutionarily conserved

from fission yeast to vertebrates and show distinctive association patterns within the

centromere, respectively.[4,5]

In normal human chromosomes, cetromeres assemble on α-satellite DNA

(alphoid DNA), a huge repetitive DNA locus consisting of 171 bp repeat units.[6,7]

CENP-A chromatin assembles on a part of α-satellite DNA, forming an epigenetic

mark of kinetochore assembly.[8–11] Some subclasses of α-satellite DNA contain a 17-

nucleotide CENP-B box sequence bound by CENP-B.[8,12,13] Thus, CENP-B is

distributed in both CENP-A chromatin (core centromere) and

pericentromeric heterochromatin. CENP-B is involving in for the assembly of CENP-A

and CENP-C in centromeres and, consequently, for functional centromere

formation.[9,10] Furthermore, CENP-B and CENP-B boxes are required for human

artificial chromosome formation and de novo centromeric CENP-A chromatin

assembly when introducing alphoid DNA into cells.[14–16] CENP-B also induces

heterochromatin formation in alphoid DNA integrated into chromosomal arms.[16]

Interestingly, CENP-B recruits many factors in a mutually exclusive manner, including

the open chromatin modifier ASH1L (H3K36 methylase) or heterochromatin factor

HP1.[17] Heterochromatin is reported to have important roles in chromosome stability

and in maintenance of sister chromatid cohesion during chromosome segregation.[18,19]

Therefore, both centromere chromatin and heterochromatin are important for accurate

chromosome segregation processes.

Pericentromeric heterochromatin insulates centromeres from the rest of the

genome and is required for the proper segregation of mitotic chromosomes.[19,20]

Additionally, heterochromatin is known to have a more condensed structure than

euchromatin and carries specific post-translational histone modifications.

Nucleosomal surfaces with charge differences contribute to the condensed structure

of heterochromatin.[21–23] Furthermore, the trimethylation of histone H3 at lysine 9

(H3K9me3) is recognized and bound by the chromodomain of the heterochromatin

protein HP1.[24] HP1, in turn, recruits the suppressor of variegation 3-9 homolog

(SUV39H), which is an H3K9 methyltransferase.[18,25] The HP1-mediated loading of

SUV39H to chromatin is required for the maintenance of heterochromatin at

pericentromeres.[26,27] Since SUV39H localizes to pericentromeres in addition to other

constitutive heterochromatin regions, heterochromatin formation at pericentromeres

involves SUV39H. The mechanisms by which heterochromatin factors are initially targeted to pericentromeres, however, remain elusive.[28]

Enzymes other than SUV39H have also been recognized as H3K9 methyltransferases. SET domain bifurcated 1 (SETDB1), for example, catalyzes H3K9me3 formation, while euchromatic histone lysine methyltransferase 1 and 2 (GLP and G9a) are responsible for H3K9me1 and H3K9me2, respectively. G9a can also methylate Lys9 and Lys27 of histone H3.[29] These H3K9 methyltransferases, including SUV39H, are selectively loaded onto distinct genomic regions through protein interactions with other factors.[30] G9a and GLP bind to the widely interspaced zinc-finger (WIZ) protein.[29,31,32] It has been observed that H3K9 methylation at chromosomal arm regions is mostly lost in G9a-deficient cells, while H3K9 methylation at pericentromeres is largely lost in SUV39H-deficient cells.[29] Moreover, the interaction between G9a and zinc-finger protein 518B (ZNF518B) was recently identified via a proteomics analysis.[32] Our previous proteomics analysis revealed a possible role for ZNF518B as a functional protein on mitotic chromosomes.[33,34] The present study demonstrates that G9a, which acts as a H3K9 methyltransferase for chromosomal arm regions, is also involved in heterochromatin formation at pericentromeres via its interaction with ZNF518B (which is targeted to pericentromeres by CENP-B). This study provides novel evidence that G9a plays a role in heterochromatin formation at the pericentromeres.

Results

Human ZNF518A/B is localized to centromeres via CENP-B

We first examined the localization of green fluorescent protein (GFP)-tagged

ZNF518B and the centromeric factors CENP-A, CENP-B, and CENP-C in human

U2OS cells. Overexpression of GFP fusion proteins allows for enhanced visualization

of even weakly-expressed proteins; with this technique, we were able to successfully

visualize of the localization of ZNF518. Surprisingly, GFP-ZNF518B co-localized

proximally to the centromeric factors (Fig. 1A–F). In addition, ZNF518B localized to

the centromeres during mitosis (Fig. S1A). This result suggests constitutive

localization of ZNF518B to the centromere throughout the cell cycle. The GFP-

ZNF518B foci were slightly shifted from the CENP-A and CENP-C foci but coincided

with the CENP-B foci, indicating that ZNF518B localized to the pericentromeric region

of the centromere, where it was bound by CENP-B (Fig. 1A–F).

To investigate the localization dependency between ZNF518B and CENP-B,

we examined the localization of GFP-ZNF518B in CENP-B knockout (KO) HeLa

cells.[17]   GFP-ZNF518B   did   not   show   centromeric   localization

(centromeres/pericentromeres) in CENP-B KO cells (Fig. 1G, H); however,

centromeric localization was restored when exogeneous CENP-B was expressed in

the CENP-B KO cells (Fig. 1I). These results indicate that the centromeric localization

of ZNF518B depended on CENP-B.

ZNF518A is a paralogue of ZNF518B present in humans (Fig. S1B). To

determine the localization of ZNF518A, GFP-ZNF518B and Halo-ZNF518A were

transiently co-expressed in U2OS cells (Fig. 1J). The GFP-ZNF518B and Halo-

ZNF518A foci co-localized at the centromeres. Again, the co-localized signals were

slightly shifted from the CENP-A foci, suggesting that both ZNF518A and ZNF518B

localize at centromeres (Fig. 1J). Moreover, GFP-ZNF518A did not show centromeric

localization in the CENP-B KO cells, and centromeric localization was restored when

exogenous CENP-B was expressed (Fig. S1C). These findings indicated that the

centromeric localization of ZNF518A was dependent on CENP-B. Taken together, our

results suggest that both ZNF518A and ZNF518B localize to the pericentromere via a

common molecular mechanism involving CENP-B or that ZNF518A and ZNF518B

interact and are recruited to pericentromeres by CENP-B (Fig. S1D).

The C-terminal segment of ZNF518A/B is required for centromeric localization

To gain insights into the molecular mechanisms underlying the recruitment of ZNF518

to centromeres, we sought to identify the segments within ZNF518B that are required

for centromeric localization. We examined the centromeric localization of 15

derivatives of ZNF518B expressed as GFP fusion proteins (Fig. 2). ZNF518B

comprises 1,074 amino acids (aa). The ZNF518B derivatives containing regions from

the N-terminus to 628 aa (ZNF518B$^{1-398}$, ZNF518B$^{344-628}$, and ZNF518B$^{1-628}$) did not

show centromeric localization (Fig. 2, S2A). Therefore, the N-terminus of ZNF518B

did not form sufficient interactions to localize the protein to centromeres.

Upon assessing the centromeric localization of the four ZNF518B derivatives

(ZNF518B$^{344\text{-}1074}$, ZNF518B$^{344\text{-}987}$, ZNF518B$^{344\text{-}877}$, and ZNF518B$^{344\text{-}769}$) with

incrementing C-terminal truncations, we found that three derivatives (ZNF518B$^{344\text{-}1074}$,

ZNF518B$^{344\text{-}987}$, and ZNF518B$^{344\text{-}877}$) showed centromeric localization in 70.7%,

58.9%, and 57.1% of the transfected cells, respectively (Fig. 2). In sharp contrast,

ZNF518B$^{344\text{-}769}$ did not show centromeric localization (Fig. 2). Similarly, among the

three derivatives ZNF518B$^{555\text{-}1074}$, ZNF518B$^{555\text{-}877}$, and ZNF518B$^{555\text{-}769}$, only

ZNF518B$^{555\text{-}769}$, which has the same C-terminal truncation as ZNF518B$^{344\text{-}769}$, did not

show centromeric localization (Fig. 2, S2A). These results suggest that residues 770–

877 of ZNF518B are deeply involved in its centromeric localization. However, only

2.2% of the transfected cells showed centromeric localization of ZNF518B$^{770\text{-}877}$ (Fig.

2). Therefore, residues 770–877 are important, but insufficient alone, for centromeric

localization.

Due to the large difference in the percentage of cells localized to the centromere

between ZNF518B$^{555\text{-}1074}$, ZNF518B$^{555\text{-}877}$, and ZNF518B$^{555\text{-}769}$ (64.5%, 31.8%, and

0% of the cells, respectively), we examined the centromeric localization of ZNF518B

derivatives that carried residues 770–1074 (Fig. 2, S2A). The three ZNF518B

derivatives ZNF518B$^{878\text{-}1074}$, ZNF518B$^{878\text{-}987}$, and ZNF518B$^{770\text{-}987}$ showed centromeric

localization to varying degrees (53.6%, 56.7%, and 89.3%, respectively; Fig. 2, S2A).

ZNF518B$^{770\text{-}987}$ showed the highest frequency of centromeric localization among all

the ZNF518B derivatives. These results collectively suggest that residues 770–987

are Involved in the centromeric localization of ZNF518B.

We next examined the centromeric localization of several ZNF518A derivatives

(Fig. S2B). The wild-type (WT) ZNF518A (ZNF518A[WT]) and two derivatives

(ZNF518A[971-1483] and ZNF518A[1317-1483]) containing the C-terminal 1317–1483-aa

segment showed high frequencies of centromeric localization, suggesting that the C-

terminal portion of ZNF518A harbors crucial segment(s) for centromeric localization

(Fig. S2B). Notably, the C-terminal segments of ZNF518A and ZNF518B required for

centromeric localization carried PxV(I)xL HP1-interaction motifs, which are widely

conserved among higher vertebrates, including fish (Fig. S3). This implied their

evolutionarily conserved role potentially through their localization at centromeres.

The C-terminal PxIxL motif is required for the centromeric localization of

ZNF518B

To elucidate the functional significance of the PxV(I)xL motifs in ZNF518B, we

constructed the GFP-ZNF518B[KSS/C] derivative and substituted the conserved

hydrophobic residues PxIxL in the C-terminal segment with the hydrophilic residues

KxSxS (Fig. 2). ZNF518B[KSS/C] exhibited centromeric localization (14.3%) less

frequently compared to ZNF518[WT] (77.5%; Fig. 2). In contrast, the centromeric

localization of ZNF518B[KSS/N], which carries a KxSxS substitution at the PxVxL motif in

the N-terminal segment, was comparable to that of ZNF518[WT] (Fig. 2). ZNF518B[KSS/NC],

which carries the KxSxS substitutions for the PxV(I)xL motifs in both the N- and C-

terminal segments, showed a similar frequency of centromeric localization (21.4%) as

the ZNF518B[KSS/C] derivative (Fig. 2). The PxIxL motif present in the C-terminal

segment, but not in the N-terminal, is therefore involved in the centromeric localization

of ZNF518B. ZNF518A^{KSS}, which carries a KxSxS substitution at the PxVxL motif in

the C-terminal segment, also showed the centromeric localization with reduced

frequency (Fig. S2B). These results demonstrate that the conserved PxV(I)xL motif in

the C-terminal segment of ZNF518 is important for centromeric localization.

Although the C-terminal PxIxL motif is important for centromeric localization,

both ZNF518B$^{KSS/C}$ and ZNF518B$^{KSS/NC}$ could localize at the centromeres in 14.3%

and 21.4% of the cells, respectively, implying the involvement of other interactions in

centromeric localization (Fig. 2, S2B). To test this notion, we used the ZNF518B$^{770\text{-}987}$

derivative that localized to centromeres with the highest frequency and substituted the

PxVxL motif in the C-terminal segment with KxSxS. This yielded the new derivative,

ZNF518B$^{770\text{-}987\ KSS/C}$. Although less than that for ZNF518B$^{770\text{-}987}$, ZNF518B$^{770\text{-}987\ KSS/C}$

showed centromeric localization in 60.6% of the transfected cells (Fig. 2, S2B). These

results reveal that both the PxIxL motif and C-terminal segment (770–987 aa)

participate in the centromeric localization of ZNF518B.

ZNF518B interacts with HP1α and HP1β

The localization of full-length ZNF518B^{WT} to centromeres and pericentromeres

requires CENP-B. The ZNF518B$^{770\text{-}987}$ derivative with the C-terminal segment (770–

987 aa) that includes the PxIxL motif also localized to centromeres (Fig. 2). In CENP-

B KO cells, ZNF518B$^{770\text{-}987}$ showed focal localization that was not observed for

ZNF518B^{WT} (Fig. S4A). However, the ZNF518B$^{770\text{-}987}$ foci were distinct from

centromeres. These foci were found to co-localize with the heterochromatin foci in the

CENP-B KO cells (Fig. S4B). Thus, ZNF518B[770-987] can be recruited to non-

centromeric heterochromatin domains through its interaction with heterochromatic

factors in the absence of CENP-B (Fig. S4C).

Given its similarity to the canonical HP1-interacting motif PxVxL, we

hypothesized that the PxIxL motif in the C-terminal segment of ZNF518B interacts with

HP1, thereby resulting in the accumulation of ZNF518B at heterochromatin regions in

CENP-B KO cells. Three isozymes of human HP1 exist: HP1α, HP1β, and HP1γ.[35]

We attempted to characterize the interaction between ZNF518B and the HP1 isoforms

using the fluorescence microscopy-based interaction-trap (FMIT) assay.[36] The FMIT

assay may be used to examine protein interactions on chromatin *in vivo*. Given the

overabundance of target proteins compared to their endogenous counterparts, this

technique allows for the detection of direct and indirect protein–protein interactions

using fluorescence microscopy.[17,20,36–40]

An FMIT assay was performed to test the interaction between ZNF518B and

HP1 isoforms in HeLa-Int-03 cells. In the HeLa-Int-03 cells, approximately 15,000

repeats of the synthetic alphoid DNA dimer unit (derived from the human chromosome

21 centromere) containing the tetO sequence (alphoid*tetO*) were ectopically integrated

into a chromosome arm region, and CENP-B was knocked out (Fig. 3A).[17] TetR-

EYFP-ZNF518B localized to the alphoid*tetO* repeats but did not exhibit centromeric

localization due to the absence of CENP-B. We next examined whether Halo-HP1 co-

expressed with TetR-EYFP-ZNF518B would be recruited to the alphoid^{tetO} array (Fig.

3B). Halo-HP1α, -HP1β, and -HP1γ were recruited to the alphoid^{tetO} in 48%, 58%, and

14% of the cells, respectively (Fig. 3C). In the negative controls where TetR-EYFP not

fused to ZNF518B was expressed, Halo-fused HP1α, HP1β, and HP1γ were barely

recruited to the alphoid^{tetO} (Fig. 3C). These results suggest that ZNF518B interacts

with HP1.

The PxIxL motif in the C-terminal segment of ZNF518B interacts with HP1

isoforms

We next investigated the segments within ZNF518B that are required to interact with

the HP1 isoforms by subjecting the ZMF518B derivatives described in Figure 1 to the

FMIT assay (Fig. 3C). Upon co-expressing Halo-HP1 and TetR-EYFP-fused ZNF518B

derivatives   (ZNF518B$^{1-398}$,   ZNF518B$^{1-628}$,   ZNF518B$^{344-628}$,   ZNF518B$^{344-877}$,

ZNF518B$^{555-877}$, and ZNF518B$^{770-877}$), Halo-HP1 was either scarcely or not recruited

to the alphoid^{tetO}. Since these derivatives are devoid of the PxIxL motif, this result

implies the importance of the PxIxL motif in the interactions between ZNF519B and

the HP1 isoforms (Fig. 3C).

To further examine the significance of the PxV(I)xL motifs in the ZNF518B-HP1

interactions, TetR-EYFP-fused ZNF518B derivatives (ZNF518B$^{KSS/N}$, ZNF518B$^{KSS/C}$,

ZNF518B$^{KSS/NC}$, and ZNF518B$^{770-987\ KSS/C}$) were used for the FMIT assay. These

derivatives carry the substitutions of the PxV(I)xL motifs with KxSxS, as depicted in

Figure 1. Halo-HP1 was scarcely recruited to the alphoid^{tetO} by the ZNF518B

derivatives, apart from ZNF518B$^{KSS/N}$. The PxIxL motif in the C-terminal segment, but

not the PxVxL motif in the N-terminal segment, therefore participates in the interaction

with HP1 (Fig. 3C).

Next, the TetR-EYFP-fused ZNF518B derivatives ZNF518B$^{344-1074}$,

ZNF518B$^{344-987}$, ZNF518B$^{555-1074}$, ZNF518B$^{878-1074}$, ZNF518B$^{878-987}$, and ZNF518B^{770-}

^{987} were used for the FMIT assay. These derivatives all carry the PxIxL motif in the C-

terminal segment of ZNF518B. We observed that Halo-HP1 was recruited to the

alphoid^{tetO} in more than 75% of the cells (Fig. 3C). Remarkably, these N-terminal-

deleted derivatives were able to recruit the three distinct HP1 proteins with similar

frequencies. Furthermore, ZNF518B$^{344-1074}$ was more efficient at recruiting HP1 than

ZNF518B^{WT}. The N-terminal segment (1–398) is therefore inhibitory for the ZNF518B

interactions with HP1 isoforms (Fig. 3C).

The PxIxL motif of ZNF518B interacts with the chromo-shadow domain of HP1

HP1 comprises a central linker flanked by chromo and chromo-shadow domains,

the latter of which interacts with the PxVxL motif in CAF1/p150 [41]. Since the PxIxL

motif in ZNF518B plays a vital role in the interactions with HP1 isoforms, we predicted

that ZNF518B may interact with the chromo-shadow domain rather than the chromo

domain of HP1 isoforms. Using the FMIT assay, we examined the interaction between

ZNF518B and the chromo domain (HP1β$^{1-92}$) or chromo-shadow domain (HP1β$^{93-185}$)

of HP1β. As expected, the chromo-shadow domain, but not the chromo domain, was

recruited to the alphoid^{tetO} by ZNF518B derivatives carrying the PxIxL motif in the C-

terminal segment. This confirmed the interaction between the PxIxL motif of ZNF518B

and chromo-shadow domain of HP1β (Fig. 3C).

ZNF518B$^{344-769}$ and ZNF518B$^{555-769}$ weakly bound HP1β despite the absence of

a PxIxL motif. Indeed, the PxIxL motif of ZNF518B had similar binding affinities for all

three HP1 isoforms. However, residues 971–1321 of ZNF518A, which correspond to

residues 555–769 of ZNF518B, did not show such specificity for the HP1 isoforms (Fig.

S5B). The 555–769-aa segment in ZNF518B is therefore a subsidiary segment for the

HP1 interactions that determine its specificity for HP1β.

ZNF518B maintains HP1 at the alphoid^{tetO}

We next examined whether HP1β loaded to the alphoid^{tetO} by TetR-EYFP-ZNF518B

is maintained at the alphoid^{tetO} at the repeat locus when ZNF518B is released from

the alphoid^{tetO} following the addition of doxycycline (Dox) (Fig. 3D). TetR-EYFP-

ZNF518 was rapidly released from the alphoid^{tetO} after introducing Dox. EYFP foci

disappeared in 96% of the cells 120 min after Dox introduction (Fig. S5C). We further

quantified the presence of Halo-HP1β foci at the alphoid^{tetO} 150 min after the

introduction of Dox (Fig. 3E). The Halo-HP1β foci disappeared from the alphoid^{tetO} in

all experiments employing ZNF518B^{WT}, ZNF518B$^{878-987}$, and ZNF518B$^{770-987}$ (Fig. 3E).

ZNF518B is therefore required to maintain HP1β at the alphoid^{tetO}.

ZNF518B interacts directly with HP1 isoforms *in vitro*

The above FMIT results strongly suggest that ZNF518B interacts with HP1 isoforms. We were able to reproduce previously published results regarding the interaction of ZNF518B with HP1s.[42] To further examine their direct interactions, we purified the ZNF518B770-987 derivative containing the PxIxL motif as maltose-binding protein (MBP) fusion protein and HP1 isoforms as glutathione S-transferase (GST) fusion protein using overexpression system in *Escherichia coli* cells , respectively. The two purified proteins were mixed, and the MBP-fused ZNF518B$^{770-987}$ proteins were pulled-down using magnetic amylose beads. We observed that the HP1 isoforms were co-purified with ZNF518B$^{770-987}$. The GST-HP1 isoforms were more abundant in the pull-down (PD) samples with MBP-ZNF518B$^{770-987}$ than in those with MBP alone (Fig. 3F). These results suggest that the PxIxL motif at the C-terminal segment of ZNF518B directly interacts with HP1α, HP1β, and HP1γ (Fig. 3G).

ZNF518B localization cannot be maintained at centromeres without CENP-B

The previous results (Fig. 1) showed that CENP-B is required for ZNF518B localization at centromeres. Based on this observation, we next investigated whether CENP-B is required to maintain ZNF518B at centromeres. We created a CENP-B conditional KO system using human U2OS cells (Fig. S6A). The gene expressing *Oryza sativa* TIR1 was inserted into an adeno-associated virus genomic integration site. In these cells, the endogenous CENP-B fused with a mini auxin-inducible degron (mAID) tag could be rapidly targeted for degradation via the ubiquitin-proteasome pathway following indole-3-acetic acid (IAA) treatment (Fig. S6A).[43]

Subsequent analysis was conducted under three conditions to examine the

localization of GFP-ZNF518B (Fig. 4A): CENP-B:mAID available at all times (CENP-

B[ON]); CENP-B:mAID removed 24 h prior to ZNF518B transfection (CENP-B[OFF]); and

CENP-B:mAID removed 4 h before fixation (CENP-B[4h-OFF]). As expected, GFP-

ZNF518B localized at the centromeres in CENP-B[ON] cells (Fig. 4B–E). In CENP-B[OFF]

cells, the CENP-B signals at the centromeres were below the detection limit, and GFP-

ZNF518B dissociated from the centromeres (Fig. 4B–E). In the CENP-B[4h-OFF] cells,

GFP-ZNF518B also dissociated from the centromeres (Fig. 4B–E). These results

demonstrated that the recruitment and maintenance of ZNF518B at centromeres

require CENP-B.

Interaction between ZNF518B and CENP-B

We examined the interaction between ZNF518B and CENP-B using the FMIT assay.

For this purpose, we constructed Halo-CENP-B[133-599], which is devoid of the helix-turn-

helix domain and thus cannot bind to α-satellites at centromeres.[17] Halo-CENP-B[133-

599] and TetR-EYFP-ZNF518B (WT and derivatives) were co-expressed in HeLa 03 Int

cells (Fig. 4F). As expected, Halo-CENP-B[133-599] did not localize to centromeres.

Therefore, CENP-B[133-599] foci reflected the recruitment of CENP-B to the alphoid*tetO* by

TetR-EYFP-ZNF518B.

We observed that CENP-B[133-599] was recruited to the alphoid*tetO* in 26% of the

cells expressing ZNF518B[WT], despite CENP-B[133-599] not being recruited to the loci in

cells expressing TetR-EYFP and lacking ZNF518B. Moreover, ZNF518B^{WT} was

recruited to the alphoid^{tetO} in cells expressing TetR:EYFP:CENP-B$^{133-599}$ (Fig. S6B).

These results were consistent with the interaction between ZNF518B and CENP-B

(Fig. 4G). In addition, CENP-B$^{133-599}$ was barely recruited at the alphoid^{tetO} by

ZNF518B$^{1-398}$, ZNF518B$^{1-628}$, and ZNF518B$^{344-628}$, suggesting that the ZNF518B

segment from 629 aa to the C-terminus is important for the interaction. Moreover,

CENP-B$^{133-599}$ was recruited to the alphoid^{tetO} at high frequencies in cells expressing

ZNF518B$^{344-1074}$, ZNF518B$^{344-987}$, ZNF518B$^{344-877}$, ZNF518B$^{555-1074}$, ZNF518B$^{555-877}$,

ZNF518B$^{770-987}$, and ZNF518B$^{770-877}$, all of which contain residues 770–877 (Fig. 4G).

Moreover, the recruitment frequencies of CENP-B$^{133-599}$ were reduced by 20% when

cells either expressing ZNF518$^{344-877}$ or ZNF518$^{344-769}$ were compared, and by 39%

when cells expressing ZNF518$^{555-877}$ or ZNF518$^{555-769}$ were compared. ZNF518B^{878-}

^{1074} and ZNF518B$^{878-987}$, which do not carry residues 770–877 of ZNF518B, did not

exhibit CENP-B$^{133-599}$ recruitment. These results support the importance of residues

770–877 of ZNF518B to its interaction with CENP-B (Fig. 4H).

We next examined the recruitment of CENP-B to the alphoid^{tetO} in cells

expressing the ZNF518 derivatives ZNF518B$^{KSS/N}$, ZNF518B$^{KSS/C}$, ZNF518B$^{KSS/NC}$,

and ZNF518B$^{770-987\ KSS/C}$ and carrying substitutions of the PxV(I)xL motifs with KxSxS

(Fig. 4G). The substitutions of the PxV(I)xL motifs generally increased the recruitment

of CENP-B to the *tetO* repeats. This implies an inhibitory role of the PxV(I)xL motifs in

the interaction between ZNF518B and CENP-B. Although the association of ZNF518B

with both HP1 and CENP-B is less effective in the full-length protein (Figs. 3C and 4G),

the derivatives with the N-terminal region deleted showed considerably stronger

interactions. We speculate that this is because the interaction domain in some ZNF518

segments is exposed on the surface of the protein, allowing for easier interactions with

the binding partner.

The C-terminal segment of ZNF518B interacts with G9a

Published proteomics data have revealed an interaction between ZNF518B and G9a

(EHMT2).[32] We therefore attempted to identify the segments of ZNF518B that interact

with G9a and vice versa. We used the FMIT assay with cells co-expressing EYFP-

TetR-ZNF518 and Halo-G9a. The control experiment showed that G9a was recruited

to $tetO$ repeats in cells expressing ZNF518B^{WT} (Fig. S6D and 5A). G9a was recruited

to the alphoid^{tetO} at high frequencies in cells expressing ZNF518B$^{KSS/C}$,

ZNF518B$^{KSS/NC}$, ZNF518B$^{344-1074}$, ZNF518B$^{555-1074}$, and ZNF518B$^{878-1074}$, all of which

carry the C-terminal end (Fig. 5A). In contrast, G9a was barely recruited to the

alphoid^{tetO} by the other ZNF518B derivatives (Fig. 5A). The C-terminal end segment

(residues 988–1074) of ZNF518B are therefore crucial for its interaction with G9a.

We also performed FMIT experiments to examine the interaction between

ZNF518A and G9a. We observed that G9a was recruited to the alphoid^{tetO} in cells

expressing ZNF518A^{WT}, ZNF518A^{KSS}, ZNF518A$^{971-1483}$, ZNF518A$^{971-1483\ KSS}$,

ZNF518A$^{1317-1487}$, and ZNF518A$^{1317-1483\ KSS}$, all of which include C-terminal end (Fig.

5B). However, G9a was less efficiently recruited to the alphoid^{tetO} in cells expressing

ZNF518A$^{1-970}$ and ZNF518A$^{971-1321}$ compared to the other derivatives (Fig. 5B). These

results suggest that the C-terminal end segment (residues 1317–1483 are important

for the interaction between ZNF518A and G9a. When combined with previous results,

our data indicate that the C-terminal end segments within ZNF518 are required for its

interaction with G9a.

The segment downstream of the TAD in G9a interacts with ZNF518A/B

Interactions between G9a and the C2H2-type zinc-finger motifs in ZNF644 and

ZNF803, also known as Widely interspaced zinc finger (WIZ)-containing proteins,

have previously been reported.[31,44] The zinc-finger motifs in WIZ-containing proteins

also interact with the transcriptional regulatory CCAAT/enhancer-binding protein-β

(C/EBPβ) transactivation domain (TAD) within the N-terminal segments of G9a and

GLP [45]. Because zinc-finger motifs that interact with G9a are highly conserved in the

C-terminal segments of ZNF518A/B in all species except *Zenopus laevis* (Fig. S6E),

we hypothesized that the C-terminal segments of ZNF518A/B may interact with the

TAD of G9a.

To test this hypothesis, we performed FMIT experiments with several G9a

derivatives (Fig. 5C). In contrast with the above prediction, we found that G9a$^{1-290}$ was

not recruited to the alphoid^{tetO} by ZNF518A/B. Instead, G9a$^{291-1210}$, which lacks a TAD,

was recruited to the alphoid^{tetO} by ZNF518A/B. Moreover, the two 291–545-aa

segments containing the G9a derivatives, G9a$^{1-545}$ and G9a$^{291-545}$, were recruited to

the alphoid^{tetO} by ZNF518A/B. These results indicate that residues 291–545 of G9a,

but not the TAD, are sufficient for the interactions with ZNF518A/B.

The N-terminal residues 1–624 aa of ZNF518B did not interact with CENP-B,

HP1, or G9a, nor did they exhibit centromeric localization. However, the N-terminal

residues 1–970 aa of ZNF518A retained their centromeric localization with an

observed interaction with HP1 and less effective interaction with CENP-B and G9a

(Fig. S6C). ZNF518A and ZNF518B may therefore differ in terms of the functions of

their N-terminal segments.

ZNF518B recruit G9a H3K9 methyltransferase to centromeres

Finally, we assessed the possibility of the recruitment of G9a to centromeres by

ZNF518. Plasmids expressing GFP-ZNF518B or ZNF518B derivatives and Halo-G9a

were co-transfected into U2OS cells, and the localization of Halo-G9a was

investigated (Fig. 5D). The localization of Halo-G9a at centromeres was observed only

when Halo-G9a was expressed with WT or ZNF518B$^{344\text{-}1074}$ carrying the C-terminal

segment, as revealed by the FMIT assay (Fig. 5D). However, the localization of Halo-

G9a at centromeres was no longer observed when ZNF518B$^{344\text{-}987}$ lacking the C-

terminal zinc-finger was expressed (Fig. 5D). Furthermore, no localization of Halo-G9a

at centromeres was observed when GFP-ZNF518B and Halo-G9a were expressed in

CENP-B KO cells (Fig. 5E). CENP-B therefore recruits ZNF518B to centromeres, and

ZNF518B in turn recruits G9a H3K9 methyltransferase (Fig. 5F).

Discussion

In this study, we show that the 770–877 segment within ZNF518B interacts with CENP-B (Fig. 4G) and is recruited to centromeres and pericentromeres in a CENP-B-dependent manner (Fig. 1, 2, 4). Furthermore, the 555–765 segment and PxIxL motif within ZNF518B are required for its interaction with the chromo-shadow domain of HP1β (Fig. 3). The pericentromere forms constitutive heterochromatin, promoting HP1 accumulation.[8,9,11] Therefore, among the ZNF518 recruited to the centromere/pericentromere by CENP-B, which assembled in the pericentromere is likely to have maintained a stable association with the chromatin through interaction with HP1. The C-terminal segment carrying the ZF motif within ZNF518B interacts with the segment downstream of TAD within G9a, and CENP-B is required for the localization of G9a at centromeres (Fig. 5). Based on these observations, we propose the following mechanisms: 1) CENP-B binds to α-satellite DNA to initiate the establishment of heterochromatin; 2) CENP-B recruits ZNF518B to centromeres and pericentromeres; 3) HP1β stabilizes ZNF518 only at pericentromeres; and 4) ZNF518B interacts with G9a to introduce H3K9 methylation to pericentromeres.

These conclusions do not disagree with the previously proposed mechanism involving SUV39H1 histone H3K9 methyltransferase.[26,27] It has been reported that HP1-dependent SUV39H1 recruitment and short RNA-mediated heterochromatinization are important for the maintenance of pericentromeric heterochromatin.[26,28] Meanwhile, we propose that G9a recruited by CENP-B and ZNF518B is required for the establishment of pericentromeric heterochromatin. We

suggest that the CENP-B/ZNF518B/HP1β/G9a axis can target heterochromatin factors to pericentromeres, where specificity is yielded by the binding of CENP-B to CENP-B box sequences in pericentromeric alphoid repeats.

In addition to ZNF518B, ZNF518A is recruited to pericentromeres in a CENP-B-dependent manner (Fig. S1) and interacts with HP1 and G9a (Fig. 5, S5). ZNF518A and ZNF518B are further evolutionarily conserved in all vertebrates higher than amphibians. Therefore, we speculate that pericentromeric heterochromatin formation by ZNF518 is conserved among many vertebrates. Although further investigation is needed, our work provides insights into the potential mechanisms underlying the targeting of heterochromatin to human pericentromeres and indicate the possibility of a similar mechanism existing in many species.

Materials and Methods

DNA constructs

To clone the target genes, DNA fragments encoding human ZNF518A, ZNF518B,

CENP-B$^{133\text{-}599}$, HP1α, HP1β, HP1γ, and G9a were amplified with the primers listed in

Table S1 using PCR on the cDNA library, which was generated by the Kazusa

ORFeome Project and Genome Network Project Clones by RIKEN BRC. The

fragments were then inserted into the entry vector (pENTR™ 4 Dual Selection Vector;

#A10465, Thermo Fisher Scientific) between the *Xmn*I (#R0194S, New England

BioLabs [NEB]) and *EcoR*V (#R0195S, NEB) sites using the In-Fusion HD Cloning Kit

(#Z9648N, Takara Bio). The entry plasmids pENTR4XE153, pENTR4XE127,

pENTR4ΔCENPB, pENTR4HP1α, pENTR4HP1β, pENTR4HP1γ, and pENTR4G9a

were then obtained, which comprised open reading frames encoding ZNF518A,

ZNF518B, CENP-B$^{133\text{-}599}$, HP1α, HP1β, HP1γ, and G9a, respectively.[46–50]

To generate derivatives of the target genes, DNA fragments encoding

ZNF518A derivatives (ZNF518A^{KSS}, ZNF518A$^{1\text{-}970}$, ZNF518A$^{971\text{-}1483}$, ZNF518A$^{971\text{-}}$

^{1483KSS}, ZNF518A$^{971\text{-}1321}$, ZNF518A$^{1317\text{-}1483}$, and ZNF518A$^{1317\text{-}1483KSS}$), ZNF518B

derivatives (ZNF518B$^{KSS/N}$, ZNF518B$^{KSS/C}$, ZNF518B$^{KSS/NC}$, ZNF518B$^{1\text{-}398}$, ZNF518B$^{1\text{-}}$

^{628}, ZNF518B$^{344\text{-}628}$, ZNF518B$^{344\text{-}1074}$, ZNF518B$^{344\text{-}987}$, ZNF518B$^{344\text{-}877}$, ZNF518B$^{344\text{-}769}$,

ZNF518B$^{555\text{-}985}$, ZNF518B$^{555\text{-}877}$, ZNF518B$^{555\text{-}769}$, ZNF518B$^{878\text{-}1074}$, ZNF518B$^{878\text{-}987}$,

ZNF518B$^{770\text{-}987}$, ZNF518B$^{770\text{-}987KSS/C}$, and ZNF518B$^{770\text{-}877}$), HP1β derivatives

(HP1$\beta^{N(1\text{-}92)}$ and HP1$\beta^{C(93\text{-}185)}$), and G9a derivatives (G9a$^{1\text{-}92}$ and G9a$^{93\text{-}185}$) were

amplified using PCR on pENTR4XN153, pENTR4XN127, pENTR4HP1β, and

pENTR4G9a, respectively, with the primers listed in Table S1.

To overexpress the target proteins in human or *E. coli* cells, plasmids were

generated by exploiting DNA recombination between the above-mentioned entry

plasmids and pDEST131NGFP (to express EGFP fusion proteins),[51] pDEST152JTY

(to express EYFP-TetR fusion proteins), pDEST155Halo (to express Halo-tagged

proteins), pDEST174HATMBP (to express HAT [KDHLIHNVHKEEHAHNK]-MBP

fusion proteins),[52] or pDEST172HGST (to express His_6-GST fusion proteins using the

Gateway LR Clonase II Enzyme mix [#11791020, Thermo Fisher Scientific]).

To introduce CRISPR-mediated double-strand break sites, 54-bp DNA

fragments were generated by annealing two complementary single-stranded

oligonucleotides. They were then inserted into the *Age*I (#R3552S, NEB) site of the

pTORA14 vector using the In-Fusion HD Cloning Kit.[53] This yielded the

pTORA14AAVS1, pTORA14CENP-B_C, pTORA14HA4, pTORA14HB7, and

pTORA14HG1 plasmids, which were then used to induce DNA cleavage at the adeno-

associated virus integration site 1 region of the *CENPB, HP1ALPHA, HP1BETA*, and

HP1GAMMA genes, respectively. The DNA sequences targeted by the guide RNAs

are listed in Table S2.

To construct the plasmid directing homologous recombination (HR) at the 3′

end of the *CENPB* gene, the DNA fragment was amplified via PCR on human genomic

DNA by using Tks Gflex DNA polymerase (#R060A, Takara Bio) with the primers listed

in Table S1. This yielded the 584-bp DNA fragment containing the 267-bp left

homological arm (HRL) and 314-bp right homological arm (HRR) regions identical to

the target locus. The amplified DNA fragment comprising the HRL and HRR regions

was cloned into pCR-Blant (#K275020, Thermo Fisher Scientific). To construct the HR

donor plasmids (pKANA1CENPB and pKANA2CENPB), the mAID cassettes

containing the mAID-encoding region and either the hygromycin or blasticidin

resistance genes were amplified using PCR on the template plasmid pMK287 or

pMK288 and were inserted between the HRL and HRR regions using the In-Fusion

HD Cloning Kit.[43] The resultant plasmids were then used to generate the human cell

line expressing the CENP-B:mAID fusion protein.

Cell lines

A HeLa-Int-03 human cell line containing approximately 15,000 repeats of the *tetO*

sequence in chromosome 21 with and without CENP-B or HP1 KO was obtained as

previously described.[17,37,54] A U2OS-*os*TIR1 cell line, which is the parental cell line of

the CENP-B cKO cells, was generated by transfecting the U2OS cells with

pTORA14AAVS1 and pMK243 using Lipofectamine LTX Reagent (#A12621, Thermo

Fisher Scientific).[43] The clone was selected by maintaining the culture with 1 µg/mL of

puromycin (#10-2100, Focus Biomolecules). The CENP-B cKO cell line was

generated by transfecting the U2OS-*os*TIR1 cell line with pTORA14CENP-B_C,

pKANA1CENPB, and pKANA2CENPB. The clone was selected by maintaining the

culture with 1 mg/mL of hygromycin B (#089-06151, Wako) and 5 µg/mL of blasticidin

S (#ant-bl-05, InvivoGen).

Cell culture

U2OS and HeLa cells were cultured in Dulbecco's modified Eagle's medium (DMEM;

#044-29765, FUJIFILM Wako) supplemented with 10% (v/v) fetal bovine serum

(Bovogen Biologicals), 100 U/mL of penicillin, and 100 µg/mL of streptomycin (#168-

23191, Wako) at 37 °C in a humidified incubator maintained at 5% CO_2. To generate

CENP-B^{OFF} cells, CENP-B conditional KO (cKO) cells were cultured with 0.2 µg/mL of

Dox (#D3447, Sigma-Aldrich) for 24 h and subsequently with 0.5 mM of IAA (#094-

07121, Wako) for 24 h prior to experimentation.

Indirect immunofluorescence and microscopy

The U2OS and HeLa cells at the exponential growth phase were seeded onto

coverslips at 30–40% confluence and allowed to grow overnight. To overexpress

EGFP-ZNF518A and EGFP-ZNF518B, pDEST131NEGFP153 and

pDEST131NEGFP127, respectively, were transfected into cells using Lipofectamine

LTX for indirect immunofluorescence experiments. Twenty-four hours after

transfection, the cells were fixed with 4% (v/v) paraformaldehyde (#163-20145, Wako)

in phosphate-buffered saline (PBS) for 10 min and permeabilized with 0.15% (v/v)

Triton X-100 (#1610407, Bio-Rad Laboratories) in PBS for 2 min. The cells on the

coverslips were treated with 1% (v/v) bovine serum albumin (BSA; #A7284, Sigma-

Aldrich) in PBS and subsequently with primary antibodies, including 1:200-diluted

mouse anti-CENP-A (#ab13939, Abcam), 1:50,000-diluted mouse anti-CENP-B

(5E6C1),[16] 1:200-diluted rabbit anti-CENP-B (#ab25734, Abcam), 1:600-diluted rabbit

anti-CENP-C (#554),[55] 1:1,000-diluted mouse anti-HP1α (#MAB3584, Merck

Millipore), 1:800-diluted rabbit anti-HP1β (#8676 D2F2, Cell Signaling Technology),

1:3,000-diluted mouse anti-HP1γ (#MAB3450 2MOD-1G6, Merck Millipore), or

1:1,000-diluted rabbit anti-histone H3K9me3 (#ab8898, Abcam). The cells were

washed three times with PBS for 5 min and incubated with 1:600-diluted Alexa-

conjugated secondary antibodies. The DNA was then stained with 0.1 µg/mL of DAPI

(#H-1200, Vector Laboratories).

Confocal images were obtained using an Olympus FV1000 microscope, which

is based on an IX81 confocal microscope system, with a UPlanSApo 60×/1.35 oil-

immersion objective lens (Olympus) and FV10-SAW2.1 software (Olympus). Images

were also obtained by the Nikon Confocal A1Rsi system, which is based on a Ti-E

inverted microscope system, with a CFI Plan Apochromat VC 60×/1.40 oil-immersion

objective lens (Nikon) and FNIS-Elements AR software (Nikon). The images were

acquired at 0.5-µm intervals in the z-axis and Kalman-filtered to suppress background

noise.

FMIT

The pDEST172JTY and pDEST155Halo plasmids encoding EYFP-TetR and Halo

fusion proteins, respectively, were transfected into HeLa cells carrying the *tetO* array

at 60% confluence using Lipofectamine 3000 (#L3000001, Thermo Fisher Scientific).

Twenty-four hours after the transfection, the cells were further incubated with 1 nM of

Halo Tag TMR Ligand (G825A, Promega) at 37 °C for 2 h. The labeled cells were

washed with DMEM and incubated with the same medium at 37 °C for 30 min. The

cells were subsequently fixed with 4% (v/v) paraformaldehyde in PBS for 10 min, and

the DNA was stained with 0.1 µg/mL of DAPI. The EYFP and TMR signals were

observed using the Olympus BX53 fluorescence microscope system with a

UPlanSApo 60×/1.35 oil-immersion objective lens and cellSens standard software

(Olympus). More than 100 cells were analyzed for each sample.

Quantification of microscopic data

Fluorescence quantitation and line plot analyses of the fluorescence signals were

performed using ImageJ 1.50e.[56] To quantitate the fluorescent signals, z-stack images

were captured at every 500 nm between the lower and upper edges of the nuclei and

merged using the maximum intensity z-projection function in ImageJ. In the projected

images, nuclei and centromeres were defined by DAPI and CENP-A signals,

respectively. The nucleus- and centromere-associated GFP signals in the respective

cells were then estimated, and a ratio between them was calculated to determine how

GFP-fused proteins are enriched at centromeres. Statistical analyses were performed

using R statistical software 3.5.1 (R Foundation for Statistical Computing, http://www.r-

project.org).

For the line plot analyses, a specific slice showing the strongest signal of a

centromeric protein (CENP-A or CENP-B) and GFP-ZNF518 was selected from the z-

stack images. Using the RGB plot profile function in ImageJ, the signal intensity of the

centromeric protein and GFP-ZNF518 was quantified for each pixel along the indicated

lines.

Protein purification

The target proteins, His$_6$-GST-HP1 and HAT-MBP-ZNF518B$^{770-987}$, were purified from

BL21 *E. coli* cells. To express the target proteins, *E. coli* cells carrying either

pDEST173HGSTHP1 for His$_6$-GST-HP1 or pDEST174HATMBP127 for HAT-MBP-

ZNF518B$^{770-987}$ were cultured at 30 °C for 2 h after adding IPTG (#096-05143, Wako)

to L broth at a final concentration of 0.1 mM. A cell extract was prepared using a protein

purification buffer containing 50 mM of Tris-HCl (pH 8.0; #201-06273, Wako), 500 mM

of NaCl (#191-01665, Wako), 10 mM of β-mercaptoethanol (#131-14571, Wako), and

10% glycerol (#075-00611, Wako) for the His$_6$-GST-HP1 protein. For the HAT-MBP-

ZNF518B$^{770-987}$ protein, a cell extract was prepared using the protein purification buffer

with 1% Tween 20 (#H5152, Promega). The target proteins were then affinity-purified

using Ni-NTA agarose (#143-09763, Wako).

MBP PD

All the procedures in the MBP PD experiments were performed with a PD buffer [25

mM of Tris-HCl (pH 8.0), 100 mM of NaCl, 1 mM of EDTA (#E5134, Sigma-Aldrich), 1

μM of DTT (#GE17-1318-02, Sigma-Aldrich), 1% Tween 20, 10% glycerol, and 0.02%

BSA]. First, 2 μg of HAT-MBP-ZNF518B$^{770-987}$ and 5 μg of His$_6$-GST-HP1 proteins,

purified as described in "Protein purification," were mixed and incubated with rotation

at 4 °C for 4 h. Next, 50 µg of amylose magnetic beads (#E8035S, NEB) were added

to the protein mixture and further incubated with rotation at 4 °C for 1 h. After washing

the beads six times with 1 mL of the PD buffer, the sample was boiled with 1× SDS

loading buffer at 95 °C for 5 min. The eluate was then subjected to immunoblotting.

Immunoblotting

Immunoblotting was carried out using primary antibodies, including 1:200-diluted

mouse anti-MBP (#sc-271524, Santa Cruz Biotechnology) and mouse anti-GST (#sc-

138, Santa Cruz Biotechnology), 1:1,000-diluted rabbit anti-HP1α (#2616, CST),

mouse anti-HP1α (#MAB3584, Merck Millipore), 1:1,000-diluted rabbit anti-HP1β

(#8676, CST), 1:1,000-diluted rabbit anti-HP1γ (#2619, CST), 1:1,000-diluted and

mouse anti-HP1γ (#MAB3450, Merck Millipore), and 1:10,000-diluted mouse anti-

GAPDH (#5A12,016-25523, Wako). The target proteins were then detected using

1:15,000-diluted IRDye 800CW donkey anti-rabbit IgG and 1:15,000-diluted anti-

mouse IgG (#926-32213, #926-32210, Li-COR Biosciences) secondary antibodies.

Amino acid sequence analysis

The Basic Local Alignment Search Tool (BLAST) was used to identify similarities

between query and amino acid sequences deposited in the DDBJ/GenBank/EMBL

databases.[57] A phylogenetic tree was constructed by the neighbor-joining method

using MEGA X software [58].

Acknowledgments

The authors thank Iyo Toramoto, Kana Yamazoe, and Fusako Kawasaki (Kochi University) for their technical assistance; Koichiro Otake (Kazusa DNA Research Institute) for providing preliminary data; Koichi Honke, Hideaki Kuge, Kaoru Miyahara, Arisa Miyagawa-Yamaguchi, and Tatsuyuki Yamashita (Kochi University) for their scientific advice; and William C Earnshaw (University of Edinburgh) for critically reading the manuscript. This work was supported by grants from JSPS KAKENHI (grant number 18K06061), the Kato Memorial Bioscience Foundation, and the Takeda Science Foundation to SO. This work was also supported by JST, CREST (grant number 18070874), partly by MEXT KAKENHI (grant number 20H00473), and by the Kazusa DNA Research Institute Foundation to HM.

Author contributions

SO, JO, and HM designed the study. SO, JO, and NS performed the experiments. SO, JO, and HM analyzed the data. SO, JO, KN, and HM wrote the manuscript.

Disclosure and competing interests statement

The authors declare that they have no conflicts of interest.

Figure Legends

Figure 1. ZNF518B and its homolog ZNF518A localize to centromeres in a CENP-

B-dependent manner.

(A, C, and E) Co-localization of GFP-ZNF518B (green) and the centromeric factors

[red; CENP-A **(A),** CENP-B **(C),** or CENP-C **(E)**]. Areas marked by the open boxes in

the left images are enlarged in the right panel. Scale bars indicate 5 μm.

(B, D, and F) Line plot analyses were carried out along the white lines in panels A, C,

and E for the quantification of GFP-ZNF518B (green) and the three centromeric factors

[red; CENP-A **(B)** CENP-B **(D)** and CENP-C **(F)**].

(G) Localization of GFP-ZNF518B (green) and CENP-A (red) in WT (upper) and

CENP-B knockout cells (lower). The areas marked by the open boxes in the left panels

are enlarged in the right panels.

(H) Quantification of the fluorescence intensities of GFP-ZNF518B signals at

centromeres in WT, CENP-B knockout (KO) [see panel **(G)**], and CENP-B KO cells

overexpressing CENP-B-Flag$_3$ [CENP-B KO + CENP-B-FL$_3$ OX; see panel **(I)**]. The y-

axis shows the relative fluorescence intensities of GFP-ZNF518B signals associated

with centromeres against total GFP signals in the nucleus. Open circles represent

measurements from the individual cells shown in panels G and I. Red bars show

medians. ***$p < 0.005$.

(I) Co-localization of GFP-ZNF518B (green), CENP-A (red), and CENP-B (blue) in

CENP-B KO cells overexpressing CENP-B-Flag$_3$ (CENP-B KO + CENP-B-FL$_3$ OX).

The area marked by the open box in the left panel is enlarged in the right panel.

(J) Co-localization of Halo-ZNF518A (red), GFP-ZNF518B (green), and CENP-A (blue). The area marked by the open box in the left panel is enlarged in the right panel.

Figure 2. Dissection of ZNF518B to identify its protein segments required for centromeric localization.

Localization of the ZFN518B derivatives at centromeres. The schematic representation of the respective ZNF518B derivatives is drawn approximately to scale. Yellow boxes indicate zinc-finger motifs. The two PxV(I)xL motifs (purple boxes) were substituted with KxSxS (light purple boxes), shown as KSS/N and KSS/C. KSS/NC indicates that both PxV(I)xLs motifs were substituted with KxSxS. GFP-fused ZNF518B signals (WT and 18 derivatives; green) were co-visualized with the CENP-A or CENP-B centromeric markers (left images). The two images (ZNF518B$^{KSS/NC}$ and ZNF518B$^{878-987}$) labeled with "A" show CENP-A foci, whereas the remaining images represent CENP-B signals. The areas marked by the open boxes in the left images are enlarged in the right panel. Numbers and colors (left) reflect % frequencies of cells with GFP-ZNF518B signals at pericentromeres. Scale bar indicates 5 μm.

Figure 3. The PxIxL motif in ZNF518B is required for its interaction with the chromo-shadow domain of HP1.

(A) Schematic diagram of the FMIT assay for investigating the interaction between ZNF518B and HP1. CENP-B KO cells were employed in this assay.

(B) Tethering YFP-ZNF518B (green) to the *tetO* repeats results in the accumulation of Halo-HP1β (top; red) but not Halo-HP1γ (bottom; red) at the locus.

(C) FMIT assay to examine the interaction between ZNF518B derivatives and HP1. Cells expressing TetR-EYFP-ZNF518B (WT or derivative) and Halo-HP1 were subjected to the FMIT assay. The ZNF518B segment predicted to interact with HP1 is indicated by shaded boxes. Numbers and colors reflect % frequencies of the co-localization of TetR-EYFP-ZNF518B with full-length HP1 (HP1α, HP1β, and HP1γ), and the chromo domain (CD) and chromo-shadow domain (CSD) of HP1β.

(D) Experimental design to examine whether ZNF518B is required to maintain HP1β localization at the *tetO* repeats. The TetR-EYFP-ZNF518 quickly dissociated from the *tetO* repeats in the presence of doxycycline (Dox). CENP-B KO cells were employed in this assay.

(E) Cells carrying TetR-EYFP-ZNF518 (WT or derivatives) and Halo-HP1β were subjected to the FMIT assay as explained in panel D.

(F) Pull-down (PD) experiments to examine the physical interactions between ZNF518B and HP1. Purified recombinant GST-HP1 (input) was incubated with MBP-ZNF518B$^{770-987}$, and MBP PD samples were subjected to immunoblotting using anti-GST. Signal intensities in the indicated samples were normalized by the intensities of the samples with MBP alone.

(G) A model depicting the central roles of ZNF518A and ZNF518B in pericentromeric heterochromatin formation via their interactions with CENP-B, HP1, and G9a. Molecules and arrows in color represent the reaction steps indicated in this figure.

Figure 4. Dissection of ZNF518B relative to its interaction with CENP-B.

(A) Schematic representation of the GFP-ZNF518B localization assay, where CENP-B was depleted by adding indole-3-acetic acid (IAA) at different times. Data are shown in panel B.

(B) Representative images of GFP-ZNF518B-transfected cells quantified in panels C and D. Localization of GFP-ZNF518B (green) relative to centromeres in CENP-B^{ON}, CENP-B^{OFF}, and CENP-B^{4h-OFF} cells. CENP-A (red), CENP-B (blue), and DAPI staining are shown. The areas marked by the open boxes in the left images are enlarged in the right panel.

(C) Quantification of the fluorescence intensities of total CENP-B signals in CENP-B^{ON}, CENP-B^{OFF}, and CENP-B^{4h-OFF} cells based on the images in panel B. ***$p < 0.005$, and red bars show medians. CENP-B signals in nuclei were measured.

(D) Quantification of the fluorescence intensities of GFP-ZNF518B signals at the pericentromeres in CENP-B^{ON}, CENP-B^{OFF}, and CENP-B^{4h-OFF} cells based on the images shown in panel B. The y-axis shows the relative fluorescence intensities of GFP-ZNF518B signals associated with pericentromeres against total GFP signals in nuclei. Open circles represent measurements from the individual cells shown in panel B, and red bars show medians. **$p < 0.05$.

(E) Percentage of GFP-ZNF518B foci co-localized with centromeres (CENP-A foci) in CENP-B^{ON}, CENP-B^{OFF}, and CENP-B^{4h-OFF} cells. The images in panel B were subjected to this analysis.

(F) Schematic diagram of the FMIT assay for investigating the interaction between ZNF518B and CENP-B. The CENP-B$^{133-599}$ derivative was used as it cannot bind to the α-satellite repeats at pericentromeres by itself.

(G) FMIT assay to quantify the co-localization between TetR-EYFP-ZNF518 and Halo-

CENP-B$^{133-599}$. Numbers and colors reflect % frequencies of Halo-CENP-B$^{133-599}$

recruitment onto TetR-EYFP-ZNF518B foci. Red shaded boxes represent the

ZNF518B segment predicted to be involved in the interaction with CENP-B$^{133-599}$.

(H) A model depicting the central roles of ZNF518A and ZNF518B in pericentromeric

heterochromatin formation via their interactions with CENP-B, HP1, and G9a.

Molecules and arrows in color represent the reaction steps indicated in the present

figure.

Figure 5. Dissection of ZNF518A and ZNF518B relative to their requirement for

the loading of G9a to centromeres.

(A) FMIT assay to quantify the co-localization between TetR-EYFP-ZNF518 (WT and

16 derivatives) and Halo-G9a. Green shaded boxes represent the ZNF518B segment

potentially involved in its interaction with G9a.

(B) FMIT assay to quantify the co-localization between TetR-EYFP-ZNF518A (WT and

seven derivatives) and Halo-G9a. Green shaded boxes represent the ZNF518A

segment potentially involved in its interaction with G9a.

(C) FMIT assay to quantify the co-localization between Halo-G9a (WT and four

derivatives) and either TetR-EYFP-ZNF518A or TetR-EYFP-ZNF518B.

(D) Co-visualization of Halo-G9a (red) and GFP-ZNF518B (WT and two derivatives)

or GFP alone (green) in WT cells. The areas marked by the open boxes in the left

images are enlarged in the right panel. Scale bar indicates 5 μm.

(E) Localization of Halo-G9a (red) and GFP-ZNF518B^{WT} (green) in CENP-B KO cells.

Scale bar indicates 5 µm.

(F) A model depicting the central roles of ZNF518A and ZNF518B in pericentromeric

heterochromatin formation via their interactions with CENP-B, HP1, and G9a (see

Discussion for details).

REFERENCES

1. Molina, O., Abad, M.A., Solé, F., and Menéndez, P. (2021). Aneuploidy in cancer: lessons from acute lymphoblastic leukemia. Trends Cancer *7*, 37–47. 10.1016/j.trecan.2020.08.008.

2. Barra, V., and Fachinetti, D. (2018). The dark side of centromeres: types, causes and consequences of structural abnormalities implicating centromeric DNA. Nat Commun *9*, 4340. 10.1038/s41467-018-06545-y.

3. Cheeseman, I.M., and Desai, A. (2008). Molecular architecture of the kinetochore-microtubule interface. Nat. Rev. Mol Cell Biol *9*, 33–46. 10.1038/nrm2310.

4. McKinley, K.L., and Cheeseman, I.M. (2016). The molecular basis for centromere identity and function. Nat Rev Mol Cell Bio *17*, 16–29. 10.1038/nrm.2015.5.

5. Amor, D.J., Kalitsis, P., Sumer, H., and Choo, K.H.A. (2004). Building the centromere: from foundation proteins to 3D organization. Trends Cell Biol *14*, 359–368. 10.1016/j.tcb.2004.05.009.

6. Waye, J.S., and Willard, H.F. (1987). Nucleotide sequence heterogeneity of alpha satellite repetitive DNA: a survey of alphoid sequences from different human chromosomes. Nucleic Acids Res *15*, 7549–7569. 10.1093/nar/15.18.7549.

7. Schueler, M.G., Higgins, A.W., Rudd, M.K., Gustashaw, K., and Willard, H.F. (2001). Genomic and genetic definition of a functional human centromere. Science *294*, 109–115. 10.1126/science.1065042.

8. Schueler, M.G., and Sullivan, B.A. (2006). Structural and functional dynamics of human centromeric chromatin. Annu Rev Genom Hum *7*, 301–313. 10.1146/annurev.genom.7.080505.115613.

9. Black, B.E., and Cleveland, D.W. (2011). Epigenetic centromere propagation and

the nature of CENP-A nucleosomes. Cell *144*, 471–479. 10.1016/j.cell.2011.02.002.

10. Cleveland, D.W., Mao, Y., and Sullivan, K.F. (2003). Centromeres and

kinetochores from epigenetics to mitotic checkpoint signaling. Cell 112, 407–421.

10.1016/s0092-8674(03)00115-6.

11. Sullivan, B.A., and Karpen, G.H. (2004). Centromeric chromatin exhibits a

histone modification pattern that is distinct from both euchromatin and

heterochromatin. Nat Struct Mol Biol *11*, 1076–1083. 10.1038/nsmb845.

kinetochores: from epigenetics to mitotic checkpoint signaling. Cell *112*, 407–421.

10.1016/s0092-8674(03)00115-6.

12. Masumoto, H., Nakano, M., and Ohzeki, J. (2004). The role of CENP-B and α-

satellite DNA: de novo assembly and epigenetic maintenance of human

centromeres. Chromosome Res *12*, 543–556.

10.1023/b:chro.0000036593.72788.99.

13. Earnshaw, W.C., Sullivan, K.F., Machlin, P.S., Cooke, C.A., Kaiser, D.A., Pollard,

T.D., Rothfield, N.F., and Cleveland, D.W. (1987). Molecular cloning of cDNA for

CENP-B, the major human centromere autoantigen. J Cell Biol *104*, 817–829.

10.1083/jcb.104.4.817.

14. Ohzeki, J., Nakano, M., Okada, T., and Masumoto, H. (2002). CENP-B box is

required for de novo centromere chromatin assembly on human alphoid DNA. J Cell

Biol *159*, 765–775. 10.1083/jcb.200207112.

15. Okamoto, Y., Nakano, M., Ohzeki, J., Larionov, V., and Masumoto, H. (2007). A

minimal CENP-A core is required for nucleation and maintenance of a functional

human centromere. EMBO J *26*, 1279–1291. 10.1038/sj.emboj.7601584.

16. Okada, T., Ohzeki, J., Nakano, M., Yoda, K., Brinkley, W.R., Larionov, V., and

Masumoto, H. (2007). CENP-B controls centromere formation depending on the

chromatin context. Cell *131*, 1287–1300. 10.1016/j.cell.2007.10.045.

9. Cleveland, D.W., Mao, Y., and Sullivan, K.F. (2003). Centromeres and

17. Otake, K., Ohzeki, J., Shono, N., Kugou, K., Okazaki, K., Nagase, T.,

Yamakawa, H., Kouprina, N., Larionov, V., Kimura, H., et al. (2020). CENP-B creates

alternative epigenetic chromatin states permissive for CENP-A or heterochromatin

assembly. J Cell Sci *133*, jcs243303. 10.1242/jcs.243303.

18. Peters, A.H.F.M., O'Carroll, D., Scherthan, H., Mechtler, K., Sauer, S., Schöfer,

C., Weipoltshammer, K., Pagani, M., Lachner, M., Kohlmaier, A., et al. (2001). Loss

of the Suv39h histone methyltransferases impairs mammalian heterochromatin and

genome stability. Cell *107*, 323–337. 10.1016/s0092-8674(01)00542-6.

19. Grewal, S.I.S., and Jia, S. (2007). Heterochromatin revisited. Nat Rev Genet *8*,

35–46. 10.1038/nrg2008.

20. Molina, O., Vargiu, G., Abad, M.A., Zhiteneva, A., Jeyaprakash, A.A., Masumoto,

H., Kouprina, N., Larionov, V., and Earnshaw, W.C. (2016). Epigenetic engineering

reveals a balance between histone modifications and transcription in kinetochore

maintenance. Nat Commun *7*, 13334. 10.1038/ncomms13334.

21. Du, J., and Patel, D.J. (2014). Structural biology-based insights into

combinatorial readout and crosstalk among epigenetic marks. Biochim Biophys Acta-

Gene Regul Mech *1839*, 719–727. 10.1016/j.bbagrm.2014.04.011.

22. Hyun, K., Jeon, J., Park, K., and Kim, J. (2017). Writing, erasing and reading

histone lysine methylations. Exp Mol Med *49*, e324–e324. 10.1038/emm.2017.11.

23. Martin, C., and Zhang, Y. (2005). The diverse functions of histone lysine

methylation. Nat Rev Mol Cell Bio *6*, 838–849. 10.1038/nrm1761.

24. Canzio, D., Chang, E.Y., Shankar, S., Kuchenbecker, K.M., Simon, M.D.,

Madhani, H.D., Narlikar, G.J., and Al-Sady, B. (2011). Chromodomain-mediated

oligomerization of HP1 suggests a nucleosome-bridging mechanism for

heterochromatin assembly. Mol Cell *41*, 67–81. 10.1016/j.molcel.2010.12.016.

25. Rea, S., Eisenhaber, F., O'Carroll, D., Strahl, B.D., Sun, Z.-W., Schmid, M.,

Opravil, S., Mechtler, K., Ponting, C.P., Allis, C.D., et al. (2000). Regulation of

chromatin structure by site-specific histone H3 methyltransferases. Nature *406*, 593–

599. 10.1038/35020506.

26. Aagaard, L., Laible, G., Selenko, P., Schmid, M., Dorn, R., Schotta, G., Kuhfittig,

S., Wolf, A., Lebersorger, A., Singh, P.B., et al. (1999). Functional mammalian

homologues of the Drosophila PEV-modifier Su(var)3-9 encode centromere-

associated proteins which complex with the heterochromatin component M31.

EMBO J *18*, 1923–1938. 10.1093/emboj/18.7.1923.

27. Muramatsu, D., Kimura, H., Kotoshiba, K., Tachibana, M., and Shinkai, Y.

(2016). Pericentric H3K9me3 formation by HP1 interaction-defective histone

methyltransferase Suv39h1. Cell Struct Funct *41*, 145–152. 10.1247/csf.16013.

28. Shirai, A., Kawaguchi, T., Shimojo, H., Muramatsu, D., Ishida-Yonetani, M.,

Nishimura, Y., Kimura, H., Nakayama, J.-I., and Shinkai, Y. (2017). Impact of nucleic

acid and methylated H3K9 binding activities of Suv39h1 on its heterochromatin

assembly. eLife *6*. 10.7554/eLife.25317.

29. Tachibana, M., Sugimoto, K., Fukushima, T., and Shinkai, Y. (2001). SET

domain-containing protein, G9a, is a novel lysine-preferring mammalian histone

methyltransferase with hyperactivity and specific selectivity to lysines 9 and 27 of

histone H3. J Biol Chem *276*, 25309–25317. 10.1074/jbc.M101914200.

30. Fritsch, L., Robin, P., Mathieu, J.R.R., Souidi, M., Hinaux, H., Rougeulle, C.,

Harel-Bellan, A., Ameyar-Zazoua, M., and Ait-Si-Ali, S. (2010). A subset of the

histone H3 lysine 9 methyltransferases Suv39h1, G9a, GLP, and SETDB1

participate in a multimeric complex. Mol Cell *37*, 46–56.

10.1016/j.molcel.2009.12.017.

31. Ueda, J., Tachibana, M., Ikura, T., and Shinkai, Y. (2006). Zinc finger protein

WIZ links G9a/GLP histone methyltransferases to the co-repressor molecule CtBP. J

Biol Chem *281*, 20120–20128. 10.1074/jbc.M603087200.

32. Maier, V.K., Feeney, C.M., Taylor, J.E., Creech, A.L., Qiao, J.W., Szanto, A.,

Das, P.P., Chevrier, N., Cifuentes-Rojas, C., Orkin, S.H., et al. (2015). Functional

proteomic analysis of repressive histone methyltransferase complexes reveals

ZNF518B as a G9A regulator. Mol Cell Proteom *14*, 1435–1446.

10.1074/mcp.M114.044586.

33. Ohta, S., Bukowski-Wills, J.-C., Sanchez-Pulido, L., Alves, F. de L., Wood, L.,

Chen, Z.A., Platani, M., Fischer, L., Hudson, D.F., Ponting, C.P., et al. (2010). The

protein composition of mitotic chromosomes determined using multiclassifier

combinatorial proteomics. Cell *142*, 810–821. 10.1016/j.cell.2010.07.047.

34. Ohta, S., Montaňo-Gutierrez, L.F., Alves, F. de L., Ogawa, H., Toramoto, I., Sato,

N., Morrison, C.G., Takeda, S., Hudson, D.F., Rappsilber, J., et al. (2016).

Proteomics analysis with a nano random Forest approach reveals novel functional

interactions regulated by SMC complexes on mitotic chromosomes. Mol Cell

Proteom *15*, 2802–2818. 10.1074/mcp.M116.057885.

35. Canzio, D., Larson, A., and Narlikar, G.J. (2014). Mechanisms of functional

promiscuity by HP1 proteins. Trends Cell Biol 24, 377–386.

10.1016/j.tcb.2014.01.002.

36. Ohzeki, J., Shono, N., Otake, K., Martins, N.M.C., Kugou, K., Kimura, H.,

Nagase, T., Larionov, V., Earnshaw, W.C., and Masumoto, H. (2016).

KAT7/HBO1/MYST2 regulates CENP-A chromatin assembly by antagonizing

Suv39h1-mediated centromere inactivation. Dev Cell *37*, 413–427.

10.1016/j.devcel.2016.05.006.

37. Ohzeki, J., Bergmann, J.H., Kouprina, N., Noskov, V.N., Nakano, M., Kimura, H.,

Earnshaw, W.C., Larionov, V., and Masumoto, H. (2012). Breaking the HAC Barrier:

histone H3K9 acetyl/methyl balance regulates CENP-A assembly. EMBO J *31*,

2391–2402. 10.1038/emboj.2012.82.

38. Bergmann, J.H., Rodríguez, M.G., Martins, N.M.C., Kimura, H., Kelly, D.A.,

Masumoto, H., Larionov, V., Jansen, L.E.T., and Earnshaw, W.C. (2011). Epigenetic

engineering shows H3K4me2 is required for HJURP targeting and CENP-A

assembly on a synthetic human kinetochore. EMBO J *30*, 328–340.

10.1038/emboj.2010.329.

39. Martins, N.M.C., Bergmann, J.H., Shono, N., Kimura, H., Larionov, V.,

Masumoto, H., and Earnshaw, W.C. (2016). Epigenetic engineering shows that a

human centromere resists silencing mediated by H3K27me3/K9me3. Mol Biol Cell

27, 177–196. 10.1091/mbc.e15-08-0605.

40. Shono, N., Ohzeki, J., Otake, K., Martins, N.M.C., Nagase, T., Kimura, H.,

Larionov, V., Earnshaw, W.C., and Masumoto, H. (2015). CENP-C and CENP-I are

key connecting factors for kinetochore and CENP-A assembly. J Cell Sci *128*, 4572–

4587. 10.1242/jcs.180786.

41. Thiru, A., Nietlispach, D., Mott, H.R., Okuwaki, M., Lyon, D., Nielsen, P.R.,

Hirshberg, M., Verreault, A., Murzina, N.V., and Laue, E.D. (2004). Structural basis

of HP1/PXVXL motif peptide interactions and HP1 localisation to heterochromatin.

EMBO J *23*, 489–499. 10.1038/sj.emboj.7600088.

42. Nozawa, R.-S., Nagao, K., Masuda, H.-T., Iwasaki, O., Hirota, T., Nozaki, N.,

Kimura, H., and Obuse, C. (2010). Human POGZ modulates dissociation of HP1α

from mitotic chromosome arms through Aurora B activation. Nat Cell Biol *12*, 719–

727. 10.1038/ncb2075.

43. Natsume, T., Kiyomitsu, T., Saga, Y., and Kanemaki, M.T. (2016). Rapid protein

depletion in human cells by auxin-inducible degron tagging with short homology

donors. Cell Rep *15*, 210–218. 10.1016/j.celrep.2016.03.001.

44. Simon, J.M., Parker, J.S., Liu, F., Rothbart, S.B., Ait-Si-Ali, S., Strahl, B.D., Jin,

J., Davis, I.J., Mosley, A.L., and Pattenden, S.G. (2015). A role for widely

interspaced zinc finger (WIZ) in retention of the G9a methyltransferase on chromatin. J Biol Chem *290*, 26088–26102. 10.1074/jbc.M115.654459.

45. Bian, C., Chen, Q., and Yu, X. (2015). The zinc finger proteins ZNF644 and WIZ regulate the G9a/GLP complex for gene repression. eLife *4*, e05606. 10.7554/eLife.05606.

46. Itoh, M., Yasunishi, A., Imamura, K., Kanamori-Katayama, M., Suzuki, H., Suzuki, M., Carninci, P., Kawai, J., and Hayashizaki, Y. (2006). Constructing ORFeome resources with removable termination codons. BioTechniques *41*, 44–50. 10.2144/000112209.

47. Nagase, T., Yamakawa, H., Tadokoro, S., Nakajima, D., Inoue, S., Yamaguchi, K., Itokawa, Y., Kikuno, R.F., Koga, H., and Ohara, O. (2008). Exploration of human ORFeome: high-throughput preparation of ORF clones and efficient characterization of their protein products. DNA Res *15*, 137–149. 10.1093/dnares/dsn004.

48. Kimura, K., Wakamatsu, A., Suzuki, Y., Ota, T., Nishikawa, T., Yamashita, R., Yamamoto, J., Sekine, M., Tsuritani, K., Wakaguri, H., et al. (2006). Diversification of transcriptional modulation: large-scale identification and characterization of putative alternative promoters of human genes. Gen Res *16*, 55–65. 10.1101/gr.4039406.

49. Otsuki, T., Ota, T., Nishikawa, T., Hayashi, K., Suzuki, Y., Yamamoto, J., Wakamatsu, A., Kimura, K., Sakamoto, K., Hatano, N., et al. (2005). Signal sequence and keyword trap in silico for selection of full-length human cDNAs encoding secretion or membrane proteins from oligo-capped cDNA libraries. DNA Res *12*, 117–126. 10.1093/dnares/12.2.117.

50. Ota, T., Suzuki, Y., Nishikawa, T., Otsuki, T., Sugiyama, T., Irie, R., Wakamatsu,

A., Hayashi, K., Sato, H., Nagai, K., et al. (2004). Complete sequencing and

characterization of 21,243 full-length human cDNAs. Nat Genet *36*, 40–45.

10.1038/ng1285.

51. Ohta, S., Hamada, M., Sato, N., and Toramoto, I. (2015). Polyglutamylated

tubulin binding protein C1orf96/CSAP is involved in microtubule stabilization in

mitotic spindles. PloS One *10*, e0142798. 10.1371/journal.pone.0142798.

52. Cℏaga, G., Bochkariov, D.E., Jokhadze, G.G., Hopp, J., and Nelson, P. (1999).

Natural poly-histidine affinity tag for purification of recombinant proteins on cobalt(II)-

carboxymethylaspartate crosslinked agarose. J Chromatogr A *864*, 247–256.

10.1016/s0021-9673(99)01008-0.

53. Ohta, S., Taniguchi, T., Sato, N., Hamada, M., Taniguchi, H., and Rappsilber, J.

(2019). Quantitative proteomics of the mitotic chromosome scaffold reveals the

association of baz1b with chromosomal axes. Mol Cell Proteom *18*, 169–181.

10.1074/mcp.ra118.000923.

54. Ruppert, J.G., Samejima, K., Platani, M., Molina, O., Kimura, H., Jeyaprakash,

A.A., Ohta, S., and Earnshaw, W.C. (2018). HP1α targets the chromosomal

passenger complex for activation at heterochromatin before mitotic entry. EMBO J

37. 10.15252/embj.201797677.

55. Ohta, S., Bukowski-Wills, J.-C., Wood, L., de Lima Alves, F., Chen, Z.,

Rappsilber, J., and Earnshaw, W.C. (2010). Proteomics of isolated mitotic

chromosomes identifies the kinetochore protein Ska3/Rama1. Cold Spring Harb

Symp Quant Biol *75*, 433–438. 10.1101/sqb.2010.75.022.

56. Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ:

25 years of image analysis. Nat Methods *9*, 671–675. 10.1038/nmeth.2089.

57. Altschul, S.F., Gish, W., Miller, W., Myers, E.W., and Lipman, D.J. (1990). Basic

local alignment search tool. J Mol Biol *215*, 403–410. 10.1016/S0022-

2836(05)80360-2.

58. Kumar, S., Stecher, G., Li, M., Knyaz, C., and Tamura, K. (2018). MEGA X:

molecular evolutionary genetics analysis across computing platforms. Mol Biol Evol

35, 1547–1549. 10.1093/molbev/msy096.