Title: Functional connectivity across the human subcortical auditory system using an
autoregressive matrix-Gaussian copula graphical model approach with partial correlations

Authors: Noirrit Kiran Chandra[1]*, Kevin R. Sitek[2]*, Bharath Chandrasekaran[2], Abhra Sarkar[3]
*=contributed equally

Affiliations: (1) The University of Texas at Dallas, Dept. of Mathematical Sciences; (2)
Northwestern University, Roxelyn and Richard Pepper Department of Communication Sciences
and Disorders; (3) The University of Texas at Austin, Dept. of Statistics and Data Sciences

Corresponding author:

Bharath Chandrasekaran (b.chandra@northwestern.edu)

Abhra Sarkar (abhra.sarkar@utexas.edu)

Abstract/Summary

The auditory system comprises multiple subcortical brain structures that process and refine
incoming acoustic signals along the primary auditory pathway. Due to technical limitations of
imaging small structures deep inside the brain, most of our knowledge of the subcortical auditory
system is based on research in animal models using invasive methodologies. Advances in ultra-
high field functional magnetic resonance imaging (fMRI) acquisition have enabled novel non-
invasive investigations of the human auditory subcortex, including fundamental features of
auditory representation such as tonotopy and periodotopy. However, functional connectivity
across subcortical networks is still underexplored in humans, with ongoing development of
related methods. Traditionally, functional connectivity is estimated from fMRI data with full
correlation matrices. However, partial correlations reveal the relationship between two regions
after removing the effects of all other regions, reflecting more direct connectivity. Partial

correlation analysis is particularly promising in the ascending auditory system, where sensory

information is passed in an obligatory manner, from nucleus to nucleus up the primary auditory

pathway, providing redundant but also increasingly abstract representations of auditory stimuli.

While most existing methods for learning conditional dependency structures based on partial

correlations assume independently and identically Gaussian distributed data, fMRI data exhibit

significant deviations from Gaussianity as well as high temporal autocorrelation. In this paper,

we developed an autoregressive matrix-Gaussian copula graphical model (ARMGCGM) approach

to estimate the partial correlations and thereby infer the functional connectivity patterns within

the auditory system while appropriately accounting for autocorrelations between successive

fMRI scans. Our results show strong positive partial correlations between successive structures

in the primary auditory pathway on each side (left and right), including between auditory

midbrain and thalamus, and between primary and associative auditory cortex. These results are

highly stable when splitting the data in halves according to the acquisition schemes and

computing partial correlations separately for each half of the data, as well as across cross-

validation folds. In contrast, full correlation-based analysis identified a rich network of

interconnectivity that was not specific to adjacent nodes along the pathway. Overall, our results

demonstrate that unique functional connectivity patterns along the auditory pathway are

recoverable using novel connectivity approaches and that our connectivity methods are reliable

across multiple acquisitions.

Keywords (6):

functional MRI, Gaussian copula graphical model, partial correlation, resting state connectivity,

subcortical auditory system, timeseries

1. Introduction

The mammalian auditory pathway conveys acoustic information between the inner ear—where sounds are mechanoelectrically transduced in the cochlea—and auditory cortex by way of multiple subcortical nuclei across the brainstem, midbrain, and thalamus. Much of our knowledge of the auditory system arises from anatomical and physiological research with non-human animal models (McIntosh and Gonzalez-Lima, 1991; Popper and Fay, 1992; Webster *et al.*, 1992). This work has contributed tremendously to our understanding of the mammalian auditory system. However, due to methodological and ethical limitations, our ability to directly assess auditory function in the human nervous system is severely constrained (as discussed in (Moerel *et al.*, 2021)).

This is particularly true for subcortical auditory structures. In mammals, the ascending central auditory pathway receives signals from the cochlea of the inner ear by way of the cochlear nerve, which principally innervates the cochlear nucleus in the brainstem. Auditory signals are then transmitted to the superior olivary complex, which is the first decussation point at which signals largely pass contralaterally from the left cochlear nucleus to the right superior olive (and similarly from right to left). From the superior olive, auditory signals travel (via the lateral lemniscus) to the inferior colliculus in the midbrain. The last subcortical auditory structure is the medial geniculate nucleus of the thalamus, which then passes information to primary auditory cortex. In addition to the ascending "lemniscal" auditory pathway, an equal number of efferent connections transmit top-down information from higher order auditory regions to earlier auditory structures (Malmierca and Ryugo, 2011; Winer, 2005). Due to the small size of the subcortical auditory structures—tightly packed in with other heterogeneous nuclei and white matter pathways—and their anatomical location deep within the cranium, the subcortical auditory structures have received limited attention in non-invasive human research.

Functional magnetic resonance imaging (fMRI) is the most popular non-invasive method for probing macroscopic network-related brain activity. While studies of the human subcortical auditory system are somewhat limited, previous task-based fMRI research has functionally localized the subcortical auditory structures (Sitek *et al.*, 2019), identified the tonotopic frequency mappings within the auditory midbrain and thalamus (De Martino *et al.*, 2013; Moerel

et al., 2015; Ress and Chandrasekaran, 2013), separated top-down and bottom-up speech-
selective subregions of auditory thalamus (Mihai *et al.*, 2019; Tabas *et al.*, 2021), and recorded
level-dependent BOLD signals throughout the auditory pathway (Hawley *et al.*, 2005; Sigalovsky
and Melcher, 2006).

In contrast to task-related BOLD activity, fMRI connectivity methods (often utilizing resting state
fMRI paradigms and full correlation analysis) are commonly used to assay cortical brain networks
(Biswal *et al.*, 1995; Gordon, Laumann, Adeyemo, *et al.*, 2017; Power *et al.*, 2011; Smith,
Beckmann, *et al.*, 2013), including the cortical auditory system (Abrams *et al.*, 2020; Cha *et al.*,
2016; Chen *et al.*, 2020; Eckert *et al.*, 2008; Maudoux *et al.*, 2012; Ren *et al.*, 2021). However,
fMRI connectivity methods have limited history in subcortical research, especially in the auditory
system, where they have—to our knowledge—only been utilized a handful of times to assess
connectivity differences between individuals with and without tinnitus percepts (Berlot *et al.*,
2020; Hofmeier *et al.*, 2018; Leaver *et al.*, 2016; Zhang *et al.*, 2015).

From the seminal resting state connectivity studies identifying human default mode and motor
networks (Biswal *et al.*, 1995; Fox and Raichle, 2007), to work linking functional connectivity with
behavioral variability (Baldassarre *et al.*, 2012; Deng *et al.*, 2016), to investigations into brain
network differences associated with disorders (Chai *et al.*, 2016; Di Martino *et al.*, 2011; Ferri *et*
al., 2018; Greicius *et al.*, 2007; Hahn *et al.*, 2011; Husain and Schmidt, 2014; Kaiser *et al.*, 2015;
Sitek *et al.*, 2016; Wilson *et al.*, 2022), full correlation analysis has contributed tremendously to
our understanding of human brain networks. However, moving beyond the traditional full
correlation analysis should enable greater specificity in assessing functional connectivity
patterns, particularly in identifying specific node-to-node connectivity patterns within an
established brain network (Marrelec *et al.*, 2006; Smith, 2012). In contrast to full correlations,
which represent both direct and indirect connections, partial correlation analyses represent the
direct association between two specific nodes after filtering out the effects of the remaining
nodes and thus hold great promise for estimating direct functional connectivity within a network
(Smith, Beckmann, *et al.*, 2013). (Please refer to the illustration in the Supplementary Materials.)
For these reasons, partial correlation approaches are increasingly used to study functional
connectivity networks in the brain (Wang *et al.*, 2016; Warnick *et al.*, 2018), including improved

identification of individualized connectivity profiles compared to full correlation methods
(Menon and Krishnamurthy, 2019), as well as improved prediction of brain disorders (Reeves *et*
al., 2023; Skåtun *et al.*, 2017; de Vos *et al.*, 2018) and identification of general cognitive ability
(Sripada *et al.*, 2021). However, we are unaware of the application of such methods to assess
functional connectivity within subcortical networks, particularly within the human auditory
system.

In this article, we build on the Bayesian precision factor model (PFM) introduced recently in
Chandra et al. (2021) to develop a novel highly robust autoregressive matrix-Gaussian copula
graphical model (ARMGCGM) to assess partial correlation-based functional connectivity in a
specific network in the human brain that spans subcortical and cortical regions, the auditory
system. The PFM decomposes the model precision matrix into a flexible low-rank and diagonal
structure, then exploits that to design very efficient estimation algorithms. However, it makes
the restrictive assumption that each variable is marginally Gaussian distributed. Several studies
in the literature also make this assumption (Yu *et al.*, 2022; Zhang *et al.*, 2014) which has the very
useful implication that a zero partial correlation between two variables (equivalent to a zero
entry in the precision matrix in the corresponding position) also means independence between
them after removing the effects of other variables. However, in many applications—including
ours—data are often not Gaussian distributed. Additionally, data from successive fMRI volumes
exhibit strong autocorrelation. The ARMGCGM extends the PFM to the case where the univariate
marginals can be any arbitrary distribution while also accounting for the autocorrelations
between successive fMRI scans. The association between the variables are modeled using a
Gaussian copula that implies conditional independence for zero partial correlation, allowing easy
interpretability of the conditional dependence graph. As developed, the ARMGCGM approach is
broadly applicable for studying undirected functional graphs using large-scale fMRI data.

We use the novel ARMGCGM to investigate functional connectivity between specific nodes
across the human auditory system. We used publicly available 7T resting state fMRI from over
one hundred participants to examine auditory connectivity. To probe connectivity within the
auditory system, we included auditory cortical regions of interest as well as subcortical auditory
regions derived from human histology (Sitek *et al.*, 2019). As the auditory pathway comprises a

chain of multiple subcortical structures, and due to the largely lateralized organization of the

lemniscal auditory system, we hypothesized that partial correlations would be greatest between

adjacent nodes in the same hemisphere, over and above the contributions from other auditory

(and non-auditory) regions of interest. In particular, because of the critical position of auditory

midbrain and thalamus as computational hubs involving bottom-up and top-down information

transfer (Mihai *et al.*, 2019; Tabas *et al.*, 2021), we hypothesized strong partial correlations

between inferior colliculus and medial geniculate. We further assessed reliability across

acquisitions by separately analyzing data with anterior–posterior and posterior–anterior phase-

encoding directions, as well as leave-10%-out cross-validation. We then compared our

ARMGCGM-based connectivity results with those from a full correlation approach as well as

alternative partial correlation methods. Overall, our consistent findings of hierarchical

connectivity within the auditory system using our novel partial correlation method—consistent

across data partitions and leave-10%-out validation— demonstrate the methodological reliability

of our ARMGCGM approach as well as the neurobiological organization of auditory structures in

the human primary auditory system.

2. Materials and methods

2.1 Magnetic resonance imaging acquisition and processing

We used resting state fMRI from 106 participants in the 7T Human Connectome Project (Elam *et*

al., 2021; Van Essen *et al.*, 2012). Specifically, we utilized the minimally preprocessed volumetric

data in common space (Glasser *et al.*, 2013). BOLD fMRI data were acquired with 1.6 mm isotropic

voxel size across four runs of a resting state paradigm (repetition time [TR] = 1 s, 900 TRs per

run). Two runs were acquired with anterior–posterior (AP) phase encoding, and two were

acquired with posterior–anterior (PA) phase encoding. For all ROIs and runs, we discarded the

first 50 TRs to increase stability.

For each individual and each run, we extracted mean timeseries from predefined regions of

interest (ROIs). Subcortical auditory ROIs were defined using the Sitek–Gulban atlas (Sitek *et al.*,

2019). Cortical ROIs were defined using FreeSurfer's implementation of the DKT atlas (Dale *et al.*,

1999; Klein and Tourville, 2012). For this study we used transverse temporal gyrus (TTG) and

superior temporal gyrus (STG) as auditory ROIs, as well as pericalcarine cortex (Calc) and superior

frontal gyrus (SFG) as non-auditory control ROIs (see Supplementary Materials). Mean timeseries

were extracted for each ROI using nilearn's [Masker] function.

2.2 Data partitioning and cross-validation

BOLD fMRI is prone to geometric distortions in the phase-encoding direction which can be largely

corrected using a variety of methods (Esteban *et al.*, 2021; Jezzard and Balaban, 1995). Adjacent

to motion-sensitive cerebrospinal fluid (CSF), the brainstem is particularly susceptible to such

geometric distortions. Although the HCP minimal preprocessing pipeline corrects for phase-

encoding distortions (Glasser *et al.*, 2013), to isolate the potential residual contribution of phase

encoding direction on functional connectivity estimates, we conducted separate analyses on data

collected with posterior–anterior (PA) phase encoding direction (runs 1 and 3) and anterior–

posterior (AP) phase encoding direction (runs 2 and 4) and compared the results. As the fMRI

data acquired in the two phases will be analyzed separately but using the same probability-

model, we use the same notations for the different phases to describe our proposed model in

the following sections.

We further evaluated our results using a leave-10%-out cross-validation (CV) approach. We

assessed the stability of the connectivity estimates by checking the correlation between the

results obtained from each acquisition scheme.

2.3 Autoregressive matrix-Gaussian copula graphical models

The Bayesian precision factor model (PFM), developed recently in (Chandra *et al.*, 2021),

provided a novel computationally efficient robust technique for estimating precision matrices.

Since partial correlations can be readily obtained from the precision matrix, the approach allowed

straightforward estimation of the underlying connectivity graphs. Previously, (Lee and Kim, 2021)

obtained the precision matrix by inverting the estimated covariance matrix. However, this

approach often tends to exhibit poor empirical performance (Pourahmadi, 2013).

Most partial correlation-based conditional dependency graph estimation procedures in the
statistical literature (Cai *et al.*, 2020; Chandra *et al.*, 2021; Friedman *et al.*, 2008; Warnick *et al.*,
2018) assume that the joint distribution of the data is independent and identically distributed
(iid) multivariate Gaussian, which implies that the univariate marginal distributions are also all
Gaussians. However, successive fMRI scans have strong autocorrelations, and their marginal
distributions exhibit substantial deviance from the Gaussian assumption. In this paper, we
therefore extend the PFM to accommodate non-Gaussian marginals, while appropriately
accounting for the temporal dependence between successive fMRI scans.

Let $Y_{t,j}^{(r,i)}$ be the fMRI timeseries corresponding to the i-th individual's j-th ROI at the t-th
timepoint in the r-th run. In our application we are interested in studying the connectivity
between $d = 12$ ROIs along the central auditory pathway using fMRI timeseries of length $T =$
850 from $N = 106$ individuals each undergoing $R = 2$ runs. We let $f_j^{(r,i)}$ be the (unknown)
marginal density of $Y_{t,j}^{(r,i)}$ with corresponding cumulative distribution function (CDF) $F_j^{(r,i)}$.

Copulas provide a broadly applicable class of tools that allow the joint distribution of $Y_{t,j}^{(r,i)}$ to be
flexibly characterized by first modeling the univariate marginals $f_j^{(r,i)}$ and then hierarchically
modeling their joint dependencies by mapping the $F_j^{(r,i)}\left(Y_{t,j}^{(r,i)}\right)$'s to a joint probability space. For
Gaussian copulas, this is done by setting $Z_{t,j}^{(r,i)} = \Phi^{-1}\left\{F_j^{(r,i)}\left(Y_{t,j}^{(r,i)}\right)\right\}$, where $\Phi(\cdot)$ is the CDF of a
standard Gaussian distribution. This implies marginally $Z_{t,j}^{(r,i)} \sim \text{N}(0,1)$ for all r, i, j, t, where
$\text{N}(\mu, \sigma^2)$ denotes a univariate Gaussian distribution with mean μ and variance σ^2. We let $\boldsymbol{Z}^{(r,i)} =$
$\left(\left(Z_{t,j}^{(r,i)}\right)\right)_{T \times d}$ denote the matrix of fMRI signals corresponding to the i-th individual in the r-th
run in the transformed Gaussian space. The Gaussian copula assumption on $F_{t,j}^{(r,i)}\left(Y_{t,j}^{(r,i)}\right)$'s
implies that the joint distribution of $\boldsymbol{Z}^{(r,i)}$ is Gaussian as well. The dependencies between the
$Z_{t,j}^{(r,i)}$'s are therefore characterized entirely by their correlations. Additionally, since the
dependence relationships between the observed $Y_{t,j}^{(r,i)}$'s are modeled only through $Z_{t,j}^{(r,i)}$'s, these
correlations also completely characterize the dependencies between the $Y_{t,j}^{(r,i)}$'s.

Note that probabilistic dependencies exist between $Z_{t,j}^{(r,i)}$'s across both j and t; dependence

across j incurs due to the interaction between the ROIs whereas dependence across t occurs due

to the temporal dependence between successive fMRI scans. We let $\boldsymbol{R}_\Omega$ denote the $d \times d$

correlation matrix accounting for the dependence across the d different ROIs across all runs and

individuals.

While our main interest lies in estimating these dependencies between the ROIs, it is also crucial

to consider the temporal dependence in the $Y_{t,j}^{(r,i)}$'s. Let $\boldsymbol{Z}_{\cdot,j}^{(r,i)} = \left(Z_{1,j}^{(r,i)}, \ldots, Z_{T,j}^{(r,i)}\right)^T$ be the j-th

column of $\boldsymbol{Z}^{(r,i)}$, i.e., the timeseries corresponding to the i-th individual's j-th ROI in the r-th run.

We develop our model in a hierarchical manner. To being with, we use autoregressive (AR)

processes of order L to model higher-order temporal dependencies in the $Z_{t,j}^{(r,i)}$'s as

$$Z_{t,j}^{(r,i)} = \sum_{t'=1}^{L} \beta_{t',j}^{(r,i)} Z_{t-t',j}^{(r,i)} + \epsilon_{t,j}^{(r,i)}, \qquad \epsilon_{1,j}^{(r,i)}, \ldots, \epsilon_{T,j}^{(r,i)} \overset{iid}{\sim} N\left(0, \varsigma_j^{2(r,i)}\right), \tag{1}$$

with $Z_{t-t',j}^{(r,i)} = 0$ if $t' \geq t$. We assume separate $(\boldsymbol{\beta}, \boldsymbol{\varsigma}^2)$ parameters across (r,i,j) in (1) to make
the model adapt to different timeseries patterns across different ROIs, individuals, and runs.

Let $\boldsymbol{R}_{\mathcal{T},j}^{(r,i)}$ be the correlation matrix of $\boldsymbol{Z}_{\cdot,j}^{(r,i)}$ induced by the AR(L) model in (1). We let $\boldsymbol{\xi}_{\cdot,j}^{(r,i)} = \left(\boldsymbol{R}_{\mathcal{T},j}^{(r,i)}\right)^{-\frac{1}{2}} \boldsymbol{Z}_{\cdot,j}^{(r,i)}$ be the AR corrected timeseries corresponding to the j-th ROI for $j = 1, \ldots, d$. We
define the $T \times d$ matrix $\boldsymbol{\Xi}^{(r,i)} = \left(\boldsymbol{\xi}_{\cdot,1}^{(r,i)} \quad \cdots \quad \boldsymbol{\xi}_{\cdot,d}^{(r,i)}\right) = \left(\boldsymbol{\xi}_{1,\cdot}^{(r,i)} \quad \cdots \quad \boldsymbol{\xi}_{T,\cdot}^{(r,i)}\right)^T$ where $\boldsymbol{\xi}_{t,\cdot}^{(r,i)}$ is the t-th row of $\boldsymbol{\Xi}^{(r,i)}$ and can be interpreted as the fMRI signals in the Gaussian copula space
subsequent to filtering out the temporal dependence at time point t. We then let

$$\boldsymbol{\xi}_{1,\cdot}^{(r,i)}, \ldots, \boldsymbol{\xi}_{T,\cdot}^{(r,i)} \overset{iid}{\sim} N(\boldsymbol{0}, \boldsymbol{R}_\Omega). \tag{2}$$

229   The formulations in (1)-(2) imply the following joint distribution on $\boldsymbol{Z}^{(r,i)}$

$$f\left(\boldsymbol{Z}^{(r,i)} \mid \boldsymbol{R}_\Omega, \boldsymbol{R}_{\mathcal{T},1}^{(r,i)}, \ldots, \boldsymbol{R}_{\mathcal{T},d}^{(r,i)}\right) = \frac{e^{-\frac{1}{2}\text{tr}\left(\boldsymbol{R}_\Omega^{-1}\boldsymbol{\Xi}^{(r,i)T}\boldsymbol{\Xi}^{(r,i)}\right)}}{(2\pi)^{\frac{dT}{2}}|\boldsymbol{R}_\Omega|^{\frac{T}{2}} \prod_{j=1}^{d}\left|\boldsymbol{R}_{\mathcal{T},j}^{(r,i)}\right|^{\frac{1}{2}}}. \tag{3}$$

Figure 1 Graphical illustration of the hierarchical structure of the autoregressive matrix-Gaussian copula graphical model (ARMGCGM): For brevity, we omit the superscript (r, i)—the run and subject indicators, respectively, in this illustration. $Y_{t,j}$ is the observed BOLD signal from the j-th ROI at the t-th time point. $Z_{t,j} = \Phi^{-1}\left(F_j\left(Y_{t,j}\right)\right)$ is the BOLD signal in the Gaussian copula space. Probabilistic dependencies exist between $Z_{t,j}$'s across both j and t; dependence across j incurs due to the interaction between the ROIs whereas dependence across t occurs due to the temporal dependence between successive fMRI scans. $\xi_{t,j}$'s are the autocorrelation corrected $Z_{t,j}$'s with $\left(\xi_{t,1}, \ldots, \xi_{t,d}\right)^T \overset{iid}{\sim} N_d(\mathbf{0}, \mathbf{R}_\Omega^{-1})$ for all $t = 1, \ldots, T.$

Although the $\boldsymbol{R}_{\mathcal{T},j}^{(r,i)}$'s are massive dimensional $T \times T$ matrices, notably they are characterized

entirely by the associated autoregressive parameters $\left\{\beta_{1,j}^{(r,i)}, \ldots, \beta_{L,j}^{(r,i)}, \varsigma_j^{2(r,i)}\right\}$, leading to their

straightforward numerically inexpensive evaluations in (3); we discuss the details in the posterior

computation section in Supplementary Materials. Let $\boldsymbol{\Omega} = \left(\left(\omega_{j,j'}\right)\right)$ be the precision matrix

corresponding to $\boldsymbol{R}_\Omega$, i.e., $\boldsymbol{R}_\Omega = \boldsymbol{\Psi}^{-\frac{1}{2}}\boldsymbol{\Omega}^{-1}\boldsymbol{\Psi}^{-\frac{1}{2}}$ with $\boldsymbol{\Psi} = \text{diag}(\boldsymbol{\Omega}^{-1})$. Then, by properties of

copula and the simple multivariate Gaussian distribution, it can be shown that, irrespective of

the form of the marginal $f_j^{(r,i)}$'s, the ROIs $Y_{t,j}^{(r,i)}$ and $Y_{t,j'}^{(r,i)}$ will be conditionally dependent on

(and hence functionally connected with) each other given the rest if and only if $\omega_{j,j'} \neq 0$. This

way, the precision matrix $\boldsymbol{\Omega}$ characterizes the functional connectivity between the different

$Y_{t,j}^{(r,i)}$'s across j.

We model the unknown univariate marginal distributions $f_j^{(r,i)}$'s using location-scale mixtures of

Gaussians. Such mixtures can flexibly estimate a very large class of unknown densities (Ghosal *et*

al., 1999). Specifically, we let

$$f_j^{(r,i)} = \sum_{h=1}^{K} \pi_{h,j}^{(r,i)} \mathrm{N}\left(\mu_{h,j}^{(r,i)}, \sigma_{h,j}^{2(r,i)}\right), \tag{4}$$

where $\pi_{h,j}^{(r,i)}$ is the weight attached to the h-th mixture component and $\sum_{h=1}^{K} \pi_{h,j}^{(r,i)} = 1$ with K

being a suitably chosen, moderately large, fixed integer.

While fMRI timeseries often showcase very distinct patterns and distributions for different

individuals as well as across different ROIs of the same individual, we expect primary sensory

region to be highly similar between healthy participants due to strongly stereotyped processing

across individuals and well-conserved auditory function in these brain regions across evolution

(Hutchison *et al.*, 2013). In (4), we therefore consider separate mixture models across different

ROIs, individuals, and runs, whereas in (3), the $\mathbf{Z}^{(r,i)}$'s share a common correlation matrix $\mathbf{R}_\Omega$

across all individuals and runs. This modeling strategy also allows borrowing of information

across individuals and runs to amplify signals for estimating resting-state functional connectivity

in adult humans while accounting for subject and run-specific variabilities using separate

marginal distributions. This is conceptually similar to approaches used in group independent

component analysis (Calhoun *et al.*, 2001) and cohort-level brain mapping (Varoquaux *et al.*,

2013). In later sections, we showed that this approach yields highly consistent estimates of

functional connectivity graphs even though BOLD signals from small deep brain regions have very

low signal-to-noise ratio (Bianciardi *et al.*, 2016; Colizoli *et al.*, 2020; de Hollander *et al.*, 2017;

Sclocco *et al.*, 2018).

To estimate the precision matrix $\mathbf{\Omega}$, we consider the framework of Chandra et al. (2021). Recall

that $\mathbf{R}_\Omega = \mathbf{\Psi}^{-\frac{1}{2}}\mathbf{\Omega}^{-1}\mathbf{\Psi}^{-\frac{1}{2}}$ with $\mathbf{\Psi} = \mathrm{diag}(\mathbf{\Omega}^{-1})$. We assume $\mathbf{\Omega}$ to admit a lower-rank plus diagonal

(LRD) decomposition $\mathbf{\Omega} = \mathbf{\Lambda}\mathbf{\Lambda}^T + \mathbf{\Delta}$ where $\mathbf{\Lambda}$ is a $d \times q$ matrix and a diagonal matrix $\mathbf{\Delta} =$

$\mathrm{diag}(\delta_1^2, \ldots, \delta_d^2)$ with positive δ_j^2's. Notably, all positive definite matrices admit such a

representation for some $q \leq d$.

We take a Bayesian route to estimation and inference, where we assign priors to the model

parameters, and then infer them based on samples drawn from the posterior using a Markov

chain Monte Carlo (MCMC) algorithm discussed in detail in the Supplementary Materials. As was

shown in Chandra et al. (2021), the LRD representation makes the MCMC sampling very efficient

via a latent variable augmentation scheme. In the Supplementary Materials, we also discuss a

multiple hypothesis testing-based edge discovery procedure that utilizes the posterior

uncertainty of the parameters and controls the false discovery rate (FDR) at 10% level in a

principled manner.

2.4 Priors on the parameters

For the autoregressive parameters in equation (*1*), for all r, i, j we assume

$\beta_{1,j}^{(r,i)}, \ldots, \beta_{L,j}^{(r,i)} \mid \varsigma_j^{2(r,i)} \overset{iid}{\sim} \mathrm{N}\left(0, v_\beta^{-1}\varsigma_j^{2(r,i)}\right), \quad \varsigma_j^{-2(r,i)} \overset{iid}{\sim} \mathrm{Ga}\left(a_\varsigma, b_\varsigma\right)$, where $v_\beta, a_\varsigma, b_\varsigma > 0$ are

fixed hyperparameters, and $\mathrm{Ga}(a, b)$ denotes a gamma distribution with mean a/b and variance

a/b^2. For the parameters of the mixture models specifying the marginals in equation (4), we

consider the following priors

$$\left(\pi_{1,j}^{(r,i)}, \ldots, \pi_{K,j}^{(r,i)}\right) \sim \mathrm{Dir}\left(\frac{\alpha_\pi}{K}, \ldots, \frac{\alpha_\pi}{K}\right), \qquad \left(\mu_{h,j}^{(r,i)}, \sigma_{h,j}^{2(r,i)}\right) \overset{iid}{\sim} \mathrm{NIG}(\mu_0, v_0, a_0, b_0),$$

where $(\mu, \sigma^2) \sim \mathrm{NIG}(\mu_0, v_0, a_0, b_0)$, implying that $\mu \mid \sigma^2 \sim \mathrm{N}(\mu_0, v_0^{-1}\sigma^2)$ and $\sigma^{-2} \sim \mathrm{Ga}(a_0, b_0)$.

We consider a shrinkage prior on the elements of $\boldsymbol{\Lambda}$ that shrinks redundant elements of $\boldsymbol{\Lambda}$ to

zero allowing additional model-based parameter reduction. In particular, we assign a two-

parameter generalization of the Dirichlet-Laplace (DL) prior from Bhattacharya et al.

(Bhattacharya *et al.*, 2015) that allows more flexible tail behavior on the elements of $\boldsymbol{\Lambda}$. On

a d-dimensional vector $\boldsymbol{\theta} = (\theta_1, \ldots, \theta_d)^T$, our DL prior with parameters a and b, denoted by

$\mathrm{DL}(a, b)$, can be specified in the following hierarchical manner.

$$\theta_j \mid \boldsymbol{\varrho}, \boldsymbol{\phi}, \tau \overset{ind}{\sim} \mathrm{N}\left(0, \varrho_j \phi_j^2 \tau^2\right), \quad \varrho_j \overset{iid}{\sim} \mathrm{Exp}(1/2), \quad \boldsymbol{\phi} \sim \mathrm{Dir}(a, \ldots, a), \quad \tau \sim \mathrm{Ga}(da, b),$$

where θ_j is the j-th element of $\boldsymbol{\theta}$, $\boldsymbol{\phi}$ and $\boldsymbol{\varrho}$ are vectors of same length as $\boldsymbol{\theta}$, $\mathrm{Exp}(a)$ is an

exponential distribution with mean $1/a$, $\mathrm{Dir}(a_1, \ldots, a_d)$ is a d-dimensional Dirichlet distribution.

The original DL prior is a special case with $b = 1/2$. We let $\mathrm{vec}(\boldsymbol{\Lambda}) \sim \mathrm{DL}(a, b)$.

We use a Dirichlet process (DP) prior (Ferguson, 1973) on the δ_j^2's as $\delta_j^2 \mid G \overset{iid}{\sim} G, G \mid \alpha \sim$

$\mathrm{DP}(\alpha, G_0)$ with $G_0 = \mathrm{Ga}(a_\delta, b_\delta), \alpha \sim \mathrm{Ga}(a_\alpha, b_\alpha)$, where α is the concentration parameter and

G_0 is the base measure to favor a smaller number of unique δ_j^2's in a model-based manner. The

DP model allows clustering the δ_j^2's facilitating additional data-adaptive parameter reduction

when necessary. Additionally a DP prior has full prior support on the range-space of the number

of unique δ_j^2's implying a fully flexible model (see Chapter 4 of (Ghosal and van der Vaart, 2017).

We discuss a Markov chain monte Carlo (MCMC) based strategy of sampling from the posterior

of ARMGCGM in Section S1.2 of the Supplementary Materials, where the involving steps are

parallelized over the subjects, allowing scalability. In Section S1.3 of the Supplementary

Materials, we discuss the choice of hyperparameters used for the analyses presented in this

paper. Our implementation using the proposed parallelized MCMC scheme ran in 125 min in a

system with 13th Gen Intel(R) Core(TM) i9-13900K CPU and 128GB RAM, fitting the ARMGCGM

to the 12-node auditory network with 7,500 MCMC iterations, *including* both phase-encoding

schemes.

3. Results

3.1 Partial correlations between regions of interest

Using our ARMGCGM approach, we first estimate the precision matrix $\boldsymbol{\Omega}$ and subsequently

compute the partial correlation matrix. We report the significant edges subject to controlling the

posterior false discovery rate (Sarkar *et al.*, 2008) at the 10% level. (Details are provided in Section

S1.4 in the Supplementary Materials.) In Figure 2 we provide the circos plots of the connectivity

graphs along with respective weighted adjacency matrices. The (j, h)-th off-diagonal element of

the adjacency matrices admit the value 0 if the edge between ROIs j and h is not statistically

significant, if the edge is significant then the corresponding partial correlation is plugged in to

indicate the strength of the edge.

We found that the strongest auditory connectivity was between adjacent structures in the same

hemisphere, particularly between the auditory midbrain (inferior colliculus, or IC) and thalamus

(medial geniculate body, or MGB) and between core and associative auditory cortex (TTG and

STG). Minimal connectivity was observed between homologous auditory structures across

hemispheres. Connectivity was also present between adjacent brainstem auditory structures

(cochlear nucleus [CN] and superior olivary complex [SOC]), largely bilaterally. Interestingly,

despite the SOC being the primary (and earliest) decussation point in the primary auditory

pathway, we only observed partial correlations between right CN and left SOC (in both data

partitions), not left CN and right SOC. (See the Discussion section for potential explanations.)

Figure 2. A: Partial correlation connectivity in data acquired with posterior-to-anterior (PA; left) and anterior-to-posterior (AP; right) phase-encoding directions using the ARMGCGM approach in subcortical and cortical auditory regions. Positive (negative) associations are represented by red

(blue) links, their opacities being proportional to the corresponding association strengths. The link widths are inversely proportional to the number of edges associated with the corresponding nodes. B: The same results as (A), viewed as adjacency matrices (left = PA runs; right = AP runs). C: Schematic of the auditory pathway from the cochlea through brainstem to cortex (https://osf.io/u2gxc/). D: Regions of interest from which functional timeseries were extracted. Top: cortical regions from FreeSurfer's DKT atlas. Bottom: subcortical auditory regions (Sitek et al., 2019).

327 *3.2 Effect of phase encoding scheme on subcortical connectivity*

Due to the anatomical location of the subcortical auditory structures—in dense, heterogeneous

subcortical regions and largely adjacent to CSF—we conducted connectivity analyses separately

on AP- and PA-acquired fMRI runs. We then compared the connectivity results from the two

acquisition schemes. Overall connectivity patterns were highly similar between the two phase-

encoding schemes, as seen in panels A and B of Figure 2. Subcortical connectivity was quite robust

between the brainstem auditory regions. To quantify the similarity between the results in PA and

AP acquisitions (plotted in Figure 3A), we computed the Pearson correlation coefficient r

between the estimated partial correlations. For the proposed ARMGCGM, $r = 0.940$ with p-

value < 0.001. To assess the similarity in sign of connectivity between acquisition schemes, we

computed the Jaccard dissimilarity on the signed off-diagonals of each adjacency matrix. For the

proposed ARMGCGM, the Jaccard dissimilarity was 0.231. Additionally, we computed the

Euclidean distance between partial correlations in the two acquisition schemes, compared it with

some selected approaches from the literature and report the results in Table 1. These results

indicate strong consistency between the findings in the two different acquisition schemes.

Figure 3. A: Pearson correlation of partial correlation values between posterior-to-anterior and anterior-to-posterior phase-encoding acquisition schemes across all edges in the graph (region-to-region connections). B: Partial correlation coefficients across cross-validation folds (10 folds for each of the two phase-encoding acquisition schemes).

3.3 Cross-validation of partial correlations

To assess the stability of our proposed ARMGCGM, we ran leave-10%-out cross-validation by

removing 10% randomly selected subjects in each fold. We repeated this for each phase encoding

scheme for a total of 20 folds; partial correlation coefficients for each fold are presented in Figure

3B. Across all folds (and both phase encoding schemes), the results are very highly similar

(intraclass correlation of edgewise partial correlations across cross-validation folds = 0.991).

3.4 Comparison with existing approaches

We compared with three standard approaches in the literature: (1) correlation analysis between

the ROIs (Biswal *et al.*, 1995; Cordes *et al.*, 2000; Fox and Raichle, 2007; Lowe *et al.*, 1998; Smith,

Vidaurre, *et al.*, 2013); and partial correlation analyses using (2) the graphical lasso (Friedman *et*

al., 2008) and (3) the precision factor model (Chandra *et al.*, 2021). In all comparisons, we did

separate analyses for each of the acquisition schemes.

Comparison 1: Full correlation approach: Here we study the marginal correlation between the

ROIs. Letting $\rho_{j,j'}$ denote the correlation between ROIs j and j' in resting state we test

$$H_{0,j,j'}: \left|\rho_{j,j'}\right| = 0 \text{ versus } H_{1,j,j'}: \left|\rho_{j,j'}\right| \neq 0 \text{ for all } 1 \leq j < j' \leq d.$$

For each acquisition scheme, we first concatenate the timeseries across all runs and individuals.

Then we perform t-tests for correlation for all (j, j') pairs followed by the Benjamini-Hochberg

false discovery rate (FDR) correction (Benjamini and Hochberg, 1995) for multiplicity adjustment

and control the FDR at level 0.10. Traditionally, full correlations are used to measure functional

"connectivity" in resting state fMRI studies (Biswal *et al.*, 1995; Cordes *et al.*, 2000; Lowe *et al.*,

1998). In Figure 4 we provide the correlation graphs and correlation matrices separately for each

acquisition schemes. We find the correlation graphs to be much denser compared to the partial

correlation graphs presented in Figure 2. Unlike in the ARMGCGM method, we observed negative

values when using full correlations (particularly in the PA acquisition scheme), although the

negative correlations are generally closer to 0 than the positive correlations are. In this full

correlation approach, the similarity between runs split by data acquisition scheme was

characterized by Pearson's r of 0.811 (p-value < 0.001) and a Euclidean distance of 1.016. The

Jaccard dissimilarity of the signed adjacency matrix was 0.338.

Figure 4. A: Full correlation connectivity in data acquired with posterior-to-anterior (PA; left) and anterior-to-posterior (AP; right) phase-encoding directions using t-tests. Positive (negative) associations are represented by red (blue) links, their opacities being proportional to the corresponding association strengths. The link widths are inversely proportional to the number of edges associated with the corresponding nodes. B: The same results as (A), viewed as adjacency matrices (left = PA runs; right = AP runs).

Comparison 2: Partial correlations with Glasso approach: We first consider the graphical lasso (Glasso) approach (Friedman *et al.*, 2008) as another alternative choice for partial correlation based conditional graph estimation. Glasso assumes iid data from a multivariate Gaussian distribution (i.e., without any correction for autocorrelation) with $\ell_1(\cdot)$ penalty on the precision

matrix. In this analysis we concatenated the $Y_{t,j}^{(r,i)}$ values across (r,i) and created a $(TR) \times d$

matrix, say Y, for each acquisition scheme, and applied the Glasso model on Y. We use 10-fold

cross-validation to choose the optimal penalty parameter. The Glasso approach provides a point

estimate of the sparse precision matrix and hence the functional connectivity network. In panels

A and B of Figure 5, we plot the connectivity graphs and weighted adjacency matrices,

respectively, separately for each acquisition scheme.

These functional connectivity graphs in Figure 5 are much denser compared to the estimates

obtained by our proposed ARMGCGM in Figure 2. To quantify the robustness and stability of the

Glasso graphs in this application, we compute Pearson correlation coefficient and Euclidean

distance between the estimated partial correlations, and the Jaccard dissimilarity between the

signs of the adjacency matrices in PA and AP acquisitions in the same manner as we did for

ARMGCGM elaborated in Section *3.2*. We reported the values in Table 1. Figure 5 indicates that

the strong positive correlations are consistent across the acquisition schemes. However,

substantial discrepancy can be observed for the weak edges, particularly for the negative

correlations. This is a common phenomenon for graphical models if the Gaussian assumption is made on non-Gaussian distributed data; see, e.g., Section 5, example 1(a) in (Guha *et al.*, 2020).

Figure 5. A: Partial correlation connectivity in data acquired with posterior-to-anterior (PA; left) and anterior-to-posterior (AP; right) phase-encoding directions using the Glasso approach. Positive (negative) associations are represented by red (blue) links, their opacities being proportional to the corresponding association strengths. The link widths are inversely proportional to the number of edges associated with the corresponding nodes. B: The same results as (A), viewed as adjacency matrices (left = PA runs; right = AP runs).

Comparison 3: Partial correlations with PFM: In this analysis we concatenated the $Y_{t,j}^{(r,i)}$ values across (r, i) and created a $(TR) \times d$ matrix, say $\boldsymbol{Y}$, for each acquisition scheme, and applied the PFM on $\boldsymbol{Y}$. Similar to Glasso, the PFM also assumes iid data from a multivariate Gaussian distribution and does not correct for temporal autocorrelations in the data. We infer on the graph using the Bayesian multiple comparison technique described in the Supplementary Materials. Results are provided in Figure 6. The connectivity graphs majorly differed with our ARMGCGM results, with the PFA model-derived graph being much denser and including more (weakly) negative edges. In the top panel of Figure 7 we plot the marginal Gaussian fits on the histograms of some BOLD signals. Clearly the simple Gaussian assumption does not hold here and a more sophisticated approach like ours is required.

To assess the robustness and stability of the PFM, we computed Pearson correlation coefficient and Euclidean distance between the estimated partial correlations, and the Jaccard dissimilarity between the signs of the adjacency matrices in PA and AP acquisitions in the same manner as we did for ARMGCGM elaborated in Section *3.2*. We reported the values in Table 1, where we see that PFM exhibits more consistency than Glasso but ARMGCGM performed best.

Figure 6. A: Partial correlation connectivity in data acquired with posterior-to-anterior (PA; left) and anterior-to-posterior (AP; right) phase-encoding directions using the PFA approach (comparison 3). Positive (negative) associations are represented by red (blue) links, their opacities being proportional to the corresponding association strengths. The link widths are inversely proportional to the number of edges associated with the corresponding nodes. B: The same results as (A), viewed as adjacency matrices (left = PA runs; right = AP runs).

Table 1 Several measures of dissimilarities between the functional connectivity networks (using full and partial correlations) estimated in each of the acquisition schemes are reported here. We provide results for all the correlation-based functional connectivity analyses, viz. ARMGCGM, full correlation, Glasso and PFA. Note that lower values of Jaccard dissimilarity and Euclidean distance implies better method.

Method	Pearson's Correlation Coefficient	Jaccard Dissimilarity	Euclidean Distance
ARMGCGM	0.940 (p-value < 0.001)	0.231	0.262
Full correlation	0.811 (p-value < 0.001)	0.338	1.016
Glasso	0.749 (p-value < 0.001)	0.414	0.959
PFM	0.747 (p-value < 0.001)	0.379	0.975

3.5 Fit of the autoregressive matrix-Gaussian copula graphical model

Density fits: We studied the goodness-of-fit of the proposed ARMGCGM. In the top panel of

Figure 7, we plotted the sample histograms and the corresponding fitted marginal densities for

some selected ROIs. The sample histograms strongly indicate the distribution of the data to

substantially deviate from Gaussian distributions, including some with multiple well-separated

modes. Figure 7 also shows that our flexible location-scale mixture of Gaussians fit the data very

well, even for the most complicated distributions.

Note that finite mixture models with reasonably large number of mixture components can

approximate nonparametric Dirichlet process mixture models (Ishwaran and Zarepour, 2002b,

2002a). To validate whether the number of mixtures ($K = 20$) in model (4) is adequate, we

computed the median number of non-empty clusters across MCMC samples for each subject and

ROI in each run. In the bottom panel of Figure 7 we plotted the histograms of the medians across

the subjects. As the number of non-empty clusters are smaller than K consistently across all

setups yielding excellent fits for complicated distributions, we conclude that our model

specifications are adequate.

Figure 7. Marginal fits for the location-scale mixture of Gaussian distributions: The sample histograms in the top panel in green showed that distribution of the ROIs substantially deviates from Gaussian assumption. The blue lines were the respective fitted densities corresponding to the mixture model in equation (4). These figures indicated excellent goodness-of-fit. The dashed brown lines corresponded to marginal Gaussian fits of the vanilla PFM which were evidently underfitted. In the bottom panel we plotted the histogram of the number of occupied clusters across subjects corresponding to the mixture model indicating that our model specifications are adequate.

Autocorrelation corrections: From Section 2.3 recall that $Z_{t,j}^{(r,i)}$'s were the transformed BOLD

signal timeseries in the Gaussian space and $\xi_{t,j}^{(r,i)}$'s were the autocorrelation corrected timeseries.

To check for autocorrelation corrections using model (1), we plotted the partial autocorrelations

between $Z_{t,j}^{(r,i)}$'s and $\xi_{t,j}^{(r,i)}$'s across time for all ROIs. Since the **Z** and **ξ** values vary across MCMC

iterations, we used their posterior expectations in this analysis. In Figure 8 we plot the partial

autocorrelations for a randomly selected individual in run 1 of posterior-to-anterior acquisition

scheme; the blue horizontal dashed lines represent the 5% band. The figure clearly indicates that

the autoregressive model of order $L = 5$ corrects for the autocorrelations. As we obtained very

similar results for other runs and acquisition schemes, we omit those in the paper.

Figure 8. Partial autocorrelation plots of the fMRI timeseries for a randomly selected subject in the posterior-to-anterior acquisition scheme across the ROIs before (orange) and after (green) autocorrelation corrections; the horizontal dashed blue lines represent the 5% band. The plot indicates that autocorrelations are corrected in our model.

4. Discussion

The mammalian auditory pathway consists of a series of obligatory and interconnected

subcortical and cortical brain structures. Assessing the connectivity of the human subcortical

auditory structures has been limited due to methodological challenges of non-invasive imaging

of the deep, small structures. Recent acquisition and analytical advances enable finer grained

investigations of connectivity throughout the brain, including the brainstem. In this paper, we

validated a novel an autoregressive matrix Gaussian copula graphical model to estimate

functional auditory connectivity patterns from a publicly available high-resolution resting state

functional MRI dataset. Using partial correlations (as opposed to full correlations) allowed us to

identify specific relationships between nodes in a connectivity graph by removing shared

variance across nodes (Supplementary Figures S1 and S2). We found highly consistent

connectivity patterns between adjacent auditory brain regions along the auditory pathway that

demonstrate the efficacy of our connectivity method as well as the potential for functional

connectivity investigations of the subcortical auditory system. Below, we separately discuss our

novel scientific findings and our novel contributions to the statistics literature.

Novel contributions to the human auditory neuroscience literature

To date, there have been only limited applications of functional MRI methods to study subcortical

auditory connectivity (Berlot *et al.*, 2020; Hofmeier *et al.*, 2018). Using our novel ARMGCGM

approach in the present study, we found strong partial correlations between cochlear nucleus

(CN) and superior olivary complex (SOC) bilaterally using resting state functional MRI data. Most

interestingly, we observed contralateral CN–SOC connectivity between right CN and left SOC (and

in both data acquisition schemes), fitting the ground truth primary auditory pathway crossing

from left to right (and vice versa) between CN and SOC (Barnes *et al.*, 1943; Schofield, 1994).

These functional connectivity patterns between CN and SOC have not been previously observed

in human auditory brainstem in vivo but follow our understanding of the mammalian primary

auditory pathway based on research in animal models (Barnes *et al.*, 1943; Doucet and Ryugo,

2003; Harrison and Irving, 1966). Principally, auditory information that is transduced by the

cochlea of each ear is transmitted via the cochlear nerve to the cochlear nucleus, the first stage

of the central auditory pathway, on each side of the brainstem. In the primary auditory pathway,

the lemniscal anteroventral subdivision of the cochlear nucleus enhances the fine temporal

precision of incoming auditory signals (Pickles, 2015). From there, auditory signals are passed to

both the ipsilateral and contralateral SOC for further auditory processing, including spatial

localization (Moore, 2000). The SOC is comprised of multiple distinct subdivisions, which receive

ipsilateral and contralateral connections from cochlear nucleus to varying degrees (Pickles,

2015), aligning with our overall bilateral connectivity results between CN and SOC.

While we observed consistent right CN and left SOC connectivity, it is unclear why similar patterns

were not observed between left CN and right SOC. One contributor is the lower signal-to-noise

ratio in fMRI data from the lower brainstem. Paired with the small size of each of the brainstem

auditory nuclei, we may still be at the edge of what is detectable using present functional

connectivity methods. Additionally, this analysis was conducted on "resting state" fMRI data,

during which no auditory stimuli of interest were presented or overt tasks were conducted.

Resting state fMRI connectivity in the cochlear nucleus and superior olivary complex has not been

examined in the previous literature to our knowledge; it is possible that sound-evoked

connectivity methods would evoke greater functional connectivity, particularly in these earliest

stages of the auditory pathway. Further, ipsilateral connections (i.e., between left CN and left

SOC and between right CN and right SOC) may be artifactually stronger due to their close physical

proximity. Even with relatively high 1.05 mm spatial resolution 7T fMRI data, CN and SOC on each

side are only separated by a few voxels. These regions are thus at increased likelihood of sharing

temporal fluctuations due to partial volume effects, wide point-spread functions, spatial

dependence, or other as-yet-unsolved fMRI confounds that are particularly acute in the lower

brainstem.

Moving up the primary auditory pathway, we observed significant partial correlations between

ipsilateral inferior colliculus (IC) in midbrain and medial geniculate body (MGB) of the thalamus

in both hemispheres and in both phase-encoding schemes. Inferior colliculus is a major

convergence point in the auditory system, with the lemniscal IC subdivision being thought to

convert distinct auditory features into discrete auditory objects for the first time in the auditory

pathway. MGB continues the refinement of auditory objects via direct lemniscal connections

from IC as well as rich corticofugal connections from auditory cortex to non-lemniscal MGB

subdivisions (Pickles, 2015). Interestingly, we found strong partial correlations between left and

right MGB in both datasets. Although not directly connected by large white matter bundles, left

and right MGB are expected to process auditory information from IC at similar levels of

abstraction. Thus, partial correlations may reflect indirect but shared neural mechanisms of

auditory processing in the thalamus.

We did not observe IC partial correlation connectivity with either brainstem or cortical structures.

IC is a key hub in the auditory system, receiving bottom-up sensory information as well as top-

down modulating signals from auditory cortex and other brain regions. The lack of partial

correlation connectivity with IC may be due to strong IC subdivision-specific functionality, with IC

core primarily serving an ascending lemniscal role and dorsal and external IC having top-down

and non-lemniscal functions. Averaging over these subdivisions may obfuscate specific

connectivity patterns. Alternatively, our results may suggest that it does not have a specialized

relationship with any one region beyond MGB but rather integrates and transforms auditory and

other neural signals.

Finally in auditory cortex, transverse temporal gyrus (TTG)—the location of primary auditory

cortex—was strongly connected with ipsilateral superior temporal gyrus (STG), which contains

secondary and associative auditory cortices. Primary auditory cortex receives direct input from

lemniscal MGB and is the last auditory structure with fine-grained tonotopicity (Pickles, 2015). In

humans, STG is hierarchically structured, with portions further away from primary auditory

cortex having increasingly wider temporal integration windows (Hamilton *et al.*, 2018; Norman-

Haignere *et al.*, 2022) and greater categorical specificity (Bhaya-Grossman and Chang, 2022; Feng

et al., 2021; Hamilton *et al.*, 2020; Keshishian *et al.*, 2023; Nourski *et al.*, 2018; Pernet *et al.*, 2015;

Rauschecker and Tian, 2000; Rupp *et al.*, 2022). While invasive recordings from human STG

suggest a potential direct connection between MGB and posterior STG (Hamilton *et al.*, 2021),

we found mixed evidence for such a direct pathway in ourOur partial correlation data (in one

hemisphere in only one of the data partitions). Our partial correlation functional connectivity

results align with a vast literature demonstrating information flow between primary and non-

primary auditory cortex (for review, see (Hackett, 2011). The lack of contralateral partial

correlation approach measures the correlation between the concerned ROIs after filtering out

the indirect effects of the remaining ROIs. Complementarily, some literature suggests that left

and right auditory cortex process auditory information at distinct timescales and levels of

abstraction (Güntürkün *et al.*, 2020; Hickok and Poeppel, 2007; Zatorre *et al.*, 2002), with left

auditory cortex being uniquely tuned to rapidly changing temporal information—such as the

phonetics of speech sounds—while right auditory cortex is more sensitive to slower changes in
the spectral domain, particularly for speech prosody as well as music.

In comparison to our ARMGCGM partial correlation approach, we computed full correlations in
the same network. Connections were much denser in the full correlation approach, aligning with
the rich interconnectedness of the auditory system (Pickles, 2015). Unlike with partial
correlations, which highlighted hierarchical connections between adjacent notes along the
auditory pathway, we observed positive full correlations between all auditory cortical regions,
regardless of hemisphere. Additionally, we found a strong positive subnetwork including IC and
MGB bilaterally, whereas many of these connections (such as between left and right IC) were
absent in the ARMGCGM partial correlation analysis. Since partial correlations characterize
connectivity between two nodes after filtering out the effects of the other nodes, our combined
results point to widespread shared information across the auditory system (per full correlation
analysis) with additional shared processing between adjacent nodes of the canonical auditory
pathway (per partial correlation analysis). This suggests distinct but complementary use of full
and partial correlations, with full correlation analysis identifying a rich network of interconnected
nodes, while partial correlations are sensitive to strong node-to-node connections.

Novel contributions to the graphical model literature

In this article, we developed an autoregressive matrix-Gaussian copula graphical model
(ARMGCGM) for non-Gaussian distributed data with temporal autocorrelation, the problem of
estimating brain connectivity patterns from resting state fMRI data being the motivating
problem. The ARMGCGM first uses higher order autoregressive models to capture the temporal
dependence in the time series for each brain region of interest, then uses flexible location-scale
mixtures of Gaussians for modeling component wise residual marginal distributions associated
with different regions, and finally uses a Gaussian copula to capture the dependence across the
different regions. The ARMGCGM allows borrowing of information across subjects to infer on a
common connectivity graph while appropriately taking into account subject and run-specific
variability via flexible mixture models. We leverage recent advances on modeling precision
matrices via a flexible but computationally efficient low-rank-diagonal decomposition method
that not only allows efficient exploration of the posterior space for estimating the connectivity

graph but also enables easy assessment of associated uncertainty. Compared to alternative

approaches to exploiting partial correlations to estimate connectivity graphs, our proposed

ARMGCGM method produces results that are more consistent across the acquisition schemes

with respect to multiple metrics. Additionally, the results remain highly stable in a leave-10%-out

cross-validation. Considering the low signal to noise ratio in BOLD signals from deep small

auditory structures, our results demonstrate the sensitivity and specificity of our model to

neurobiologically plausible connections. Although the proposed ARMGCGM is a more nuanced

approach and potentially capable of fitting arbitrary complicated distributions, it can be

numerically expensive compared to simplistic parametric models. However, our parallelized

Markov chain Monte Carlo implementation ran across all participants and runs in just over two

hours, demonstrating a feasible computation time given the size of the data.

Comparisons to connectivity literature

Our study is the first to systematically assess connectivity across the human auditory pathway

using multiple connectivity measures, with previous subcortical connectivity studies limited to

full correlation analysis (Berlot *et al.*, 2020; Hofmeier *et al.*, 2018; Leaver *et al.*, 2016; Zhang *et*

al., 2015). Given the anatomical and methodological constraints with subcortical fMRI, the

limited fMRI connectivity literature is not too surprising. First, the deep location of the subcortical

structures places them far from MRI transmit and receive coils, limiting the signal-to-noise ratio

from these regions (Miletić *et al.*, 2020). Because the brainstem is relatively centrally located

relative to the multiple receiver coils, accelerated acquisition techniques that are based on phase

differences between receiver coils are less effective (Preibisch *et al.*, 2015). Second, subcortical

nuclei can be quite small, requiring higher resolution imaging protocols (which unfortunately

trade off SNR in order to achieve greater spatial resolution). Third, subcortical nuclei are densely

organized adjacent to nuclei with heterogeneous functions, so voxels immediately next to those

containing core auditory structures could contain visual, motor, or sensory nuclei, white matter,

CSF, or a combination of any of these. Ultimately, each of these constraints limits the SNR from

subcortical auditory nuclei.

Constraints in human subcortical auditory research have translational consequences beyond

basic science. For instance, while cochlear implants have been widely successful at providing

sensory information to individuals with sensorineural hearing impairments with an intact

cochlear nerve (Kral *et al.*, 2019; Reiss, 2020), auditory prostheses in the central auditory system

have been less successful (Lim *et al.*, 2009; Shetty *et al.*, 2021), due in no small part to our limited

understanding of the complexity of sound representation in the ascending auditory pathway.

Limitations and future directions

As the canonical neuroanatomy of the primary auditory pathway is consistent across individuals

and well-described in the literature (Sitek *et al.*, 2019), we have the *a priori* expectation of a

shared auditory graph across all participants. As the goal of this study was to map a network that

is strongly expected based on anatomy and non-human neurophysiology, we built a joint model

that includes data from all participants. This is similar to approaches used in group independent

component analysis (Calhoun *et al.*, 2001) and cohort-level brain mapping (Varoquaux *et al.*,

2013). However, subject-specific differences in the distributions of the BOLD signals as well as

autocorrelations between successive scans can induce artifactual and noisy edges in functional

connectivity graphs, as seen in the Glasso and PFMs; we take care of these issues in ARMGCGM.

Nevertheless, we did not investigate differences in functional connectivity between participants

in the current article. Building on ARMGCGM to explore how functional connectivity varies

between individuals and groups or as a function of behavior is a priority for future work.

In general, resting state fMRI connectivity measures become more reliable with longer scans (Zuo

et al., 2019). Measurement correlations increase as time in the scanner increases, from Pearson's

$r = 0.82$ at 9 min to $r = 0.92$ at 27 min to $r = 0.97$ at 90 min (Laumann *et al.*, 2015). Others

described improved intraclass correlation coefficients with datasets beyond 20 minutes and up

to 50 minutes (Xu *et al.*, 2016). The Midnight Scan Club group (Gordon, Laumann, Gilmore, *et al.*,

2017) computed a range of network connectivity metrics and found that reliability generally

required at least 30 minutes of resting state data per subject. One paper (Greene *et al.*, 2020)

specifically investigated functional connectivity in subcortical structures and found even longer

scan requirements (up to 100 minutes) for subcortical structures due to decreased signal-to-

noise ratios deeper in the brain. Additionally, primary sensory networks are among the most

stable within and across participants (Gratton *et al.*, 2018; Hutchison *et al.*, 2013). We therefore

believe it is appropriate and necessary to use datasets with longer scans of resting state data in
order to investigate even static subcortical auditory connectivity.

Further, while many brain networks exhibit temporal dynamics that can tell us about mental state
(Fornito *et al.*, 2012) or disease ((Sakoğlu *et al.*, 2010); see (Hutchison *et al.*, 2013) for a review),
functional connectivity within primary sensory networks are among the most stable over time
(Gratton *et al.*, 2018), as they share bidirectional physical connections, share contributions to the
same physiological tasks, and are evolutionarily conserved across species (Hutchison and
Everling, 2012). In the present work, we were interested in characterizing the stationary
connectivity in the primary auditory pathway that is present at rest across individuals. Adapting
time-varying dynamics into this model is a promising future direction, particularly if we are
interested in higher level cognitive brain networks that vary as a function of task or mental state.

5. Conclusions

In this article, we validated a novel autoregressive matrix Gaussian copula graphical model for
partial correlation estimation while appropriately correcting for temporal autocorrelations. Using
this approach, we identified functional connectivity in the human auditory system using resting
state functional MRI. Whereas a complementary approach using full correlations identified a rich
network of interconnected auditory regions, partial correlations highlighted direct connections
between adjacent structures along auditory pathways. In particular, subcortical connectivity was
highly consistent across acquisitions, demonstrating the utility and applicability of functional
connectivity methods in deep brain structures. In the future, we plan to investigate whole-brain
partial correlation connectivity across sensory, motor, and higher cognitive networks using the
proposed models and their relationship to behavior across individuals.

Acknowledgements:

Data were provided [in part] by the Human Connectome Project, WU-Minn Consortium (Principal
Investigators: David Van Essen and Kamil Ugurbil; 1U54MH091657) funded by the 16 NIH

Institutes and Centers that support the NIH Blueprint for Neuroscience Research; and by the
McDonnell Center for Systems Neuroscience at Washington University.

Ethics:

Data usage adhered to the Open Access Data Use Terms from the Human Connectome Project
(WU-Minn HCP).

Data and Code Availability:

Data are publicly available through the Human Connectome Project
(https://www.humanconnectome.org/study/hcp-young-adult). Analysis code is available in the
Supplementary Materials.

Author Contributions:

Noirrit Chandra: conceptualization (equal); formal analysis (lead); software (lead); visualization
(equal); writing – original draft (equal); writing – review and editing (equal).

Kevin R. Sitek: conceptualization (equal); formal analysis (equal); visualization (equal); writing –
original draft (lead); writing – review and editing (equal).

Bharath Chandrasekaran: conceptualization (equal); writing – review and editing (equal).

Abhra Sarkar: conceptualization (equal); writing – review and editing (equal).

Funding:

This work was supported by the National Science Foundation grant NSF DMS- 1953712 (to AS and
BC) and the National Institutes of Health/National Institute on Deafness and Other
Communication Disorders (K01DC019421 to KRS and R01DC01550 to BC).

Declaration of Competing Interests:

The authors have no competing interests.

References

Bibliography
678

Abrams DA, Kochalka J, Bhide S, Ryali S, Menon V. Intrinsic functional architecture of the
human speech processing network. Cortex 2020; 129: 41–56.

Baldassarre A, Lewis CM, Committeri G, Snyder AZ, Romani GL, Corbetta M. Individual
variability in functional connectivity predicts performance of a perceptual task. Proc. Natl. Acad.
Sci. USA 2012; 109: 3516–3521.

Barnes WT, Magoun HW, Ranson SW. The ascending auditory pathway in the brain stem of the
monkey. J. Comp. Neurol. 1943; 79: 129–152.

Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful
approach to multiple testing. Journal of the Royal Statistical Society: Series B (Methodological)
1995; 57: 289–300.

Berlot E, Arts R, Smit J, George E, Gulban OF, Moerel M, et al. A 7 Tesla fMRI investigation of
human tinnitus percept in cortical and subcortical auditory areas. Neuroimage Clin. 2020; 25:
102166.

Bhattacharya A, Pati D, Pillai NS, Dunson DB. Dirichlet-Laplace priors for optimal shrinkage. J.
Am. Stat. Assoc. 2015; 110: 1479–1490.

Bhaya-Grossman I, Chang EF. Speech computations of the human superior temporal gyrus.
Annu. Rev. Psychol. 2022; 73: 79–102.

Bianciardi M, Toschi N, Eichner C, Polimeni JR, Setsompop K, Brown EN, et al. In vivo
functional connectome of human brainstem nuclei of the ascending arousal, autonomic, and
motor systems by high spatial resolution 7-Tesla fMRI. MAGMA 2016; 29: 451–462.

Biswal B, Yetkin FZ, Haughton VM, Hyde JS. Functional connectivity in the motor cortex of
resting human brain using echo-planar MRI. Magn. Reson. Med. 1995; 34: 537–541.

Cai MB, Shvartsman M, Wu A, Zhang H, Zhu X. Incorporating structured assumptions with
probabilistic graphical models in fMRI data analysis. Neuropsychologia 2020; 144: 107500.

Calhoun VD, Adali T, Pearlson GD, Pekar JJ. A method for making group inferences from
functional MRI data using independent component analysis. Hum. Brain Mapp. 2001; 14: 140–
151.

Cha K, Zatorre RJ, Schönwiesner M. Frequency Selectivity of Voxel-by-Voxel Functional
Connectivity in Human Auditory Cortex. Cereb. Cortex 2016; 26: 211–224.

Chai XJ, Hirshfeld-Becker D, Biederman J, Uchida M, Doehrmann O, Leonard JA, et al. Altered
intrinsic functional brain architecture in children at familial risk of major depression. Biol.

Psychiatry 2016; 80: 849–858.

Chandra NK, Mueller P, Sarkar A. Bayesian Scalable Precision Factor Analysis for Massive
Sparse Gaussian Graphical Models. arXiv 2021: arXiv:2107.11316.

Chen J, Hu B, Qin P, Gao W, Liu C, Zi D, et al. Altered brain activity and functional
connectivity in unilateral sudden sensorineural hearing loss. Neural Plast. 2020; 2020: 9460364.

Colizoli O, de Gee JW, van der Zwaag W, Donner TH. Comparing fMRI responses measured at
3 versus 7 Tesla across human cortex, striatum, and brainstem. BioRxiv 2020

Cordes D, Haughton VM, Arfanakis K, Wendt GJ, Turski PA, Moritz CH, et al. Mapping
functionally related regions of brain with functional connectivity MR imaging. AJNR Am. J.
Neuroradiol. 2000; 21: 1636–1644.

Dale AM, Fischl B, Sereno MI. Cortical surface-based analysis. I. Segmentation and surface
reconstruction. Neuroimage 1999; 9: 179–194.

De Martino F, Moerel M, van de Moortele P-F, Ugurbil K, Goebel R, Yacoub E, et al. Spatial
organization of frequency preference and selectivity in the human inferior colliculus. Nat.
Commun. 2013; 4: 1386.

Deng Z, Chandrasekaran B, Wang S, Wong PCM. Resting-state low-frequency fluctuations
reflect individual differences in spoken language learning. Cortex 2016; 76: 63–78.

Di Martino A, Kelly C, Grzadzinski R, Zuo X-N, Mennes M, Mairena MA, et al. Aberrant
striatal functional connectivity in children with autism. Biol. Psychiatry 2011; 69: 847–856.

Doucet JR, Ryugo DK. Axonal pathways to the lateral superior olive labeled with biotinylated
dextran amine injections in the dorsal cochlear nucleus of rats. J. Comp. Neurol. 2003; 461: 452–
465.

Eckert MA, Kamdar NV, Chang CE, Beckmann CF, Greicius MD, Menon V. A cross-modal
system linking primary auditory and visual cortices: evidence from intrinsic fMRI connectivity
analysis. Hum. Brain Mapp. 2008; 29: 848–857.

Elam JS, Glasser MF, Harms MP, Sotiropoulos SN, Andersson JLR, Burgess GC, et al. The
Human Connectome Project: A retrospective. Neuroimage 2021; 244: 118543.

Esteban O, Adebimpe A, Markiewicz CJ, Goncalves M, Blair RW, Cieslak M, et al. The
Bermuda Triangle of d- and f-MRI sailors - software for susceptibility distortions (SDCFlows).
2021

Feng G, Gan Z, Yi HG, Ell SW, Roark CL, Wang S, et al. Neural dynamics underlying the
acquisition of distinct auditory category structures. Neuroimage 2021; 244: 118565.

Ferguson TS. A Bayesian Analysis of Some Nonparametric Problems on JSTOR [Internet].
1973[cited 2022 May 10] Available from: https://www.jstor.org/stable/2958008

Ferri J, Ford JM, Roach BJ, Turner JA, van Erp TG, Voyvodic J, et al. Resting-state thalamic
dysconnectivity in schizophrenia and relationships with symptoms. Psychol. Med. 2018; 48:
2492–2499.

Fornito A, Harrison BJ, Zalesky A, Simons JS. Competitive and cooperative dynamics of large-
scale brain functional networks supporting recollection. Proc. Natl. Acad. Sci. USA 2012; 109:
12788–12793.

Fox MD, Raichle ME. Spontaneous fluctuations in brain activity observed with functional
magnetic resonance imaging. Nat. Rev. Neurosci. 2007; 8: 700–711.

Friedman J, Hastie T, Tibshirani R. Sparse inverse covariance estimation with the graphical
lasso. Biostatistics 2008; 9: 432–441.

Ghosal S, Ghosh JK, Ramamoorthi RV. Posterior consistency of Dirichlet mixtures in density
estimation. Ann. Statist. 1999; 27

Ghosal S, van der Vaart A. Fundamentals of nonparametric bayesian inference. Cambridge:
Cambridge University Press; 2017.

Glasser MF, Sotiropoulos SN, Wilson JA, Coalson TS, Fischl B, Andersson JL, et al. The
minimal preprocessing pipelines for the Human Connectome Project. Neuroimage 2013; 80:
105–124.

Gordon EM, Laumann TO, Adeyemo B, Petersen SE. Individual Variability of the System-Level
Organization of the Human Brain. Cereb. Cortex 2017; 27: 386–399.

Gordon EM, Laumann TO, Gilmore AW, Newbold DJ, Greene DJ, Berg JJ, et al. Precision
functional mapping of individual human brains. Neuron 2017; 95: 791–807.e7.

Gratton C, Laumann TO, Nielsen AN, Greene DJ, Gordon EM, Gilmore AW, et al. Functional
brain networks are dominated by stable group and individual factors, not cognitive or daily
variation. Neuron 2018; 98: 439–452.e5.

Greene DJ, Marek S, Gordon EM, Siegel JS, Gratton C, Laumann TO, et al. Integrative and
Network-Specific Connectivity of the Basal Ganglia and Thalamus Defined in Individuals.
Neuron 2020; 105: 742–758.e6.

Greicius MD, Flores BH, Menon V, Glover GH, Solvason HB, Kenna H, et al. Resting-state
functional connectivity in major depression: abnormally increased contributions from subgenual
cingulate cortex and thalamus. Biol. Psychiatry 2007; 62: 429–437.

Guha N, Baladandayuthapani V, Mallick BK. Quantile Graphical Models: a Bayesian Approach.
Journal of Machine Learning Research 2020

Güntürkün O, Ströckens F, Ocklenburg S. Brain lateralization: A comparative perspective.
Physiol. Rev. 2020; 100: 1019–1063.

Hackett TA. Information flow in the auditory cortical network. Hear. Res. 2011; 271: 133–146.

Hahn A, Stein P, Windischberger C, Weissenbacher A, Spindelegger C, Moser E, et al. Reduced
resting-state functional connectivity between amygdala and orbitofrontal cortex in social anxiety
disorder. Neuroimage 2011; 56: 881–889.

Hamilton LS, Edwards E, Chang EF. A spatial map of onset and sustained responses to speech in
the human superior temporal gyrus. Curr. Biol. 2018; 28: 1860–1871.e4.

Hamilton LS, Oganian Y, Chang EF. Topography of speech-related acoustic and phonological
feature encoding throughout the human core and parabelt auditory cortex. BioRxiv 2020

Harrison JM, Irving R. Ascending connections of the anterior ventral cochlear nucleus in the rat.
J. Comp. Neurol. 1966; 126: 51–63.

Hawley ML, Melcher JR, Fullerton BC. Effects of sound bandwidth on fMRI activation in

human auditory brainstem nuclei. Hear. Res. 2005; 204: 101–110.

Hickok G, Poeppel D. The cortical organization of speech processing. Nat. Rev. Neurosci. 2007;
8: 393–402.

Hofmeier B, Wolpert S, Aldamer ES, Walter M, Thiericke J, Braun C, et al. Reduced sound-
evoked and resting-state BOLD fMRI connectivity in tinnitus. Neuroimage Clin. 2018; 20: 637–
649.

de Hollander G, Keuken MC, van der Zwaag W, Forstmann BU, Trampel R. Comparing
functional MRI protocols for small, iron-rich basal ganglia nuclei such as the subthalamic
nucleus at 7 T and 3 T. Hum. Brain Mapp. 2017; 38: 3226–3248.

Husain FT, Schmidt SA. Using resting state functional connectivity to unravel networks of
tinnitus. Hear. Res. 2014; 307: 153–162.

Hutchison RM, Everling S. Monkey in the middle: why non-human primates are needed to
bridge the gap in resting-state investigations. Front. Neuroanat. 2012; 6: 29.

Hutchison RM, Womelsdorf T, Allen EA, Bandettini PA, Calhoun VD, Corbetta M, et al.
Dynamic functional connectivity: promise, issues, and interpretations. Neuroimage 2013; 80:
360–378.

Ishwaran H, Zarepour M. DIRICHLET PRIOR SIEVES IN FINITE NORMAL MIXTURES .
Stat Sin 2002a

Ishwaran H, Zarepour M. Exact and approximate sum representations for the Dirichlet process.
Can. J. Statistics 2002b; 30: 269–283.

Jezzard P, Balaban RS. Correction for geometric distortion in echo planar images from B0 field
variations. Magn. Reson. Med. 1995; 34: 65–73.

Kaiser RH, Andrews-Hanna JR, Wager TD, Pizzagalli DA. Large-Scale Network Dysfunction in
Major Depressive Disorder: A Meta-analysis of Resting-State Functional Connectivity. JAMA
Psychiatry 2015; 72: 603–611.

Keshishian M, Akkol S, Herrero J, Bickel S, Mehta AD, Mesgarani N. Joint, distributed and
hierarchically organized encoding of linguistic features in the human auditory cortex. Nat. Hum.
Behav. 2023

Klein A, Tourville J. 101 labeled brain images and a consistent human cortical labeling protocol.
Front. Neurosci. 2012; 6: 171.

Kral A, Dorman MF, Wilson BS. Neuronal development of hearing and language: cochlear
implants and critical periods. Annu. Rev. Neurosci. 2019; 42: 47–65.

Laumann TO, Gordon EM, Adeyemo B, Snyder AZ, Joo SJ, Chen M-Y, et al. Functional system
and areal organization of a highly sampled individual human brain. Neuron 2015; 87: 657–670.

Leaver AM, Turesky TK, Seydell-Greenwald A, Morgan S, Kim HJ, Rauschecker JP. Intrinsic
network activity in tinnitus investigated using functional MRI. Hum. Brain Mapp. 2016; 37:
2717–2735.

Lee N, Kim J-M. Dynamic functional connectivity analysis based on time-varying partial
correlation with a copula-DCC-GARCH model. Neurosci. Res. 2021; 169: 27–39.

Lim HH, Lenarz M, Lenarz T. Auditory midbrain implant: a review. Trends Amplif. 2009; 13:
149–180.

Lowe MJ, Mock BJ, Sorenson JA. Functional connectivity in single and multislice echoplanar
imaging using resting-state fluctuations. Neuroimage 1998; 7: 119–132.

Malmierca MS, Ryugo DK. Descending connections of auditory cortex to the midbrain and brain
stem. In: Winer JA, Schreiner CE, editor(s). The Auditory Cortex. Boston, MA: Springer US;
2011. p. 189–208.

Marrelec G, Krainik A, Duffau H, Pélégrini-Issac M, Lehéricy S, Doyon J, et al. Partial
correlation for functional brain interactivity investigation in functional MRI. Neuroimage 2006;
32: 228–237.

Maudoux A, Lefebvre P, Cabay J-E, Demertzi A, Vanhaudenhuyse A, Laureys S, et al. Auditory
resting-state network connectivity in tinnitus: a functional MRI study. PLoS One 2012; 7:
e36222.

McIntosh AR, Gonzalez-Lima F. Structural modeling of functional neural pathways mapped
with 2-deoxyglucose: effects of acoustic startle habituation on the auditory system. Brain Res.
1991; 547: 295–302.

Menon SS, Krishnamurthy K. A comparison of static and dynamic functional connectivities for
identifying subjects and biological sex using intrinsic individual brain connectivity. Sci. Rep.
2019; 9: 5729.

Mihai PG, Moerel M, de Martino F, Trampel R, Kiebel S, von Kriegstein K. Modulation of
tonotopic ventral medial geniculate body is behaviorally relevant for speech recognition. Elife
2019; 8

Miletić S, Bazin PL, Weiskopf N, Van der Zwaag W, Forstmann BU, Trampel R. fMRI protocol
optimization for simultaneously studying small subcortical and cortical areas at 7 T. Neuroimage
2020: 116992.

Moerel M, De Martino F, Uğurbil K, Yacoub E, Formisano E. Processing of frequency and
location in human subcortical auditory structures. Sci. Rep. 2015; 5: 17048.

Moerel M, Yacoub E, Gulban OF, Lage-Castellanos A, De Martino F. Using high spatial
resolution fMRI to understand representation in the auditory network. Prog. Neurobiol. 2021;
207: 101887.

Moore JK. Organization of the human superior olivary complex. Microsc. Res. Tech. 2000

Norman-Haignere SV, Long LK, Devinsky O, Doyle W, Irobunda I, Merricks EM, et al.
Multiscale temporal integration organizes hierarchical computation in human auditory cortex.
Nat. Hum. Behav. 2022; 6: 455–469.

Nourski KV, Steinschneider M, Rhone AE, Kawasaki H, Howard MA, Banks MI. Processing of
auditory novelty across the cortical hierarchy: An intracranial electrophysiology study.
Neuroimage 2018; 183: 412–424.

Pernet CR, McAleer P, Latinus M, Gorgolewski KJ, Charest I, Bestelmeyer PEG, et al. The
human voice areas: Spatial organization and inter-individual variability in temporal and extra-
temporal cortices. Neuroimage 2015; 119: 164–174.

Pickles JO. Auditory pathways: anatomy and physiology. Handb Clin Neurol 2015; 129: 3–25.

Pourahmadi M. High-Dimensional Covariance Estimation: With High-Dimensional Data.
illustrated. John Wiley & Sons; 2013.

Power JD, Cohen AL, Nelson SM, Wig GS, Barnes KA, Church JA, et al. Functional network
organization of the human brain. Neuron 2011; 72: 665–678.

Preibisch C, Castrillón G JG, Bührer M, Riedl V. Evaluation of Multiband EPI Acquisitions for
Resting State fMRI. PLoS One 2015; 10: e0136961.

Rauschecker JP, Tian B. Mechanisms and streams for processing of "what" and "where" in
auditory cortex. Proc. Natl. Acad. Sci. USA 2000; 97: 11800–11806.

Reeves WD, Ahmed I, Jackson BS, Sun W, Brown ML, Williams CF, et al. Characterization of
Resting-State Functional Connectivity Changes in Hypertension by a Modified Difference
Degree Test. Brain Connect. 2023

Reiss LA. Cochlear implants and other inner ear prostheses: today and tomorrow. Curr. Opin.
Physiol. 2020; 18: 49–55.

Ren J, Xu T, Wang D, Li M, Lin Y, Schoeppe F, et al. Individual variability in functional
organization of the human and monkey auditory cortex. Cereb. Cortex 2021; 31: 2450–2465.

Ress D, Chandrasekaran B. Tonotopic organization in the depth of human inferior colliculus.
Front. Hum. Neurosci. 2013; 7: 586.

Rupp K, Hect JL, Remick M, Ghuman A, Chandrasekaran B, Holt LL, et al. Neural responses in
human superior temporal cortex support coding of voice representations. PLoS Biol. 2022; 20:
e3001675.

Sakoğlu U, Pearlson GD, Kiehl KA, Wang YM, Michael AM, Calhoun VD. A method for
evaluating dynamic functional network connectivity and task-modulation: application to
schizophrenia. MAGMA 2010; 23: 351–366.

Sarkar SK, Zhou T, Ghosh D. A General Decision Theoretic Formulation of Procedures
Controlling FDR and FNR from a Bayesian Perspective. Stat Sin 2008; 18

Schofield BR. Projections to the cochlear nuclei from principal cells in the medial nucleus of the
trapezoid body in guinea pigs. J. Comp. Neurol. 1994; 344: 83–100.

Sclocco R, Beissner F, Bianciardi M, Polimeni JR, Napadow V. Challenges and opportunities for
brainstem neuroimaging with ultrahigh field MRI. Neuroimage 2018; 168: 412–426.

Shetty KR, Ridge SE, Kanumuri V, Zhu A, Brown MC, Lee DJ. Clinical and scientific
innovations in auditory brainstem implants. World J. Otorhinolaryngol. Head Neck Surg. 2021;
7: 109–115.

Sigalovsky IS, Melcher JR. Effects of sound level on fMRI activation in human brainstem,
thalamic and cortical centers. Hear. Res. 2006; 215: 67–76.

Sitek KR, Cai S, Beal DS, Perkell JS, Guenther FH, Ghosh SS. Decreased Cerebellar-
Orbitofrontal Connectivity Correlates with Stuttering Severity: Whole-Brain Functional and
Structural Connectivity Associations with Persistent Developmental Stuttering. Front. Hum.
Neurosci. 2016; 10: 190.

Sitek KR, Gulban OF, Calabrese E, Johnson GA, Lage-Castellanos A, Moerel M, et al. Mapping
the human subcortical auditory system using histology, postmortem MRI and in vivo MRI at 7T.
Elife 2019; 8

Skåtun KC, Kaufmann T, Doan NT, Alnæs D, Córdova-Palomera A, Jönsson EG, et al.
Consistent functional connectivity alterations in schizophrenia spectrum disorder: A multisite
study. Schizophr. Bull. 2017; 43: 914–924.

Smith SM. The future of FMRI connectivity. Neuroimage 2012; 62: 1257–1266.

Smith SM, Beckmann CF, Andersson J, Auerbach EJ, Bijsterbosch J, Douaud G, et al. Resting-
state fMRI in the Human Connectome Project. Neuroimage 2013; 80: 144–168.

Smith SM, Vidaurre D, Beckmann CF, Glasser MF, Jenkinson M, Miller KL, et al. Functional
connectomics from resting-state fMRI. Trends Cogn. Sci. (Regul. Ed.) 2013; 17: 666–682.

Sripada C, Angstadt M, Taxali A, Clark DA, Greathouse T, Rutherford S, et al. Brain-wide
functional connectivity patterns support general cognitive ability and mediate effects of
socioeconomic status in youth. Transl. Psychiatry 2021; 11: 571.

Tabas A, Kiebel S, Marxen M, von Kriegstein K. Fast frequency modulation is encoded
according to the listener expectations in the human subcortical auditory pathway. arXiv 2021

Van Essen DC, Ugurbil K, Auerbach E, Barch D, Behrens TEJ, Bucholz R, et al. The Human
Connectome Project: a data acquisition perspective. Neuroimage 2012; 62: 2222–2231.

Varoquaux G, Schwartz Y, Pinel P, Thirion B. Cohort-level brain mapping: learning cognitive
atoms to single out specialized regions. Inf. Process. Med. Imaging 2013; 23: 438–449.

de Vos F, Koini M, Schouten TM, Seiler S, van der Grond J, Lechner A, et al. A comprehensive
analysis of resting state fMRI measures to classify individual patients with Alzheimer's disease.
Neuroimage 2018; 167: 62–72.

Wang Y, Kang J, Kemmer PB, Guo Y. An efficient and reliable statistical method for estimating
functional connectivity in large scale brain networks using partial correlation. Front. Neurosci.
2016; 10: 123.

Warnick R, Guindani M, Erhardt E, Allen E, Calhoun V, Vannucci M. A Bayesian Approach for
Estimating Dynamic Functional Network Connectivity in fMRI Data. J. Am. Stat. Assoc. 2018;
113: 134–151.

Wilson KC, Kornisch M, Ikuta T. Disrupted functional connectivity of the primary auditory
cortex in autism. Psychiatry Research: Neuroimaging 2022: 111490.

Winer JA. Decoding the auditory corticofugal systems. Hear. Res. 2005; 207: 1–9.

Xu T, Opitz A, Craddock RC, Wright MJ, Zuo X-N, Milham MP. Assessing variations in areal
organization for the intrinsic brain: from fingerprints to reliability. Cereb. Cortex 2016; 26:
4192–4211.

Yu Z, Guindani M, Grieco SF, Chen L, Holmes TC, Xu X. Beyond t test and ANOVA:
applications of mixed-effects models for more rigorous statistical analysis in neuroscience
research. Neuron 2022; 110: 21–35.

Zatorre R, Belin P, Penhune VB. Structure and function of auditory cortex: music and speech.
Trends Cogn. Sci. (Regul. Ed.) 2002; 6: 37–46.

Zhang J, Chen Y-C, Feng X, Yang M, Liu B, Qian C, et al. Impairments of thalamic resting-state
functional connectivity in patients with chronic tinnitus. Eur. J. Radiol. 2015; 84: 1277–1284.

Zhang L, Guindani M, Versace F, Vannucci M. A spatio-temporal nonparametric Bayesian
variable selection model of fMRI data for clustering correlated time courses. Neuroimage 2014;
95: 162–175.

Zuo X-N, Xu T, Milham MP. Harnessing reliability for neuroscience research. Nat. Hum. Behav.
2019; 3: 768–771.

The mammalian auditory pathway: neurophysiology. New York, NY: Springer New York; 1992.

The Mammalian Auditory Pathway: Neuroanatomy. New York, NY: Springer New York; 1992.