Title Page

Emergence of plasmid-mediated RND-type efflux pump gene cluster *tmexCD-*

toprJ in *Shewanella xiamenensis* in a water environment

Trung Duc Dao[1], Taichiro Takemura[1#], Ikuro Kasuga[2,3], Aki Hirabayashi[4],

Nguyen Thi Nga[1], Pham Hong Quynh Anh[1], Nguyen Dong Tu[5], Le Thi Trang[5],

Hoang Huy Tran[5], Keigo Shibayama[6], Futoshi Hasebe[1], and Masato Suzuki[4#]

[1]Vietnam Research Station, Center for Infectious Disease Research in Asia and

Africa, Institute of Tropical Medicine, Nagasaki University, Nagasaki, Japan

[2]Vietnam-Japan University, Vietnam National University, Hanoi, Vietnam

[3]Research Center for Advanced Science and Technology, The University of

Tokyo, Tokyo, Japan

[4]Antimicrobial Resistance Research Center, National Institute of Infectious

Diseases, Tokyo, Japan

[5]National Institute of Hygiene and Epidemiology, Hanoi, Vietnam

[6]Nagoya University Graduate School of Medicine, Nagoya, Japan

[#]Corresponding authors. E-mail address: taichiro@nagasaki-u.ac.jp (T.

Takemura); suzuki-m@nih.go.jp (M. Suzuki)

Keywords: tigecycline, *tmexCD-toprJ*, carbapenem, *bla*$_{NDM-1}$, *Shewanella*

Abstract

The emergence of the mobile resistance-nodulation-division (RND)-type efflux

pump *tmexCD-toprJ* gene cluster that confers multidrug resistance (MDR),

including tigecycline resistance, in gram-negative bacteria poses a global public

health threat. However, the spread of such clinically important antimicrobial

resistance genes (ARGs) in the natural environment has not yet been well

investigated. In this study, we investigated MDR aquatic bacteria in Vietnam. A

carbapenem- and tigecycline-resistant *Shewanella xiamenensis* isolate NUITM-

VS2 was obtained from urban drainage in Hanoi, Vietnam, in October 2021. *S.*

xiamenensis NUITM-VS2 showed resistance to most antimicrobials tested,

including   tigecycline,   tetracyclines,   carbapenems,   cephalosporins,

fluoroquinolone, and aminoglycosides. Whole-genome analysis was performed

by long- and short-read sequencing, resulting in the complete genome sequence

consisting of one chromosome and five plasmid sequences. ARGs and plasmid

replicons in the genome were detected using ResFinder with the custom ARG

database, including all known tigecycline resistance genes, and PlasmidFinder,

respectively. A 152.2-kb IncC plasmid, pNUITM-VS2_2, co-carried two mobile

tigecycline resistance genes, *tet*(X4) and *tmexC3.2D3.2-toprJ1*. In addition, a

24.8-kb untypeable plasmid, pNUITM-VS2_4, carried the carbapenemase gene

bla$_{NDM-1}$. pNUITM-VS2_2 was transferred to *Escherichia coli* by conjugation,

which   simultaneously   conferred   high-level   resistance   against   many

antimicrobials, including tigecycline. To the best of our knowledge, this is the first

report of the detection of the mobile RND-type efflux pump gene cluster *tmexCD-*

toprJ in *Shewanella* species. Our results provide genetic evidence of the

complexity of the dynamics of clinically important ARGs among aquatic bacteria,

which could be important reservoirs for ARGs in the natural environment.

Main text

Introduction

Tigecycline is a semi-synthetic antimicrobial agent, which was the first drug of the glycylcycline class developed for the last-line treatment of infections caused by multidrug-resistant (MDR) bacterial pathogens, especially carbapenem-resistant *Enterobacterales*. However, due to the widespread use of this antimicrobial in clinical settings, novel antimicrobial resistance genes (ARGs), including the mobile tigecycline-inactivation enzyme gene *tet*(X) and the resistance-nodulation-division (RND)-type efflux pump gene cluster *tmexCD-toprJ*, both conferring high-level tigecycline resistance, have rapidly been disseminated between diverse gram-negative bacteria (1). This not only poses an economic burden to the healthcare system but also increases mortality risks in patients by disabling the availability of last-resort antimicrobial agents, such as tigecycline.

Shewanella species are gram-negative anaerobes found in marine and freshwater environments, which belong to the family Shewanellaceae within the class Gammaproteobacteria. *Shewanella* are natural reservoirs of *bla*$_{OXA-48}$-like carbapenemase genes, such as *bla*$_{OXA-48}$ and *bla*$_{OXA-181}$, encoded on the chromosomes (2, 3). In 2022, we reported a *Shewanella xiamenensis* isolate obtained from a water environment in Vietnam, which harbored *bla*$_{OXA-48}$ and *tet*(X4) on the chromosome (4).

In 2021, we reported a *Klebsiella aerogenes* isolate harboring a self-transferable incompatibility group C (IncC)–IncX3 hybrid plasmid co-carrying *bla*$_{NDM-4}$, *tet*(X4), and *tmexCD3-toprJ1* [initially named *tmexCD3-toprJ3* (5) and

later renamed (6)] from a water environment in Vietnam (7). Although the

Flavobacteriaceae family is estimated as the natural reservoir of *tet*(X) genes (8),

limited information is available on the distribution of *tmexCD-toprJ* in the natural

environment.

In this study, we report an *S. xiamenensis* isolate obtained from the aquatic

environment in Vietnam harboring *tmexCD-toprJ*, representing the first such

report for *Shewanella* species. The isolate showed carbapenem and tigecycline

resistance, and harbored *bla*$_{OXA-181}$ on the chromosome, *bla*$_{NDM-1}$ on a 24.8-kb

untypeable plasmid, and *tmexC3.2D3.2-toprJ1* together with *tet*(X4) on a self-

transferable 152.2-kb IncC plasmid.

Results and discussion

An environmental carbapenem- and tigecycline-resistant bacterial isolate, NUITM-VS2, was identified as *Shewallena* species using MALDI Biotyper. The minimum inhibitory concentrations (MICs) of tigecycline, tetracyclines, carbapenems, cephalosporins, and gentamycin were all above 128 mg/L (R); and those of ciprofloxacin, tobramycin, and streptomycin were 4, 16, and 64 mg/L (R), respectively; whereas that of amikacin was 0.5 mg/L (S) (Table 1).

Whole-genome analysis using the Illumina and Oxford Nanopore Technologies (ONT) sequencing platforms resulted in the complete genome sequence consisting of one chromosome and five plasmids, designated pNUITM-VS2_1, pNUITM-VS2_2, pNUITM-VS2_3, pNUITM-VS2_4, and pNUITM-VS2_5 (accession nos. AP026732, AP026733, AP026734, AP026735, AP026736, and AP026737), respectively. Genome annotation and average nucleotide identity (ANI) analyses revealed that the 4.9-Mb chromosome with a 46.4% GC content contains 4,100 coding sequences (CDSs), in which $bla_{OXA-181}$ was detected by ResFinder, showed 97.3% identity to the genome sequence of *S. xiamenensis* JCM 16212^T (accession no. BMOP01000000).

The 152.2-kb pNUITM-VS2_2 plasmid of *S. xiamenensis* NUITM-VS2 contained a replicon classified to IncC and co-carried two mobile tigecycline resistance genes, *tet*(X4) and a variant of *tmexCD3-toprJ1*, whose sequence was a perfect match to that reported in the *K. aerogenes* pNUITM-VK5_mdr plasmid (accession no. LC633285) isolated from the same environmental location as NUITM-VS2 in Hanoi, Vietnam, in February 2021 (7) (Fig. 1). The CDSs of *tmexCD-toprJ* in pNUITM-VS2_2 and pNUITM-VK5_mdr showed high identity to

tmexCD3-toprJ1 of *Proteus mirabilis* RGF134-1 (accession no. CP066833) (5)

isolated from a pig in China in 2019. The identity for *tmexC3* was 99.7%

(1,161/1,164 nt), with three amino acid substitutions (Q187H, T256M, and

A386T). For *tmexD3*, the identity was 99.9% (3,133/3,135 nt), including two

amino acid substitutions (V610L and L611F). For *toprJ1*, the identity was 100%

(1,434/1,434 nt). Therefore, we named the variant identified in this and previous

studies *tmexC3.2D3.2-toprJ1*, consisting of *tmexC3.2*, *tmexD3.2*, and *toprJ1*.

Moreover, the carbapenemase gene bla_{NDM-1}, fluoroquinolone resistance gene

qnrVC6, and aminoglycoside resistance genes *aac(3)-IId* and *aac(6')-IIa* were

encoded in another 24.8-kb untypeable plasmid, pNUITM-VS2_4, that is highly

identical to the *S. xiamenensis* 26.8-kb untypeable plasmid pSx1 (100% identity

with 96% of the region of accession no. CP013115) isolated from the environment

in Algeria in 2012 (9). pSx1 was previously classified as an IncP-2 plasmid (10),

which was subsequently found to be a misidentification (11). The replicon type of

pNUITM-VS2_4 and pSx1 could not be identified by PlasmidFinder; thus, this

plasmid type is presumed to be unique to *Shewanella* species.

On pNUITM-VS2_2 in *S. xiamenensis* NUITM-VS2, *tet*(X4) was flanked by two

mobile genetic elements, IS*Vsa3* and IS*91* family transposase, while

tmexC3.2D3.2-toprJ1 was flanked by Δ*tnpA* and IS*L3*. Interestingly, pNUITM-

VS2_2 was almost identical to the IncC backbone of the *K. aerogenes* IncC–

IncX3 hybrid plasmid pNUITM-VK5_mdr (100% identity with 38% of the region of

accession no. LC633285) (7) (Fig. 1). The rest of the IncX3 backbone of pNUITM-

VK5_mdr carried additional ARGs, such as bla_{NDM-4} and *qnrS1*, conferring

resistance to carbapenem and fluoroquinolone, respectively, suggesting that the

MDR IncC–IncX3 hybrid plasmid pNUITM-VK5_mdr resulted from fusion of a

tigecycline-resistance IncC plasmid such as pNUITM-VS2_2 with a carbapenem-

resistance IncX3 plasmid.

A bacterial conjugation assay using *Escherichia coli* J53 as the recipient strain

showed that *S. xiamenensis* NUITM-VS2 transferred pNUITM-VS2_2 to J53 at a

frequency of 6.4×10^{-5} after overnight co-culture at 37°C. PCR confirmed that the

transconjugant   strain   (J53/pNUITM-VS2_2)   co-harbored   *tet*(X4)   and

tmexC3.2D3.2-toprJ1, and was resistant to tigecycline and tetracyclines (Table

1).

Conclusions

In conclusion, the complete genome sequence of the carbapenem- and

tigecycline-resistant isolate *S. xiamenensis* NUITM-VS2 obtained from a water

environment in Vietnam was characterized. This isolate co-carried two mobile

tigecycline resistance genes, *tet*(X4) and *tmexC3.2D3.2-toprJ1*, in the self-

transferable 152.2-kbp IncC plasmid. This plasmid corresponds to the IncC

backbone of the IncC–IncX3 hybrid plasmid pNUITM-VK5_mdr identified in *K.*

aerogenes in our previous report that was isolated from the same area (7),

suggesting that they might share a common ancestor. Our results provide genetic

evidence that clinically important ARGs have been disseminated though plasmids,

such as broad-host-range IncC plasmids, and their fusion plasmids among

environmental bacteria, such as *Shewanella* species, in aquatic environments.

Materials and methods

Bacterial isolation and antimicrobial susceptibility testing

Carbapenem- and tigecycline-resistant *S. xiamenensis* NUITM-VS2 was isolated from the Kim-Nguu River in Hanoi, Vietnam, in October 2021. Environmental water sample was collected and cultured using Luria-Bertani (LB) broth containing 4 mg/L of meropenem at 37°C overnight, and then further selected and isolated using CHROMagar COL-APSE (CHROMagar Microbiology) containing 4 mg/L of tigecycline. Bacterial species identification was performed using MALDI Biotyper (Bruker). Antimicrobial susceptibility testing using *Escherichia coli* ATCC 25922 as quality control was performed with agar dilution according to the CLSI 2020 guidelines. MIC breakpoints of antimicrobials (S: susceptible; I: intermediate; R: resistant) for other non-*Enterobacterales* were also determined according to the CLSI 2020 guidelines.

Whole-genome sequencing and subsequent bioinformatics analysis

Whole-genome sequencing of *S. xiamenensis* NUITM-VS2 was performed using MiSeq (Illumina) with MiSeq Reagent Kit v2 (300-cycle) and MinION (Oxford Nanopore Technologies; ONT) with the R9.4.1 flow cell. The library for Illumina sequencing (paired-end, insert size of 300–800 bp) was prepared using the Nextera XT DNA Library Prep Kit, and the library for MinION sequencing was prepared using the Rapid Barcoding Kit (SQK-RBK004). ONT reads were base-called using Guppy v5.0.11, in the super-accuracy mode. Hybrid de novo assembly with both Illumina and ONT reads was performed using Unicycler v0.4.8.0 (https://github.com/rrwick/Unicycler) (12) with default parameters.

Genome annotation and average nucleotide identity (ANI) analyses were

performed using the DFAST server (https://dfast.nig.ac.jp) (13). ARGs and

plasmid   replicons   were   detected   using   ResFinder   v4.1

(https://bitbucket.org/genomicepidemiology/resfinder/) (14) with the custom ARG

database, including all known tigecycline resistance genes, and PlasmidFinder

v2.1 (https://bitbucket.org/genomicepidemiology/plasmidfinder/) (15) with the

default parameters, respectively. Type IV secretion system (T4SS)-associated

genes   involved   in   conjugation   were   detected   by   CONJscan   v1.0.5

(https://github.com/macsy-models/TXSS) with default parameters (16). Linear

comparison of sequence alignment was performed using BLAST and visualized

by Easyfig v.2.2.2 (http://mjsull.github.io/Easyfig/) (17).

Bacterial conjugation assay

A bacterial conjugation assay was performed as follows. LB broth cultures of

the donor *S. xiamenensis* NUITM-VS2 and the recipient azide-resistant *E. coli*

J53 (ATCC BAA-2731, F⁻ *met pro* Aziʳ) were mixed in a 1:10 ratio, spotted onto

MacConkey agar, and then incubated at 37°C overnight. Subsequently, the

mixed cells, including transconjugants, were suspended in LB broth and then

plated onto MacConkey agar containing 1 mg/L of tigecycline and 100 mg/L of

sodium azide after 10-fold serial dilution and incubated at 37°C overnight.

tet(X) and *tmexCD-toprJ* in transconjugants were detected by colony PCR

using the following primer sets: tetX_F, CCCGAAAATCGWTTTGACAATCCTG;

tetX_R,   GTTTCTTCAACTTSCGTGTCGGTAAC;   tmexC_F,

TGGCGGGGATCGTGCTCAAGCGCAC;   tmexC_R,

CAGCGTGCCCTTGCKCTCGATATCG. The PCR amplification consisted of the

initial denaturation for 3 min at 95°C, followed by 35 cycles of denaturation for

30 s at 95°C, annealing for 30 s at 54°C, and extension for 30 s at 72°C, and then

the final extension for 5 min at 72°C. Genomic DNAs of *K. aerogenes* strain

NUITM-VK5 co-harboring *tet*(X4) and *tmexC3.2D3.2-toprJ1* (7) was used as

positive controls. Purified water was used as a negative control.

Nucleotide sequence accession numbers

The complete genome sequence of *S. xiamenensis* NUITM-VS2 has been deposited at GenBank/EMBL/DDBJ under accession nos. AP026732 (chromosome), AP026733 (pNUITM-VS2_1), AP026734 [pNUITM-VS2_2 co-carrying *tet*(X4) and *tmexC3.2D3.2-toprJ1*], AP026735 (pNUITM-VS2_3), AP026736 (pNUITM-VS2_4 carrying bla_{NDM-1}), and AP026737 (pNUITM-VS2_5).

Funding

This work was supported by grants (JP22gm1610003, JP22fk0108133, JP22fk0108139, JP22fk0108642, JP22wm0225004, JP22wm0225008, JP22wm0225022, JP22wm0325003, JP22wm0325022, and JP22wm0325037 to M. Suzuki; JP22fk0108132 and JP22wm0225008 to I. Kasuga; JP22wm0125006 and JP22wm0225008 to F. Hasebe; JP22fk0108604 and JP22gm1610003 to K. Shibayama) from the Japan Agency for Medical Research and Development (AMED), grants (20K07509 and 21K18742 to M. Suzuki; 21K18742 to T. Takemura; 19K21984 and 21K18742 to I. Kasuga; 21K15440 to A. Hirabayashi) from the Ministry of Education, Culture, Sports, Science and Technology (MEXT), Japan, and a grant (MS.108.02-2017.320 to H. H. Tran) from the National Foundation for Science and Technology Development (NAFOSTED), Vietnam.

Competing interests

None declared.

Ethical approval

235   Not required.

237 **Table 1**

Strain	MIC (mg/L)												
	TIG	MIN	DOX	TET	IPM	MEM	CTX	CAZ	CIP	AMK	GEN	TOB	STR
S. xiamenensis NUITM-VS2	>128	>128	>128	>128	>128	>128	>128	>128	4	0.5	>128	16	64
E. coli J53/pNUITM-VS2_2	128	128	128	128	0.25	0.016	0.032	0.25	0.016	0.5	0.5	0.25	0.5
E. coli J53	0.25	2	4	1	0.25	0.016	0.032	0.25	0.016	0.5	0.5	0.25	0.5
E. coli ATCC 25922	0.25	1	2	1	0.25	0.016	0.064	0.5	0.008	1	0.5	0.5	0.5

Legends

Table 1. MICs of antimicrobials against *S. xiamenensis* NUITM-VS2 and the

transconjugant of *E. coli* J53. Abbreviations: TIG, tigecycline; MIN, minocycline;

DOX, doxycycline; TET, tetracycline; IPM, imipenem; MEM, meropenem; CTX,

cefotaxime; CAZ, ceftazidime; CIP, ciprofloxacin; AMK, amikacin; GEN,

gentamicin; TOB, tobramycin; STR, streptomycin.

Fig. 1. Linear comparison of *tet*(X4) and *tmexC3.2D3.2-toprJ1*–carrying plasmids

S. xiamenensis pNUITM-VS2_2 and *K. aerogenes* pNUITM-VK5_mdr isolated in

Vietnam in 2021. Red, yellow, blue, green, and gray indicate carbapenem and

tetracycline resistance genes (CRG/TRG), other AMR genes (ARG), mobile gene

elements (MGE), type IV secretion system (T4SS)-associated genes involved in

conjugation, and other coding sequences (Other), respectively. The blue color in

comparison of sequences indicates % of identity.

References

References

1. Anyanwu MU, Nwobi OC, Okpala COR, Ezeonu IM.

Mobile Tigecycline Resistance: An Emerging Health Catastrophe Requiring

Urgent One Health Global Intervention.

Front Microbiol. 2022 Aug 1;13:808744

2. Tacão M, Araújo S, Vendas M, Alves A, Henriques I.

Shewanella species as the origin of bla_{OXA-48} genes: insights into gene

diversity, associated phenotypes and possible transfer mechanisms.

Int J Antimicrob Agents. 2018 Mar;51(3):340-348.

3. Nguyen NT, Takemura T, Pham AHQ, Tran HT, Vu KCT, Tu ND, Huong LT,

Cuong NT, Kasuga I, Hasebe F, Suzuki M.

Whole-genome sequencing and comparative genomic analysis of *Shewanella*

xiamenensis strains carrying bla_{OXA-48}-like genes isolated from a water

environment in Vietnam.

J Glob Antimicrob Resist. 2020 Jun;21:272-274.

4. Dao TD, Kasuga I, Hirabayashi A, Nguyen DT, Tran HT, Vu H, Pham LTN, Vu

TMH, Hasebe F, Nguyen HT, Thi TL, Tran HH, Shibayama K, Takemura T,

Suzuki M.

Emergence of mobile tigecycline resistance gene *tet*(X4)-harbouring

Shewanella xiamenensis in a water environment.

J Glob Antimicrob Resist. 2022 Mar;28:140-142.

5. Wang Q, Peng K, Liu Y, Xiao X, Wang Z, Li R.

Characterization of TMexCD3-TOprJ3, an RND-Type Efflux System

Conferring Resistance to Tigecycline in *Proteus mirabilis*, and Its Associated

Integrative Conjugative Element.

Antimicrob Agents Chemother. 2021 Jun 17;65(7):e0271220.

6. Wang CZ, Gao X, Lv LC, Cai ZP, Yang J, Liu JH.

Novel tigecycline resistance gene cluster *tnfxB3-tmexCD3-toprJ1b* in *Proteus*

spp. and *Pseudomonas aeruginosa*, co-existing with *tet*(X6) on an SXT/R391

integrative and conjugative element.

J Antimicrob Chemother. 2021 Nov 12;76(12):3159-3167.

7. Hirabayashi A, Dao TD, Takemura T, Hasebe F, Trang LT, Thanh NH, Tran HH,

Shibayama K, Kasuga I, Suzuki M.

A Transferable IncC-IncX3 Hybrid Plasmid Cocarrying *bla*$_{NDM-4}$, *tet*(X), and

tmexCD3-toprJ3 Confers Resistance to Carbapenem and Tigecycline.

mSphere. 2021 Aug 25;6(4):e0059221.

8. Zhang R, Dong N, Shen Z, Zeng Y, Lu J, Liu C, Zhou H, Hu Y, Sun Q, Cheng

Q, Shu L, Cai J, Chan EW, Chen G, Chen S.

Epidemiological and phylogenetic analysis reveals *Flavobacteriaceae* as

potential ancestral source of tigecycline resistance gene *tet*(X).

Nat Commun. 2020 Sep 16;11(1):4648.

9. Yousfi K, Touati A, Lefebvre B, Fournier É, Côté JC, Soualhine H, Walker M,

Bougdour D, Tremblay C, Bekal S.

A Novel Plasmid, pSx1, Harboring a New Tn*1696* Derivative from Extensively

Drug-Resistant *Shewanella xiamenensis* Encoding OXA-416.

Microb Drug Resist. 2017 Jun;23(4):429-436.

10. Jiang X, Yin Z, Yuan M, Cheng Q, Hu L, Xu Y, Yang W, Yang H, Zhao Y, Zhao

X, Gao B, Dai E, Song Y, Zhou D.

Plasmids of novel incompatibility group Inc$_{pRBL16}$ from *Pseudomonas* species.

J Antimicrob Chemother. 2020 Aug 1;75(8):2093-2100.

11. Shintani M, Suzuki H, Nojiri H, Suzuki M.

Precise classification of antimicrobial resistance-associated IncP-2

megaplasmids for molecular epidemiological studies on *Pseudomonas*

species.

J Antimicrob Chemother. 2022 Mar 31;77(4):1203-1205.

12. Wick RR, Judd LM, Gorrie CL, Holt KE.

Unicycler: Resolving bacterial genome assemblies from short and long

sequencing reads.

PLoS Comput Biol. 2017 Jun 8;13(6):e1005595.

13. Tanizawa Y, Fujisawa T, Nakamura Y.

DFAST: a flexible prokaryotic genome annotation pipeline for faster genome

publication.

Bioinformatics. 2018 Mar 15;34(6):1037-1039.

14. Bortolaia V, Kaas RS, Ruppe E, Roberts MC, Schwarz S, Cattoir V, Philippon

A, Allesoe RL, Rebelo AR, Florensa AF, Fagelhauer L, Chakraborty T,

Neumann B, Werner G, Bender JK, Stingl K, Nguyen M, Coppens J, Xavier

BB, Malhotra-Kumar S, Westh H, Pinholt M, Anjum MF, Duggett NA, Kempf I,

Nykäsenoja S, Olkkola S, Wieczorek K, Amaro A, Clemente L, Mossong J,

Losch S, Ragimbeau C, Lund O, Aarestrup FM.

ResFinder 4.0 for predictions of phenotypes from genotypes.

J Antimicrob Chemother. 2020 Dec 1;75(12):3491-3500.

15. Carattoli A, Hasman H.

PlasmidFinder and In Silico pMLST: Identification and Typing of Plasmid

Replicons in Whole-Genome Sequencing (WGS).

Methods Mol Biol. 2020;2075:285-294.

16. Cury J, Abby SS, Doppelt-Azeroual O, Néron B, Rocha EPC.

Identifying Conjugative Plasmids and Integrative Conjugative Elements with

CONJscan.

Methods Mol Biol. 2020;2075:265-283.

17. Sullivan MJ, Petty NK, Beatson SA.

Easyfig: a genome comparison visualizer.

Bioinformatics. 2011 Apr 1;27(7):1009-10.

Fig. 1