Pupillary Dynamics of Mice Performing a Pavlovian Delay Conditioning Task Reflect Reward-Predictive Signals.

AUTHORS:

Kota Yamada[1,2,*], Koji Toda[1*]

AFFILIATIONS:

1 Department of Psychology, Keio University, Tokyo, JAPAN

2 Japan Society for Promotion of Science, Tokyo, JAPAN

*CORRESPONDENCE:

Kota Yamada[†]

Department of Psychology, Keio University

Mita 2-15-45, Minato-ku, Tokyo, JAPAN

Email: haroldthebarrel.yk@gmail.com

Koji Toda, Ph.D.

Department of Psychology, Keio University

Mita 2-15-45, Minato-ku, Tokyo, JAPAN

Email: koji@keio.jp

[†]Lead contact

Abstract

Pupils can signify various internal processes and states, such as attention, arousal, and working memory. Changes in pupil size have been associated with learning speed, prediction of future events, and deviations from the prediction in human studies. However, the detailed relationships between pupil size changes and prediction are unclear. We explored pupil size dynamics in mice performing a Pavlovian delay conditioning task. A head-fixed experimental setup combined with deep-learning-based image analysis enabled us to reduce spontaneous locomotor activity and to track the precise dynamics of pupil size of behaving mice. By setting up two experimental groups, one for which mice were able to predict reward in the Pavlovian delay conditioning task and the other for which mice were not, we demonstrated that the pupil size of mice is modulated by reward prediction and consumption, as well as body movements, but not by unpredicted reward delivery. Furthermore, we clarified that pupil size is still modulated by reward prediction even after the disruption of body movements by intraperitoneal injection of haloperidol, a dopamine D2 receptor antagonist. These results suggest that changes in pupil size reflect reward prediction signals. Thus, we provide important evidence to reconsider the neuronal circuit involved in computing reward prediction error. This integrative approach of behavioral analysis, image analysis, pupillometry, and pharmacological manipulation will pave the way for understanding the psychological and neurobiological mechanisms of reward prediction and the prediction errors essential to learning and behavior.

Manuscript contributions to the field

Predicting upcoming events is essential for the survival of many animals, including humans. Accumulating evidence suggests that pupillary responses reflect autonomic activity and are modulated by noradrenergic, cholinergic, and serotonergic neurotransmission. However, the relationships between pupillary responses, reward prediction, and reward prediction errors remain unclear. This study examined changes in pupil size while water-deprived mice performed a Pavlovian delay conditioning task using a head-fixed setup. The head-fixed experimental setup, combined with deep-learning-based image analysis, enabled us to reduce spontaneous locomotor activity and to track the precise dynamics of the licking response and the pupil size of behaving mice. A well-controlled, rigid behavioral experimental design allowed us to investigate the modulation of behavioral states induced by reward prediction. While pharmacological manipulation might affect pupil size, the combined approach of pupillometry and pharmacological manipulation allowed us to differentiate reward prediction signals and signals modulated by body movements. We revealed that the changes in pupil size (1) reflect reward prediction signals and (2) do not reflect signals of reward prediction error. These results provide novel insights into the neuronal circuitry potentially involved in computing reward prediction errors. The integrative approach of behavioral analysis, image analysis, pupillometry, and pharmacological manipulation used in this study will pave the way for understanding the psychological and neurobiological mechanisms of prediction and the prediction errors essential in learning and behavior.

71 ## Keywords

72 dopamine, reward prediction error, pupil, licking, Pavlovian conditioning, mice

Introduction

Predicting future events from current observations helps organisms to obtain rewards and avoid aversive events in a given environment. Pavlovian conditioning is a widely used experimental procedure for investigating the predictive abilities of animals. For example, water-restricted mice are exposed to an auditory stimulus, followed by the water reward. After several training sessions, mice develop anticipatory responses to the auditory stimulus. Pavlovian conditioning involves both behavioral and physiological responses. In appetitive conditioning, a conditioned approach response to a stimulus that signals food (Hearst and Jenkins, 1974) or to the location where the food is presented (Boakes, 1977) is observed. In fear conditioning, freezing responses (Estes and Skinner, 1941) are induced by a stimulus that signals aversive events. Physiological responses, such as salivary response, changes in skin conductance, heart rate, pupil dilation, body temperature, and respiration, are also acquired through Pavlovian conditioning (Esteves et al., 1994; Leuchs et al., 2017; Lonsdorf et al., 2017; Notterman et al., 1952; Öhman et al., 1976; Ojala and Bach, 2020; Pavlov, 1927; Pietrock et al., 2019; Wood and Obrist, 1964). Pavlovian conditioning includes several response types: preparatory, consummatory, and opponent responses to unconditioned responses (Konorski, 1967; Solomon and Corbit, 1974). Thus, accumulating evidence in the field of psychological and physiological studies of animal learning demonstrates that Pavlovian conditioning is a valuable technique for studying the function and underlying mechanism of prediction.

Although the use of pupillometry in Pavlovian conditioning dates back more than half a century, its reliability as an indicator of learning has recently been reevaluated (Finke et al., 2021). It has been reported that changes in pupil size occur as a reactive response to a conditioned stimulus in fear and appetitive conditioning in humans (Leuchs et al., 2017; Lonsdorf et al., 2017; Ojala and Bach, 2020; Pietrock et al., 2019). The relationship between pupil size and theories of learning, such as

prediction errors in temporal difference learning (Sutton and Barto, 2018), the
Mackintosh and Rescorla-Wagner models (Mackintosh, 1975; Rescorla and Wagner,
1972), as well as attention to the stimuli in the Pearce-Hall model (Pearce and Hall,
1980) have also been discussed (Koenig et al., 2018; Pietrock et al., 2019; Vincent
et al., 2019). Changes in pupil size are associated with various internal states,
including arousal level, attention, working memory, social vigilance, the value of
alternatives in choice tasks, and uncertainty in diverse research fields (Ebitz et al.,
2014; Ebitz and Platt, 2015; Finke et al., 2021; Joshi and Gold, 2020; Larsen and
Waters, 2018; Van Slooten et al., 2018; Vincent et al., 2019; Zénon, 2019). These
findings suggest that pupil size is a reactive response to a conditioned stimulus and
an active modulator of sensorimotor processing that affects prediction (Ebitz and
Moore, 2019).

Despite the potential usefulness of pupillometry in understanding the
neurobiological mechanisms underlying behavior, there have been only a few
attempts to record pupillary changes in rodent research (Cazettes et al., 2021; Lee
and Margolis, 2016; Nelson and Mooney, 2016; Privitera et al., 2020; Reimer et al.,
2014; Wang et al., 2022). This can be attributed to two technical issues. First,
conventional behavioral tasks designed for rodents use experimental apparatuses
in which animals move freely, making it impossible to precisely record pupil size.
Second, body movements also modulate pupil size (Cazettes et al., 2021; Nelson
and Mooney, 2016). This makes its interpretations more complex than human
studies that allow participants to remain in the experimental setup. Recent
experimental setups and advances in machine learning have enabled researchers
to overcome these technical limitations. By combining a head-fixed setup with image
analysis techniques such as DeepLabCut (Mathis et al., 2018; Nath et al., 2019),
several studies have successfully quantified pupils and eyelid size of mice
performing behavioral tasks (Kaneko et al., 2022; Privitera et al., 2020).

This study explored the dynamics of licking and pupillary responses of mice
performing a Pavlovian delay conditioning task with a head-fixed experimental setup.

Pupil size is known to be increased by the presentation of the cue in appetitive and
aversive conditioning in human participants as well as rodent subjects, supporting
the view that animals gain their arousal by the cue presentation (Finke et al., 2021;
Pietrock et al., 2019). In Experiment 1, we trained head-fixed mice on the Pavlovian
delayed conditioning task. An auditory stimulus was presented before a sucrose
solution reward was delivered while recording their licking and pupil response. In this
task, we designed contingent and non-contingent groups to manipulate the
predictability of the delivery of the sucrose solution by the auditory stimulus. In the
contingent group, the auditory stimulus signaled the arrival of the sucrose solution,
such that the delivery of the sucrose solution immediately followed the auditory
stimulus. In the non-contingent group, the auditory stimulus provided no predictive
information about the arrival of the sucrose solution, as the presentation of the
auditory stimulus and the delivery of the sucrose solution were randomized. We
investigated the dynamics of licking and pupillary responses in predictable and
unpredictable situations by measuring licking and pupillary responses while the mice
performed the Pavlovian delay conditioning tasks. In addition, bout analysis of the
licking responses allowed us to unveil the detailed relationship between the licking
and pupillary responses. In Experiment 2, we examined the pupil dynamics by
suppressing body movements with systemic administration of haloperidol, an
antagonist of dopamine D2 receptors that have been reported to inhibit anticipatory
and consummatory licking (Fowler and Mortell, 1992; Liao and Ko, 1995) and
spontaneous movements in an open-field experiment (Arruda et al., 2008; Bernardi
et al., 1981; Conceição and Frussa-Filho, 1996; Strömbom, 1977).

Methods

Subjects

Eight adult male C57BL/6J mice were used. All mice were purchased from Nippon Bio-Supp.Center and bred in the breeding room provided in the laboratory. All mice were naive and eight weeks old at the start of the experiment. The mice were maintained on a 12:12 h light cycle. All the experiments were conducted during the dark phase of the light cycle. The mice had no access to water in their home cage and were provided with water only during experimental sessions. The mice were allowed to consume sufficient sucrose solution during the experiment. The mice's body weight was monitored daily (21.7 ± 2.1 g before the experiment). They were provided additional access to water at their home cage if their body weight fell below 85% of their normal body weight measured before the experiment. The mice were allowed to feed freely in their home cages. The experimental and housing protocols adhered to the Japanese National Regulations for Animal Welfare and were approved by the Animal Care and Use Committee of Keio University.

Surgery

Mice were anesthetized with 1.0% to 2.5% isoflurane mixed with room air and placed in a stereotactic frame (942WOAE, David Kopf Instruments, Tujunga, CA, USA). A head post (H.E. Parmer Company, Nashville, TN, USA) was fixed at the surface of the skull, aligning the midline using dental cement (Product #56849, 3M Company, Saint Paul, MN, USA) to head-fix the mice during the experiment. The mice were group-housed (four mice per cage) before the experiments, and a recovery time of two weeks was scheduled between the surgery and experiment commencement.

Procedure

Mice were habituated to a head-fixed experimental setup (Figure 1A; Kaneko et al.,
2022; Toda et al., 2017; Yamamoto et al., 2022) the day before the experiment
commenced. During habituation, mice were head-fixed in the apparatus with dim
light and randomly presented with 10% sucrose solution through a drinking steel
spout and a pure tone of 6000 Hz at 80 dB from a set of two speakers placed 30 cm
in front of the platform. We conducted habituation to rewards and the auditory
stimulus separately. The number of reward presentations during the habituation
phase was not strictly defined; we determined that the mice were habituated by
confirming that they consumed the reward stably from the spout by visibly checking
the video. Auditory stimuli during the habituation phase were presented 120 times.
Mice were head-fixed on a tunnel-like, covered platform by clamping a surgically
implemented head plate on both sides (i.e., left and right from the anteroposterior
axis of the skull). The clamps were placed on a slide bar next to the platform and
adjusted to an appropriate height for each mouse. The platform floor was covered
with a copper mesh sheet, and a touch sensor was connected to the sheet and steel
spout.

After habituation, we conducted a Pavlovian delay conditioning task. Figure 1
(B and C) shows the experimental procedure. Mice were assigned to two
experimental groups, contingent (Figure 1B) and non-contingent (Figure 1C), with
eight mice in each group. In the contingent group, a pure tone of 6000 Hz at 80 dB
was randomly presented for 1 s as the conditioned stimulus (CS), followed
immediately by a 4µl drop of 10% sucrose solution (Figure 1B). The CS presentation
interval was random, ranging from 10 s to 20 s, and the mean value was set to 15 s.
In the non-contingent group, the CS and reward were independently presented
(Figure 1C). The CS and reward presentation intervals were random, ranging from
10 s to 20 s. We showed the percentages of CS and US overlapping trials for all
individuals and sessions in the non-contingent group (Supplementary Figure 1), and
there are no large differences between individuals. One session comprised 120
reward presentations for both groups. The training lasted for eight days. The CS and

reward presentation, response, and video recording were controlled using a custom-
made program written in Python 3 (3.7.8). The experiment was conducted in a
soundproof box with 75 dB of white noise in the laboratory to mask external sounds.
211

A. Head-fixed experimental setup

10% sucrose solution (Reward)

Personal computer

Solenoid

Speaker

Head-fixed mouse

Copper mesh sheet

Location of the infrared camera

45°

45 mm

View from the top

Arduino UNO

Touch sensor

B. Contingent group : reward is predictable (N = 8)

Tone ♪ x 120

Reward 💧 x 120

1s   1s   1s   1s

Durations of the inter-trial interval is randomized with average of 15s

C. Non-contingent group : reward is unpredictable (N = 8)

Durations of the inter-stimulus interval is randomized with average of 15s

Tone ♪ x 120

Reward 💧 x 120

1s   1s   1s   1s

Durations of the inter-reward interval is randomized with average of 15s

D. Quantification of pupil size with DeepLabCut

| Original image | Tracking with DeepLabCut | Ellipse fitting |

Figure 1. Schematic representation of head-fixed apparatus, Pavlovian delay
conditioning task, and pupillometry.

(A) Schematic representation of the head-fixed experimental apparatus and the
custom-made experimental control system. (B) Contingent group. In this group, the
1 s auditory stimulus (6000 Hz tone) was followed by a reward delivery of a 4 µL
drop of 10% of sucrose solution, and the auditory stimulus signaled the upcoming
reward. (C) Non-contingent group. In this group, the auditory stimulus and the reward
were presented independently and semi-randomly to prevent the development of the
reward-predictive value of the auditory stimulus. (D) The left panel shows an image
of a mouse's eye taken by the infrared camera in mice performing the Pavlovian
delay conditioning task. The center panel shows an image with eight tracked points
using DeepLabCut. The right panel shows an image of an ellipse fitted to the points
and an example of the temporal change in pupil size.

226　Drug

Pharmacological manipulations were conducted after training the Pavlovian
conditioning task to suppress the licking response in mice. Six blocks were
conducted for all individuals, each lasting three days. On Day 1, all mice were
intraperitoneally administered saline solution 15 min before the experiment
commenced. On Day 2, 15 minutes before the experiment commenced, haloperidol
(Serenace, Sumitomo Pharma) 0.1, 0.2, and 0.5 mg/kg was administered
intraperitoneally. This has been reported to inhibit licking (Fowler and Mortell, 1992;
Liao and Ko, 1995) and spontaneous movements (Arruda et al., 2008; Bernardi et
al., 1981; Conceição and Frussa-Filho, 1996; Strömbom, 1977) dose-dependently.
After Day 2, mice were allowed to drink water freely for one hour. On Day 3, mice
were not allowed access to water at all, and the experiment was not conducted to
avoid the residual effects of the drug. Therefore, we set approximately 48 hours to
wash out the effects of haloperidol. All individuals received each concentration of
haloperidol twice. The administration followed an ascending (0.1, 0.2, 0.5, 0.1, 0.2,

0.5) and descending (0.5, 0.2, 0.1, 0.5, 0.2, 0.1) order. Four mice experienced the

ascending order, and four experienced the descending order in each group.

Haloperidol was diluted in a saline solution. We administered haloperidol to mice via

intraperitoneal injection with a 10 mL/kg dose. After the injection, mice were returned

to their home cage until the start of the experiment.

246   Pupillometry

To measure the pupil size of mice performing the Pavlovian delay conditioning task,

we used an infrared camera (Iroiro1, Iroiro House) to capture a video of the mice's

heads during the task. The camera was placed at 45° from the midline of the mouse

(anteroposterior axis) and 45 mm from the top of the head (Figure 1A). The room's

brightness was set to 15 lux using a luminaire device (VE-2253, Etsumi). The pupil

size was extracted from videos. Figure 1D shows the flow of the pupil size analysis.

DeepLabCut, a deep-learning tracking software (Mathis et al., 2018; Nath et al.,

2019), was used to track the pupil edge at eight points. Using the "EllipseModel"

provided by scikit-image (Van der Wal et al., 2014), an ellipse was fitted to the eight

points obtained by tracking, and the estimated parameters (major and minor

diameters) were used to calculate the area of the ellipse. This area was used as

pupil size. After fitting ellipses to the tracked points and calculating pupil area, each

session's data for all subjects were independently transformed to the standard

normal distribution using the "scale" function in R. We employed "resnet50" as a

backbone network of a model and used default parameters. We annotated 8 points,

top, top-right, right, bottom-right, bottom, bottom-left, left, and top-left, for each frame.

The dataset contains 1650 frames (15 frames per video and 110 videos). We trained

the model with 1030000 iterations, and the train and test errors were 0.92 and 0.95

pixels, respectively.

266   Licking bout analysis

Animal responses occur as bouts, characterized by bursts of responses and pauses
that separate each bout (Gilbert, 1958; Shull et al., 2001). Conditioned responses
(CR) also occur as bouts (Harris, 2015; Kirkpatrick, 2002; Toda et al., 2017). Since
the CR has such a temporal pattern, individual licking can be classified into two
types: those that occur within bursts and during pauses. In previous studies, such a
bout-and-pause pattern was described by the mixture distribution of two exponential
distributions (Killeen et al., 2002): $P(IRT = \tau) = qe^{-b\tau} + (1-q)e^{-w\tau}$. In the
equation, q denotes the mixture ratio of the two types of responses, and w and b
denote the speed of the responses within bouts and the length of the pauses,
respectively. These parameters were free, and we fitted the equation to the empirical
data to estimate the parameters q, w, and b using a custom-made script and Turing
(Ge et al., 2018), a Bayesian inference software in Julia language. Under the
estimated parameters, individual licking was classified based on the likelihood of
whether it occurred within bursts or during pauses.

Statistical analysis

We collected data from all subjects repeatedly through our experiments and had
missing values caused by the failure of video recording, and we excluded those data
from the analysis. Given that our data was repeated-measurement data, including
missing values, assumptions employed in standard statistical analysis (i.e., that data
is independently and identically distributed) were violated. To account for repeated-
measurement data, we employed a linear-mixed model that can be used in the same
way as standard analysis of variance (ANOVA). Important aspects of the linear-
mixed model are fixed and random effects. Fixed effects are of primary interest to
researchers. In our experiments, group (contingent or non-contingent) and
pharmacological treatment (dose of haloperidol) are fixed effects. Fixed effects can
be interpreted analogously to effects examined in standard statistical analysis such
as ANOVA. Random effects comprise additional variability from other sources, such
as repeated measures clustered within subjects. By considering random effects, we

can account for the effects of repeated measurement and assess the fixed effect
more precisely (Singmann and Kellen, 2019). In statistical methodology, the maximal
random effect structure is usually recommended to be modeled directly to reduce
the Type I error when examining fixed effects (Barr et al., 2013). Thus, we included
factors measured repeatedly within subjects (i.e., time windows, sessions, and
pharmacological treatments) as random slopes. However, complex models have the
risk of failing to converge and of overfitting (Bates et al., 2015). In such cases,
several solutions have been proposed, and we have modified our implementation of
the model in cases where we were facing convergence or overfitting problems
(Brauer and Curtin, 2018). Therefore, we implemented a maximal random effect
structure at first. When the model failed to converge or was overfitted to the data, we
removed the random effect with the smallest variance. Finally, we employed the
model in which the model converged without overfitting problems. The specific model
we employed will be presented in the Results section. We fitted a linear mixed model
to our data using R (4.1) and the *lme4* package (Bates et al., 2014). We also used
the *emmeans* package (Lenth et al., 2018) to examine simple effects and employed
Tukey's method to adjust p-values for multiple comparisons.

Results

In the contingent group, the stimulus signaled reward delivery. Licking and pupil responses increased after the auditory stimulus presentation and the sucrose solution delivery ("Contingent"; Figure 2B and 2C). In the non-contingent group, the auditory stimulus did not signal reward delivery. Licking and pupil responses did not change after the auditory stimulus presentation but increased after the sucrose solution delivery ("Non-contingent"; in Figure 2B and 2C). We set three periods for analysis of licking and pupil size, 1 s before the presentation of the auditory stimulus (Pre-CS period; shown in green the shade in Figure 2B), 1 s during the presentation of the auditory stimulus (CS period; shown in the red shade in Figure 2B), and 1 s after the reward presentation (US period; shown in the blue shade in Figure 2B). We computed linear-mixed models to examine the effects of the respective procedure on the amount of licking and pupil size at each time window. In the respective model, we included the time window (levels: Pre, CS, and US), group (levels: contingent and non-contingent), and their interaction as fixed effects, as well as random intercepts and random slopes for the variable time window on subject level. Linear-mixed for the amount of licking revealed significant interaction between the time window and the procedure (Left panel of Figure 2D; $F(2, 13.999) = 105.8967$, p <.0001). Subsequently, multiple comparisons revealed significant differences in the amount of licking between each time window in the contingent group (Contingent in the left panel of Figure 2D; Pre vs. CS: $t(14) = -20.071$, p <.0001; Pre vs. US: $t(136)$ = -17.000, p <.0001; CS vs. US: $t(136) = -4.033$, p =.0033) and between Pre and US, US and CS in the non-contingent group (Non-contingent in the left panel of Figure 2D; Pre vs. US: $t(14) = -15.275$, p <.0001; CS vs. US: $t(14) = -13.521$, p <.0001), suggesting that the amount of licking increased after the auditory stimulus and reward presentation in the contingent group, but only after reward presentation in the non-contingent group. We also analyzed pupil size using an equivalent model, and the respective analysis revealed a significant main effect of the variable time

window (Right panel of Figure 2D; F (2, 14.023) = 5.4407, p =.0178). Subsequently,

we examined simple effects of the time window, which revealed significant

differences in pupil size between Pre and US, CS and US in the contingent group

(Right panel of Figure 2D; Pre vs. CS: t (13.8) = -2.563, p =.0557; Pre vs. US: t (14)

= -3.346, p =.0125; CS vs. US: t (13.9) = -3.061, p =.0217), suggesting that pupil

size increased after the reward presentation in the contingent group.

A. Schematic representation of analysis

B. Examples of licking and pupil responses

C. Average of licking and pupil responses

D. Licking and pupil responses in each period

Figure 2. Results of the Pavlovian conditioning training.

(A) Schematic representation of analyzed time windows. The presentation of CS or

US was set as 0, and the 3 s before and after the presentation were used for analysis.

(B). An example of licking responses and pupil response during each group's

Pavlovian delay conditioning task. Raster plot (top), temporal change in licking and

pupil size (middle and low) of a representative individual from each of the contingent

and non-contingent groups (N = 1 for each group, 120 trials each) 1 s before the

auditory stimulus presentation (Pre), during the auditory stimulus presentation (CS),

and immediately after the reward presentation (US) are shown in green, red, and

blue, respectively. (C) Mean temporal changes in licking frequency and pupil size

before and after CS and US presentations. Solid black lines indicate means; gray-

covered areas indicate the standard error of the mean (N = 8 for each group, 3

sessions each). Thin, colored lines indicate individual data. (D) Licking frequency

(left) and pupil size (right) at 1 s before, during, and immediately after CS

presentations (N = 8 for each group, 3 sessions each). Each time window

corresponds to the area covered by green, red, and blue in (A).

Figure 3. Temporal changes in licking responses and pupil size aligned with onsets
of bouts.
(A) Schematic representation of the response bout analysis. Initiation of the bout
was set as 0, and the 3 s before and after bout initiation was used for the analysis.
(B) Examples of raster plots before and after the start of the licking bout (top) and
temporal changes in licking responses and pupil size (middle and low) in contingent
and non-contingent groups ($N = 1$ for each group). (C) Average temporal changes
in licking and pupil size in contingent and non-contingent groups ($N = 8$ for each
group, 3 sessions each). (D) Mean amount of licking (left) and pupil size (right) at 3
s before and after the bout initiation ($N = 8$ for each group, 3 sessions each). In both
(B) and (C), data, including CS and US presentations within 3 s before and after the
initiation of the bout, were excluded. Individual data are shown as colored lines.

To examine the effect of licking responses on pupil size, we analyzed
temporal changes in licking and pupil responses around the onset of the licking bout
(Figure 3A). When we aligned the licking and pupil responses with the licking bout
onset, both groups' licking responses and pupil size increased with the bout onset
(Figure 3B–D). Licking responses recorded a phasic increase at the bout onset, and
the pupil size increased slightly after the bout onset. The pupil size slightly decreased
before the bout onset and increased after the bout onset. These results indicate that
pupil size increased after the initiation of licking responses. We examined the
amount of licking and pupil size before and after bout onsets using a linear mixed
model, which had the time window (Pre and Post) and group (contingent and non-
contingent) as a fixed effect and random intercepts and random slopes of time
window on subject level. The linear-mixed model revealed significant effects of the
time window and group on the amount of licking (Left panel of Figure 3D; Pre vs.
Post, $F(1, 13.481) = 38.358$, $p < .0001$; Contingent vs. Non-contingent, $F(1, 13.948)$
$= 5.5768$, $p = .0333$), suggesting that the amount of licking increased after bout onset.
The linear-mixed analysis also revealed a significant interaction of time window and
group (Right panel of Figure 3D; $F(1, 14.226) = 14.226$, $p = .0087$) regarding pupil

size. We examined the simple effects of time window on pupil size (Right panel of

Figure 3D; Pre vs. Post, t (13.7) = -6.223, p < .0001 in the contingent group),

suggesting that pupil size increased after bout onsets in the contingent group. We

provide the results of fitting the mixture distribution of two exponential distributions

in Supplementary Figure 2.

To further investigate whether the increase in pupil size resulted solely from

licking responses or reward prediction independently of licking responses, we

attempted to suppress licking responses in the same task. We thus intraperitoneally

injected haloperidol, a dopamine D2 receptor antagonist known to suppress licking

responses and locomotor activity (Fowler and Mortell, 1992; Liao and Ko, 1995;

Strömbom, 1977; Bernardi et al., 1981; Conceição and Frussa-Filho, 1996; Arruda

et al., 2008). After saline administration, licking responses and pupil size increased

after the auditory stimulus presentation in the contingent group (Contingent in CS-

aligned in Figure 4A and 4B) but remained unchanged in the non-contingent group

(Non-contingent in CS-aligned in Figure 4A and 4B). We observed an increase in

the licking frequency and pupil size at the reward delivery in both the contingent and

non-contingent groups (US-aligned in Figure 4A and 4C). Systemic administration

of haloperidol suppressed licking responses and pupil size in both the contingent

and non-contingent groups (Figure 4A–D). In particular, the increase in pupil size

after the reward delivery was suppressed in the contingent group (Contingent in US-

aligned in Figure 4A and the solid line in Figure 4C). We examined the effects of

group, time window, and dose of haloperidol on the amount of licking and pupil size

using linear-mixed modeling where the time window (Pre, CS, and US), group

(contingent and non-contingent), dose (saline, 0.0, 0.1, 0.2, and 0.5 mg/kg) and their

interactions were considered fixed effects, while subject-level random intercepts and

random sloped for the variables time window and dose were included. The linear-

mixed model revealed significant interactions between all variables (Left panel of

Figure 4D; F (6, 479.34) = 11.512, p < .0001), and we examined simple effects of

the time window. In the contingent group, the amount of licking differed between Pre

and CS, and Pre and US in all dose conditions (Upper left panel of Figure 4D; Saline:

Pre vs. CS, t (23.7) = -19.044, p < .0001; Pre vs. US, t (16.0) = -12.864, p < .0001;

0.1 mg/kg: Pre vs. CS, t (83.2) = -9.364, p < .0001; Pre vs. US, t (25.2) = -6.877, p

< .0001; 0.2 mg/kg Pre vs. CS, t (83.2) = -7.953, p < .0001; Pre vs. US, t (25.2) = -

5.127, p = .0001; 0.5 mg/kg: Pre vs. CS, t (83.2) = -3.324, p = .0037; Pre vs. US, t

(25.2) = -3.036, p = .0147), suggesting that the amount of licking increased after

auditory stimulus presentations. In the non-contingent group, the amount of licking

differed between Pre and US, and CS and US in all conditions (Bottom-left panel in

Figure 4D; Saline: Pre vs. US, t (16.0) = -11.311, p < .0001; CS vs. US, t (16.2) = -

11.729, p < .0001; 0.1 mg/kg: Pre vs. US, t (25.2) = -8.085, p < .0001; CS vs. US, t

(26.4) = -8.038, p < .0001; 0.2 mg/kg: Pre vs. US, t (26.3) = -5.833, p < .0001; CS

vs. US, t (27.7) = -5.603, p < 0.0001; 0.5 mg/kg: Pre vs. US, t (25.2) = -4.444, p

= .0004; CS vs. US, t (26.4) = -4.413, p = .0004), suggesting that the amount of

licking increased after reward presentations. We also analyzed pupil size with linear-

mixed analysis, revealing a significant interaction between all variables (Right panel

of Figure 4D; F (6, 479.52) = 3.009, p = .0068). We examined the simple effects of

the time window and found significant differences between all the time windows in

the contingent group (Upper right panel of Figure 4D; Saline: Pre vs. CS, t (20.7) =

-5.450, p = .0001; Pre vs. US, t (17.6) = -8.389, p < .0001; 0.1 mg/kg: Pre vs. CS, t

(59.0) = -3.452, p = .0029; Pre vs. US, t (36.0) = –7.569, p < .0001; 0.2 mg/kg Pre

vs. CS, t (59.0) = -3.000, p = .0109; Pre vs. US, t (36.0) = -7.817, p < .0001; 0.5

mg/kg: Pre vs. CS, t (59.0) = -4.344, p = .0002; Pre vs. US, t (36.0) = 9.585, p

< .0001), and Pre and CS in the non-contingent group (Bottom-right panel of Figure

4D; 0.1 mg/kg: Pre vs. CS, t (59.0) = -3.069, p = .0090; 0.2 mg/kg Pre vs. CS, t (63.8)

= -2.579, p = .0324; 0.5 mg/kg: Pre vs. CS, t (59.0) = -2.496, p = .0402). The increase

in licking responses and pupil size after the auditory stimulus presentation was

examined by calculating the difference between the mean values of licking

responses and pupil size for 3 s before and after the auditory stimulus presentation.

We performed a linear-mixed analysis to the change in the amount of licking and

pupil size. We assigned the group (contingent and non-contingent) and dose (saline,
0.1–0.5 mg/kg) to fixed effects and subject to random effect in licking analysis.
Linear-mixed analysis revealed a significant interaction between the group and the
dose. Subsequently, we examined the simple effect of dose and found that injection
of haloperidol decreased the amount of licking in a dose-dependent manner in the
contingent group (Upper left panel of Figure 4E; Saline vs. 0.1 mg/kg, t (169) = 7.741,
p < .0001; Saline vs. 0.2 mg/kg, t (169) = 10.704, p < .0001; Saline vs. 0.5 mg/kg, t
(169) = 16.205, p < .0001; 0.1 mg/kg vs. 0.5 mg/kg, t (169) = 6.910, p < .0001; 0.2
mg/kg vs. 0.5 mg/kg, t (169) = 4.492, p = .0001). We also analyzed pupil size using
a linear mixed model, where group (contingent and non-contingent), dose (saline,
0.1–0.5 mg/kg), and the change in the amount of licking were considered fixed
effects to assess pupil increase keeping out the effect of the increase of licks, and
subject to the random intercept. Linear-mixed analysis revealed a significant effect
of group, dose, and increase of licks (Upper right panel of Figure 4E; Group, F (1,
23.195) = 5.4185, p = .029; Dose, F (3, 173.005) = 3.1752, p = .0256; Licking, F (1,
184.886) = 4.7037, p = .0314), suggesting that pupil size showed stronger increases
in the contingent group than the non-contingent group, even after ruling out effects
of licks. We also examined the increase in licking responses and pupil size after
examining the reward presentation by calculating the difference between the mean
values of licking responses and pupil size for 3 s before and after the reward
presentation. We performed linear-mixed analysis considering changes in the
amount of licking and pupil size. We considered the variables group (contingent and
non-contingent) and dose (saline, 0.1–0.5 mg/kg) as fixed effects and included
subject-level random effects. Linear-mixed analysis revealed a significant interaction
between the group and the dose. Subsequently, we examined the simple effects of
dose and found that injection of haloperidol decreased the amount of licking in a
dose-dependent manner (Contingent in the bottom-left panel of Figure 4E; Saline vs.
0.2 mg/kg, t (169) = 3.878, p = .0009; Saline vs. 0.5 mg/kg, t (169) = 4.220, p = .0002;
Non-contingent in the bottom-left panel of Figure 4E; Saline vs. 0.1 mg/kg, t (169) =

3.052, p = .0139; Saline vs. 0.2 mg/kg, t (169) = 6.093, p < .0001; Saline vs. 0.5

mg/kg, t (169) = 8.476, p < .0001; 0.1 vs. 0.5 mg/kg, t (169) = 4.429, p = .0001). We

also analyzed pupil size using a linear mixed model, where group (contingent and

non-contingent), dose (saline, 0.1–0.5 mg/kg), and the change in the amount of

licking were considered fixed effects to assess pupil increase keeping out the effect

of the increase of licks, and subject to the random intercept. Linear-mixed analysis

revealed a significant effect of group (Bottom-right panel of Figure 4E; Group, F (1,

18.395) = 8.683, p = .0085), suggesting that pupil size significantly more strongly

increased in the contingent group than the non-contingent group.

Figure 4. Effects of haloperidol injection on the licking and pupil responses after

Pavlovian conditioning training.

(A) Representative raster plot (top), temporal change in licking, and pupil responses

(middle and low) of individuals in contingent and non-contingent groups. Periods of

1 s before the auditory stimulus presentation (Pre-CS), during the auditory stimulus

presentation (CS), and after the reward presentation (US) are shown in green, red,

and blue, respectively ($N = 1$ for each group, 120 trials each). (B) Mean temporal

changes in licking and pupil size before and after CS presentations ($N = 8$, 6

sessions for saline condition and 2 sessions for all haloperidol conditions). The upper

panel indicates licking responses. The horizontal axis indicates the time from the

reward onset. The vertical axis indicates frequencies of licking responses. The lower

panel indicates the data on pupil size. The horizontal axis indicates the time from the

reward onset. The vertical axis indicates the normalized pupil size. (C) Mean

temporal changes in licking and pupil responses before and after US presentations.

(D) Licking responses at Pre-CS, CS, and US periods ($N = 8$, 6 sessions for saline

condition and 2 sessions for all haloperidol conditions). The pupil size at Pre-CS, CS,

and US periods. (E) Difference between the mean values of licking responses and

the normalized pupil size during a 3 s before and after CS presentation (upper panel)

and reward presentation (bottom panel). HAL indicates haloperidol.

Discussion

This study explored the dynamics of the licking response and pupil size while mice performed a Pavlovian delay conditioning task to investigate the relationship between reward prediction and pupil size. The head-fixed experimental setup combined with deep-learning-based image analysis enabled us to reduce mice's spontaneous locomotor activity and to track the precise dynamics of licking responses and pupil size of the behaving mice. By manipulating the predictability of the reward in the Pavlovian delay conditioning task, we demonstrated that the pupil size of mice was modulated by reward prediction, consumption of the reward, and body movements associated with reward processing. Additionally, we found that the pupil size was modulated by reward prediction even after dose-dependent disruption of body movements by intraperitoneal injection of haloperidol, a dopamine D2 receptor antagonist.

In Experiment 1, we trained head-fixed mice on the Pavlovian delay conditioning task while recording licking and pupil responses. In this task, we designed contingent and non-contingent conditions to manipulate the predictability of the delivery of the sucrose solution by the auditory stimulus. In the contingent group, the auditory stimulus signaled the sucrose solution delivery. The mice showed increased licking responses and pupil size after the auditory stimulus presentation, suggesting that they could predict the outcome in this group. In the non-contingent group, the auditory stimulus did not signal the reward delivery. Licking responses and the pupil size of mice remained unchanged by the auditory stimulus presentation, suggesting that they did not associate the auditory stimulus with the reward in this group. In addition, the behavioral results obtained from the non-contingent group demonstrated that the sensory stimulus itself did not affect changes in licking responses and pupil size. The frequencies of the auditory stimulus presentation and reward delivery were identical between the contingent and non-contingent groups, with the only difference being the predictability of the outcome following the auditory

stimulus. This well-controlled rigid behavioral design allowed us to investigate the

modulation of behavioral states induced by reward prediction with the same sensory

signals.

Detailed bout analysis of licking responses revealed that pupil size increased

after the licking bout initiation in both the contingent and non-contingent groups,

suggesting that licking responses may modulate pupil size. Bout-aligned pupil size

also showed a clear decrease before the increase in pupil size. Before the bout

initiation, there was no licking response for approximately 0.5 s (Figure 3B and 3C).

This result also confirms the close relationship between pupil size and licking

responses. Many kinds of anticipatory behaviors occur when the stimulus signals a

future outcome. Thus, whether changes in pupil size reflect signals related to reward

prediction or are simply modulated by the motor-related signals accompanied by the

predictive movement remains unclear.

To examine whether the changes in pupil size reflect the modulations by the

prediction irrespective of motor-related signals, we examined the effects of

intraperitoneal injection of haloperidol, a dopamine D2 receptor antagonist, on the

dynamics of the pupil size of mice performing the Pavlovian delay conditioning task

in Experiment 2. Intraperitoneal injection of haloperidol suppressed licking

responses in a dose-dependent manner, supporting previous findings (Fowler and

Mortell, 1992; Liao and Ko, 1995). Although haloperidol administration decreased

pupil size, the effect was not as drastic as that of licking responses (Figure 4D and

4E). The highest dose of haloperidol injection almost completely disrupted licking

responses; however, we still observed pupil dilation after the auditory stimulus

presentation in the contingent group. This result implies that changes in pupil size

reflect reward-predictive signals irrespective of movement-related modulations.

In our experiments, the pupil size and licking responses were larger in the

non-contingent group, even if CSs were not presented compared to the contingent

group (Figure 4C and 4D). The timing of the reward presentations was completely

unpredictable in the non-contingent group, but the context predicted the possibility

of reward presentations. Pupil size was not increased after unpredictable reward
presentation in the non-contingent condition, suggesting that pupil size did not reflect
reward prediction errors in our experimental context. In our experiments, subjects
were trained extensively in a non-contingent context. This overtraining condition
might have caused the subjects to have no prediction errors at reward presentation,
even if the rewards were unpredicted in the non-contingent group. In further studies,
investigating the effect of reward prediction errors or uncertainty on pupil size by
presenting or omitting the reward will be important to understand the relationship
between pupil size changes and reward prediction errors.

In this study, we explored the dynamics of pupil size of mice performing the
Pavlovian delay conditioning task. We found that pupil dynamics reflected reward
prediction signals, irrespective of modulations by body movements. Pupil size is
modulated by autonomic nervous system activity. Sympathetic and parasympathetic
activation lead to pupil expansion and contraction, respectively. The sympathetic
control of the pupil is mediated by neuronal activity in the intermediolateral cell
column (IML) of the cervical and thoracic regions of the spinal cord. Cholinergic
neurons mediate the parasympathetic control in the Edinger-Westphal nucleus
(EWN). Most locus coeruleus (LC) neurons are noradrenergic, and their direct
projections to the IML stimulate sympathetic activation via noradrenergic α1
receptors. Direct projections to the EWN are thought to suppress the
parasympathetic nervous system by acting in an inhibitory manner via α2 receptors
(Joshi and Gold, 2020). Simultaneous measurements of LC neuronal activity and
pupil size in monkeys and rats have been reported to correlate (Joshi et al., 2016;
Liu et al., 2017). Therefore, pupil size measurement can be interpreted as an indirect
measure of LC activity.

Considering that LC neuronal activity is highly correlated with pupil size (Joshi
et al., 2016; Liu et al., 2017), our results that (1) the pupil was dilated to the auditory
stimulus that predicted the reward, and (2) the pupil size was unchanged when the
stimulus signaled no information about the reward, are consistent with existing

findings from electrophysiological experiments of LC neurons (Aston-Jones et al., 1997, 2005; Bouret and Richmond, 2015; Bouret and Sara, 2004). LC neurons show a burst of activity when the stimuli that predict biologically important events, such as reward and aversive events, are presented (Aston-Jones ad Bloom, 1981; Aston-Jones et al., 1997, 2005; Bouret and Richmond, 2015; Bouret and Sara, 2004). LC neurons also show similar activities to dopaminergic neurons in the ventral tegmental area (VTA), such as increased phasic activity in response to unpredicted reward and decreased activity through repeated experience and transfer to a reward-predicting stimulus (Amo et al., 2022; Bouret and Sara, 2004; Schultz et al., 1997). However, the neuronal activities of LC neurons in the study of Bouret and Sara (2003) were examined with the reversal of the contingency between the stimulus and the outcome or the re-acquisition after the extinction in the Go/No-Go task. Although the phasic activities to unpredicted reward found in LC neurons (Bouret and Sara, 2004) may be slightly different from those neurons found in dopaminergic neurons found in VTA (Amo et al., 2022; Schultz et al., 1997), both phasic activities of LC and DA neurons are known to show phasic responses to unpredictable events. Moreover, LC neurons show phasic activity in response to a novel stimulus and decreased activity when the stimulus ceases to predict biologically important events (Berridge and Waterhouse, 2003; Vankov et al., 1995). LC activity might be related to the salience-related dopaminergic activity found in the midbrain (Matsumoto and Hikosaka, 2009). As shown in Figures 2B and 4B, pupil size dilated for the reward-predictive stimulus but not for the reward-non-predictive stimulus. Although the reward prediction error did not modulate pupil size, the dynamics of pupil size observed in our experiments could be partially interpreted as reflecting LC activity.

In the canonical view of the reward prediction error hypothesis, neuronal activities of dopamine neurons in the VTA are modulated by the reward prediction errors and this signal is considered as teaching signals (Bayer and Glimcher, 2005; Eshel et al., 2016; Hollerman and Schultz, 1998; Satoh et al., 2003; Schultz et al., 1997). Learning also involves several other components, such as modulation of

motor outputs. Pupil size is considered to reflect internal states of organisms
involving arousal and/or attention and is modulated by noradrenergic neurons in the
LC. We found that haloperidol suppressed licking responses but not pupil size,
suggesting that dopamine D2 receptors are not involved in the modulation of the
reward prediction itself or attention/arousal modulated by the reward prediction. In
contrast, the fact that licking responses are suppressed by haloperidol suggests that
dopamine D2 receptors play a crucial role in the motor output based on the reward
prediction. Theoretically, if output signals of the reward prediction error are
modulated by the manipulation, the prediction itself as "associative strength"
assumed in the associative learning theory or "value" assumed in the reinforcement
learning theory would be also updated. In this sense, our results suggest that the
output of dopaminergic signals from the midbrain to D2 receptors in the brain areas
that receive dopaminergic projections might have an important role in modulating the
motor output irrespective of updating the reward-predictive signals. This conclusion
from our study supports recent findings that showed the neuronal activities of
dopamine neurons in the midbrain encode information about movement kinematics
(Barter et al., 2015; Hughes et al., 2020). However, we did not examine the effect of
the other dopamine receptor, dopamine D1 receptor. In the future, examining the
pharmacological manipulation of dopamine D1 receptors is an important step for
better circuit-level understanding of the neuronal mechanism of the reward
prediction and the reward prediction error.

Dopamine neurons in VTA show phasic activity to unexpected reward
presentations, but phasic activity to the reward decreases as learning progresses,
and the neurons show phasic activity to reward-predictive cues (Amo et al., 2022;
Schultz et al., 1997). Thus, in our experiments, increases in pupil size may reflect
reward prediction errors in the presentation of the auditory stimulus. However,
increases in pupil size for reward delivery were small in the non-contingent group
(Figure 2C, D and 4C, D). In addition, if the pupil size is modulated by the reward
prediction errors, pupil dilation should occur after the reward presentation only in the

 that the pupil size did not reflect reward prediction errors in our experiments. In addition, human studies indicated that the pupil size dilated for the reward-predictive cue in a delay conditioning task where the cue was presented 5 s before the reward (Pietrock et al., 2019). Taken together, increases in pupil size caused by the presentation of auditory stimuli in the contingent group could be interpreted as reward-predictive signals.

Considering the neurobiological mechanisms underlying the pupillary control system, the present findings of changes in pupil size being reflective of reward prediction signals invite us to reconsider the neuronal circuits computing reward prediction error signals. Cohen et al. (2012) reported that neuronal activities of GABAergic neurons in the rodent's VTA reflect the prediction of upcoming reward values. These activities are considered the source of the prediction for computing reward prediction errors encoded in dopamine neurons in the VTA. In the study of Cohen et al. (2012), they recorded neuronal activity while the mice performed a Pavlovian trace conditioning task, in which each odor cue was associated with different upcoming outcomes, e.g., small and large amounts of liquid rewards and air puffs. GABAergic neurons in the VTA showed persistent ramping activity during the delay between the presentation of cues and reward. However, CR, such as licking the reward spout, occurred during the delay between the cue and the reward delivery. In such cases, it is difficult to assess whether the neuronal activity reflects the reward value or behavioral expression, for example, the motor activity involved in licking responses modulated by the reward value. In the present study, we attempted to overcome this problem by suppressing body movements with haloperidol and found that the changes in pupil size reflected reward prediction signals independent of licking movements. The integrative approach of behavioral analysis, image analysis, pupillometry, and pharmacological manipulations

employed in the present study will pave the way for understanding the psychological
and neurobiological mechanisms involved in the computation of reward prediction
and reward prediction errors, which are essential features of learning and behavior.
We identified three limitations in this study: (1) the influence of body
movements other than licking responses, (2) the pharmacological selectivity to
haloperidol, and (3) the properties of CS and US, such as duration, magnitude, and
timing, being determinants of learned responses. In appetitive Pavlovian
conditioning, the presentation of the cue that predicts the outcome leads to the
observation of approach behavior to the cue or to the location where the reward is
presented (Boakes, 1977; Hearst and Jenkins, 1974). The locomotor activity also
occurs in mice under a head-fixed situation and has been reported to affect pupil
size (Cazettes et al., 2021). Intraperitoneal injection of haloperidol has been known
to dose-dependently decrease spontaneous activities, including locomotor activities.
Therefore, we hypothesized that the effect of locomotion on pupil size would be low
in our experiments. However, we found that licking responses were not suppressed
in all subjects. We cannot exclude this possibility because we could only measure
licking responses and no other motor expressions in our head-fixed setup. Taken
together, our experiments are limited mainly due to the potential effect of body
movement on pupil size. Second, we used haloperidol to suppress mice's body
movements, but haloperidol might affect pupil size due to its non-selective nature.
Haloperidol is a non-selective dopamine D2 antagonist that binds to D2-like
receptors, including D3 and D4 receptors, and others, such as adrenergic α1
receptors. Adrenergic α1 receptors are involved in pupil dilation, and haloperidol has
been reported to suppress pupil dilation produced by adrenaline administration in
mice (Korczyn and Keren, 1980). Furthermore, electrical stimulation of the LC
triggers the activity of dopamine cells in the midbrain via adrenergic α1 and
dopamine release in the nucleus accumbens (Grenhoff et al., 1993; Park et al., 2017).
Since pupil size is highly correlated with LC activity (Joshi et al., 2016; Liu et al.,
2017), the injection of haloperidol may affect the activity of dopamine neurons in the

midbrain and nucleus accumbens, which are modulated by LC activity. This
suggests that haloperidol may consequently affect reward prediction and the
calculation of reward prediction error. Furthermore, since it has been reported that
body movements suppress the activity of the auditory cortex in mice (Nelson et al.,
2013), it is possible that the injection of haloperidol suppresses body movement and
consequently modulates pupillary responsiveness to the CS. The results of
Experiment 2 may reflect this factor, where pupil dilation to CS was observed in the
non-contingent group after haloperidol injection.

In this study, we used haloperidol, a non-selective Dopamine D2 antagonist.
Although haloperidol does not increase pupil size, we might obtain cleaner results if
a more selective antagonist is used. In future investigations, the use of selective
dopamine D2 antagonists, such as eticlopride, and in combination with selective
dopamine D1 antagonists, such as SCH-23390, may more definitely prevent
modulations by the motor output and refine our understanding of the neurobiological
mechanisms underlying the relationship between the pupil size and reward
prediction.

Properties of CS and US, such as duration, intensity, and timing, affect
learned responses (Fanselow, 1994; Holland, 1977; Solomon and Corbit, 1974;
Timberlake, 1994). Taking such characteristics of conditioning, these factors may
have affected the result of our study. In our experiments, the duration of the auditory
stimulus was short, and the reward was followed by the auditory stimulus
immediately. In such a situation, mice would gain their arousal immediately after the
auditory stimulus presentation to consume the reward immediately. However, if the
auditory stimulus duration was long, the CS presentation was not followed by the
reward immediately, mice do not need to prepare to consume the reward
immediately after the CS presentation, and this kind of difference in the task structure
may lead to a different result. Although we did not use different types of US in our
experiments, pupil size may show that CR differs depending on CS and US
properties.

To verify that organisms predict future outcomes, behavioral evidence of
preparatory or anticipatory responses is necessary. In general, anticipatory
responses are accompanied by motor expressions; thus, dissociating whether the
physiological changes related to reward prediction encode the signal of the
prediction itself or are simply modulated by motor-related signals is difficult. Here,
we successfully measured changes in pupil size in mice performing the Pavlovian
delay conditioning task in the head-fixed situation using image processing. We
revealed that dynamic changes in pupil size reflect reward-predictive signals.
Pharmacological intervention experiments using haloperidol demonstrated that pupil
size increased even when licking responses were suppressed, supporting that the
changes in pupil size reflect reward prediction. Considering the brain circuits
involved in controlling pupil size, the predictive feature of pupil size suggests that
reward prediction is encoded in regions other than those reported by Cohen et al.
(2012) and Tian et al. (2016). These results pave the way for our understanding of
reward prediction signals in the brain by neutralizing the factor of motor expression
and suggest a different hypothesis for the neuronal circuits of predictive learning.
Future studies are expected to identify the neuronal circuit that computes the reward
prediction and reward prediction errors by eliminating the modulation of motor
expressions.

Data availability statement

762 Data supporting the findings of this study are available from the corresponding
763 author upon reasonable request. The original codes written for the analysis are
764 available from the corresponding author upon reasonable request.

Ethics statement

766 The experimental and housing protocols adhered to the Japanese National
767 Regulations for Animal Welfare and were approved by the Animal Care and Use
768 Committee of Keio University.

Author contributions

YK and KT designed the experiments. YK conducted stereotaxic surgery and pharmacological manipulations for mice and collected all the data from the head-fixed Pavlovian conditioning experiment with the help of KT. YK and KT analyzed the data and created all figures. YK and KT wrote the manuscript.

Funding

This research was supported by JSPS KAKENHI 18KK0070(KT), 19H05316 (KT), 19K03385 (KT), 19H01769 (KT), 20J21568 (KY), 22H01105 (KT), Keio Academic Development Fund (KT), and Keio Gijuku Fukuzawa Memorial Fund for the Advancement of Education and Research (KT).

Conflict of interest

The authors declare that they have no conflict of interest.

Acknowledgments

We thank Kohei Yamamoto, Saya Yatagai, Yusuke Ujihara, Daiki Nasukawa, Yasuyuki Niki, Haruka Hirakata, Shohei Kaneko, Ryuto Tamura, Lingcheng Kong, and Kazuko Hayashi for their assistance with animal care and Youcef Bouchekioua, Akihiro Funamizu, and Takaaki Ozawa for valuable discussions. We also thank the anonymous reviewers for their constructive comments on the manuscript.

References

Amo, R., Matias, S., Yamanaka, A., Tanaka, K. F., Uchida, N., & Watabe-Uchida, M. (2022). A gradual temporal shift of dopamine responses mirrors the progression of temporal difference error in machine learning. *Nature Neuroscience, 25*(8), 1082-1092.

Arruda, M. D. O. V., Soares, P. M., Honório, J. E. R., Lima, R. C. D. S., Chaves, E. M. C., Lobato, R. D. F. G.,... & Vasconcelos, S. M. M. (2008). Activities of the antipsychotic drugs haloperidol and risperidone on behavioural effects induced by ketamine in mice. *Scientia Pharmaceutica*, *76*(4), 673-688.

Aston-Jones, G., Rajkowski, J., & Kubiak, P. (1997). Conditioned responses of monkey locus coeruleus neurons anticipate acquisition of discriminative behavior in a vigilance task. *Neuroscience, 80*(3), 697-715.

Aston-Jones, G., Rajkowski, J., Kubiak, P., & Alexinsky, T. (1994). Locus coeruleus neurons in monkey are selectively activated by attended cues in a vigilance task. *Journal of Neuroscience, 14*(7), 4467-4480.

Barr, D. J., Levy, R., Scheepers, C., & Tily, H. J. (2013). Random effects structure for confirmatory hypothesis testing: Keep it maximal. *Journal of memory and Language*, *68*(3), 255-278.

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2014). Fitting linear mixed-effects models using lme4. *arXiv preprint arXiv:1406.5823*.

Bates, D., Kliegl, R., Vasishth, S., & Baayen, H. (2015). Parsimonious mixed models. arXiv preprint arXiv:1506.04967.

Brauer, M., & Curtin, J. J. (2018). Linear mixed-effects models and the analysis of
nonindependent data: A unified framework to analyze categorical and
continuous independent variables that vary within-subjects and/or within-items.
Psychological Methods, *23*(3), 389.

Barter, J. W., Li, S., Lu, D., Bartholomew, R. A., Rossi, M. A., Shoemaker, C. T.,
Salas-Meza, D., Gaidis, E., & Yin, H. H. (2015). Beyond reward prediction
errors: the role of dopamine in movement kinematics. *Frontiers in Integrative*
Neuroscience, 9, 39.

Bayer, H. M., & Glimcher, P. W. (2005). Midbrain dopamine neurons encode a
quantitative reward prediction error signal. *Neuron*, *47*(1), 129-141.

Bernardi, M. M., De Souza, H., & Neto, J. P. (1981). Effects of single and long-term
haloperidol administration on open field behavior of rats. *Psychopharmacology*,
73(2), 171-175.

Berridge, C. W., & Waterhouse, B. D. (2003). The locus coeruleus–noradrenergic
system: modulation of behavioral state and state-dependent cognitive
processes. *Brain Research Reviews, 42*(1), 33-84.

Boakes, R. A. (1977). "Performance on learning to associate a stimulus with positive
reinforcement," in *Operant-Pavlovian Interactions*, eds H. Davis and H. M. B.
Hurwitz (Hillsdale, NJ: Lawrence Erlbaum Associates), 67–97.

Bouret, S., & Richmond, B. J. (2015). Sensitivity of locus ceruleus neurons to reward
value for goal-directed actions. *Journal of Neuroscience, 35*(9), 4005-4014.

Bouret, S., & Sara, S. J. (2004). Reward expectation, orientation of attention and
locus coeruleus-medial frontal cortex interplay during learning. *European*
Journal of Neuroscience, 20(3), 791-802.

Cazettes, F., Reato, D., Morais, J. P., Renart, A., & Mainen, Z. F. (2021). Phasic
activation of dorsal raphe serotonergic neurons increases pupil size. *Current*
Biology, *31*(1), 192-197.

Cohen, J. Y., Haesler, S., Vong, L., Lowell, B. B., & Uchida, N. (2012). Neuron-type-
specific signals for reward and punishment in the ventral tegmental area.
Nature, *482*(7383), 85-88.

Conceição, I. M., & Frussa-Filho, R. (1996). Effects of microgram doses of
haloperidol on open-field behavior in mice. *Pharmacology Biochemistry and*
Behavior, *53*(4), 833-838.

Ebitz, R. B. & Platt, M. L. (2015). Neuronal activity in primate dorsal anterior cingulate
cortex signals task conflict and predicts adjustments in pupil-linked arousal.
Neuron, 85(3), 628-40.

Ebitz, R. B. & Moore, T. (2019). Both a gauge and a filter: Cognitive modulations of
pupil size. *Frontiers in Neurology*, *9*, 1190.

Ebitz, R. B., Pearson, J. M., & Platt, M. L. (2014). Pupil size and social vigilance in
rhesus macaques. *Frontiers in Neuroscience*, *8*, 100.

Eshel, N., Tian, J., Bukwich, M., & Uchida, N. (2016). Dopamine neurons share
common response function for reward prediction error. *Nature Neuroscience*,
19(3), 479-486.

Estes, W. K., & Skinner, B. F. (1941). Some quantitative properties of anxiety.
Journal of Experimental Psychology, *29*(5), 390.

Esteves, F., Parra, C., Dimberg, U., & Öhman, A. (1994). Nonconscious associative
learning: Pavlovian conditioning of skin conductance responses to masked
fear-relevant facial stimuli. *Psychophysiology*, *31*(4), 375-385.

Fanselow, M. S. (1994). Neural organization of the defensive behavior system
responsible for fear. *Psychonomic Bulletin & Review*, *1*(4), 429-438.

Finke, J. B., Roesmann, K., Stalder, T., & Klucken, T. (2021). Pupil dilation as an
index of Pavlovian conditioning. A systematic review and meta-analysis.
Neuroscience & Biobehavioral Reviews, *130*, 351-368.

Fowler, S. C., & Mortell, C. (1992). Low doses of haloperidol interfere with rat tongue
extensions during licking: a quantitative analysis. *Behavioral Neuroscience*,
106(2), 386.

Ge, H., Xu, K., & Ghahramani, Z. (2018, March). Turing: a language for flexible
probabilistic inference. Proceedings of the Twenty-First International
Conference on Artificial Intelligence and Statistics, in *Proceedings of Machine*
Learning Research, 84,1682-1690.

Gilbert, T. F. (1958). Fundamental dimensional properties of the operant.
Psychological Review, *65*(5), 272.

Grenhoff, J., Nisell, M., Ferre, S., Aston-Jones, G., & Svensson, T. H. (1993).
Noradrenergic modulation of midbrain dopamine cell firing elicited by
stimulation of the locus coeruleus in the rat. *Journal of Neural*
Transmission/General Section, 93(1), 11-25.

Harris, J. A. (2015). Changes in the distribution of response rates across the CS-US
interval: Evidence that responding switches between two distinct states.

Journal of Experimental Psychology: Animal Learning and Cognition, *41*(3),
217.

Hearst, E., & Jenkins, H. M. (1974). *Sign-tracking: The stimulus-reinforcer relation*
and directed action. Psychonomic Society.

Holland, P. C. (1977). Conditioned stimulus as a determinant of the form of the
Pavlovian conditioned response. *Journal of Experimental Psychology: Animal*
Behavior Processes, *3*(1), 77.

Hollerman, J. R., & Schultz, W. (1998). Dopamine neurons report an error in the
temporal prediction of reward during learning. *Nature Neuroscience*, *1*(4), 304-
309.

Hughes, R. N., Bakhurin, K. I., Petter, E. A., Watson, G. D. R., Kim, N., Friedman, A.
D., & Yin, H. H. Ventral Tegmental Dopamine Neurons Control the Impulse
Vector during Motivated Behavior. *Current Biology*, 30(14), 2681-2694.

Joshi, S., & Gold, J. I. (2020). Pupil size as a window on neural substrates of
cognition. *Trends in Cognitive Sciences*, *24*(6), 466-480.

Joshi, S., Li, Y., Kalwani, R. M., & Gold, J. I. (2016). Relationships between pupil
diameter and neuronal activity in the locus coeruleus, colliculi, and cingulate
cortex. *Neuron*, *89*(1), 221-234.

Kaneko, S., Niki, Y., Yamada, K., Nasukawa, D., Ujihara, Y., & Toda, K. (2022).
Systemic injection of nicotinic acetylcholine receptor antagonist mecamylamine
affects licking, eyelid size, and locomotor and autonomic activities but not
temporal prediction in male mice. *Molecular Brain, 17*, 77.

Killeen, P. R., Hall, S. S., Reilly, M. P., & Kettle, L. C. (2002). Molecular analyses of
the principal components of response strength. *Journal of the Experimental*
Analysis of Behavior, *78*(2), 127-160.

Kirkpatrick, K. (2002). Packet theory of conditioning and timing. *Behavioural*
Processes, *57*(2-3), 89-106.

Koenig, S., Uengoer, M., & Lachnit, H. (2018). Pupil dilation indicates the coding of
past prediction errors: Evidence for attentional learning theory.
Psychophysiology, *55*(4), e13020.

Konorski, J. (1967). *Integrative activity of the brain*. Chicago: University of Chicago
Press.

Korczyn, A. D., & Keren, O. (1980). The effect of dopamine on the pupillary diameter
in mice. *Life Sciences*, *26*(10), 757-763.

Larsen, R. S., & Waters, J. (2018). Neuromodulatory correlates of pupil dilation.
Frontiers in Neural Circuits, *12*, 21.

Lee, C. R., & Margolis, D. J. (2016). Pupil dynamics reflect behavioral choice and
learning in a go/nogo tactile decision-making task in mice. *Frontiers in*
Behavioral Neuroscience, *10*, 200.

Lenth, R., Singmann, H., Love, J., Buerkner, P., & Herve, M. (2018). Emmeans:
Estimated marginal means, aka least-squares means. *R package version*, *1*(1),
3.

Leuchs, L., Schneider, M., Czisch, M., & Spoormaker, V. I. (2017). Neural correlates
of pupil dilation during human fear learning. *Neuroimage*, *147*, 186-197.

Liao, R. M., & Ko, M. C. (1995). Chronic effects of haloperidol and SCH23390 on
operant and licking behaviors in the rat. *Chinese Journal of Physiology*, *38*, 65-
74.

Liu, Y., Rodenkirch, C., Moskowitz, N., Schriver, B., & Wang, Q. (2017). Dynamic
lateralization of pupil dilation evoked by locus coeruleus activation results from
sympathetic, not parasympathetic, contributions. *Cell Reports*, *20*(13), 3099-
3112.

Lonsdorf, T. B., Menz, M. M., Andreatta, M., Fullana, M. A., Golkar, A., Haaker, J.,...
& Merz, C. J. (2017). Don't fear 'fear conditioning': Methodological
considerations for the design and analysis of studies on human fear acquisition,
extinction, and return of fear. *Neuroscience & Biobehavioral Reviews*, *77*, 247-
285.

Mackintosh, N. J. (1975). A theory of attention: Variations in the associability of
stimuli with reinforcement. *Psychological Review*, *82*(4), 276.

Mathis, A., Mamidanna, P., Cury, K. M., Abe, T., Murthy, V. N., Mathis, M. W., &
Bethge, M. (2018). DeepLabCut: markerless pose estimation of user-defined
body parts with deep learning. *Nature Neuroscience*, *21*(9), 1281-1289.

Matsumoto, M., & Hikosaka, O. (2009). Two types of dopamine neuron distinctly
convey positive and negative motivational signals. *Nature*, *459*(7248), 837-841.

Nath, T., Mathis, A., Chen, A. C., Patel, A., Bethge, M., & Mathis, M. W. (2019).
Using DeepLabCut for 3D markerless pose estimation across species and
behaviors. *Nature Protocols*, *14*(7), 2152-2176.

Nelson, A., & Mooney, R. (2016). The basal forebrain and motor cortex provide
convergent yet distinct movement-related inputs to the auditory cortex. *Neuron*,
90(3), 635-648.

Nelson, A., Schneider, D. M., Takatoh, J., Sakurai, K., Wang, F., & Mooney, R.
(2013). A circuit for motor cortical modulation of auditory cortical activity.
Journal of Neuroscience, 33(36), 14342-14353.

Notterman, J. M., Schoenfeld, W. N., & Bersh, P. J. (1952). Conditioned heart rate
response in human beings during experimental anxiety. *Journal of*
Comparative and Physiological Psychology, 45(1), 1–8.

Öhman, A., Fredrikson, M., Hugdahl, K., & Rimmö, P.-A. (1976). The premise of
equipotentiality in human classical conditioning: Conditioned electrodermal
responses to potentially phobic stimuli. *Journal of Experimental Psychology:*
General, 105(4), 313–337.

Ojala, K. E., & Bach, D. R. (2020). Measuring learning in human classical threat
conditioning: Translational, cognitive and methodological considerations.
Neuroscience & Biobehavioral Reviews, *114*, 96-112.

Park, J. W., Bhimani, R. V., & Park, J. (2017). Noradrenergic modulation of
dopamine transmission evoked by electrical stimulation of the locus coeruleus
in the rat brain. *ACS Chemical Neuroscience, 8*(9), 1913-1924.

Pavlov, I. P. (1927). Conditioned reflexes: an investigation of the physiological
activity of the cerebral cortex. Oxford Univ. Press.

Pearce, J. M., & Hall, G. (1980). A model for Pavlovian learning: variations in the
effectiveness of conditioned but not of unconditioned stimuli. *Psychological*
Review, *87*(6), 532.

Pietrock, C., Ebrahimi, C., Katthagen, T. M., Koch, S. P., Heinz, A., Rothkirch, M., &
Schlagenhauf, F. (2019). Pupil dilation as an implicit measure of appetitive
Pavlovian learning. *Psychophysiology*, *56*(12), e13463.

Privitera, M., Ferrari, K. D., von Ziegler, L. M., Sturman, O., Duss, S. N., Floriou-
Servou, A.,... & Bohacek, J. (2020). A complete pupillometry toolbox for real-
time monitoring of locus coeruleus activity in rodents. *Nature Protocols*, *15*(8),
2301-2320.

Reimer, J., Froudarakis, E., Cadwell, C. R., Yatsenko, D., Denfield, G. H., & Tolias,
A. S. (2014). Pupil fluctuations track fast switching of cortical states during quiet
wakefulness. *Neuron*, *84*(2), 355-362.

Rescorla, R A., & Wagner, A. R. (1972). A theory of Pavlovian conditioning:
Variations in the effectiveness of reinforcement and nonreinforcement. In A. H.
Black & W. F. Proktsy (Eds.), *Classical conditioning II: Current research and*
theory, 64-99. Appleton-Century-Crofts.

Satoh, T., Nakai, S., Sato, T., & Kimura, M. (2003). Correlated coding of motivation
and outcome of decision by dopamine neurons. *Journal of Neuroscience*,
23(30), 9913-9923.

Schultz, W., Dayan, P., & Montague, P. R. (1997). A neural substrate of prediction
and reward. *Science*, *275*(5306), 1593-1599.

Shull, R. L., Gaynor, S. T., & Grimes, J. A. (2001). Response rate viewed as
engagement bouts: Effects of relative reinforcement and schedule type.
Journal of the Experimental Analysis of Behavior, *75*(3), 247-274.

Singmann, H., & Kellen, D. (2019). An introduction to mixed models for experimental
psychology. In *New Methods in Cognitive Psychology*, 4-31. Routledge.

992 Solomon, R. L., & Corbit, J. D. (1974). An opponent-process theory of motivation: I.
993   Temporal dynamics of affect. *Psychological Review*, *81*(2), 119.

994 Strömbom, U. (1977). Antagonism by haloperidol of locomotor depression induced
995   by small doses of apomorphine. *Journal Neural Transmission 40*, 191–194.

996 Sutton, R. S., & Barto, A. G. (2018). *Reinforcement learning: An introduction*. MIT
997   press.

998 Tian, J., Huang, R., Cohen, J. Y., Osakada, F., Kobak, D., Machens, C. K.,... &
999   Watabe-Uchida, M. (2016). Distributed and mixed information in monosynaptic
1000   inputs to dopamine neurons. *Neuron*, *91*(6), 1374-1389.

1001 Timberlake, W. (1994). Behavior systems, associationism, and Pavlovian
1002   conditioning. *Psychonomic Bulletin & Review*, *1*(4), 405-420.

1003 Toda, K., Lusk, N. A., Watson, G. D., Kim, N., Lu, D., Li, H. E.,... & Yin, H. H. (2017).
1004   Nigrotectal stimulation stops interval timing in mice. *Current Biology*, *27*(24),
1005   3763-3770.

1006 Vankov, A., Hervé-Minvielle, A., & Sara, S. J. (1995). Response to novelty and its
1007   rapid habituation in locus coeruleus neurons of the freely exploring rat.
1008   *European Journal of Neuroscience, 7*(6), 1180-1187.

1009 Van der Walt, S., Schönberger, J. L., Nunez-Iglesias, J., Boulogne, F., Warner, J. D.,
1010   Yager, N.,... & Yu, T. (2014). scikit-image: image processing in Python. *PeerJ,*
1011   *2*, e453.

1012 Van Slooten, J. C., Jahfari, S., Knapen, T., & Theeuwes, J. (2018). How pupil
1013   responses track value-based decision-making during and after reinforcement
1014   learning. *PLoS Computational Biology*, *14*(11), e1006632.

Vincent, P., Parr, T., Benrimoh, D., & Friston, K. J. (2019). With an eye on
uncertainty: Modelling pupillary responses to environmental volatility. *PLoS*
Computational Biology, *15*(7), e1007126.

Wang, H., Ortega, H. K., Atilgan, H., Murphy, C. E., & Kwan, A. C. (2022). Pupil
correlates of decision variables in mice playing a competitive mixed-strategy
game. *eNeuro*, *9*(2), 0457-21.

Wood, D. M., & Obrist, P. A. (1964). Effects of controlled and uncontrolled respiration
on the conditioned heart rate response in humans. *Journal of Experimental*
Psychology, 68(3), 221–229.

Yamamoto, K., Yamada, K., Yatagai, S., Ujihara, Y., and Toda, K. (2022).
Spatiotemporal Pavlovian head-fixed reversal learning task for mice. *Molecular*
Brain, 15, 78.

Zénon, A. (2019). Eye pupil signals information gain. *Proceedings of the Royal*
Society B, 286(1911), 20191593.

1029

Supplementary Materials

Supplementary Figure 1. Empirical contingency in the non-contingent group.

We calculated the percentage of CS and US overlapping trials for all individuals and sessions in the non-contingent group. The definition of overlap was determined by whether reward presentations were presented during auditory stimulus presentations.

The log survivor plot is a method to visualize the bout-and-pause patterns, and when
responses have bout-and-pause patterns, the plot shows the broken-stick curve.
The left side line denotes the within-bout inter-licking intervals, and the right-side line
denotes the bout initiation intervals. The intercept of the right-side line denotes the
bout length, the amount of licking contained in one bout. Log survivor plots of
empirical and simulated data showed broken-stick curves, suggesting licks have
bout-and-pause patterns. As shown in the figure, the bend points were
approximately 0.1–1.0s, suggesting that the boundaries separating the within-bout
licking from the bout initiation licking were in the range and corresponded to Figure
3B and C.

We found that the log survivor plot in the contingent group showed a clear
bend point in training and saline conditions; in contrast, the plot did not show a clear
bend point and showed a gradual curve in the non-contingent group. As the dose of
haloperidol increased, the bend point became clearer. In the contingent group, the
slope of the right lines became gradual as the dose of haloperidol increased. Taken
together, mice did not show spontaneous licking during the inter-reward intervals.

1054

Supplementary Figure 2. Fitting results of the mixture exponential distribution.

(A) Solid lines show log survivor plots of empirical inter-licking intervals, and dashed

lines show log survivor plots of simulated data using fitted parameters. Model fitting

was performed independently for individual and session data, and we generated

random numbers from the distribution of the estimated parameters. Each line

denotes the average over subjects and sessions, and gray shades denote standard

errors. (B) Average value and range of estimated parameters, *w*, *b*, and *p,* in each

group and dose condition. T denotes the data in the last three sessions of training.

1063

We compared the dynamics of licking and pupil size before/after the auditory
stimulus presentation between the first and last training sessions. However, we failed
to record several data in the first session, showing only 4 and 3 subjects for
contingent and non-contingent groups. In the first session, the amount of licking
slightly increased after the auditory stimulus in the contingent group but not in the
non-contingent group. In the last session, the increase in the amount of licking
became larger in the contingent group. In the non-contingent group, the amount of
licking did not increase after the auditory stimulus presentation, but the baseline was
larger in the last session compared to the first session. In both groups, mice showed
pupil dilation to the auditory stimulus in the first session, but in the last session, it
decreased in the non-contingent group but not in the contingent group.

The amount of licking was acquired by Pavlovian conditioning, but the pupil
size showed an increase in the very first session. The pupil size is highly correlated
with LC activity, and LC shows the phasic activity to a novel stimulus. It also shows
the activity when the environmental rule, such as stimulus-reward contingency,
changes. The increase in pupil size in the first session may reflect the novelty of
stimulus or change in the environmental rule, such as the transition from habituation
to contingent or non-contingent procedure. At least, pupil size does not reflect only
reward prediction but also novelty, uncertainty, and environmental changes, so that
a well-known learning curve may not be drawn.

1084

Supplementary Figure 3. Comparison of dynamics of licking and pupil size

between early and last session in training.

We analyzed the relationship between the amount of licking and pupil size in each

trial. Although licking increased pupil size, as shown in Figure 3, we could not find

any relationship between the amount of licking and pupil size. The large temporal

variance in pupil size may mask the relationship between licking and pupil size in

this time-scale analysis.

1092

Supplementary Figure 4. Scatter plots of the amount of licking and pupil size.

The upper panels show relationships between the amount of licking and pupil size

at 3 s before the auditory stimulus presentation. The bottom panels show those of 3

s after the auditory stimulus presentation. Blue and red points denote the non-

contingent and contingent groups, respectively. We analyzed the saline condition

separately for previous haloperidol dose conditions to examine whether the effects

of haloperidol were washed-out. We found no difference between the previous dose

in the amount of licking and pupil size at any time window, suggesting that the effects

of haloperidol had been washed-out until the saline condition.

1102

1103 Supplementary Figure 5. The amount of licking and pupil size at the time windows.

1104 The amount of licking and pupil size in Pre-CS, CS, and US periods of saline

1105 condition were separately shown by the previous dose of haloperidol injection.

1106