Molecular Acclimation of *Halobacterium salinarum* to Halite Brine Inclusions

C. Favreau[1], A. Tribondeau[2], M. Marugan[1], F. Guyot[3], B. Alpha-Bazin[4], A. Marie[1], R. Puppo[1],

T. Dufour[5], A. Huguet[6], S. Zirah[1], A. Kish[1*]

[1]Unité Molécules de Communication et Adaptation des Microorganismes (MCAM), Muséum National
d'Histoire Naturelle (MNHN), CNRS, 57 rue Cuvier, 75005 Paris; France

[2]Unité Physiologie Moléculaire et Adaptation (PhyMA), MNHN, CNRS, 57 rue Cuvier, 75005 Paris;
France

[3]Institut de Minéralogie, de Physique des Matériaux et de Cosmochimie (IMPMC), MNHN, Sorbonne
Université, CNRS, IRD, 4 place Jussieu, 75005 Paris; France

[4] Département Médicaments et Technologies pour la Santé (DMTS), Université Paris-Saclay, CEA,
INRAE, SPI, 30200 Bagnols-sur-Cèze; France

[5]Laboratoire de Physique des Plasma (LPP), Sorbonne Université, CNRS, Ecole Polytechnique,
Université Paris-Sud, Observatoire de Paris, 4 Place Jussieu, 75252 Paris; France

[6]Unité Milieux Environnementaux Transferts et Interactions dans les hydosystèmes et les Sols
(METIS), Sorbonne Université, CNRS, EPHE, PSL 4 place Jussieu, 75005 Paris; France

* Correspondence:
Adrienne Kish, Unité MCAM, Muséum National d'Histoire Naturelle (MNHN), CNRS, 57 rue
Cuvier, 75005 Paris; France
adrienne.kish@mnhn.fr

Keywords: *Halobacterium*, halophile, halite, proteomics

1   Abstract

Halophilic microorganisms have long been known to survive within the brine inclusions of salt crystals,

as evidenced by their pigmentation. However, the molecular mechanisms allowing this survival has

remained an open question for decades. While protocols for the surface sterilization of halite (NaCl)

have enabled isolation of cells and DNA from within halite brine inclusions, "-omics" based

approaches have faced two main technical challenges: (1) removal of all contaminating organic

biomolecules (including proteins) from halite surfaces, and (2) performing selective biomolecule

extractions directly from cells contained within halite brine inclusions with sufficient speed to avoid

modifications in gene expression during extraction. In this study, we present methods to resolve these

two technical challenges. In addition, we apply these methods to perform the first examination of the

early acclimation of a model haloarchaeon (*Halobacterium salinarum* NRC-1) to halite brine

inclusions. Examinations of the proteome of *Halobacterium* cells two months post-evaporation
revealed a high degree of similarity with stationary phase liquid cultures, but with a sharp down-
regulation of ribosomal proteins. Low quantities of RNA from halite brine inclusions corroborate the
hypothesis of low transcriptional and translational activities. While proteins for central metabolism
were part of the shared proteome between liquid cultures and halite brine inclusions, proteins involved
in cell mobility (archaellum, gas vesicles) were either absent or less abundant in halite samples.
Proteins unique to cells within brine inclusions included transporters, suggesting modified interactions
between cells and the surrounding brine inclusion microenvironment. The methods and hypotheses
presented here enable future studies of the survival of halophiles in both culture model and natural
halite systems.

2   Introduction

Extremely halophilic archaea are adapted to the highest possible salinity conditions, thriving in
saturated brines. Microbial diversity in such environments is highly reduced and tends to be dominated
by haloarchaea (Oren, 2006). Saturated brines are characterized by low dissolved oxygen content
despite contact with the air and are often exposed to elevated temperatures and high levels of solar
radiation (Oren, 2006; Merino *et al*., 2019) leading to evaporation. During evaporation events, the total
contents of the brine, including salts, organics, some atmospheric gases, and any microorganisms
present, are trapped within inclusions in the salt crystals. Viable halophilic bacteria and archaea have
been isolated from halite (NaCl) of various ages, from months to years (Norton and Grant, 1988;
Gramain *et al*., 2011; Huby *et al*., 2020), to geologically relevant timescales. Some studies have even
reported the observation or isolation of microorganisms from halite dated to several hundreds of
millions of years (Vreeland *et al*., 2000; Schreder-Gomes *et al*., 2022). These observations have raised
two important questions: (1) are these truly 'ancient' cells? (2) what physiological changes are required
for microbial cells to remain viable within halite brine inclusions?

The first question has been studied in greater detail, and while many questions still remain,
these studies provided valuable insights into the potential for surface-contaminant microorganisms
(Graur and Pupko, 2001; Maughan *et al*., 2002; Nickle *et al*., 2002). In response to these criticisms,
appropriate surface sterilization and cleaning procedures for microbial cultivation and DNA isolation
for micro-biodiversity studies from dissolved ancient halite (Gramain *et al*., 2011; Sankaranarayanan
et al., 2011) were developed. These approaches enabled more accurate identification of only those
microorganisms trapped within halite inclusions. Another opened question is whether the age

determined from the mineral halite can be applied directly to the microorganisms in the brine
inclusions. Some studies have examined the possibility that microbial cell-like biomorphs are
preserved within brine inclusions (Nims *et al*., 2021).

Experimentally confirming how microbial life is supported within brine inclusions has proven
more difficult. Determining the cellular functions expressed by viable halophiles within halite brine
inclusions requires the direct isolation of biomolecules such as proteins and nucleic acids for "-omics"
analyses, including proteomics and transcriptomics. However, significant technical challenges are
presented by the closed system of brine inclusions within halite. Accessing the contents of brine
inclusions by rapid crystal dissolution leads to cell lysis by osmotic shock and biomolecule
degradation, while the time required for gradual crystal dissolution preserving cellular integrity is also
sufficient for alterations in the transcriptomes and proteomes of viable cells away from their state
within the brine inclusions. In addition, the direct extraction of biomolecules from brine inclusions
involves a large amount of NaCl, which is incompatible with mass spectrometry due to the lack of
sufficient desalination steps. Although current hypotheses concerning the modifications of halophile
physiology during entrapment within halite brine inclusions include a change to anaerobic metabolism
(Winters *et al.,* 2015) along with potential for cell envelope modifications depending on conditions
(Fendrihan *et al*. 2012; Kixmüller *et al*., 2012), these remain largely unverified at the molecular level
due to these experimental challenges. New methods are needed to isolate proteins and RNA directly
from halite brine inclusions, without allowing the microorganisms to alter their gene expression during
salt dissolution and processing. The development of a new analytical workflow compatible with multi-
omics analyses is best conducted with a known model organism for which there is a large repertoire of
physiological studies and multi-omics data for liquid cultures exposed to a range of conditions
applicable to halite brine inclusions (different oxygen availabilities, salinities, nutrient availabilities,
etc.). For these reasons, we choose the model haloarchaeon *Halobacterium salinarum*, the type strain
of the *Halobacteriales* family (Gruber, 2004; Oren, 2009).

H. salinarum is an appropriate model as it is found both in contemporary NaCl-saturated
aqueous environments such as Great Salt Lake (Post, 1977) and has been detected in halite and closely
related to isolates from ancient salt deposit (Mormile *et al*., 2003). The accumulation of high
intracellular concentrations of potassium (K^+) and chloride (Cl^-) (Engel and Catchpole, 2005), needed
to maintain osmotic homeostasis, induces both biochemical adaptations (in the form of a strongly acidic
proteome) as well as technical challenges to desalt cellular extracts. *H. salinarum* survives in changing

environments by using a complex variable energetic metabolism with a preference for aerobic

respiration but capable to switch under low oxygen conditions to phototrophy via bacteriorhodopsin,

arginine fermentation or anaerobic respiration using dimethyl sulfoxide (DMSO) and trimethylamine

oxide (TMAO) if available (Hartmann *et al*., 1980; Müller and DasSarma, 2005; Falb *et al*., 2008;

Gonzalez *et al*., 2009). Established multi-omics protocols for liquid cultures of *H. salinarum* have

permitted analyses of the cellular responses to a broad range of environmental conditions, including

variations in NaCl (Leuko *et al*., 2009), pH (Moran-Reyna and Coker, 2014), oxygen (Schmid *et al*.,

2007), and temperature (Coker *et al*., 2007), all relevant to halite brine inclusions. This existing

knowledge base far exceeds that developed to date for halophiles isolated directly from ancient halite.

Using the established techniques and known physiological responses of liquid cultures *H. salinarum*,

we were able to develop new analytical techniques.

Here we present this new toolkit for the study of microorganisms within halite, validated using

H. salinarum entrapped in lab-grown halite to answer the question of what physiological changes are

required for microbial cells to remain viable within halite brine inclusions, focusing on the initial phase

of halite entrapment. The workflow developed includes removal of not only surface-attached cells and

nucleic acids but also proteins, with subsequent extraction and desalting of proteins directly from brine

inclusions compatible with mass spectrometry analyses. Applying these methods, we determined the

molecular acclimation of *H. salinarum* cells to inclusions within lab-grown halite by analyzing the

differences in the expressed proteome prior to evaporation and two months after culture evaporation.

Analyses focused on the characterization of cellular activity compared to stationary cells not trapped

within halite, as well as the interactions of cells with halite brine inclusion environment.

3   Material and Methods

All reagents used were analytical grade and suitable for molecular biology.

3.1   Strain and Culture Conditions

Halobacterium salinarum strain NRC-1 (JCM 11081) was grown under oxic conditions in autoclaved

complex medium (CM: 4.28 M NaCl, 81 mM $MgSO_4.7H_2O$, 27 mM KCl, 10 mM trisodium-

citrate.$2H_2O$, 1 % (w/v) peptone Oxoïd® LP0034, pH adjusted to 7.4) following Oesterhelt and

Stoeckenius, 1974 at 37 °C, 180 rpm. Growth was monitored by spectrophotometry at 600 nm (OD_{600}).

Cultures for crystallizations, proteins and RNA extractions were grown to stationary phase ($OD_{600} =$

1.0 to 1.6) to approximate the physiological condition of haloarchaea under natural conditions during

evaporation.

3.2   Laboratory-Grown Halite

3.2.1 Internally Inoculated Halite

Modeling entombment of haloarchaea within halite was already done in laboratory by Norton and

Grant (1988), Gramain *et al*. (2011) and Kixmüller *et al*. (2012). In this study, laboratory-grown halite

crystals were produced following a modified version of protocols of Fendrihan *et al.* (2012), by adding

nutrients to their Tris-buffered NaCl solution (TN buffer; 100 mM Tris-HCl pH 7.4, 4.28 M NaCl) to

simulate cell lysis and metabolite release from less extreme halophiles concomitant with increasing

NaCl concentrations in the natural environment just prior to halite precipitation (Winters *et al.*, 2015).

Briefly, *H. salinarum* cells in stationary growth phase were harvested by centrifugation at 7 500 x g,

10 min, 20 °C and the growth medium removed by washing with sterile TN buffer. Cells were then

resuspended in sterile TNPA buffer (TN buffer with 1 % (w/v) Oxoïd® peptone LP0034 and 0.5 %

(w/v) L-Arg HCl, adjusted pH 7.4) with a ratio of 10 mL TNPA buffer per 500 mg of cells (wet weight).

A 20 mL volume of cell resuspension was evaporated in each sterile 90/14.3 mm Petri dish in a 37 °C

with a 12 h:12 h light:dark photoperiod (66,75 µmol photons.m^{-2}.s^{-2}, verified by a Li-250A, Li-Cor

Inc., Germany) to model natural brine-surfaces. Complete evaporation and drying of precipitated halite

were obtained after ~22 days, followed by a further 60 days (two months) of incubation to study the

early phase of *H. salinarum* entombment within halite brine inclusions.

3.2.2 Externally Inoculated Halite

To produce halite with *H. salinarum* cells localized exclusively at the halite surface, 20 mL of sterile

TN buffer was first evaporated as described above until complete drying (~ 22 days). The resultant

crystals were then collected aseptically using sterilized forceps and inoculating the surfaces of each

crystal with 4.5x10^9 *H. salinarum* cells in stationary phase, applied as a highly concentrated cell

solution after centrifugation (7 500 x g, 10 min, room temperature, RT°C). Crystals were inoculated

on the largest faces (designated here as 'top' and 'bottom', see Figure 2k) by first inoculating the 'top'

face drop-by-drop and spreading out using a sterile inoculating loop. Halite were then dried overnight

at 37 °C, followed by the 'bottom' face the next day with the same protocol.

3.3   Scanning Electron Microscopy (SEM)

Observations and analyses of evaporite crystals were performed by first attaching the halite directly to
aluminum supports with carbon tape followed by carbon coating (7-8 nm thickness). Observations
were performed using a Zeiss Ultra 55 field emission gun scanning electron microscope (SEM-FEG)
equipped with a Brucker EDS QUANTAX detector installed at IMPMC (Sorbonne Université, Paris,
France). Secondary electron images were obtained using an in-column detector (InLens) at 10 kV and
a working distance of 7.5 mm. Energy dispersive X-ray spectroscopy (EDX) analyses and mapping
were performed at 15 kV at a working distance of 7.5 mm.

3.4 In-Brine Cellular Viability Tests

To assess cell viability after halite inclusion, salt crystals were weighted to determine the appropriated
NaCl concentration required for the complex medium to obtain a final concentration of 4.28 M after
halite dissolution as previously described by Gramain *et al.* (2011). Halite and complex medium were
then incubated at 37 °C, 180 rpm. After crystal dissolution, cultures were visually inspected then OD_{600}
and morphology by optical microscopy was measured. Red colored cultures reaching a normal
stationary phase culture ($OD_{600} > 1.0$) were classified as viable and cultures showing no increase in
OD_{600} after 1 month ($OD_{600} < 0.1$) were classified as non-viable.

3.5 Halite Surface Cleaning (Removal of Cells and Proteins)

3.5.1 Cold Atmospheric Plasma Treatment

Cold atmospheric plasmas (*i.e.*, weakly ionized gases) were generated were generated using either a
dielectric barrier device (DBD) or an atmospheric pressure plasma jet (APPJ). They were polarized to
the high voltage using two types of electrical power supply: (1) an alternative current (AC) generator
composed of a function generator (ELC, GF467AF) and a power amplifier (Crest Audio, CC5500) as
well as (2) a pulse generator (RLC electronic Company, NanoGen1 model) coupled with a high voltage
power supply (Spellman company, SLM 10 kV 1200W model). DBD and APPJ were supplied with
different a carrier gas (helium or argon) with/without oxygen. Hence several plasma conditions were
performed to produce active species (radicals, reactive oxygen species but also electrons and photons).
The Table 1 details the experimental plasma conditions that were investigated as well as the resulting
proteolysis to assess the efficiency of proteins removing.

3.5.2 Chemical Treatments

Previous studies applied surface cleaning protocols developed for removal of cells and nucleic acids,

as detailed in Supplementary Table 1. To adapt these protocols for the removal of surface-bound

proteins, we modified the protocol of Sankaranarayanan *et al*. (2011) using shorter time for baths to

avoid dissolution of smaller laboratory-grown crystals. Briefly, crystals were incubated in successive

5 min baths: 4.00 - 4.99 % NaOCl (Honeywell Research Chemicals; avoiding commercial bleach

solutions that induced rapid and near-complete crystal dissolution) in saturated NaCl and 10 M NaOH

in saturated NaCl supplied or not with 10 N HCl followed by 100 mM Na_2CO_3 in saturated NaCl. A

sterile saturated NaCl was used to wash off each treatment solution between each successive bath,

testing both passive wash baths and active-spray washes using a hemolytic pump.

3.6 Protein Extraction and Preliminary Desalting

All solvents used for protein extractions were HPLC-grade and suitable for mass spectrometry analysis.

Extractions were performed with TRIzol Reagent™, based on the halophilic protein extraction protocol

of Kirkland, *et al*. (2006). This protocol was modified for protein extraction directly from halite crystals

with high salt content and proteomics analysis by mass spectrometry. Compatibility for liquid cultures

was kept as described below. All steps described were performed with autoclaved glassware to avoid

any organics contamination from plastics (bench top protocol see Supplementary Section 2). Briefly,

crystals were fully immersed in 5 mL of TRIzol Reagent™ in 30 mL glass centrifuge tube and crushed

using autoclaved glass stir-rod. After 20 min incubation at 60 °C, total RNA was extracted from the

resulting cell lysate by adding 1 mL of 100 % chloroform, incubating for 5 min at RT°C followed by

phase separation by centrifugation (10 000 x g, 20 min, 4 °C). The chloroform-containing aqueous

phase was removed with autoclaved glass Pasteur pipet and 1.5 mL of 100 % ethanol was added to

precipitate DNA. After 3 min at RT°C, supernatant was collected by centrifugation (2 000 x g, 10 min,

4 °C). Proteins were precipitated with 7.5 mL of 100 % isopropanol and collected by centrifugation

(10 000 x g, 20 min, 4 °C). The resulting protein pellet was washed twice to remove phenol traces

using 5 mL of 0.3 M guanidine-HCl in 95 % ethanol to denature the proteins, followed by a third wash

step with 5 mL of 100 % ethanol to remove any residual guanidine-HCl. Each wash step was performed

by incubating the cell pellet in the solution for 20 min at RT°C followed by centrifugation (10 000 x

g, 20 min, 4 °C). Protein desalting was accomplished by two successive protein precipitations with 2

mL of 100 % glacial acetone (-20 °C) and centrifugation (10 000 x g, 20 min, 4 °C). After acetone

removal, the pellet was completely dried under laminar flow. Then, proteins were solubilized in 1 M

NaHCO$_3$ with 0.1 % SDS at RT°C for two days and quantified with a bicinchoninic acid (BCA)

proteins assay (Pierce) using bovine serum albumin (BSA) standards concentration from

manufacturer's instruction in the case of quantitative mass spectrometry and adapted BSA standard

concentrations (enable low protein concentration assay, see Supplementary Section 3) for surface

proteins removal procedure. Proteolysis rate was determined by proteins quantity comparison with and

without treatments.

Total proteins were extracted from *H. salinarum* cultures in stationary growth stage using the

same procedure. For this, 2.0 x10^{10} cells from liquid cultures were pelleted by centrifugation (7 500 x

g, 10 min, 20 °C) and the cell pellets directly resuspended in 5 mL TRIzol Reagent™ solution, prior

to following all steps described above for halite samples. After solubilization of the protein pellet,

solubilization with 1M NaHCO$_3$ with 0.1 % SDS at RT°C, protein quantification was performed using

the BCA assay (Pierce) protein assays as per the manufacturer's instructions.

3.7   iTRAQ® Isobaric Labelling and Mass Spectrometry

Aliquots of 100 µg of proteins for each sample condition and replicate were reduced using 2 mM of

tris-(2-carboxyethyl) phosphine (TCEP) at 37 °C for 1 h, and alkylated with 5 mM of iodoacetamide

30 min in the dark at RT°C prior to digestion with 5 µg of trypsin Gold (Promega) for 15 h at 37 °C.

After digestion, peptides were directly desalted by solid phase extraction (SPE) using C18 cartridges

(Sep-Pak C18 Plus Short 400 mg Sorbent, Waters). The resulting peptides were dried by speed-vac

and resuspended in tetraethylammonium bromide (TEAB) 0.5 M prior to labelling. iTRAQ® labelling

was performed according manufacturer's instructions (Applied Biosystems). Briefly, each of the

iTRAQ® isobaric labelling reagents were reconstituted with isopropanol and then added to the 50 µg

of protein digest (113, 114, 115 and 116 iTRAQ® isobaric labels for proteins from liquid controls and

117, 118, 119 and 121 for halite brine inclusion protein extractions). After 2 h at RT°C, samples were

desalted again with C18 SPE. The labelled peptides eluted were then dried by speed-vac and

resuspended in 2 % acetonitrile, 98 % H$_2$O with 0.1 % formic acid, see Supplementary Section 5.

Labelled peptide samples were analyzed by mass spectrometry as previously described (Pinel-

Cabello *et al.,* 2021) on a Q Exactive HF tandem mass spectrometer (Thermo Scientific) coupled to an

UltiMate 3000 Nano LC System. Peptides were desalted online on an AcclaimPepmap100 C18

precolumn (5µm, 100 Å, 300 µm i.d x 5 mm) and further resolved on a nanoscale AcclaimPepmap100

C18 column (3 µm, 100 Å, 75 µm i.d. x 500 mm) at a flow rate of 200 nL min^{-1} using a 120-min

gradient of 4-32% acetonitrile. A Top 20 strategy was applied in data dependent acquisition mode. Full

scan mass spectra were acquired from 350 to 1800 *m/z* at a resolution of 60,000 with an automatic gain

control (AGC) target set at 3×10^6 ions. MS/MS fragmentation was initiated when the ACG target

reached 10^5 ions with an intensity threshold of 9×10^4. Only precursor ions with potential charge states

of 2^+ and 3^+ were selected for fragmentation applying a dynamic exclusion time of 10 s.

3.8   Mass Spectrometry Data Analyses

3.8.1 Protein Identification

Protein identifications were performed using PEAKS® X-Pro software (64 bits version, 2020,

BioInformatics solutions). It allows database search assisted *de novo* sequencing against the protein

coding sequences from *H. salinarum* NRC-1 (8 533 entries from NCBI, downloaded 2021/08/03).

MS/MS assignment was searched with the following parameters: (1) mass tolerance of 10 ppm on the

parent ion, (2) mass tolerance of 0.005 Da for fragment ions from MS/MS; (3) carbamidomethylated

Cys (+57.0215) and iTRAQ® isobaric tag Lys and N-terminal (+304.2054) as static modifications; and

(4) oxidized Met (+15.9949), deamidated Asn and Gln (+0.9840) and iTRAQ® isobaric tag Tyr

(+304.2054) as variable modification. The false discovery rate (FDR) was estimated with decoy-fusion

option.

Then, resulting proteins lists were then filtered allowing FDR < 1 % (corresponding to a -

10logP score above 25) for peptide-spectrum matches (PSMs), protein identification using minimum

2 unique peptides per proteins (a unique peptide is a peptide with a -10logP score above the peptide

filtering threshold that can be mapped to only one protein group).

3.8.2 Protein iTRAQ® Quantitation

The eight labelled samples (four replicates each of proteins from liquid stationary cultures and from

halite brine inclusions) were mixed in equimolar ratios and injected in nanoLC-MS/MS in triplicate to

reduce instrument variability. Quantitation was performed using PEAKS Q (quantitation program)

iTRAQ 8-plex type with 10 ppm mass tolerance and only peptides with a score above FDR threshold

1 % are used to quantify the identified proteins. Resulted quantitation was filtered accepting only

proteins groups with fold change ≥ 2, at least two unique peptides and FDR adjusted ≤ 1 % for both

proteins identification and fold change significance using ANOVA significance method. Mass

spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE

partner repository with the dataset identifier (*under submission*).

3.8.3 Sample Comparisons and Functional Annotation

Potential biases during analysis were avoided by filtering protein identifications produced using the

PEAKS software to group proteins having multiple annotations in the NCBI database. To do so, we

developed custom BASH and PYTHON scripts to merge all descriptions for one proteins identifier

(favoring in order AAG-type, DAC-type, WP-type and then Q9H-type identifier). Venn diagrams were

computed with VennDiagram and ggplot2 packages in R script. Functional proteins annotation was

done with Kyoto Encyclopaedia of Genes and Genomes (KEGG, Kanehisa and Goto, 2000) using

blastKOALA tool (Kanehisa *et al*., 2016; https://www.kegg.jp/blastkoala/), assigning K numbers to

identified proteins (see Supplementary Table 6-2). K number were then used for mapping identified

proteins onto pre-existing metabolic pathways using KEGG Mapper search tools

(https://www.kegg.jp/kegg/mapper/search.html ; see Supplementary Table 6-3).

3.9   Total RNA extraction and analysis

Total RNA for transcriptomics analyses were extracted using TRIzol Reagent™ method described

above for protein extractions with the following modifications. To avoid RNA degradation, the TRIzol

Reagent™ mixture with pulverized halite crystals was incubated for 5 min at RT°C rather than 20 min

at 65 °C. The chloroform phase was then collected by centrifugation and RNA purification was perform

using RNeasy MinElute cleanup columns (Qiagen). Resulting RNA were quantified using QuBit

Q32857 (Invitrogen) the purity assayed using Nanodrop 2000/2000c (Thermo Scientific), and the

quality was determined with the BioAnalyzer 2100 system (Agilent). All steps after RNA extraction

were performed according manufacturer's instructions with RNAase-free materials and solutions.

4   Results

4.1   Characterization of Laboratory-Grown Halite Containing *H. salinarum*

In order to study the physiology of *H. salinarum* cells after entombment within brine inclusions, we

first needed to characterize the laboratory-grown halite, including resultant crystal size, cell

localization, and surface-bound biomolecules. This enabled the establishment of selection criteria for

downstream proteomics analyses and the development of protocols to exclude any 'contaminant'

proteins from the surface of the crystal.

4.1.1 Variability of *H. salinarum* cells from Inoculated Laboratory-Grown Halite

Laboratory-grown halite crystals produced from TNPA solutions containing *H. salinarum* cells using

a slow evaporative process exhibited heterogeneous crystallization with respect to crystal size and

coloration. As shown in Figure 1, even replicates produced under the same conditions produced

different quantities of crystals of varying lengths and widths, but with relatively uniform thicknesses

(±1 mm) of hopper and fishtail crystals. Color variations observed for halite in this study were likely

the result of heterogeneities in the number of brine inclusions containing pigmented *Halobacterium*

cells. The red color indicative of *H. salinarum* cells tended to concentrate in the center of hopper

crystals where inclusions were most abundant (Roedder *et al*., 1984). Individual crystals of similar size

and coloration were collected aseptically for further study, selecting only halite without visible defects

signaling potential rupture of near-surface brine inclusions.

4.1.2 Contaminating Surface-Bound Cells and Proteins

To evaluate the potential for 'contaminating' microbial cells or residual biomolecules at the crystal

surface, the elemental composition of halite with and without *H. salinarum* cells were determined by

SEM-EDX. Elemental analyses of control halite derived from evaporation of the TNPA solution only

exhibited the expected sodium chloride crystal surface, along with dense, irregular C agglomerates

(Figure 2c-f) derived from the presence of organics (peptone, L-Arg) in the TNPA solution. The

addition of *H. salinarum* cells to the TNPA solution prior to evaporation produced similar structures

to control crystals, with no intact cells of *H. salinarum* visible on halite surfaces (Figure 2i,j). However,

EDX analyses revealed differences in the distribution of carbon compared to control samples, as well

as the presence of microscopic potassium crystals at the halite surface (Figure 2j). These differences

were likely due to lysis of surface adhered *H. salinarum* cells releasing high concentration of cytosolic

K^+ and Cl^- accumulated as part of the "salt-in" osmoadaptation strategy. In order to assess both the

potential for surface bound contaminants, including viable cells as well as proteins, externally

inoculated halite were produced (Figure 2k). After two months incubation, halite surface-attached cells

were revealed to be (at least partially) viable according positive viability test. After growth back,

culture observations by optical microscopy showed rod-shaped cells with low motility. Additionally,

SEM-EDX analysis of externally inoculated crystals revealed a similar elemental composition and

distribution to that of internally inoculated halite (bearing cells both within brine inclusions as well as

on the halite surface) after two months post-evaporation (Figure 2m,n). Taken together, these results

confirm the presence of both viable cells and cellular debris on the surface of halite after two months

of desiccation, validating the need for halite surface cleaning procedure to remove any residual

proteins.

4.2   Development of Total Protein Isolation Methods for Halite

For proteomics approaches inside crystals without prior dissolutions, a new methodology for direct

proteins extraction from brine inclusions was developed. This method was based on the protocol of

Kirkland *et al.* (2006) for liquid cultures to allow mass spectrometry analysis compatible both for

crystals and liquid cultures with TRIzol Reagent™ (see Supplementary Section 4 for details on the

parameters tested during protocol development and optimization). Briefly, protocol was modified to

optimize desalting steps (due to direct halite extraction with high NaCl content), proteinaceous pellets

dissolution and proteinaceous digestion.

4.3   Development of Proteolysis Protocols for Removal of Halite Surface-Bound Contaminants

Previous halite cleaning procedures from Gramain *et al.* (2011) and Sankaranarayanan *et al.* (2011)

(see Supplementary Table 1) were developed to deactivate surface-contaminating microorganisms and

remove surface-contaminating DNA. However, these methods were not designed for the removal of

other types of biomolecule contaminants, such as proteins. A new, modified surface cleaning procedure

was therefore needed to remove surface protein contamination to enable extraction of proteins only

from within halite brine inclusions while avoiding crystal dissolution during the cleaning process. To

meet these requirements, both cold plasma treatments and chemical protocols were tested using

externally inoculated halite.

Cold plasma exposure is routinely used in astrobiology to sterilize spacecraft surfaces of

microbial contaminants prior to launch. For this reason, a cold plasma approach was investigated here

for the decontamination of halite surface. Two plasma sources – a DBD and an APPJ – were tested as

an alternative to conventional liquid solvents. The expected benefit was to minimize halite dissolution-

recrystallization events, hence ensuring the survival of the microorganisms within the halite. However,

owing to their insulating properties and irregular topography, halite crystals attenuate plasma electric

field and/or bend its field lines. As a result, and whatever the implemented plasma treatment (Ar, He,

He mixed with O_2), the externally inoculated halite crystals still retained the pink coloration which is

indicative of the protein and lipid pigments of *H. salinarum* cells (data not shown). As shown in Table

1, the most efficient plasma treatment drove to partial cell lysis (as determined by culture-based

viability tests following growth by OD_{600}) and to a proteolysis of only 39.1%: a value that remains

insufficient for the application.

As an alternative, chemical treatments were tested, using protocols modified from the

microorganism deactivation and DNA removal treatments of Sankaranarayanan *et al.* (2011) to extend

to proteolysis. First, reduction of the exposure times in each chemical bath from previously published

protocols were tested to avoid crystal dissolution, while maintaining the use of NaCl-saturated

chemical solutions to avoiding crystal dissolution. Subsequent, 5-minute sequential NaOCl and NaOH

treatments, either with or without additional HCl treatments, resulted in insufficient halite surface

proteolysis (Supplementary Figure 5-1a). This was likely due to the rough structure of hopper crystals,

which could allow proteins to remain attached during passive chemical baths. After chemical

treatments have partially degraded the contaminating biomolecules and reduce their attachment to the

halite surface, active-spray was used for wash steps to increase the efficiency of biomolecule removal

and prevent re-association with halite surfaces, compared to the use of passive wash baths

(Supplementary Figure 5-1b). Externally inoculated halite post-treatments with and without active-

spray, present reduced coloration. This active-spray method also allowed for both deactivation of halite

surface-bound microorganisms (as indicated by a lack of culture growth in post-treatment viability

tests) and better protein lysis compared to passive chemical bath washes. Using the protein extraction

method described above, the efficacy of active-spray method for removal of surface-bound proteins

was tested comparing NaOCl-NaOH and NaOCl-NaOH-HCl protocols. A shown in Figure 3a, addition

of an acid wash results in a proteolysis efficiency of 93.0 ±4.3 %, compared to from of 83.7 ±15.9 %

without the HCl treatment step (residual proteins 51.6 ±31.6 µg and 120.5 ±117,6 µg, respectively,

n=3). These results suggest that NaOCl-NaOH-HCl active-spray washes protocol (Figure 3b) present

optimal and less variable surface proteolysis for the case of this study, working on small laboratory-

grown halite crystals.

To ensure that this active-spray chemical treatment did not affect *H. salinarum* cells within the

halite brine inclusions, growth was monitored for crystals containing cells trapped in brine inclusions

after surface chemical cleaning with active-spray NaOCl-NaOH-HCl treatment. Viable cultures were

obtained from dissolution of the crystals in CM growth medium, thereby validating the active-spray

chemical surface protein removal protocol for use in studying biomolecules within halite brine

inclusions.

These new methods for removal of surface-bound proteins followed by extraction of total

proteins from halite brine inclusions using a modified TRIzol Reagent™ method with desalting were

then applied to answer the question of how *H. salinarum* acclimates to halite brine inclusions over the

initial phase post-evaporation using proteomics

4.4   Proteomic Shifts in *H. salinarum* Induced by Acclimation to Halite Brine Inclusions

Total proteins were extracted from the brine inclusions of halite after two months of complete dryness,

with surface-bound proteins removed using these new methods detailed above. These brine inclusion

extracts were compared to total proteins extracted from *H. salinarum* cultures in stationary growth

phase, representing conditions prior to halite formation. The proteins unique to each condition, as well

as the common proteome between liquid cultures and from halite interiors were analyzed to determine

how *H. salinarum* acclimates to halite brine inclusions.

Using mass spectrometry of four replicates from each condition, 1 249 total proteins were

identified from stationary phase liquid cultures and 1 052 from halite brine inclusions, representing 1

342 unique proteins. Identifications were obtained with high confidence using FDR < 1 % and at least

2 unique peptides. Internal comparisons of the four replicate samples per condition revealed a core

proteome of 839 proteins common to all sample replicates for liquid cultures and 715 proteins for halite

brine inclusion extracts (Figure 4a, b). Of these, 655 proteins were expressed by cells both in liquid

cultures and within halite brine inclusions (after   referred to "shared proteins"), while 60 specific to

halite brine inclusion samples and another 184 only identified from liquid cultures (Figure 4c,

Supplementary Table 6-1). KEGG blastKOALA searches provided matches for roughly 75 % of these

shared proteins (488 of 655 shared proteins), as well as 62 % of proteins unique to halite brine inclusion

samples (37 of 60 proteins) and 52 % of proteins unique to liquid cultures (95 of 184 proteins) (see

Supplementary Table 6-2), allowing for functional pathway reconstruction. The halite brine inclusion

proteome contained genome maintenance and repair proteins (Nth, UvrB, MutS2) not found expressed

by cells from liquid cultures, along with multiple enzymes implicated in amino acid metabolism,

metabolism of cofactors, and membrane transport (see Table 2 and Supplementary Table 6-3).

A qualitative approach was insufficient to determine differential regulation of protein

expression for *H. salinarum* cells within halite brine inclusions. Instead, a quantitative mass

spectrometry analysis was also performed. Results from all three injections were pooled to reduce

instrument variability (Figure 5). Subsequent 68 proteins from halite brine inclusions were identified

with statistically significant changes in expression levels compared to liquid cultures (fold change ≥ 2;

see Supplementary Table 6-4 and Supplementary Figure 6-1). KEGG blastKOALA searches provided

functional pathway results for 80 % of down-regulated proteins (35 of 44 proteins) and 71 % of up-

regulated proteins (17 of 24 proteins) (see Supplementary Table 6-2). Up-regulated proteins included

metal transporters, redox reactions, acetyl-CoA production, and the UgpB glycerol-3-phosphate-

binding protein precursor. Down-regulated proteins were predominantly ribosomal subunits, indicating

reduced translational activity (see Table 2 and Supplementary Table 6-3). To determine whether there

was a corresponding downwards shift in transcriptional activity for cells inside halite brine inclusions

compared to liquid cultures, RNA extractions were performed.

4.5   Reduced Total RNA Concentration Within Halite Brine Inclusions

The advantage of a TRIzol Reagent™-based extraction method is that it allows for extraction of not

only proteins, but also nucleic acids (DNA, RNA), and lipids (Podechard *et al.,* 2018) for complete

multi-omics analyses. To this end, the new methodology presented here was applied to extract and

purify RNA from surface cleaned, internally inoculated halite after two months of incubation at 37 ˚C

to complement the proteomic analyses. Consistent with the reduced quantities of ribosomal proteins

observed for halite brine inclusion extracts, low quantities of RNA were obtained from brine extracts

compared to stationary cells extracts (Table 3). In addition, the quality and purity of the resultant RNA

from halite was lower than that from liquid cultures (Supplementary Figure 6-2). To determine if this

low-quality RNA was due to the age of the samples (two months after drying) or degrading effects of

halite salts on RNA during extraction, extractions of RNA were performed from fresh externally

inoculated halite (*i.e.*, approximatively one week after external cells addition) using the same protocol.

RNA extractions from these 'young' crystals presented higher quantity and purity compared to those

from halite brine inclusions after two months, but still lower quality and purity than RNA extracted

from liquid cultures.

5   Discussion

The initial objective of this study was to determine the molecular acclimation of haloarchaea to

entrapment within halite brine inclusions using a proteomics approach. To this aim, we first needed to

(1) better characterize the process of laboratory-based evaporation for halite entrapment of

haloarchaeal and its effects on both the resulting halite and the microbial cells, (2) develop efficient

methods for removal of halite surface-bound cells and biomolecules in order to isolate only proteins

contained within the brine inclusions, (3) develop effective methods for protein extraction directly from

halite brine inclusions in a rapid time period to avoid alterations in protein expression that can occur

during gradual salt dissolution, and finally (4) combine these approaches to analyze total proteins and

RNA from *H. salinarum* extracted from halite brine inclusions and compare these to control samples

from stationary phase liquid cultures in order to propose a model of acclimation within halite. A slow

evaporative process was chosen to generate the internally inoculated halite to model natural processes

that occur over time scales sufficient for cellular acclimation to the environmental changes, rather than

a rapid "shock" evaporative process.

5.1 Biotic/Abiotic Interactions Influencing Halite Formation and *H. salinarum* Viability

Similar to previous studies of survival of halophiles within halite brine inclusions, a NaCl-based

buffered evaporation solution was used in this study (Norton and Grand, 1988; Fendrihan *et al*., 2012;

Kixmüller *et al*., 2012). However, here we chose to add nutrient sources (1 % peptone, 0.5 % L-Arg-

HCl) to simulate organics present in the natural environment. This enabled us to study the role of

metabolism in the survival and acclimation processes of *H. salinarum* cells to the early phases of

entrapment within halite brine inclusions. The presence of microbial biomass and additional organics

can affect halite precipitation processes and also explain the observed crystallization heterogeneity

observed. For example, Norton and Grant (1988) previously demonstrated a positive correlation

between the initial microbial cell density in liquid culture and the quantity of halite brine inclusions

formed during evaporation.

In addition, the presence of nutrients seemed to diminish rather than increase the duration of

survival. While Gramain *et al*. (2011) observed growth of *H. salinarum* NRC-1 after 27 months post-

entombed in halite, there were some notable differences in the experimental parameters used. Gramain

et al. (2011) used evaporation buffers with different compositions of nutrients to the evaporation buffer

(either without nutrients or with addition of 0.01 % Difco yeast extract and 0.0075 % Merck casein

hydrolysate diluted from the standard concentrations in modified Payne's medium of 1 % and 0.75 %,

respectively, Payne *et al.,* 1960), incubated the resulting salt crystals in the dark at RT°C rather than

with a 12 h:12 h diurnal cycle at 37 ˚C as used here, and did not appear to employ halite surface

sterilization or cleaning protocols prior to these survival tests. In contrast, our viability tests of *H.*

salinarum cells from internally inoculated crystals incubated at 37 ˚C over 80 days prior to subjected

to surface organics cleaning treatments, dissolution and culturing showed no growth. The lower

duration of survival observed for *H. salinarum* cells inside halite in our study may be the result of

increased metabolic activity due to the higher concentration of nutrients in TNPA solution (1%

peptone) compared to the 100-time diluted modified Payne's medium used in Gramain *et al*. (2011).

This could result in an inhibitory effect due to the accumulation of toxic waste products within the

closed microenvironment of halite brine inclusions. On the other hand, in the absence of surface

sterilization of halite used in the viability studies in Gramain *et al*. (2011), some of the surviving cells

observed in the study may have been the result of surface-adhered cells rather than cells within the

halite brine inclusions. These results also suggest a possible survival advantage for cells on halite

surfaces rather than those within the brine inclusions over the early stages of evaporation and

acclimation, an hypothesis supported by the findings of Gramain *et al.* (2011) showing no difference

in growth for *H. salinarum* cells evaporated in salt buffer without nutrients and those containing the

diluted Payne's medium nutrients. This epilithic lifestyle stay probable also due to organics observed

form surface halite, as well as microscopic sylvite (KCl) crystals derived from the lysis of *H. salinarum*

cells were also observed on the surface of the laboratory-grown halite while the vacuum conditions for

SEM observations may have resulted in rupture of unfixed cells.

All together, these results of surface contamination confirm that the isolation of proteins

exclusively from halite brine requires the removal of halite surface-bound cells and biomolecules.

5.2   Removal of Halite Surface-Bound Microorganisms and Biomolecules

The small sizes of laboratory-grown halite do not allow using treatment processes that could result in

salt dissolution. Cold atmospheric plasmas were therefore tested in an effort to avoid the use of liquid

cleaners based on their effectiveness for the sterilization of spacecraft surfaces for planetary protection

(Shimizu *et al.*, 2013) and agricultural applications (Judée *et al.*, 2021). Unfortunately, none of the

plasma conditions tested in this study (gas mixtures, power delivery modalities) enabled a complete

removal of proteins from the rough textured halite surfaces. However, the results remain encouraging

insofar as cold plasma has demonstrated effects on proteolysis which must now be amplified. Further

experiments will be required to innovate an *ad hoc* plasma process delivering active species at higher

densities while treating the whole surface of halites regardless their roughness surface, dimensions or

dielectric permittivity.

Therefore, the deactivation of microorganisms and removal of nucleic acids from halite

surfaces have been performed by adapting chemical wash methods first developed by Rosenzweig *et*

al. (2000) and then optimized by Gramain *et al.* (2011) and Sankaranarayanan *et al.* (2011). By

reducing the exposure times in sequential NaOCl, NaOH, and HCl treatments, and replacing passive

chemical baths with an active spray process, we were able to achieve sufficient proteolysis of surface-

bound proteins for downstream isolation of proteins exclusively from within halite brine inclusions,

while avoiding halite dissolution during treatment. Compared to total proteins extracted from surface

sterilized internally inoculated crystals (± 1 mg order), residual surface bound proteins (51.6 µg) were

marginal. However, observations of residual pigmentation after halite treatment suggest the presence

of non-protein pigments such as carotenoids. Further refinements are therefore needed before future

studies can isolate lipids exclusively from within halite brine inclusions.

5.3   Limitations of the TRIzol Reagent™ Based Method for Direct Extraction of Biomolecules

From Halite Brine Inclusions

TRIzol Reagent™ has the distinct benefit of enabling selective extraction of proteins, lipids, and

nucleic acids. While TRIzol-based protein extraction methods have previously been used and validated

to study proteins from liquid cultures of haloarchaea (Kirkland *et al*., 2006; Bidle *et al*., 2008). In this

study procedure was modified for direct proteins extraction on salts crystals and subsequent

desalinations steps were developed to allow mass spectrometry analysis possible from these high salt

extracts. The results obtained after modifications of the procedure by Kirkland *et al*. (2006) procedure

appear compatible and efficient for quantitative mass spectrometry approach directly on halite brine

inclusions which confirm optimal modifications. In addition, compared to crystal dissolution method

prior to proteins extraction, this methodology avoid bias related to dissolution-associated proteome

changes.

However, while peripheral membrane proteins were identified, transmembrane peptides were

absent from the mass spectrometry dataset. We hypothesize that this was due to retention of

transmembrane protein domains with the lipid membrane fraction during TRIzol Reagent™ extraction

in residual phenol phase after protein precipitation with isopropanol. Importantly, some transmembrane

proteins could be identified by peptides outside the membrane-spanning domains *e.g.*, the ArcD

arginine-ornithine antiporter protein, for which seven cytosolic- and extracellular-facing peptides were

identified (see Supplementary Figure 7-1). One explanation of this phenomenon is the desalting effect

of isopropanol (as suggested by Kirkland *et al*., 2006) that can result in increased protein instability,

leading to loss of conformation and subsequent peptide fragmentation for peptides outside the

transmembrane regions. While these results indicate that further membrane disruption steps may be

needed for complete extraction of all transmembrane proteins, similar to the protocol used by

Podechard *et al.,* 2018 to isolate membrane lipids using TRIzol, the potential bias introduced by our

methodology is limited, as shown in by the identification of peptides extending to either side of the

lipid membrane.

While the low quantities of RNA extracted from halite brine inclusions were likely the result of

low transcriptional activity based on the down regulation of ribosomal proteins observed during

proteomics analyses (see following section), it might also be due to the age of the halite with subsequent

degradation of RNA after cells trapping. In addition, RNA quantity, purity and quality were insufficient

to allow transcriptomics analysis for brine extracts. The quantity, quality and purity scores for fresh

externally inoculated crystals were higher than those for surface sterilized crystals incubated for two

months. Further desalting steps may help to improve future RNA extraction directly from halite.

5.4   Acclimation of *H. salinarum* to Halite Brine Inclusions

We applied the slow laboratory evaporation method to *H. salinarum* cultures in TNPA buffer followed

by removal of halite surface-bound cells and organics (particularly proteins) and selective extraction

of proteins from within halite brine inclusions described in the preceding sections. This allowed us to

study the early (two month) acclimation of the haloarchaeal cells to halite brine inclusions using a

proteomics approach. Stationary growth phase cultures were used as a control to approximate the cell

physiology prior to a slow evaporative process. Our analyses were focused on two main themes: (1)

cell activity within halite brine inclusions, and (2) interactions between cells and local

microenvironment. The investigations of the cell activity targeted proteins involved in central

metabolism, energy production, cell division, replication, transcription and translation pathways. The

cell interactions with the closed halite brine inclusion microenvironment were examined via cell

envelope proteins (surface layer cell wall proteins and transporters) as well as proteins involved in the

motility processes (chemotaxis, gas vesicle and archaellum).

5.4.1 Cellular Origin of Total Brine Extracted Proteome

The proteins extracted using the protocol presented here represent the total protein complement of the

halite brine inclusion. These proteins may be components of viable or non-viable cells, or even proteins

released by cells into the extracellular environment of the brine inclusion. Many questions remain about

the potential for salts to preserve proteins over time as molecular biosignatures, particularly within the

protected evaporite brine inclusion microenvironment. However, as the *H. salinarum* cells studied here

remained viable during the two months of this study, we can hypothesize a cellular origin for the

extracted proteins.

5.4.2 Cell Activities Within Halite Brine Inclusions

Table 2 shown similar proteins between cells from the stationary growth stage and brine extracts,

suggesting reduced cell division in both conditions. Proteins involved in DNA replication, reparation

and transcriptional pathways were shown to be shared without down-regulation, which suppose active

cells or at least few changes comparing late growth stage and brine entombment. However,
translational proteins were not detected in brine extracts, suggesting that protein synthesis decrease
compared to stationary cells. This surprising result can be explained by a transcriptional activity
directed not to mRNA synthesis but to regulatory RNA. However, this hypothesis stays open to
criticism, as tRNA synthesis and ribosomal biogenesis proteins remains similar for both conditions.

5.4.3 Halite brine Inclusions Cells Retain Proteins for Central Metabolism

Proteins involved in central metabolism, including key enzymes involved in
glycolysis/gluconeogenesis and the TCA cycle, formed part of the shared proteome between halite
brine inclusion extracts and stationary phase liquid cultures (Table 2). Additionally, proteins involved
in phototrophy, arginine fermentation and aerobic respiration were shared between halite brine
inclusions and stationary growth phase liquid cultures. The microenvironment of halite brine inclusions
was assumed to be oxygen poor due to the combination of low DO_2 at high salinities as well as the
closed microenvironment of brine inclusions. The metabolism of *Halobacterium* cells trapped within
such inclusions is therefore hypothesized to shift from aerobic metabolism by respiratory chain to
anaerobic fermentation or phototrophy through arginine deiminase pathway (ADI). Incubations of
halite were performed with a 12 h:12 h light:dark photoperiod, allowing for phototrophy under oxygen-
limited conditions.

Quantitative analyses showed that the majority of proteins involved in pyruvate metabolism
maintained consistent expression levels after two months of halite inclusion with the exception of up-
regulation of proteins involved in acetyl-CoA production. Surprisingly, transporters and enzymes
involved in ADI pathway were expressed by both cells in stationary phase liquid cultures and brine
inclusion extracts, including the arginine deiminase, ornithine-carbamoyl transferase, carbamate
kinase, arginosuccinase, arginosuccinate synthetase, arginine-ornithine antiporter and an IclR family
transcriptional regulator. Only arginine deaminase was found significantly up-regulated for cells from
halite brine inclusions. Proteins for key components of the electron transport chain, including NADH
dehydrogenase/oxidoreductase subunits B, C, D and I, NAD(P)/FAD-dependent oxidoreductase,
succinate dehydrogenases and ATP synthase subunits A, B, C, D, E, H and I were not found to be
differentially regulated between cells from free-living (liquid cultures) or trapped within halite brine
inclusions (with the exception of with F subunit lacked in one brine inclusion extract). The bacterio-
opsin activator-like protein and Htr signal transducer were common to all replicates and conditions,
while bacterio-opsin activator was only found in two of the four halite brine inclusion replicate

samples. Confirmation of the precise metabolic pathways used by *H. salinarum* within halite brine

inclusions will require further analyses, overcoming the challenges of accessing the closed halite

system.

Acclimation to the halite brine inclusions microenvironment could presumably involve reduced

or silenced cell division, DNA replication and repair pathways. However, eight proteins involved in

chromosome partitioning were found shared between halite brine inclusion extracts and stationary

growth phase cultures, along with 11 other proteins involved in DNA replication including matches

for DNA primase, polymerase, gyrase, topoisomerase and replication factors. Only four DNA

replication proteins were found to be specific to stationary growth phase proteomes. This suggests that

either such proteins are constitutively expressed, or that DNA replication and cell division occur in a

similar fashion for both stationary growth phase cells and those within halite brine inclusions.

DNA repair processes were represented both within halite brine inclusions and liquid cultures

by the DNA photolyase, Rad50, the UvrD repair helicase, UvrA excinuclease, RadA recombinase,

Rad25 DNA repair helicase, Fen1 repair endonuclease and the Hel208 helicase are found shared

between liquids and brine extracts. Other DNA repair proteins were specific to either halite or liquid

samples. Stationary growth phase cultures contained additional repair proteins including the RmeR

site-specific deoxyribonuclease, endonuclease IV, RecJ single-strand exonuclease, MutL mismatch

repair protein, DNA gyrase and N-glycosylase, whereas the UvrB, MutS and endonuclease III proteins

were exclusive to halite brine inclusion proteomes.

Investigations of DNA transcription identified a shared proteome between liquid cultures and

brine inclusions including proteins for transcriptional initiation factors (TFIIE and TFIID) and

transcriptional machinery (RNA polymerase subunits rpoA2, rpoB1, rpoB2, rpoD and rpoE1). Proteins

involved in transfer RNA biogenesis such as tRNA ligases for 18 different amino acids along with two

ribonucleases completed the shared transcriptional proteome of liquid cultures cells and halite extracts.

Only tRNA ligases specific for tryptophan and threonine, four unique transcriptional proteins (rpoF,

rpoH, rpoN and nusA termination protein) and one unique transcription initiation factor (TFIIB) were

only found only in stationary growth phase cell extracts.

While early acclimation to the halite brine inclusion microenvironment resulted in nuanced

differences in the proteome of *H. salinarum* cells regarding cell division, replication, DNA repair and

transcriptional processes, stark differences were observed for proteins involved in translation activities.

Translational activities were severely restricted for cell within halite brine inclusions, as evidenced by

the down-regulation of 27 of the 42 ribosomal proteins shared by cells from both conditions tested.

Only one ribosomal protein was up-regulated. The shared proteome between cells in halite brine

inclusions and those in liquid culture included an additional 10 proteins involved in ribosome

biogenesis, and 10 translation initiation factors for which non showed any significant up- or down-

regulation. Chaperones proteins were showed a high degree of conservation between conditions, with

thioredoxin, chaperone DnaJ, DnaK along with the GrPE stimulator, Hsp20, thermosome subunit and

prefoldin all shared by both for liquid and halite extracts. Only prefoldin beta subunit was down-

regulated for cells from halite brine inclusions.

Taken together, these data indicate a high level of similarity between stationary growth phase

cells in liquid culture and those trapped within halite brine inclusions. Acclimation to the closed brine

inclusion microenvironment appears to severely limit translational activity. However, expressed

proteins are preserved indicating low levels of protein turnover. Cells maintain at least the capacity for

central metabolism, DNA repair, replication, and even cell division comparable to stationary growth

phase cells.

5.4.4 Interactions Between Cells and the Brine Inclusion Microenvironment

the potential interactions between *H. salinarum* cells and the local microenvironment within halite

brine inclusions focusing on cell surface and mobility processes involved in responses to external

stimuli. Previous investigations of the survival of *Halobacterium* cells trapped within salt crystals have

proposed potential for cell envelope modifications with surface proteinaceous layer (S-layer) shed

(Fendrihan *et al*., 2012). However, this result seems dependent on conditions (Kixmüller *et al.,* 2012).

S-layer shedding thought to improve exchanges between cells and the local microenvironment.

However, in this study S-layer proteins were identified in all liquid culture and halite samples without

significant up- or down-regulation. This suggests that the S-layer is maintained by *H. salinarum* cells

within halite brine inclusions. Additionally, of the 20 membrane transporters shared between the

proteomes of liquid cultures and cells inside halite brine inclusions, four were found to be up-regulated

in halite brine inclusion extract, including the glycerol-3-phosphate-binding protein precursor.

Moreover, six proteins were found to be unique to the halite brine inclusion proteome including

phosphate, iron and glycerol-3-phosphate transporters. This suggests that transport capacity is not only

maintained but enhanced in cells during acclimation to halite brine inclusions, likely to maximize

extraction of essential molecules and trace elements from the local microenvironment.

Previous published transcriptomics and proteomics studies on *H. salinarum* cell shown impact

of salinity and dissolved oxygen concentrations on motility (Schmid *et al*., 2007; Leuko *et al*., 2009).

We therefore examined the expression of archaellum, gas vesicles and chemotaxis proteins. A high

degree of variability was noted in the expression of these proteins between halite brine inclusion

extracts and stationary growth phase cultures. Some proteins were common to both proteomes without

any differences in protein quantities, including for the gas vesicle protein (Gvp) subunits F, C, N of

cluster A, while GvpA was found only in two of the four liquid cultures and two of four halite samples.

GvpD and I subunits of A cluster and F of B cluster were found punctually in liquid samples. Only

GvpO cluster A and B seems to be impacted by crystallization with significant down-regulation in

halite brine inclusion extracts. For the archaellum proteins, only archaellin B1 was down-regulated in

halite samples whereas achaellin A1 and prearchaellin peptidase were identified punctually in liquid

culture samples. Archaellin B2, A2 and flagellin related H protein were found in all liquid culture

replicates but only in one halite replicate. An examination of chemotaxis proteins showed that all 18

identified proteins including CheR, CheA, CheW, CheC, CheB and mcp family proteins were shared

between halite brine inclusions extract and stationary growth phase liquid cultures.

These data show a general down-regulation of archaellin proteins involved in motility. A

reduced motility may be induced in halite brine inclusions to preserve energy. Gas vesicle down-

regulation suggests a lack of needed capacity for vertical movement in a water column in response to

oxygen levels, consistent with the restricted confines of a microscopic halite brine inclusions. Although

no swim or floating occur for brine inclusions, cells remain able to receive environmental signals due

to chemotaxis proteins founded both in stationary cells and brine extract without significant differential

expression.

6   Conclusion

The methods presented in this study expand on previous protocols for deactivation of halite surface-

attached cells removal of nucleic acids and allow for the first time for isolation of proteins directly

from halite brine inclusions that both excludes surface-bound contaminating proteins and avoids

changes in cell protein expression during salt dissolution. These methodologies were applied here to

provide the first sights into the molecular mechanisms involved in early acclamation of *H. salinarum*

within brine inclusions. While further functional investigations are needed to confirm the proteomics

hypotheses generated by the data in this study, the methodologies presented here could also have future

applications for the study of microorganisms in halite inclusions in natural settings.

7   Data Availability Statement

Proteomics data are available to the ProteomeXchange Consortium via the PRIDE partner repository

with the dataset identifier (*under submission*).

8   Author Contributions

Study was designed by CF, AK, AH and SZ under supervision of AK, SZ and AH. Experiments were

conducted by CF with the help of MM for crystals forming; AM and RP for mass spectrometry and

improvements of extraction methods; BAB for nanoLC-MS/MS data acquisition; FG for MEB-EDX

analyses; AT for RNA extraction and TD for plasma treatments. Proteomics data treatments were

conducted by AT and CF. Manuscript was written by CF and then revised and approved by all co-

authors.

9   Acknowledgments

This work was supported by Sorbonne Université, the X-life program of CNRS-MITI, the ATM

program of the Muséum National d'Histoire Naturelle and the ANR program "ExocubeHALO" (ANR-

21-CE49-0017-01_ACT). The UHPLC MS/MS data were acquired at the Plateau technique de

spectrométrie de masse bio-organique, UMR 7245 Molécules de Communication et d'Adaptation des

Microorganismes, Muséum national d'Histoire naturelle, Paris, France. The nanoLC-MS/MS data

were acquired at ProGenoMix MS-platform at CEA-Marcoule. The SEM-EDX images were acquired

at the Plateforme de microscopie électronique from IMPMC. Parts of the figure were drawn by using

pictures from Servier Medical Art. Servier Medical Art by Servier is licensed under a Creative

Commons Attribution 3.0 Unported License (https://creativecommons.org/licenses/by/3.0/).

10   Supplementary

The supplementary for this article are available online.

11   References

Bidle, K. A., Kirkland, P. A., Nannen, J. L., and Maupin-Furlow, J. A. (2008). Proteomic analysis of
Haloferax volcanii reveals salinity-mediated regulation of the stress response protein PspA. Microbiol
Read Engl 154, 1436–43. doi: 10.1099/mic.0.2007/015586-0.

Coker, J. A., DasSarma, P., Kumar, J., Müller, J. A., and DasSarma, S. (2007). Transcriptional profiling
of the model Archaeon *Halobacterium* sp. NRC-1: responses to changes in salinity and temperature.
Saline Syst 3, 6. doi: 10.1186/1746-1448-3-6.

Engel, M. B., and Catchpole, H. R. (2005). A microprobe analysis of inorganic elements in
Halobacterium salinarum. Cell Biol Int 29, 616–622. doi: 10.1016/j.cellbi.2005.03.024.

Falb, M., Müller, K., Königsmaier, L., Oberwinkler, T., Horn, P., Gronau, S. von, et al. (2008).
Metabolism of halophilic archaea. Extremophiles 12, 177–196. doi: 10.1007/s00792-008-0138-x.

Fendrihan, S., Dornmayr-Pfaffenhuemer, M., Gerbl, F. W., Holzinger, A., Grösbacher, M., Briza, P.,
et al. (2012). Spherical particles of halophilic archaea correlate with exposure to low water activity –
implications for microbial survival in fluid inclusions of ancient halite. Geobiology 10, 424–433. doi:
10.1111/j.1472-4669.2012.00337.x.

Gonzalez, O., Gronau, S., Pfeiffer, F., Mendoza, E., Zimmer, R., and Oesterhelt, D. (2009). Systems
Analysis of Bioenergetics and Growth of the Extreme Halophile *Halobacterium salinarum*. Plos
Comput Biol 5, e1000332. doi: 10.1371/journal.pcbi.1000332.

Gramain, A., Díaz, G. C., Demergasso, C., Lowenstein, T. K., and McGenity, T. J. (2011). Archaeal
diversity along a subterranean salt core from the Salar Grande (Chile). Environ Microbiol 13, 2105–
2121. doi: 10.1111/j.1462-2920.2011.02435.x.

Graur, D., and Pupko, T. (2001). The Permian Bacterium that Isn't.

Gruber, C., Legat, A., Pfaffenhuemer, M., Radax, C., Weidler, G., Busse, H.-J., et al. (2004).
Halobacterium noricense sp. nov., an archaeal isolate from a bore core of an alpine Permian salt
deposit, classification of *Halobacterium* sp. NRC-1 as a strain of *H. salinarum* and emended
description of *H. salinarum*. Extremophiles 8, 431–439. doi: 10.1007/s00792-004-0403-6.

Hartmann, R., Sickinger, H. D., and Oesterhelt, D. (1980). Anaerobic growth of halobacteria. Proc
National Acad Sci 77, 3821–3825. doi: 10.1073/pnas.77.7.3821.

Huby, T. J. C., Clark, D. R., McKew, B. A., and McGenity, T. J. (2020). Extremely halophilic archaeal
communities are resilient to short-term entombment in halite. Environ Microbiol. doi: 10.1111/1462-
2920.14913.

Judée, F., Simon, S., Bailly, C., and Dufour, T. (2018). Plasma-activation of tap water using DBD for
agronomy applications: Identification and quantification of long lifetime chemical species and
production/consumption mechanisms. Water Res 133, 47–59. doi: 10.1016/j.watres.2017.12.035.

Kanehisa, M., and Goto, S. (2000). KEGG: Kyoto Encyclopedia of Genes and Genomes. Nucleic Acids
Res 28, 27–30. doi: 10.1093/nar/28.1.27.

Kanehisa, M., Sato, Y., and Morishima, K. (2016). BlastKOALA and GhostKOALA: KEGG Tools for
Functional Characterization of Genome and Metagenome Sequences. J Mol Biol 428, 726–731. doi:
10.1016/j.jmb.2015.11.006.

Kirkland, P. A., Busby, J., Stevens, S., and Maupin-Furlow, J. A. (2006). Trizol-based method for
sample preparation and isoelectric focusing of halophilic proteins. Anal Biochem 351, 254–259. doi:
10.1016/j.ab.2006.01.017.

Kixmüller, D., and Greie, J. (2012). An ATP-driven potassium pump promotes long-term survival of
Halobacterium salinarum within salt crystals. Env Microbiol Rep 4, 234–241. doi: 10.1111/j.1758-
2229.2012.00326.x.

Leuko, S., Raftery, M. J., Burns, B. P., Walter, M. R., and Neilan, B. A. (2009). Global Protein-Level
Responses of *Halobacterium salinarum* NRC-1 to Prolonged Changes in External Sodium Chloride
Concentrations. J Proteome Res 8, 2218–2225. doi: 10.1021/pr800663c.

Maughan, H., Birky, W., Nicholson, W., Rosenzweig, W., and Vreeland, R. (2002). The paradox of
the "ancient" bacterium rich contains "modern" protein-coding genes.

Merino, N., Aronson, H. S., Bojanova, D. P., Feyhl-Buska, J., Wong, M. L., Zhang, S., et al. (2019).
Living at the Extremes: Extremophiles and the Limits of Life in a Planetary Context. Front
Microbiol 10, 780. doi: 10.3389/fmicb.2019.00780.

Moran-Reyna, A., and Coker, J. A. (2014). The effects of extremes of pH on the growth and
transcriptomic profiles of three haloarchaea. F1000research3, 168. doi:
10.12688/f1000research.4789.2.

Mormile, M. R., Biesen, M. A., Gutierrez, M. C., Ventosa, A., Pavlovich, J. B., Onstott, T. C., et al.
(2003). Isolation of *Halobacterium salinarum* retrieved directly from halite brine inclusions. Environ
Microbiol 5, 1094–1102. doi: 10.1046/j.1462-2920.2003.00509.x.

Müller, J. A., and DasSarma, S. (2005). Genomic Analysis of Anaerobic Respiration in the Archaeon
Halobacterium sp. Strain NRC-1: Dimethyl Sulfoxide and Trimethylamine N-Oxide as Terminal
Electron Acceptors. J Bacteriol 187, 1659–1667. doi: 10.1128/jb.187.5.1659-1667.2005.

Nickle, D. C., Learn, G. H., Rain, M. W., Mullins, J. I., and Mittler, J. E. (2002). Curiously Modern
DNA for a "250 Million-Year-Old" Bacterium. J Mol Evol 54, 134–137. doi: 10.1007/s00239-001-
0025-x.

Nims, C., Lafond, J., Alleon, J., Templeton, A. S., and Cosmidis, J. (2021). Organic biomorphs may
be better preserved than microorganisms in early Earth sediments. Geology 49, 629–634. doi:
10.1130/g48152.1.

Norton, C. F., and Grant, W. D. (1988). Survival of Halobacteria Within Fluid Inclusions in Salt
Crystals. Journal of General Microbiology, 1365–1373.

Oesterhelt, D., and Stoeckenius, W. (1974). Methods in Enzymology. Methods Enzymol 31, 667–678.
doi: 10.1016/0076-6879(74)31072-5.

Oren, A., Arahal, D. R., and Ventosa, A. (2009). Emended descriptions of genera of the family
Halobacteriaceae. Int J Syst Evol Micr 59, 637–642. doi: 10.1099/ijs.0.008904-0.

Oren, A. (2006). Life at high salt concentrations. The prokaryotes 3, 263–282. doi: 10.1007/0-387-
30742-7-9.

Payne, J. I., Sehgal, S. N., and Gibbons, N. E. (1960). Immersion Refractometry of some halophilic
bacteria. Can J Microbiol 6, 9–15. doi: 10.1139/m60-002.

Pinel-Cabello, M., Chapon, V., Ruiz-Fresneda, M. A., Alpha-Bazin, B., Berthomieu, C., Armengaud,
J., et al. (2021). Delineation of cellular stages and identification of key proteins for reduction and
biotransformation of Se(IV) by Stenotrophomonas bentonitica BII-R7. J Hazard Mater 418, 126150.
doi: 10.1016/j.jhazmat.2021.126150.

Podechard, N., Ducheix, S., Polizzi, A., Lasserre, F., Montagner, A., Legagneux, V., et al. (2018). Dual
extraction of mRNA and lipids from a single biological sample. Sci Rep-uk 8, 7019. doi:
10.1038/s41598-018-25332-9.

Post, F. J. (1977). The microbial ecology of the Great Salt Lake. Microbial Ecol 3, 143–165. doi:
10.1007/bf02010403.

Roedder, E. (1984). The fluids in salt. American Mineralogist 69, 413–439.

Rosenzweig, W. D., Peterson, J., Woish, J., and Vreeland, R. H. (2000). Development of a Protocol to
Retrieve Microorganisms from Ancient Salt Crystals. Geomicrobiol J 17, 185–192. doi:
10.1080/01490450050121152.

Sankaranarayanan, K., Timofeeff, M. N., Spathis, R., Lowenstein, T. K., and Lum, J. K. (2011).
Ancient Microbes from Halite Fluid Inclusions: Optimized Surface Sterilization and DNA
Extraction. Plos One 6, e20683. doi: 10.1371/journal.pone.0020683.

Schmid, A. K., Reiss, D. J., Pan, M., Koide, T., and Baliga, N. S. (2009). A single transcription factor
regulates evolutionarily diverse but functionally linked metabolic pathways in response to nutrient
availability. Mol Syst Biol 5, 282. doi: 10.1038/msb.2009.40.

Schreder-Gomes, S. I., Benison, K. C., and Bernau, J. A. (2022). 830-million-year-old microorganisms
in primary fluid inclusions in halite. Geology. doi: 10.1130/g49957.1.

Shimizu, S., Barczyk, S., Rettberg, P., Shimizu, T., Klaempfl, T., Zimmermann, J. L., et al. (2014).
Cold atmospheric plasma – A new technology for spacecraft component decontamination. Planet Space
Sci 90, 60–71. doi: 10.1016/j.pss.2013.10.008.

Tebbe, A., Schmidt, A., Konstantinidis, K., Falb, M., Bisle, B., Klein, C., et al. (2009). Life-style
changes of a halophilic archaeon analyzed by quantitative proteomics. Proteomics 9, 3843–3855. doi:
10.1002/pmic.200800944.

Vreeland, R. H., Rosenzweig, W. D., and Powers, D. W. (2000). Isolation of a 250 million-year-old
halotolerant bacterium from a primary salt crystal. Nature 407, 897–900. doi: 10.1038/35038060.

Winters, Y. D., Lowenstein, T. K., and Timofeeff, M. N. (2015). Starvation-Survival in
Haloarchaea. Life 5, 1587–1609. doi: 10.3390/life5041587.

12   Table and Figures

Table 1: Experimental parameters of cold plasma sources used to remove surface-bound proteins. The
evaluation is based of proteolysis results. DBD means dielectric barrier device; APPJ for atmospheric
pressure plasma jet. Slm stands for standard liters per minute and sccm standard centimeters cube per
minute.

Plasma source	Excitation mode	Electrical parameters			Gas and flow rate	Proteolysis
		Ampl.	Freq.	Duty cycle		
DBD	Alternative Current	7 kV	500 Hz	-	He (1 slm)	0.0 %
					He-O_2 (1slm - 150 sccm)	0.0 %
					He-O_2 (1slm - 3 sccm)	0.0 %
					Ar (1 slm)	0.0 %
APPJ	Alternative Current	7 kV	700 Hz	-	He (6 slm)	0.0 %
	Pulse	7 kV	700 Hz	1 %	He (6 slm)	29.3 %
					He-O_2 (6 slm - 100 sccm)	39.1 %

835 **Table 2**: Targeted proteins involved in early acclimation of *Halobacterium salinarum* through halite
836 brine inclusion (for extended matched proteins, see Supplementary Table 6-5). Shared mean protein
837 found for all the fourth brine extract and the fourth liquids stationary extracts.

Central metabolism - glycolysis/gluconeogenesis

		Samples
AAG20336.1	phosphohexomutase (phosphoglucomutase /phosphomannomutase)	Shared
AAG20164.1	glucose-6-phosphate isomerase	Shared
AAG19177.1	fructose-bisphosphatase	Shared
AAG19176.1	class I fructose-bisphosphate aldolase	Shared
AAG19439.1	triosephosphate isomerase	Shared
AAG18725.1	glyceraldehyde 3-phosphate dehydrogenase	Shared
AAG18584.1	phosphoglycerate kinase <=> glucose-6-phosphate isomerase	Shared
AAG20083.1	phosphoglycerate mutase	Shared
AAG19526.1	phosphopyruvate hydratase	Shared
AAG18902.1	pyruvate kinase	Shared
AAG18906.1	phosphoenolpyruvate synthase	Shared

Central metabolism - tricarboxylic acid (TCA) cycle

AAG20246.1	citrate synthase	Shared
AAG20620.1	aconitase	Shared
AAG20073.1	isocitrate dehydrogenase NADP	Shared
DAC78228.1	pyruvate--ferredoxin oxidoreductase alphasubunit	Shared
AAG19513.1	2-ketoglutarate ferredoxin oxidoreductase	Shared
AAG19826.1	succinate-CoA synthetase beta chain	Shared
AAG19827.1	succinyl-CoA synthetase alpha chain	Shared
AAG19651.1	succinate dehydrogenase subunit A	Shared
AAG19652.1	succinate dehydrogenase subunit B	Shared
AAG19688.1	fumarate hydratase	Shared
AAG20463.1	L-malate dehydrogenase	Shared

Central metabolism - acetyl-CoA metabolism

AAG19414.1	acetyl-CoA synthetase	Shared
AAG19020.1	acetyl-CoA synthetase	Shared
DAC78075.1	acetate--CoA ligase	Shared
AAG19243.1	aldehyde dehydrogenase (retinol)	Shared

Arginine deiminase pathway

AAG20948.1	arginine deiminase (plasmid)	Up-regulated
AAG20946.1	ornithine carbamoyltransferase (plasmid)	Shared
AAG20947.1	carbamate kinase (plasmid)	Shared
AAG20518.1	argininosuccinate lyase	Shared
AAG20519.1	argininosuccinate synthetase	Shared
AAG20945.1	Arginine/ornithine antiporter ArcD	Shared
AAG20949.1	IclR family transcription regulator ArcR (plasmid)	Shared

Aerobic respiration

AAG19138.1	NADH dehydrogenase/oxidoreductase	Shared
AAG19139.1	NADH dehydrogenase/oxidoreductase	Shared
AAG19141.1	NADH dehydrogenase/oxidoreductase-like protein	Shared
AAG19331.1	NADH dehydrogenase	Shared
AAG19651.1	succinate dehydrogenase subunit A	Shared
AAG19652.1	succinate dehydrogenase subunit B	Shared
AAG20249.1	succinate dehydrogenase subunit	Shared
AAG20277.1	H+-transporting ATP synthase subunit A	Shared
AAG20276.1	H+-transporting ATP synthase subunit B	Shared
AAG20279.1	H+-transporting ATP synthase subunit C	Shared
AAG20273.1	H+-transporting ATP synthase subunit D	Shared
AAG20278.1	H+-transporting ATP synthase subunit E	Shared
AAG20283.1	ATP synthase archaeal subunit H	Shared
AAG20282.1	H+-transporting ATP synthase subunit I	Shared
AAG20278.1	H+-transporting ATP synthase subunit F	Lacked in brine inclusion 4

Bacteriorhodopsin-mediated phototrophy

AAG18778.1	bacterio-opsin activator-like protein	Shared
AAG19769.1	bacterio-opsin activator	Specific brine inclusion 1, 4
AAG19313.1	sensory rhodopsin I transducer; Htr1 transducer	Shared
AAG19989.1	sensory rhodopsin II transducer; Htr2 transducer	Shared

Cell division and chromosome partitioning associated

AAG20773.1	chromosome partitioning protein ParA	Shared
AAG18937.1	cell division protein FtsZ type I	Shared
AAG18804.1	cell division protein FtsZ type II	Shared
AAG19919.1	Cell division cycle protein; AAA-type ATPase (CDC48 subfamily)	Shared
AAG19776.1	Cell division cycle protein; AAA-type ATPase (CDC48 subfamily)	Shared
AAG19516.1	Mrp protein homolog	Shared
AAG20269.1	cell division inhibitor MinD/ParA family protein	Shared
AAG20524.1	DNA starvation/stationary phase protection protein; ferritin DspA	Shared
AAG18755.1	archaeal histone A1	Shared
AAG20951.1	chromosome segregation protein SMC	Shared

DNA replication and repair

AAG19883.1	replication factor C large subunit	Shared
AAG20394.1	replication factor C small subunit	Shared
AAG20005.1	DNA primase DnaG	Shared
AAG19049.1	DNA-directed DNA polymerase B	Shared
AAG19329.1	DNA gyrase subunit A	Shared
AAG19326.1	DNA topoisomerase VI subunit A	Shared
AAG19327.1	DNA topoisomerase VI subunit B	Shared
DAC77610.1	DNA topoisomerase I	Shared
AAG20376.1	DNA polymerase sliding clamp	Shared
AAG20387.1	Orc1-type DNA replication protein	Shared
AAG19323.1	ATP-dependent DNA ligase	Shared
AAG19328.1	DNA gyrase subunit B	Lacked in brine inclusion 4
AAG18851.1	ribonuclease H type I	All stationary and brine inclusion 2
AAG20362.1	replication factor C small subunit	Lacked in brine inclusion 1
AAG20498.1	Orc1-type DNA replication protein	All stationary and brine inclusion 1
AAG19899.1	ribonucleoside reductase large chain	Shared
AAG19981.1	DNA photolyase family protein	Shared
AAG19671.1	DNA photolyase family protein	Shared
AAG19045.1	DNA double-strand break repair Rad50 ATPase	Shared
AAG20657.1	repair helicase UvrD	Shared
AAG20669.1	excinuclease ABC subunit UvrA	Shared
AAG20465.1	ATP-dependent DNA helicase Hel308	Shared
DAC79307.1	DNA repair and recombination protein RadA	Shared
AAG19690.1	DNA repair protein; flap endonuclease Fen1	Shared
DAC78445.1	DNA repair helicase Rad25	Shared
AAG20482.1	excinuclease ABC subunit UvrB	Lacked in stationary extract 1
AAG20386.1	DNA mismatch repair protein MutS	Lacked in stationary extract 2
AAG19103.1	endonuclease III	Lacked in stationary extract 1
AAG20477.1	type I site-specific deoxyribonuclease subunit RmeR	All stationary and brine extract 1, 3
AAG18797.1	endonuclease IV	All stationary extract
AAG19328.1	DNA gyrase subunit B	Lacked in brine inclusion 4
AAG18777.1	DNA mismatch repair protein MutL	Lacked in brine inclusion 1
AAG19222.1	DNA N-glycosylase	Lacked in brine inclusion 2
AAG20439.1	single-stranded-DNA-specific exonuclease RecJ1	Lacked in brine inclusion 2

Transcriptional machinery and factors

AAG20691.1	DNA-directed RNA polymerase subunit C	Shared
AAG20693.1	DNA-directed RNA polymerase subunit B'	Shared
AAG20694.1	DNA-directed RNA polymerase subunit B''	Shared
AAG19520.1	DNA-directed RNA polymerase subunit A'	Shared
AAG20029.1	DNA-directed RNA polymerase subunit E	Shared
AAG20743.1	transcription initiation factor IID (plasmid); TATA-box-binding protein B	Shared
AAG20365.1	transcription initiation factor IID; TATA-box-binding protein E	Shared
AAG19546.1	transcription initiation factor TFE alpha subunit	Shared
DAC77935.1	NusA-like transcription termination signal-binding factor	Lacked in brine inclusion 4
AAG19546.1	DNA-directed RNA polymerase subunit F	Shared
AAG20695.1	DNA-directed RNA polymerase subunit H	Down-regulated
AAG19524.1	DNA-directed RNA polymerase subunit N	All stationary extracts
AAG18850.1	transcription initiation factor IIB	Lacked in brine inclusion 1

Transfert RNA biogenesis

AAG20361.1	ribonuclease Z	Shared
AAG20359.1	tyrosine-tRNA ligase	Shared
AAG20355.1	tryptophanyl-tRNA ligase	Shared
AAG20349.1	leucine-tRNA ligase	Shared
AAG20324.1	isoleucine-tRNA ligase	Shared
AAG20397.1	alanine-tRNA ligase	Shared
AAG20602.1	Valine-tRNA ligase	Shared
AAG18903.1	methionine-tRNA ligase	Shared
AAG20222.1	serine-tRNA ligase	Shared
AAG20454.1	glycine-tRNA ligase	Shared
DAC77651.1	proline-tRNA ligase	Shared
AAG19494.1	cysteine-tRNA ligase	Shared
AAG19534.1	glutamate-tRNA ligase	Shared
AAG20944.1	arginine-tRNA ligase	Shared
AAG20567.1	phenylalanine-tRNA ligase alpha chain	Shared
AAG20568.1	phenylalanine-tRNA ligase beta chain	Shared
AAG20175.1	histidine-tRNA ligase	Shared
AAG20183.1	lysine-tRNA ligase	Shared
AAG19001.1	nondiscriminating Aspartyl—tRNA(Asp/Asn) ligase	Shared
AAG20252.1	tRNA(Ile2) 2-agmatinylcytidine synthetase TiaS	Shared
AAG18757.1	tRNA adenylyltransferase CCA-adding	Shared
AAG20422.1	L-threonylcarbamoyladenylate synthase	Shared
AAG20156.1	tRNA pseudouridine synthase Pus10	Shared
AAG20680.1	ribonuclease R	Shared
AAG19760.1	tRNA-splicing ligase RtcB	Shared
AAG19031.1	tRNA/rRNA cytosine-C5-methylase	Shared
AAG20338.1	tryptophan-tRNA ligase	All stationary extracts
AAG20043.1	threonine-tRNA ligase	All stationary extracts
AAG20339.1	tRNA-intron endonuclease	All stationary extracts
AAG19436.1	PtpS family protein	All stationary extracts
AAG18873.1	L-isoaspartyl protein carboxyl methyltransferase	All stationary extracts
AAG20139.1	archaeosine tRNA-ribosyltransferase	All stationary extracts

Ribosome biogenesis

AAG20650.1	ribosome biogenesis/translation initiation ATPase RLI	Shared
AAG20352.1	probable ribosome biogenesis protein Dim2	Shared
AAG19629.1	ribosome maturation protein Sdo1	Shared
DAC78346.1	ribosome-associating GTPase HflX	Shared
AAG19036.1	GTP-binding protein	Shared
AAG19552.1	fibrillarin-like rRNA/tRNA 2'-O-methyltransferase	Shared
AAG19551.1	rRNA/tRNA 2'-O-methyltransferase complex proteinNop5	Shared
AAG20422.1	L-threonylcarbamoyladenylate synthase	Shared
AAG19112.1	ribonucleoside-diphosphate reductase beta subunit	Shared
AAG20662.1	probable ribonuclease FAU-1	Shared

Translational machinery and factors

AAG19500.1	50S ribosomal protein L1P	Shared
AAG19501.1	50S ribosomal protein L11P	Shared
AAG19946.1	50S ribosomal protein L4P	Shared
AAG18792.1	50S ribosomal protein L15E	Shared
AAG19941.1	50S ribosomal protein L22P	Shared
AAG19938.1	50S ribosomal protein L23P	Shared
AAG19943.1	50S ribosomal protein L29P	Shared
AAG19936.1	50S ribosomal protein L13P	Shared
AAG19952.1	50S ribosomal protein L6P	Shared
AAG20681.1	30S ribosomal protein S10P	Shared
AAG19256.1	30S ribosomal protein S15P	Shared
AAG19945.1	30S ribosomal protein S17P	Shared
AAG20178.1	30S ribosomal protein S19E	Shared
AAG20206.1	30S ribosomal protein S24E	Shared
AAG19442.1	30S ribosomal protein S3P	Shared
AAG19253.1	30S ribosomal protein S3E	Shared
AAG19956.1	30S ribosomal protein S5P	Shared
AAG20576.1	30S ribosomal protein S6E	Shared
AAG20688.1	30S ribosomal protein S7P	Shared
AAG19951.1	30S ribosomal protein S8P	Shared
AAG19499.1	50S ribosomal protein L10P	Down-regulated
AAG18728.1	50S ribosomal protein L10E	Down-regulated
AAG19522.1	50S ribosomal protein L13P	Down-regulated
AAG19521.1	50S ribosomal protein L18E	Down-regulated
AAG19954.1	50S ribosomal protein L19E	Down-regulated
AAG19939.1	50S ribosomal protein L2P	Down-regulated
AAG19947.1	50S ribosomal protein L24P	Down-regulated
AAG19957.1	50S ribosomal protein L30P	Down-regulated
AAG19953.1	50S ribosomal protein L32E	Down-regulated
AAG19937.1	50S ribosomal protein L4E	Down-regulated
AAG19949.1	50S ribosomal protein L5P	Down-regulated
AAG19537.1	50S ribosomal protein L7Ae	Down-regulated
AAG19519.1	30S ribosomal protein S11P	Down-regulated
AAG20689.1	30S ribosomal protein S12P	Down-regulated
AAG19517.1	30S ribosomal protein S13P	Down-regulated
AAG19950.1	30S ribosomal protein S14P	Down-regulated
AAG19940.1	30S ribosomal protein S19P	Down-regulated
AAG19527.1	30S ribosomal protein S2P	Down-regulated
AAG19518.1	30S ribosomal protein S4P	Down-regulated
AAG19948.1	30S ribosomal protein S4E	Down-regulated
AAG19920.1	30S ribosomal protein S8E	Down-regulated
DAC78237.1	30S ribosomal protein S9	Down-regulated
AAG19070.1	translation initiation factor eIF-2 subunit alpha	Shared
AAG19619.1	translation initiation factor eIF-2 subunit beta	Shared
AAG20211.1	translation initiation factor eIF-2 subunit gamma	Shared
AAG19992.1	translation initiation factor eIF-5A	Shared
AAG19758.1	translation initiation factor eIF-2B subunit delta	Shared
AAG20682.1	translation elongation factor eEF-1A subunit alpha	Shared
AAG19926.1	homolog to translation elongation factor aEF-1alpha subunit	Shared
DAC79456.1	translation elongation factor aEF-2	Shared
AAG20628.1	translation initiation factor aIF-1	Down-regulated
AAG20168.1	translation initiation factor aIF-5B	Up-regulated

Chaperonnes and folding catalysts

AAG20641.1	thioredoxin	Shared
AAG19025.1	molecular chaperone DnaJ	Shared
AAG19026.1	Hsp70-type molecular chaperone DnaK	Shared
AAG19028.1	DnaJ/DnaK ATPase stimulator GrpE	Shared
AAG19995.1	Hsp20-type molecular chaperone	Shared
AAG18869.1	Hsp20-type molecular chaperone	Shared
AAG20240.1	thermosome subunit beta	Shared
AAG20539.1	prefoldin alpha subunit	Shared
AAG18836.1	prefoldin subunit beta	Down-regulated

Surface proteinaceous layer

AAG20702.1	cell surface glycoprotein	Shared

Transporters

AAG18929.1	arsenical pump-driving ATPase ArsA	Shared
AAG19127.1	propionyl-CoA carboxylase carboxyltransferasecomponent	Shared
AAG20473.1	copper transport ATP-binding protein	Shared
AAG19180.1	ABC transporter ATP-binding protein	Shared
AAG19353.1	spermidine/putrescine ABC transporter ATP-binding	Shared
AAG20071.1	spermidine/putrescine ABC transporter	Shared
AAG20612.1	ABC-type transport system periplasmic substrate-binding protein	Shared
AAG20603.1	ABC-type transport system periplasmic substrate-binding protein	Shared
AAG20587.1	dipeptide ABC transporter ATP-binding	Shared
AAG20588.1	dipeptide ABC transporter ATP-binding	Shared
AAG20852.1	dipeptide ABC transporter dipeptide-binding	Shared
AAG20449.1	oligopeptide ABC transporter	Shared
AAG20448.1	oligopeptide ABC transporter ATP-binding	Shared
AAG20457.1	oligopeptide binding protein	Shared
AAG18995.1	phosphate import ATP-binding protein PstB 1	Shared
AAG20556.1	phosphate ABC transporter binding	Shared
AAG18998.1	phosphate ABC transporter substrate-binding protein PstS family	Shared
AAG20233.1	phosphate/phosphite/phosphonate ABC transporter substrate-binding protein	Shared
AAG20486.1	thiamine ABC transporter substrate-binding protein	Shared
AAG20882.1	TrkA family potassium uptake protein	Shared
AAG20878.1	TrkA family potassium uptake protein	Shared
DAC78851.1	TrkA family potassium uptake protein	Shared
AAG20419.1	hemolysin family protein	Shared
AAG19086.1	Mrp-type sodium/proton antiporter system subunit	Shared
AAG18746.1	ABC-type transport system ATP-binding/permease protein	Shared
AAG19338.1	BMP family protein	Shared
DAC77962.1	chemotactic signal transduction system periplasmic substrate-binding protein; BMP family protein	Shared
AAG19051.1	Fe-S cluster assembly ATPase SufC	Shared
AAG20945.1	arginine/ornithine antiporter ArcD	Shared
DAC77887.1	ABC-type transport system permease protein	Lacked in stationary 2, 3
AAG19355.1	iron ABC transporter substrate-binding protein	All brine inclusion
AAG20553.1	phosphate import ATP-binding protein PstB 2	All brine inclusion
AAG20937.1	hemolysin family protein	All brine inclusion
AAG20918.1	glycerol-3-phosphate-binding protein precursor (plasmid)	All brine inclusion
AAG20070.1	PotD/PotF family extracellular solute-binding protein	All brine inclusion
AAG19271.1	spermidine/putrescine-binding protein	All brine inclusion
AAG20451.1	ABC transporter permease	All brine inclusion extracts
AAG20450.1	dipeptide ABC transporter permease	All stationary extracts
AAG19633.1	TrkA family potassium uptake protein	All stationary extracts
AAG18790.1	TrkA family potassium uptake protein	All stationary extracts
AAG19697.1	iron (III) ABC transporter ATP-binding / VitB12-binding protein precursor	All stationary extracts

Archaellum proteins

AAG19424.1	flagellin A1 precursor	Stationary extracts 1, 2
AAG19384.1	flagellin B2 precursor	Lacked in brine inclusion 1, 2, 4
AAG19383.1	flagellin B1 precursor	Down-regulated
AAG19425.1	flagellin A2 precursor	Lacked in brine inclusion 1, 2, 3
AAG19380.1	flagella-related protein E	Shared
AAG19377.1	flagella-related protein H	Stationary extract 3; brine inclusions 3, 4
DAC79152.1	prepilin/prearchaelin peptidase	Stationary extracts 1, 3, 4

Gas vesicle proteins

AAG20723.1	Gas vesicle protein GvpF cluster A (plasmid)	Shared
AAG20727.1	Gas vesicle protein GvpC cluster A (plasmid)	Shared
AAG20898.1	Gas vesicle protein GvpO cluster A (plasmid)	Down-regulated
AAG20729.1	Gas vesicle protein GvpO cluster B (plasmid)	Down-regulated
AAG20721.1	Gas vesicle protein GvpH cluster A (plasmid)	All stationary extracts
AAG20728.1	Gas vesicle protein GvpN cluster A (plasmid)	Shared
AAG20717.1	Gas vesicle protein GvpL cluster A (plasmid)	Stationary extracts 1, 2, 4
AAG20725.1	Gas vesicle protein GvpD cluster A (plasmid)	Stationary extract 1
AAG20726.1	Gas vesicle protein GvpA cluster A (plasmid)	Stationary extract 2, 3; brine inclusions 1, 2
AAG20720.1	Gas vesicle protein GvpI cluster A (plasmid)	Stationary extract 2
AAG20892.1	Gas vesicle protein GvpF cluster B (plasmid)	Stationary extract 3, 4

Chemotaxis proteins

AAG19389.1	chemotaxis protein, protein-glutamate O-methyltransferase CheR	Shared
AAG19394.1	chemotaxis protein, protein-glutamate methylesterase /protein-glutamine glutaminase CheB	Shared
AAG19393.1	taxis sensor histidine kinase CheA	Shared
AAG19396.1	purine-binding taxis protein CheW	Shared
AAG19371.1	purine-binding taxis protein CheW	Shared
AAG19392.1	chemotaxis protein CheC	Shared
AAG19399.1	taxis cluster protein CheC	Shared
AAG19986.1	transducer protein CosT; Htr5 transducer	Shared
AAG19812.1	methyl-accepting chemotaxis protein; Htr8 transducer	Shared
AAG19257.1	methyl-accepting chemotaxis protein; Htr13	Shared
AAG19381.1	Transducer protein Htr15	Shared
AAG19270.1	transducer protein Htr18	Shared
AAG19266.1	methyl-accepting chemotaxis protein; Htr4 transducer	Shared
AAG19257.1	methyl-accepting chemotaxis protein; Htr6 transducer	Shared
AAG18922.1	transducer protein MpcT; Htr14 transducer	Shared
DAC78795.1	transducer protein BasT	Shared

839 **Table 3**: Total purified RNA quality check results after TRIzol Reagent™ -based extraction. [a]See
840 Supplementary Figure 6-2 for BioAnalyzer results. Purity score below 2.0 suggest purity unsuitable
841 for RNAseq analyses.

	Liquid stationary cells			Internally inoculated halite (2 months)			Externally inoculated halite (~1 week)		
RNA quantity (µg)	1040.0	1045.2	1190.8	<20	4.4	13.9	327.6	444.0	129.6
Purity (absorbance 260/280nm)	2.10	2.09	2.08	1.95	1.54	1.61	1.99	2.03	1.98
Purity (absorbance 260/230nm)	2.44	2.38	2.47	4.67	0.96	1.44	1.89	1.78	2.05
Quality[a]	+	++	++	--	--	--	-	-	-

844 **Figure 1**: TNPA laboratory-grown halite crystals internally inoculated with *Halobacterium salinarum*
845 cells observations after two months incubated at 37 °C. a) crystallization replicate 1, b) crystallization
846 replicate 2 and c) crystallization replicate 3.

Figure 2: Laboratory-grown halite surface observation. a-f) TNPA crystals without cells. g-j) Internally inoculated TNPA crystals with *Halobacterium salinarum* cells incubated two months at 37 °C. k-n) Externally inoculated TNPA crystals with *Halobacterium salinarum* cells after approximatively one week incubated at 37 °C. a, g, k) Schematic representation of halite crystals without cells, internally inoculated and externally inoculated respectively. b-h-l) Visual observation of crystals without cells, internally inoculated and externally inoculated respectively. c, e, i, m, d, f, j, n) SEM observations and elemental composition analysis by EDX.

856 **Figure 3**: Surface-bounded proteins removal by chemical treatments with active-spray washes. a)
857 Percentage of proteins lysis after NaOCl-NaOH treatment with or without additional HCl step and
858 subsequent residual protein quantity. b) Overview of optimal NaOCl-NaOH-HCl chemical treatment
859 with active-spray washes.

Molecular Acclimation to Halite Brine Inclusions

Figure 4: Venn diagrams of proteins identified by mass spectrometry for brine inclusions and liquid stationary samples. a) Comparison of biological replicates for liquid stationary extracts. b) Comparison of biological replicates for brine inclusion extracts. Black circles in a) and b) represent "core" proteins shared by the four replicates in both cases. c) Comparison of core proteins between brine inclusion extract and liquid stationary cells.

868   **Figure 5**: Venn diagram of the 68 proteins quantified by iTRAQ® with triple injection and exhibit
869   significative differential expression in brine extracts compared to liquid stationary cultures.