Female chacma baboons modulate their sexual receptivity in response to male coercion

Darmis Fragkiskos[1], Huchard Élise[2], Cowlishaw Guy[3] & Carter J. Alecia[1]

[1]Department of Anthropology, University College London, London, UK

[2]Institut des Sciences de L'Evolution de Montpellier, UMR 5554, CNRS, Université de Montpellier, Montpellier, France

[3]Institute of Zoology, Zoological Society of London, London, UK

Research in social mammals has revealed the complexity of female counter-strategies to
reproductive competition and sexual conflict. For example, comparative research has shown
that the length of female sexual receptivity varies with infanticide risk, but whether individuals
can strategically adjust their period of receptivity from cycle to cycle remains unknown. This
study addresses this gap by exploring whether wild female chacma baboons (*Papio ursinus*)
modulate their receptivity in response to the social environment. Given that female chacma
baboons (a) compete for paternal care and (b) that infanticide risk and coercion are high, we
predicted that: females could (a) shorten their receptive period to reduce intrasexual aggression
and (b) male coercion, or (c) increase their conceptive period to access multiple or their
preferred male. We quantified 158 receptivity cycles from 47 females recorded over 15 years.
Females' swelling duration, but not maximal-swelling, had low repeatability between females,
but most variation in both phases stemmed from within females. We found evidence that
females decrease their oestrous duration in response to an increasing number of in-group males,
possibly to decrease their exposure to sexual coercion. We thus present preliminary evidence
for an unexplored mechanism under sexual selection: female cycle length manipulation.
25
26 Keywords: cycle length manipulation, sexual selection, sexual coercion, social environment,
27 female competition

Introduction

Recent advances in sexual selection theory underline the importance of both mate choice and intrasexual female competition in generating variance in female reproductive success [1–4]. In the first case, it may be advantageous for females to concentrate paternity into a single male who can provide benefits in the form of good genes or of paternal services such as protection to her or her offspring. However, in group-living species, a number of social pressures may decrease the possibility for females to express their mating preference. For example, recent studies highlight the role of sexual conflict [5], which frequently manifests via infanticide in mammals. Indeed, infanticide risk may push females to mate with multiple rather than single partners, in order to dilute paternity among multiple males who subsequently restrain from attacking an offspring that they could have sired [6,7]. Female choice may be further constrained by sexual coercion, when males direct aggression to females [3,5,8], either to increase the probability that they will mate with them or to decrease the probability that they will mate promiscuously [9]. In such cases, females will mate with their aggressors if resisting is more costly than giving in (i.e., convenience polyandry) [5]. In groups of water striders (*Aquarius remigis*), aggressive males acquire more matings than less aggressive ones [10]. In primates, male baboons (*Papio ursinus*) that attack pre-ovulatory females are more likely to monopolize them when receptive [11] and females do not copulate close to adult males to avoid bystanders' aggression [11,12]. Also, female chimpanzees (*Pan troglodytes*) initiate copulations most frequently with those males that were most aggressive towards them throughout their cycle [13,14] and male lemurs (*Lemur catta*) that are not preferred can force copulations [15]. However, male coercion can lead to female counterstrategies and antagonistic co-evolution between the sexes [5] since it is associated to multiple costs for females: for example when they are forced to mate with subordinate or incompatible males, when coercion manifested as direct aggression/harassment increases stress hormone secretion [14], when it leads caring males to reduce parental care [8] or when it impairs female choice [16].

Females also face intrasexual competition over access to males and reproductive resources [17–20]. This competition for access to males increases when there is synchrony of females' fertility (i.e., ovarian synchrony), which creates a female-biased operational sex ratio (OSR) and increases aggression received by receptive females [21,22]. In general, females compete with consexuals for (a) their preferred mates (e.g., chacma baboons: [21]; red-fronted lemurs: [23]), for (b) paternal services for their offspring (e.g., chacma baboons: [24,25]) and (c) other reproductive resources [26], such as breeding territories (e.g., lek center [27]) or food

resources [28]. Those benefits are achieved either through monopolisation of resources directly

from females (i.e., contest competition for food) [29] or indirectly (e.g., nutrient-rich

spermatophores in insects:[30]; low predation risk in lek centers [27]). Antagonism for (a), (b)

and (c) can manifest through non-aggressive interactions (e.g., signals: [30]) and escalate up to

physical aggression, which can be detrimental to the receiver's short- or long-term fitness [31].

When facing coercion and intrasexual competition, females may use cycle

manipulation by adjusting the length of the receptive period of the menstrual cycle (cycle

length manipulation, hereafter CLM) as one counterstrategy. Such a mechanism could be under

selection as it can potentially decrease the rate of aggression females receive from (1)

intrasexual competition and (2) sexual coercion, or (3) alternatively it might increase females'

probability of mating with their preferred or multiple males (increase female mate choice).

Theoretical and experimental evidence for cycle manipulation exists [32,33], with studies

detecting physiological changes in female cycling patterns (e.g., changes to cycle length,

resumption of cycling) in response to the social environment. Socially-stressed primate

females, for example by being harassed near menstruation, have longer menstrual cycles [34].

Cycling female yellow baboons that receive aggression from consexuals tend to have more

cycles to conception and longer interbirth intervals [31], while pregnant females that are

systematically harassed experience reduced fitness (through non-normal gestation length,

spontaneous abortion, and premature delivery) [31]. Psychosocial stress stemming from social

instability, in the form of male take-overs, induces female physiological responses too: the

presence of unfamiliar males can mediate early sexual maturation [35,36] or abortion [37] in

female mammals, in a likely effort to limit infanticide risk. Novel males in geladas prompt

females of any state (immature, cycling, lactating, or pregnant) to start or resume cycling

[36,38], while female hamadryas with young infants develop swellings to advertise a reduced

postpartum amenorrhea that results in copulations with novel males during take-overs [33].

However, these swellings are deceptive signals that ultimately decrease infanticide risk (four

out of five infants survived in [33]) since females did not conceive during the first cycles after

the take-over, their reproduction was not accelerated and the interbirth intervals were not

shorter compared to times with no male take-overs. Despite the above results though, no study

to date has specifically proposed CLM, or tested for it, as a further potential mechanism under

selection.

In this study, we explored whether females adjust the duration of their oestrus in

response to the social environment to either reduce aggression from (1) coercive males or (2)

other females or to increase their access to (3) their preferred or (4) multiple males. We focused

on female chacma baboons because they exhibit elevated female-female competition,

compared to other primates [25,39,40], live in multimale-multifemale groups, mate

promiscuously [11,41], and have exaggerated swellings that reliably indicate females' stage of

the follicular phase of the oestrus cycle [32]. Female chacma baboons are an ideal model to

study female intra- and intersexual competition, as well as sexual coercion, since they target

the dominant male as a sexual partner [25,40], but also mate promiscuously [21], outside their

peak swelling, to establish more males with a nonzero paternity probability or to initiate male-

male competition (e.g., sperm competition) [12,32]. In addition, male's monopolisation of only

one receptive female at a time (i.e., mate-guarding: [19,24]) results in female-female

competition over sexual access to particular males [21,25], who might be better able to protect

them from aggressive consexuals or provide future postpartum care for the offspring [42].

To address our overarching aim, we proceeded in two steps. First, we explored the

variation within and among females in the durations of swelling and maximal-swelling stages

of the cycle. We aimed to quantify the repeatability of females' receptivity phases to determine

their flexibility, predicting large variation within females. Second, we tested whether

intersexual or intrasexual selection, or sexual conflict in the form of male coercion, could

explain the remaining variation in the duration of specific cycle phases (summarised in Table

1). First, considering female intrasexual selection, we tested whether female competition for

paternal care explains variation in the duration of receptivity (the avoidance of female

aggression hypothesis [H1]). Pregnant and lactating females aggressively target swollen ones

to decrease competition for paternal care of their own offspring, or to induce reproductive

suppression so that their own infants are not born at the same time as many others [43]. For

this reason, we predicted that more in-group pregnant and lactating females result in higher

stress levels in cycling females, which in turn leads receptive females to deflate quickly. This

response should, ultimately, decrease their exposure to consexual aggression. Our second

hypothesis deals with intersexual selection and sexual conflict (H2): we hypothesized that

females would manipulate their swelling and maximal-swelling duration to access the

dominant mate (the paternity concentration hypothesis, H2a). We predicted that, when

competition for access to the alpha male was likely because there were simultaneously-swollen

individuals, females would delay ovulation by elongating their swelling or maximal-swelling

duration to increase the probability of mating with this male. By doing so females may benefit

from (a) good genes [25,32] and from (b) concentrating paternity certainty in their preferred

male, usually the dominant, who is also the most likely to pose a future risk of infanticide if

they fail to mate with him [25,44]. We also tested whether the risk of infanticide from multiple

males explains patterns of variation in female baboons' cycle-phases (the paternity confusion

hypothesis, H2b). We predicted that females would increase their swelling or maximal-

swelling length as the operational sex ratio (OSR) becomes female-skewed—and female-

female mating competition increases accordingly—so they still manage to mate with many

males and reduce infanticide risk. Finally, we tested whether the effect of social instability in

the form of high numbers of in-group males (i.e., strong male-male competition) explains

patterns of variation in a female's receptivity phases. We predicted that a male-skewed OSR

or a high number of in-group males induces high male-male competition, which translates into

an increased risk of sexual coercion for receptive females [11,24]. Females may thus shorten

their oestrous period to escape or minimize the aggression received from coercive males (the

avoidance of male coercion hypothesis, H2c).

Table 1

Hypotheses, predictions and support for the hypotheses proposed to affect the duration of female cycle-specific phases

Hypotheses		Predictions	Support?[1]	
Hypothesis	Explanation		SD[2]	MSD[3]
H1: Intrasexual selection	*H1. Avoidance of female aggression hypothesis* = Females shorten their sexual receptivity to decrease their exposure to aggression from pregnant and lactating females who compete for access to paternal care .	↑ PL[4] -> ↓ (M)SD to ↓ consexual aggression.	—	—
H2: Intersexual selection & Sexual conflict	H2a: *Paternity concentration hypothesis* = Females lengthen their sexual receptivity to increase their chances of mating with the alpha male when several other females are simultaneously swollen.	↑ simultaneously receptive females -> ↑ (M)SD to ↑ access to preferred (dominant) male.	—	—
	H2b: *Paternity confusion hypothesis* = Females lengthen their sexual receptivity to mate with multiple males and decrease infanticide risk when the operational sex-ratio is female-biased.	Female-biased OSR -> ↑ (M)SD to mate with many males and confuse paternity.	—	—
	H2c: *Avoidance of male coercion hypothesis* = Females shorten their sexual receptivity to decrease their exposure to sexual coercion when there is strong male-male competition and social instability because of high numbers of in-group males.	Male-biased OSR or ↑ in-group males -> ↓ (M)SD to avoid coercive behaviour.	+	—

142 [1] + = support models, — = no support; [2]SD= swelling duration, [3]MSD= maximal-swelling duration; [4]PL = the sum of in-group pregnant and lactating females.

Materials and Methods

Study site and animals

Data were collected from identifiable females within two habituated groups, J and L, of a wild chacma baboon population at Tsaobis Leopard Park (22º22´S, 15º 44´E) in central Namibia. Studies on this population, which have been ongoing since 2000 [45], have revealed that the naturally-foraging baboons [45] exhibit low predation risk [46] and high female-female aggression rates [21,25] that can result in reproductive suppression [25]. Over the course of the study, troop numbers fluctuated around a median of 55 (44–69) in J troop and 52 (21–71) in L troop.

The data used in this study were collected periodically over 15 years, from 2005 to 2019, during field seasons lasting between 2 and 8 months *per annum*, usually during the austral winter. During field seasons, troops were visited daily, when possible, with some interruptions for other tasks or unforeseen circumstances. During troop visits, we collected data on troop composition (see below) and individual behaviour *ad libitum*.

Individuals' ages were known from observing births or from patterns of dentition wear following capture. Individuals' relative ranks were calculated annually from *ad libitum* recorded dominance interactions in the package Matman 1.1.4 (Noldus Information Technology, 2013). Absolute ranks were converted to a relative scale ranging from 0 (lowest rank) to 1 (highest rank), to control for group size, using the formula *1-((1-r)/(1-n))*, with *r* being individual's absolute rank ranging from 1 to the total group size *n*.

Troop composition and reproductive state data

Each day that the troop was contacted, a census was completed that recorded the identities of the individuals present and the reproductive states of each adult female. Females' reproductive states were recorded as: (1) pregnant (determined *post hoc* based on lack of resumption of swelling, reddening of the paracallosal skin and subsequent birth); (2) lactating (i.e., period following the birth of an infant until cycle resumption); (3) oestrous/swollen (i.e., exhibiting periovulatory swelling of the anogenital region); (4) non-swollen (i.e., deturgescent but not pregnant). For females in oestrus, we also recorded swelling size on a scale from 0.5 (smallest)-4 (largest), on an absolute (*cf.* the population) rather than relative (*cf.* for that particular female) scale [47]. As mentioned above, baboon troops were not followed daily, and as such there were some missing data during some females' cycles. We discarded any

observations where the start or end of the swelling duration was missing and thus could introduce inaccuracy.

For females in oestrus, swelling duration was calculated as the number of continuous days that a female was recorded with a swelling of any size. Because different mechanisms could act on the different parts of the cycle, in addition to swelling durations, we calculated the maximal-swelling duration as the number of days the focal female exhibited the largest swelling size during that respective cycle. We excluded from the initial data 8 observations of maximal-swelling duration which were calculated as 0, i.e., observations that showed that females were only swollen, and this resulted in 158 swellings and 158 maximal-swellings from 47 females (median number of swellings / female = 5, range = 1-14).

We calculated five variables describing aspects of the social environment relevant to our hypotheses: for H1 we calculated (1) the sum of pregnant and lactating females ("PL") during the time a focal female was either swollen or maximally-swollen. For hypothesis H2a, we measured the number of other females that were (2) not maximally-swollen and (3) maximally-swollen, both when the focal female started swelling. To test hypothesis H2b and part of H2c, we calculated the (4) operational sex ratio ("OSR") as the ratio of sexually active females (i.e., swollen at any level) to adult males. For H2c, we also calculated the (5) the number of in-group adult males.

Statistical analyses

All statistical analyses were performed using R (version 4.0.3, 2020-10-10).

Question 1: Repeatability of receptivity

To determine how much variation in females' receptivity part of the cycle was due to differences between individuals, we calculated the repeatability (R) of the swelling and maximal-swelling durations. R is calculated as $\frac{\sigma_\alpha^2}{\sigma_\alpha^2 + \sigma_\varepsilon^2}$, where σ_α^2 represents the between-group variance, σ_ε^2 the within-group variance ($\sigma_\alpha^2 + \sigma_\varepsilon^2$ = the total phenotypic variance, V_p [48]). A relatively high R estimate would indicate that the dependent variable exhibits high between- and low within-female variance; low values of R indicate traits with high within-female or low between-female variation [48].

We calculated R using the `rpt` function (`rptR` package), which can calculate repeatability for Poisson-distributed data, such as we had, while controlling for the following confounding fixed effects [49]: relative rank and age. Female identity and troop were included

as random effects. The permutations and bootstrapping were set to 1000 and confidence
intervals at 95%.

Question 2: Social determinants of swelling duration

To determine whether the social environment affected swelling durations, we used
generalized linear mixed models (GLMMs; `glmer` function, lme4 package [50]) to control
for repeated measures on individuals and the Poisson distribution of the duration counts. All
models followed the same basic structure and were specified to use "bobyqa" as an optimizer:
they included as fixed effects female age and relative rank, female identity and troop as random
effects, and specific predictor variables that targeted each hypothesis. To test for H1, H2a, H2b
and part of H2c—the effect of a male-skewed OSR—we ran two models (Table 2, models 1
and 2) with the following predictors: to test for H1 we included the number of pregnant and
lactating females specified with a rank interaction because higher-ranking females primarily
harass lower-ranking ones. To test for H2a, we used the number of not maximally-swollen
females when a focal female started swelling when the response was swelling duration; and the
number of maximally-swollen females when a focal female started swelling when the response
was maximal-swelling duration. Thirdly, to test for H2b and part of H2c—the effect of a male-
skewed OSR—we included the operational sex ratio (OSR, i.e., no. of receptive females/no. of
adult males) as a predictor. Finally, to test for H2c we ran two more models (Table 2, models
3 and 4, one for swelling and one for maximal-swelling duration) with the same predictors as
models 1 and 2, but with the number of in-group males instead of OSR, since the former is a
proxy of social instability in the sense that male ranks are more volatile when there is more
competition (i.e., more males) (A. Carter, E. Huchard, pers. Obs.).

Results

Females were swollen for an average (median) of 22 days (mean = 22.29, range 8-44)
and maximally-swollen for 6 days (median, mean = 6.6, range 1-24). There were no differences
between troops (Mann-Whitney U test: swelling duration: W = 3458.5, p = 0.233; maximal-
swelling duration: W = 3097, p = 0.949).

Question 1: Repeatability of receptivity

Swelling duration was repeatable at the individual level (R = 0.148, SE = 0.083, CI =
[0, 0.284], p = 0.012) (Table A1; Figure A1a), but most variation stemmed from within females

(Figure A2). Maximal-swelling duration was not repeatable (R = 0, SE = 0.03, CI = [0, 0.113],

p = 1; Table A1; Figure A1b) and all variation was detected within females (Figure A2).

Question 2: Social determinants of swelling duration

Age was the only fixed effect predictor that consistently predicted swelling (model 1:

β = -0.009, SE = 0.004, p = 0.030; model 3: β = -0.008, SE = 0.004, p = 0.064) and maximal-

swelling duration (model 2: β = -0.038, SE = 0.011, p < 0.001; model 4: β = -0.035, SE = 0.011,

p = 0.001) in three models. Rank was not significant in any model.

H1: Intrasexual selection

Avoidance of female aggression hypothesis

Neither swelling nor maximal-swelling duration responded to an increase in the number

of pregnant and lactating individuals (swelling: β = -0.008, SE = 0.023, p > 0.05; maximal-

swelling: β = -0.051, SE = 0.043, p > 0.05; Table 2; Figure 1).

H2: Intersexual selection

H2a Paternity concentration hypothesis

We tested whether female-female competition for the dominant mate contributed to

variation in swelling or maximal-swelling duration. However, we found no support for this:

greater numbers of simultaneously (non-maximally and maximally) swollen females did not

induce an increase in either duration (β = -0.015, SE = 0.017, p > 0.05; β = 0.12, SE = 0.078,

p > 0.05, respectively; Table 2; Figure 1).

H2b Paternity confusion hypothesis

We predicted that females should elongate their swelling or maximal-swelling phase in

response to a female-biased OSR to be able to mate with many males and reduce infanticide

risk. However, neither the length of swelling nor of maximal-swelling duration was

significantly impacted by this predictor (GLMMs, swelling: β = 0.053, SE = 0.066, p > 0.05;

maximal-swelling: β = -0.179, SE = 0.107, p > 0.05; Table 2; Figure 1).

Figure 1: Standardised forest plot estimates of models 1 and 2 with swelling (blue) and maximal-swelling duration

(red) as response variables. Only age shows a significant effect.

H2c Avoidance of male coercion hypothesis

When testing our last 'avoidance of sexual coercion' hypothesis, we found that swelling

duration followed the predicted decrease with an increasing number of males (Figure 2; Table

2). While the OSR did not affect swelling duration, the number of males did: each additional

male decreased swelling duration by 1.021 days (β = -0.021, SE = 0.009, p = 0.013; Figure 2).

The length of maximal-swelling period, however, was unaffected by the number of males (β =

-0.025, SE = 0.017, p = 0.134; Table 2).

Figure 2: Effect of the number of males on female swelling duration. The line indicates the predicted effect on

swelling duration by varying the focal variable, number of males. The shaded area represents the 95% CI, the dots

show the predicted values, with grey dots being single observations and black dots multiple overlapping

observations.

Table 2

Results of GLMMs with swelling or maximal-swelling duration as response variables. Age and rank were fixed

effects, and troop and female identity were random effects in all models. Estimate measures the effect of the

predictor on the response variable and is given in log-scale.

Models	Hypothesis tested	Response[1]	Predictor for specific hypothesis	Estimate	SE	z-value	p-value
Model 1→ SD[1]: PL[2]*rank + OSR[3] + not-mxm ♀[4]	H1	SD	PL*rank	-0.008	0.023	-0.334	0.738
	H2a	SD	not-mxm ♀	-0.015	0.017	-0.841	0.400
	H2b & H2c	SD	OSR(♀/♂)	0.053	0.066	0.809	0.419
Model 3→ SD: PL*rank + males[5] + not-mxm	H2c	SD	males	-0.021	0.009	-2.473	**0.013**
Model 2→ MSD[1]: PL*rank + OSR + mxm-sw ♀[6]	H1	MSD	PL*rank	-0.051	0.043	-1.173	0.241
	H2a	MSD	mxm-sw ♀	0.12	0.078	1.539\	0.124
	H2b & H2c	MSD	OSR(♀/♂)	-0.179	0.107	-1.678	0.093

Model 4→ MSD: PL*rank + males + mxm-sw ♀	H2c	MSD	males	-0.025	0.017	-1.499	0.134

286 [1]Swelling duration (SD), maximal-swelling duration (MSD), [2]sum of pregnant (P) and lactating females (L),
287 [3]operational sex ratio, [4]the number of not-maximally-swollen females when a focal female started swelling, [5]the
288 number of adult males, [6]the number of maximally-swollen females when a focal female started swelling.

Discussion

The present study investigated the variation in the length of receptivity within versus
across females. We tested whether females could manipulate receptivity to decrease aggression
from competitors or coercive males or increase their mating opportunities. First, we detected
high within-female variation in the length of sexual receptivity. Second, we found support for
the 'avoidance of male coercion hypothesis', suggesting that females may be able to
strategically adjust the length of their sexual receptivity to their social environment.

Our results support CLM as a sexually-selected mechanism that reduces the costs of male
coercion. Our results show that the duration of the swollen part of the cycle is reduced, but not
of the maximally-swollen phase; the number of non-swollen days between two cycles seems
unaffected (data not shown). Overall, this suggests that females may start swelling at the same
time of their oestrus cycle, but may reach maximal-swelling size slightly earlier when they are
more males in the group, and then show the same duration of maximal-swelling. Chacma
baboon males start to coerce females even before they start to swell [11], which may explain
why swelling duration, but not maximal-swelling duration, changed in response to the number
of adult males in the group. Shorter swelling durations when male-male competition and risk
of coercion is high could lead to shorter periods of aggression and a lower risk of injury [11],
an adaptive response from females to male coercion. Perhaps surprisingly, a male-biased OSR
did not elicit a similar response. This may be because it does not capture social instability and
male-male competition as accurately as the number of in-group males: a male-skewed OSR
can be either due to a low number of swollen females or to a high number of in-group males,
but only the latter was expected to affect females' cycles.

We propose two possible mechanisms that may produce the effect of CLM. First,
previous studies in baboons have shown that socially-induced stress may result in a decrease
of the swollen period [34,43], which is probably mediated by increased female cortisol levels
[51]. The social dynamics involving male immigration events, high male-male competition and
unstable male hierarchies are stressful for females because they typically increase the intensity
of coercion [16,25,52] and harassment [34,43,53]. It is possible that the same mechanism is

responsible for the shortening of the cycle we observe in this population. Indeed, higher
circulating cortisol could compromise the timing of the expression of follicle stimulating
hormone, a mechanism seen in ewes [54]. However, glucocorticoids in cycling chacma baboon
females did not rise during male take-over events [52], suggesting that cortisol may not be the
mechanism. A second possible mechanism is that females might arrive faster at maximal-
swelling—thus showing the reported decrease in the swelling period—via a male-mediated
increase in estrogen levels, similar to the Vandenbergh effect, i.e., the earlier than normal
female maturation in the presence of an unrelated adult male [35]. Estrogen is positively
correlated with swelling size in baboons [55], and estrogen has been implicated in earlier sexual
maturation in geladas following the arrival of a novel male [36].

Our results also suggest that the costs of coercion may, in this population, be
comparatively higher than the risk of infanticide for females. At a comparative level, increased
infanticide risk is associated with elongated receptivity periods across Cercopithecine primates
[32]. It would thus seem adaptive for females to elongate, rather than shorten, their receptivity
in response to a high number of males which increases infanticide risk [25]. At Tsaobis, the
costs of coercion for females may be high [11] and the risk of infanticide relatively low
compared to some other chacma baboon populations such as Moremi, with a much lower
frequency of dominance take-overs, possibly due to the lower predation risk [56]. In addition,
mate-guarding episodes are long in chacma baboons, and as a consequence paternity certainty
towards the alpha-male is high, suggesting that females either fail to dilute paternity certainty
due to indirect coercion [57], or alternatively chose to concentrate paternity in order to gain
paternal services [44]. This may also explain the lack of evidence for H2b: paternity dilution
is costly or not a necessary strategy for Tsaobis females.

We predicted that higher numbers of receptive females would prompt an elongation of
oestrous duration so that females can concentrate paternity in the alpha male but found no
support for such prediction. At least two explanations may account for this absence of evidence.
First, because female chacma baboons are non-seasonal breeders [25], oestrous synchrony in
this population is low, with a median of two concurrently swollen females. Consequently,
mating with the preferred male is possible for most cycling females and females may not need
to elongate their fertile period. Second, the substantial aggression among cycling females in
Tsaobis [21,25] may limit females' propensity to extend their receptivity period. This is
predominantly true for lower-ranking oestrous females and females who are not mate-guarded,
which tend to be the usual target of aggression from higher-ranking females.

One reason why the number of pregnant and lactating females did not affect receptivity
duration could be that female reproductive suppression in the form of aggression was given by
the female friends of a particular male [24]. Specifically, aggression was initiated by pregnant
and lactating females who were friends of the male that was mate-guarding the receptive
female. A more targeted approach that considers as a predictor the number of female friends a
swollen female's mate-guarding male has might therefore provide evidence for CLM.
Alternatively, as for the other explanations, sexual coercion from males might have larger
consequences on female fitness and be under stronger selection.

Lastly, we found that maximal-swelling duration was not repeatable, whereas almost
15% of the variance in the swelling phase was repeatable. Our results do not mean that there
are no between-individual differences in maximal-swelling duration, but rather that those
differences are not large relative to within-individual differences and are not consistent
between females. The failure to detect significant repeatability in maximal-swelling duration
could also reflect individual variation in the physiological responses to aggression and stress
stemming from changes in the social environment. Our findings show that cycle phases are
labile traits that females are potentially manipulating. At a comparative level, high variation in
oestrous duration has been detected in animals (e.g., captive jaguars, with a mean duration of
10.42 and a range of 7 to 15 days [58]). Women, in contrast, show repeatable cycle length,
even though there is some variation within [59] and across women [60], as well as extensive
variation in average cycle length across human populations [61]. Here, it may be that baboons'
low repeatability of swelling duration reflects a greater selective pressure on females' ability
to manipulate cycle length than women, who conceal ovulation [62].

In summary, this study correlated variation in the length of female sexual receptivity
with aspects of the social environment that may influence the intensity of female reproductive
competition and of intersexual conflict. We reported empirical evidence for cycle length
manipulation as a possible female counter-strategy to male coercion. Specifically, we found
that receptive females respond to an increment in the number of in-group males by decreasing
their swelling duration, presumably to reduce male coercion. In contrast, we failed to find any
evidence that female-female competition, or female counter-strategies to infanticide, may
influence the length of sexual receptivity, while we confirmed previous research that suggested
the OSR is not an accurate predictor of rates of female-female aggression [25,63]. Following
these results, future investigations and cross-population studies of the scale and sources of
variation of sexual receptivity within and among females, and of the underlying physiological
mechanisms responsible for the lengthening of swelling duration, will further confirm the

extent to which females can modulate it in response to changes in their social environment. In any case, it seems that CLM will favor an extension of receptivity in cases of increased infanticide and low coercion risk, and the opposite when infant survival is more or less secure.

Ethics

Our research procedures were evaluated and approved by the Ethics Committee of the Zoological Society of London and the Ministry of Environment and Tourism, Namibia (MET Research/ Collecting Permits 886/2005, 1039/2006, 1186/2007, 1302/2008, 1379/2009, 1486/2010, 1486/2011, 1696/2012, 1786/2013, 1892/ 2014, 2009/2015, 2147/2016, 2303/2017, RPIV00392018/2019) and adhered to the ASAB/ABS Guidelines for the Treatment of Animals in Behavioural Research and Teaching.

Data accessibility

Authors' contributions

AJC, EH and FD contributed to the conceptualization. AJC and EH designed the study. GC established the study system and oversaw data collection, with assistance from AJC and EH. FD performed the statistical analyses. All authors contributed to draft the manuscript. All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Competing interests

We declare we have no competing interests.

Funding

Long-term data collection was partially funded by grants from: the Agence Nationale de la Recherche (ANR ERS-17-CE02-0008, 2018-2021) awarded to EH; a Fenner School of Environment and Society Scholarship awarded to A.J.C., a Ministère de l'Education et de la Recherche Studentship awarded to E.H; a NERC Advanced Research Fellowship awarded to G.C.; Leakey Foundation grants awarded to AJC; and a Templeton World Charity Foundation grant (TWCF0502) awarded to AJC. This paper is a contribution ISEM (n° TBD) and a publication of the ZSL Institute of Zoology's Tsaobis Baboon Project, supported by Research England.

Acknowledgments

We would like to thank the past Tsaobis Baboon Project leaders, students, volunteers and interns for contributing to the long-term data collection at the field site. This research was carried out with the permission of the Ministry of Environment and Tourism, the Ministry of

Land Reform, and the National Commission on Research, Science, and Technology. We

further thank the Tsaobis beneficiaries for permission to work at Tsaobis, the Gobabeb Namib

Research Institute for affiliation, and Johan Venter and the Snyman and Wittreich families for

permission to work on their land. Some findings formed part of a dissertation submitted in

partial fulfilment of the requirements of the degree of MSc of the University College London

in September 2021. This paper is a publication of the ZSL Institute of Zoology's Tsaobis

Baboon Project. Contribution ISEM n°202x-xx.

1. Kleindorfer S, Wasser SK. 2004 Infant handling and mortality in yellow baboons (Papio
cynocephalus): evidence for female reproductive competition? *Behav Ecol Sociobiol* **56**.
(doi:10.1007/s00265-004-0798-1)

2. Drea CM. 2005 Bateman Revisited: The Reproductive Tactics of Female Primates1.
Integrative and Comparative Biology **45**, 915–923. (doi:10.1093/icb/45.5.915)

3. Stumpf RM, Martinez-Mota R, Milich KM, Righini N, Shattuck MR. 2011 Sexual conflict
in primates. *Evol. Anthropol.* **20**, 62–75. (doi:10.1002/evan.20297)

4. Tang-Martínez Z. 2016 Rethinking Bateman's Principles: Challenging Persistent Myths of
Sexually Reluctant Females and Promiscuous Males. *The Journal of Sex Research* **53**,
532–559. (doi:10.1080/00224499.2016.1150938)

5. Watson-Capps JJ. 2009 2 Evolution of Sexual Coercion with Respect to Sexual Selection
and Sexual Conflict Theory. In *2 Evolution of Sexual Coercion with Respect to Sexual
Selection and Sexual Conflict Theory*, pp. 23–41. Harvard University Press.
(doi:10.4159/9780674054349-002)

6. van Schaik CP, Pradhan GR, van Noordwijk MA. 2004 Mating conflict in primates:
infanticide, sexual harassment and female sexuality. In *Sexual Selection in Primates: New
and Comparative Perspectives* (eds CP van Schaik, PM Kappeler), pp. 131–150.
Cambridge: Cambridge University Press. (doi:10.1017/CBO9780511542459.010)

7. Qi X-G, Grueter CC, Fang G, Huang P-Z, Zhang J, Duan Y-M, Huang Z-P, Garber PA, Li
B-G. 2020 Multilevel societies facilitate infanticide avoidance through increased extrapair
matings. *Animal Behaviour* **161**, 127–137. (doi:10.1016/j.anbehav.2019.12.014)

8. Clutton-Brock TH, Parker GA. 1995 Sexual coercion in animal societies. *Animal
Behaviour* **49**, 1345–1365. (doi:10.1006/anbe.1995.0166)

9. Stumpf RM, Boesch C. 2010 Male aggression and sexual coercion in wild West African
chimpanzees, Pan troglodytes verus. *Animal Behaviour* **79**, 333–342.
(doi:10.1016/j.anbehav.2009.11.008)

10.   Eldakar OT, Gallup AC. 2011 The Group-Level Consequences of Sexual Conflict in
Multigroup Populations. *PLOS ONE* **6**, e26451. (doi:10.1371/journal.pone.0026451)

11.   Baniel A, Cowlishaw G, Huchard E. 2017 Male Violence and Sexual Intimidation in a
Wild Primate Society. *Current Biology* **27**, 2163-2168.e3. (doi:10.1016/j.cub.2017.06.013)

12.   Baniel A, Delaunay A, Cowlishaw G, Huchard E. 2019 Oestrous females avoid
mating in front of adult male bystanders in wild chacma baboons. *R. Soc. open sci.* **6**,
181009. (doi:10.1098/rsos.181009)

13.   Muller MN, Thompson ME, Kahlenberg SM, Wrangham RW. 2011 Sexual coercion
by male chimpanzees shows that female choice may be more apparent than real. *Behav
Ecol Sociobiol* **65**, 921–933. (doi:10.1007/s00265-010-1093-y)

14.   Muller MN, Kahlenberg SM, Emery Thompson M, Wrangham RW. 2007 Male
coercion and the costs of promiscuous mating for female chimpanzees. *Proceedings of the
Royal Society B: Biological Sciences* **274**, 1009–1014. (doi:10.1098/rspb.2006.0206)

15.   Parga JA, Henry AR. 2008 Male aggression during mating: evidence for sexual
coercion in a female dominant primate? *Am. J. Primatol.* **70**, 1187–1190.
(doi:10.1002/ajp.20609)

16.   Polo P, Hernández-Lloreda V, Colmenares F. 2014 Male Takeovers Are
Reproductively Costly to Females in Hamadryas Baboons: A Test of the Sexual Coercion
Hypothesis. *PLoS ONE* **9**, e90996. (doi:10.1371/journal.pone.0090996)

17.   Clutton-Brock T. 2007 Sexual Selection in Males and Females. *Science* **318**, 1882–
1885. (doi:10.1126/science.1133311)

18.   Clutton-Brock T. 2009 Sexual selection in females. *Animal Behaviour* **77**, 3–11.
(doi:10.1016/j.anbehav.2008.08.026)

19.   Huchard E, Courtiol A, Benavides JA, Knapp LA, Raymond M, Cowlishaw G. 2009
Can fertility signals lead to quality signals? Insights from the evolution of primate sexual
swellings. *Proc. R. Soc. B.* **276**, 1889–1897. (doi:10.1098/rspb.2008.1923)

20.   Vaillancourt T. 2013 Do human females use indirect aggression as an intrasexual
competition strategy? *Phil. Trans. R. Soc. B* **368**, 20130080. (doi:10.1098/rstb.2013.0080)

21.   Huchard E, Cowlishaw G. 2011 Female–female aggression around mating: an extra
cost of sociality in a multimale primate society. *Behavioral Ecology* **22**, 1003–1011.
(doi:10.1093/beheco/arr083)

22.   Tobler R, Pledger S, Linklater W. 2010 No evidence for ovarian synchrony or
asynchrony in hamadryas baboons. *Animal Behaviour* **80**, 829–837.
(doi:10.1016/j.anbehav.2010.07.018)

23.   Kappeler PM, Fichtel C. 2012 Female reproductive competition in *Eulemur rufifrons* :
eviction and reproductive restraint in a plurally breeding Malagasy primate: EVICTION
AND REPRODUCTIVE RESTRAINT IN A LEMUR. *Molecular Ecology* **21**, 685–698.
(doi:10.1111/j.1365-294X.2011.05255.x)

24.   Baniel A, Cowlishaw G, Huchard E. 2018 Jealous females? Female competition and
reproductive suppression in a wild promiscuous primate. *Proc. R. Soc. B.* **285**, 20181332.
(doi:10.1098/rspb.2018.1332)

25.   Baniel A, Cowlishaw G, Huchard E. 2018 Context dependence of female reproductive
competition in wild chacma baboons. *Animal Behaviour* **139**, 37–49.
(doi:10.1016/j.anbehav.2018.03.001)

26.   Watson NL, Simmons LW. 2010 Reproductive competition promotes the evolution of
female weaponry. *Proc. R. Soc. B.* **277**, 2035–2040. (doi:10.1098/rspb.2009.2335)

27.   Bro-Jørgensen J. 2002 Overt female mate competition and preference for central
males in a lekking antelope. *Proceedings of the National Academy of Sciences* **99**, 9290–
9293. (doi:10.1073/pnas.142125899)

28.   Robinson MR, Kruuk LEB. 2007 Function of weaponry in females: the use of horns
in intrasexual competition for resources in female Soay sheep. *Biol. Lett.* **3**, 651–654.
(doi:10.1098/rsbl.2007.0278)

29.   Stockley P, Bro-Jørgensen J. 2011 Female competition and its evolutionary consequences in mammals. *Biol Rev Camb Philos Soc* **86**, 341–366. (doi:10.1111/j.1469-185X.2010.00149.x)

30.   Rosvall KA. 2011 Intrasexual competition in females: evidence for sexual selection? *Behavioral Ecology* **22**, 1131–1140. (doi:10.1093/beheco/arr106)

31.   Wasser SK, Starling AK. 1988 Proximate and ultimate causes of reproductive suppression among female yellow baboons at Mikumi National Park, Tanzania. *Am. J. Primatol.* **16**, 97–121. (doi:10.1002/ajp.1350160202)

32.   Nunn CL. 1999 The evolution of exaggerated sexual swellings in primates and the graded-signal hypothesis. *Animal Behaviour* **58**, 229–246. (doi:10.1006/anbe.1999.1159)

33.   Deschner T. 2000 Sexual swellings in female hamadryas baboons after male take-overs: 'deceptive' swellings as a possible female counter-strategy against infanticide. *Am J Primatol* **52**, 157–168. (doi:10.1002/1098-2345(200012)52:4<157::AID-AJP1>3.0.CO;2-L)

34.   Rowell TE. 1970 BABOON MENSTRUAL CYCLES AFFECTED BY SOCIAL ENVIRONMENT. *Reproduction* **21**, 133–141. (doi:10.1530/jrf.0.0210133)

35.   VANDENBERGH JG. 1967 Effect of the Presence of a Male on the Sexual Maturation of Female Mice. *Endocrinology* **81**, 345–349. (doi:10.1210/endo-81-2-345)

36.   Lu A, Feder JA, Snyder-Mackler N, Bergman TJ, Beehner JC. 2021 Male-Mediated Maturation in Wild Geladas. *Current Biology* **31**, 214-219.e2. (doi:10.1016/j.cub.2020.10.003)

37.   Bruce HM. 1959 An Exteroceptive Block to Pregnancy in the Mouse. *Nature* **184**, 105–105. (doi:10.1038/184105a0)

38.   Tinsley Johnson E, Snyder-Mackler N, Lu A, Bergman TJ, Beehner JC. 2018 Social and ecological drivers of reproductive seasonality in geladas. *Behavioral Ecology* **29**, 574–588. (doi:10.1093/beheco/ary008)

39.   Palombit RA, Cheney DL, Seyfarth RM. 2001 Female–female competition for male 'friends' in wild chacma baboons (Papio cynocephalus ursinus). *Animal Behaviour* **61**, 1159–1171. (doi:10.1006/anbe.2000.1690)

40.   Cheney DL, Silk JB, Seyfarth RM. 2012 Evidence for intrasexual selection in wild female baboons. *Animal Behaviour* **84**, 21–27. (doi:10.1016/j.anbehav.2012.03.010)

41.   Clutton-Brock T, Huchard E. 2013 Social competition and its consequences in female mammals. *J Zool* **289**, 151–171. (doi:10.1111/jzo.12023)

42.   Baniel A, Cowlishaw G, Huchard E. 2016 Stability and strength of male-female associations in a promiscuous primate society. *Behav Ecol Sociobiol* **70**, 761–775. (doi:10.1007/s00265-016-2100-8)

43.   Wasser S. 1983 13 – Reproductive Competition and Cooperation among Female Yellow Baboons*. (doi:10.1016/B978-0-12-735950-2.50018-9)

44.   Huchard E, Alvergne A, Féjan D, Knapp LA, Cowlishaw G, Raymond M. 2010 More
than friends? Behavioural and genetic aspects of heterosexual associations in wild chacma
baboons. *Behav Ecol Sociobiol* **64**, 769–781. (doi:10.1007/s00265-009-0894-3)

45.   Carter AJ, Baniel A, Cowlishaw G, Huchard E. 2020 Baboon thanatology: responses
of filial and non-filial group members to infants' corpses. *R. Soc. open sci.* **7**, 192206.
(doi:10.1098/rsos.192206)

46.   Carter AJ, Torrents Ticó M, Cowlishaw G. 2016 Sequential phenotypic constraints on
social information use in wild baboons. *eLife* **5**, e13125. (doi:10.7554/eLife.13125)

47.   Gesquiere LR, Wango EO, Alberts SC, Altmann J. 2007 Mechanisms of sexual
selection: sexual swellings and estrogen concentrations as fertility indicators and cues for
male consort decisions in wild baboons. *Horm Behav* **51**, 114–125.
(doi:10.1016/j.yhbeh.2006.08.010)

48.   Nakagawa S, Schielzeth H. 2010 Repeatability for Gaussian and non-Gaussian data: a
practical guide for biologists. *Biological Reviews* , no-no. (doi:10.1111/j.1469-
185X.2010.00141.x)

49.   Stoffel MA, Nakagawa S, Schielzeth H. 2017 rptR: repeatability estimation and
variance decomposition by generalized linear mixed-effects models. *Methods Ecol Evol* **8**,
1639–1644. (doi:10.1111/2041-210X.12797)

50.   Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting Linear Mixed-Effects Models
Using **lme4**. *J. Stat. Soft.* **67**. (doi:10.18637/jss.v067.i01)

51.   Alberts SC, Sapolsky RM, Altmann J. 1992 Behavioral, endocrine, and
immunological correlates of immigration by an aggressive male into a natural primate
group. *Horm Behav* **26**, 167–178. (doi:10.1016/0018-506x(92)90040-3)

52.   Beehner JC, Bergman TJ, Cheney DL, Seyfarth RM, Whitten PL. 2005 The effect of
new alpha males on female stress in free-ranging baboons. *Animal Behaviour* **69**, 1211–
1221. (doi:10.1016/j.anbehav.2004.08.014)

53.   Dunbar RIM, Sharman M. 1983 Female Competition for Access to Males Affects
Birth Rate in Baboons. *Behavioral Ecology and Sociobiology* **13**, 157–159.

54.   Breen KM, Billings HJ, Wagenmaker ER, Wessinger EW, Karsch FJ. 2005 Endocrine
basis for disruptive effects of cortisol on preovulatory events. *Endocrinology* **146**, 2107–
2115. (doi:10.1210/en.2004-1457)

55.   Stevens VC. 1997 Some reproductive studies in the baboon. *Hum Reprod Update* **3**,
533–540. (doi:10.1093/humupd/3.6.533)

56.   Cheney DL *et al.* 2004 Factors Affecting Reproduction and Mortality Among
Baboons in the Okavango Delta, Botswana. *International Journal of Primatology* **25**, 401–
428. (doi:10.1023/B:IJOP.0000019159.75573.13)

57.   Clarke P, Pradhan G, Schaik C van. 2009 3 Intersexual Conflict in Primates:
Infanticide, Paternity Allocation, and the Role of Coercion. In *3 Intersexual Conflict in*

Primates: Infanticide, Paternity Allocation, and the Role of Coercion, pp. 42–78. Harvard University Press. (doi:10.4159/9780674054349-003)

58. Viau P, Rodini DC, Sobral G, Martins GS, Morato RG, de Oliveira CA. 2020 Puberty and oestral cycle length in captive female jaguars Panthera onca. *Conservation Physiology* **8**, coaa052. (doi:10.1093/conphys/coaa052)

59. Shea AA, Vitzthum VJ. 2020 The extent and causes of natural variation in menstrual cycles: Integrating empirically-based models of ovarian cycling into research on women's health. *Drug Discovery Today: Disease Models* **32**, 41–49. (doi:10.1016/j.ddmod.2020.11.002)

60. Mihm M, Gangooly S, Muttukrishna S. 2011 The normal menstrual cycle in women. *Anim Reprod Sci* **124**, 229–236. (doi:10.1016/j.anireprosci.2010.08.030)

61. Alvergne A, Högqvist Tabor V. 2018 Is Female Health Cyclical? Evolutionary Perspectives on Menstruation. *Trends Ecol Evol* **33**, 399–414. (doi:10.1016/j.tree.2018.03.006)

62. Benshoof L, Thornhill R. 1979 The evolution of monogamy and concealed ovulation in humans. *Journal of Social and Biological Structures* **2**, 95–106. (doi:10.1016/0140-1750(79)90001-0)

63. Patterson SK, Strum SC, Silk JB. 2021 Resource competition shapes female–female aggression in olive baboons, Papio anubis. *Animal Behaviour* **176**, 23–41. (doi:10.1016/j.anbehav.2021.03.013)