1   Cycle length flexibility: is the duration of sexual receptivity associated with changes

2   in the social environment?

3   Darmis Fragkiskos[1,2], Huchard Élise[3], Cowlishaw Guy[4] & Carter J. Alecia[1]

[1]Department of Anthropology, University College London, London, UK.

[2]Current address: Max Planck Institute for Evolutionary Biology, Plön, Germany

[3]Institut des Sciences de L'Evolution de Montpellier, UMR 5554, CNRS, Université de Montpellier,

Montpellier, France

[4]Institute of Zoology, Zoological Society of London, London, UK

Research in social mammals has revealed the complexity of strategies females use in
response to female-female reproductive competition and sexual conflict. One point at which
competition and conflict manifests acutely is during sexual receptivity, indicated by swellings
in some primates. Whether females can adjust their sexual receptivity from cycle to cycle to
decrease reproductive competition and sexual conflict in response to changes in the social
environment has not been tested. As a first step, this study explores whether sexual
receptivity duration is predicted by the social environment in wild female chacma baboons
(*Papio ursinus*). Given that female baboons face intense reproductive competition and sexual
coercion, we predicted that: females could shorten the duration of their sexual receptive
period to reduce female-female aggression and male coercion or increase it to access multiple
or their preferred male(s). We quantified 157 ovulatory cycles from 46 wild females living in
central Namibia recorded over 15 years. We found no support for our hypothesis; however,
we found a negative correlation between maximal-swelling duration and group size, a proxy
of within-group competition. This study provides further evidence that swelling is costly for
females as well as a testable framework for future investigations of 'cycle length
manipulation'.
Keywords: cycle length manipulation, female-female competition, sexual coercion, sexual
selection, social environment

Introduction

Recent advances in sexual selection theory underline the importance of female-female

competition, along with mate choice, in generating variance in female reproductive success

[1–4]. In the first case, females tend to face competition with consexuals over access to males

and reproductive resources [5–8]. This competition for access to males increases when there

is synchrony of females' fertility (i.e., ovarian synchrony), which creates a female-biased

operational sex ratio (OSR) and increases aggression received by receptive females [9,10]. In

general, females compete with consexuals for (a) their preferred mates (e.g., chacma baboons

Papio Ursinus: [9]; red-fronted lemurs *Eulemur rufifrons*: [11]), (b) paternal services for their

offspring (e.g., chacma baboons: [12,13]) and (c) other reproductive resources [14], such as

breeding territories (e.g., lek center [15]) or food resources [16]. Those benefits are achieved

either through monopolisation of resources directly from females (i.e., contest competition

for food) [17] or indirectly (e.g., nutrient-rich spermatophores in insects [18], low predation

risk in lek centers [15]). Antagonism with consexuals can manifest through non-aggressive

interactions (e.g., signals [18]) and escalate up to physical aggression, which can be

detrimental to the receiver's short- or long-term fitness [19].

Considering mate choice, it may be advantageous for females to concentrate paternity

into a single male who can provide benefits in the form of good genes or paternal services

such as protection to her or her offspring. However, in group living species, a number of

social pressures may decrease the possibility for females to express their mating preference.

For example, recent studies highlight the role of sexual conflict [20], which frequently

manifests via infanticide in mammals. Indeed, infanticide risk may push females to mate with

multiple rather than single partners, to dilute paternity certainty among multiple males who

subsequently restrain from attacking an offspring that they could have sired [21,22]. Female

choice may be further constrained by sexual coercion, when males direct aggression to

females [3,20,23], either to increase the probability that they will mate with them or to

decrease the probability that they will mate promiscuously [24]. In such cases, females will

mate with their aggressors if resisting is more costly than giving in [20]. In groups of water

striders (*Aquarius remigis*), aggressive males acquire more matings than less aggressive ones

[25]. In primates, male baboons (*Papio ursinus*) that attack pre-ovulatory females are more

likely to monopolize them when receptive [26] and females do not copulate close to adult

males to avoid bystanders' aggression [26,27]. Also, female chimpanzees (*Pan troglodytes*)

initiate copulations most frequently with those males that were most aggressive towards them

throughout their cycle [28,29] and male lemurs (*Lemur catta*) that are not preferred can force

copulations [30]. However, male coercion can lead to female counterstrategies and

antagonistic co-evolution between the sexes [20] since coercion is associated with multiple

costs for females. For example, the social dynamics involving male immigration events, high

male-male competition and unstable male hierarchies are stressful for females because they

typically increase the intensity of coercion [13,31,32] and harassment [33–35]. Moreover,

females suffer costs from coercion when: they are forced to mate with subordinate or

incompatible males; direct aggression or harassment increases stress hormone secretion [29];

males reduce their preferred level of parental care [23]; or coercion impairs female choice

[31].

By decreasing the length of their receptive period, females could potentially decrease

(1) time that they are subject to aggression they receive from other females (female-female

competition) (2) and/or males (sexual coercion). By contrast, by increasing the length of their

receptive period, females could potentially (3) increase their probability of mating with a

preferred male (female choice) or (4) multiple males (paternity confusion). Evidence for

manipulation of female cycles exists [19,35–37], with studies detecting physiological changes

in female cycling patterns (e.g., changes to cycle length, resumption of cycling) as stress

responses to the social environment. Socially-stressed primate females, for example by being

harassed near menstruation, have longer menstrual cycles [35]. Cycling female yellow

baboons (*Papio cynocephalus*) that receive aggression from consexuals tend to have more

cycles to conception and longer interbirth intervals [19], while pregnant females that are

systematically harassed experience reduced fitness (through abnormal gestation length,

spontaneous abortion, and premature delivery) [19]. Psychosocial stress stemming from

social instability, in the form of male take-overs, also induces female physiological

responses: the presence of unfamiliar males can mediate early sexual maturation [38,39] or

abortion [40,41] in female mammals, in a likely effort to limit infanticide risk. Novel males in

geladas (*Theropithecus gelada*) prompt females of any reproductive state (immature, cycling,

lactating, or pregnant) to start or resume cycling [39,42], while female hamadryas (*Papio*

hamadryas) with young infants develop swellings to advertise a reduced postpartum

amenorrhea that results in copulations with novel males during take-overs [37]. However,

these swellings are deceptive signals that ultimately decrease infanticide risk (four out of five

infants survived in [37]) since females did not conceive during the first cycles after the take-

over, their reproduction was not accelerated and the interbirth intervals were not shorter

compared to times with no male take-overs. Despite the above results, no study to date has

proposed a theoretical framework that considers cycle length manipulation as a
counterstrategy to sexual selection pressures. Additionally, the aforementioned studies
provide evidence of cycle manipulation as an acute reaction to social pressures but have not
explicitly tested or explored the fluctuations in sexual receptivity in response to changes in
females' social environment.

In this study, we explored whether there is a correlation between females' oestrous
duration and the social environment. We assumed that females could adjust their oestrous
cycles to either reduce aggression from (1) coercive males or (2) females or to increase their
access to (3) their preferred or (4) multiple males. We focused on female chacma baboons
because they exhibit elevated female-female competition compared to other primates
[13,43,44], live in multimale-multifemale groups, mate promiscuously [26,45], and have
exaggerated swellings that reliably indicate females' stage of the follicular phase of the
oestrus cycle [36]. Female chacma baboons are an ideal model to study female-female
competition, intersexual competition, as well as sexual coercion, since they target the
dominant male as a sexual partner [13,44] but also mate promiscuously [9] outside their peak
swelling to establish more males with a nonzero paternity probability or to initiate male-male
competition (e.g., sperm competition) [27,36]. In addition, males' monopolisation of only one
receptive female at a time (i.e., mate-guarding: [7,12]) results in female-female competition
over sexual access to particular males [9,13], who might be better able to protect them from
aggressive consexuals or provide future postpartum care for the offspring [46].

To address our overarching aim, we proceeded in two steps. First, we explored the
variation within and among females in the durations of swelling (SD) and maximal-swelling
(MSD) stages of the cycle and we predicted that different mechanisms could act on each part
of the cycle. We predicted that there would be more variation within females' cycles than
between females across each of their cycles. Second, we tested whether variation in the
duration of specific cycle phases was associated with four aspects of the social environment
(summarised in Table 1).

First, considering female intrasexual selection, we tested whether female competition
for paternal care was associated with variation in the duration of receptivity (the avoidance of
female aggression hypothesis [H1]). Pregnant and lactating females aggressively target
swollen females to decrease competition for paternal care of their own offspring, or to induce
reproductive suppression so that their own infants are not born at the same time as many
others [33]. For this reason, we predicted a positive correlation between in-group pregnant
and lactating females and aggression in cycling females that would result in receptive females

terminating the swelling earlier and, ultimately, decreasing their exposure to consexual

aggression.

Our second hypothesis dealt with intersexual selection and sexual conflict (H2): we

hypothesized that females' SD and MSD would show a correlation with competition for

access the dominant mate (the paternity concentration hypothesis, H2a). We predicted that,

when competition for access to the alpha male was likely because there were simultaneously-

swollen individuals, females would respond by delaying ovulation—through an elongation of

their swelling or MSD—to increase the probability of mating with this male. By doing so

females may benefit from (a) good genes [13,36] and (b) concentrating paternity certainty in

their preferred male, usually the dominant, who is also the most likely to pose a future risk of

infanticide if they fail to mate with him [13,47].

We also tested whether the risk of infanticide from multiple males correlated with

patterns of variation in female baboons' cycle-phases (the paternity confusion hypothesis,

H2b). We predicted that an increase in swelling duration (SD) and maximal-swelling duration

(MSD) will be associated with a female-skewed operational sex ratio (OSR) so that females

can mate with many males to reduce infanticide risk. Finally, we tested whether the effect of

social instability in the form of high numbers of in-group males (i.e., strong male-male

competition) correlated with patterns of variation in a female's receptivity phases. We

predicted that a male-skewed OSR or a high number of in-group males translates into high

male-male competition, which in turn increases the risk of sexual coercion and male-initiated

aggression for receptive females [12,26]. Under this scenario, a negative association between

oestrous duration and the number of males or a male-biased OSR might provide evidence for

females' attempts to escape or minimize the aggression received from coercive males (the

avoidance   of   male   coercion   hypothesis,   H2c).

Table 1

Hypotheses, predictions and support for the hypotheses proposed to affect the duration of female cycle-specific phases.

Hypotheses		Predictions	Support?[1]	
Hypothesis	Explanation		SD[2]	MSD[3]
H1: Intrasexual selection	*H1. Avoidance of female aggression hypothesis* = Pregnant and lactating females predict shorter durations of female sexual receptivity, possibly because those females could attempt to decrease their exposure to aggression.	↑ P females + L females[4] -> ↓ SD to ↓ consexual aggression. ↑ PL[5] -> ↓ MSD to ↓ consexual aggression.	—	—
H2: Intersexual selection & Sexual conflict	H2a: *Paternity concentration hypothesis* = Females sexual receptivity is predicted to increase when several other females are simultaneously swollen, possibly because this increase could correlate with higher chances of mating with the alpha male.	↑ simultaneously receptive females -> ↑ (M)SD to ↑ access to preferred (dominant) male.	—	—
	H2b: *Paternity confusion hypothesis* = Females sexual receptivity is predicted to increase in response to a female-biased operational sex-ratio, possibly because females attempt to mate with multiple males and decrease infanticide risk.	Female-biased OSR -> ↑ (M)SD to mate with many males and confuse paternity.	—	—
	H2c: *Avoidance of male coercion hypothesis* = Females their sexual receptivity is predicted to decrease in response to strong male-male competition and social instability, possibly because females attempt to decrease their exposure to sexual coercion. Strong male-male competition and social instability can be inferred by an increasing numbers of in-group males.	Male-biased OSR or ↑ in-group males -> ↓ (M)SD to avoid coercive behaviour.	Males: — OSR: —	Males: — OSR: —

155 [1] — = no support, [2]SD= swelling duration, [3]MSD= Maximal-swelling duration; [4]PL = the number of in-group pregnant and of lactating females, [5]The sum of

156 in-group   pregnant   and   lactating.

Materials and Methods

Study site and animals

Data were collected from identifiable females within two habituated groups, J and L, of a wild chacma baboon population at Tsaobis Leopard Park (22º22´S, 15º 44´E) in central Namibia. Studies on this population, which have been ongoing since 2000 [48], have revealed that the naturally-foraging baboons [48] exhibit low predation risk [49] and high female-female aggression rates [9,13] that can result in reproductive suppression [13]. Over the course of the study, troop numbers fluctuated around a median of 55 (44–69) in J troop and 52 (21–71) in L troop.

The data used in this study were collected periodically over 15 years, from 2005 to 2019, during field seasons lasting between 2 and 8 months *per annum*, usually during the austral winter. During field seasons, troops were visited daily, when possible, with some interruptions for other tasks or unforeseen circumstances. During troop visits, we collected data on troop composition (see below) and individual behaviour *ad libitum*.

Individuals' ages were known from observing births or from patterns of dentition wear following capture. Individuals' relative ranks were calculated annually from *ad libitum* recorded dominance interactions in the package Matman 1.1.4 (Noldus Information Technology, 2013). Absolute ranks were converted to a relative scale ranging from 0 (lowest rank) to 1 (highest rank), to control for group size, using the formula *1-((1-r)/(1-n))*, with *r* being individual's absolute rank ranging from 1 to the total group size *n*. Group size was calculated as the total number of individuals in the troop for a particular year. Individuals who were present for less than half of a field season (due to emigration or death) were not included.

Troop composition and reproductive state data

Each day that the troop was contacted, a census was completed that recorded the identities of the individuals present and the reproductive states of each adult female. Females' reproductive states were recorded as: (1) pregnant (determined *post hoc* based on lack of resumption of swelling, reddening of the paracallosal skin and subsequent birth); (2) lactating (i.e., period following the birth of an infant until cycle resumption); (3) oestrous/swollen (i.e., exhibiting periovulatory swelling of the anogenital region); (4) non-swollen (i.e., deturgescent but not pregnant). For females in oestrus, we also recorded swelling sizes on a semi-quantitative scale from 0 (smallest) to 4 (largest), in order to capture within-individual

variation in swelling size across successful oestrus cycles [50]. As mentioned above, baboon
troops were not followed daily, and as such there were some missing data during some
females' cycles. We discarded any observations of the entire oestrous cycle where the start or
end of the swelling duration was missing and thus could introduce inaccuracy.

For females in oestrus, swelling duration was calculated as the number of continuous
days that a female was recorded with a swelling of any size. Because different mechanisms
could act on the different parts of the cycle, in addition to swelling durations, we calculated
the MSD as the number of days the focal female exhibited the largest swelling size during
that respective cycle. In summary, we used 157 receptivity observations for swelling duration
from 46 females (median number of swellings / female = 5, range = 1-14) and 150 receptivity
observations for maximal-swelling duration (median number of swellings / female = 5, range
= 1-14). The difference between the number of observations of these two variables stems
from the fact that for seven observations of swelling duration the start and end date of
receptivity were accurately known, yet this was not the case for the maximal-swelling
duration of those cycles.

We calculated eight predictor variables describing aspects of the social environment:
to consider (H1) female-female competition for paternal care, we calculated (1) the number
of pregnant and (2) of lactating females, as well as (3) their sum (PL), during the time a focal
female was started swelling. To consider (H2a) competition for females' access to the
preferred male, for the first day of a focal female's swelling we calculated the number of
simultaneously swollen females that were (4) not maximally-swollen and (5) maximally-
swollen. To consider sexual conflict in terms of (H2b) infanticide risk and (H2c) sexual
coercion, we calculated the (6) operational sex ratio (OSR) as the ratio of sexually active
females (i.e., swollen at any level) to adult males and the (7) the number of in-group adult
males. Finally, to account for food and resource competition we included group size in our
models (8) if adding this covariate improved model performance.

Statistical analyses

All statistical analyses were performed using R (version 4.0.3, 2020-10-10).

Question 1: Repeatability of receptivity

To determine how much variation in females' receptivity part of the cycle was due to
differences between individuals, we calculated the repeatability (R) of the SD and MSD of all

the females that had at least two measurements for each response (n = 38). R is calculated as

$\frac{\sigma_\alpha^2}{\sigma_\alpha^2 + \sigma_\varepsilon^2}$, where σ_α^2 represents the between-group variance, σ_ε^2 the within-group variance

($\sigma_\alpha^2 + \sigma_\varepsilon^2$ = the total phenotypic variance, V_p [51]). A relatively high R estimate would

indicate that the dependent variable exhibits high between- and low within-female variance;

low values of R indicate traits with high within-female or low between-female variation [51].

We calculated R using the `rpt` function (`rptR` package), which can calculate repeatability

for Poisson-distributed data, and the models included female identity and troop as random

effects. Finally, both models controlled for the possible confounding fixed effects [52] of age

and rank. The permutations and bootstrapping were set to 1000 and confidence intervals at

95%. However, we are aware that the repeatability we present in the results might be

somewhat flawed as our response variables are overdispersed (the `rptR` package offers no

way to correct for that excess variance though).

Question 2: Social determinants of oestrous length

To determine whether the social environment was correlated with females'

fluctuations in sexual receptivity duration, we used two generalised linear mixed models

(GLMMs; `glmer` function, `lme4` package [53]) with the response variables: swelling

duration (SD) and maximal-swelling duration (MSD). Both SD and MSD were modelled

using a negative binomial distribution to correct for overdispersion. Female identity nested in

troop identity was included as a random effect for SD models and female identity was a

random effect for MSD models, since variation between troops was absent for MSD. In both

models, we included rank and age as control variables. To control for possible type I errors

[54], we tested whether a model with random slopes for each of our predictors performed

better than one without (Supp. Mat. Tables A2 & A3), using the corrected AIC criterion,

AICc to compare models (`aictab` function, `AICcmodavg` package [55]). We included

this analysis as some authors argue that random slopes are crucial to avoid false positive rates

[56,57]. We also performed full-null model comparisons using a likelihood-ratio test. The

null models contained only the control variables age and relative rank and, as random factors,

female ID nested in troop for SD and female ID for MSD (see above). In both cases, the full

models performed better than the null model (Supp. Mat. Table A4), so we proceeded with

these full models for our analysis. Residuals and Q-Q plots were also checked for normality

(for model diagnostics see Supp. Mat. Figures A1-A4).

We used variables to test our hypotheses. Firstly, we determined, with the use of
AICc (as above), whether group size—the number of adults and sub-adults present in a
troop—should be included as a control variable in the models and which predictors should be
used to test for the effect of pregnant and lactating consexuals on sexual receptivity duration
(H1). For group size, larger groups could result increased competition for limited resources
(i.e., feeding competition), which could impact females' condition and their swelling duration
[7]; while for H1 we did not have an *a priori* prediction of whether the number of pregnant
and of lactating females or the combined number (i.e., sum) of pregnant and lactating females
(PL) is a better predictor. The results of model selection led us to include pregnant and
lactating females separately for SD (but not group size), and the sum, as well as group size,
for MSD (Supp. Mat. Table A1). In both SD and MSD models, we included an interaction
with rank because rank usually determines who can harass whom. To explore the effect of the
number of receptive females on sexual receptivity duration (H2a), we used the number of not
maximally-swollen females when a focal female started swelling when the response was SD;
and the number of maximally-swollen females when a focal female started swelling when the
response was MSD. Thirdly, to determine if the risk of infanticide (H2b) or male-male
competition (part of H2c) predicts fluctuations in female sexual receptivity durations, we
included the OSR as a predictor. Lastly, to determine the response of sexual receptivity
durations to male-male competition (part of H2c), we included the number of in-group males
because it is a proxy of social instability in the sense that male ranks are more volatile when
there is more competition (i.e., more males) (A. Carter, E. Huchard, pers. obs.). Lastly, we
tested whether group size should be included as a control variable.

The best-fit model for SD and MSD, carrying 97% and 69% of the cumulative model
weight (AICcWt), respectively, and showing the lowest AICc values, did not include random
slopes. For the two resulting models we checked for multicollinearity of model terms using
variance inflation factors (VIFs; Supp. Mat. Figures A2 & A4 respectively;
`check_collinearity` function, `performance` package [58], a VIF < 5 indicates a
low correlation of the predictor with other predictors and a VIF > 10 a high correlation [59]).
All VIFs were < 5, except interaction terms, which is expected [60], so these variables were
retained.
285
286 **Results**

287   Females were swollen for an average (median) of 22 days (range 8-44) and
288 maximally-swollen for a median of 6 days (range 1-24). There were no differences between
289 troops (Mann-Whitney U test: SD, W = 3481.5, p = 0.10; MSD, W = 2651.5, p = 0.59).
290
291 *Question 1: Repeatability of receptivity*
292   Neither SD nor MSD were repeatable at the individual level (SD: R = 0.036, SE =
293 0.049, CI = [0, 0.167], p = 0.251 & MSD: R = 0.054, SE = 0.058, CI = [0, 0.2], p = 0.283)
294 (Supp. Mat. Table A5; Figure A5).
295
Question 2: Social determinants of SD & MSD
We included two control variables in both our models, while for MSD we also
included group size as a predictor of feeding and contest competition (Table 2). Rank was not
significantly associated with either of our responses (SD: β = 0.25, SE = 0.26, p > 0.05;
MSD: β = 1.02, SE = 0.62, p > 0.05; Figure 1), and neither was age (for SD, β = -0.01, SE =
0.004, p = 0.069; for MSD, β = -0.02, SE = 0.01, p = 0.056; Figure 1). Group size, on the
other hand, predicted a decrease in MSD (β = -0.18, SE = 0.07, p = 0.013; Figure 1; Figure 2;
Table 2). Specifically, a one-unit increase in group size negatively correlated with a drop of
almost 16% in MSD (approximately 1.22 fewer days of MSD for each additional group
member).
306

H1: Intrasexual selection

Avoidance of female aggression hypothesis

Under our first hypothesis, we tested whether the number of pregnant and lactating
females was associated negatively with females' SD or MSD. There were on average 10
(median, range = 3-28) pregnant and lactating females (PL), 5 pregnant (median, range = 0-
11) as well as 6 lactating (median, range = 0-18) during females' swelling and maximal-
swelling phase. There was no relationship between the latter two variables and SD (pregnant:
β = -0.01, SE = 0.02, p > 0.05; lactating: β = 0.01, SE = 0.02, p > 0.05) or between PL and
MSD (β = 0.04, SE = 0.04, p > 0.05; summary of statistics at Table 2; Figure 1). There was
also no effect of the interaction of these variables with rank.

H2: Intersexual selection and sexual conflict

H2a Paternity concentration hypothesis

To test our second hypothesis, we tested whether there was positive association
between SD and MSD and the number of swollen females. There were on average 2 (median,
range = 0-6) not maximally-swollen females and 0 (median, range = 0-3) maximally-swollen
females when a focal female started swelling. Neither response was associated with swelling
durations (SD, not-maximally-swollen females: β = -0.01, SE = 0.02, p > 0.05; MSD,
maximally-swollen females: β = 0.08, SE = 0.07, p > 0.05; Table 2; Figure 1).

H2b Paternity confusion hypothesis
We predicted a positive correlation between females' SD or MSD and a female-biased
OSR. The mean OSR value was 0.64 (range 0.1–2.5): on average there were more sexually
active males than females. Our analyses showed no correlation between OSR and SD (β =
0.003, SE = 0.09, p > 0.05) or MSD (β = -0.26, SE = 0.17, p > 0.05; Table 2; Figure 1).

Figure 1**:** Forest plot of standardised estimates of the models with swelling duration (red) and maximal-swelling
duration (blue) as response variables. The level of significance is indicated by an unfilled circle (p > 0.05) or a
filled circle (p < 0.05). Only group size significantly predicted a decreased maximal-swelling duration.

H2c Avoidance of male coercion hypothesis

336   We tested whether there was a negative correlation between the number of males and
337 receptivity durations. There were on average 5 males (median, range = 1-10) when females
338 started to swell. We found support for this hypothesis either (SD: β = -0.01, SE = 0.02, p >
339 0.05; MSD: β = -0.04, SE = 0.03, p > 0.05; Figure 1; Table 2).

340
Figure 2**:** The effect of group size on maximal-swelling duration. The line indicates the predicted effect by
varying the respective focal variable and by setting all the other covariates to their mean. The shaded area
represents the 95% CI.

Table 2

Results of the two models with SD and MSD as response variables. "Log-Mean" measures the effect of the

specific predictor on the response variable and is given in log-scale. "SE" indicates the standard error and CI

shows the 95% confidence intervals. Significant results are indicated with bold.

Coefficients	Model 1 (SD)				Model 2 (MSD)			
	Log-mean of estimates	SE	CI (95%)	p-value	Log-mean of estimates	SE	CI (95%)	p-value
Intercept	3.28	0.19	2.90 – 3.66	**<0.001**	2.00	0.55	0.92 – 3.07	**<0.001**
Rank	0.25	0.26	-0.26 – 0.76	0.330	1.02	0.62	-0.20 – 2.23	0.100
Number of pregnant and lactating females	—	—	—	—	0.04	0.04	-0.04 – 0.11	0.314
Rank*Number of pregnant and lactating females	—	—	—	—	-0.08	0.06	-0.20 – 0.03	0.143
Rank*Number of pregnant females	-0.03	0.03	-0.07 – 0.02	0.317	—	—	—	—
Rank*Number of lactating females	-0.02	0.03	-0.08 – 0.03	0.444	—	—	—	—
Number of lactating females	0.01	0.02	-0.02 – 0.04	0.524	—	—	—	—
Number of pregnant females	-0.01	0.02	-0.04 – 0.02	0.483		—	—	—
Age	-0.01	0.004	-0.02 – 0.00	0.069	-0.02	0.01	-0.04 – 0.00	0.056
Number of not maximally-swollen ♀	-0.01	0.02	-0.06 – 0.04	0.681	—	—	—	—
Number of maximally-swollen ♀	—	—	—	—	0.08	0.07	-0.06 – 0.23	0.271
Number of ♂	-0.01	0.02	-0.04 – 0.02	0.627	-0.04	0.03	-0.10 – 0.03	0.261
OSR (♀/♂)	0.003	0.09	-0.17 – 0.18	0.976	-0.26	0.17	-0.60 – 0.08	0.141
Group size	—	—	—	—	-0.18	0.07	-0.33 – -0.04	**0.013**

Discussion

350   The present study investigated the variation in the length of receptivity within versus

351 across females. We provided a theoretical framework for why females could evolve to

352 manipulate their receptivity—a mechanism we termed cycle length manipulation. We further

tested whether variation in female sexual receptivity could be associated with changes in the social environment. We detected high within-female variation in the length of sexual receptivity. However, we found no evidence that variation in females' sexual receptivity is strategically adjusted to decrease intra- or inter-sexual competition. We did, however, find that group size, and presumably within-group competition, negatively affected females' maximal swelling durations.

We found that SD and MSD were not repeatable. Our results do not mean that there are no between-individual differences in SD and MSD, but that those differences are not large relative to within-individual differences and not consistent between females. Our findings show that cycle phases are labile traits that females could potentially manipulate. At a comparative level, high variation in oestrous duration has been detected in animals (e.g., captive jaguars, with a mean duration of 10.42 and a range of 7 to 15 days [61]). Women, in contrast, show repeatable cycle lengths, even though there is some variation within [62] and across women [63], as well as extensive variation in average cycle length across human populations [64]. It may be that baboons' low repeatability of receptivity reflects the harsh environmental conditions at Tsaobis, with females' condition fluctuating in response.

We found no evidence that the numbers of pregnant and lactating consexuals correlated with receptivity duration. One reason why our predictor did not affect receptivity duration could be that female reproductive suppression in the form of aggression was given by the female friends of a particular male [12]. Specifically, aggression was initiated by pregnant and lactating females who were friends of the male that was mate-guarding the receptive female. A more targeted approach that considers as a predictor the number of female friends a swollen female's mate-guarding male has might provide evidence for cycle length manipulation. Alternatively, it may be that the particular stage of lactation and pregnancy might be a better predictor than just the total number of individuals in each phase. For example, a study in yellow baboons revealed an increasing attack rate initiated from lactating females as lactation stages progressed [19].

We predicted that higher numbers of receptive females would prompt an elongation of oestrous duration, so that females could concentrate paternity in the alpha male but found no support for this. At least two explanations may account for this absence of evidence. First, because female chacma baboons are non-seasonal breeders [13], oestrous synchrony in this population is low, with a median of two simultaneously swollen females (i.e., swollen at any

phase of their cycle). Consequently, mating with the preferred male is possible for most cycling females and therefore elongation of the fertile period is not necessary. Second, the substantial aggression among cycling females in Tsaobis may limit females' propensity to extend their receptivity period [9,13]. This is predominantly true for lower-ranking oestrous females and females who are not mate-guarded, which tend to be the usual target of aggression from higher-ranking females.

We found no support for hypothesis 2b, that there would be a positive association between an OSR with receptivity durations. Our results could suggest that paternity dilution through extended receptivity comes at a cost for females, which we believe is an increased rate of female-female competition and aggression. Under this scenario, it seems that females entering the mating pool are primarily aggressive towards consexuals at peak swelling, which are most likely monopolised by the dominant male.

The number of adult males, as well as the OSR, were not associated with receptivity durations, as suggested by H2c. While male aggression, intimidation and coercion are costly for females in this population [26,27], it could be that a decrease in the length of swelling might incur higher costs, such as a smaller swelling, which could deter higher-ranking males from mate-guarding. Alternatively, selection to elongate receptivity due to increased infanticide risk (seen at a comparative level across Cercopithecine primates [36]) but also to decrease receptivity to avoid coercion could result in no effect overall.

We found that group size correlated negatively with MSD. There are at least two possible explanations for this relationship that are not mutually exclusive and could operate simultaneously. Firstly, as group size increases, competition for limited resources intensifies and maintaining energetically-costly maximal swellings could become more difficult for females [7]. In addition, larger baboon groups tend to travel longer distances over greater areas, which could generate higher energetic demands [65]. This interpretation recognizes the energetic limitation that arises due to elevated levels of feeding competition. Secondly, in baboons, socially-induced stress may generate higher cortisol levels [66] and subsequently result in a decrease of the swollen period [33,35]. As the number of in-group individuals increases, receptive females might be subject to higher aggression rates, both from consexuals (intrasexual aggression) and heterosexuals (intersexual aggression). Female primates of larger groups tend to show higher fecal glucocorticoid concentrations (fGC) [65,67], which could decrease females' receptivity durations indirectly. Under this scenario, social stress is interlinked with social competition, and future studies can directly test for that by including hormonal data and behavioural observations. Group size, however, was not a

good predictor in the case of SD. This might be because (1) SD is comparatively less costly
to maintain, even under stress, compared to MSD; or/and (2) competition with females and
males is comparatively less during the whole SD compared to MSD.

In summary, this study correlated variation in the length of female sexual receptivity
with aspects of the social environment that may influence the intensity of female reproductive
competition and intersexual conflict. We found no support for sexual conflict or female-
female competition driving receptivity duration. We did, however, find that one aspect of the
social environment—group size, and presumably within-group competition—correlated with
MSD in females. This suggests that maintaining large swellings is costly for females. Further
investigations may benefit from including behavioral and hormonal data in order to integrate
more closely the proximate mechanisms with ultimate explanations when testing this or
related hypotheses.

Ethics

Our research procedures were evaluated and approved by the Ethics Committee of the
Zoological Society of London and the Ministry of Environment and Tourism, Namibia (MET
Research/ Collecting Permits 886/2005, 1039/2006, 1186/2007, 1302/2008, 1379/2009,
1486/2010, 1486/2011, 1696/2012, 1786/2013, 1892/ 2014, 2009/2015, 2147/2016,
2303/2017, RPIV00392018/2019) and adhered to the ASAB/ABS Guidelines for the
Treatment of Animals in Behavioural Research and Teaching.

Data accessibility

Authors' contributions

440 AJC, EH and FD contributed to the conceptualization. AJC, EH designed the study. GC
441 established the study system and oversaw data collection, with assistance from AJC and EH.
442 FD performed the statistical analyses. All authors contributed to draft the manuscript. All
443 authors gave final approval for publication and agreed to be held accountable for the work
444 performed therein.

Competing interests

446 We declare we have no competing interests.

Funding

Long-term data collection was partially funded by grants from: the Agence Nationale de la
Recherche (ANR ERS-17-CE02-0008, 2018-2021) awarded to EH; a Fenner School of
Environment and Society Scholarship awarded to A.J.C., a Ministère de l'Education et de la

Recherche Studentship awarded to E.H; a NERC Advanced Research Fellowship awarded to

G.C.; Leakey Foundation grants awarded to AJC; and a Templeton World Charity

Foundation grant (TWCF0502) awarded to AJC. This paper is a contribution ISEM (n° TBD)

and a publication of the ZSL Institute of Zoology's Tsaobis Baboon Project, supported by

Research England.

Acknowledgments

We would like to thank the past Tsaobis Baboon Project leaders, students, volunteers and

interns for contributing to the long-term data collection at the field site. This research was

carried out with the permission of the Ministry of Environment and Tourism, the Ministry of

Land Reform, and the National Commission on Research, Science, and Technology. We

further thank the Tsaobis beneficiaries for permission to work at Tsaobis, the Gobabeb

Namib Research Institute for affiliation, and Johan Venter and the Snyman and Wittreich

families for permission to work on their land. Some findings formed part of a dissertation

submitted in partial fulfilment of the requirements of the degree of MSc of the University

College London in September 2021. This paper is a publication of the ZSL Institute of

Zoology's Tsaobis Baboon Project. Contribution ISEM n°202x-xx.

References

1. Kleindorfer S, Wasser SK. 2004 Infant handling and mortality in yellow baboons (Papio
cynocephalus): evidence for female reproductive competition? *Behav Ecol Sociobiol* **56**.
(doi:10.1007/s00265-004-0798-1)

2. Drea CM. 2005 Bateman Revisited: The Reproductive Tactics of Female Primates1.
Integrative and Comparative Biology **45**, 915–923. (doi:10.1093/icb/45.5.915)

3. Stumpf RM, Martinez-Mota R, Milich KM, Righini N, Shattuck MR. 2011 Sexual conflict
in primates. *Evol. Anthropol.* **20**, 62–75. (doi:10.1002/evan.20297)

4. Tang-Martínez Z. 2016 Rethinking Bateman's Principles: Challenging Persistent Myths of
Sexually Reluctant Females and Promiscuous Males. *The Journal of Sex Research* **53**,
532–559. (doi:10.1080/00224499.2016.1150938)

5. Clutton-Brock T. 2007 Sexual Selection in Males and Females. *Science* **318**, 1882–1885.
(doi:10.1126/science.1133311)

6. Clutton-Brock T. 2009 Sexual selection in females. *Animal Behaviour* **77**, 3–11.
(doi:10.1016/j.anbehav.2008.08.026)

7. Huchard E, Courtiol A, Benavides JA, Knapp LA, Raymond M, Cowlishaw G. 2009 Can
fertility signals lead to quality signals? Insights from the evolution of primate sexual
swellings. *Proc. R. Soc. B.* **276**, 1889–1897. (doi:10.1098/rspb.2008.1923)

8. Vaillancourt T. 2013 Do human females use indirect aggression as an intrasexual
competition strategy? *Phil. Trans. R. Soc. B* **368**, 20130080.
(doi:10.1098/rstb.2013.0080)

9. Huchard E, Cowlishaw G. 2011 Female–female aggression around mating: an extra cost of
sociality in a multimale primate society. *Behavioral Ecology* **22**, 1003–1011.
(doi:10.1093/beheco/arr083)

10. Tobler R, Pledger S, Linklater W. 2010 No evidence for ovarian synchrony or
asynchrony in hamadryas baboons. *Animal Behaviour* **80**, 829–837.
(doi:10.1016/j.anbehav.2010.07.018)

11. Kappeler PM, Fichtel C. 2012 Female reproductive competition in Eulemur rufifrons:
eviction and reproductive restraint in a plurally breeding Malagasy primate. *Mol Ecol* **21**,
685–698. (doi:10.1111/j.1365-294X.2011.05255.x)

12. Baniel A, Cowlishaw G, Huchard E. 2018 Jealous females? Female competition and
reproductive suppression in a wild promiscuous primate. *Proc. R. Soc. B.* **285**, 20181332.
(doi:10.1098/rspb.2018.1332)

13. Baniel A, Cowlishaw G, Huchard E. 2018 Context dependence of female reproductive
competition in wild chacma baboons. *Animal Behaviour* **139**, 37–49.
(doi:10.1016/j.anbehav.2018.03.001)

14. Watson NL, Simmons LW. 2010 Reproductive competition promotes the evolution of
female weaponry. *Proc. R. Soc. B.* **277**, 2035–2040. (doi:10.1098/rspb.2009.2335)

15. Bro-Jørgensen J. 2002 Overt female mate competition and preference for central males in
a lekking antelope. *Proceedings of the National Academy of Sciences* **99**, 9290–9293.
(doi:10.1073/pnas.142125899)

16. Robinson MR, Kruuk LEB. 2007 Function of weaponry in females: the use of horns in
intrasexual competition for resources in female Soay sheep. *Biol. Lett.* **3**, 651–654.
(doi:10.1098/rsbl.2007.0278)

17. Stockley P, Bro-Jørgensen J. 2011 Female competition and its evolutionary consequences
in mammals. *Biol Rev Camb Philos Soc* **86**, 341–366. (doi:10.1111/j.1469-
185X.2010.00149.x)

18. Rosvall KA. 2011 Intrasexual competition in females: evidence for sexual selection?
Behavioral Ecology **22**, 1131–1140. (doi:10.1093/beheco/arr106)

19. Wasser SK, Starling AK. 1988 Proximate and ultimate causes of reproductive
suppression among female yellow baboons at Mikumi National Park, Tanzania. *Am. J.*
Primatol. **16**, 97–121. (doi:10.1002/ajp.1350160202)

20. Watson-Capps JJ. 2009 2 Evolution of Sexual Coercion with Respect to Sexual Selection
and Sexual Conflict Theory. In *2 Evolution of Sexual Coercion with Respect to Sexual*
Selection and Sexual Conflict Theory, pp. 23–41. Harvard University Press.
(doi:10.4159/9780674054349-002)

21. van Schaik CP, Pradhan GR, van Noordwijk MA. 2004 Mating conflict in primates:
infanticide, sexual harassment and female sexuality. In *Sexual Selection in Primates:*
New and Comparative Perspectives (eds CP van Schaik, PM Kappeler), pp. 131–150.
Cambridge: Cambridge University Press. (doi:10.1017/CBO9780511542459.010)

22. Qi X-G, Grueter CC, Fang G, Huang P-Z, Zhang J, Duan Y-M, Huang Z-P, Garber PA,
Li B-G. 2020 Multilevel societies facilitate infanticide avoidance through increased
extrapair matings. *Animal Behaviour* **161**, 127–137. (doi:10.1016/j.anbehav.2019.12.014)

23. Clutton-Brock TH, Parker GA. 1995 Sexual coercion in animal societies. *Animal*
Behaviour **49**, 1345–1365. (doi:10.1006/anbe.1995.0166)

24. Stumpf RM, Boesch C. 2010 Male aggression and sexual coercion in wild West African
chimpanzees, Pan troglodytes verus. *Animal Behaviour* **79**, 333–342.
(doi:10.1016/j.anbehav.2009.11.008)

25. Eldakar OT, Gallup AC. 2011 The Group-Level Consequences of Sexual Conflict in
Multigroup Populations. *PLOS ONE* **6**, e26451. (doi:10.1371/journal.pone.0026451)

26. Baniel A, Cowlishaw G, Huchard E. 2017 Male Violence and Sexual Intimidation in a
Wild Primate Society. *Current Biology* **27**, 2163-2168.e3.
(doi:10.1016/j.cub.2017.06.013)

27. Baniel A, Delaunay A, Cowlishaw G, Huchard E. 2019 Oestrous females avoid mating in
front of adult male bystanders in wild chacma baboons. *R. Soc. open sci.* **6**, 181009.
(doi:10.1098/rsos.181009)

28. Muller MN, Thompson ME, Kahlenberg SM, Wrangham RW. 2011 Sexual coercion by
male chimpanzees shows that female choice may be more apparent than real. *Behav Ecol*
Sociobiol **65**, 921–933. (doi:10.1007/s00265-010-1093-y)

29. Muller MN, Kahlenberg SM, Emery Thompson M, Wrangham RW. 2007 Male coercion
and the costs of promiscuous mating for female chimpanzees. *Proceedings of the Royal*
Society B: Biological Sciences **274**, 1009–1014. (doi:10.1098/rspb.2006.0206)

30. Parga JA, Henry AR. 2008 Male aggression during mating: evidence for sexual coercion
in a female dominant primate? *Am. J. Primatol.* **70**, 1187–1190. (doi:10.1002/ajp.20609)

31. Polo P, Hernández-Lloreda V, Colmenares F. 2014 Male Takeovers Are Reproductively
Costly to Females in Hamadryas Baboons: A Test of the Sexual Coercion Hypothesis.
PLoS ONE **9**, e90996. (doi:10.1371/journal.pone.0090996)

32. Beehner JC, Bergman TJ, Cheney DL, Seyfarth RM, Whitten PL. 2005 The effect of new
alpha males on female stress in free-ranging baboons. *Animal Behaviour* **69**, 1211–1221.
(doi:10.1016/j.anbehav.2004.08.014)

33. Wasser S. 1983 13 – Reproductive Competition and Cooperation among Female Yellow
Baboons*. (doi:10.1016/B978-0-12-735950-2.50018-9)

34. Dunbar RIM, Sharman M. 1983 Female Competition for Access to Males Affects Birth
Rate in Baboons. *Behavioral Ecology and Sociobiology* **13**, 157–159.

35. Rowell TE. 1970 Baboon menstrual cycles affected by social environment. *J Reprod*
Fertil **21**, 133–141. (doi:10.1530/jrf.0.0210133)

36. Nunn CL. 1999 The evolution of exaggerated sexual swellings in primates and the
graded-signal hypothesis. *Animal Behaviour* **58**, 229–246. (doi:10.1006/anbe.1999.1159)

37. Zinner D, Deschner T. 2000 Sexual swellings in female hamadryas baboons after male
take-overs: "Deceptive" swellings as a possible female counter-strategy against
infanticide. *American Journal of Primatology* **52**, 157–168. (doi:10.1002/1098-
2345(200012)52:4<157::AID-AJP1>3.0.CO;2-L)

38. Vandenbergh JG. 1967 Effect of the presence of a male on the sexual maturation of
female mice. *Endocrinology* **81**, 345–349. (doi:10.1210/endo-81-2-345)

39. Lu A, Feder JA, Snyder-Mackler N, Bergman TJ, Beehner JC. 2021 Male-Mediated
Maturation in Wild Geladas. *Current Biology* **31**, 214-219.e2.
(doi:10.1016/j.cub.2020.10.003)

40. Bruce HM. 1959 An Exteroceptive Block to Pregnancy in the Mouse. *Nature* **184**, 105–
105. (doi:10.1038/184105a0)

41. Roberts EK, Lu A, Bergman TJ, Beehner JC. 2012 A Bruce Effect in Wild Geladas.
Science **335**, 1222–1225. (doi:10.1126/science.1213600)

42. Tinsley Johnson E, Snyder-Mackler N, Lu A, Bergman TJ, Beehner JC. 2018 Social and
ecological drivers of reproductive seasonality in geladas. *Behavioral Ecology* **29**, 574–
588. (doi:10.1093/beheco/ary008)

43. Palombit RA, Cheney DL, Seyfarth RM. 2001 Female–female competition for male
'friends' in wild chacma baboons (Papio cynocephalus ursinus). *Animal Behaviour* **61**,
1159–1171. (doi:10.1006/anbe.2000.1690)

44. Cheney DL, Silk JB, Seyfarth RM. 2012 Evidence for intrasexual selection in wild
female baboons. *Animal Behaviour* **84**, 21–27. (doi:10.1016/j.anbehav.2012.03.010)

45. Clutton□Brock T, Huchard E. 2013 Social competition and its consequences in female
mammals. *J Zool* **289**, 151–171. (doi:10.1111/jzo.12023)

46. Baniel A, Cowlishaw G, Huchard E. 2016 Stability and strength of male-female
associations in a promiscuous primate society. *Behav Ecol Sociobiol* **70**, 761–775.
(doi:10.1007/s00265-016-2100-8)

47. Huchard E, Alvergne A, Féjan D, Knapp LA, Cowlishaw G, Raymond M. 2010 More
than friends? Behavioural and genetic aspects of heterosexual associations in wild
chacma baboons. *Behav Ecol Sociobiol* **64**, 769–781. (doi:10.1007/s00265-009-0894-3)

48. Carter AJ, Baniel A, Cowlishaw G, Huchard E. 2020 Baboon thanatology: responses of
filial and non-filial group members to infants' corpses. *R. Soc. open sci.* **7**, 192206.
(doi:10.1098/rsos.192206)

49. Carter AJ, Torrents Ticó M, Cowlishaw G. 2016 Sequential phenotypic constraints on
social information use in wild baboons. *eLife* **5**, e13125. (doi:10.7554/eLife.13125)

50. Gesquiere LR, Wango EO, Alberts SC, Altmann J. 2007 Mechanisms of sexual selection:
sexual swellings and estrogen concentrations as fertility indicators and cues for male
consort decisions in wild baboons. *Horm Behav* **51**, 114–125.
(doi:10.1016/j.yhbeh.2006.08.010)

51. Nakagawa S, Schielzeth H. 2010 Repeatability for Gaussian and non-Gaussian data: a
practical guide for biologists. *Biological Reviews* , no-no. (doi:10.1111/j.1469-
185X.2010.00141.x)

52. Stoffel MA, Nakagawa S, Schielzeth H. 2017 rptR: repeatability estimation and variance
decomposition by generalized linear mixed□effects models. *Methods Ecol Evol* **8**, 1639–
1644. (doi:10.1111/2041-210X.12797)

53. Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting Linear Mixed-Effects Models
Using **lme4**. *J. Stat. Soft.* **67**. (doi:10.18637/jss.v067.i01)

54. Barr DJ, Levy R, Scheepers C, Tily HJ. 2013 Random effects structure for confirmatory
hypothesis testing: Keep it maximal. *Journal of memory and language* **68**.
(doi:10.1016/j.jml.2012.11.001)

55. Mazerolle MJ. 2020 AICcmodavg: Model Selection and Multimodel Inference Based on
(Q)AIC(c).

56. Bates D, Kliegl R, Vasishth S, Baayen H. 2018 Parsimonious Mixed Models.
(doi:10.48550/arXiv.1506.04967)

57. Matuschek H, Kliegl R, Vasishth S, Baayen H, Bates D. 2017 Balancing Type I error and
power in linear mixed models. *Journal of Memory and Language* **94**, 305–315.
(doi:10.1016/j.jml.2017.01.001)

58. Lüdecke D, Ben-Shachar MS, Patil I, Waggoner P, Makowski D. 2021 performance: An
R Package for Assessment, Comparison and Testing of Statistical Models. *Journal of
Open Source Software* **6**, 3139. (doi:10.21105/joss.03139)

59. James G, Witten D, Hastie T, Tibshirani R. 2013 *An Introduction to Statistical Learning*.
New York, NY: Springer New York. (doi:10.1007/978-1-4614-7138-7)

60. Francoeur RB. 2013 Could Sequential Residual Centering Resolve Low Sensitivity in
Moderated Regression? Simulations and Cancer Symptom Clusters. *Open Journal of
Statistics* **3**, 24–44. (doi:10.4236/ojs.2013.36A004)

61. Viau P, Rodini DC, Sobral G, Martins GS, Morato RG, de Oliveira CA. 2020 Puberty
and oestral cycle length in captive female jaguars Panthera onca. *Conservation
Physiology* **8**, coaa052. (doi:10.1093/conphys/coaa052)

62. Shea AA, Vitzthum VJ. 2020 The extent and causes of natural variation in menstrual
cycles: Integrating empirically-based models of ovarian cycling into research on
women's health. *Drug Discovery Today: Disease Models* **32**, 41–49.
(doi:10.1016/j.ddmod.2020.11.002)

63. Mihm M, Gangooly S, Muttukrishna S. 2011 The normal menstrual cycle in women.
Anim Reprod Sci **124**, 229–236. (doi:10.1016/j.anireprosci.2010.08.030)

64. Alvergne A, Högqvist Tabor V. 2018 Is Female Health Cyclical? Evolutionary
Perspectives on Menstruation. *Trends Ecol Evol* **33**, 399–414.
(doi:10.1016/j.tree.2018.03.006)

65. Markham AC, Gesquiere LR, Alberts SC, Altmann J. 2015 Optimal group size in a
highly social mammal. *Proceedings of the National Academy of Sciences* **112**, 14882–
14887. (doi:10.1073/pnas.1517794112)

66. Alberts SC, Sapolsky RM, Altmann J. 1992 Behavioral, endocrine, and immunological
correlates of immigration by an aggressive male into a natural primate group. *Horm
Behav* **26**, 167–178. (doi:10.1016/0018-506x(92)90040-3)

67. Stevenson PR, Castellanos MC. 2000 Feeding rates and daily path range of the
Colombian woolly monkeys as evidence for between- and within-group competition.
Folia Primatol (Basel) **71**, 399–408. (doi:10.1159/000052737)