GPCR Signaling Promotes a Stress-induced Genetic Program for Phenoptosis in

C. elegans

Changnan Wang[1], Yong Long[3], Bingying Wang[1], Chao Zhang[2], Dengke K. Ma[1, 4, *]

[1]Cardiovascular Research Institute and Department of Physiology, University of

California San Francisco, San Francisco, USA

[2]Department of Plastic and Reconstructive Surgery, Shanghai Institute of Precision

Medicine, Shanghai Ninth People's Hospital, Shanghai Jiao Tong University School of

Medicine, Shanghai, China

[3]State Key Laboratory of Freshwater Ecology and Biotechnology, Institute of

Hydrobiology, Chinese Academy of Sciences, Wuhan, China.

[4]Innovative Genomics Institute, University of California, Berkeley, USA.

* Correspondence: Dengke.Ma@ucsf.edu

Abstract

How an organism dies is a fundamental yet poorly understood question in biology. An organism can die of many causes, including stress-induced phenoptosis, also defined as organismic death that is regulated by its genome-encoded programs. The mechanism of stress-induced phenoptosis is still largely unknown. Here we show that transient but severe freezing-thaw stress (FTS) in *C. elegans* induces rapid and robust phenoptosis that is regulated by G-protein coupled receptor (GPCR) signaling. RNAi screens identify the GPCR-encoding *fshr-1* in mediating transcriptional responses to FTS. FSHR-1 increases ligand interaction upon FTS and activates a cyclic AMP-PKA cascade leading to a genetic program to drive phenoptosis, more so in aged than young populations. FSHR-1 signaling up-regulates the bZIP-type transcription factor ZIP-10, linking FTS to expression of genes involved in lipid remodeling, proteostasis and aging. Our studies reveal roles of FSHR-1/GPCR-mediated signaling in stress-induced gene expression and phenoptosis in *C. elegans*, providing empirical new insights into mechanisms of stress-induced phenoptosis with evolutionary implications.

INTRODUCTION

Organisms of all life forms on earth live with varying but limited lifespans. While the natural aging process and regulation of lifespan in many different organisms have been extensively studied (Antebi, 2007; Fontana and Partridge, 2015; Kenyon, 2010; López-Otín et al., 2013; Mutlu et al., 2021; Shore and Ruvkun, 2013), little is known about how

a multicellular organism dies at the end of its life after normal aging or in the middle of

its life. In nature, the death of an organism can have various causes, including severe

abiotic stress from ever-changing environments. Prolonged severe abiotic stresses can

lead to the death of organisms, in many cases likely because of failure in adapting to

the stress, or alternatively through stress-induced phenoptosis (Longo et al., 2005;

Skulachev, 1999, 2002, 2019). Using the nematode *C. elegans* as a model organism,

we have previously discovered that severe hypothermic stress (e.g. cold shock, CS)

promotes phenoptosis through a gene transcriptional pathway (Jiang et al., 2018).

Discovery of such rapid cold stress-induced phenoptosis and the genetic tractability of

C. elegans afford unprecedented access and opportunities to identify the underlying cell

signal transduction pathway and mechanisms of stress-induced phenoptosis.

With its small body size, *C. elegans* experiences varying temperature depending on the

ambient environment and exhibits stereotypic behavioral and cell physiological

response to chronic or acute mild (10-15 ^{o}C) hypothermia (Al-Fageeh and Smales,

2006; Garrity et al., 2010; Xiao et al., 2013). *C. elegans* can live and reproduce normally

within 15-25 ^{o}C and increases lifespan when grown under chronic mild hypothermia (15

^{o}C) (Lee et al., 2019; Xiao et al., 2013). Temperature beyond this range causes stress,

sterility and organismic death. By studying *C. elegans* mutants abnormally responding

to severe hypothermic (4 ^{o}C) stress stimuli, we identified ZIP-10, a bZIP-type

transcription factor that promotes cold stress-induced genes and phenoptosis (Jiang et

al., 2018). We showed that cold shock-induced ZIP-10 activates a genetic program and

up-regulates protease-encoding genes in intestinal cells to promote phenoptosis (Jiang

et al., 2018). Both the cold-induced *zip-10* expression and the death-promoting effect of

ZIP-10 appeared more prominent in old adults than larvae (Jiang et al., 2018). We

postulate that genetic programs underlying stress-induced phenoptosis may have

evolved because of antagonistic pleiotropic effects and/or "kin selection" at the

population level so that the biased phenoptosis of adult and aged animals after cold

stress would benefit young and reproductively more privileged populations to facilitate

the spreading of genes by young populations under resource-limiting and high-stress

conditions (Hamilton, 1963; Jiang et al., 2018; Smith, 1964). While evolutionary

mechanisms and significance of phenoptosis still remain debatable in the field (Galimov

et al., 2019; Kirkwood and Melov, 2011; Longo et al., 2005; Sapolsky, 2004; Skulachev,

2019), many empirical studies suggest that organisms with clonal population structures

such as *C. elegans* may indeed chronically age or die under stresses by mechanisms

(terminal investment, antagonistic pleiotropy, consumer sacrifice or disposable soma)

that can benefit its progeny or reproductive success of the organism itself (Ezcurra et

al., 2018; Galimov and Gems, 2020; Gulyas and Powell, 2021; Wu et al., 2022).

Despite critical roles of ZIP-10 in cold stress-induced phenoptosis in *C. elegans*, its

upstream regulators linking severe hypothermic stress to the ZIP-10-dependent genetic

program remain unidentified. By transcriptome profiling, RNAi screens and genotype-to-

phenotype mechanistic analysis, we uncovered a GPCR (FSHR-1, follicular stimulating

hormone receptor related) cascade that activates ZIP-10-dependent transcriptional

response to freeze-thaw stress (FTS). Activation of this pathway promotes organismic

phenoptosis under severe FTS conditions, more prominently in aged than young

animals. Our findings reveal an essential role of FSHR-1 and its GPCR signaling

cascade in stress-induced gene expression, providing mechanistic understanding of

phenoptosis and empirical evidence with implications for the evolutionary kin selection

theory and the "disposable soma" hypothesis of organismic aging.

RESULTS

RNAseq identifies transcriptomic changes to freeze-thaw stress in *C. elegans*

By exploring various severe hypothermic conditions in *C. elegans*, we identified an

environmental stress scheme (-20 °C freezing for 45 mins followed by recovery at 20 °C

for 24 hrs, FTS) that triggers robust ZIP-10-dependent phenoptosis with higher

penetrance than a previous scheme using 4 °C cold shock (Jiang et al., 2018) (Figure

1a). We measured the survival rates of both wild-type and *zip-10* mutants to quantify

organismic phenoptosis under such scheme for *C. elegans* at different developmental

stages (larval L1, L4, young and older adults). We found that wild-type animals

exhibited progressively increased rates of FTS-induced phenoptosis along with age,

with the young (48 hrs post L4) and older adult-stage (5 days post L4) populations

reaching nearly 100% death rates after FTS treatment (Figure 1b). By contrast, FTS-

induced phenoptosis was markedly attenuated in *zip-10* mutants at all stages. These

data establish a robust FTS scheme to induce rapid, age-dependent and environmental

stress-triggered phenoptosis in *C. elegans* and also confirm a critical role of ZIP-10 in

mediating severe hypothermic stress-induced phenoptosis.

To identify genes acutely regulated by FTS in *C. elegans* that may participate in

phenoptosis, we conducted RNA sequencing (RNAseq) of wild-type *C. elegans*

populations after 25-minute exposure to -20 °C freezing followed by recovery at 20°C

for 1 hr. We used this regime to identify genes that specifically and rapidly respond to

FTS rather than those that respond to general organismic deterioration once

phenoptosis occurs. After differential expression analyses of triplicate samples, we

identified 277 genes that are significantly up- or down-regulated by FTS conditions

(Figure 1c-d). Tissue enrichment analysis (TEA, Wormbase) revealed that intestine is

the most prominent site of gene regulation by FTS (Figure S1a). FTS-regulated genes

do not apparently overlap with those regulated by common stress-responding

transcription factors DAF-16 (nutritional stress), SKN-1 (oxidative stress), HSF-1 (heat

stress) or HIF-1 (hypoxic stress), and are involved in biological processes including lipid

remodeling (e.g. *oac-31*, *oac-14* and *oac-6*), responses to abiotic and pathogen

stresses (e.g. *kreg-1*, *irg-1* and *lys-3*) (Figure S1b-d). We generated transgenic *C.*

elegans strains in which GFP is driven by promoters of the top-ranked FTS-inducible

genes. *oac-31*p::GFP emerged as a robust FTS-inducible reporter with low baseline

expression and high-fold high-penetrance induction by FTS (Figure 1e). We confirmed

by quantitative RT-PCR (qRT-PCR) that FTS up-regulates endogenous *oac-31* in a ZIP-

10 dependent manner (Figure 1f). *oac-31* encodes a *C. elegans* homolog of sterol O-

acyltransferases and is also the most up-regulated member among the *oac* gene family

(Figure S1g). These results identify transcriptomic changes to FTS and led to *oac-*

*31*p::GFP as a robust transcriptional reporter for the genetic response to FTS.

RNAi screens identify FSHR-1/GPCR signaling in regulating responses to FTS

With an integrated *oac-31*p::GFP transgenic reporter strain, we performed RNAi (Figure

2a) and forward genetic screens to identify genes that are required for FTS induction of

*oac-31*p::GFP. For RNAi screen, we assembled a library of RNAi clones that target

genes with adequate intestinal expression (Transcript per million reads, TPM > 2.0) and

encoding transmembrane receptors or transcription factors representing a

comprehensive set of signal transduction pathways in *C. elegans* (Table S1; Figure

S2a). We found that two gene RNAi clones (*sbp-1* and *tra-1*) activated baseline *oac-*

*31*p::GFP, whereas another gene RNAi clone (*fshr-1*) blocked FTS-induced *oac-*

*31*p::GFP (Figure S2b-c). *fshr-1* encodes a glycoprotein hormone receptor homolog in

C. elegans and has been implicated in mediating oxidative and innate immune

responses (Cho et al., 2007; Kim and Sieburth, 2020; Kudo et al., 2000; Miller et al.,

2015; Powell et al., 2009). A genetic deletion allele *ok778* (back-crossed to N2 for 5

times) of *fshr-1* fully recapitulated the RNAi phenotype in blocking FTS-induced *oac-*

*31*p::GFP (Figure 2b). Neither FTS nor *fshr-1* RNAi affected *unc-54*p::mCherry, the

transgenic co-injection marker for *oac-31*p::GFP. By qRT-PCR, we confirmed that FTS

induction of *oac-31* (and also another gene *ncr-1* identified from RNAseq) requires *fshr-*

1 (Figure S2d). However, at least two other genes, *W09G12.7* and *tsp-1*, do not

apparently require *fshr-1* for induction by FTS (Figure S2e). These results reveal an

essential and specific role of *fshr-1* in transcriptional up-regulation of FTS-induced

genes including *oac-31*.

fshr-1 encodes the sole ortholog of glycoprotein hormone receptor family proteins in the

C. elegans genome (Cho et al., 2007; Kudo et al., 2000). FSHR-1 family proteins are

GPCRs that couple to the small G-protein Gαs to stimulate cAMP production and

downstream protein kinase A (PKA) signaling. We found that a gain-of-function (GOF)

mutation (Schade et al., 2005) in the *C. elegans* Gαs gene *gsa-1* strongly activated *oac-*

*31*p::GFP even without FTS (Figure 2c). GOF of the Gαs GSA-1 leads to cAMP

elevation and activation of PKA, which comprises the catalytic subunit KIN-1 and the

inhibitory regulatory subunit KIN-2 in *C. elegans* (Lu et al., 1990). We found that *kin-2*

RNAi led to constitutive *oac-31*p::GFP activation as *gsa-1* GOF did (Figure 2d), and

both phenotypes can be suppressed by *kin-1* but not *fshr-1* RNAi (Figure 2e). These

results indicate that FTS up-regulates *oac-31* via a classic FSHR-1/GPCR-mediated

cAMP and KIN-1/PKA signaling cascade (Figure 2f).

FSHR-1 acts in the intestine to regulate ZIP-10 and cold-induced phenoptosis

We next examined the site of expression/action of FSHR-1 and its physiological

consequences in FTS-induced phenoptosis. A transcriptional reporter with GFP driven

by the promoter of *fshr-1* revealed its predominant intestinal site of expression (Figure

3a). A translational reporter with mCherry driven by the promoter and the genomic

protein-coding sequence of *fshr-1* also revealed an intestinal pattern of FSHR-

1::mCherry (Figure 3b). Transgenic wild-type *fshr-1* driven by the intestine-specific *ges-*

1 promoter rescued *oac-31* induction by FTS in *fshr-1* mutants, whereas RNAi against

fshr-1 specifically in the intestine nearly abolished *oac-31* induction by FTS (Figure 3c).

FTS-induced *oac-31* also requires *zip-10* (Figure 1f), consistent with its role in mediating

transcriptional responses to cold stress in intestinal cells. We found that FTS strongly

induced *zip-10* expression in control RNAi but not intestine-specific *fshr-1* RNAi-treated

animals (Figure 3d). Although FTS transcriptionally up-regulates *zip-10* through FSHR-

1, high baseline expression of ZIP-10 likely also contributes to *oac-31* expression given

its rapid induction kinetics. These results indicate that FSHR-1 acts in the intestine to

promote FTS-induced *zip-10* and *oac-31* gene expression.

We next assessed the role of the FSHR-1 signaling cascade in FTS-induced

phenoptosis. Compared to wild type, *fshr-1* loss-of-function mutants showed markedly

increased rates of survival after FTS for both larvae and adults (Figure 3e). We

observed death-characteristic blue fluorescence after the FTS treatment in wild-type

animals as previously reported for naturally dying animals (Coburn et al., 2013), but

markedly less so in *zip-10* or *fshr-1* mutants (Figure 3f). In addition, FLR-2 is the sole *C.*

elegans ortholog of glycoprotein hormones and putative FSHR-1 ligand (Oishi et al.,

2009) that may signal through the FSHR/GSA-1/PKA cascade (Figure 2). Like *fshr-1*

deficient animals, loss of *flr-2* or *kin-1* caused defective *oac-31* induction by FTS (Figure

4a). To determine the epistatic relationship of the genes in the FLR-2/FSHR-1 pathway

for FTS-induced phenoptosis, we examined how their loss-of-function or gain-of-

function impacted FTS-induced phenoptosis at L4 stages, in which either positive or

negative effects on phenoptosis can be quantitatively measured (Figure 4b). Both *flr-2*

single and *flr-2; fshr-1* double loss-of-functions (by mutant or RNAi) showed defective

FTS-induced phenoptosis (Figure 4b). Consistent with their roles in regulation of *oac-31*

downstream of FSHR-1, *gsa-1* gain-of-function or *kin-2* loss-of-function exhibited

enhanced FTS-induced phenoptosis, while such enhancement can be suppressed with

zip-10 loss-of-function mutations (Figure 4b). Loss of ZIP-10 strongly attenuated FTS-

induced phenoptosis (Figure 1a), while individual loss of its target genes *oac-31* or *asp-*

17 (previously identified as a ZIP-10-dependent gene encoding a protease) decreased

FTS-induced phenoptosis to a lesser extent (Figure 4b). These results indicate that FTS

promotes phenoptosis through an FSHR-1/GSA-1/PKA/ZIP-10 cascade (Figure 4c).

FSHR-1 has been previously implicated in responses to oxidative and pathogen

stresses (Miller et al., 2015; Powell et al., 2009). Among the membrane lipids that can

be oxidized, cholesterol is most prone to oxidation by reactive oxygen species (Murphy

and Johnson, 2008). Consistent with previous studies supporting a role of FSHR-1 in

protecting against oxidative stress associated with cell membrane lipids, we found that

fshr-1 RNAi caused higher sensitivity than wild type to exposure of exogenous

cholesterol when supplemented with the oxidizing agent Paraquat (Figure S3a). In

addition, we found that the cold stress or FTS resilience phenotypes caused by RNAi

against *fshr-1* can be modulated by the amount of exogenous (supplemented in media)

or endogenous (manipulated by *ncr* expression) cholesterol availability (Figure S3b, c,

d). These results suggest that resilience to severe hypothermic stress can come with a

physiological trade-off in organismic resilience to cholesterol oxidative stresses.

We next explored how FTS may regulate FLR-2/FSHR-1. Based on RNAseq results,

FTS did not affect expression levels of *flr-2* or *fshr-1* (Figure S4a). While *flr-2* is

expressed predominantly in neurons based on single-cell RNA results (Taylor et al.,

2021), we generated a translational reporter for FLR-2 fused with GFP and identified its

localization to coelomocytes, which likely uptake secreted forms of FLR-2::GFP from

neurons (Figure S4b). However, *flr-2* reporters showed no apparent changes after FTS

treatment (Figure S4b). These data suggest that neuron-derived systemically-acting

FLR-2 is required for FTS induction of *oac-31* and phenoptosis, but not directly

regulated by FTS transcriptionally. Nonetheless, we observed that FTS induced marked

elevation of intracellular cAMP levels in wild type but not *flr-2* or *fshr-1* mutants (Figure

4d). In addition, co-immunoprecipitation experiments using transgenic animals show

that GFP-tagged FLR-2 can increasingly bind to mCherry-tagged FSHR-1 extracellular

domains after FTS in a time-dependent manner (Figure 4e), consistent with an increase

in cAMP levels after FTS. As glycoprotein hormones typically bind to its obligatory

GPCRs to stimulate cAMP levels, these results further support the notion that FSHR-1-

dependent GPCR signaling actively mediates response to FTS, rather than plays a

permissive role, to drive stress-induced gene expression and phenoptosis.

224 A mathematical model to explain plausible adaptive functions of phenoptosis

At first glance, existence of phenoptosis and a genetic pathway promoting phenoptosis

would seem incompatible with the evolutionary theory as genes promoting phenoptosis

would compromise individual fitness, thus not be subject to selection pressure. Why

would a genetic program be possibly evolved to promote phenoptosis? We established

a mathematical model (Figure 5) built upon the evolutionary concept of kin selection that

may explain adaptive or beneficial functions of phenoptosis at the population level. We

define three key parameters in the model that determine the growth of reproduction-

competent haploid genomes in a population, including the parameter of α (aging rate), β

(genome replication rate), and μ (cohort competition/infection cost after stress). More

stress sensitivity (i.e. phenoptosis susceptible) in old (v.s. young) animals would

decrease (v.s. increase) μ, since old animals contribute less than young animals in the

growth of replication-competent haploid genomes in the population. Simulated results

from our model assuming constant parameters in a given population show that the

number of healthy individuals in a starting population can increase faster than a

population without stress-inducible phenoptosis, under conditions where there is

competition for limited resources to reproduce or infectious spreading of pathogens

among individuals (Figure 5). According to the model, stress-induced phenoptosis may

act against the unfit group members (less reproductive old adults or sick individuals

shedding infectious pathogens) to benefit the whole group at the population level.

DISCUSSION

While genes can control how cells die in well-known processes including apoptosis,

necroptosis, and pyroptosis (Metzstein et al., 1998; Shi et al., 2017; Weinlich et al.,

2017), how genes might control the death of an organism remains largely unclear.

Based on comprehensive genetic analyses and stress-responding organismic

phenotypes, we propose a model in which FSHR-1 signaling may mediate FTS-induced

phenoptosis in *C. elegans*. After exposure to severe hypothermic stress conditions,

FLR-2 may promote FSHR-1 activation and signaling through a GPCR-dependent Gαs-

cAMP-PKA pathway, leading to increased expression of *zip-10*, *oac-31* and other genes

in a genetic program that coordinately drive phenoptosis. Site-of-expression and site-of-

action analyses indicate that this GPCR pathway operates in the intestine, consistent

with intestinal expression and cell-autonomous regulation of *zip-10* and *oac-31*.

Although the precise mechanism of how FTS activates FLR-2/FSHR-1 is unknown, it is

conceivable that sensory neurons may detect thermal stress signals to promote release

of FLR-2 to act systemically, while intestinal cells may also adjust membrane properties

upon FTS to modulate FLR-2/FSHR-1 activation (Ernst et al., 2016). Various types of

stress can trigger acute organismic death in *C. elegans* and have been postulated to

elicit systemic necrosis, yet by unknown mechanisms (Coburn et al., 2013; Ezcurra et

al., 2018). As ZIP-10 appears both required and partially sufficient to drive cold-induced

phenoptosis, it may represent a master transcriptional regulator downstream of stress-

sensing GPCR signaling, leading to systemic necrosis and organismic death through its

target gene effectors, including those encoding proteases we previously identified

(Jiang et al., 2018). Our results suggest that the ISY-1/*mir-60* axis plays a gatekeeping

role for ZIP-10 activation and phenoptosis, whereas the FLR-2/FSHR-1/GSA/KIN-2 axis

is likely instructive by linking FTS stress to genes driving phenoptosis or promoting

organismic sensitivity to lethal effects of FTS. FSHR-1 and ZIP-10 also mediate host

cell responses to oxidative stress, various types of pathogens and immune aging (Kato

et al., 2016; Lee et al., 2021; Miller et al., 2015; Powell et al., 2009). Indeed, we found

that *fshr-1* loss-of-function mutants, while being more FTS resistant than wild type, were

less tolerant of cholesterol oxidative stress (Figure S4). How FSHR-1 or ZIP-10 may

function in other stress paradigms similarly in promoting organismic phenoptosis or play

context-specific roles in physiological trade-off awaits further studies.

We note that *oac-31*, like several other *oac* genes (*oac-3, 6, 14, 20*) strongly up-

regulated by FTS (Figure S1g), encodes a *C. elegans* homolog of sterol O-

acyltransferase (SOAT). SOAT converts accessible cholesterol into ester forms for

storage in lipid droplets (Luo et al., 2020; Sevanian and Peterson, 1986). While the

biochemical function of OAC-31 remains further characterized, a recent study showed

that cold shock can promote lipid movement from the intestine to the germline (Gulyas

and Powell, 2021). Interestingly, natural aging of *C. elegans* is accompanied with

autophagy-mediated conversion of intestinal biomass into yolk, contributing to late-life

mortality (Ezcurra et al., 2018). Thus, it is plausible that FSHR-1-dependent up-

regulation of SOAT may be part of a genetic program that converts somatic biomass

into nutrient/energy reserves for trafficking to the germline to benefit the progeny at the

expense of parental death. In early life, such genetic programs may have evolved to

promote reproductive health, while exhibiting antagonistic pleiotropic effects promoting

organismic death/aging in late life. Phenoptosis-promoting effects of FSHR-1/ZIP-10 are

also dependent on stress types and severity, reflecting on their age-dependent

pleiotropic effects. At the population level, phenoptosis of relatively aged adults can also

be beneficial as limiting resources can be funneled to reproductively active younger

individuals under high stress conditions. These two scenarios ("disposable soma" and

"kin selection") are not mutually exclusive and may have been subject to selection

pressure and explain the conservation of phenoptosis and its pathway of regulation in

certain species. Together, our findings identify key signaling mediators of severe cold

stress-induced phenoptosis in *C. elegans* and provide empirical evidence of how

phenoptosis can be genetically regulated with plausible evolutionary significance.

MATERIALS AND METHODS

C. elegans strains

C. elegans strains were maintained with standard procedures unless otherwise

specified. The N2 Bristol strain was used as the reference wild type, and the

polymorphic Hawaiian strain CB4856 was used for genetic linkage mapping and SNP

analysis (Brenner, 1974; Davis et al., 2005). Forward genetic screen for *oac-31*p::GFP

mutants after ethyl methanesulfonate (EMS)-induced random mutagenesis was

performed as described previously (Ma et al., 2012, 2015). Feeding RNAi was

performed as previously described (Kamath and Ahringer, 2003). Customized sub

libraries of RNAi clones were constructed by manual selection from the main Ahringer

library based on gene expression abundance (TPM>2) in the intestine and encoded

proteins as putative transmembrane receptors, signaling proteins and transcription

factors (Table S2). Transgenic strains were generated by germline transformation as

described (Mello et al., 1991). Transgenic constructs were co-injected (at 10 - 50 ng/µl)

with dominant *unc-54p*::mCherry or *rol-6* markers, and stable extrachromosomal lines of

mCherry+ or roller animals were established. Genotypes of strains used are: *gsa-*

1(ce81) I, ncr-2(nr2023) III, fshr-1(ok778) V, flr-2(ut5) V, ncr-1(nr2022) X. dmaEx616

[fshr-1p::GFP; unc-54p::mCherry], vjEx1449[ges-1p::fshr-1], dmaEx617 [fshr-1p::fshr-

1::GFP; unc-54p::mCherry], dmaEx620 [rpl-28p::fshr-1ECD::mCherry], rde-1(ne219);

Is[Pges-1::RDE-1::unc54 3'UTR; Pmyo2::RFP3] (gift of Dr. Meng Wang's laboratory)*,*

dmaIs136 [flr-2p::flr-2::GFP], dmaIs117 IV [oac-31p::GFP; unc-54p::mCherry].

Sample preparation for RNA sequencing and data analysis

Control N2 animal were maintained at 20 °C. For freeze-thaw stress, N2 animals were

exposed to -20 °C for 25 mins followed by 1 hr recovery at 20 °C. Upon sample

collection, the animals were washed down from NGM plates using M9 solution and

subjected to RNA extraction using the RNeasy Mini Kit from Qiagen. 1 µg total RNA

from each sample was used for sequencing library construction. RNA-seq Library

preparation and data analysis were performed as previously described (Jiang et al.,

2018). Three biological replicates were included for each treatment. The libraries were

sequenced to 51 bp at single end by the Center for Advanced Technology (CAT) of the

University of California, San Francisco using a Hiseq-3000 system. The cleaned

RNAseq reads were mapped to the genome sequence of *C. elegans* using hisat2 (Kim

et al., 2015). The mapped reads were assigned to the genes using featureCounts (Liao

et al., 2014). Abundance of genes was expressed as RPKM (Reads per kilobase per

million mapped reads). Identification of differentially expressed genes was performed

using the DESeq2 package (Love et al., 2014).

Quantitative RT-PCR

25 μl pellet animals were resuspended in 400 μl lysis buffer of Quick-RNA MiniPrep kit

(Zymo Research, R1055) then lysed by TissueRuptor (Motor unit '8' for 1 min). Total

RNA was extracted following the instruction (Zymo Research, R1055). 1 μg

RNA/sample was reverse transcribed into cDNA (Thermo Fisher Scientific, 18080051).

Real-time PCR was performed by using Roche LightCycler96 (Roche, 05815916001)

system and SYBR Green (Thermo Fisher Scientific, FERK1081) as a dsDNA-specific

binding dye. qRT-PCR condition was set to 95°C for denaturation, followed by 45 cycles

of 10 s at 95°C, 10 s at 60°C, and 20 s at 72°C. Melting curve analysis was performed

after the final cycle to examine the specificity of primers in each reaction. Relative

mRNA was calculated by $\Delta\Delta$CT method and normalized to actin. Primers for qRT-PCR:

oac-31 (Forward, ATGGCTTAAATCGCTGGAAA; Reverse,

ATCTTTCGCCATCAATACGG), *W09G12.7*, Forward, CGAAGATCACTCGAACACGA;

Reverse, CCATTTTTACGAAGGCTGGA), *tsp-1* (Forward,

CTTTGATTGCCGTTGGATTT; Reverse, CCCAAAGAAAGGCCGATAAT), *act-3*

(Forward, TCCATCATGAAGTGCGACAT; Reverse, TAGATCCTCCGATCCAGACG).

Freezing resilience assay

Animals were cultured under non-starved conditions for at least 4 generations at 20 °C

before cold and freezing resilience assay. For freezing resilience assay, bleach-

synchronized L1, L4, young (48 hrs post L4) and older adult (5 days post L4)

populations were kept at -20 °C for 45 mins and then recovered for 24 hrs at 20 °C. For

such experiments, NGM plates spread with equal agar thickness seeded with equal

amounts of OP50 were used while hypothermic temperature readings were monitored

by thermometers to ensure minimal fluctuation. After freezing shock, animals were

moved to 20 °C for recovery and scored as dead if they showed no pumping and

movement upon light touch with the body necrosis subsequently confirmed.

cAMP assay

N2, *fshr-1* and *flr-2* mutants were freeze shocked (-20 °C) for 25 mins, followed by

recovery at 20 °C for 0, 1, 2 or 4 hrs. Animals were harvested and washed three times

with M9 and 20 µl pellet animals were lysed directly in ELISA lysis buffer. Then worm

extracts were collected by centrifugation. The cAMP levels in the supernatant were

determined using the cyclic AMP ELISA kit (Cayman Chemical, Ann Arbor, MI)

according to the manufacturer's instructions. BCA kit was used to measure the

concentration of protein as normalizing control.

Co-immunoprecipitation assay

Transgenic animals expressing *flr-2p::flr-2::GFP* and *fshr-1::ECD::mcherry* were

cultured under non-starved conditions at 20 °C on NGM plates before freezing

treatment. For freezing treatment, mixed stage worms were kept at -20 °C for 25 mins

and then recovery for 0, 1, 2, 4 hrs at 20 °C. Upon sample collection, the animals were

washed down from NGM plates using M9 solution and subjected to protein extraction

using ice-cold 2x lysis buffer (60 mM HEPES, pH = 7.4, 100 mM potassium chloride,

0.1% Triton X, 4 mM magnesium chloride, 10% glycerol, 2 mM DTT with RNase

inhibitor, protease inhibitor, and phosphatase inhibitors) and lysed with sonication. The

lysate was then centrifuged at 15,000 rpm for 15 mins to remove cell debris. The

supernatant was incubated with 4 µl of pre-clean magnetic beads at 4 °C for 15

minutes. To precipitate *fshr-1*::ECD::mcherry proteins, the sample was placed into a

1.5 ml RNase-free tube containing RFP-trap magnetic beads at 4 °C overnight. The

beads were then washed with 400 µl of lysis buffer for 5 times and incubated with *flr-*

2::flr-2::GFP lysate samples at 4 °C overnight. The beads were then washed with 400 µl

of lysis buffer for 5 times and resuspended in 40 µl of 1×Laemmli Sample Buffer. The

samples were then boiled at 95 °C for 10 minutes and subject to Western blot analysis.

Statistical analysis

Data were analyzed using GraphPad Prism 9.2.0 Software (Graphpad, San Diego, CA)

and presented as means ± S.D. unless otherwise specified, with *P* values calculated by

unpaired two-tailed t-tests (comparisons between two groups), one-way or two-way

ANOVA (comparisons across more than two groups) followed by Tukey's post-hoc tests

and adjusted with Bonferroni's corrections.

Data availability

The RNAseq read datasets were deposited in NCBI SRA (Sequence Read Archive)

under the BioProject accession PRJNA763790. All other data newly generated for this

study are included in this article.

Acknowledgments

Some strains were provided by the CGC, which is funded by NIH Office of Research

Infrastructure Programs (P40 OD010440), and from Drs. Derek Sieburth and Meng

Wang laboratories. The work was supported by NIH grant 1R35GM139618 and the

Packard Fellowship in Science and Engineering (D.K.M).

Author contributions

C.W., B.W. and D.K.M. designed, performed and analyzed the experiments and wrote

the manuscript. Y.L. performed RNA sequencing and analysis. D.K.M. supervised the

project.

Competing interests

The authors declare no competing interests.

Materials & Correspondence.

Correspondence and material requests should be addressed to Dengke K. Ma, Ph.D.

(dengke.ma@ucsf.edu).

Figures and Figure legends

Figure 1: RNAseq identifies transcriptomic changes to freeze-thaw in *C. elegans*.

(a), Schematic of experimental flow to assay FTS-induced phenoptosis. **(b)**,

Quantification of phenoptosis based on percentages of animals survived after FTS in

each condition (young adults, 2 days post L4; older adults, 5 days post L4). Values are

means ± S.D with ***$P < 0.001$ (one-way ANOVA with post-hoc Tukey HSD, N = 5

independent experiments, n > 50 for each experiment). **(c)**, Volcano plot showing

identified genes that are regulated by FTS based on RNAseq. Red: up-regulated. Blue:

down-regulated. **(d)**, Heat map and hierarchical clustering analysis showing the top 50-

ranked genes regulated by FTS. **(e)**, Representative epifluorescence images showing

*oac-31*p::GFP but not co-injection marker *unc-54*p::mCherry induction by FTS. **(f)**, qRT-

PCR showing induction of endogenous *oac-31* by FTS in wild type but attenuated in *zip-*

10(ok3462) loss-of-function mutants. Values are means ± S.D with ***$P < 0.001$ (two-

way ANOVA for genotype-effect interaction and post-hoc Tukey HSD, N = 3

independent experiments, n > 50 for each experiment). Scale bar, 100 μm.

Figure 2: Genetic screens identify FSHR-1/GPCR signaling in regulating freeze-

thaw induced *oac-31*. (a). Schematic of RNAi and EMS screens. **(b)**, Representative

epifluorescence images showing FTS induction of *oac-31*p::GFP is blocked in *fshr-1*

deletion mutants. **(c)**, Representative epifluorescence images showing constitutive

induction of *oac-31*p::GFP in *gsa-1* gain-of-function mutants without FTS. **(d)**,

Representative epifluorescence images showing constitutive induction of *oac-31*p::GFP

in *kin-2* RNAi-treated animals without FTS. **(e)**, Summary table listing the *oac-31*p::GFP

phenotype (with the numbers of "+" signs qualitatively indicating the GFP fluorescence

levels) and penetrance of FTS induction in animals with genotypes indicated. **(f)**,

Schematic of the FSHR-1/GSA/PKA pathway in regulating *oac-31* expression in

response to FTS, based on genetic epistasis analysis. Scale bar, 100 μm.

Figure 3: ***fshr-1* is expressed and acts in the intestine to regulate FTS induction of**

***oac-31* and phenoptosis. (a)**, Representative epifluorescence image of the

transcriptional *fshr-1*p::GFP reporter showing predominant expression of GFP in the

intestine. **(b)**, Representative epifluorescence image of the translational *fshr-1*p::*fshr-*

1::mCherry reporter showing intestinal expression pattern. **(c)**, Tissue-specific rescue

(*fshr-1* cDNA driven by the intestine-specific *ges-1* promoter) and *fshr-1* RNAi (in the

strain *rde-1(ne219); Is[Pges-1::RDE-1::unc54 3'UTR; Pmyo2::RFP3]*) analysis showing

fshr-1 acting primarily in the intestine for FTS induction of *oac-31*. Values are means ±

S.D with **$P < 0.01$ and *$P < 0.05$ (two-way ANOVA for genotype-effect interaction and

post-hoc Tukey HSD, N > 3 independent experiments, n > 50 for each experiment). **(d)**,

qRT-PCR results showing induction of *zip-10* by FTS was attenuated in animals with

intestine-specific *fshr-1* RNAi. Values are means ± S.D with **P < 0.01, ***$P < 0.001$

(two-way ANOVA for genotype-effect interaction and post-hoc Tukey HSD, N = 3

independent experiments, n > 50 for each experiment). **(e)**, Quantification of

phenoptosis based on percentages of larval (L1 and L4) and adult (YA: 48 hrs post L4,

OA: 5 days post L4) stage animals survived after FTS treatment. Values are means ±

S.D with **$P < 0.01$ and ***$P < 0.001$ (two-way ANOVA for genotype-effect interaction

and post-hoc Tukey HSD, N > 3 independent experiments, n > 50 for each experiment).

(f), Representative bright field and fluorescence images showing characteristic blue

"death fluorescence" in wild type but not *zip-10* or *fshr-1* mutants (***$P < 0.001$, one-way

ANOVA with post-hoc Tukey HSD, N = 10) 24 hrs after FTS. Scale bar, 100 μm.

Figure 4: FLR-2/FSHR-1-mediated GPCR/cAMP signaling drives phenoptosis. (a),

qRT-PCR results showing abolished *oac-31* induction by FTS in *flr-2* reduction-of-

function mutants, mimicking *fshr-1* mutants. **$P < 0.01$ (two-way ANOVA for genotype-

effect interaction and post-hoc Tukey HSD, N = 3 independent experiments, n > 50 for

each experiment). **(b)**, Table summary of percentages of larval L4-stage animals with

indicated genotypes survived after FTS treatment *$P < 0.05$ and ***$P < 0.001$ (one-way

ANOVA and post-hoc Tukey HSD N = 5 independent experiments, n > 50 for each

experiment). **(c)**, Schematic of the genetic pathway leading to FTS-induced phenoptosis

based on epistasis and gene expression analysis. Known genes or proteins promoting

(green) or antagonizing (red) cold stress-induced phenoptosis are noted and with other

contributors (dashed line). **(d)**, Normalized ELISA results showing FTS-induced (no FTS

control, 1 hr , 2hrs, 4 hrs of recovery after freezing) elevation of cAMP was abolished in

flr-2 and *fshr-1* mutants. Values are means ± S.D with ***P < 0.001 (two-way ANOVA

for genotype-effect interaction and post-hoc Tukey HSD, N > 3 independent

experiments, n > 50 for each experiment). N.S., non-significant. **(e)**, Representative

western blots of GFP-tagged FLR-2, mCherry-tagged FSHR-1::ECD(extracellular

domain)::mCherry from input and co-immunoprecipitation lysate samples showing time-

dependent (no FTS control, 0 hr, 1 hr, 2 hrs, and 4 hrs of recovery after freezing)

binding of FLR-2::GFP and FSHR-1::ECD::mCherry upon FTS treatment (tubulin as

loading control).

Figure 5: A simplified mathematical model to simulate potential adaptive function of phenoptosis. Shown is a graph simulating proliferation of reproducing genomes following the equation $N(t) = N(0) \times [(1-\alpha)(1+\beta)(1-\mu)]^t$, where N (0) = Initial number of reproduction-competent haploid genomes in a population, N (t) = Number of reproducing haploid genomes in a population at the generation time of t, α = aging rate constant, β = genome replication rate constant, μ = cohort competition/infection cost constant. For simulation, we used $N(0) = 100$; $\alpha = 0.2$; $\beta = 0.5$; $\mu = 0.04$ for the growth of numbers of reproducing genomes without phenoptosis or decreased phenoptosis (e.g. in *fshr-1* mutants), while $N(0) = 100$; $\alpha = 0.2$; $\beta = 0.5$; $\mu = 0.01$ with phenoptosis (e.g. in the wild-type *C. elegans*). For simplicity, we make assumptions on the clonal population structure and growth characteristics in a resource-limiting competitive environment and that α and β remain constant. Note that μ is lower with increased phenoptosis given limited resources for a population, since increased stress-induced phenoptosis in less reproduction-competent individuals leads to reduced competition for resources or vulnerability to pathogen infection at the population level.

(a)

Tissue Enrichment Analysis (TEA) for genes up-regulated by freeze-thaw stress:

Term	Expected	Observed	Enrichment	P value
intestine WBbt:0005772	95	178	1.9	5.3e-16
head mesodermal cell WBbt:0004697	43	84	2	4.9e-09
cephalic sensillum WBbt:0006920	6.7	23	3.4	1.4e-07
PVD WBbt:0006831	39	74	1.9	1.4e-07
outer labial sensillum WBbt:0005501	40	74	1.9	2.7e-07
cephalic sheath cell WBbt:0008406	10	27	2.7	1.7e-06
anterior ganglion WBbt:0005375	10	27	2.6	3e-06
ASE WBbt:0005663	25	43	1.7	0.00037
muscular system WBbt:0005737	95	123	1.3	0.0017
thermosensory neuron WBbt:0005838	31	47	1.5	0.0019

(b)

Gene Ontology Enrichment Analysis (GOEA) for genes up-regulated by freeze-thaw stress:

Term	Expected	Observed	Enrichment	P value
immune system process GO:0002376	3.7	48	13	1.8e-39
response to biotic stimulus GO:0009607	4.8	53	11	5.2e-39
interspecies interaction GO:0044419	4.9	53	11	6e-39
defense response GO:0006952	4.9	53	11	7.8e-39
membrane microdomain GO:0098857	0.6	8	13	8.4e-09
transmembrane transporter GO:0015399	1	5	5	0.00052
protein folding GO:0035966	2.4	8	3.3	0.00074
response to axon injury GO:0048678	0.43	3	6.9	0.00091
extracellular stimulus GO:0009991	0.92	4	4.4	0.0023

Figure S1: Transcriptome profiling and analysis of genes up-regulated by FTS.

(a), Tissue Enrichment Analysis (TEA) of genes up-regulated by FTS showing intestine as a major site of gene regulation. **(b)**, Gene Ontology Enrichment (GOE) Analysis of genes up-regulated by FTS showing abiotic stress and immune response as major processes involved. **(c-f)**, Cumulative probability analysis of genes regulated by SKN-1 (**c**), DAF-16 (**d**), HIF-1 (**e**), HSF-1 (**f**) (Brunquell et al., 2016; Kumar et al., 2015; Oliveira et al., 2009; Steinbaugh et al.) does not show marked enrichment of those also regulated by FTS (NCBI Sequence Read Archive BioProject accession PRJNA763790). **(g)**, Normalized fold induction of gene expression for the entire *oac* gene family based on RNAseq results, showing *oac-31* as the most up-regulated family member.

(a)

Signal transduction pathways (Cytoplasmic membrane proteins transcription factors)	Genes tested for RNAi phenotype in *oac-31*p::GFP (Selected with intestinal TPM expression > 2.0) (Blue: baseline GFP expression increased) (Red: freeze-thaw stress induction decreased)
Neuropeptide GPCRs	*npr-2, npr-8, npr-21, npr-25, npr-30*
Wnt receptors	*mom-5, mig-1, cam-1, lin-18*
Shh receptor-like proteins	*ptc-1, ptr-2, ptr-5, ptr-17, ptr-18, ptr-21*
TGF/BMP receptors	*daf-1*
TRP channels	*trpa-1, trpa-2, trp-1, trpl-5, gon-2, cup-5, gtl-1, gtl-2*
Voltage-gated calcium channels	*egl-19, unc-2, nca-2*
Cyclic nucleotide-gated ion channels	*cng-1*
Mechanosensory ion channels	*pezo-1, flr-1*
Innate immune receptors	*octr-1, fshr-1, tol-1*
Progestin/AdipoQ receptors	*paqr-1, paqr-2*
Receptor tyrosine kinases (Insulin, Eph, EGF, FGF receptors)	*daf-2, vab-1, let-23, kin-15*
Transcription factors	298 genes tested including: *hif-1, hsf-1, daf-16, skn-1, sbp-1, tra-1, zip-2, zip-10*

(b) Bright field / *oac-31*p::GFP / *unc-54*p::mCherry
Freeze-thaw + RNAi — Control, *fshr-1* (100%, n=50)

(c) No freeze-thaw + RNAi — Control, *sbp-1* (100%, n=50), *tra-1* (90%, n=50)

(d) Fold induction by qPCR — *oac-31*, *ncr-1*; Control, FTS; Ctr, *fshr-1*, Ctr; **

(e) Fold induction by qPCR — *W09* (n.s.), *tsp-1* (n.s.); Control, FTS; Ctr, *fshr-1*, Ctr, *fshr-1*

Figure S2: RNAi screens identify genes that regulate *oac-31*p::GFP induction by

FTS. (a), Table listing RNAi clones screened for *oac-31*p::GFP induction phenotype.

RNAi cloned were selected based on intestinal expression and genes encoding

transmembrane receptors and representing major signal transduction pathways. **(b)**,

Representative fluorescence images showing suppression of FTS-induced *oac-*

*31*p::GFP by RNAi against *fshr-1*. **(c)**, Representative fluorescence images showing

constitutive activation of *oac-31*p::GFP by RNAi against *tra-1* or *sbp-1* in the absence of

FTS treatment. Scale bar, 100 µm. **(d),** qRT-PCR results showing abolished induction of

ncr-1, *oac-31* by FTS in animals deficient in *fshr-1*. **(e),** qRT-PCR results showing

normal induction of the genes *W09G12.7* and *tsp-1*, by FTS in *fshr-1* mutants. Values

are means ± S.D with **$P < 0.01$ and *$P < 0.05$ (two-way ANOVA for genotype-effect

interaction and post-hoc Tukey HSD, N = 3 independent experiments, n > 50 for each

experiment). n.s., non-significant. Scale bar, 100 µm.

Figure S3: FSHR-1 regulates trade-off of organismic resilience to cholesterol

oxidative and hypothermic stresses. (a), Survival rates of control and *fshr-1* RNAi-

treated animals grown on increasing doses of Paraquat (3, 5, 10, 50 and 100 mM in

NGM). **(b),** Survival rates of wild type or *ncr-1; ncr-2* double mutant animals treated with

control or *fshr-1* RNAi after severe FTS (-20 °C for 45 minutes, followed by recovery for

24 hrs at 25 °C) or cold shock (CS, 4 °C for 24 hrs, followed by recovery for 24 hrs at 25

°C) grown on cholesterol (5 µg/ml). **(c),** Survival rates of control and *fshr-1* RNAi-treated

533 animals after severe CS grown on increasing doses of cholesterol (0, 5, 25, 50 μg/ml).

534 **(d),** Survival rates of control and *fshr-1* RNAi-treated animals after severe FTS grown on

535 increasing doses of cholesterol (0, 5, 25, 50 μg/ml). Values are means ± S.D with ***P <

536 0.001, **P < 0.01 and *P < 0.05 (two-way ANOVA for genotype-effect interaction and

537 post-hoc Tukey HSD, N = 3 independent experiments, n > 50 for each experiment).

Figure S4

Figure S4: *flr-2*, which encodes the putative ligand of FSHR-1, is essential for FTS

induction of *oac-31* but not transcriptionally regulated by FTS. (a), RNAseq results

showing no apparent up-regulation of *flr-2* or *fshr-1* by FTS (two-way ANOVA for

genotype-effect interaction and post-hoc Tukey HSD, N = 3 independent experiments,

n > 50 for each experiment). N.S., non-significant. **(b)**, Representative fluorescence

images for the translational reporter *flr-2*p::*flr-2*::GFP showing its normal secretion and

enrichment in coelomocytes but without apparent regulation by FTS (unpaired student-t

test, N.S., non-significant).

REFERENCES

Al-Fageeh, M.B., and Smales, C.M. (2006). Control and regulation of the cellular responses to cold shock: the responses in yeast and mammalian systems. Biochem. J. *397*, 247–259. https://doi.org/10.1042/BJ20060166.

Antebi, A. (2007). Genetics of aging in Caenorhabditis elegans. PLoS Genet. *3*, 1565–1571. https://doi.org/10.1371/journal.pgen.0030129.

Brenner, S. (1974). The genetics of Caenorhabditis elegans. Genetics *77*, 71–94. .

Brunquell, J., Morris, S., Lu, Y., Cheng, F., and Westerheide, S.D. (2016). The genome-wide role of HSF-1 in the regulation of gene expression in Caenorhabditis elegans. BMC Genomics *17*. https://doi.org/10.1186/s12864-016-2837-5.

Cho, S., Rogers, K.W., and Fay, D.S. (2007). The C. elegans glycopeptide hormone receptor ortholog, FSHR-1, regulates germline differentiation and survival. Curr. Biol. CB *17*, 203–212. https://doi.org/10.1016/j.cub.2006.12.027.

Coburn, C., Allman, E., Mahanti, P., Benedetto, A., Cabreiro, F., Pincus, Z., Matthijssens, F., Araiz, C., Mandel, A., Vlachos, M., et al. (2013). Anthranilate Fluorescence Marks a Calcium-Propagated Necrotic Wave That Promotes Organismal Death in C. elegans. PLoS Biol. *11*, e1001613. https://doi.org/10.1371/journal.pbio.1001613.

Davis, M.W., Hammarlund, M., Harrach, T., Hullett, P., Olsen, S., and Jorgensen, E.M. (2005). Rapid single nucleotide polymorphism mapping in C. elegans. BMC Genomics *6*, 118. https://doi.org/10.1186/1471-2164-6-118.

Ernst, R., Ejsing, C.S., and Antonny, B. (2016). Homeoviscous Adaptation and the Regulation of Membrane Lipids. J. Mol. Biol. *428*, 4776–4791. https://doi.org/10.1016/j.jmb.2016.08.013.

Ezcurra, M., Benedetto, A., Sornda, T., Gilliat, A.F., Au, C., Zhang, Q., van Schelt, S., Petrache, A.L., Wang, H., de la Guardia, Y., et al. (2018). C. elegans Eats Its Own Intestine to Make Yolk Leading to Multiple Senescent Pathologies. Curr. Biol. CB *28*, 2544-2556.e5. https://doi.org/10.1016/j.cub.2018.06.035.

Fontana, L., and Partridge, L. (2015). Promoting health and longevity through diet: from model organisms to humans. Cell *161*, 106–118. https://doi.org/10.1016/j.cell.2015.02.020.

Galimov, E.R., and Gems, D. (2020). Shorter life and reduced fecundity can increase colony fitness in virtual Caenorhabditis elegans. Aging Cell *19*, e13141. https://doi.org/10.1111/acel.13141.

Galimov, E.R., Lohr, J.N., and Gems, D. (2019). When and How Can Death Be an Adaptation? Biochem. Biokhimiia *84*, 1433–1437. https://doi.org/10.1134/S0006297919120010.

Garrity, P.A., Goodman, M.B., Samuel, A.D., and Sengupta, P. (2010). Running hot and cold: behavioral strategies, neural circuits, and the molecular machinery for thermotaxis in C. elegans and Drosophila. Genes Dev. *24*, 2365–2382. https://doi.org/10.1101/gad.1953710.

Gulyas, L., and Powell, J.R. (2021). Cold shock induces a terminal investment reproductive response in C.
elegans.

Hamilton, W.D. (1963). The Evolution of Altruistic Behavior. Am. Nat. *97*, 354–356.
https://doi.org/10.1086/497114.

Jiang, W., Wei, Y., Long, Y., Owen, A., Wang, B., Wu, X., Luo, S., Dang, Y., and Ma, D.K. (2018). A genetic
program mediates cold-warming response and promotes stress-induced phenoptosis in C. elegans. ELife
7. https://doi.org/10.7554/eLife.35037.

Kamath, R.S., and Ahringer, J. (2003). Genome-wide RNAi screening in Caenorhabditis elegans. Methods
San Diego Calif *30*, 313–321. .

Kato, M., Kashem, M.A., and Cheng, C. (2016). An intestinal microRNA modulates the homeostatic
adaptation to chronic oxidative stress in C. elegans. Aging *8*, 1979–2005.
https://doi.org/10.18632/aging.101029.

Kenyon, C.J. (2010). The genetics of ageing. Nature *464*, 504–512. https://doi.org/10.1038/nature08980.

Kim, S., and Sieburth, D. (2020). FSHR-1/GPCR Regulates the Mitochondrial Unfolded Protein Response
in Caenorhabditis elegans. Genetics *214*, 409–418. https://doi.org/10.1534/genetics.119.302947.

Kim, D., Langmead, B., and Salzberg, S.L. (2015). HISAT: a fast spliced aligner with low memory
requirements. Nat. Methods *12*, 357–360. https://doi.org/10.1038/nmeth.3317.

Kirkwood, T.B.L., and Melov, S. (2011). On the programmed/non-programmed nature of ageing within
the life history. Curr. Biol. CB *21*, R701-707. https://doi.org/10.1016/j.cub.2011.07.020.

Kudo, M., Chen, T., Nakabayashi, K., Hsu, S.Y., and Hsueh, A.J. (2000). The nematode leucine-rich repeat-
containing, G protein-coupled receptor (LGR) protein homologous to vertebrate gonadotropin and
thyrotropin receptors is constitutively active in mammalian cells. Mol. Endocrinol. Baltim. Md *14*, 272–
284. https://doi.org/10.1210/mend.14.2.0422.

Kumar, N., Jain, V., Singh, A., Jagtap, U., Verma, S., and Mukhopadhyay, A. (2015). Genome-wide
endogenous DAF-16/FOXO recruitment dynamics during lowered insulin signalling in C. elegans.
Oncotarget *6*, 41418–41433. .

Lee, D., An, S.W.A., Jung, Y., Yamaoka, Y., Ryu, Y., Goh, G.Y.S., Beigi, A., Yang, J.-S., Jung, G.Y., Ma, D.K., et
al. (2019). MDT-15/MED15 permits longevity at low temperature via enhancing lipidostasis and
proteostasis. PLoS Biol. *17*, e3000415. https://doi.org/10.1371/journal.pbio.3000415.

Lee, Y., Jung, Y., Jeong, D.-E., Hwang, W., Ham, S., Park, H.-E.H., Kwon, S., Ashraf, J.M., Murphy, C.T., and
Lee, S.-J.V. (2021). Reduced insulin/IGF1 signaling prevents immune aging via ZIP-10/bZIP-mediated
feedforward loop. J. Cell Biol. *220*, e202006174. https://doi.org/10.1083/jcb.202006174.

Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinforma. Oxf. Engl. *30*, 923–930.
https://doi.org/10.1093/bioinformatics/btt656.

Longo, V.D., Mitteldorf, J., and Skulachev, V.P. (2005). Programmed and altruistic ageing. Nat. Rev.
Genet. *6*, 866–872. https://doi.org/10.1038/nrg1706.

López-Otín, C., Blasco, M.A., Partridge, L., Serrano, M., and Kroemer, G. (2013). The Hallmarks of Aging.
Cell *153*, 1194–1217. https://doi.org/10.1016/j.cell.2013.05.039.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol. *15*, 550. https://doi.org/10.1186/s13059-014-0550-8.

Lu, X.Y., Gross, R.E., Bagchi, S., and Rubin, C.S. (1990). Cloning, structure, and expression of the gene for
a novel regulatory subunit of cAMP-dependent protein kinase in Caenorhabditis elegans. J. Biol. Chem.
265, 3293–3303. .

Luo, J., Yang, H., and Song, B.-L. (2020). Mechanisms and regulation of cholesterol homeostasis. Nat.
Rev. Mol. Cell Biol. *21*, 225–245. https://doi.org/10.1038/s41580-019-0190-7.

Ma, D.K., Vozdek, R., Bhatla, N., and Horvitz, H.R. (2012). CYSL-1 interacts with the O2-sensing
hydroxylase EGL-9 to promote H2S-modulated hypoxia-induced behavioral plasticity in C. elegans.
Neuron *73*, 925–940. https://doi.org/10.1016/j.neuron.2011.12.037.

Ma, D.K., Li, Z., Lu, A.Y., Sun, F., Chen, S., Rothe, M., Menzel, R., Sun, F., and Horvitz, H.R. (2015). Acyl-
CoA Dehydrogenase Drives Heat Adaptation by Sequestering Fatty Acids. Cell *161*, 1152–1163.
https://doi.org/10.1016/j.cell.2015.04.026.

Mello, C.C., Kramer, J.M., Stinchcomb, D., and Ambros, V. (1991). Efficient gene transfer in C.elegans:
extrachromosomal maintenance and integration of transforming sequences. EMBO J. *10*, 3959–3970. .

Metzstein, M.M., Stanfield, G.M., and Horvitz, H.R. (1998). Genetics of programmed cell death in C.
elegans: past, present and future. Trends Genet. TIG *14*, 410–416. .

Miller, E.V., Grandi, L.N., Giannini, J.A., Robinson, J.D., and Powell, J.R. (2015). The Conserved G-Protein
Coupled Receptor FSHR-1 Regulates Protective Host Responses to Infection and Oxidative Stress. PloS
One *10*, e0137403. https://doi.org/10.1371/journal.pone.0137403.

Murphy, R.C., and Johnson, K.M. (2008). Cholesterol, Reactive Oxygen Species, and the Formation of
Biologically Active Mediators. J. Biol. Chem. *283*, 15521–15525.
https://doi.org/10.1074/jbc.R700049200.

Mutlu, A.S., Duffy, J., and Wang, M.C. (2021). Lipid metabolism and lipid signals in aging and longevity.
Dev. Cell *56*, 1394–1407. https://doi.org/10.1016/j.devcel.2021.03.034.

Oishi, A., Gengyo-Ando, K., Mitani, S., Mohri-Shiomi, A., Kimura, K.D., Ishihara, T., and Katsura, I. (2009).
FLR-2, the glycoprotein hormone alpha subunit, is involved in the neural control of intestinal functions in
Caenorhabditis elegans. Genes Cells Devoted Mol. Cell. Mech. *14*, 1141–1154.
https://doi.org/10.1111/j.1365-2443.2009.01341.x.

Oliveira, R.P., Porter Abate, J., Dilks, K., Landis, J., Ashraf, J., Murphy, C.T., and Blackwell, T.K. (2009).
Condition-adapted stress and longevity gene regulation by Caenorhabditis elegans SKN-1/Nrf. Aging Cell
8, 524–541. https://doi.org/10.1111/j.1474-9726.2009.00501.x.

Powell, J.R., Kim, D.H., and Ausubel, F.M. (2009). The G protein-coupled receptor FSHR-1 is required for
the Caenorhabditis elegans innate immune response. Proc. Natl. Acad. Sci. U. S. A. *106*, 2782–2787.
https://doi.org/10.1073/pnas.0813048106.

Sapolsky, R.M. (2004). Why Zebras Don't Get Ulcers, Third Edition (New York: Holt Paperbacks).

Schade, M.A., Reynolds, N.K., Dollins, C.M., and Miller, K.G. (2005). Mutations that rescue the paralysis
of Caenorhabditis elegans ric-8 (synembryn) mutants activate the G alpha(s) pathway and define a third
major branch of the synaptic signaling network. Genetics *169*, 631–649.
https://doi.org/10.1534/genetics.104.032334.

Sevanian, A., and Peterson, A.R. (1986). The cytotoxic and mutagenic properties of cholesterol oxidation
products. Food Chem. Toxicol. Int. J. Publ. Br. Ind. Biol. Res. Assoc. *24*, 1103–1110.
https://doi.org/10.1016/0278-6915(86)90295-4.

Shi, J., Gao, W., and Shao, F. (2017). Pyroptosis: Gasdermin-Mediated Programmed Necrotic Cell Death.
Trends Biochem. Sci. *42*, 245–254. https://doi.org/10.1016/j.tibs.2016.10.004.

Shore, D.E., and Ruvkun, G. (2013). A cytoprotective perspective on longevity regulation. Trends Cell
Biol. *23*, 409–420. https://doi.org/10.1016/j.tcb.2013.04.007.

Skulachev, V.P. (1999). Phenoptosis: programmed death of an organism. Biochem. Biokhimiia *64*, 1418–
1426. .

Skulachev, V.P. (2002). Programmed death phenomena: from organelle to organism. Ann. N. Y. Acad.
Sci. *959*, 214–237. .

Skulachev, V.P. (2019). Phenoptosis as a Phenomenon Widespread among Many Groups of Living
Organisms Including Mammals (Commentary to the Paper by E. R. Galimov, J. N. Lohr, and D. Gems
(2019) Biochemistry (Moscow), 84, 1433-1437). Biochem. Biokhimiia *84*, 1438–1441.
https://doi.org/10.1134/S0006297919120022.

Smith, J.M. (1964). Group Selection and Kin Selection. Nature *201*, 1145–1147.
https://doi.org/10.1038/2011145a0.

Steinbaugh, M.J., Narasimhan, S.D., Robida-Stubbs, S., Moronetti Mazzeo, L.E., Dreyfuss, J.M., Hourihan,
J.M., Raghavan, P., Operaña, T.N., Esmaillie, R., and Blackwell, T.K. Lipid-mediated regulation of SKN-
1/Nrf in response to germ cell absence. ELife *4*, e07836. https://doi.org/10.7554/eLife.07836.

Taylor, S.R., Santpere, G., Weinreb, A., Barrett, A., Reilly, M.B., Xu, C., Varol, E., Oikonomou, P.,
Glenwinkel, L., McWhirter, R., et al. (2021). Molecular topography of an entire nervous system. Cell *184*,
4329-4347.e23. https://doi.org/10.1016/j.cell.2021.06.023.

Weinlich, R., Oberst, A., Beere, H.M., and Green, D.R. (2017). Necroptosis in development, inflammation
and disease. Nat. Rev. Mol. Cell Biol. *18*, 127–136. https://doi.org/10.1038/nrm.2016.149.

Wu, D., Wang, Z., Huang, J., Huang, L., Zhang, S., Zhao, R., Li, W., Chen, D., and Ou, G. (2022). An
antagonistic pleiotropic gene regulates the reproduction and longevity tradeoff. Proc. Natl. Acad. Sci. U.
S. A. *119*, e2120311119. https://doi.org/10.1073/pnas.2120311119.

Xiao, R., Zhang, B., Dong, Y., Gong, J., Xu, T., Liu, J., and Xu, X.Z.S. (2013). A genetic program promotes C.
elegans longevity at cold temperatures via a thermosensitive TRP channel. Cell *152*, 806–817.
https://doi.org/10.1016/j.cell.2013.01.020.