1

Title: Nutrient addition increases grassland resistance during dry and wet climate extremes

despite functional diversity loss

Authors

Qingqing Chen[1], Shaopeng Wang[1*], Eric W. Seabloom[2], Forest Isbell[2], Elizabeth T. Borer[2],

Jonathan D. Bakker[3], Siddharth Bharath[2], Christiane Roscher[4,5], Pablo Luis Peri[6], Sally A.

Power[7], Ian Donohue[8], Carly Stevens[9], Anne Ebeling[10], Carla Nogueira[11], Maria C Caldeira[11],

Andrew S. MacDougall[12], Joslin L. Moore[13], Sumanta Bagchi[14], Anke Jentsch[15], Michelle

Tedder[16], Kevin Kirkman[16], Juan Alberti[17], and Yann Hautier[18]

Affiliations

[1]Institute of Ecology, College of Urban and Environmental Sciences, Peking University, Beijing,

China.

[2]Department of Ecology, Evolution, and Behavior, University of Minnesota, St. Paul, Minnesota,

USA.

[3]School of Environmental and Forest Sciences, University of Washington, Seattle, USA.

[4]German Centre for Integrative Biodiversity Research (iDiv) Halle- Jena- Leipzig, Leipzig,

Germany.

[5]Department of Physiological Diversity, Helmholtz Centre for Environmental Research – UFZ,

Leipzig, Germany.

[6]Instituto Nacional de Tecnología Agropecuaria (INTA)- Universidad Nacional de la Patagonia

Austral (UNPA), Santa Cruz, Argentina.

[7]Hawkesbury Institute for the Environment, Western Sydney University, New South Wales,

Australia.

[8]Department of Zoology, School of Natural Sciences, Trinity College Dublin, Dublin, Ireland.

[9]Lancaster Environment Centre, Lancaster University, Lancaster, UK.

[10]Institute of Ecology and Evolution, University Jena, Jena, Germany.

[11]Forest Research Centre, School of Agriculture, University of Lisbon, Lisbon, Portugal.

[12]Department of Integrative Biology, University of Guelph, Ontario, Canada.

[13]Hawkesbury Institute for the Environment, Western Sydney University, New South Wales,

Australia.

[14]Centre for Ecological Sciences, Indian Institute of Science, Bangalore, India.

[15]Disturbance Ecology and Vegetation Dynamics, Bayreuth Center of Ecology and Environmental

Research (BayCEER).

[16]School of Life Sciences, University of Kwazulu-Natal, Pietermaritzburg, South Africa.

[17]Instituto de Investigaciones Marinas y Costeras (IIMyC), Mar del Plata, Argentina.

[18]Ecology and Biodiversity Group, Department of Biology, Utrecht University, Utrecht, The

Netherlands.

Corresponding author: shaopeng.wang@pku.edu.cn

Total word count:5160

Word count for Introduction: 946

Word count for Materials and Methods: 2176

Word count for Results: 715

Word count for Discussion: 1323

Number of figures: 4 (Fig. 1 and Fig. 4 should be published in color)

Number of tables: 1

Number of supporting information: 1

Summary

Eutrophication impacts plant diversity such as species richness, functional trait diversity and composition of grassland communities globally, but whether and how these changes affect the functional stability of grasslands under increasing climate extremes is unknown.

We quantify the direct and diversity-mediated effects of nutrient addition on functional stability as measured by resistance during and resilience after dry and wet growing seasons, and temporal invariability of community productivity across 19 grasslands spanning four continents.

We found that nutrient addition increased resistance during dry and wet growing seasons, although the increased resistance during wet growing seasons was partly offset by the indirect negative effects of nutrient addition promoting fast-growing communities and reducing 'slow-fast' functional diversity. Nutrient addition had, however, weak net effects on resilience and invariability.

Thus, eutrophication strongly impacts plant diversity, but diversity-mediated indirect effects may be weaker on stability.

Keywords

Eutrophication; climate extremes; grasslands; biodiversity; Nutrient Network (NutNet)

Introduction

Earth is undergoing multiple global environmental changes including eutrophication and increased climatic variability, which threaten both the diversity and stability of ecosystems (IPCC, 2019). Mounting evidence shows that declines in diversity reduce the stable provisioning of essential ecosystem functions and services to humanity. Understanding the links between global environmental changes, diversity, and ecosystem stability has thus become a central goal in ecology. However, this understanding is limited because diversity and stability are multifaceted concepts, yet most studies have analyzed one or a few facets in isolation (Donohue *et al.*, 2013; Hillebrand *et al.*, 2018; Kéfi *et al.*, 2019). Stability characterizes ecosystems' responses to different types of perturbations (Pimm, 1984; Donohue *et al.*, 2013). In ecological studies, stability of an ecosystem function (e.g. productivity) can be defined as resistance during and resilience after perturbations and temporal invariability (Pimm, 1984). Resistance is the ability of a function to remain close to normal levels during a pulse perturbation (e.g. a climate extreme), and resilience is the rate of recovery towards normal levels after a pulse perturbation (Pimm, 1984). Invariability is often quantified as the ratio of the temporal mean of a function to its standard deviation (Pimm, 1984; Tilman, 1996). Because both higher resistance and resilience indicate the proximity of a function to normal levels, higher resistance, higher resilience, or both are expected to increase invariability ((Ives & Carpenter, 2007; Arnoldi *et al.*, 2016). Yet, this intuitive hypothesis has not been rigorously tested. Investigating multiple facets of stability simultaneously is critical to improve our mechanistic understanding and prediction for ecosystem stability.

In grasslands, nutrient addition has been shown to decrease resistance in aboveground primary productivity during dry climate extremes as well as temporal invariability, while its effects on resilience after dry climate extremes are mixed (Tilman & Downing, 1994; Xu *et al.*, 2014; Hautier *et al.*, 2015; Bharath *et al.*, 2020; Meng *et al.*, 2021; Van Sundert *et al.*, 2021; Carroll *et al.*, 2021). However, to date, few empirical studies have investigated the impact of nutrient addition on resistance during and resilience after wet climate extremes, which are becoming increasingly frequent just as dry climate extremes globally (IPCC, 2019). Nutrient addition may decrease resistance during dry climate extremes through increasing nutrients-to-water imbalance for plants, which reduces productivity and increases its deviation from normal levels (Meng *et al.*, 2021). Additionally, increased nutrients-to-water imbalance increase plant mortality, thus nutrient addition may decrease resilience after dry climate extremes (Southon *et al.*, 2012; Meng *et al.*, 2021). In contrast, nutrient addition may decrease resistance during wet climate extremes because

increased nutrients and water availability increase plant growth and productivity, which also increase the magnitude of its deviation from normal levels (Wright *et al.*, 2015; Wang *et al.*, 2017; Eskelinen *et al.*, 2020). Additionally, increased productivity due to increased nutrients and water availability may persist or even amplify in later years (Sala *et al.*, 2012; Wheeler *et al.*, 2021), thus nutrient addition may decrease resilience after wet climate extremes (Fig. 1; Table S1).

Fig. 1. Conceptual framework illustrating how nutrient addition impacts facets of stability through its direct and diversity-mediated indirect effects (A) and how nutrient addition alters facets of plant diversity (B-F). Shapes represent different species, darker color represents faster-growing species. Thick boxes represent nutrient addition conditions. Black and red arrows represent increase and decrease (darker color represents a larger change). Resistance.d/resistance.w: resistance during dry/wet growing seasons; resilience.d/ resilience.w: resilience after dry/wet growing seasons. CWM (slow-fast): community-weighted mean of the 'slow-fast' leaf economics spectrum; FD (slow-fast): functional diversity of the 'slow-fast' leaf economics spectrum. See Table 1 for the calculation and interpretation of each variable. See Table S1 for detailed potential mechanisms for each link in A.

In addition to these direct effects, nutrient addition may impact facets of stability indirectly by altering different facets of plant diversity including functional trait composition and diversity, and species richness. Nutrient addition may decrease resistance during dry and wet climate extremes but increase resilience after the extremes by promoting fast communities (communities dominated by fast-growing species; Fig.1). Its impact on invariability may, however, be low due to contrasting effects of fast communities on resistance and resilience (Craven *et al.*, 2018). Fast-growing (acquisitive) species are typically associated with high leaf nutrients and specific leaf area, which can make better use of resources by taking up nutrients faster and growing faster, thus these species have higher resilience after climate extremes. In contrast, slow-growing (conservative) species invest more in cell walls and secondary metabolites that may help them endure unfavorable environments, thus they have higher resistance during climate extremes. Nutrient addition may promote fast communities through shifting intraspecific traits related to fast growth, promoting fast-growing species, or both (Lepš *et al.*, 2011; Siefert & Ritchie, 2016; Tatarko & Knops, 2018; Zhou *et al.*, 2018; Pichon *et al.*, 2022). Additionally, nutrient addition may decrease all stability facets investigated here by decreasing the diversity of slow-fast trait that represents leaf economic spectrum and species richness. Species richness may capture diversity in other traits (e.g. root and size-related traits) that cannot be captured by the leaf economic spectrum. Communities with higher diversity in slow-fast trait or species richness are more likely to include species that are better adapted to perturbations, thus decreases in some species may be compensated by increases in others during and after climate extremes. This leads to less productivity deviation from normal levels, which increases resistance and resilience (Tilman *et al.*, 2014), as well as invariability (Craven *et al.*, 2018). Together, the net effects of nutrient addition on different facets of stability depend on the direction and relative importance of its direct and diversity-mediated indirect effects.

Here, we use a globally distributed nutrient addition experiment to investigate whether and how eutrophication impacts resistance during and resilience after climate extremes and temporal invariability directly and indirectly through altering the community composition and diversity of slow-fast trait, and species richness.

Materials and Methods

Experimental Design

We used a globally replicated nutrient and herbivore manipulation experiment (NutNet;

www.nutnet.org) initiated in 2007 for this analysis (Borer *et al.*, 2014, 2017). At most sites, each

of the ten treatments (Control, N, P, K, NP, NK, PK, NPK, Fence, and NPK+Fence) was

randomly imposed to a 25 m^2 plot and was replicated in at least three blocks. N was supplied as

time-release urea ((NH_2)$_2$CO), P was supplied as triple superphosphate (Ca(H_2PO_4)$_2$), and K as

potassium sulfate (K_2SO_4). N, P, K were added annually before the start of the growing season at

a rate of 10 g m^{-2} y^{-1}. A micronutrient and macronutrient mix (Fe, S, Mg, Mn, Cu, Zn, B, and Mo

in combination) was applied at a rate of 100 g m^{-2} once at the start of the experiment, as part of

the K addition. Fencing is designed to exclude large herbivores; fences were around 230 cm tall

and the lower 90 cm were covered by 1 cm woven wire mesh. To further exclude digging animals

such as voles and rabbits, an additional 30 cm outward-facing flange was stapled to the soil. Four

strands of barbless wire were strung at similar vertical distances above the wire mesh. Further

details on the design are available in (Borer *et al.*, 2014). We focus on exploring stability in the

control and nutrient addition (NPK) treatments, but we used traits measured in all treatments.

Specifically, we separated leaf traits measured in ambient (control and Fence) and nutrient-

enriched conditions (N, P, K, NP, NK, PK, NPK, NPK+Fence). Additionally, multiple trait

measurements for the same species at one site (e.g. from different blocks) were averaged. We did

this to maximize the number of species with available trait data and because previous NuNet

results found little variation in leaf traits among blocks within the non-fertilized and fertilized

conditions at 27 NutNet sites (Firn *et al.*, 2019).

Site selection

We selected sites meeting the following criteria:1) at least 3 years of data for the control and

nutrient addition (NPK) treatments; 2) 3 blocks, most sites have 3 blocks, for those with more

than 3 blocks, we selected the first 3 blocks following (Hautier *et al.*, 2020); 3) at least 3 common

to dominant plant species measured for leaf traits in both ambient and nutrient-enriched

conditions; 4) average across blocks and experimental years, summed cover of species with traits

accounted for >50% of the total cover; 5) during experimental years, at least one climate extreme

occurred (see "Defining climate extreme" for more details). Nineteen sites met these criteria (Fig.

S1; Table S2). These sites are distributed in North America, Australia, Europe, and Africa. Sites

include many grassland types such as montane, alpine, semiarid grasslands, prairies, and old

fields. See table S2 for geolocation, climate, and experimental durations for these sites. See table

S3 for the number of species, percent cover of species measured for traits at each site.

 Sampling protocol

 All sites followed standard NutNet sampling protocols. A 1×1m subplot was permanently marked

 within a 25 m^2 plot. Number of species, species identity, and their covers were recorded once per

 year in these 1×1 m subplots at most sites. At a few sites with strong seasonality, cover and

 biomass were recorded twice per year to include a full list of species and follow typical

 management procedures. For those sites, the maximum cover for each species and total biomass

 for a community were used in analyses. The taxonomy was checked and adjusted within sites to

 ensure consistent naming over time. For instance, when individuals could not be identified as

 species in all years, they were aggregated at the genus level but referred to as taxa for simplicity.

 Meanwhile, productivity was measured adjacent to these permanent subplots within two 1×0.1

 m strips (in total 0.2 m^2), which were moved from year to year to avoid resampling the same

 location. All aboveground biomass was clipped within strips. Biomass strips were sorted into

 dead and live and dried separately at 60 °C to constant mass before weighing to the nearest 0.01

 g. Dried biomass was multiplied by 5 to estimate grams per square meter. Following previous

 analyses, we used dried live aboveground biomass as a measure of primary productivity (Hautier

 et al., 2020; Seabloom *et al.*, 2021).

 Leaf trait measurements

 Leaf traits were generally measured after 3 to 4 years of nutrient addition (Table S3). Traits were

 measured for 3 to 5 of the most dominant species in each subplot (although 8 most dominant

 species were measured at one site). Traits were measured according to standard protocols (Pérez-

 Harguindeguy *et al.*, 2016). A detailed description of these trait measurements can be found in

 (Firn *et al.*, 2019). Briefly, for each species measured for leaf traits, five fully grown leaves

 without clear grazing marks were randomly selected. Leaf area (mm^2) was measured using leaf

 area meters or scanner. After that, dry weight (g) of leaves was measured after oven-drying at 60

 °C for 48 h. Specific leaf area (SLA; mm^{2}g^{-1}) was calculated as the leaf area divided by the dry

 weight. Then, leaf nutrient concentrations (%) including C, N, P, and K were determined. Leaf P,

 K, and C were determined using laser ablation inductively coupled plasma mass spectrometry

 following (Duodu *et al.*, 2015). Leaf N was determined using a LECO TruMac, based on a

 combustion technique using thermal conductivity relative to pure gas with an error $< 1\%$.

Table 1. Variables estimated in this study with their mathematical definitions and

interpretations.

Variables	Methods	Parameters in the methods explained	Interpretations	References
CWM (slow-fast)	$\frac{\sum a_i x_i}{\Sigma a_i}$	a_i is the cover of species i, x_i is the slow-fast trait for species i.	A higher value indicates that a plant community is dominated by fast-growing species.	Garnier *et al*. 2004)
CWM induced by replacement among existing species	$CWM_{Nut*} - CWM_{control}$	CWM_{Nut} and $CWM_{control}$ are CWM in the nutrient addition and control plots. CWM_{Nut*} is the CWM in the nutrient addition plots recalculated by using slow-fast trait in the control plots.	A higher value of replacement indicates that a change in CWM is more induced by a change in relative cover among existing species while less induced by intraspecific trait shift.	(Jung *et al*. 2014)
CWM induced by intraspecific trait shift	$CWM_{Nut} - CWM_{Nut*}$			
FD (slow-fast)	$\frac{\sum a_i z_i}{\Sigma a_i}$	a_i is cover of species i. z_i is the distance of slow-fast trait of species i to community-weighted mean trait (CWM).	A higher value indicates that a plant community has higher dispersion in slow-fast trait.	(Laliberte & Legendre 2010)
Invariability	$\frac{u}{\sigma}$	μ is the mean of aboveground primary productivity, σ is the standard deviation after detrending productivity over time.	A higher value indicates that a plant community varies less in productivity over time.	(Pimm 1984)
Resistance.d/resistance.w	$\frac{\bar{Y}_n}{\lvert Y_e - \bar{Y}_n \rvert}$	$\bar{Y}_n$, Y_e, and Y_{e+1} are aboveground productivity during normal growing seasons (average over all non-extreme growing seasons), during a dry or wet growing season, and one year after a dry or wet growing season, respectively.	A higher value of resistance.d or resistance.w indicates that productivity deviates less under a dry or wet growing season to that of normal growing seasons.	(Isbell *et al*. 2015)
Resilience.d/resilience.w	$\frac{\lvert Y_e - \bar{Y}_n \rvert}{\lvert Y_{e+1} - \bar{Y}_n \rvert}$		A higher value of resilience.d or resilience.w indicates that productivity deviates less one year after a dry or wet growing season relative to that under a dry or wet growing season.	

The slow-fast leaf economic spectrum, community-weighted mean traits, and functional trait

diversity

Overall, 179 plant species were measured for leaf traits (leaf C, leaf N, leaf P, leaf K, and SLA).

We used these five leaf traits to construct the "slow-fast" leaf economic spectrum using principal

component analysis (PCA). We used the function "PCA" from the R package "FactoMineR" (Lê

et al., 2008). We extracted the first axis of PCA, which explained around 41% of the variance, to

represent the "slow-fast" spectrum (i.e. slow-fast trait). Higher values (i.e. lower leaf C, higher

leaf N, leaf P, leaf K, and SLA) indicate faster-growing species (Fig. S2). We used this integrated

slow-fast trait to calculate CWM and FD. CWM is the average of the slow-fast trait weighted by

the relative cover of that species in a community (Garnier, Eric *et al.*, 2004). FD is cover-

weighted dispersion of the slow-fast trait relative to CWM (Laliberte & Legendre, 2010). We

calculated the slow-fast CWM and FD using the function "dbFD" from the R package "FD"

(Laliberté *et al.*, 2014). See table 1 for mathematical formulas for calculating the slow-fast CWM

and FD and their interpretations. These variables were calculated annually for each plot and then

averaged across years. Given that leaf traits were not measured for all plant species that occurred

within a subplot, we carried out a sensitivity analysis to evaluate how much slow-fast CWM and

FD would be affected by ignoring species with traits. Specifically, we simulated 50 species with

abundance ranging from 0 to 100 and slow-fast trait values ranging from 0 to 8 similar to our

data. For each simulated community, we calculated the slow-fast CWM and FD using all species.

We then randomly (1000 samplings) picked 1~48 species, ignoring their trait values (to simulate

lack of trait data) to examine how much slow-fast CWM and FD were influenced by losing these

species. Simulation results suggest that the relative deviation is smaller than 20% if species with

trait information contribute to > 60% of the total cover (Fig. S3).

Partitioning the slow-fast CWM into intraspecific trait change and species replacement

We first recalculated the slow-fast CWM and FD using shared species (species that were present

in both the control and NPK plots) instead of all species with trait data. We then partitioned the

slow-fast CWM based on shared species into intraspecific trait variation and replacement of

existing species following (Jung *et al.*, 2014). See Table 1 for mathematical formulas for

calculating and interpreting these variables.

Defining climate extreme and stability facets

Following (Isbell *et al.*, 2015), we used the standardized precipitation–evapotranspiration index

(SPEI) to classify climate extremes for each site. SPEI was calculated as the standardized (z-

score) water balance (precipitation – evapotranspiration; mm) over the growing season from 1901

to 2020. Precipitation and potential evapotranspiration used to calculate SPEI were downloaded

from https://crudata.uea.ac.uk/cru/data/hrg/cru_ts_4.05/ (Harris *et al.*, 2020). We initially

categorized each growing season into extreme dry, moderate dry, normal, moderate wet, and

extreme wet using the cutoff of 1.28 and 0.67 sd (around 10 % and 25% quantile). Extreme dry:

SPEI ≤ -1.28 sd; moderate dry: -1.28 sd < SPEI ≤ -0.67 sd; normal growing season: -0.67 sd <

SPEI < 0.67 sd; moderate wet: 0.67 sd ≤ SPEI < 1.28 sd; and extreme wet: SPEI ≥ 1.28 sd. For

some years at some sites, dry and wet growing seasons occurred consecutively, limiting our

ability to disentangle their separate impact on diversity and stability; these years were therefore

removed. For instance, at comp.pt, year 1 was eliminated because year 0 had a dry growing

season, which may bias resistance during wet growing season in year 1. Sites and experimental

years included in analyses can be found in Fig. S1 and Table S1. In total, 18, 22, 70, 14, and 18

extreme dry, moderate dry, normal, moderate wet, and extreme wet growing seasons across 19

sites were used in our analyses. We compared productivity under the moderate and extreme growing seasons relative to that of normal growing seasons. Productivity under moderate growing seasons was generally more similar to that under extreme growing seasons than normal growing seasons (Fig. S4). Thus, we focus on the cutoff of 0.67 in the following analyses. That is, we included moderate and extreme dry growing seasons as dry growing seasons, while moderate and extreme wet growing seasons as wet growing seasons.

We quantified invariability as the ratio of the temporal mean to the standard deviation of productivity (Pimm, 1984; Tilman *et al.*, 2006). We quantified resistance as the inverse of the proportional displacement of productivity from normal levels during a dry or wet growing season, a measure of the ability of a community to maintain a similar amount of productivity under perturbations, following (Isbell *et al.*, 2015). Also, we quantified resilience as the inverse of the proportional lack of recovery in productivity during the year after a dry or wet growing season, following (Isbell *et al.*, 2015). There are many ways to quantify resistance and resilience, here we chose this measure to facilitate comparison among sites with different levels of productivity (Ingrisch & Bahn, 2018). We calculated resistance during and resilience after dry and wet growing seasons separately. See Table 1 for mathematical formulas for calculating stability metrics and their interpretations.

Statistical analysis

All analyses were performed in R v.4.1.6 (R Core Team., 2022). We used linear mixed-effects models (function "lme") from the R package "nlme" (Pinheiro *et al.*, 2017) for the following analyses. We built models where site and block nested within sites were the random effects, treatments were the fixed effects. First, we tested whether nutrient addition impacted stability facets. To facilitate among-site comparison, we present raw data of productivity under different climate extremes over the experimental years at each site in Fig. S5. Next, we aggregated productivity difference to normal levels in control and nutrient addition treatments during and one year after dry and wet growing seasons at each site in Fig. S6. We then aggregated average productivity and the magnitude of productivity deviation to normal levels during and one year after dry and wet growing seasons in control and nutrient addition treatments across sites in Fig. S7. Furthermore, we present the effects of nutrient addition on each stability facet at each site in Fig. S8. Second, we tested whether nutrient addition impacted the slow-fast CWM and FD using all species with traits and the shared species. We also tested whether nutrient addition impacted

the component parts of the slow-fast CWM based on shared species, that is, intraspecific trait shift

and replacement of existing species.

To evaluate whether nutrient addition impacted stability facets through the slow-fast CWM and

FD, and species richness, we built a structural equation model (SEM) using the function "psem"

from the R package piecewiseSEM (Lefcheck, 2016). An initial model was built based on prior

knowledge (Fig. 1A, rationale for each link can be found in Table S1). For each component model

in SEM, we used the function "lme" with site and block nested within sites as the random effects.

We estimated variance inflation for each component model to make sure that multi-collinearity

did not affect parameter estimates (variance inflation < 4). We used 12 sites at which both dry and

wet growing seasons occurred. Sites with only one dry or wet growing season in the last year of

the experiment were excluded because resilience could not be calculated in this case. See Table

S1 for 12 sites and experimental years included. To check whether our results held using the

shared species instead of all species with traits, we ran another SEM using the slow-fast CWM

and FD calculated using shared species. The link between each plant diversity facet and each

stability facet in the SEM was evaluated across the control and nutrient addition treatments, to

check whether these relationships differ under these two treatments, we ran bivariate analyses

using function "lme" from the R package "nlme" (Pinheiro *et al.*, 2017). In each model, a stability

facet was the response variable, a plant diversity facet and its interaction with treatment were

fixed variables, and site was included as a random variable.

Results

Although adding nutrient did not change productivity in a consistent manner under different

growing seasons across sites (Fig. S5-S6), on average, nutrient addition increased productivity by

54% during normal growing seasons (control: 320.93; nutrient addition: 496.44; $g.m^{-2}$; Fig. S7).

During dry growing seasons, nutrient addition decreased productivity more from their normal

levels by 45% (control: -79; nutrient addition: -114; $g.m^{-2}$). Thus, nutrient addition also increased

the magnitude of productivity deviation by 18% (control: 185.90; nutrient addition: 248.61; $g.m^{-}$

^{2}). During wet growing seasons, nutrient addition slightly increased productivity from their

normal levels (control: -2.8; nutrient addition: 14.1; $g.m^{-2}$) and it increased the magnitude of

productivity deviation by 38% (control: 180.20; nutrient addition: 248.61; $g.m^{-2}$). Because the

increased deviations were relatively small compared with the increased productivity during

normal growing seasons under nutrient addition, nutrient addition increased resistance during dry

and wet growing seasons (dry: df = 53, t-value = 2.43, p = 0.02; wet: df = 41, t-value = 1.68, p =

0.1; Fig. 2; Table S4). One year after dry growing seasons, nutrient addition increased

productivity from their normal levels (control: 4.3; nutrient addition: 46.1; g.m^{-2}; Fig. S7), and it

increased the magnitude of productivity deviation by 42% (control: 149; nutrient addition: 105;

g.m^{-2}). Similarly, one year after wet growing seasons, nutrient addition also increased

productivity from their normal levels (control: -9.8; nutrient addition: 31.4; g.m^{-2}), and increased

the magnitude of productivity deviation by 79% (control: 213; nutrient addition: 119; g.m^{-2}).

However, nutrient addition did not impact resilience after dry and wet growing seasons, nor did it

impact invariability (Fig. 2; Table S4; also see stability facets at individual sites in Fig. S8).

Fig. 2. Effects of nutrient addition on facets of stability. Small dots are raw values from each

site. Large black dots are mean over experimental years and over 19 sites estimated from linear

mixed effect models, error bars are 95% confidence intervals. See Table S4 for test statistics. All

stability facets are on the log scale. Resistance.d/resistance.w: resistance during dry/wet growing

seasons; resilience.d/ resilience.w: resilience after dry/wet growing seasons.

Nutrient addition decreased species richness (t-value = -4.98, df = 56, p < 0.001). Using all

species with traits, nutrient addition decreased the slow-fast FD (t-value = -3.55, df = 56, p =

0.001) but increased the slow-fast CWM (t-value = 7.51, df = 56, p < 0.001). Using shared

species (species occurring in both control and nutrient addition treatment), the effects of nutrient

addition were similar for the slow-fast FD (paired t-test: t = 0.67, df = 56, p = 0.51), but the

effects were weaker on the slow-fast CWM compared with that using all species with traits

(paired t-test: t = 1.83, df = 18, *p* = 0.08). Using the shared species, we further found that

increased slow-fast CWM under nutrient addition was largely driven by intraspecific trait

variation but not by replacement among existing species (Fig. 3; Table S5).

Fig. 3. Effects of nutrient addition on species richness, slow-fast CWM and FD. For shared

species (i.e. species encountered in both control and nutrient addition treatments), change in slow-

fast CWM was further attributable to species replacement among existing species and

intraspecific trait variation (ITV). Small dots are raw values from each site. Large black dots are

mean values over experimental years and over 19 sites estimated from linear mixed effect models,

error bars are 95% confidence intervals. See Table S5 for test statistics.

The SEM revealed that nutrient addition impacted facets of stability directly and indirectly by

affecting different facets of plant diversity (Fig. 4). Nutrient addition directly increased resistance

during dry and wet growing seasons (path coefficient: 0.21 and 0.31). However, the direct effects

of nutrient addition on resistance during wet growing seasons were partly offset by its indirect

effects through promoting fast communities and reducing slow-fast FD (compound path

coefficient: -0.18). As a result, nutrient addition increased invariability due to increased resistance

(compound path coefficient: 0.13). Nutrient addition directly decreased resilience after wet

growing seasons (path coefficient: -0.18) but had no detectable impact on resilience after dry

growing seasons. However, the direct effects of nutrient addition on resilience after wet growing

seasons were partly offset by indirect effects through reducing species richness and slow-fast FD

(compound path coefficient: 0.09). As a result, nutrient addition decreased invariability due to

decreased resilience (compound path coefficient: -0.02). Moreover, nutrient addition decreased

invariability through decreasing species richness (compound path coefficient: -0.04). Summing up

the pathways through resistance, resilience, and species richness, nutrient addition weakly

increased invariability (net effects: 0.07). These results were generally similar when using shared

species to calculate the slow-fast CWM and FD, except that the link between the slow-fast CWM

and resistance during wet growing seasons and the link between the slow-fast FD and resilience

after wet growing seasons became non-significant (Fig. S9). The relationships between diversity

and stability in the SEM were generally consistent with the results from bivariate analyses (Fig.

S10).

Fig. 4. Direct and indirect effects of nutrient addition on facets of stability. The model

described the data well (F = 9.217, p = 0.512, df = 10). Results are based on 12 sites where both

dry and wet growing seasons occurred. See Fig. S1 and Table S2 for sites and experimental years

included. White box represents nutrient addition by NPK, blue boxes represent measured

variables, and arrows represent relationships among variables. The displayed numbers are

standardized path coefficients. The width of the arrows indicates the strength of the pathways.

Line color represents positive (black) and negative (red) effects. Grey lines and text show

correlated errors. Non-significant paths are not shown. Asterisks indicate significant paths: * $p \leq$

0.1; ** $p \leq 0.05$; *** $p \leq 0.001$. All stability facets were on the log scale to improve normality

and homogeneity of variance.

Discussion

Our analyses revealed idiosyncratic responses of different facets of stability to nutrient addition:

whereas nutrient addition only impacted resistance during dry and wet growing seasons, it did not

affect resilience and invariability. Additionally, nutrient addition increased resistance mainly

through its direct effects. Although nutrient addition also impacted facets of stability indirectly

through different facets of plant diversity, these diversity-mediated indirect effects were relatively

weak and were overwhelmed by the direct effects. Our results highlight the necessity of

considering multiple facets of stability and the importance of disentangling the direct and

diversity-mediated indirect effects to improve our prediction for ecosystem stability in the face of

global change.

Effects of nutrient addition on stability facets

To enable comparison among sites with varying productivity and to link resistance and with other

stability facets such as temporal invariability, we used the proportional change of productivity

from normal levels to quantify resistance. Relative to normal levels, nutrient addition strongly

decreased productivity compared with ambient conditions during dry growing seasons (Fig. S7),

consistent with previous studies that used change in productivity to quantify resistance (Bharath *et*

al., 2020; Meng *et al.*, 2021; Van Sundert *et al.*, 2021). In contrast, nutrient addition increased

productivity, albeit slightly, during wet growing seasons (Fig. S7), consistent with previous

single-site studies showing that increased nutrients and rainfall increase productivity (Xu *et al.*,

2014; Wang *et al.*, 2017). Here we found that nutrient addition increased resistance mainly

because it increased productivity during normal growing seasons. This suggests that, on average,

grassland sites investigated are nutrient-limited (Fay *et al.*, 2015). We found that diversity-

mediated indirect effects were evident for resistance during wet but not dry growing seasons

suggesting that effects of diversity may depend on climate. During wet growing seasons, an

increase in resources may intensify plant-plant competition and thereby compensatory dynamics

(Gonzalez & Loreau, 2009). During dry growing seasons, plant species are often under stress,

thus abiotic factors and facilitation, which depend less on plant diversity, may be more important

than competition in structuring plant communities (de Bello *et al.*, 2021). Moreover, the indirect

effects of nutrient addition on resistance during wet growing seasons through the slow-fast CWM

were weaker than that through the slow-fast FD. This is in contrast with a previous greenhouse

experiment showing that nutrient addition impacts resistance during flooding more through

functional trait composition rather than species richness (Oram *et al.*, 2020). Our study suggests

that compensatory dynamics among species may be more important than dominant species in

driving community resistance.

One year after dry or wet growing seasons, productivity often recovered to normal levels.

Although productivity generally deviated more under nutrient addition than ambient conditions

(Fig. S7), productivity changed in a similar manner during and one year after dry and wet

growing seasons in both conditions. Thus, nutrient addition did not impact resilience. Theories on

how functional trait composition and diversity would impact resilience are well developed, but

empirical tests lag behind (Reich, 2014; Díaz *et al.*, 2016; Craven *et al.*, 2018). To our

knowledge, no study has yet examined the relationships between slow-fast FD and resilience after

dry and wet growing seasons. Our analysis showed that, similar to resistance, functional trait

diversity and species richness were better associated with resilience than functional trait

composition. Additionally, both species richness and the slow-fast FD were negatively associated

with resilience after wet but not after dry growing seasons. Furthermore, this negative association

was mainly present in the nutrient addition but not ambient conditions (Fig. S10). This may be

because communities with higher diversity in slow-fast trait and higher species richness enhanced

resource use efficiency, resulting in higher productivity one year after wet growing seasons under

nutrient addition, which increased productivity deviation from normal levels and decreased

resilience (Wright *et al.*, 2015).

Our analyses confirmed the intuitive hypothesis that community invariability is jointly determined

by resistance and resilience (Fig. 4). Consistent with previous analyses, we found that nutrient

addition may decrease invariability through decreasing species richness (Hautier *et al.*, 2015;

Carroll *et al.*, 2021), but this indirect negative effect, together with the negative effects through

resilience were offset by the positive effects through resistance. Combined, nutrient addition did

not impact invariability.

Effects of nutrient addition on diversity facets and the weak diversity-mediated effects

Nutrient addition promoted fast communities while decreasing the slow-fast FD through

intraspecific trait variation, changing species richness and identities (Fig. 1F). Researchers often

focus on shared species (that occur both under ambient and nutrient addition conditions) to test

the relative importance of inter- and intraspecific trait shifts in shaping community traits (Fig. 1B,

C, D, and E). But under natural settings, some species may only be present under ambient or

nutrient addition conditions (Fig. 1B and F), and these species have rarely been taken into

account. We showed here that these species, though contributing to a small proportion of plant

communities (Table S3), can drive community trait change too (Fig. 3). Moreover, these

subordinate or rare species may still play a role in mediating nutrient effects on stability facets.

For instance, some links between CWM, FD and stability facets weakened when only considering

the shared species (Fig. S9).

When only considering shared species, change in slow-fast CWM was mainly induced by

intraspecific trait variation. Other studies partitioning CWM using shared species under other

global change factors reported similar results (Jung *et al.*, 2014; Duodu *et al.*, 2015; Niu *et al.*,

2016; Tatarko & Knops, 2018; Oram *et al.*, 2020). These results indicate that plant species can

adapt to multiple global change factors including flooding, drought, and nutrient enrichment

through a high level of phenotypic plasticity. Thus, ecological studies using traits measured in

situ, in which intraspecific trait variation is well captured, may improve the prediction for

community stability. At many NutNet sites, leaf traits were measured after 3 to 4 years of nutrient

addition, possibly resulting in higher intraspecific trait variation. Although there is a lack of

empirical test, in the longer term, species replacement, which can take years to manifest, may

become more important than intraspecific trait variation in determining community traits.

Using SEM, we evaluated the relative importance of species richness, slow-fast FD, and slow-fast

CWM in explaining stability under nutrient addition. FD and CWM were strongly confounded

(Fig. 4), it is difficult or even impossible to fully separate the effects of diversity from that of

composition under natural settings. Unexpectedly, we found overall weak diversity-mediated

indirect effects on stability. A few other studies also show that traits are poor predictors for

ecosystem properties including productivity and its stability (van der Plas *et al.*, 2020; Pichon *et*

al., 2022). These weak effects may be due to the following three reasons. First, plants have many

traits, although the slow-fast trait continuum representing the leaf economic spectrum is well

established and widely used in predicting community stability (de Bello *et al.*, 2021), we may

have missed some traits that can directly and strongly impact stability of productivity such as

hydrological traits (Schnabel *et al.*, 2021). Second, we did not measure traits for all plants, and

these unmeasured traits may strengthen or alter the results found here. Third and perhaps most

important, we did not consider intraspecific trait variation over time, which can be a major source

of community trait change and further impact the prediction of community traits on ecosystem

stability. This temporal intraspecific trait variation is ignored in many ecological studies using

traits to predict ecosystem functions and stability, likely because of the extreme effort required.

Digital cameras or spectrometers may be used to automatically record traits in the field, which

will help capture temporal intraspecific trait variation and greatly improve trait-based prediction

for stability.

Our study fills an important knowledge gap by showing that nutrient addition imposed different

impacts on different facets of stability, both directly and indirectly through mediating plant

diversity. However, the diversity-mediated indirect effects on facets of stability were weak and

overwhelmed by the direct effects of nutrient addition. This is not to say that diversity is not

important because diversity-mediated indirect effects and their relative importance, when multiple

diversity facets are considered simultaneously, on ecosystem stability remains unclear,

particularly under increased environmental perturbations. Thus, whereas preserving and restoring

plant diversity should still be one of the priorities in conservation and restoration, other

processes/mechanisms (e.g. changing biotic and abiotic growing environments) that impact

stability through pathways other than plant diversity should be carefully taken into account.

References

Arnoldi JF, Loreau M, Haegeman B. **2016**. Resilience, reactivity and variability: A mathematical comparison of ecological stability measures. *Journal of Theoretical Biology* **389**: 47–59.

de Bello F, Lavorel S, Hallett LM, Valencia E, Garnier E, Roscher C, Conti L, Galland T, Goberna M, Májeková M, *et al.* 2021. Functional trait effects on ecosystem stability: assembling the jigsaw puzzle. *Trends in Ecology and Evolution*: 1–15.

Bharath S, Borer ET, Biederman LA, Blumenthal DM, Fay PA, Gherardi LA, Knops JMH, Leakey ADB, Yahdjian L, Seabloom EW. **2020**. Nutrient addition increases grassland sensitivity to droughts. *Ecology* **101**: 1–11.

Borer ET, Grace JB, Harpole WS, MacDougall AS, Seabloom EW. **2017**. A decade of insights into grassland ecosystem responses to global environmental change. *Nature Ecology and Evolution* **1**: 1–7.

Borer ET, Harpole WS, Adler PB, Lind EM, Orrock JL, Seabloom EW, Smith MD. **2014**. Finding generality in ecology: A model for globally distributed experiments. *Methods in Ecology and Evolution* **5**: 65–73.

Carroll O, Batzer E, Bharath S, Borer ET, Campana S, Esch E, Hautier Y, Ohlert T, Seabloom EW, Adler PB, *et al.* 2021. Nutrient identity modifies the destabilising effects of eutrophication in grasslands (J Penuelas, Ed.). *Ecology Letters*.

Craven D, Polley HW, Wilsey B. **2018**. Multiple facets of biodiversity drive the diversity – stability relationship. *Nature Ecology & Evolution* **2**: 1579–1587.

Díaz S, Kattge J, Cornelissen JHC, Wright IJ, Lavorel S, Dray S, Reu B, Kleyer M, Wirth C, Colin Prentice I, *et al.* 2016. The global spectrum of plant form and function. *Nature* **529**: 167–171.

Donohue I, Petchey OL, Montoya JM, Jackson AL, Mcnally L, Viana M, Healy K, Lurgi M, O'Connor NE, Emmerson MC. 2013. On the dimensionality of ecological stability. *Ecology Letters* **16**: 421–429.

Duodu GO, Goonetilleke A, Allen C, Ayoko G. 2015. Determination of refractive and volatile elements in sediment using laser ablation inductively coupled plasma mass spectrometry. *Anal. Chim. Acta* **898**: 19–27.

Eskelinen A, Gravuer K, Harpole WS, Harrison S, Virtanen R, Hautier Y. 2020. Resource-enhancing global changes drive a whole-ecosystem shift to faster cycling but decrease diversity. *Ecology* **101**: e03178.

Fay PA, Prober SM, Harpole WS, Knops JMH, Bakker JD, Borer ET, Lind EM, MacDougall AS, Seabloom EW, Wragg PD, *et al.* 2015. Grassland productivity limited by multiple nutrients. *Nature Plants* **1**: 1–5.

Firn J, McGree JM, Harvey E, Flores-Moreno H, Schütz M, Buckley YM, Borer ET, Seabloom EW, La Pierre KJ, MacDougall AM, *et al.* 2019. Leaf nutrients, not specific leaf area, are consistent indicators of elevated nutrient inputs. *Nature Ecology and Evolution* **3**: 400–406.

Garnier, Eric, Cortez J, Billès G, Navas M-L, Roumet C, Debussche M, Laurent G, Blanchard A, Aubry D, Bellmann A, *et al.* 2004. Plant Functional Markers Capture Ecosystem Properties During Secondary Succession. *Ecology* **85**: 2630–2637.

Gonzalez A, Loreau M. 2009. The causes and consequences of compensatory dynamics in ecological communities. *Annual Review of Ecology, Evolution, and Systematics* **40**: 393–414.

Harris I, Osborn TJ, Jones P, Lister D. 2020. Version 4 of the CRU TS monthly high-resolution gridded multivariate climate dataset. *Scientific Data* **7**: 1–18.

Hautier Y, Tilman D, Isbell F, Seabloom EW, Borer ET, Reich PB. 2015. Anthropogenic environmental changes affect ecosystem stability via biodiversity. *Science* **348**: 336–340.

Hautier Y, Zhang P, Loreau M, Wilcox KR, Seabloom EW, Borer ET, Byrnes JEK, Koerner SE, Komatsu KJ, Lefcheck JS, *et al.* 2020. General destabilizing effects of eutrophication on grassland productivity at multiple spatial scales. *Nature Communications* **11**: 1–9.

Hillebrand H, Blasius B, Borer ET, Chase JM, Downing JA, Eriksson BK, Filstrup CT, Harpole WS, Hodapp D, Larsen S, *et al.* 2018. Biodiversity change is uncoupled from species richness trends: Consequences for conservation and monitoring. *Journal of Applied Ecology* **55**: 169–184.

Ingrisch J, Bahn M. 2018. Towards a Comparable Quantification of Resilience. *Trends in Ecology and Evolution* **33**: 251–259.

IPCC. (2019). Summary for policymakers. In P. R. Shukla, J. Skea, E. Calvo Buendia, V. Masson-Delmotte, H.-O. Pörtner, D. C. Roberts, P. Zhai, R. Slade, S. Connors, R. van Diemen, M. Ferrat, E. Haughey, S. Luz, S. Neogi, M. Pathak, J. Petzold, J. Portugal Pereira, P. Vyas, E. Huntley, K. Kissick, M. Belkacemi, & J. Malley (Eds.), Climate change and land: An IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems (p. 5). World Meterological Organization.

Isbell F, Craven D, Connolly J, Loreau M, Schmid B, Beierkuhnlein C, Bezemer TM, Bonin C, Bruelheide H, de Luca E, *et al.* 2015. Biodiversity increases the resistance of ecosystem productivity to climate extremes. *Nature* **526**: 574–577.

Ives AR, Carpenter SR. 2007. Stability and diversity of ecosystems. *Science* **317**: 58–62.

Jung V, Albert CH, Violle C, Kunstler G, Loucougaray G, Spiegelberger T. 2014. Intraspecific trait variability mediates the response of subalpine grassland communities to extreme drought events. *Journal of Ecology* **102**: 45–53.

Kéfi S, Domínguez-García V, Donohue I, Fontaine C, Thébault E, Dakos V. 2019. Advancing our understanding of ecological stability. *Ecology Letters* **22**: 1349–1356.

Laliberte E, Legendre P. **2010**. A distance-based framework for measuring functional diversity from multiple traits. *Ecology* **91**: 299–305.

Laliberté E, Legendre P, Shipley B. **2014**. Package 'FD'. : 85–91.

Lê S, Josse J, Husson F. **2008**. FactoMineR: An R package for multivariate analysis. *Journal of Statistical Software* **25**: 1–18.

Lefcheck JS. **2016**. piecewiseSEM: Piecewise structural equation modelling in r for ecology, evolution, and systematics. *Methods in Ecology and Evolution* **7**: 573–579.

Lepš J, de Bello F, Šmilauer P, Doležal J. **2011**. Community trait response to environment: disentangling species turnover vs intraspecific trait variability effects. *Ecography* **34**: 856–863.

Meng B, Li J, Maurer GE, Zhong S, Yao Y, Yang X, Collins SL, Sun W. **2021**. Nitrogen addition amplifies the nonlinear drought response of grassland productivity to extended growing□season droughts. *Ecology* **0**: 1–10.

Niu K, He JS, Lechowicz MJ. **2016**. Grazing-induced shifts in community functional composition and soil nutrient availability in Tibetan alpine meadows. *Journal of Applied Ecology* **53**: 1554–1564.

Oram NJ, De Deyn GB, Bodelier PLE, Cornelissen JHC, van Groenigen JW, Abalos D. **2020**. Plant community flood resilience in intensively managed grasslands and the role of the plant economic spectrum. *Journal of Applied Ecology* **57**: 1524–1534.

Pérez-Harguindeguy N, Díaz S, Garnier E, Lavorel S, Poorter H, Jaureguiberry P, Bret-Harte MS, Cornwell WK, Craine JM, Gurvich DE, *et al.* **2016**. Corrigendum to: New handbook for standardised measurement of plant functional traits worldwide. *Australian Journal of Botany* **64**: 715.

Pichon NA, Cappelli SL, Allan E. **2022**. Intraspecific trait changes have large impacts on community functional composition but do not affect ecosystem function. *Journal of Ecology*.

Pimm SL. **1984**. The complexity and stability of ecosystems. *Nature* **307**: 321–326.

Pinheiro J, Bates D, DebRoy S, Sarkar D, Team R-C. **2017**. nlme: Linear and Nonlinear Mixed Effects Models. *R package version 3.1-131*.

van der Plas F, Schröder-Georgi T, Weigelt A, Barry K, Meyer S, Alzate A, Barnard RL, Buchmann N, de Kroon H, Ebeling A, *et al.* **2020**. Plant traits alone are poor predictors of ecosystem properties and long-term ecosystem functioning. *Nature Ecology and Evolution* **4**: 1602–1611.

R Core Team. 2022. *R: A language and environment for statistical computing.* R Foundation for Statistical Computing.

Reich PB. **2014**. The world-wide 'fast-slow' plant economics spectrum: A traits manifesto. *Journal of Ecology* **102**: 275–301.

Sala OE, Gherardi LA, Reichmann L, Jobbágy E, Peters D. **2012**. Legacies of precipitation fluctuations on primary production: Theory and data synthesis. *Philosophical Transactions of the Royal Society B: Biological Sciences* **367**: 3135–3144.

Schnabel F, Liu X, Kunz M, Barry KE, Bongers FJ, Bruelheide H, Fichtner A, Härdtle W, Li S, Pfaff CT, *et al.* **2021**. Species richness stabilizes productivity via asynchrony and drought-tolerance diversity in a large-scale tree biodiversity experiment. *Science Advances* **7**: 11–13.

Seabloom EW, Adler PB, Alberti J, Biederman L, Buckley YM, Cadotte MW, Collins SL, Dee L, Fay PA, Firn J, *et al.* **2021**. Increasing effects of chronic nutrient enrichment on plant diversity loss and ecosystem productivity over time. *Ecology* **102**: 1–11.

Siefert A, Ritchie ME. **2016**. Intraspecific trait variation drives functional responses of old-field plant communities to nutrient enrichment. *Oecologia* **181**: 245–255.

Southon GE, Green ER, Jones AG, Barker CG, Power SA. **2012**. Long-term nitrogen additions increase likelihood of climate stress and affect recovery from wildfire in a lowland heath. *Global Change Biology* **18**: 2824–2837.

Tatarko AR, Knops JMH. **2018**. Nitrogen addition and ecosystem functioning: Both species abundances and traits alter community structure and function. *Ecosphere* **9**.

Tilman D. **1996**. Biodiversity: Population Versus ecosystem stability. *Ecology* **77**: 350–363.

Tilman D, Downing JA. **1994**. Biodiversity and stability in grasslands. *Nature* **367**: 363–365.

Tilman D, Isbell F, Cowles J. **2014**. Biodiversity and Ecosystem functioning. *Annual Review of Ecology, Evolution, and Systematics* **45**: 471–493.

Tilman D, Reich PB, Knops JMH. **2006**. Biodiversity and ecosystem stability in a decade-long grassland experiment. *Nature* **441**: 629–632.

Van Sundert K, Arfin Khan MAS, Bharath S, Buckley YM, Caldeira MC, Donohue I, Dubbert M, Ebeling A, Eisenhauer N, Eskelinen A, *et al.* **2021**. Fertilized graminoids intensify negative drought effects on grassland productivity. *Global Change Biology* **27**: 2441–2457.

Wang J, Knops JMH, Brassil CE, Mu C. **2017**. Increased productivity in wet years drives a decline in ecosystem stability with nitrogen additions in arid grasslands. *Ecology* **98**: 1779–1786.

Wheeler MM, Collins SL, Grimm NB, Cook EM, Clark C, Sponseller RA, Hall SJ. **2021**. Water and nitrogen shape winter annual plant diversity and community composition in near-urban Sonoran Desert preserves. *Ecological Monographs* **91**.

Wright AJ, Ebeling A, De Kroon H, Roscher C, Weigelt A, Buchmann N, Buchmann T, Fischer C, Hacker N, Hildebrandt A, *et al.* **2015**. Flooding disturbances increase resource availability and productivity but reduce stability in diverse plant communities. *Nature Communications* **6**: 1–6.

Xu Z, Ren H, Cai J, Wang R, Li MH, Wan S, Han X, Lewis BJ, Jiang Y. **2014**. Effects of experimentally-enhanced precipitation and nitrogen on resistance, recovery and resilience of a semi-arid grassland after drought. *Oecologia* **176**: 1187–1197.

Zhou X, Guo Z, Zhang P, Du G. **2018**. Shift in community functional composition following nitrogen fertilization in an alpine meadow through intraspecific trait variation and community composition change. *Plant and Soil* **431**: 289–302.

Acknowledgments

We thank researchers from the NutNet who contributed data to our analysis, supplementary Table S6 lists these researchers. We thank the Minnesota Supercomputing Institute for hosting project data and the Institute on the Environment for hosting Network meetings. Nitrogen fertilizer was donated to NutNet by Crop Production Services, Loveland, CO. This experiment is funded by individual researchers at the site scale.

Funding:

National Science Foundation grant NSF-DEB-1042132 (ETB, EWS; for NutNet coordination and data management)

National Science Foundation grant NSF-DEB-1234162 (ETB, EWS; for Long-Term Ecological Research at Cedar Creek).

National Science Foundation grant NSF-DEB-1831944 (ETB, EWS; for Long-Term Ecological Research at Cedar Creek)

National Natural Science Foundation of China grant 31988102, 32122053 (SW).

Author contributions:

Conceptualization: QC, SW, YH

Methodology: QC, SW, FI, S(i)B, JA, YH

Investigation: EWS, ETB, JDB, CR, CS, AE, CN, MCC, ASM, JLM, MT, KK

Visualization: QC

Supervision: YH, SW

Writing—original draft: QC, YH, SW

Writing—review & editing: QC, SW, EWS, FI, ETB, JDB, SB, CR, PLP, SAP, ID, CS, AE, CN, MCC, ASM, JLM, SB, AJ, MT, KK, JA, YH

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: Data are stored in Figshare (https://figshare.com/s/cae10e93c9029db4c347) and will be made publicly available once the manuscript is accepted.

Supplementary Materials for

Nutrient addition increases grassland resistance during dry and wet climate extremes despite functional diversity loss

Qingqing Chen[1], Shaopeng Wang[1*], Eric W. Seabloom[2], Forest Isbell[2], Elizabeth T. Borer[2], Jonathan D. Bakker[3], Siddharth Bharath[2], Christiane Roscher[4,5], Pablo Luis Peri[6], Sally A. Power[7], Ian Donohue[8], Carly Stevens[9], Anne Ebeling[10], Carla Nogueira[11], Maria C Caldeira[11], Andrew S. MacDougall[12], Joslin L. Moore[13], Sumanta Bagchi[14], Anke Jentsch[15], Michelle Tedder[16], Kevin Kirkman[16], Juan Alberti[17], and Yann Hautier[18]

Corresponding author: shaopeng.wang@pku.edu.cn

This PDF file includes:

Figs. S1 to S10
Tables S1 to S6

Fig. S1.

Categorize climate extremes at each site during experimental years. To reduce confounding effects of a dry and a wet growing season occurred consecutively, years were eliminated at a few sites. For instance, at comp.pt, year 1 was eliminated because year 0 had a dry growing season, which may bias resistance during a wet growing season in the next year. Grey dots represent years that were not included in analyses.

731
732

Fig. S2.

PCA using five leaf traits for 179 species from the ambient and nutrient enriched conditions across 19 sites.

Fig. S3.

Sensitivity test. We first simulated communities with full information (50 species with abundance ranging from 0 to 100 and slow-fast trait ranging from 0 to 8). We then randomly (1000 samplings) picked 1~48 species, ignoring their trait values (to simulate lack of trait data) to examine how much slow-fast CWM and FD were influenced by losing these species. Linear regressions with Tau 0.9 and 0.5 are shown in thick and thin lines, respectively. Simulation results suggested that the deviation would be smaller than 20 % when species with trait information contributed to > 60% cover.

Fig. S4.

Average aboveground biomass (g.m^{-2}) under different growing seasons. Growing seasons correspond to Fig. S1. Dots are means, error bars are 95% bootstrapped confidence intervals. Lines indicate aboveground biomass during normal growing seasons over experimental years over 19 sites. In general, biomass during moderate and extreme dry/wet growing seasons are similar in both treatments, except for that biomass increased during moderate wet growing seasons while decreased during extreme wet growing seasons in nutrient addition treatment.

Fig. S5.

Aboveground biomass (g.m^{-2}) over the experimental years in control and nutrient addition plots at each site.
Growing seasons correspond to Fig. S1. Dots indicate raw live biomass data averaged over three blocks in each year
at each site. Lines indicate average live biomass over normal growing seasons over three blocks at each site. Error
bars are 95% bootstrapped confidence interval.

Fig. S6.

Biomass difference (g.m^{-2}) from normal levels in control and nutrient addition treatments at each site. Dots indicate biomass difference (relative from normal level at each block) and then averaged over dry or wet growing season and over three blocks at each site. Error bars are 95% bootstrapped confidence interval.

Fig. S7.

Aboveground biomass (g.m^{-2}) and the magnitude of biomass deviation (g.m^{-2}) from normal levels in control (red dots and line) and nutrient addition plots (blue dots and line) under and one year after dry or wet growing seasons. Dots indicate average values over dry or wet growing seasons and over 19 sites. Lines indicate average live biomass under normal growing seasons over 19 sites. Error bars are 95% bootstrapped confidence interval.

Fig. S8.

Stability facets in control and nutrient addition plots at each site. Growing seasons correspond to Fig. S1. Dots indicate stability facets averaged over blocks and years at each site. Error bars are 95% bootstrapped confidence interval.

Fig. S9.

Direct and indirect effects of nutrient addition on stability facets. This figure is the same as
Fig. 4, but CWM and FD were calculated using the shared species. The model fits the data well (F
= 12.182, p = 0.273, df = 10). Boxes represent measured variables and arrows represent
relationships among variables. The displayed numbers are standardized path coefficients. The
width of the arrows indicates the strength of the pathways. Line color represents positive (black)
and negative (red) effects. Grey lines and text show correlated errors. Non-significant paths are
not shown. Asterisks indicate significant paths: * p ≤ 0.1; ** p ≤ 0.05; *** p ≤ 0.001. All stability
facets were on the log scale to improve normality and homogeneity of variance.
796

Fig. S10.

Relationships between each facet of plant diversity and stability. black lines indicate relationship over control and nutrient addition treatments as the interaction between plant diversity facet and treatment was not significant, treatment was dropped from the models. Solid lines show significant relationships, dashed lines show non-significant relationships.

 Table S1.

 Pathways and potential mechanisms for each link in SEM.

Pathways	Potential mechanisms	References
NPK->invariability	Nutrient addition may decrease invariability through decreasing resistance and resilience.	(Hautier *et al.*, 2015)
NPK->resistance.d	Nutrient addition may decrease resistance during dry climate extremes through increasing nutrients-to-water imbalance for plants, which reduces productivity and increases its deviation from normal levels.	(Xu *et al.* 2014; Bharath *et al.* 2020; Meng *et al.* 2021)
NPK->resistance.w	Nutrient addition may decrease resistance during wet climate extremes because increased nutrients and water availability increase plant growth and productivity, which also increase the magnitude of its deviation from normal levels.	(Wright *et al.*, 2015)
NPK->resilience.d	Nutrient addition may decrease resilience after dry climate extremes because increased nutrients-to-water imbalance increases plant mortality.	(Southon *et al.* 2012; Meng *et al.* 2021)
NPK->resilience.w	Increased productivity due to increased nutrients and water availability may persist or even amplify in later years, thus nutrient addition may decrease resilience after wet climate extremes.	(Sala *et al.* 2012; Wheeler *et al.* 2021)
CWM->ressistance.d, resistance.w, resilience.d, and resilience.w	Plant communities dominated by fast growing species may not respond fast during perturbations but may recovery fast after perturbations. That is, CWM decreases resistance during dry and wet climate extremes, but increases resilience after dry and wet climate extremes.	(Craven *et al.*, 2018; Oram *et al.*, 2020)
CWM->invariability	Plant communities dominated by fast growing species may vary more under perturbations, thus reducing invariability.	(Craven *et al.*, 2018; Lepš *et al.*, 1982)
FD->ressistance.d, resistance.w, resilience.d, and resilience.w	Communities with higher functional trait diversity may have a higher chance of harboring more species that are better adapted to these perturbations, thus decreases in some species may be compensated by increases in others. This leads to less productivity deviation from normal levels, which increases resistance and resilience.	NA
FD->invariability	Functional trait diversity increases compensatory dynamics and results in more complete use of resources, thus increasing invariability.	(Craven *et al.*, 2018; Polley *et al.*, 2013)
Species richness->resistance.d, resistance.w, resilience.d, and resilience.w	Species richness may capture functional trait diversity other than leaf economic spectrum such as root and size-related traits, thus higher species richness may increase resistance and resilience in a similar way to that of a higher FD.	(Craven *et al.* 2018).
Species richness->invariability	Increased number of species increases compensatory dynamics among species, which increases invariability.	(Hector *et al.* 2010)
Ressistance.d, resistance.w, resilience.d, and resilience.w->invariability	Both higher resistance and resilience indicate the proximity of a function from normal levels, higher resistance, higher resilience, or both contribute to higher invariability.	(Ives & Carpenter, 2007)
NPK->CWM	Nutrient addition increases CWM by promoting fast-growing species (with higher leaf nutrients and specific leaf area) and intraspecific trait variation.	(Siefert & Ritchie, 2016; Tatarko & Knops, 2018; Firn *et al.*, 2019)
NPK->FD	Nutrient addition decreases functional trait diversity because it increased the dominance of fast-growing species. Also, nutrient addition often decreases species richness, which is often but not always positively correlated with FD.	(De Bello *et al.* 2006; Borer *et al.* 2014)

NPK -> Species richness	Nutrient addition promotes nitrogen-loving plants, increasing dominance, decreasing light availability at the ground level, and thus reducing diversity due to loss of shade-intolerant species.	(Hautier *et al*. 2009)

307
308
309

Table S2.

19 sites selected and their geolocation and experimental years used. Growing season refers to from the start to the end of growing month at each site. Water balance is calculated as the mean over 15 growing seasons (2005-2020) at each site.

Site_code	Habitat	Continent	Latitude	Longitude	Growing season	Water balance	Pre-experiment year	Experimental years used for Fig. 2 - Fig. 3	Experimental years used for Fig. 4
bnch.us	montane grassland	North America	44.28	-121.97	4-8	-385.13	2008	1,3-11	1,3-11
bogong.au	alpine grassland	Australia	-36.87	147.25	10-1	-254.46	2010	1-10	1-10
burrawan.au	semiarid grassland	Australia	-27.73	151.14	10-5	-767.42	2009	1-11	1-11
cbgb.us	tallgrass prairie	North America	41.79	-93.39	5-10	-69.76	2010	1-8	1-8
comp.pt	annual grassland	Europe	38.83	-8.79	10-5	107.07	2013	2-8	
cowi.ca	old field	North America	48.81	-123.63	4-7	-178.57	2008	4-9	4-9
gilb.za	montane grassland	Africa	-29.28	30.29	10-4	-39.97	2011	1-3	
hopl.us	annual grassland	North America	39.01	-123.06	11-4	586.26	2008	1-12	1-12
jena.de	grassland	Europe	50.94	11.53	3-10	-250.06	2014	1-7	
kiny.au	semiarid grassland	Australia	-36.20	143.75	5-10	-176.02	2008	1-3	
lancaster.uk	mesic grassland	Europe	53.99	-2.63	3-8	111.23	2009	1-3	
look.us	montane grassland	North America	44.21	-122.13	3-8	-237.43	2008	2-9	2-9
mcla.us	annual grassland	North America	38.86	-122.41	11-4	355.41	2008	1,3-12	1,3-12
sage.us	montane grassland	North America	39.43	-120.24	4-7	-474.58	2008	1-6	1-6
sgs.us	shortgrass prairie	North America	40.82	-104.77	4-8	-506.28	2008	1-11	1-11
shps.us	shrub steppe	North America	44.24	-112.20	4-9	-628.74	2008	1-5	1-5
sier.us	annual grassland	North America	39.24	-121.28	11-4	501.67	2008	1-12	1-12
smith.us	mesic grassland	North America	48.21	-122.62	10-6	1406.28	2008	1-5	
unc.us	old field	North America	36.01	-79.02	4-9	-114.62	2008	2-4	

Table S3.

Information for leaf traits data at 19 sites used.

site_code	Years in which traits were measured	N(spp) with traits under ambient/nutrient addition	N(spp) occurred under both control and NPK	Proportional cover of species with traits under control/NPK	Proportional cover of shared species under control/NPK	Shift in CWM based on all species with traits	Shift in CWM based on shared species	Shift in CWM due to replacement among existing species	Shift in CWM due to intraspecific variation
bnch.us	4	6/9	6	0.66/0.87	0.66/0.84	0.53	0.50	0.01	0.49
bogong.au	2	6/12	6	0.82/0.81	0.82/0.65	0.86	0.48	0.09	0.40
burrawan.au	3	6/7	5	0.87/0.84	0.87/0.82	0.71	0.74	0.17	0.57
cbgb.us	3	9/17	7	0.77/0.91	0.74/0.71	0.26	0.18	-0.06	0.24
comp.pt	2	11/19	10	0.69/0.47	0.67/0.41	0.20	0.14	-0.15	0.29
cowi.ca	4	5/4	3	0.94/0.99	0.93/0.99	0.19	0.19	-0.03	0.22
gilb.za	2	4/7	4	0.48/0.56	0.48/0.44	0.32	0.28	0.05	0.23
hopl.us	4	9/14	6	0.53/0.86	0.49/0.62	1.31	0.52	0.12	0.40
jena.de	2	10/20	9	0.86/0.92	0.74/0.75	0.17	-0.11	-0.34	0.23
kiny.au	4	3/8	3	0.63/0.72	0.63/0.68	0.25	0.18	-0.09	0.27
lancaster.uk	3	5/7	5	0.8/0.91	0.8/0.84	0.42	0.52	0.04	0.48
look.us	4	6/10	4	0.82/0.88	0.71/0.82	-0.13	-0.41	-0.37	-0.04
mcla.us	4	8/13	8	0.83/0.93	0.83/0.79	0.42	0.42	0.19	0.23
sage.us	4	7/9	6	0.57/0.58	0.49/0.37	-0.55	-0.17	-0.09	-0.08
sgs.us	4	4/6	4	0.72/0.78	0.72/0.77	0.61	0.58	0.02	0.56
shps.us	4	5/10	3	0.59/0.7	0.44/0.53	0.81	0.29	0.02	0.27
sier.us	4	6/13	6	0.61/0.85	0.61/0.51	0.27	0.58	-0.05	0.63
smith.us	4	5/9	5	0.58/0.55	0.53/0.32	0.49	0.12	-0.21	0.33
unc.us	4	8/14	7	0.77/0.93	0.69/0.85	1.29	1.19	0.12	1.07

Table S4.

Model output for the fixed effects for Fig. 2. All stability facets were log-transformed to improve normality and homogeneity of variance.

variable	terms	Value	Std.Error	DF	t-value	p-value	r2m	r2c
invariability	(Intercept)	0.97	0.12	56.00	8.42	0.00	0.00	0.81
invariability	trtNPK	0.02	0.05	56.00	0.31	0.76	0.00	0.81
resistance.d	(Intercept)	0.99	0.13	53.00	7.45	0.00	0.03	0.27
resistance.d	trtNPK	0.32	0.14	53.00	2.24	0.03	0.03	0.27
resistance.w	(Intercept)	1.04	0.22	41.00	4.84	0.00	0.02	0.37
resistance.w	trtNPK	0.31	0.19	41.00	1.68	0.10	0.02	0.37
resilience.d	(Intercept)	0.73	0.26	50.00	2.76	0.01	0.00	0.36
resilience.d	trtNPK	-0.15	0.23	50.00	-0.66	0.51	0.00	0.36
resilience.w	(Intercept)	0.92	0.31	38.00	2.99	0.01	0.01	0.49
resilience.w	trtNPK	-0.29	0.23	38.00	-1.27	0.21	0.01	0.49

Table S5.

Model output for the fixed effects for Fig. 3.

variable	species	terms	Value	Std.Error	DF	t-value	p-value	r2m	r2c
Log (Species richness)	All species	(Intercept)	2.42	0.11	56.00	21.58	0.00	0.02	0.90
Log (Species richness)	All species	trtNPK	-0.16	0.03	56.00	-5.01	0.00	0.02	0.90
FD (slow-fast)	All species with traits	(Intercept)	0.59	0.05	56.00	11.61	0.00	0.05	0.62
FD (slow-fast)	All species with traits	trtNPK	-0.11	0.03	56.00	-3.66	0.00	0.05	0.62
CWM (slow-fast)	All species with traits	(Intercept)	-0.63	0.22	56.00	-2.91	0.01	0.05	0.90
CWM (slow-fast)	All species with traits	trtNPK	0.44	0.06	56.00	7.48	0.00	0.05	0.90
FD (slow-fast)	Shared species	(Intercept)	0.57	0.05	56.00	12.35	0.00	0.06	0.49
FD (slow-fast)	Shared species	trtNPK	-0.13	0.04	56.00	-3.68	0.00	0.06	0.49
CWM (slow-fast)	Shared species	(Intercept)	-0.62	0.22	56.00	-2.84	0.01	0.03	0.92
CWM (slow-fast)	Shared species	trtNPK	0.33	0.05	56.00	6.26	0.00	0.03	0.92
CWM induced by replacement	Shared species	(Intercept)	-0.62	0.23	56.00	-2.71	0.01	0.00	0.96
CWM induced by replacement	Shared species	trtNPK	-0.03	0.04	56.00	-0.80	0.43	0.00	0.96
CWM induced by ITV	Shared species	(Intercept)	-0.65	0.21	56.00	-3.12	0.00	0.04	0.96
CWM induced by ITV	Shared species	trtNPK	0.36	0.04	56.00	9.80	0.00	0.04	0.96

Table S6.
Principal investigators contributing data but are not authors; site names match that in Table S1.
Their effort in providing data is critical to this manuscript.

Site_code	Firstname	Lastname
bogong.au	John	Morgan
burrawan.au	Jennifer	Firn
cbgb.us	Lori	Biederman
cbgb.us	W.	Harpole
cbgb.us	Kirsten	Hofmockel
cbgb.us	Lauren	Sullivan
comp.pt	Miguel	Bugalho
gilb.za	Peter	Wragg
hopl.us	W.	Harpole
kiny.au	John	Morgan
mcla.us	W.	Harpole
sage.us	Daniel	Gruner
sage.us	Louie	Yang
sgs.us	Dana	Blumenthal
sgs.us	Cynthia	Brown
sgs.us	Julia	Klein
sgs.us	Alan	Knapp
shps.us	Peter	Adler
sier.us	W.	Harpole
smith.us	Janneke	Hille Ris Lambers
unc.us	Charles	Mitchell
unc.us	Justin	Wright