Pan-cancer genomic analysis shows hemizygous *PTEN* loss tumors are associated

with immune evasion and poor outcome

Vidotto T [1#], Melo CM [2#], Lautert-Dutra W [2], Chaves LP [2], Reis RB [3], Squire JA [* 2,4]

[#]*Joint first authors*

[1] *Department of Pathology, Johns Hopkins University School of Medicine, Baltimore,*

MD, USA

[2] *Department of Genetics, Medicine School of Ribeirão Preto, University of São Paulo,*

Ribeirão Preto, Brazil.

[3] *Division of Urology, Department of Surgery and Anatomy, Medicine School of*

Ribeirão Preto, University of São Paulo, Ribeirão Preto, Brazil.

[4] *Department of Pathology and Molecular Medicine, Queen's University, Kingston,*

Canada.

* Corresponding author: Jeremy Andrew Squire. E-mail: jsquireinsp@gmail.com

ABSTRACT

In tumors, somatic mutations of the *PTEN* suppressor gene are associated with advanced disease, chemotherapy resistance, and poor survival. *PTEN* loss of function may occur by inactivating mutation, by deletion, either affecting one copy (hemizygous loss) leading to reduced gene expression or loss of both copies (homozygous) with expression absent. Various murine models have shown that minor reductions in PTEN protein levels strongly influence tumorigenesis. Most *PTEN* biomarker assays dichotomize PTEN (i.e. presence vs. absence) ignoring the role of one copy loss. We performed a *PTEN* copy number analysis of 9,793 TCGA cases from 30 different tumor types. There were 419 (4.28%) homozygous and 2484 (25.37%) hemizygous *PTEN* losses. Hemizygous deletions led to reduced PTEN gene expression, accompanied by increased levels of instability and aneuploidy across tumor genomes. Outcome analysis of the pan-cancer cohort showed that losing one copy of *PTEN* reduced survival and was associated with transcriptomic changes controlling immune response and the tumor microenvironment. Immune cell abundances were significantly altered for *PTEN* loss, with changes in head and neck, cervix, stomach, prostate, brain, and colon more evident in hemizygous loss tumors. These data suggest that reduced expression of *PTEN* in tumors with hemizygous loss leads to tumor progression and influences anticancer immune response pathways.

Keywords PI3K/AKT signaling, haploinsufficiency; tumor suppressor gene; large deletions; somatic mutation; tumor microenvironment; genomic instability; gene dosage; immunotherapy; immune evasion; *in silico* analysis

INTRODUCTION

The PTEN tumor suppressor functions as a phosphatase, which metabolizes PIP3, the lipid product of PI3-Kinase that antagonizes the activation of the oncogenic PI3K/AKT/mTOR signaling axis [1,2]. In addition to its role as a primary regulator of cancer cell growth, recent findings also implicate PTEN in mediating aspects of the immune response against cancer [3,4].

PTEN is frequently lost, either partially affecting one copy of the gene (hemizygous loss; HemDel) or entirely affecting both copies of the gene (homozygous loss; HomDel), in a range of sporadic tumor types. A variety of mechanisms can cause *PTEN* gene inactivation, including missense and truncation mutations and deletions, as well as reduced expression of active PTEN protein mediated by promoter methylation, the effects of miRNAs, and the suppression of PTEN enzyme activity [5]. Loss of function of the *PTEN* tumor suppressor has been observed as one of the most common events in many types of cancer [6], having the greatest selection pressure for deletion based on an analysis of 746 human cancer genomes [7].

The *Pten* conditional knockout mouse model closely mimics human prostate initiation and progression [8]. Studies showed that loss of one copy of the *Pten* allele dramatically increases the rate of tumors [9]. It was proposed that haploinsufficiency of the *Pten* gene resulted in reduced levels of the functional protein that were insufficient for preventing tumor formation. Similar studies using a brain-tumor-specific model also showed that *Pten* haploinsufficiency accelerated the formation of glioblastomas [10]. Analysis of hypomorphic *Pten* mice demonstrated that as little as a 20% reduction of PTEN protein contributed to the development of prostate cancer in mice [11]. These and other mouse studies [12] show that minor reductions in PTEN protein levels can

profoundly influence tumorigenesis [13]. The findings from *Pten* murine tumor models are consistent with the precise cellular controls needed to regulate *PTEN* stability and function [14], implying that even subtle changes in *PTEN* expression may have profound effects on cancer development and progression [5].

Álvarez-Garcia et al. [15] recently surveyed the solid tumor literature on *PTEN* loss resulting from a mutation within the coding region of the gene; hemizygous or homozygous deletions of the gene as assessed by loss of heterozygosity (LOH), reduced copy number in array CGH analysis or loss of *PTEN* by FISH. Most tumor-associated missense mutations resulted in a loss or significantly reduced phosphatase activity or protein truncations. *PTEN* was found to demonstrate distinct patterns of somatic mutation leading to loss of function across different tumor types. Most glioblastomas display loss of function of *PTEN* in agreement with data from TCGA showing *PTEN* deletions in 143/170 (85%) of glioblastomas [16]. Other human tumor types found to have frequent *PTEN* deletions and/or loss of protein are cancers of prostate (25-50%), endometrium (50%), ovary (30-50%), lung (35-55%), breast (30-40%) and colon (10-40%) [17]. Moreover, it was found that single-copy inactivation of *PTEN* was far more common than mutation or deletion of both copies [15]. The PI3K/AKT/mTOR consortium study further suggested that partial copy number *PTEN* loss was associated with a less favorable outcome than complete loss across all available TCGA data [18]. These data, together with the murine *Pten* studies of tumorigenesis, formed the rationale for this PanCancer study that aimed to address the idea that partial loss of function of *PTEN* by hemizygous loss may confer a greater selective advantage for a subset of tumors than complete loss of function by homozygous loss (reviewed in [19]).

Our previous study examining the genomic impact of *PTEN* loss using TCGA data demonstrated that primary prostate tumors harboring *PTEN* deletions had enhanced

levels of aneuploidy and non-synonymous mutations [20]. However, it was unclear

whether the genomic instability generated by loss of one copy of *PTEN* was associated

with tumor progression. Other studies examining the impact of *PTEN* loss using TCGA

and other public domain datasets do not distinguish between one copy loss and

complete loss of the gene [21–24]. Similarly, most *PTEN* cancer biomarker studies using

immunohistochemistry and fluorescence *in situ* hybridization (FISH) have not been

rigorously standardized [15,17,25,26]. For these reasons, the impact of partial loss of function

due to one copy loss of *PTEN* is unknown for most cancers.

This study analyzed the TCGA copy-number and transcriptomic data, comparing

oncogenic and immunophenotypic features that distinguish hemizygous *PTEN* loss from

homozygous loss across the pan-cancer cohort. Our analysis also examines the

relationships between *PTEN* intact, hemi-, and homozygous loss with genomic features

of tumor progression and disease outcome.

RESULTS

The *PTEN* study group comprised 30 tumor types, including 9,793 cases analyzed from

the TCGA Pan-cancer cohort. *PTEN* somatic copy number alterations (SCNA) were

present in 3,619 (36.96%) tumors. Copy number deletions were observed in 2,903

(29.65%) tumors, with 2,484 (25.37%) having hemizygous deletions resulting in one

copy and 419 (4.28%) having homozygous loss of both copies. Copy number gains as a

duplication of the *PTEN* gene were evident in 716 (7.31%) tumors (Table 1).

Hemizygous *PTEN* deletions (HemDel) are more common than homozygous

deletions (HomDel) across the pan-cancer cohort. For most of the common solid

tumors, hemizygous *PTEN* deletions were much more frequent than homozygous

deletions (Figure 1). For example, melanoma (SKCM) has 205 (55.86%) hemizygous

deletion compared to 28 (7.63%) homozygous loss. Breast cancer (BRCA) has a

hemizygous deletion frequency of 269 (25.28%), but homozygous deletions are less

common at 54 (5.08%) of tumors. There are interesting differences in lung subtypes,

with squamous cell carcinoma (LUSC) having 217 (44.83%) hemizygous and 48

(9.92%) homozygous losses. However, adenocarcinoma (LUAD) had fewer

hemizygous deletions, 140 (27.50%) but only 5 (0.98%) homozygous losses. For

prostate cancer (PRAD), the rate of hemizygous loss was 73 (14.96%), slightly less than

the homozygous loss rate of 85 (17.42%). The four tumor types (KIRP, THYM, THCA,

and PCPG) with the lowest frequency of *PTEN* SCNA are characterized by a relatively

more favorable outcome in the literature. Pheochromocytoma and paraganglioma

(PCPG) are tumors of the autonomic nervous system. This tumor type has 11 (6.83%)

PTEN duplication gains suggesting instability leading to *PTEN* acquisition rather than

loss may be a feature of this tumor. Other tumors with more marked levels of *PTEN*

SCNA gain are ovarian (OV) 81 (19.66%), esophageal carcinoma (ESCA) 23 (12.71%),

lung adenocarcinoma (LUAD) 73 (14.34%), stomach adenocarcinoma (STAD) 48

(11.59%) and hepatocellular carcinoma of the liver (LIHC) 35 (9.70%).

PTEN expression levels closely correlate to gene copy number. To determine how

closely *PTEN* deletions and gains affect *PTEN* expression, we analyzed transcript

counts across the pan-cancer cohort to estimate gene expression levels related to *PTEN*

copy number (Figure 2a). We observed that tumors with *PTEN* HomDel had the lowest

expression levels, with mean transcript counts that were one order of magnitude lower

than the mean count observed in intact tumors (P<0.0001). The HemDel tumors showed

transcript counts lower than intact (P<0.0001), and *PTEN* gains were associated with

slightly higher expression levels than *PTEN* intact tumors (P<0.0001). We compared

PTEN intact to HemDel and HomDel in the cohort's 23 most common solid tumors

(Figure 2b). The same reduced expression pattern is apparent in all tumor types, with *PTEN* HomDel having the lowest expression levels for most tumors. Similarly, *PTEN* HemDel exhibited intermediate expression levels between HomDel and *PTEN* intact tumors. These data agree with the general correlations observed between gene copy number and RNA expression levels using TCGA transcriptomic data reported recently [27].

Associations between hemizygous *PTEN* deletions and poor outcome. Analysis of the overall survival of the entire cohort stratified by *PTEN* deletion status showed that tumors harboring *PTEN* HemDel had a worse outcome than those with HomDel and *PTEN* intact (Figure 3a). We next performed outcome analysis comparing *PTEN* intact to HomDel and HemDel for four solid tumors (Figure 3b-c and Supplementary Figure 1a-b) to illustrate the diversity of effects of *PTEN* loss on survival. For colorectal carcinomas and head and neck cancers, the proportion of tumors with HomDel *PTEN* loss was relatively low, so these patients were not considered for comparative studies. In brain tumors, the *PTEN* intact have a steadily decreased survival probability, but both *PTEN* HomDel and HemDel were associated with a rapid decline in survival probabilities (Figure 3b). In cervical tumors, there is a lower survival probability for *PTEN* HemDel compared to HomDel and intact (Figure 3c). However, in colorectal tumors, both *PTEN* HemDel and *PTEN* intact showed a worse outcome. In head and neck squamous cell carcinoma (Supplementary Figure 1a), the HemDel tumors were associated with reduced survival probabilities. In stomach tumors (Supplementary Figure 1b), HemDel was associated with intermediate survival risk. Further analysis considering the co-occurrence of *TP53* mutation showed that *PTEN* HemDel and *TP53* monoallelic mutation tumors were associated with reduced survival probabilities for the

entire cohort (Supplementary Table 1a). HNSCC tumors also showed worse outcomes

in *PTEN* HemDel and *TP53* monoallelic mutation (Supplementary Table 1b).

Relationships between hemizygous *PTEN* deletions and genomic instability. To

understand the impact of *PTEN* loss on genomic features of cancer progression, we

analyzed the total number of mutations, percent genome altered by losses and gains,

intratumor heterogeneity, aneuploidy, and homologous repair defects across the pan-

cancer cohort (Figure 4a-d). Our combined analysis revealed that cancers harboring

PTEN HemDel had the highest levels of nonsilent mutations, intratumor heterogeneity,

and percent genome altered. Moreover, PTEN HomDel and HemDel presented higher

levels of homologous repair defect compared with *PTEN* intact, but no difference was

found between the two deletion types. When stratified by tumor type, many of the

tumors harboring *PTEN* deletions had increased levels of nonsilent mutations,

intratumor heterogeneity, and homologous repair defects (Supplementary Figure 3a-d).

To compare the percentage of genome altered, we found that 65% (13/20) of the

investigated tumor types presented elevated genomic alterations when harboring *PTEN*

HemDel compared to *PTEN* Intact tumors (Supplementary Figure 3d). Pan-cancer

analysis showed aneuploidy levels were significantly elevated in HemDel compared to

HomDel and both deletion types showed significantly increased aneuploidy compared

to all intact tumors (Supplementary Figure 4a). Similar intratumor heterogeneity and

aneuploidy patterns were found in Dupl cases (Supplementary Figure 5a-b). Analysis of

aneuploidy in a subset of twenty solid tumors showed a general increase in aneuploidy

associated with PTEN deletion. Hemizygous losses in bladder, cervix, colon, head and

neck, liver, ovarian, pancreatic, and stomach cancer promoted increased levels of

aneuploidy (Supplementary Figure 4b). Collectively, these findings led us to

hypothesize that *PTEN* HemDel may be associated with an altered transcriptome,

potentially driving more genomic alterations.

Transcriptomic changes associated with *PTEN* loss activate immune pathways. We

next investigated the transcriptome of seven representative solid tumors with increased

frequencies of *PTEN* deletions and presented more diversity in genomic changes.

Differential gene expression analyses showed prostate tumors with *PTEN* HemDel

presented 2076 upregulated and 2966 downregulated genes compared to *PTEN* Intact

tumors (Supplementary Table 2). Prostate tumors with HemDel of *PTEN* presented

several enriched pathways compared with *PTEN* intact, including microtubule-based

movement and chromosome segregation (Supplementary Figure 6). *PTEN* HemDel was

also associated with distinct enrichment of immune pathways across the cancer types. In

prostate tumors with *PTEN* HomDel the high number of DEGs was linked to cell cycle,

DNA repair, and vascular process pathways (Supplementary Figure 7). Interestingly, for

HomDel in prostate cancer, the genes closely flanking *PTEN,* such as *ATAD1, RNLS*

and *KLLN,* are concurrently downregulated with the *PTEN* gene consistent with the

large genomic deletions of chromosome 10 previously observed in prostate cancer by

FISH [28] (see the vertical bar in Supplementary Figure 7).

Brain tumors presented the most distinct pathway enrichment results, including

the upregulation of several immune pathways when *PTEN* was deleted. We found 3562

up- and 10902 downregulated genes in these tumors when compared with *PTEN*

HemDel vs. *PTEN* intact (Supplementary Table 2, Supplementary Figure 8, 9). GO-

enriched analysis showed that several immune-related pathways were upregulated for

PTEN HemDel tumors, such as granulocyte migration and chemotaxis, neutrophil

migration, acute inflammatory response, humoral immune homeostasis, and many

others (Supplementary Figure 8). Similarly, when we compared the transcriptome of

tumors with *PTEN* HomDel vs. those with PTEN intact, we observed 4106 up- and

7332 downregulated genes (Supplementary Table 2, Supplementary Figure 9). We also

observed that several immune-related pathways were significantly upregulated in brain

tumors harboring *PTEN* HomDel, namely myeloid leucocyte migration, humoral

immune response, leucocyte chemotaxis, and acute inflammatory response

(Supplementary Figure 9). In summary, most investigated tumors had immune-related

pathways up- or downregulated when *PTEN* was lost. The RNA-sequencing expression

level of immunomodulatory genes like CD274 (PD-L1, Programmed Cell Death),

CTLA4 (Cytotoxic T-Lymphocyte Associated Protein 4), IDO1 (Indoleamine 2,3-

Dioxygenase 1), and LAG3 (Lymphocyte Activating 3) are altered in breast,

glioblastoma, head and neck, liver, ovarian, prostate, sarcoma, stomach, and uterine

tumors (Supplementary Figure 10). These findings led us to hypothesize whether *PTEN*

genomic deletions may be associated with an altered tumor microenvironment, as

determined from higher or lower abundance of immune cells surrounding cancerous

lesions.

Stomach tumors harboring *PTEN* HemDel show downregulated immune

pathways, such as adaptive immune response, regulation of lymphocyte activation, T

cell activation, positive regulation of immune response, leukocyte-mediated immunity,

lymphocyte proliferation, and several other pathways (Supplementary Figure 11).

Stomach tumors with *PTEN* HemDel also had downregulation of Indoleamine 2,3-

Dioxygenase 1 (*IDO1*) gene (Supplementary Figure 11). In contrast, activation of

immune pathways was not apparent in stomach tumors harboring *PTEN* HomDel, but

the sample size (N=20) for homozygous loss tumors was reduced compared to HemDel

tumors (N=90) (Supplementary Figure 12, Supplementary Table 2). For head and neck

tumors, we also observed that tumors with *PTEN* HemDel had downregulated immune-

related pathways, including adaptive immune response, activation of T-cell receptor,

regulation of lymphocyte-mediated immunity, regulation of leukocyte-mediated

immunity, and regulation of immune effector process (Figure 5a-c). However, these

results were not observed for the comparison between *PTEN* HomDel and *PTEN* intact,

in which the sample size from homozygous loss tumors was limited (N=15). However,

the linked passenger genes *MINPP1* and *KLLN* from chromosome 10 showed

concurrent reduced expression with *PTEN* (Supplementary Figure 13). Lastly, for colon

tumors, we found that tumors with *PTEN* HemDel had downregulated immune-related

pathways, including response to interferon-gamma, lymphocyte-mediated immunity,

adaptive immune response, and natural killer cell-mediated immunity (Supplementary

Figure 14). However, effects on immune pathways were not observed for the

comparison between *PTEN* HomDel and *PTEN* intact. As with other homozygous

losses, there was reduced expression of a group of *PTEN*-linked passenger genes

mapping to chromosome 10, including *ALDH18A1, DNMBP, KLLN, WAPL, MINPP1,*

and *ATAD1* (see the vertical bar in Supplementary Figure 15). Collectively, the pathway

analysis results draw attention to the role hemizygous *PTEN* loss may play in the

anticancer immune response.

PTEN deletions are associated with an altered tumor immune microenvironment.

The results from the tumor microenvironment comparison of *PTEN* HomDel, HemDel,

and intact tumors for all 22 immune cell types from brain, prostate, head and neck, and

stomach tumors were investigated. *PTEN* HemDel in brain tumors was linked to higher

CD4 T memory cells and M2 macrophage abundance and lower M1 macrophages,

monocytes, and CD8 T cells (Supplementary Figure 16a). Similar results were seen in

head and neck tumors, where *PTEN* HemDel showed higher M0 and M2 macrophages

and mast cells and lower CD4 T-cell, CD8+ T-cell, T-regulatory cells, and NK cells

(Figure 5d). While in prostate tumors with *PTEN* HemDel, we observed an increase in

M1 macrophages and a decrease in M0 macrophages and monocytes abundance

(Supplementary Figure 16b). Stomach tumors with *PTEN* HemDel had an increased

plasma and M0 macrophages density and decreased density of CD8 T-cells

(Supplementary Figure 16c). Cervical tumors with *PTEN* HemDel were linked to higher

M0 macrophages and an abundance of activated mast cells (Supplementary Figure 16d).

Similar results were seen in colorectal tumors, where *PTEN* HemDel showed a higher

M0 macrophage density and lower CD8 T-cell abundance (Supplementary Figure 16e).

DISCUSSION

The presence of the PTEN protein is essential for controlling the tumor

promoting activities of the PI3K/AKT pathway [2]. Deregulation of this signaling axis

confers a strong selective advantage to tumors, so that loss of *PTEN* control by

inactivating mutation or deletion is one of the most common somatic events in human

cancer [6]. PTEN mutation is significantly related to advanced disease, chemotherapy

resistance, and poor survival in patients with head and neck cancers, breast, melanoma,

colorectal, esophageal, and prostate [19,29–34] so that its role as a predictive and prognostic

cancer biomarker is increasingly gaining prominence [35–37]. Moreover, there is increasing

awareness of the role PTEN deficiency plays in shaping the immune landscape in

different cancers [3,4].

Various murine models have shown that the tumor-promoting abilities of loss of

the *Pten* gene are strongly dose-dependent [38]. In some models, the *Pten* gene loss has

been found to exhibit haploinsufficiency with reduced levels of the normal Pten protein

resulting in a diversity of tumors in mice [2]. Thus, in contrast to the classical definition

of a tumor suppressor gene that requires complete loss of both copies of the gene [39], it

appears that reduced amounts of PTEN protein caused by either monoallelic loss or by

one inactivating mutation is sufficient to cause cancer (reviewed in [19]).

301   Similarly, partial loss of function of PTEN activity has been observed amongst

302   *PTEN* hamartoma tumour syndrome (PHTS) patients [40]. These rare germline mutations

303   appear to have less severe developmental features, but it is unclear whether the cancer

304   susceptibility risk is altered in PHTS patients with partial PTEN activity.

305   A recent review across all solid tumors showed that hemizygous *PTEN* loss due

to monoallelic deletion or point mutation occurred at a much higher frequency than

biallelic loss [15]. Our *PTEN* analysis of TCGA data also shows that most tumors have a

much greater frequency of HemDel than HomDel and that hemizygous loss is less

favorable than homozygous loss for many tumor types. These data agree with previous

genetic and proteomic studies of PI3K/AKT/mTOR axis mutations using TCGA data

that showed outcomes associated with partial loss of *PTEN* were less favorable than

outcomes for complete loss of the gene [18]. Our analysis of *PTEN* RNA transcript levels

across the TCGA cohort shows good agreement with gene copy number, as reported

recently by others [27], so the HemDel losses we analyzed in this study would be expected

to reduce RNA expression levels. Moreover, since one somatic *PTEN* deletion or point

mutations leads to a reduction in RNA levels, then partial PTEN protein expression is

expected in HemDel tumors.

318   *PTEN* deletions and protein loss are known to affect DNA damage responses

and disrupt mitotic spindle architecture resulting in accumulation of mutations in other

genes and increased chromosome instability in both murine models and human cancers

[41]. As expected, our TCGA analysis showed both hemi- and homozygous losses

increased genomic instability features in many tumors. However, HemDel tumors had

significantly higher rates of non-silent mutations, intratumor heterogeneity, percentage

genome altered, and aneuploidy compared to HomDel tumors. This diversity of genetic

alteration is likely to facilitate genomic evolution and increase cancer progression,

treatment resistance, and poor outcome [42]. Collectively, these data suggest that the

elevated levels of genomic instability and somatic mutations in the tumors with

hemizygous loss could provide a selective advantage for the acquisition of other driver

genes and the less favorable outcomes observed across the TCGA cohort.

PTEN loss may be sufficient to cause tumorigenesis in some tissues but not in

others. A recent TCGA analysis of collaborating somatic mutations in *PTEN*-defective

solid tumors identified cancer genes such as *TP53, APC, TTN, MUC16, PIK3CA,*

BRAF, CDH1, and *KMT2D,* irrespective of the tumor type [37]. When surveying the

outcome in different cancers across the TCGA cohort, we found the consequences of

HemDel loss were more pronounced in some tumor types, suggesting that other cellular

contexts or unknown tumor-specific molecular features may contribute to progression

when *PTEN* is partially deficient.

The most obvious effects on outcome for HemDel *PTEN* losses were observed

in head and neck squamous cell carcinomas (HNSCC) harboring a mono-allelic somatic

loss-of-function mutation in *TP53*. For this type, the hemizygous loss was associated

with a less favorable overall survival with a significant HR of 1.67 (P=0.023)

(Supplementary table 1). *PTEN* loss is associated with poor outcomes in HNSCC [30].

HNSCC also has a high frequency of *TP53* somatic mutations [43]. We used cBioPortal to

investigate the co-occurrence frequencies of *TP53* mutations with *PTEN* intact,

HemDel, and HomDel in HNSCC. Interestingly, *PTEN* HemDel was significantly

(P<0.001) associated with *TP53* mutation, and *PTEN* HomDel was under-represented in

HNSCC with mutated *TP53* (data not shown). These data suggest that there may be

preferential selection for one copy of a *PTEN* gene in HNSCC so that reduced levels of

the functional PTEN protein allow tumor cells to bypass the *TP53*-induced cell

senescence pathway [44]. These observations in HNSCC are in keeping with the *PTEN*

literature that suggests that single-copy inactivation of *PTEN* may actually be selected

for during the progression of some tumor types since bi-allelic inactivation of *PTEN* has

been shown to lead to senescence or cell death when *TP53* is mutated [5]. Recent

functional studies of the relationship between *TP53* mutation and *PTEN* loss in HNSCC

indicate that loss of regulation of both pathways could influence radio- and

chemosensitivity [45]. These findings also agree with recent studies showing that *PTEN*

loss mediates resistance to cetuximab in HNSCC, indicating the need for more

treatment response *PTEN* predictive biomarker studies that also examines the role of

HemDel events [46].

Mouse models of HNSCC draw attention to the role of *Pten* in the evasion of

cellular senescence and activation of cancer-related inflammation [47]. Our gene set

enrichment analysis for HNSCC indicates that various pathways involved with immune

activation are downregulated in *PTEN* HemDel tumors. Pathways indicated in our

analysis were antigen presentation, interferon, interleukins, toll-like, neutrophil, B cell,

and inflammasome signaling. More marked alterations to the immune response in

HemDel HNSCC were also evident in our CIBERSORT analysis. There were decreased

CD8 T cells and alterations to tumor-associated macrophages (TAMs) consistent with

the immune tumor microenvironment being less favorable in hemizygous HNSCC.

These findings are consistent with the emerging recognition that *PTEN* loss could be a

useful predictive immune biomarker in HNSCC [48].

Biomarker analysis of colorectal cancer showed weak expression of the PTEN

protein in primary tumors with metastasis[33]. Our study found that *PTEN* HemDel

colorectal tumors had downregulated immune-related pathways. Functional activities

associated with immune response included regulation of granulocyte-macrophage colony-stimulating factor production, neutrophil migration, and regulation of leukocyte-mediated cytotoxicity. These findings are consistent with recent studies that showed that *PTEN* mutation was associated with microsatellite instability subtypes and tumor mutational burden in colon cancer [49]. Both *in vitro* studies [50] and recent analyses of colon tumors [51] suggest that *PTEN* loss in colorectal carcinoma may activate PI3K and upregulate *PD-L1* expression. These data are in keeping with other studies indicating that the PTEN/ PI3K axis may promote immune escape through regulating *PDL1/PD1* expression [3,52].

Brain tumors also presented distinct pathway enrichment findings, including the upregulation of several immune pathways for *PTEN* HemDel and HomDel losses. These data are consistent with a recent study showing *PTEN* loss in glioblastoma was associated with immunosuppressive expression signatures and failed response to anti-PD-1 immunotherapy [53].

Our TCGA analysis of prostate tumors with *PTEN* HemDel identified elevated levels of genomic instability and increased M1 macrophages but decreased abundances of M0 macrophages and monocytes. Pathway analysis showed enrichment for response to type I-interferon, consistent with other recent findings showing that *PTEN* loss leads to an immunosuppressive microenvironment in prostate cancer (PCa) [54]. These conclusions also agree with our previous study that showed increased IDO1 and Tregs in *PTEN*-deficient prostate cancer tumor microenvironment [55]. Interestingly, analysis of PCa showed that the genes flanking *PTEN,* such as *ATAD1, RNLS,* and *KLLN* are concurrently down-regulated when the *PTEN* gene is homozygously deleted. Similar observations were made in both colon and head and neck tumors with homozygous deletions. These data suggest that large somatic deletions of chromosome 10 may often

encompass closely linked genes whose expression changes could contribute to clinical

phenotypes [36].

Unfortunately, most previous *PTEN* cancer biomarker studies have not

systematically analyzed mutations or deletions using molecular genetic methods that

can distinguish between HemDel and HomDel mutations. Similarly, loss of the protein

using immunohistochemistry as an alternative way of determining PTEN status has also

not been studied using uniform, reproducible methods that can detect subtle changes in

protein levels [15,17,25,26]. The end result has been that majority of published *PTEN* cancer

biomarker studies have depended on a variety of simple two-way classification assays

(*PTEN* loss vs. intact) that ignore the distinction between hemi- and homozygous losses.

Our study also has some limitations that need to be addressed in future *PTEN*

analyses of the highlighted tumors. Our work was based entirely on *in silico* analysis of

copy number data derived TCGA, and all the pathway and immune parameters were

inferred from the associated transcriptome data. The analysis presented here cannot

fully capture the impact of PTEN mutations based on the limitations of the TCGA

dataset, which lacks prognostic or treatment information. Also, our expression analyses

associated with *PTEN* status were, like all *in silico* TCGA studies, obtained from the

bulk tumors containing heterogeneous tumor cells intermingled with a small subset of

stromal cells and varying quantities of immune cells. Lastly, our analysis regarding

PTEN status did not include *PTEN* inversions, fusions, inactivating somatic mutations,

promoter methylations, and large-scale structural variants, which may be a source of

bias in our evaluation of the impact of HemDel and HomDel effects.

Collectively, our findings based on TCGA, together with the extensive murine

Pten studies by others [8,9], suggest that loss of one copy of *PTEN* by mutation or deletion

may be sufficient to promote tumorigenesis and anti-cancer immune evasion. However,

for most human cancers, there have been limited biomarker studies on how hemizygous

PTEN deletion and partial protein loss may influences outcome and treatment response.

METHODS

Data download and processing

We initially downloaded raw and level 3 PanCancer normalized RNAseq, array-

CGH, and SNV from 10,713 samples derived from 31 tumor types from The Cancer

Genome Atlas (TCGA) cohort (www.gdc.cancer.gov). These initial cases were used to

determine outcome, genomic alterations, and relative immune abundances. For the

differential gene expression analysis (DEG), TCGA biolinks had 9793 cases were

available from 30 primary tumors as detailed in Table 1.

Raw RNAseq experiments were performed through Illumina HiSeq 2000 RNA

Sequencing platform. Copy number data were obtained from array-CGH experiments

performed with the Affymetrix Genome-Wide Human SNP Array 6.0. DNA sequencing

was performed in Illumina Genome Analyzer DNA sequencing. Clinical data for

progression-free intervals and overall survival were obtained for all patients from the

TCGA cohort. All analyses were conducted with data derived from primary tumors.

Defining *PTEN* deletions

PTEN deletions were characterized by using GISTIC-derived categorical SCNA.

In brief, GISTIC identifies a cutoff for copy number loss or gain using the log2

continuous data from each individual segments, such as genes and other regions [56].

These analyses were performed by the TCGA group for the PanCancer analysis. We

obtained, for our study, the Level 3 copy number calls for *PTEN* gene resulting from

whole genome detection of copy number changes or point mutations inactivating either

one or both copies of the gene [20]. The calls were defined as HomDel (two copies with

loss/mutation), HemDel (one copy loss or mutation), intact *PTEN* (both copies present

without mutation), and duplications of the *PTEN* gene [57]. *PTEN* duplications consisted

of the presence of gain of an extra copy of the entire gene or regions at 10q23.31 that

included *PTEN*. Our initial analysis of *PTEN* somatic mutations showed that in keeping

with the literature [15], this class of inactivation of the gene was infrequent in the TCGA

cohort (Supplementary Figure 2). We, therefore, choose to focus our analysis solely on

SCNA affecting *PTEN*.

Gene expression and enrichment analyses.

Analysis of gene expression was performed on the RNAseq level. Raw read

counts   for   TCGA-tumors   were   downloaded   using   FANTOM-CAT/recount2

(https://jhubiostatistics.shinyapps.io/recount/) [58,59]. The subset of samples was compiled

using each patient's clinical information, and only primary tumors were used in the

analysis. Study groups were divided based on *PTEN* status, HomDel (two copies with

loss/mutation), HemDel (one copy loss or mutation), and Intact *PTEN* (both copies

present without mutation). Differential expression (DE) analysis was carried out using

DESeq2 (v 1.32.0) [59,60], and *PTEN* status was used as the design factor. For enrichment

analysis of DE genes, we used the clusterProfile package (v 3.18.1) [61,62]. We used a P-

adjusted value < 0.05 as the cutoff for the enrichment results. All analyses were

conducted in RStudio software (R Foundation for Statistical Computing, R v4.1.2) and

all results were plotted by *pheatmap* (v 1.0.12) and *ggplot2* [64]. All results are displayed

with *PTEN* Intact as a reference for statistical analysis.

Immunogenic and genomic effects of PTEN deficiency

To determine the association between *PTEN* inactivation and acquired DNA

changes in tumors, we investigated the presence of genome doublings, homologous

recombination defects (HRD), aneuploidy levels, tumor mutational burden, and

intratumoral heterogeneity. These data were downloaded from the PanCanAtlas database

for The Immune Landscape of Cancer from TCGA (https://gdc.cancer.gov/about-

data/publications/panimmune) [65].Genome doubling status was determined through

ABSOLUTE algorithm [66], which measures tumor ploidy and purity based on copy number

and mutational signatures. Ploidy levels were determined by the quantity of DNA that each

cancer cell presents after undergoing several numerical and structural chromosome

aberrations. Clonality calls were employed to determine intratumor heterogeneity scores,

which determine tumor copy number and point mutations as aggregates of clonal and

subclonal components having varying ploidy levels. Tumor purity is a measurement that

considers the fractions of normal cells within the bulk mass of tumor cells. Aneuploidy

levels were calculated based on the total length of gained and deleted chromosome arms

divided by the genome size [67]. Similarly, HRDs were calculated based on the copy number

calls in with large (>15 Mb) non-arm-level chromosome imbalances with loss of

heterozygosity, breaks between genes larger than >10 Mb, and subtelomeric regions

harboring allelic imbalances. Groups with distinct *PTEN* inactivation statuses were

compared by employing the Kruskal-Wallis test in R.

Prognostic effect of PTEN-deficiency

To determine if *PTEN* HemDel and HomDel status has a significant impact in

predicting recurrence and death events, we performed survival analysis using *Survival*

package in R. Log-rank tests and Kaplan Meier curves were generated through the

Survival and *Survminer* packages in R. In addition, Cox Regression univariate models

were obtained for all tumors, grouped and separately by tumor type.

***In silico* analysis of immune abundance in tumor types**

To estimate the immune cell landscape in each tumor, we used the CIBERSORT deconvolution method (https://cibersort.stanford.edu/), which is an established tool to determine the abundance of immune cells using whole☐transcriptome data [68]. We separately imputed normalized gene expression data of brain, prostate, head and neck, stomach, cervical and colorectal tumors from the TCGA cohort. All results are displayed with *PTEN* Intact as a reference.

REFERENCES

1. Liu, T., Wang, Y., Wang, Y. & Chan, A. M. Multifaceted regulation of PTEN subcellular distributions and biological functions. *Cancers (Basel).* **11**, (2019).

2. Chalhoub, N. & Baker, S. J. PTEN and the PI3-kinase pathway in cancer. *Annu. Rev. Pathol.* **4**, 127–150 (2009).

3. Vidotto, T. *et al.* Emerging role of PTEN loss in evasion of the immune response to tumours. *Br. J. Cancer* **122**, 1732–1743 (2020).

4. Bezzi, M. *et al.* Diverse genetic-driven immune landscapes dictate tumor progression through distinct mechanisms. *Nat. Med.* **24**, 165–175 (2018).

5. Lee, Y., Chen, M. & Pandolfi, P. P. The functions and regulation of the PTEN tumour suppressor: new modes and prospects. *Nat. Rev. Mol. Cell Biol.* **19**, 547–562 (2018).

6. The ICGC/TCGA Pan-Cancer Analysis of Whole Genomes Consortium. Pan-cancer analysis of whole genomes. *Nature* **578**, 82–93 (2020).

7. Bignell, G. R. *et al.* Signatures of mutation and selection in the cancer genome. *Nature* **463**, 893–898 (2010).

8. Wang, S. *et al.* Prostate-specific deletion of the murine Pten tumor suppressor gene leads to metastatic prostate cancer. *Cancer Cell* **4**, 209–221 (2003).

9. Kwabi-Addo, B. *et al.* Haploinsufficiency of the Pten tumor suppressor gene

promotes prostate cancer progression. *Proc. Natl. Acad. Sci. U. S. A.* **98**, 11563–

11568 (2001).

10.   Kwon, C. H. *et al.* Pten haploinsufficiency accelerates formation of high-grade

astrocytomas. *Cancer Res.* **68**, 3286–3294 (2008).

11.   Alimonti, A. *et al.* Subtle variations in Pten dose determine cancer susceptibility.

Nat. Genet. **42**, 454–458 (2010).

12.   Shen-Li, H., Koujak, S., Szablocs, M. & Parsons, R. Reduction of Pten dose

leads to neoplastic development in multiple organs of PtenshRNA mice. *Cancer*

Biol. Ther. **10**, 1194–1200 (2010).

13.   Carracedo, A., Alimonti, A. & Pandolfi, P. P. PTEN level in tumor suppression:

How much is too little? *Cancer Res.* **71**, 629–633 (2011).

14.   Masson, G. R. & Williams, R. L. Structural mechanisms of PTEN regulation.

Cold Spring Harb. Perspect. Med. **10**, (2020).

15.   Álvarez-Garcia, V., Tawil, Y., Wise, H. M. & Leslie, N. R. Mechanisms of

PTEN loss in cancer: It's all about diversity. *Semin. Cancer Biol.* **59**, 66–79

(2019).

16.   Brennan, C. W. *et al.* The somatic genomic landscape of glioblastoma. *Cell* **155**,

462–77 (2013).

17.   Pulido, R. *et al.* Precise immunodetection of PTEN protein in human neoplasia.

Cold Spring Harb. Perspect. Med. **9**, 1–30 (2019).

18.   Zhang, Y. *et al.* A Pan-Cancer Proteogenomic Atlas of PI3K/AKT/mTOR

Pathway Alterations. *Cancer Cell* **31**, 820-832.e3 (2017).

19.   Berger, M. F. *et al.* The genomic complexity of primary human prostate cancer.

Nature **470**, 214–220 (2011).

20.   Vidotto, T., Tiezzi, D. G. & Squire, J. A. Distinct subtypes of genomic PTEN

deletion size influence the landscape of aneuploidy and outcome in prostate

cancer. *Mol. Cytogenet.* **11**, 1 (2018).

21.   Fan, C., Zhao, C., Shugen Li, F. W. & Wang, J. Significance of PTEN mutation

in cellular process, prognosis, and drug selection in clear cell renal cell

carcinoma. *Front. Oncol.* **9**, 1–9 (2019).

22.   Sun, J., Li, S., Wang, F., Fan, C. & Wang, J. Identification of key pathways and

genes in pten mutation prostate cancer by bioinformatics analysis. *BMC Med.*

Genet. **20**, 1–9 (2019).

23.   Imada, E. L. *et al.* Transcriptional landscape of PTEN loss in primary prostate

cancer. *BMC Cancer* **21**, 1–13 (2021).

24.   Lin, Z. *et al.* PTEN loss correlates with T cell exclusion across human cancers.

BMC Cancer **21**, 1–14 (2021).

25.   Jamaspishvili, T. *et al.* Clinical implications of PTEN loss in prostate cancer.

Nat. Rev. Urol. **15**, 222–234 (2018).

26.   Hocking, C. *et al.* Can we accurately report PTEN status in advanced colorectal

cancer? *BMC Cancer* **14**, 1–7 (2014).

27.   Shao, X. *et al.* Copy number variation is highly correlated with differential gene

expression: A pan-cancer study. *BMC Med. Genet.* **20**, 1–14 (2019).

28.   Yoshimoto, M. *et al.* PTEN genomic deletions that characterize aggressive

prostate cancer originate close to segmental duplications. *Genes, Chromosom.*

Cancer **51**, 149–160 (2012).

29.   da Costa, A. A. B. A. *et al.* Low PTEN expression is associated with worse

overall survival in head and neck squamous cell carcinoma patients treated with

chemotherapy and cetuximab. *Int. J. Clin. Oncol.* **20**, 282–289 (2015).

30.   Nagata, Y. *et al.* PTEN activation contributes to tumor inhibition by trastuzumab,

574 and loss of PTEN predicts trastuzumab resistance in patients. *Cancer Cell* **6**,

575 117–127 (2004).

576 31. Bucheit, A. D. *et al.* Complete loss of PTEN protein expression correlates with

577 shorter time to brain metastasis and survival in stage IIIB/C melanoma patients

578 with BRAFV600 mutations. *Clin. Cancer Res.* **20**, 5527–5536 (2014).

579 32. Tachibana, M. *et al.* Expression and Prognostic Significance of PTEN Product

580 Protein in Patients with Esophageal Squamous Cell Carcinoma. *Cancer* **94**,

581 1955–1960 (2002).

582 33. Sawai, H. *et al.* Loss of PTEN expression is associated with colorectal cancer

583 liver metastasis and poor patient survival. *BMC Gastroenterol.* **8**, 1–7 (2008).

584 34. Sircar, K. *et al.* PTEN genomic deletion is associated with p-Akt and AR

585 signalling in poorer outcome, hormone refractory prostate cancer. *J. Pathol.* **218**,

586 505–513 (2009).

587 35. Bazzichetto, C. *et al.* Pten as a prognostic/predictive biomarker in cancer: An

588 unfulfilled promise? *Cancers (Basel).* **11**, 1–22 (2019).

589 36. Poluri, R. T. K. & Audet-Walsh, É. Genomic deletion at 10q23 in prostate

590 cancer: More than PTEN loss? *Front. Oncol.* **8**, 1–7 (2018).

591 37. Fusco, N. *et al.* Pten alterations and their role in cancer management: Are we

592 making headway on precision medicine? *Genes (Basel).* **11**, 1–19 (2020).

593 38. Carnero, A. & Paramio, J. M. The PTEN/PI3K/AKT Pathway in vivo, cancer

594 mouse models. *Front. Oncol.* **4**, 1–10 (2014).

595 39. Wang, L. H., Wu, C. F., Rajasekaran, N. & Shin, Y. K. Loss of tumor suppressor

596 gene function in human cancer: An overview. *Cell. Physiol. Biochem.* **51**, 2647–

597 2693 (2018).

598 40. Leslie, N. R. & Longy, M. Inherited PTEN mutations and the prediction of

phenotype. *Semin. Cell Dev. Biol.* **52**, 30–38 (2016).

41.   Hou, S. Q., Ouyang, M., Brandmaier, A., Hao, H. & Shen, W. H. PTEN in the

maintenance of genome integrity: From DNA replication to chromosome

segregation. *BioEssays* **39**, 1–9 (2017).

42.   Salk, J. J., Fox, E. J. & Loeb, L. A. Mutational Heterogeneity in Human Cancers:

Origin and Consequences. *Annu. Rev. Pathol. Mech. Dis.* **5**, 51–75 (2010).

43.   Lawrence, M. S. *et al.* Comprehensive genomic characterization of head and

neck squamous cell carcinomas. *Nature* **517**, 576–582 (2015).

44.   Chen, Z. *et al.* Crucial role of p53-dependent cellular senescence in suppression

of Pten-deficient tumorigenesis. *Nature* **436**, 725–730 (2005).

45.   Vahabi, M. *et al.* MiR-96-5p targets PTEN expression affecting radio-

chemosensitivity of HNSCC cells. *J. Exp. Clin. Cancer Res.* **38**, 1–16 (2019).

46.   Izumi, H. *et al.* Pathway-specific genome editing of PI3K/mTOR tumor

suppressor genes reveals that PTEN loss contributes to cetuximab resistance in

head and neck cancer. *Mol. Cancer Ther.* **19**, 1562–1571 (2020).

47.   Bian, Y. *et al.* Loss of TGF-β signaling and PTEN promotes head and neck

squamous cell carcinoma through cellular senescence evasion and cancer-related

inflammation. *Oncogene* **31**, 3322–3332 (2012).

48.   dos Santos, L. V., Abrahão, C. M. & William, W. N. Overcoming Resistance to

Immune Checkpoint Inhibitors in Head and Neck Squamous Cell Carcinomas.

Front. Oncol. **11**, 1–16 (2021).

49.   Serebriiskii, I. G. *et al.* Comprehensive characterization of PTEN mutational

profile in a series of 34,129 colorectal cancers. *Nat. Commun.* **13**, (2022).

50.   Song, M. *et al.* PTEN Loss Increases PD-L1 Protein Expression and Affects the

Correlation between PD-L1 Expression and Clinical Parameters in Colorectal

Cancer. *PLoS One* **8**, (2013).

51.   Ugai, T. *et al.* Association of PIK3CA mutation and PTEN loss with expression

of CD274 (PD-L1) in colorectal carcinoma . *Oncoimmunology* **10**, 1956173

(2021).

52.   Lastwika, K. J. *et al.* Control of PD-L1 expression by oncogenic activation of the

AKT-mTOR pathway in non-small cell lung cancer. *Cancer Res.* **76**, 227–238

(2015).

53.   Zhao, J. *et al.* Immune and genomic correlates of response to anti-PD-1

immunotherapy in glioblastoma. *Nat. Med.* **25**, 462–469 (2019).

54.   Imada, E. L. *et al.* Transcriptional landscape of PTEN loss in primary prostate

cancer. *bioRxiv* 1–42 (2020) doi:10.1101/2020.10.08.332049.

55.   Vidotto, T. . *et al.* PTEN-deficient prostate cancer is associated with an

immunosuppressive tumor microenvironment mediated by increased expression

of IDO1 and infiltrating FoxP3+ T regulatory cells. *Prostate* **1**, 1–11 (2019).

56.   Mermel, C. H. *et al.* GISTIC2.0 facilitates sensitive and confident localization of

the targets of focal somatic copy-number alteration in human cancers. *Genome*

Biol. **12**, 1–14 (2011).

57.   Zack, T. I. *et al.* Pan-cancer patterns of somatic copy number alteration. *Nat.*

Genet. **45**, 1134–1140 (2013).

58.   Collado-Torres, L. *et al.* Reproducible RNA-seq analysis using recount2. *Nat.*

Biotechnol. **35**, 319–321 (2017).

59.   Imada, E. L. *et al.* Recounting the FANTOM CAGE-Associated Transcriptome.

Genome Res. **30**, 1073–1081 (2020).

60.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 1–21 (2014).

61.   Zhu, A., Ibrahim, J. G. & Love, M. I. Heavy-Tailed prior distributions for

650   sequence count data: Removing the noise and preserving large differences.

651   *Bioinformatics* **35**, 2084–2092 (2019).

652   62.   Liberzon, A. *et al.* The Molecular Signatures Database Hallmark Gene Set

653   Collection. *Cell Syst.* **1**, 417–425 (2015).

654   63.   Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting

655   omics data. *Innov.* **2**, 100141 (2021).

656   64.   Wickham, H. *ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New*

657   *York. Media* vol. 35 (2016).

658   65.   Thorsson, V. *et al.* The Immune Landscape of Cancer. *Immunity* **48**, 812-830.e14

659   (2018).

660   66.   Carter, S. L. *et al.* Absolute quantification of somatic DNA alterations in human

661   cancer. *Nat. Biotechnol.* **30**, 413–421 (2012).

662   67.   Wellenstein, M. D. & de Visser, K. E. Cancer-Cell-Intrinsic Mechanisms

663   Shaping the Tumor Immune Landscape. *Immunity* **48**, 399–416 (2018).

664   68.   Newman, A. M. *et al.* Robust enumeration of cell subsets from tissue expression

665   profiles. *Nat. Methods* 1–10 (2015) doi:10.1038/nmeth.3337.

DECLARATIONS

Authors' contributions: All authors contributed equally to the development of this

study. All authors read and approved the final manuscript.

Competing interests: The authors declare that they have no competing interests

Availability of data and materials: All data used in this analysis can be found at the

GDC data portal. The code and processed data were provided as supplementary

materials.

Ethics approval and consent to participate: Research Ethics Committee of the

Clinical Hospital of the Faculty of Medicine of Ribeirão Preto, number 3.088.034.

Funding: This research was funded by FAPESP, grant numbers 2019/22912-8 (JAS),

2021/12271-5 (WLD), and 2020/12816-9 (LPC) and by CNPq grant PQ-1D to JAS.

Figures

Figure 1. Relative percentages of *PTEN* Somatic Copy Number Alterations (*PTEN* SCNA) from the 23 most common solid tumors taken from the 30 tumor types analyzed in the Pan-cancer cohort (shown in Table 1). Hemizygous deletions (HemDel) = blue; Homozygous deletions (HomDel) = red, Intact (no *PTEN* SCNA) = green, and gain/duplication = brown. The tumors are ranked by their combined frequencies of PTEN deletion (HemDel + HomDel) with melanomas (SKCM) having the highest overall frequency of *PTEN* deletion (top) and Pheochromocytoma and Paraganglioma (PCPG) having the lowest (bottom). Duplication or *PTEN* gain (Dupl) was not included in the rank order determination. Plotted using *ggplot2* (https://ggplot2.tidyverse.org/).

Figure 2. (**a**) Combined analysis of the association between *PTEN* SCNA status and differential gene expression of PTEN for all 30 tumors in the Pan-cancer cohort. Boxplots show the relationship between PTEN normalized transcript counts Log^2 (Y-axis) versus *PTEN* SCNA status (HemDel, HomDel, Dupl and Intact). The mean transcript count for *PTEN* intact was 18.78, HomDel was the lowest expression level at 17.15, and HemDel showed reduced expression at 18.26, but Dupl was slightly higher than intact at 19.20. The mean transcript counts for all three classes of SCNA significantly deviated from intact (P<0.0001, t-test). (**b**) Association between *PTEN* SCNA status and differential gene expression of PTEN for the 23 most common solid tumors in the Pan-cancer cohort. Boxplots show the relationship between PTEN transcript counts Log^2 (Y-axis) versus *PTEN* SCNA status HomDel (red), HemDel (blue), and Intact (green) for each tumor. For most of tumors, the mean transcript counts for both HomDel and HemDel deviated significantly from intact (P<0.05). Plotted using *ggplot2* (https://ggplot2.tidyverse.org/).

Figure 3. Kaplan-Meier curves illustrate overall survival for both *PTEN* intact and *PTEN* deletions groups (HemDel and HomDel). Boxplots (**a**) Pan-cancer cohort shows a lower survival probability for *PTEN* HemDel compared to HomDel and intact (Kaplan-Meier, P<0.0001; *PTEN* HemDel HR: 2.02 (95% CI, 1.87-2.18), P<0.001; *PTEN* HomDel HR: 1.5 (95% CI, 1.27-1.77), P<0.001). (**b**) Brain tumors (GBM + LGG) show that both HomDel and HemDel are equally associated with earlier events (Kaplan-Meier, P<0.0001; *PTEN* HemDel HR: 8.38 (95% CI, 6.04-11.62), P<0.001; *PTEN* HomDel HR: 7.77 (95% CI, 4.7-12.86), P<0.001). (**c**) Cervical tumors show a lower survival probability for *PTEN* HemDel compared to HomDel and intact (Kaplan-Meier, P<0.0001; *PTEN* HemDel HR: 1.57 (95% CI, 0.92-2.66), P=0.1; *PTEN* HomDel HR: 0.88 (95% CI, 0.31-2.51), P=0.8). HR, Hazard Ratio. Plotted using *Survminer* (https://rpkgs.datanovia.com/survminer/index.html).

Figure 4. *PTEN* deletions are associated with (**a**) increased mutation levels, (**b**) intra-tumor heterogeneity (ITH), (**c**) homologous repair defects (HRD), and (**d**) percent genome alteration. Hemizygous deletions of *PTEN* are associated with increased levels of genomic alterations compared with homozygous deletions. Intact *PTEN* showed the lowest levels of genomic alterations. Plotted using *ggplot2* (https://ggplot2.tidyverse.org/).

Figure 5. *PTEN* HemDel in head and neck squamous cell carcinoma (HNSC) shows distinct transcriptome activity. (**a**) Transcriptome heatmap exhibiting clustering of top 50 DEGs (Log2Fold change >0.58). The color scale in the heatmap represents the Z-score of the normalized read counts for each gene, where the red scale color indicates upregulated, and blue low expressed genes. Enrichment analyses were performed using *clusterProfiler* and *p*-adjusted value = 0.05 as the cutoff. (**b**) Enriched GO (BP) pathways of the upregulated genes. (**c**) Enriched GO (BP) pathways of the downregulated genes. The tables shown in (b) and (c) indicate the top 10 most significant Gene Ontology (BP) terms. (**d**) CIBERSORT□derived immune□cell abundance of 22 distinct cell subsets based on *PTEN* status and head and neck tumors. Y□axis shows CIBERSORT relative scores. *PTEN* HemDel in head and neck tumors was linked to higher M0 and M2 macrophages, mast cells, and lower CD4 T, CD8 T, T regs, and NK cells. *: *p*-value < 0.05; **: *p*-value < 0.01, by Mann-Whitney test. n=492. DC, dendritic cell; NK, natural□killer cell; PCa, prostate cancer; Treg, regulatory T cell; M0, M0 macrophage; M1, M1 macrophage; M2, M2 macrophage. Plotted using *pheatmap* (https://rdocumentation.org/packages/pheatmap/versions/1.0.12).

Table 1. *PTEN* somatic copy number status (SCNA). HemDel: *PTEN* hemizygous deletion. HomDel: *PTEN* homozygous deletion. *Dupl* Duplication or gain across the Pan-cancer cohort of 30 tumors. ACC - Adrenocortical carcinoma, BLCA - Bladder Urothelial Carcinoma, BRCA - Breast invasive carcinoma, Brain (LGG - Brain Lower Grade Glioma combined with glioblastoma GBM), CESC - Cervical squamous cell carcinoma and endocervical adenocarcinoma, COAD - Colon adenocarcinoma, DLBC - Lymphoid Neoplasm Diffuse Large B - cell Lymphoma, ESCA - Esophageal carcinoma, HNSC - Head and Neck squamous cell carcinoma, KICH – kidney chromophobe, KIRC - Kidney renal clear cell carcinoma, KIRP - Kidney renal papillary cell carcinoma, LIHC - Liver hepatocellular carcinoma, LUAD - Lung adenocarcinoma, LUSC – Lung Squamous Squamous Cell Carcinoma, MESO - Mesothelioma, OV - Ovarian serous cystadenocarcinoma, PAAD - Pancreatic adenocarcinoma, PCPG - Pheochromocytoma and Paraganglioma, PRAD, Prostate Adenocarcinoma, READ - Rectum adenocarcinoma, SARC - Sarcoma, SKCM - Skin Cutaneous Melanoma, STAD - Stomach adenocarcinoma, TGCT - Testicular Germ Cell Tumors, THCA - Thyroid carcinoma, THYM - Thymoma, UCEC - Uterine Corpus Endometrial Carcinoma, UCS - Uterine Carcinosarcoma, UVM - Uveal Melanoma.

Figure 1

Figure 2

a

b

Figure 3

Figure 4

Figure 5

a

b

ID	Description	qvalue
GO:0002250	adaptive immune response	2,90E-53
GO:0002449	lymphocyte mediated immunity	1,34E-27
GO:0050778	positive regulation of immune response	2,30E-27
GO:0042110	T cell activation	2,75E-27
GO:0051249	regulation of lymphocyte activation	4,96E-22
GO:0002697	regulation of immune effector process	9,23E-20
GO:0007159	leukocyte cell-cell adhesion	3,95E-17
GO:0032943	mononuclear cell proliferation	9,05E-17
GO:0001819	positive regulation of cytokine production	1,26E-15
GO:0046651	lymphocyte proliferation	1,38E-15

d

c

ID	Description	qvalue
GO:0050804	modulation of chemical synaptic transmission	0.0012
GO:0099177	regulation of trans-synaptic signaling	0.0012
GO:0009953	dorsal/ventral pattern formation	0.0012
GO:0006749	glutathione metabolic process	0.0029
GO:0007409	axonogenesis	0.0097
GO:0071466	cellular response to xenobiotic stimulus	0.0116
GO:0030900	forebrain development	0.0183
GO:0007224	smoothened signaling pathway	0.0183
GO:0060571	morphogenesis of an epithelial fold	0.0183
GO:0060572	morphogenesis of an epithelial bud	0.0183

Table 1

TCGA Study	PTEN CNA Status								
	HemDel	%	HomDel	%	Dupl	%	Intact	%	Total
ACC	8	10,39	1	1,30	22	28,57	46	59,74	77
BLCA	153	37,87	14	3,47	30	7,43	207	51,24	404
BRCA	269	25,28	54	5,08	87	8,18	654	61,47	1064
Brain	235	35,34	22	3,31	2	0,30	406	61,05	665
CESC	74	25,43	14	4,81	17	5,84	186	63,92	291
COAD	86	19,72	16	3,67	15	3,44	319	73,17	436
DLBC	4	8,33	3	6,25	5	10,42	36	75,00	48
ESCA	61	33,70	7	3,87	23	12,71	90	49,72	181
HNSC	117	23,78	13	2,64	33	6,71	329	66,87	492
KICH	47	72,31	1	1,54	2	3,08	15	23,08	65
KIRC	86	17,03	5	0,99	11	2,18	403	79,80	505
KIRP	18	6,36	1	0,35	6	2,12	258	91,17	283
LIHC	82	22,71	13	3,60	35	9,70	231	63,99	361
LUAD	140	27,50	5	0,98	73	14,34	291	57,17	509
LUSC	217	44,83	48	9,92	35	7,23	184	38,02	484
MESO	25	28,74	2	2,30	3	3,45	57	65,52	87
OV	143	34,71	23	5,58	81	19,66	165	40,05	412
PAAD	30	17,05	1	0,57	13	7,39	132	75,00	176
PCPG	1	0,62	1	0,62	11	6,83	148	91,93	161
PRAD	73	14,96	85	17,42	6	1,23	324	66,39	488
READ	48	30,97	1	0,65	3	1,94	103	66,45	155
SARC	120	47,81	15	5,98	10	3,98	106	42,23	251
SKCM	205	55,86	28	7,63	6	1,63	128	34,88	367
STAD	90	21,74	20	4,83	48	11,59	256	61,84	414
TGCT	67	44,97	1	0,67	9	6,04	72	48,32	149
THCA	8	1,62	5	1,01	2	0,40	480	96,97	495
THYM	4	3,36	0	0,00	2	1,68	113	94,96	119
UCEC	53	10,23	19	3,67	108	20,85	338	65,25	518
UCS	19	33,93	1	1,79	18	32,14	18	32,14	56
UVM	1	1,25	0	0,00	0	0,00	79	98,75	80
	2484	25,37	419	4,28	716	7,31	6174	63,05	9793